High-resolution simulations of chromatin folding at genomic rearrangements in malignant B-cells provide mechanistic insights on proto-oncogene deregulation

Daniel Rico[*1], Daniel Kent[2], Aneta Mikulasova[1], Nefeli Karataraki[2], Rolando Berlinguer-Palmini[3], Brian A. Walker[4], Biola M. Javierre[5], Lisa J. Russell[*2], and Chris A. Brackley[*6]

[1] Biosciences Institute, Newcastle University, Newcastle upon Tyne NE2 4HH, UK
[2] Translational and Clinical Research Institute, Newcastle University, Newcastle upon Tyne NE2 4HH, UK
[3] Bio-Imaging Unit, Newcastle University, Newcastle upon Tyne NE2 4HH, UK
[4] Melvin and Bren Simon Comprehensive Cancer Center, Division of Hematology Oncology, Indiana University, Indianapolis, IN, USA
[5] Josep Carreras Leukaemia Research Institute (IJC), IJC Building, Campus ICO-Germans Trias i Pujol, Ctra de Can Ruti, Camí de les Escoles s/n, 08916 Badalona, Barcelona, Spain
[6] SUPA, School of Physics and Astronomy, University of Edinburgh, Peter Guthrie Tait Road, Edinburgh EH9 3FD, UK
[*] Senior authors. Address correspondence to: C.Brackley@ed.ac.uk, lisa.russell@newcastle.ac.uk, or daniel.rico@newcastle.ac.uk

Genomic rearrangements are known to result in proto-oncogene deregulation in many cancers, but the link to genome 3D structure remains poorly understood. Here we used the highly-predictive heteromorphic polymer (HiP-HoP) model to predict chromatin conformations at the proto-oncogene *CCND1* in healthy and malignant B-cells. After confirming that the model gives good predictions of Hi-C data for the non-malignant human B-cell derived cell line GM12878, we generated predictions for two cancer cell lines, U266 and Z-138. These possess genome rearrangements involving *CCND1* and the immunoglobulin heavy chain locus (*IGH*), which we mapped using targeted genome sequencing. Our simulations showed that a rearrangement in U266 cells where a single *IGH* super-enhancer is inserted next to *CCND1* leaves the local topologically associated domain (TAD) structure intact. We also observed extensive changes in enhancer-promoter interactions within the TAD, suggesting that it is the downstream chromatin remodelling which gives rise to the oncogene activation, rather than the presence of the inserted super-enhancer DNA sequence *per se*. Simulations of the *IGH-CCND1* reciprocal translocation in Z-138 cells revealed that an oncogenic fusion TAD is created, encompassing *CCND1* and the *IGH* super-enhancers. We predicted how the structure and expression of *CCND1* changes in these different cell lines, validating this using qPCR and fluorescence *in situ* hybridization microscopy. Our work demonstrates the power of polymer simulations to predict differences in chromatin interactions and gene expression for different translocation break-points.

Chromatin structure, nuclear organisation and the epigenome are intimately linked to gene regulation and cell function, and are tightly controlled during differentiation. Genome rearrangements leading to structural variants (including deletions, insertions, duplications and translocations) can disturb these, leading to gene mis-regulation and cancer (Norton and Phillips-Cremins, 2017). An important example occurs during B-cell differentiation, where programmed breakage and recombination of the genome takes place in order to generate the broad heterogeneity of immunoglobulins (Igs) required for immune system function (Jung *et al.*, 2006). Errors in this process can lead to repositioning of Ig regulatory elements which then drives proto-oncogene activation (Kontomanolis *et al.*, 2020). The diversity in translocations and the difficulties in accessing and handling patient samples means that detection, accurate mapping, and characterisation of their functional consequences are inherently problematic.

Advances in molecular probes of genomic and epigenomic structure, such as ChIP-seq and chromosome-conformation-capture methods like Hi-C, have transformed our understanding of the regulatory link between structure and function. For example, Hi-C, which probes chromosome interactions genome-wide, has revealed that the genome can be partitioned into topologically associated domains (TADs) (Dixon *et al.*,

2012). These are contiguous chromosome regions which show enriched self-interactions, and are thought to be associated with cis-regulatory mechanisms; they are often bounded by binding sites for the CCCTC-binding factor CTCF (Rao *et al.*, 2014). ChIP-seq profiling of histone modifications and protein binding has identified super-enhancers, clusters of enhancer elements thought to drive expression patterns responsible for cell identity (Hnisz *et al.*, 2013). Super-enhancers have been found to preferentially sit within TADs which are highly insulated from their surroundings (Gong *et al.*, 2018), and have been identified as drivers of oncogene expression in many tumours (Thandapani, 2019; Mikulasova *et al.*, 2020b).

A full epigenetic and three-dimensional (3D) structural characterisation of genome rearrangements is crucial for our understanding of proto-oncogene activation. However, the small size of patient samples and their inherent variability means that routinely applying methods like Hi-C to primary cancer samples remains challenging, despite recent progress in reducing the amount of material required (Díaz *et al.*, 2018). This is why much work to date has focused on the biogenesis of chromosome translocations (Engreitz *et al.*, 2012; Roukos and Misteli, 2014), but comparatively less attention has been paid to the structural *consequences* of translocations and other genome rearrangements (Bianco *et al.*, 2018; Akdemir *et al.*, 2020). In

A. Wild Type (GM12878 cell line)

B. Multiple Myeloma (U266 cell line)

C. Mantle Cell Lymphoma (Z-138 cell line)

Figure 1. Cartoons showing the layout of the *CCND1* and *IGH* loci in **A.** Healthy cells. For *CCND1* we show the approximate positions and orientations of four genes within the TAD (*TPCN2* lies just outside the TAD which encloses the other three). For *IGH* we show four previously identified enhancer elements within the constant region. **B.** A rearrangement which has been observed in multiple myeloma (and in the U266 cell line) is an *IGH* insert bringing an enhancer element into the vicinity of *CCND1*. **C.** A t(11;14) translocation often found in mantle cell lymphoma (and in the Z-138 cell line) brings *CCND1* next to the entire *IGH* constant region.

this paper we show how computer simulations can help us to understand the effects of common genome rearrangements in B-cell malignancies, and how these lead to the changes in the 3D structure of gene loci which in turn leads to deregulation.

We used our "highly-predictive heteromorphic polymer" (HiP-HoP) model (Buckle *et al.*, 2018; Brackey *et al.*, 2020) to study genomic rearrangements involving the Ig heavy chain locus (*IGH*) (Watson and Breden, 2012) and the *CCND1* proto-oncogene encoding the cell cycle protein cyclin D1. *IGH-CCND1* is a common translocation observed in mantle cell lymphoma (MCL) and multiple myeloma (MM); when translocated together the *IGH* super-enhancers drive *CCND1* overexpression. Figure 1 shows a cartoon of the two loci in healthy cells and with two different rearrangements which are studied here; specifically we consider an MM cell line, U266, where a single *IGH* super-enhancer is inserted into the *CCND1*-TAD, and an MCL cell line, Z-138, possessing a translocation which repositions *CCND1* next to the *IGH* joining region (Watson and Breden, 2012). In previous work (Mikulasova *et al.*, 2020b) we analysed the changes in patterns of histone modifications associated with super-enhancer translocation events, finding that relocation of the *IGH* super-enhancer results in local chromatin remodelling and the emergence of new patterns of histone modifications including H3K4me3 broad domains (Park *et al.*, 2020). Here we study how these epigenetic changes alter 3D chromosome structure.

HiP-HoP uses data on DNA accessibility, histone modifications, and protein binding to generate an ensemble of 3D chromosome structures. We previously showed that the method gives good predictions of both population level data (*e.g.*, Hi-C or 4C) and single cell measurements [*e.g.*, from fluorescence microscopy; Buckle *et al.* (2018)]. Importantly, HiP-HoP does not use Hi-C (or any other 3D genome data) as an input [unlike other popular models which use fitting-based methods; Serra

et al. (2015)]. This means that we can make predictions about the 3D structure of cell types or tissues where Hi-C data is not available.

Here we found that HiP-HoP gives good predictions of Hi-C data at the *CCND1* locus in a human cell line derived from healthy B-cells. Simulations of the MM cell line U266 (Fig. 1B) predicted that while the TAD remains intact, chromatin interactions among gene promoters and emergent regulatory regions are radically altered. Predictions for the MCL cell line Z-138 (Fig. 1C) showed an oncogenic fusion TAD encompassing *CCND1* and four previously identified *IGH* super-enhancer regions (Mills *et al.*, 1983; Chen and Birshtein, 1997; Mills *et al.*, 1997; Mikulasova *et al.*, 2020b). Further simulations suggest that it is the downstream epigenetic changes which drive the 3D changes, rather than the genomic rearrangements itself. This points to a scenario where the translocated super-enhancers first recruit chromatin remodellers to the region; this then drives epigenomic changes which in turn disrupts the wider 3D structure leading to overexpression of *CCND1*. Importantly, the simulations allowed us to investigate different possible rearrangements, to infer the pathway through which the *CCND1* proto-oncogene becomes activated, and to generate testable hypotheses which will direct new experiments suggesting targets for therapeutic intervention. This points to a future role for the method in a clinical setting, *e.g.* in personalised medicine, where structural predictions based on available or easily obtainable data could be extremely valuable.

Results

HiP-HoP predicts the domain structure at *CCND1* in a B-cell derived cell line

We previously showed that in malignant B-cells, genome rearrangements involving *CCND1* lead to local changes in histone modifications (Mikulasova *et al.*, 2020b). Here we hypothesised that this in turn leads to changes in 3D structure and chromatin interactions which drive *CCND1* expression. To understand these malignant structural changes it is first necessary to study the healthy case. To this end we considered the human B-cell derived lymphoblastoid cell line GM12878, adapting our HiP-HoP method to study chromatin structure around *CCND1*. This is an ideal cell line due to the abundance of publicly available data.

In the model, a 3 Mbp chromosome region was represented by a chain of beads [a common approach in polymer physics; Brackey *et al.* (2020)]. The region was chosen because it is large enough to encompass seven TADs surrounding *CCND1* such that the local chromatin context is captured, but small enough to give feasible simulation run times. The model combines three mechanisms which drive locus structure: (i) diffusing spheres representing complexes of chromatin binding proteins which can bind to multiple points at the same time to form molecular bridges (*e.g.*, between promoters and enhancers) (Brackley *et al.*, 2013, 2016b; Brackley, 2020); (ii) a heteromorphic polymer structure, where different sections of the bead chain have different properties (thickness/compaction and flexibility); and (iii) the loop extrusion mechanism (Fudenberg *et al.*, 2016; Sanborn *et al.*, 2015). Loop extrusion is a molecular mechanism where chromatin is pushed onto loops by factors [the cohesin complex; Davidson *et al.* (2019)], which are halted by CTCF proteins bound in a convergent orientation (Rao *et al.*, 2014).

A way to identify different types of chromatin is via high-

Figure 2. A. A map of the genes in the vicinity of *CCND1* is shown above heat maps for: (i) Publicly Hi-C data for GM12878 cells; and (ii) simulated Hi-C from the same region (obtained from 4400 independent configurations). TAD boundary positions are indicated with blue dashed lines; green dashed lines indicate a "subTAD" region (see text). Some unmappable regions are visible as gaps in the data (white stripes). **B.** Simulation snapshots of the 3D structure of the *CCND1* region, including the *CCND1*-TAD (pale orange) and the two neighbouring TADs (pale green and blue) (a 1.3 Mbp section of the 3 Mbp region simulated is shown). One representative snapshot is shown large, with five other smaller examples. Several genes are shown in bright colours as indicated. For clarity proteins are not shown. Box: the same configuration is shown but coloured according to the input data used in the simulation, as indicated by label colours; gene positions are circled. **C.** Simulated 4C interaction profiles are shown from four viewpoints [blue triangles; these are positioned at promoters of *TPCN2*, *MYEOV*, *CCND1* and *LTO1*, and at a DNase hypersensitive site (DHS) downstream of *MYEOV*]. Data used as simulation input [obtained from ENCODE; see Supplemental Methods and Supplemental Table S3] is shown as coloured blocks above the plots. Red blocks indicate DHS, used to infer binding sites for active proteins. Blue and black blocks indicate regions with chromatin states corresponding to polycomb and heterochromatin respectively, used to infer binding for the corresponding proteins. Yellow blocks indicate regions with chromatin states associated with H3K27ac, and indicate regions which have a more open chromatin structure in the model. Orange and purple arrowheads indicate the position and directionality of CTCF sites, used to infer loop extruder anchors. Some features are labelled with numbered arrows as referred to in the text.

throughput analysis of histone modification data and hidden Markov modelling (HMM) (Ernst *et al.*, 2011; Carrillo-de Santa-Pau *et al.*, 2017); this approach classifies chromosome regions into a number of states (*e.g.*, promoter/enhancer associated, polycomb repressed, and heterochromatin states, see Supplemental Table S1). In the model we had two different types of chromatin structure (the heteromorphic polymer): more open regions (thinner, more flexible polymer) and more compact regions (thicker, "crumpled" polymer, see Supplemental Fig. S1 for a schematic). We used the chromatin states to identify these regions: H3K27ac associated states have the more open structure (Risca *et al.*, 2017). The model included three different species of bridge forming protein: a general "active binder", and two types of repressive binders [this is an extension of the scheme in Buckle *et al.* (2018) which only included active proteins]. To identify active protein binding sites we used DNA accessibility measured via DNase-seq experiments, assuming that DNase-hypersensitive sites (DHS) are binding sites (see Methods and Supplemental Methods). Repressive protein binding sites were identified using the chromatin states, with one species binding to H3K9me3 states, and

one species representing polycomb repressive complexes and binding to H3K27me3 states. Finally, ChIP-seq for CTCF was used to identify "anchor" sites where loop extrusion is halted. Full details of the simulations are given in Supplemental Methods, and a schematic is shown in Supplemental Fig. S1.

To verify that HiP-HoP gives good predictions for *CCND1*, we generated simulated Hi-C interaction maps and compared them with publicly available data (Rao *et al.*, 2014) (Fig. 2A; Supplemental Fig. S1 shows a similar plot but for the entire 3 Mbp region which was simulated, along with the experimental data used as an input). The simulations gave a good prediction of the TAD pattern across the region. *CCND1* sits at the far right of a TAD which is bounded by convergent CTCF sites, starts just to the left of *TPCN2*, and ends between *LTO1* and *FGF19*. We shall refer to the three neighbouring domains as the *TPCN2*, *CCND1*, and *FGF19*-TADs. The gene *MYEOV* sits just left of the centre of the *CCND1*-TAD.

Simulated Hi-C maps tend to show more structure within the domains than experiments (visible as dark spots and stripes); typically, these features arise in the model due to protein medi-

ated enhancer-promoter interactions. This was also observed in previous HiP-HoP studies, and is likely because the experimental data lacks sufficient resolution to reveal these features. Many of the predicted 'within-TAD' interactions are present in publicly available Hi-ChIP and ChIA-PET data which has higher resolution [Supplemental Fig. S1; promoter-enhancer interactions are also more apparent in targeted 4C (Zhao *et al.*, 2006), nucleosome resolution MicroC (Krietenstein *et al.*, 2020), and recent Hi-C data which has been treated using a method for removing biases (Lu *et al.*, 2020)]. In the simulated Hi-C several smaller 'subTADs' are also visible; for example, one subTAD is generated by looping interactions between a CTCF site just upstream of *MYEOV* and a CTCF within the *CCND1* gene body [green dashed lines in Fig 2A(ii)]. A typical simulated structure is shown in Fig. 2B; also shown are 5 other representative structures (from a population of 4400 generated).

A more focused view of chromatin interactions at gene promoters is shown in simulated 4C interaction profiles [Fig. 2C; we take a 1 kbp region immediately upstream of a gene's transcription start site (TSS) to be its promoter]. These reveal that the *CCND1* promoter often interacts with polycomb associated regions (arrow 'i' in the figure); this is because there is enrichment of H3K27me3 at the promoter. There are also 'active' H3K4me1/3 modifications, which could indicate variation between alleles or across the population, with the gene sometimes expressed and sometimes repressed. Consistent with this, publicly available RNA-seq data (The ENCODE Project Consortium, 2012) shows weak expression of *CCND1* in GM12878 cells (see Supplemental Methods).

To more quantitatively assess the expression of four of the genes within the locus we performed qPCR experiments (see Supplemental Methods and Supplemental Fig. S2). We found that in these cells *TPCN2* and *LTO1* are expressed 0.6-fold and 0.7-fold lower than *CCND1* respectively; both have DHS at their promoters, and simulations show these interacting with several other nearby DHS. Despite *TPCN2* being in the adjacent TAD, its promoter interacts with a number of DHS between it and *MYEOV*; there is a CTCF site just upstream of *MYEOV* (arrow 'ii' in Fig. 2C), which seems to isolate *TPCN2* from further downstream interactions. The *LTO1* promoter interacts with a DHS within the *CCND1* gene body (which has chromatin state indicative of weak transcription, arrow 'iii'), and with a DHS between *MYEOV* and *CCND1* which has an enhancer chromatin state (arrow 'iv' in Fig. 2C). *MYEOV* shows on average 0.045-fold lower expression than the other three genes (Supplemental Fig. S2A), and is covered by enhancer-associated H3K4me1 and H3K27ac states. While there is no DHS at the *MYEOV* promoter to drive strong interactions, there is a general enrichment of interactions within a broad region downstream of the gene (arrow 'v' in Fig. 2C), driven by the CTCF site immediately upstream of the gene (arrow 'ii'). Interestingly, there is a DHS just downstream of *MYEOV* (arrow 'vi') which interacts strongly with DHS across the locus (including at several promoters, see bottom track in Fig. 2C); this, together with the chromatin states, suggests that this region may have enhancer function.

A super-enhancer insertion near *CCND1* drives changes in 3D structure within the TAD

We next used HiP-HoP to study *CCND1* in the MM cell line U266. These cells possess a genomic rearrangement where an enhancer region from *IGH* is inserted upstream of *CCND1*, within its TAD boundaries. A number of super-enhancers have

previously been described in the non-variable *IGH* region, including the long recognised Eα1, Eα2 and Eμ enhancers (Mills *et al.*, 1983; Chen and Birshtein, 1997; Mills *et al.*, 1997). More recently we proposed that a fourth element Eδ should be identified as distinct from Eμ, based on high quality chromatin state mapping (Mikulasova *et al.*, 2020b). The ∼100 kbp inserted region in U266 includes Eα1 [accurately mapped by Mikulasova *et al.* (2020b)]. U266 is a useful model because it allows study of the effect of relocating a single *IGH* enhancer within an existing TAD. Strikingly, it leads to significant changes in expression which are similar to those resulting from a reciprocal translocation.

In order to simulate the 3D genome structure U266 cells we generated an alternative genome build which includes the insert, denoting this hg19_u266 (compared to the unaltered hg19 genome used for GM12878 above). We obtained U266 DNA accessibility and chromatin states from the BLUEPRINT project (Stunnenberg *et al.*, 2016), and used CTCF binding site data for B-cells (The ENCODE Project Consortium, 2012). Figure 3A shows simulated Hi-C results for U266 along with the input data; the insert region is marked in green. For comparison, Fig. 3B shows the intact locus from the GM12878 simulations (no insert). Example configurations are shown in Fig. 3C.

The overall TAD structure is unchanged by the presence of the insert, however, interactions within the domains show significant differences between the two cell lines. Particularly, in U266 there are several strongly interacting regions (dark spots in the Hi-C map, some highlighted with green circles), including close to *CCND1* and within the *IGH* insert itself. Many of these are between DHS, between regions with heterochromatin states, or between regions with polycomb states. DHS often coincide with regulatory elements such as promoters and enhancers; particularly in U266 there is a cluster of DHS near and within *CCND1* which is not present in GM12878. Together these observations suggest that loop extruders and CTCF sites drive TAD formation, while bridging proteins drive interactions within TADs. Extrusion is unchanged by the presence of the insert (which does not contain any CTCF sites), while significant remodelling of chromatin in U266 compared to GM12878 leads to changes within the TAD.

Figure 3D shows simulated 4C with viewpoints positioned at gene promoters (top four rows). We found that there were new (compared to GM12878) strong interactions between the *CCND1* promoter and several DHS across the TAD, including three prominent peaks within the insert (arrow 'i' in Fig. 3D), as well as several peaks at DHS between *MYEOV* and the insert (arrows 'ii'; some of these DHS were not present in GM12878). Reciprocal interactions were observed as expected (Fig. 3D, bottom four rows). Many of the regions around the new DHS also gained enhancer or promoter chromatin states compared to GM12878. In U266, *CCND1* also interacts with nearby *LTO1* (arrow 'iii' in Fig. 3D), as well as showing weak interactions with the promoters of *MYEOV* and *TPCN2* (arrows 'iv' and 'v'). These changes result from the fact that *CCND1* gains DHS and active promoter and enhancer chromatin states (H3K27ac and H3K4me3) at the promoter and across its body (while losing H3K4me1 and H3K27me3 at the promoter). These features are consistent with H3K4me3 broad domains, which have previously been associated with super-enhancers (Thibodeau *et al.*, 2017; Dhar *et al.*, 2018). Their appearance has been implicated in cancer-specific super-enhancer hijacking at a number of oncogenes [including *CCND1*; Mikulasova *et al.* (2020b)].

Figure 3. A. Simulated Hi-C data is shown below a map of the genes within the rearranged *CCND1* locus as found in the U266 cell line (hg19_u266 genome; a region encompassing the *CCND1* and *FGF19*-TADs is shown). Green rings highlight additional interactions not present in GM12878 cells. The green block and dashed lines show the position of the inserted region. Data used as simulation input [obtained from BLUEPRINT and ENCODE] is shown as coloured blocks under the Hi-C map using the same scheme as in Fig. 2C. **B.** Simulated Hi-C data is shown for the intact *CCND1* locus for GM12878 cells (hg19 genome). The green dashed line indicates the insertion point for the U266 rearrangement. **C.** Example snapshots from simulations showing the *CCND1*-TAD in GM12878 (left) and U266 (right). In the latter the *IGH* insert containing Eα1 is shown in green; for comparison in the GM12878 snapshot the position of the insert is shown as a single green bead. Four further example snapshots are shown underneath. **D.** Simulated 4C plots are shown for U266 at various viewpoints (blue triangles). In the top four tracks, viewpoints are at promoters of *TPCN2*, *MYEOV*, *CCND1* and *LTO1*. The bottom four tracks show reciprocal viewpoints from interacting regions. The green block and dashed lines show the position of the insert, red blocks show DHS, and the purple block shows Eα1. Some features are labelled with numbered arrows as detailed in the text.

The promoter of *MYEOV* also gains a DHS in U266 compared to GM12878, and the chromatin state changes such that H3K27ac and H3K4me3 now covers the gene, extending several kbp downstream (another broad domain). It also shows a number of new interaction peaks, particularly in a region downstream where there is a cluster of new DHS with active enhancer and promoter chromatin state (arrows 'vi' in Fig. 3D). The *MYEOV* broad domain region as a whole interacts with the *CCND1* broad domain in ∼3% of configurations, compared to ∼2% in GM12878. Surprisingly, *MYEOV* shows more frequent interactions with the *CCND1* promoter that the *IGH* insert itself, despite the latter being closer genomically. This is likely because the DHS within the insert are alone and embedded in very short H3K27ac regions, whereas both *CCND1* and *MYEOV* have a cluster of DHS in a broad domain. Taken together these observations suggest that this new *MYEOV* broad domain is active as an enhancer (as noted above the *MYEOV* proximal region also displayed some enhancer-like features in the healthy cell line).

HiP-HoP generates a population of structures (representing a population of cells), and provides full three-dimensional details. We can also therefore measure, for example, separations between specific points or the overall 3D size of a given region. We find that the mean separation of *TPCN1* and *CCND1* tends to be larger in U266 where the insert is present. This is expected because not only is the genomic separation (along the chromosome) of the probes larger, but in the model more of the region has the open (H3K27ac associated) structure. The latter effect is highlighted by examining the 3D size of a region of fixed genomic length around *CCND1* in the two cell lines: the 3D size is larger in U266 cells (Supplemental Fig. S2D, and see also Fig. 5A below).

The reciprocal translocation t(11;14)(q13;q32) generates an oncogenic TAD fusion

The MCL cell line Z-138 harbours a translocation that dramatically changes the local environment of *CCND1*: it relocates to chromosome 14 where it becomes juxtaposed with the joining

A. t(14;11) rearrangement in Z-138 cells.

B. Domain pattern.

C. Simulated 4C interaction profiles

Figure 4. A. Simulated Hi-C for Z-138 cells which possess a t(11;14) translocation (hg19_z138 genome build). Positions of the *IGH* super-enhancers are shown as purple blocks above the map. Simulation input data (DHS, chromatin states and CTCF peaks) are shown under the map as in previous figures. **B.** Positions of TADs obtained from inspection of the simulated Hi-C. **C.** Simulated 4C from viewpoints positioned at DHS across the region (blue triangles). Some features are highlighted with numbered arrows as discussed in the text.

region of the *IGH* locus. While the t(11;14) rearrangement is common, the precise break-point has not previously been mapped in this cell line. Using paired-end read targeted DNA sequencing, we precisely mapped the chromosomal changes (Supplemental Fig. S3). With this information we could then perform simulations of the rearranged genome, which we denote hg19_z138.

Figure 4 shows results from simulations using Z-138 epigenomic data [again obtained from BLUEPRINT, Stunnenberg *et al.* (2016), and ENCODE, The ENCODE Project Consortium (2012)]. The simulated Hi-C map (Fig. 4A) predicts that a new

TAD will form, encompassing *CCND1*, *LTO1* and the *IGH* super-enhancers (Eα2, Eα1, Eμ, and Eδ). The right-hand boundary of the TAD is formed by a pair of divergent CTCF sites, and is conserved from the chromosome 11 boundary observed in GM12878 and other cell types. The left-hand boundary is formed by three inward pointing CTCF sites; a boundary is also seen at this position on chromosome 14 in some other cell types which do not have a translocation [including human umbilical vein endothelial cells, Rao *et al.* (2014)]. In summary, the simulations predict that these two boundaries are conserved after the translocation, and this leads to formation of a new *IGH-CCND1* TAD. This new TAD has a length of about 500 kbp, about 20% smaller than the ∼ 640 kbp *CCND1*-TAD observed in GM12878 cells (the 3D volume of the TAD in Z-138 is also correspondingly smaller). Interestingly a stronger subTAD (length ∼ 407 kbp, Fig. 4B) forms due to looping between a CTCF site just upstream of Eα2 and a CTCF within *CCND1*; this is the same CTCF site which forms the subTAD in GM12878 (length ∼ 390 kbp).

Figure 4C shows simulated 4C with viewpoints at *CCND1* and several positions within the translocated *IGH* super-enhancers. *CCND1* shows prominent interaction peaks in both Eδ and Eμ; these coincide with DHS within H3K27ac regions (arrows 'i' and 'ii' in the figure; the interaction is strongest where the DHS are clustered together). There are also smaller interaction peaks within Eα1 and Eα2, and at a number of other DHS (notably a cluster to the left of Eα2, arrow 'iii'). Similar to U266, there are a number of DHS at the *CCND1* promoter and within the gene body which are not present in the GM12878 cell line. In the simulations the interactions of the promoter are driven by binding (and molecular bridging) of active proteins; this bridging is likely further promoted by CTCF/cohesin driven loop extrusion. Viewpoints positioned at DHS within the enhancers show reciprocal interactions with *CCND1*.

Here we have simulated the break-point and rearrangement as mapped for the Z-138 cell line (about 60 kbp upstream of *CCND1*). It is also possible to simulate alternative rearrangements to better understand the effect of break-point variation between patients. Up to 50% of t(11;14) MCL break-points are within the so-called 'major translocation cluster' about 120 kbp upstream of *CCND1* (Jares and Campo, 2008), with the rest scattered throughout a broader 380 kbp region around this. Supplemental Figure S4 shows simulated 4C from two different examples where the chromosome 14 break-point is kept the same but we move the break-point on chromosome 11 to different positions between 11 and 100 kbp upstream of *CCND1*. We found that in general, interactions between the gene and the Eμ enhancer are weaker if their genomic separation is larger.

Chromatin remodelling and *CCND1* 3D structures differ in U266 and Z-138

The changes to chromatin states and DNA accessibility at *CCND1* differ subtly between the two cancer cell lines. This is visible in Figs. 3A and 4A, but is more evident if the data are mapped back to the hg19 reference genome (Supplemental Fig. S6 shows the chromatin states and DHS at three regions within the *CCND1* locus where there is significant remodelling compared to GM12878). In both U266 and Z-138 the *CCND1* gene body gains several DHS and a H3K4me3 broad domain appears. From 10 distinct DHS within *CCND1*, 5 are common to both cancer cell lines, 3 are specific to Z-138 and 2 are specific to U266. In other words, the different

Figure 5. A. Left: box plot showing the distribution of the radius of gyration of the *CCND1* gene body (a measure of its 3D size) for each of the three cell types (440 configurations for each case). Right: Typical snapshots of the gene and a 5 kbp flaking region on each side. The gene body is shown as a solid tube, with flanking regions shown as an outline. The arrow shows the transcription start site. The polymer is coloured according to the simulation input data: red indicates active protein binding sites (DHS), blue polycomb protein binding sites, and yellow are open chromatin (H3K27ac). **B.** Left: bar plot showing the change in expression level in U266 cells compared to GM12878 obtained from qPCR measurements (see Supplemental Methods and Supplemental Fig. S2 for further details). The bar height gives the $\log_2$ fold change based on an average over three replicates; the error bar shows the standard error in the mean. Right: similar plot for Z-138 cells compared to GM12878. No *MYEOV* expression was detected in these cells. **C.** Plot comparing fold change in expression (U266 *vs.* GM12878) obtained from qPCR and the prediction from simulations based on the frequency of interaction of the gene promoter and chromatin with enhancer state. We do not expect a linear relationship, but a good prediction is where the points sit on an arbitrary monotonically increasing curve (such a curve is shown as a guide to the eye; see Supplemental Methods for further details). Error bars again show standard error in the mean (from three replicates from the qPCR and 440 measurements from simulations). **D.** Map of the *CCND1* locus (hg19) showing the positions matching the fosmid probes used in FISH experiments. **E.** Example FISH images. The left image shows a whole cell with red, green and yellow probes covering different genes as indicated. Middle images show the yellow probe spots only (regions indicated by orange boxes in the whole-cell image). Right images show a 3D reconstruction of the gold probes as used to measure the volume (see Supplemental Methods for details). **F.** Box plot showing the distributions of the volume of the probe covering *CCND1* in the three cell lines (see text and Supplemental Methods for details). The number of probe sports measured in each case (n) is indicated. **G.** Box plot showing a similar measurement but extracted from $n = 440$ simulated configurations from each cell line (see Supplemental Methods for details of the calculation).

rearrangements lead to similar, but not identical, changes to the chromatin structure at *CCND1*. In U266 cells two other regions within the TAD also show substantial changes: a region covering and extending downstream of *MYEOV* and a region between *MYEOV* and *CCND1* (location of arrows 'ii' and 'vi' in Fig. 3D) both gain a H3K4me3 broad domain and several new DHS. We note that the pattern of chromatin states and accessibility at the *IGH* enhancers also varies between the U266 and Z-138 cell lines; this is discussed in more detail in Supplemental Methods (and see Supplemental Fig. S7).

The changes to the chromatin states at *CCND1* has a striking effect on the 3D structure of the gene: in Fig. 5A we plot the distribution of the simulated 3D size of the gene body (quantified by the radius of gyration, R_g, see Supplemental Methods). In GM12878 the gene is on average more compact than in the other cell lines. The snapshots in Fig. 5A show typical configurations: the differences in the size of the gene, the chromatin states, and the DHS pattern are clear. In GM12878 the gene has a crumpled structure, while in U266 it is more stretched out. Protein mediated loops can form between the DHS within the gene body which reduce its 3D size: variation in the number of such loops present at any one time leads to the large variation in R_g. The snapshot for the Z-138 case shows a configuration where a loop forms between a DHS within the gene body and a DHS in the upstream region. Since there are more DHS across the gene in Z-138 there are more possibilities for loops to form: smaller R_g values are therefore more likely and

there is less variation.

To see how this large change in 3D structure correlates with gene expression, we performed additional qPCR experiments comparing the expression of the four genes in the locus between the cell lines. Figure 5B shows their fold change in expression in U266 compared to GM12878 (see also Supplemental Fig. S2B). Notably, while all of the genes show an increase, that of *CCND1* is by far the largest, more than 500-fold. The next largest change is for *LTO1* which shows a ~ 9.8-fold increase. We considered whether these changes could be predicted from the simulated structures. In our simulations the active proteins represent complexes of transcription factors and polymerases, and in previous work using a simpler model we showed that the fraction of time a promoter is bound by an active protein during the simulation is correlated with gene expression (Brackley *et al.*, 2020). Crucially, this depends on the local 3D structure, since promoter-protein binding is typically due to protein clustering driven by chromatin looping. This measure does not, however, take into account enhancer 'strength' or compatibility, and can only have a value between 0 and 1; therefore we only expect a correlation and not a definitive mapping to expression. Nevertheless, the predictions are largely consistent with the qPCR data (see Supplemental Methods and Supplemental Fig. S2Ei for further details). We get an even better correlation between simulations and data if, rather than considering protein binding, we instead ask how often the promoter interacts with (forms loops to) chromatin which has

an enhancer associated state (Fig. 5C).

In Z-138 cells, *CCND1* expression shows an even larger 1,200-fold increase compared to GM12878 (qPCR data shown in Supplemental Fig. S2C); *LTO1* and *TPCN2* also show increased expression in Z-138 compared to GM12878 cells, but we did not detect *MYEOV* expression. Only *CCND1* and *LTO1* were present within the region simulated, and simulations correctly predicted the former would display the greater fold change compared to GM12878 cells (Supplemental Fig. S2Eii).

To verify that the change in 3D gene size is observed *in vivo*, we performed fluorescence *in situ* hybridisation (FISH) microscopy using BAC probes [see Methods and Supplemental Methods]. Figure 5D shows a typical image from the Z-138 cells. First we note that the positioning of the spots is consistent with the translocation. One of each of the red, green and gold spots are together in a group (on the non-translocated chromosome 11); the other gold (*CCND1*) probe is alone, while the other red and green probes (*TPNC2* and *MYEOV* respectively) are together. From the images we obtain the 3D volume of the probe covering *CCND1* (see Supplemental Methods), and we plot the distribution of volumes across multiple cells in Fig. 5E (we limit our analysis to cells where two probes can be unambiguously identified, see Supplemental Methods). A similar measure is obtained from the simulations [Fig. 5F]; we note that the probes cover a region larger than the *CCND1* gene body (42.8 kbp compared to the 13.4 kbp gene), and the predicted differences between the cell lines is not as striking as in Fig. 5A. The simulations correctly predict that the *CCND1* probe is on average larger in the cancer cell lines (and largest in Z-138 cells), however the predicted differences are smaller than measured from the FISH (particularly for Z-138). This suggests that while they reproduce the trends, the simulations do not fully capture the volumes; this is consistent with previous work where predictions for 3C-based data (interactions) were better than those for microscopy data (Buckle *et al.*, 2018).

The oncogene structure of *CCND1* is driven by chromatin remodelling

We have observed that in these cell lines the genome rearrangement is accompanied by an extensive chromatin remodelling. In particular, in both U266 and Z-138 a H3K4me3 broad domain containing several DHS appears over *CCND1*. A likely scenario is that after the rearrangement, the *IGH* super-enhancers recruit transcription factors, chromatin remodellers, and architectural proteins, to the region. The resulting changes then in turn disrupt the wider 3D structure, leading to dysregulation of *CCND1*. The HiP-HoP computational framework provides a unique opportunity to examine these two effects in isolation: we can rearrange the genome without otherwise changing the chromatin states, or we can use chromatin states from a cancer cell line without including any genomic rearrangement.

Figure 6A shows results from a simulation where chromatin state and DHS data for GM12878 cells were used as input, but the hg19_u266 genomic insert rearrangement was included (as depicted in the cartoon). In Fig. 6C we compare 4C results from GM12878 simulations with and without the insert; this shows that adding the insert has only a small effect on the interaction profile of the gene promoters. The profiles for *TPCN2* and *MYEOV* are unchanged. Also, despite the insert containing a DHS within a H3K27ac region, there is little in-

teraction with the nearby *CCND1* promoter (which has a polycomb chromatin state). On the other hand, there is some interaction between the insert and *LTO1*. Using the simulations to predict gene expression as above, these results suggest that without any subsequent chromatin remodelling, the presence of the insert alone would not lead to CCND1 up-regulation.

Figures 6B and D show similar results from simulations where U266 epigenomic input data is used with the intact hg19 genome (no insert, as shown in the cartoon). Here we see the effect of the chromatin remodelling in the absence of the insert. We find that in this case the *CCND1* promoter still shows interaction peaks across the *CCND1* gene body, at regions downstream (in and around *LTO1*), and to a lesser extent the H3K4me3 broad domain regions upstream. Beyond the loss of interactions with the insert itself, the main effect of removing the insert is an increase in interactions between *CCND1* and upstream DHS, consistent with their reduced genomic separation. The simulations also predict that very little change in gene expression level would result from removing the insert while keeping the U266 chromatin states. Together these results suggest that it is chromatin remodelling which drives the changes in the 3D structure of the locus in terms of promoter-enhancer interactions, and this in turn drives up-regulation of cyclin D1.

As well as simulating different rearranged genomes, it is also possible to directly edit the input data, *i.e.*, we can edit the epigenome *in silico*. For example, we found little change in *CCND1* interactions if we take chromatin states from GM12878 cells for chr11, and add an insert from chr14 with states from the U266 cells (Supplemental Fig. S8A-B). In a simulation of the U266 cells with the insert where we "switch off" protein binding at four DHS within the promoter and gene body of *CCND1*, there is a loss of *CCND1* interactions compared to the previous U266 simulations (Supplemental Figs. S8C-E).

Discussion

In this work we have adapted the HiP-HoP simulation model to study chromatin 3D structure and the effect of genome rearrangements in malignant and non-malignant B-cells. HiP-HoP simulations predict 3D structures from DNA accessibility, chromatin states, and CTCF binding data. We used these structures to simulate Hi-C and 4C data, and to make single cell-like measurements. Importantly, Hi-C data are *not* used as an input, so in this sense the model is truly predictive for 3D structures. Here, by "rearranging" the input data we generated predictions for the effect of genome rearrangements which are found in MCL and MM cell lines.

We first confirmed that the HiP-HoP model gives good predictions for the *CCND1* gene locus by performing simulations of the healthy B-cell derived GM12878 lymphoblastoid cell line. The TAD pattern observed in Hi-C (Rao *et al.*, 2014) is clearly reproduced by the simulations, with boundaries at CTCF sites. In these cells *CCND1* is only expressed at very low levels; the promoter region has a polycomb associated chromatin state, and our simulations predicted interactions between the promoter and other polycomb regions.

Further simulations predicted that the TAD structure around *CCND1* is preserved in the U266 MM cell line which possesses a rearrangement where a super-enhancer from the *IGH* locus is inserted upstream of *CCND1*. This is consistent with previously published low-resolution Hi-C data in that cell line (Wu *et al.*, 2017). In these cells the chromatin around *CCND1* is

Figure 6. A. Top: cartoon showing an *in silico* genome rearrangement where an *IGH* region is inserted into the GM12878 *CCND1* locus, and the GM12878 (healthy) chromatin states are retained (*i.e.*, the hg19_u266 reference genome is used). Bottom: Hi-C map generated from simulations with this *in silico* rearrangement. The input data are shown below the map. **B.** Top: cartoon showing an *in silico* genome rearrangement where the *IGH* insert is removed from the U266 *CCND1* locus, and the U266 chromatin states are retained (*i.e.*, the hg19 reference genome is used). Bottom: simulated Hi-C generated from simulations with this *in silico* rearrangement. **C.** Simulated 4C are shown for simulations using GM12878 chromatin states both with and without the insert. **D.** Simulated 4C are shown for simulations using U266 chromatin states both with and without the insert. **E.** Cartoon showing a model where first the genomic rearrangement brings the *IGH* super-enhancer into the *CCND1* TAD; this then leads to recruitment of chromatin remodellers *etc.* to the region and local chromatin states and DNA accessibility are altered. This could be described as an "epigenomic rearrangement".

remodelled: several new DHS are established, and a broad region gains an active promoter chromatin state (H3K4me3 and H3K27ac). The simulations predicted that the *CCND1* promoter interacts with these new DHS in the gene body, as well as with a number of DHS within the inserted *IGH* region and the neighbouring gene *LTO1* (interactions which are not present in GM12878 cells).

We also performed simulations of the MCL cell line Z-138, where *CCND1* translocates with chromosome 14 becoming juxtaposed with the *IGH* Eμ enhancer. In this cell line a TAD boundary formed by a cluster of CTCF binding sites downstream of *CCND1* was preserved after the translocation, as was

a boundary to the left of the *IGH* super-enhancers (presumably in healthy cells this isolates the super-enhancers from other nearby genes). In other words, the simulations predict a fusion of the *CCND1* and *IGH* regions into a new TAD – an oncogenic TAD fusion (Valton and Dekker, 2016). In this new arrangement the promoters of *CCND1* and *LTO1* readily interact with each other and with the proximate Eμ and Eδ super-enhancers, while weaker interactions were observed with the more distant Eα1 and Eα2.

In vivo, genomic rearrangements are accompanied by chromatin remodelling; within our simulation scheme it was also possible to examine the effect of a rearrangement in the ab-

sence of remodelling. Simulations of GM12878 cells with the Eα1 enhancer DNA inserted *in silico* upstream of *CCND1* (with GM12878 chromatin states otherwise unchanged) showed only very minor changes in terms of 3D structure compared with the unaltered genome. We also performed a simulation using U266 chromatin states, but without the insert (*i.e.*, the epigenomic rearrangement is included, but not the genomic rearrangement, an experiment that is only possible to do *in silico*). This showed little change compared to the U266 case with the insert. Together this suggests that it is the local chromatin remodelling, rather than the proximity of *CCND1* to the *IGH* enhancers *per se*, which drives gene deregulation. Or in other words, a genomic translocation leads to an epigenomic translocation, which drives cyclin D1 overexpression. The remodelling includes the appearance of an H3K4me3 broad domain over *CCND1*; the results here support our previous work suggesting that such broad domains are associated with super-enhancer hijacking (Mikulasova *et al.*, 2020b). In U266 two other nearby regions become H3K4me3 broad domains, one covering the gene *MYEOV* (this region also appears to have some enhnacer-like properties in the healthy cell line). Importantly, the chromosomal locations which are predicted to strongly interact with *CCND1* could be used for targeting in future experiments which aim to uncover the mechanisms through which broad domains are generated.

In principle, our method can be applied to simulate any genome rearrangement for which input data is available. One current limitation is that the input data comes from both alleles of the locus, so for example in the Z-138 case the chromatin states are based on a combination of histone modifications from the translocated and non-translocated *CCND1* loci. We note that this is a limitation of the input data rather than the simulation scheme; if the full sequence of a given cell line or patient sample were available, ChIP-seq reads could be aligned in an allele specific way. It may also be possible to use the simulations to explore hypotheses about differences between alleles, *e.g.*, if one were to assume one copy of the locus has active chromatin states and the other inactive states, how does the 3D structure differ? A simplifying assumption in the model is that all DHS are binding sites for a general active factor, and it would be interesting to incorporate more detail in the future, *e.g.*, if transcription factors specific to a particular oncogene are known (Brackley *et al.*, 2016a). Other possible improvements to the model include adding further "compaction levels" in the heteromorphic polymer (which might improve microscopy predictions), or adding protein-protein interactions promoting larger liquid-like phase separated protein droplet formation [augmenting the small protein clusters which form via bridging-induced attraction in the present model; Brackley *et al.* (2013)].

In summary, our work strongly suggests that genome rearrangements drive a subsequent epigenomic rearrangement, which in turn leads to deregulation and proto-oncogene activation. We have demonstrated that polymer physics-based modelling can be useful for understanding the structural consequences of genome rearrangements, and can help to focus future experimental work. It would be interesting to see if in the future, such simulations could also help us understand the mechanisms behind the epigenomic rearrangement. This would clearly be important for the development of any therapies which seek to interfere with that process.

Methods

HiP-HoP simaultions

We used the HiP-HoP model as detailed in Buckle *et al.* (2018); full details are given in Supplemental Methods. In brief, chromatin is represented as a chain of beads (each representing a 1 kbp region), and we evolve the configuration of this chain using a molecular dynamics scheme and the LAMMPS simulation software (Plimpton, 1995). To improve efficiency, in each simulation we include 40 Mbp of chromatin (40,000 beads) which includes 11 copies of the region of interest; for each region we perform two such simulations to generate 4400 individual configurations each of which can be said to represent a single cell. The chromatin concentration is roughly matched to that of a typical nucleus. From the configurations we generate simulated Hi-C and 4C and single cell-like measurements. As detailed in the text, three different mechanisms are included to drive the chain into specific structures: diffusing bridge forming proteins, loop extrusion, and a heteromorphic polymer structure. DNase hypersensitive sites (DHS) are used to identify binding sites for an active protein (we use the simplifying assumption that all DHS are the same and bind this protein, which represents a general complex of polymerase and transcription factors). We use chromatin state data to identify binding sites for two species of repressive protein (*e.g.*, representing HP1 and polycomb repressive complexes), and to identify regions which have an open chromatin structure. ChIP-seq data for CTCF is used to identify direction-dependent anchor sites for loop extrusion. Full details of the input data treatment are given in the Supplemental Methods.

Quantification of gene expression

For gene expression measurements, total RNA was extracted using RNeasy kit (QIAGEN, D usseldorf, Germany) from between 5×10^6 and 10×10^6 Z-138, U266 and GM12878 cells. For qPCR, cDNA was synthesised from 1 μg of total RNA using random primers (Promega, Wisconsin, USA) then treated with M-MLV Reverse Transcriptase (Promega, Wisconsin, USA). 2 μg cDNA reaction were amplified with SYBR green master mix (Invitrogen, Massachusetts, USA), 10 μM forward and reverse primers for *LTO1*, *TPCN2*, *CCND1* and *MYEOV* (Sigma-Aldrich, Missouri, USA) with 40 cycles of PCR (95°C for 15 s, 60°C for 60 s) after initial denaturation (95°C for 10 min). Cycle threshold values were normalized by comparison with *GAPDH* expression. See Supplemental Methods for further details (oligonucleotide sequences are shown in Supplemental Table S2).

Genome rearrangement breakpoint mapping

Breakpoint mapping in U266 and Z-138 cells was performed as described previously (Mikulasova *et al.*, 2020b,a); further details are given in Supplemental Methods.

FISH analysis

For FISH microscopy, briefly, fosmid clones found to cover *TPCN2* (G248P87917D11/WI2-1721H21), *MYEOV* (G248P87014D2/WI2-2222G4) and *CCND1* (G248P86668E2/WI2-2191I3) loci were obtained from BACPAC resources (California, USA). DNA was fluorescently labelled by nick translation. Probes and cell line DNA were denatured at 75°C for five minutes followed by hybridisation at 37°C overnight. Further details are given in Supplemental Methods.

Data access

Existing publicly available data used in this study is detailed in Supplemental Table S3 and Supplemental Methods. For new data generated in this study, sequencing data are available via the NCBI SRA (accession number pending), while qPCR, FISH and simulation data are available via the Edinburgh DataShare repository (https://datashare.ed.ac.uk accession number pending). Further details are given in Supplemental Table S4.

Competing interest statement

The authors declare no competing interests.

Acknowledgements

Work in the DR lab is supported by a Wellcome Trust Seed Award in Science (206103/Z/17/Z). BMJ acknowledges funding from FEDER / Ministry of Science and Innovation - Spanish State Research Agency under the project RTI2018-094788-A-I00, and La Caixa Banking Foundation under the r project LCF/BQ/PI19/11690001. Work in the LJR lab was supported by CCLG Little Princess Trust (NK) and an MRC DiMeN DTP studentship (DK). CAB acknowledges support from European Research Council (ERC CoG 648050 THREEDCELLPHYSICS).

Author contributions: AM and BAW performed the Z-138 break-point mapping and sequencing; DK performed bioinformatics analysis and cell line qPCR experiments. NK performed cell line qPCR experiments and FISH probe generation. LJR performed FISH probe generation, hybridisation and post capture analysis of images. RB-P performed confocal microscopy and post-capture deconvolution and image analysis. CAB designed and performed the simulations. DR, LJR and CAB conceived and designed the research, interpreted the results, and wrote the paper. All authors contributed to writing the paper.

References

Akdemir, KC, Le, VT, Chandran, S, Li, Y, Verhaak, RG, Beroukhim, R, Campbell, PJ, Chin, L, Dixon, JR, Futreal, PA, *et al*. 2020. Disruption of chromatin folding domains by somatic genomic rearrangements in human cancer. *Nature Genetics* **52**: 294–305. doi:10.1038/s41588-019-0564-y

Bianco, S, Lupiáñez, DG, Chiariello, AM, Annunziatella, C, Kraft, K, Schöpflin, R, Wittler, L, Andrey, G, Vingron, M, Pombo, A, *et al*. 2018. Polymer physics predicts the effects of structural variants on chromatin architecture. *Nature Genetics* **50**: 662–667. doi:10.1038/s41588-018-0098-8

Brackey, CA, Marenduzzo, D, Gilbert, N. 2020. Mechanistic modeling of chromatin folding to understand function. *Nature Methods* **17**: 767–775. doi:10.1038/s41592-020-0852-6

Brackley, CA. 2020. Polymer compaction and bridging-induced clustering of protein-inspired patchy particles. *J Phys Condens Matter* **32**: 314002. doi:10.1088/1361-648X/ab7f6c

Brackley, CA, Brown, JM, Waithe, D, Babbs, C, Davies, J, Hughes, JR, Buckle, VJ, Marenduzzo, D. 2016a. Predicting the three-dimensional folding of cis-regulatory regions in mammalian genomes using bioinformatic data and polymer models. *Genome Biology* **17**: 59. doi:10.1186/s13059-016-0909-0

Brackley, CA, Gilbert, N, Michieletto, D, Papantonis, A, Pereira, MCF, Cook, PR, Marenduzzo, D. 2020. Complex small-world regulatory networks emerge from the 3d organisation of the human genome. *bioRxiv* 2020.05.12.091041. doi:10.1101/2020.05.12.091041

Brackley, CA, Johnson, J, Kelly, S, Cook, PR, Marenduzzo, D. 2016b. Simulated binding of transcription factors to active and inactive regions folds human chromosomes into loops, rosettes and topological domains. *Nucleic Acids Res* **44**: 3503–3512. doi:10.1093/nar/gkw135

Brackley, CA, Taylor, S, Papantonis, A, Cook, PR, Marenduzzo, D. 2013. Nonspecific bridging-induced attraction drives clustering of DNA-binding proteins and genome organization. *Proc Natl Acad Sci USA* **110**: E3605–E3611. doi:10.1073/pnas.1302950110

Buckle, A, Brackley, CA, Boyle, S, Marenduzzo, D, Gilbert, N. 2018. Polymer simulations of heteromorphic chromatin predict the 3-d folding of complex genomic loci. *Molecular Cell* **72**: 786–797. doi:10.1016/j.molcel.2018.09.016

Carrillo-de Santa-Pau, E, Juan, D, Pancaldi, V, Were, F, Martin-Subero, I, Rico, D, Valencia, A, on behalf of The BLUEPRINT Consortium. 2017. Automatic identification of informative regions with epigenomic changes associated to hematopoiesis. *Nucleic Acids Research* **45**: 9244–9259. doi:10.1093/nar/gkx618

Chen, C, Birshtein, BK. 1997. Virtually identical enhancers containing a segment of homology to murine 3'igh-e(hs1,2) lie downstream of human ig c alpha 1 and c alpha 2 genes. *Journal of Immunology* **159**: 1310–1318

Davidson, IF, Bauer, B, Goetz, D, Tang, W, Wutz, G, Peters, JM. 2019. Dna loop extrusion by human cohesin. *Science* **366**: 1338–1345. doi:10.1126/science.aaz3418

Dhar, SS, Zhao, D, Lin, T, Gu, B, Pal, K, Wu, SJ, Alam, H, Lv, J, Yun, K, Gopalakrishnan, V, *et al*. 2018. Mll4 is required to maintain broad h3k4me3 peaks and super-enhancers at tumor suppressor genes. *Molecular Cell* **70**: 825–841.e6. doi:10.1016/j.molcel.2018.04.028

Díaz, N, Kruse, K, Erdmann, T, Staiger, AM, Ott, G, Lenz, G, Vaquerizas, JM. 2018. Chromatin conformation analysis of primary patient tissue using a low input hi-c method. *Nature Communications* **9**: 4938. doi:10.1038/s41467-018-06961-0

Dixon, JR, Selvaraj, S, Yue, F, Kim, A, Li, Y, Shen, Y, Hu, M, Liu, JS, Ren, B. 2012. Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**: 376–380. doi:10.1038/nature11082

Engreitz, JM, Agarwala, V, Mirny, LA. 2012. Three-dimensional genome architecture influences partner selection for chromosomal translocations in human disease. *PLOS ONE* **7**: 1–9. doi:10.1371/journal.pone.0044196

Ernst, J, Kheradpour, P, Mikkelsen, TS, Shoresh, N, Ward, LD, Epstein, CB, Zhang, X, Wang, L, Issner, R, Coyne, M, *et al*. 2011. Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature* **473**: 43–49. doi:10.1038/nature09906

Fudenberg, G, Imakaev, M, Lu, C, Goloborodko, A, Abdennur, N, Mirny, L. 2016. Formation of chromosomal domains by loop extrusion. *Cell Reports* **15**: 2038–2049. doi:10.1016/j.celrep.2016.04.085

Gong, Y, Lazaris, C, Sakellaropoulos, T, Lozano, A, Kambadur, P, Ntziachristos, P, Aifantis, I, Tsirigos, A. 2018. Stratification of tad boundaries reveals preferential insulation of super-enhancers by strong boundaries. *Nature Communications* **9**: 542. doi:10.1038/s41467-018-03017-1

Hnisz, D, Abraham, BJ, Lee, TI, Lau, A, Saint-André, V, Sigova, AA, Hoke, HA, Young, RA. 2013. Super-enhancers in the control of cell identity and disease. *Cell* **155**: 934–947. doi:doi.org/10.1016/j.cell.2013.09.053

Jares, P, Campo, E. 2008. Advances in the understanding of mantle cell lymphoma. *British journal of haematology* **142**: 149–65. doi:10.1111/j.1365-2141.2008.07124.x

Jung, D, Giallourakis, C, Mostoslavsky, R, Alt, FW. 2006. Mechanism and control of v(d)j recombination at the immunoglobulin heavy chain locus. *Annual Review of Immunology* **24**: 541–570. doi:10.1146/annurev.immunol.23.021704.115830

Kontomanolis, EN, Koutras, A, Syllaios, A, Schizas, D, Mastoraki, A, Garmpis, N, Diakosavvas, M, Angelou, K, Tsatsaris, G, Pagkalos, A, *et al*. 2020. Role of oncogenes and tumor-suppressor genes in carcinogenesis: A review. *Anticancer Research* **40**: 6009–6015. doi:10.21873/anticanres.14622

Krietenstein, N, Abraham, S, Venev, SV, Abdennur, N, Gibcus, J, Hsieh, THS, Parsi, KM, Yang, L, Maehr, R, Mirny, LA, *et al*. 2020. Ultrastructural details of mammalian chromosome architecture. *Molecular Cell* **78**: 554–565.e7. doi:10.1016/j.molcel.2020.03.003

Lu, L, Liu, X, Huang, WK, Giusti-Rodríguez, P, Cui, J, Zhang, S, Xu, W, Wen, Z, Ma, S, Rosen, JD, *et al*. 2020. Robust hi-c maps of enhancer-promoter interactions reveal the function of non-coding genome in neural development and diseases. *Molecular Cell* **79**: 521 – 534.e15. doi:10.1016/j.molcel.2020.06.007

Mikulasova, A, Ashby, C, Tytarenko, RG, Qu, P, Rosenthal, A, Dent, JA, Ryan, KR, Bauer, MA, Wardell, CP, Hoering, A, *et al*. 2020a. Microhomology-mediated end joining drives complex rearrangements and overexpression of myc and pvt1 in multiple myeloma. *Haematologica* **105**: 1055–1066. doi:10.3324/haematol.2019.217927

Mikulasova, A, Trevisan-Herraz, M, Fung, K, Ashby, C, Karataraki, N, Morgan, GJ, Corcoran, A, Hambleton, S, Walker, BA, Rico, D, *et al*. 2020b. Epigenomic translocation of h3k4me3 broad domains following super-enhancer hijacking. *bioRxiv* 2020.02.12.938563. doi:10.1101/2020.02.12.938563

Mills, FC, Fisher, LM, Kuroda, R, Ford, AM, Gould, HJ. 1983. Dnase i hypersensitive sites in the chromatin of human mu immunoglobulin heavy-chain genes. *Nature* **306**: 809–12. doi:10.1038/306809a0

Mills, FC, Harindranath, N, Mitchell, M, Max, EE. 1997. Enhancer complexes located downstream of both human immunoglobulin calpha genes. *The Journal of Experimental Medicine* **186**: 845–58. doi:10.1084/jem.186.6.845

Norton, HK, Phillips-Cremins, JE. 2017. Crossed wires: 3d genome misfolding in human disease. *J Cell Biol* **216**: 3441–3452. doi:10.1083/jcb.201611001

Park, S, Kim, GW, Kwon, SH, Lee, JS. 2020. Broad domains of histone h3 lysine 4 trimethylation in transcriptional regulation and disease. *The FEBS Journal* **287**: 2891–2902. doi:10.1111/febs.15219

Plimpton, S. 1995. Fast Parallel Algorithms for Short-Range Molecular Dynamics. *J Comp Phys* **117**: 1–19. doi:10.1006/jcph.1995.1039

Rao, SS, Huntley, MH, Durand, NC, Stamenova, EK, Bochkov, ID, Robinson, JT, Sanborn, AL, Machol, I, Omer, AD, Lander, ES, *et al*. 2014. A 3d map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* **159**: 1665–1680. doi:10.1016/j.cell.2014.11.021

Risca, VI, Denny, SK, Straight, AF, Greenleaf, WJ. 2017. Variable chromatin structure revealed by in situ spatially correlated dna cleavage mapping. *Nature* **541**: 237–241. doi:10.1038/nature20781

Roukos, V, Misteli, T. 2014. The biogenesis of chromosome translocations. *Nature Cell Biology* **16**: 293–300. doi:10.1038/ncb2941

Sanborn, AL, Rao, SSP, Huang, SC, Durand, NC, Huntley, MH, Jewett, AI, Bochkov, ID, Chinnappan, D, Cutkosky, A, Lia, J, *et al*. 2015. Chromatin extrusion explains key features of loop and domain formation in wild-type and engineered genomes. *Proc Natl Acad Sci USA* **112**: E6456–E6465. doi:10.1073/pnas.1518552112

Serra, F, Di Stefano, M, Spill, YG, Cuartero, Y, Goodstadt, M, Baù, D, Marti-Renom, MA. 2015. Restraint-based three-dimensional modeling of genomes and genomic domains. *FEBS Letters* **589**: 2987–2995. doi:10.1016/j.febslet.2015.05.012

Stunnenberg, HG, Abrignani, S, Adams, D, de Almeida, M, Altucci, L, Amin, V, Amit, I, Antonarakis, SE, Aparicio, S, Arima, T, *et al*. 2016. The international human epigenome consortium: A blueprint for scientific collaboration and discovery. *Cell* **167**: 1145–1149. doi: 10.1016/j.cell.2016.11.007

Thandapani, P. 2019. Super-enhancers in cancer. *Pharmacology & Therapeutics* **199**: 129–138. doi: 10.1016/j.pharmthera.2019.02.014

The ENCODE Project Consortium. 2012. An integrated encyclopedia of dna elements in the human genome. *Nature* **489**: 57–74. doi: 10.1038/nature11247

Thibodeau, A, Márquez, EJ, Shin, DG, Vera-Licona, P, Ucar, D. 2017. Chromatin interaction networks revealed unique connectivity patterns of broad h3k4me3 domains and super enhancers in 3d chromatin. *Scientific Reports* **7**: 14466. doi:10.1038/s41598-017-14389-7

Valton, AL, Dekker, J. 2016. Tad disruption as oncogenic driver. *Current Opinion in Genetics & Development* **36**: 34–40. doi: https://doi.org/10.1016/j.gde.2016.03.008

Watson, CT, Breden, F. 2012. The immunoglobulin heavy chain locus: genetic variation, missing data, and implications for human disease. *Genes & Immunity* **13**: 363–373. doi:10.1038/gene.2012.12

Wu, P, Li, T, Li, R, Jia, L, Zhu, P, Liu, Y, Chen, Q, Tang, D, Yu, Y, Li, C. 2017. 3d genome of multiple myeloma reveals spatial genome disorganization associated with copy number variations. *Nature Communications* **8**: 1937. doi:10.1038/s41467-017-01793-w

Zhao, Z, Tavoosidana, G, Sjölinder, M, Göndör, A, Mariano, P, Wang, S, Kanduri, C, Lezcano, M, SinghSandhu, K, Singh, U, *et al*. 2006. Circular chromosome conformation capture (4c) uncovers extensive networks of epigenetically regulated intra- and interchromosomal interactions. *Nature Genetics* **38**: 1341–1347. doi:10.1038/ng1891