Crosstalk in oxygen homeostasis networks: SKN-1/NRF inhibits the HIF-1 hypoxia-inducible factor in *Caenorhabditis elegans*

Dingxia Feng*¶, Zhiwei Zhai¶, Zhiyong Shao, Yi Zhang, and Jo Anne Powell-Coffman*

Department of Genetics, Development and Cell Biology, Iowa State University, Ames, Iowa,

United States of America

*corresponding author:

E-mail: japc@iastate.edu (JP)
E-mail: fengdi@indiana.edu (DF)

¶co-first authors contributed equally

Abstract

During development, homeostasis, and disease, organisms must balance responses that allow adaptation to low oxygen (hypoxia) with those that protect cells from oxidative stress. The evolutionarily conserved hypoxia-inducible factors are central to these processes, as they orchestrate transcriptional responses to oxygen deprivation. Here, we employ genetic strategies in *C. elegans* to identify stress-responsive genes and pathways that modulate the HIF-1 hypoxia-inducible factor and facilitate oxygen homeostasis. Through a genome-wide RNAi screen, we show that RNAi-mediated mitochondrial or proteasomal dysfunction increases the expression of hypoxia-responsive reporter *Pnhr-57:GFP* in *C. elegans*. Interestingly, only a subset of these effects requires *hif-1*. Of particular importance, we found that *skn-1* RNAi increases the expression of hypoxia-responsive reporter *Pnhr-57:GFP* and elevates HIF-1 protein levels. The SKN-1/NRF transcription factor has been shown to promote oxidative stress resistance. We present evidence that the crosstalk between HIF-1 and SKN-1 is mediated by EGL-9, the prolyl hydroxylase that targets HIF-1 for oxygen-dependent degradation. Treatment that induces SKN-1, such as heat, increases expression of a *Pegl-9:GFP* reporter, and this effect requires *skn-1* function and a putative SKN-1 binding site in *egl-9* regulatory sequences. Collectively, these data support a model in which SKN-1 promotes *egl-9* transcription, thereby inhibiting HIF-1. We propose that this interaction enables animals to adapt quickly to changes in cellular oxygenation and to better survive accompanying oxidative stress.

Introduction

Oxygen homeostasis has profound effects on health and fitness. Oxygen serves as the terminal electron acceptor in the oxidative phosphorylation processes that generate energy for life. When

oxygen levels are low (hypoxia), cells and tissues must adapt quickly by increasing oxygen

delivery, adjusting the levels of key metabolic enzymes, and limiting the accumulation of

misfolded proteins. While oxygen is essential, it is also highly reactive. The reactive oxygen

species (ROS) generated by cellular metabolism and signaling processes can damage

macromolecules, and excess ROS are thought to contribute to cellular aging and deterioration [1-

3]. One of the central challenges of aerobic life is to coordinate the biological networks that

control disparate aspects of oxygen homeostasis.

This balance between surviving hypoxic stress and mitigating the potential damage caused by

reactive oxygen species is especially important in cardiovascular development and disease.

When ischemia blocks circulation to a mammalian tissue, oxygen levels drop, and cells induce

hypoxia-inducible transcription factors (HIFs). Upon reperfusion and reoxygenation of the tissue,

mammalian cells respond by rapidly degrading HIF and inducing the NRF2 transcription factor

[4, 5]. However, intermittent hypoxia has been shown to induce both HIF-1α and NRF2 [6, 7].

NRF2 activates phase II detoxification genes to mitigate the effects of oxidative insults [8, 9].

Although mammalian HIF and NRF2 share some common target genes such as aldehyde

dehydrogenase 1A1 or heme oxygenase-1 HO-1, the genes induced by re-oxygenation are

largely distinct from those that respond to oxygen deprivation [5, 10, 11]. Crosstalk between

these two pathways is complex and context specific in mammals [12], as these important

transcription factors facilitate the rapid changes in gene expression needed to limit reperfusion

injury and regulate oxygen-dependent developmental processes.

C. elegans has been proven to be an excellent model system for studying the regulatory networks

that govern oxygen homeostasis. The *C. elegans* genome encodes a single hypoxia-inducible

factor alpha subunit (HIF-1), and the *hif-1* gene has been shown to have important roles in stress

response and in aging [13-17]. HIF protein levels and HIF activity are tightly regulated. When

oxygen is abundant, the HIF alpha subunit is hydroxylated by the PHD/EGL-9 enzymes. Once

modified, HIFα protein interacts with the Von Hippel-Lindau tumor suppressor (VHL) and is

targeted for ubiquitination and proteasomal degradation [18-23]. Thus, in hypoxic conditions,

HIF-1 protein is stable, and the transcription factor complex can activate the expression of a

battery of genes that enable adaptation to low oxygen. This pathway for oxygen-dependent

degradation of HIF protein is evolutionarily conserved. The *C. elegans hif-1, aha-1, egl-9,* and

vhl-1 genes are orthologous to mammalian *HIFα*, *HIFβ*, *PHD*, and *VHL*, respectively [24-26].

The targets of *C. elegans* HIF-1 include *egl-9* and *rhy-1*, genes that inhibit HIF-1 expression and

activity [27-30]. In wild-type animals, these negative feedback loops keep HIF-1 activity in

check and limit the potentially adverse effects of HIF-1 over-activation.

The *C. elegans skn-1* gene is homologous to mammalian *NRF1/2/3* [31]. SKN-1 regulates the

expression of a battery of genes with cytoprotective functions, including phase II detoxification

genes [32, 33]. SKN-1 is activated by a range of stresses or toxicants that cause oxidative stress,

and SKN-1 promotes resistance to these insults [32, 34-37].

Here, we investigate the cellular processes and transcriptional networks that regulate HIF-1

function. We describe an unbiased genetic screen to identify genes that inhibit *C. elegans* HIF-1.

We discover that SKN-1/NRF represses HIF-1 protein levels. Hence, SKN-1-mediated repression

of HIF-1 may provide a mechanism by which cells can rapidly respond to specific environmental

stresses and optimize gene expression to achieve oxygen homeostasis. We investigate the

hypothesis that this cross talk is mediated by EGL-9, the oxygen-sensing prolyl hydroxylase that

modulates HIF-1 stability and activity.

Results

To identify genes and cellular processes that attenuated HIF-1-mediated gene expression, we conducted a genome-wide RNAi screen. This experimental strategy relied on the *Pnhr-57:GFP* reporter gene, which had been shown to be responsive to HIF-1 and hypoxia [28, 29]. Through chromatin immunoprecipitation experiments, we confirmed that *nhr-57* was a direct target of HIF-1 (S1 Fig). We screened a bacterial RNAi library representing ~80% of *C. elegans* genes [38], and we identified 179 genes for which RNAi increased *Pnhr-57:GFP* expression, as assayed by inspection under a fluorescent stereomicroscope (screen design illustrated in Fig 1A). These genes are listed in S1 Table and their functions are summarized in Fig 1B. Remarkably, 89 genes were predicted to have mitochondrial or metabolic functions, and the majority of these were electron transport chain components (42 genes) or subunits of mitochondrial ribosomes (24 genes). The second largest category of the genes was protein folding or turnover (33 genes), and the majority of these were proteasomal components. The remaining major categories include vesicular transport, transporters and channels, signaling and cytoskeleton, transcription and DNA or RNA processing.

Fig 1. Genome-wide RNAi screen to identify negative regulators of HIF-1-mediated gene expression. (A) Illustration of screen design. The screen identified RNAi treatments that increased the expression of *Pnhr-57:GFP*, a reporter that has been shown to be induced by hypoxia and positively regulated by HIF-1. (B) Functional summary of the 179 genes identified from the RNAi screen.

Recognizing that most eukaryotic genes are coordinately regulated by multiple transcription

factors, we did a secondary screen to identify those RNAi treatments that had a clear *hif-1-*

dependent effect on *Pnhr-57:GFP* expression. To do this, we compared the *Pnhr-57:GFP*

induction of these 179 RNAi treatments in wild-type animals and *hif-1*-deficient animals. Most

of the 179 RNAi treatments increased the expression of *Pnhr-57:GFP* independent of *hif-1*.

However, the *Pnhr-57:GFP* induction by 13 RNAi treatments showed a strong *hif-1*-dependent

effect (S2 Table). Among these 13 genes, as expected, RNAi for *egl-9, rhy-1*, and *vhl-1*,

previously characterized negative regulators of *C. elegans* HIF-1 [26, 29], increased *Pnhr-*

57:GFP expression in wild-type animals, but not in *hif-1* mutants. These results validated the

efficacy of our screen approach, and gave us the confidence to continue investigating the

potentially new negative regulators of HIF-1 among these 13 genes.

skn-1 attenuates HIF-1 protein levels and HIF-1 function

We were especially intrigued by the finding that *skn-1* RNAi increased the expression of HIF-1-

responsive reporter. Transcription factor SKN-1 has been shown to have critical roles in enabling

C. elegans to respond to oxidative stress [32-34, 39]. Our finding suggested a potential crosstalk

between hypoxia response and oxidative stress response. To quantify the effect of *skn-1* RNAi

on this HIF-1-responsive reporter, we examined *Pnhr-57:*GFP levels using protein blots. In the

normal room air culture conditions, *Pnhr-57:*GFP was 40% higher in *skn-1* RNAi compared to

control RNAi (Fig 2A) (**p < 0.01, from three independent experiments). To gain insight to the

effects of this interaction in hypoxic conditions, we moved the animals to 0.5% oxygen. After 4

hours of hypoxia treatment, *Pnhr-57:*GFP was 50% higher in *skn-1* RNAi compared to control

RNAi (Fig 2A) (**p < 0.01, from three independent experiments). Thus, *skn-1* RNAi increased

*Pnhr-57:*GFP levels under normoxic and hypoxic conditions.

We next asked whether *skn-1* RNAi increased HIF-1 protein levels. We tested two *skn-1* RNAi

constructs, and each resulted in an increase of HIF-1 protein levels by 2 to 3-fold as shown in Fig

2B (**$p < 0.01$, from three independent experiments). In sum, these results showed that *skn-1*

RNAi increased HIF-1 protein level and HIF-1 reporter expression.

Fig 2. Identification of SKN-1 as a regulator of HIF-1. (A) *skn-1* RNAi increased expression

of the *Pnhr-57:GFP* reporter. Reporter gene expression was quantitated in L4-stage animals in

normal culture conditions or after 2 or 4 hours of hypoxia treatment (0.5% oxygen). The protein

levels were calculated from three independent experiments and normalized to 0 hour hypoxia

control RNAi. A representative western blot is shown. In each lane, lysates from 80 L4-satge

worms were loaded. Asterisks indicate significant differences between control RNAi and *skn-*

1 RNAi at any given time point. *: $p < 0.05$; **: $p < 0.01$. (B) *skn-1* RNAi increased HIF-1

protein levels. These animals expressed an epitope-tagged HIF-1 protein [40]. The protein levels

were calculated from three independent experiments and normalized to control RNAi. Two

different RNAi clones were assayed, and they are designated as *skn-1(1)* and *skn-1(2)* here. A

representative western blot is shown. In each lane, lysates from 100 L4-satge worms were

loaded. Asterisks indicate significant differences between control RNAi and *skn-1* RNAi at any

given time point. **: $p < 0.01$.

Differential requirements for *skn-1* and *hif-1*

The finding that SKN-1 repressed HIF-1 protein levels suggested that there might be conditions

in which it would be beneficial for the animal to express one of these two stress-responsive

transcription factors, but not the other. To address this, we examined the relative requirements

for *skn-1* and *hif-1* more closely.

In previous studies, we and others had shown that *hif-1* was required for survival in moderate

hypoxia [25, 41]. As validated in the experiments described in Table 1, loss of *hif-1* impaired

animal development and survival in 0.5% oxygen: after 24 hours of hypoxia treatment, only

75.8% of *hif-1*-deficient eggs hatched, and only 25.6% developed to adulthood within 72 hours.

In contrast, *skn-1* RNAi had no effect on *C. elegans* development and survival in hypoxic

conditions: after 24 hours of hypoxia treatment, 99.4% of *skn-1* RNAi treated eggs hatched and

completed normal development to adulthood within 72 hours.

Table 1. Relative requirements for *skn-1* and *hif-1*: survival in 0.5% oxygen.
165

Genotype	% hatched ± SEM	% survive to adult ± SEM	n [a]
N2 (wild type)	99.1 ± 0.25	99.1± 0.3	672
hif-1(ia04)	75.8 ± 2.5	25.6 ± 0.9	616
N2;control(RNAi)	99.6 ± 0.19	99.6 ± 0.2	558
N2;*skn-1(RNAi)*	99.4 ± 0.42	99.4 ± 0.4	640

166 [a] n is the total number of animals assayed in three independent experiments.
167

Prior studies also suggested that there were differential requirements for HIF-1 and SKN-1 in

oxidative stress conditions. Mutants carrying loss-of-function mutations in *skn-1* have been

shown to decrease the ability of *C. elegans* to survive exposure to agents that cause oxidative

stress [34, 42-44]. In contrast, *C. elegans* carrying loss-of-function mutations in *hif-1* have been

reported to be relatively resistant to peroxide [40]. We compared these phenotypes directly, and

the data are provided in Table 2. These experiments confirmed that, while *skn-1*-deficient

animals were sensitive to t-butyl peroxide, mutants lacking *hif-1* were remarkably resistant to

this oxidative stress: while none of the *skn-1*-deficient mutants survived 6 hours of t-butyl

peroxide treatment, 97.5% of *hif-1*-deficient mutants survived 10 hours of t-butyl peroxide

treatment.

Table 2. Relative requirements for *skn-1* and *hif-1*: survival on t-butyl-peroxide.
180

Exposure time	Genotype	Mean Survival ± SEM	N [a]
6 hours	N2 (wild type)	95.2 ± 2.6	106
	skn-1(zu67)	0.0 ± 0	117
	hif-1(ia04)	100.0 ± 0	120
8 hours	N2 (wild type)	29.08 ± 7.7	106
	skn-1(zu67)	0.0 ± 0	117
	hif-1(ia04)	99.17 ± 0.8	120
10 hours	N2 (wild type)	2.78 ± 2.8	106
	skn-1(zu67)	0.0 ± 0	117
	hif-1(ia04)	97.50 ± 2.5	120

181 [a] n is the total number of animals assayed in three independent experiments.
182

SKN-1/NRF promotes *egl-9* expression

We next sought to discover the mechanism by which SKN-1 inhibits HIF-1 protein levels. In

silico analyses identified a potential SKN-1 binding site in the *egl-9* promoter region (Fig 3A).

EGL-9 is a central inhibitor of HIF-1 protein levels [26, 27] and of HIF-1 transcriptional activity

[29, 30]. This suggested a model in which the SKN-1 DNA-binding complex bound directly to

the *egl-9* regulatory sequences to promote *egl-9* expression, which, in turn, would ultimately

decrease HIF-1 protein levels. To test this, we employed real-time quantitative RT-PCR to

compare *egl-9* mRNA levels in worms fed with *skn-1* RNAi versus control RNAi. To produce

reliable and reproducible results, *egl-9* mRNA levels were quantitated in three independent real-

time quantitative RT-PCR experiments in L4-stage animals in room air or hypoxic conditions

(0.5% oxygen). Each sample was performed with three technical replicates, and they produced

similar C_t values. There are seven isoforms of *egl-9* mRNA transcripts

(https://wormbase.org/species/c_elegans/gene/WBGene00001178#0-9f-10). The real-time

quantitative PCR primer set used in this study can detect six *egl-9* mRNA isoforms. In room air,

skn-1 RNAi decreased *egl-9* mRNA levels by 30% compared to control RNAi (**p < 0.01, from

three independent experiments) (Fig 3B). HIF-1 has been shown to activate *egl-9* mRNA

expression under hypoxia, creating a negative feedback loop [27, 28]. In accordance with this,

the inhibition effects of *skn-1* RNAi on *egl-9* mRNA levels were minimized by placing the

animals in hypoxic conditions (Fig 3B).

Fig 3. Identification of *egl-9* as a potential transcriptional target of SKN-1. (A) Sequence

from the *egl-9* promoter was aligned with established SKN-1 binding sites in *gcs-1*, *med-1*,

and *med-2*. Asterisks identify sequence identities shared by all four promoter regions in this

interval, and predicted SKN-1 binding sites are in red. (B) *skn-1* RNAi decreased *egl-9* mRNA

levels. *egl-9* mRNA levels were quantitated from three independent real-time quantitative RT-

PCR experiments. The values at each time point were normalized to the 0 hour hypoxia control

RNAi. Asterisks indicate significant differences between control RNAi and *skn-1* RNAi at any

given time point. *: *p* < 0.05; **: *p* < 0.01.

To test the hypothesis that conditions that activate SKN-1 can promote *egl-9* promoter activity,

we generated a reporter construct in which 1.6 kb of *egl-9* regulatory sequence directed the

expression of GFP (Fig 4A). To distinguish the effects of SKN-1 on *egl-9* expression from those

of HIF-1, we conducted these experiments in a *hif-1* mutant background. In agreement with prior

studies [45], *Pegl-9:*GFP was visible in several tissues, including the body muscle, vulva,

pharynx, anterior intestine, rectal cells and additional cells in the tail in standard culture

conditions (20 °C) (Figs 4B and 4C). When the animals were treated with heat shock conditions

that had been shown to activate SKN-1 (29 °C for 20 hours) [34], we observed dramatic

induction of *Pegl-9:*GFP in the intestine. (Figs 4D and 4E).

Fig 4. Heat shock alters *Pegl-9:GFP* expression. (A) The *Pegl-9:GFP* construct includes 1.6

kb of sequence 5' to the *egl-9* translational start. GFP coding sequence is diagramed as a green

box. The red oval indicates the position of the putative SKN-1 binding site. (B-E) *Pegl-*

9:GFP expression in L4-stage animals under normal culture conditions and heat shock. Animals

are shown as DIC images (B and D) and corresponding images of GFP fluorescence (C and E).

In all images, the head is to the right. (B and C) Under normal conditions, *Pegl-9:GFP* was

expressed in the body muscle, vulva, pharynx, anterior intestine, rectal cells and additional cells

in the tail. (D and E) After heat shock treatment (29°C for 20 hours), *Pegl-9:GFP* was strongly

induced in the intestine. The*Pegl-9:GFP* expression patterns in the L1, L2, L3 and adults were

similar to that in the L4 worms, under both normal and heat shock conditions (data not shown).

We next asked whether heat shock induction of *Pegl-9:GFP* required *skn-1* function. We found

that heat shock increased *Pegl-9:*GFP by 2.5-fold in animals carrying the wild-type *skn-1* allele.

However, the heat shock induction of *Pegl-9:GFP* was abolished in *skn-1(zu67)* loss-of-function

mutants (Fig 5A). Analyses of another independent *Pegl-9:GFP* transgenic line yielded similar

results (data not shown).

To test the hypothesis that the putative SKN-1 binding site in the *egl-9* promoter was required for

heat shock induction of *Pegl-9:GFP*, we generated the *P(m)egl-9:GFP* construct, which

contained mutations in the putative SKN-1 binding site (in red type in Fig 3A). Heat shock

increased *Pegl-9:*GFP by 2.1-fold. However, heat shock failed to induce the expression of

P(m)egl-9:GFP (Fig 5B). Experiments with a second *P(m)egl-9:GFP* transgenic line gave

similar results (data not shown). Collectively, these data demonstrated that heat shock induction

of *Pegl-9:GFP* required *skn-1* function and the putative SKN-1 binding site in the *egl-9*

promoter.

Fig 5. SKN-1 acts through the putative SKN-1 binding site in the *egl-9* promoter to

activate *egl-9* expression. (A) Heat shock induced *Pegl-9:GFP* in animals carrying the wild-

type s*kn-1* allele, but did not induce the reporter in animals carrying the *skn-1(zu67)* loss-of-

function mutation. The vertical axis shows the log2 fold changes of GFP caused by heat shock

for each strain with standard errors, as determined by four biological replicates. A representative

western blot is shown. For each sample, 20 L4-satge worms were boiled and lysates

corresponding to 10 worms were loaded to each lane. (B) Heat shock increased the expression

of *Pegl-9*:*GFP*, but did not increase the expression of the reporter in which the putative SKN-1

binding site was mutated (*P(m)egl-9:GFP*). The vertical axis shows the log2 fold changes of

GFP caused by heat shock for each strain with standard errors, as determined by five biological

replicates. A representative western blot is shown. For each sample, 20 L4-satge worms were

boiled and lysates corresponding to 10 worms were loaded to each lane.

Discussion

This study provides new insights to the mechanisms that allow animals to respond appropriately to diverse stresses. While HIF-1 and SKN-1 are both stress responsive transcription factors, they have distinct functions. For example, *skn-1*-deficient animals are less able to survive exposure to peroxide, while *hif-1-deficient* mutants are relatively resistant to this oxidizing agent [40]. Conversely, while a deletion mutation in *hif-1* dramatically impairs survival in 0.5% oxygen [25], *skn-1* RNAi has little effect (Table 1). Thus, animals may benefit from cross-talk between these two transcription factors as they are challenged by oxygen deprivation and oxidative stress. Here, we provide evidence that SKN-1 promotes *egl-9* expression, thereby attenuating HIF-1 function.

SKN-1 promotes *egl-9* expression, thereby inhibiting HIF-1

Our data support a model in which SKN-1 binds directly to the *egl-9* promoter to increase *egl-9* expression. This conclusion is further substantiated by the genome wide chromatin immunoprecipitation experiments that determined that SKN-1:GFP associated with DNA sequences in the *egl-9* 5' regulatory region [46]. EGL-9 functions as a cellular oxygen sensor, and it mediates oxygen-dependent degradation of HIF-1 [26]. Hence, *skn-1* RNAi results in increased HIF-1 protein levels (Fig 2B). We expect that this regulatory interaction, in which SKN-1 can quickly down-regulate HIF-1, would allow animals to adapt quickly to changes in cellular conditions.

Prior studies have identified genes that are positively regulated by either SKN-1 or HIF-1, and these gene lists are largely non-overlapping [28, 32, 47-49]. Genes that are commonly regulated by both SKN-1 and HIF-1 include K10H10.2/*cysl-2*, F57B9.1, M05D6.5, and *rhy-1*. Two lines of evidence suggest that *rhy-1* may be a direct target of SKN-1. First, Oliveira et al. (2009)

identified a potential SKN-1 binding site in the *rhy-1* promoter region. Second, the

modENCODE project found that *rhy-1* 5' regulatory sequences associated with SKN-1 *in vivo*

[46]. Like *egl-9, rhy-1* is also a negative regulator of HIF-1 [28, 29]. Collectively, these data

suggest that SKN-1 may act through both *egl-9* and *rhy-1* to reduce HIF-1 function.

Interactions between HIF-1 and SKN-1 are likely to be different in select cells, developmental

stages, or environmental contexts. HIF-1, EGL-9 and SKN-1 each have other developmental

functions, and some of these are specific to certain cell types [34, 50-54]. The downstream

effects of SKN-1 or HIF-1 activation are further influenced by cellular or environmental

contexts. For example, *C. elegans* SKN-1 can be activated by arsenite or by t-butyl

hydroperoxide, but only a subset of SKN-1 targets are activated by either toxicant [32]. Also,

while *C. elegans* SKN-1 and HIF-1 have distinct roles in peroxide and hypoxia stress responses,

their functions overlap in selenium and hydrogen sulfide stress responses [13, 15, 35-37].

Similarly, the mammalian NRF and HIF transcription factors are very sensitive to environmental

and physiological cues or stresses, and their regulatory relationships are context specific.

Ischemia causes tissue hypoxia, which stabilizes HIF transcription factors. Once the ischemic

tissue is reperfused, HIF transcription factors are degraded quickly and NRF2 is up-regulated,

presumably to limit the oxidative damage [4, 5]. However, intermittent hypoxia has been shown

to induce both HIF-1α and NRF2 [6, 7]. While NRF2 signaling activates HIF-1 in several cancer

types [12], studies of the anti-inflammatory drug andrographolide in endothelial cells revealed

interactions between NRF2 and the PHDs that modulate HIF-1 [55]. NRF2 and the HIF

transcription factors have key roles in angiogenesis and iron regulation, and their functions can

converge on developmental processes or feedback loops that modulate their activities [12, 56-

58].

Pnhr-57:GFP as a marker for hypoxia-induced gene expression: discoveries and insights from a genome-wide screen

Prior studies have demonstrated that *nhr-57* was induced by hypoxia in a *hif-1*-dependent

manner and that over-expression of *Pnhr-57*:*GFP* in *egl-9* mutants required *hif-1* [13, 27-30].

Moreover, Bellier *et al*. (2009) found that HIF-1-mediated induction of *nhr-57* helped to protect

C. elegans from the lethal effects of pore-forming toxins [59]. While these studies showed that

nhr-57 is a direct target of HIF-1, other transcription factors must also contribute to its

expression. The studies presented here show that, while genes such as *egl-9* or *rhy-1* clearly

regulate HIF-1 to control *Pnhr-57*:*GFP* expression, many other RNAi treatments can activate

Pnhr-57:*GFP* through *hif-1*-independent pathways. The *Pnhr-57*:*GFP* reporter will continue to

be a valuable marker, but these data inform our interpretations of studies that employ this

reporter. While *Pnhr-57*:*GFP* is clearly regulated by HIF-1, it is important to compare

expression of the reporter in wild-type and *hif-1*-deficient animals before drawing conclusions

about HIF-1 activity.

We found that diverse RNAi treatments that compromise metabolic function or protein

homeostasis increased *Pnhr-57*:*GFP* expression, and this effect did not require a functional *hif-1*

gene. Interestingly, many of the genes integral to these processes have been shown to have roles

in stress response and aging [60-65]. RNAi-mediated depletion of proteasomal components has

also been shown to impact resistance to polyglutamine toxicity and to induce expression of

Pgpdh-1:*GFP*, a marker for osmotic stress and glycerol production [66, 67].

Further characterization of the 179 RNAi treatments that increased *Pnhr-57*:GFP identified 13

genes that had much stronger effects in animals carrying a wild-type *hif-1* gene. These genes

included *vhl-1, egl-9,* and *rhy-1*. These three genes had all been identified in prior studies as

negative regulators of HIF-1 [26, 29]. The succinate dehydrogenase subunit *sdhb-1* was also

found to have *hif-1*-dependent effects. This is especially interesting, since studies in cancer cell

lines have shown that succinate can inhibit the enzymatic activities of HIF prolyl hydroxylases

[68, 69]. *sams-1* and *sbp-1* encode the *C. elegans* S-adenosyl methionine synthetase and the

SREBP homologs, respectively. The RNAi treatments of sams-1 and sbp-1 have a lesser impact

on *Pnhr-57*:GFP levels in *hif-1* mutants, suggesting that the effects of *sams-1* and *sbp-1* RNAi on

the reporter are mediated by HIF-1 (S2 Fig). Both of these genes have key roles in methionine

metabolism and fatty acid biosynthesis [70], and it will be interesting to investigate the ways in

which these important processes intersect with hypoxia response.

Materials and methods

Strains

The following strains were used in this study: wild-type N2 Bristol; ZG430: *Pnhr-*

*57:GFP(iaIs07)*IV; *egl-9(sa307)*V; *hif-1(ia04)*V; *Phif-1:hif-1a:Myc:HA (iaIs28)*; ZG120: *Pnhr-*

*57:GFP(iaIs07)*IV; ZG509: *rrf-3(pk1426)*II; *Pnhr-57:GFP(iaIs07)*IV; ZG508: *rrf-3(pk1426)*II;

*Pnhr-57:GFP(iaIs07)*IV; *hif-1(ia04)*V; ZG429: *hif-1(ia04)*V; *Phif-1:hif-1a:Myc:HA(iaIs28)*;

ZG472: *hif-1(ia04)*V; *Pegl-9:GFP(iaEx84)*; ZG487: *hif-1(ia04)*V; *P(m)egl-9:GFP(iaEx96)*;

ZG488: *skn-1(zu67)*IV; *hif-1(ia04)*V; *Pegl-9:GFP(iaEx84)*. The transgenes expressing epitope-

tagged HIF-1 protein were described and characterized previously [40].

RNAi experiments

The RNAi screen was conducted as previously described [71], with few modifications. Each

bacterial clone (expressing double-stranded RNA for one gene) was cultured in L-broth with 50

ug/mL ampicillin and 12.5 ug/mL tetracycline overnight at 37°C. The following morning, the

bacteria were inoculated into new L-broth with 100 ug/mL ampicillin for 6 hours at 37°C before

seeding on 24-well NGM agar plates with 25 ug/mL carbenicillin and 2 mM IPTG. Each RNAi

clone was plated in duplicate. The following day, 15-25 L1-stage worms were added to each

well. The plates were incubated at 15°C for 5-6 days, and then the worms were screened for

positive *Pnhr-57:*GFP green fluorescence by stereomicroscopy. Bacterial RNAi clones that

increased the reporter were rescreened in two independent replicates, and the plasmid inserts

were validated by sequencing.

skn-1 RNAi causes maternal-effect lethality, in this study we examined the effects of first

generation *skn-1* RNAi. Dead egg percentages given by the first generation *skn-1* RNAi adults

were measured to check the *skn-1* RNAi efficiency. We routinely achieved as high as 90% dead

egg percentages from the first generation *skn-1* RNAi adults, indicating high *skn-1* RNAi

efficiency.

Hypoxia stress and oxidative stress assays

To assess the relative effects of t-butyl-peroxide exposure, animals in the first day of adulthood

were placed on NGM plates containing 7.5 mM t-butyl-peroxide, in the presence of bacterial

food. The survival was scored after treating the animals for 6, 8 or 10 hours.

For hypoxia experiments, adults were allowed to lay eggs on standard NGM plates with OP50

bacterial food for 2 hours. The adults were then removed, and the plates with embryos were

placed in a hypoxia chamber (0.5% oxygen; 21°C) for 24 hours. After 24 hours, the plates were

removed from the hypoxia chamber, and the un-hatched eggs were counted immediately. The

plates were then maintained in room air (21°C). The adult worms were counted at 72 hour since

the eggs had been laid. Wild-type control animals hatched within 24 hours and reached

adulthood within 72 hours.

Pegl-9:GFP expression constructs

To generate the *Pegl-9:GFP* construct, a fragment that contained 1.6 kb of sequence upstream of the initiation ATG of *egl-9* gene was amplified by PCR using the forward primer 5'-CGCGCATGCGTGTATGTGTGTGAAAGAG-3' and the reverse primer 5'-GCGGTCGACGCAACTTTTTTCTGTCACATTCAG-3'. The PCR product was cloned into the green fluorescence protein (GFP) vector pPD95.75 (gift from Andrew Fire). To create the *P(m)egl-9:GFP* point mutation construct, the predicted SKN-1 binding site TTTGTCAT [34, 72]was altered to CGACGGGC. Transgenic animals were generated by injection of DNA into the gonadal syncitium, using standard methods with *rol-6* (pRF4) as the co-injection marker [73]. For each construct, two independent transgenic lines were generated and assayed.

Protein Blots

To assay the expression of GFP or HIF-1 proteins, 20-100 L4-stage worms were collected and boiled for 5 min in 1X SDS sample buffer, and the lysates were size fractionated on polyacrylamide gels and analyzed by Western blots. The GFP-specific mouse monoclonal antibody (from Roche) was used at 1:500. The HA-specific mouse monoclonal antibody (from Cell Signaling) was used at 1:250. The secondary antibody (goat anti-mouse IgG+IgM from Biorad) was used at 1:2000 dilutions. The western blot images were analyzed by the Image J software. For each assay, three to five independent biological replicates were included.

RNA extraction and real-time quantitative RT-PCR

Total RNA was isolated from synchronized L4-stage animals using Trizol (Invitrogen) and RNeasy Mini Kit (Qiagen). After being treated by RNase free DNase (Promega), total RNA was reverse transcribed to complementary DNA using Oligo dT_{18} primer and AffinityScript reverse

transcriptase (Stratagene). Real-time quantitative PCR was performed using the iQ SYBR

GREEN supermix (Bio-Rad) and Stratagene Mx4000 multiplex PCR system. In the assay, three

biological replicates were included. And for each sample, three technical replicates were

performed and they gave similar C_t values. And cDNA from 62.5 ng of total RNA was added to

each PCR reaction. Relative mRNA quantification was performed using the efficiency-corrected

comparative quantification method [74]. *inf-1*, a gene not regulated by hypoxia, was used as the

reference gene [28]. The primer sequences for *inf-1* real-time quantitative PCR are included in

S1 Fig. The primers for *egl-9* real-time quantitative PCR are the forward primer 5'-

GCCGACTTTCAATCCACTTC-3' and reverse primer 5'- AATGATCGGAGATCGACTGG-

3'. There are seven isoforms of *egl-9* mRNA transcripts

(https://wormbase.org/species/c_elegans/gene/WBGene00001178#0-9f-10). This primer set can

detect six out of seven *egl-9* mRNA isoforms. The isoform d.1 will not be detected by this

primer set, because the forward primer is located within the eighth intron of the unspliced

isoform d.1.

References

1.   Beckman KB, Ames BN. The free radical theory of aging matures. Physiol Rev. 1998;78(2):547-
81. Epub 1998/04/30. PubMed PMID: 9562038.
2.   Finkel T. Oxidant signals and oxidative stress. Current opinion in cell biology. 2003;15(2):247-
54. Epub 2003/03/22. PubMed PMID: 12648682.
3.   Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing. Nature.
2000;408(6809):239-47. Epub 2000/11/23. doi: 10.1038/35041687. PubMed PMID: 11089981.
4.   Kim YJ, Ahn JY, Liang P, Ip C, Zhang Y, Park YM. Human prx1 gene is a target of Nrf2 and is
up-regulated by hypoxia/reoxygenation: implication to tumor biology. Cancer Res. 2007;67(2):546-54.
Epub 2007/01/20. doi: 67/2/546 [pii]
10.1158/0008-5472.CAN-06-2401. PubMed PMID: 17234762.
5.   Leonard MO, Kieran NE, Howell K, Burne MJ, Varadarajan R, Dhakshinamoorthy S, et al.
Reoxygenation-specific activation of the antioxidant transcription factor Nrf2 mediates cytoprotective
gene expression in ischemia-reperfusion injury. FASEB journal : official publication of the Federation of
American Societies for Experimental Biology. 2006;20(14):2624-6. Epub 2006/12/05. doi: 10.1096/fj.06-
5097fje. PubMed PMID: 17142801.
6.   Malec V, Gottschald OR, Li S, Rose F, Seeger W, Hanze J. HIF-1 alpha signaling is augmented
during intermittent hypoxia by induction of the Nrf2 pathway in NOX1-expressing adenocarcinoma A549

cells. Free radical biology & medicine. 2010;48(12):1626-35. Epub 2010/03/30. doi:
10.1016/j.freeradbiomed.2010.03.008. PubMed PMID: 20347035.
7.   Martinive P, Defresne F, Bouzin C, Saliez J, Lair F, Gregoire V, et al. Preconditioning of the
tumor vasculature and tumor cells by intermittent hypoxia: implications for anticancer therapies. Cancer
Res. 2006;66(24):11736-44. doi: 10.1158/0008-5472.CAN-06-2056. PubMed PMID: 17178869.
8.   Thimmulappa RK, Mai KH, Srisuma S, Kensler TW, Yamamoto M, Biswal S. Identification of
Nrf2-regulated genes induced by the chemopreventive agent sulforaphane by oligonucleotide microarray.
Cancer Res. 2002;62(18):5196-203. Epub 2002/09/18. PubMed PMID: 12234984.
9.   Hayes JD, McMahon M. Molecular basis for the contribution of the antioxidant responsive
element to cancer chemoprevention. Cancer Lett. 2001;174(2):103-13. Epub 2001/11/02. doi:
S0304383501006954 [pii]. PubMed PMID: 11689285.
10.   Hough RB, Piatigorsky J. Preferential transcription of rabbit Aldh1a1 in the cornea: implication
of hypoxia-related pathways. Molecular and cellular biology. 2004;24(3):1324-40. Epub 2004/01/20.
PubMed PMID: 14729976; PubMed Central PMCID: PMC321433.
11.   Yeligar SM, Machida K, Kalra VK. Ethanol-induced HO-1 and NQO1 are differentially regulated
by HIF-1alpha and Nrf2 to attenuate inflammatory cytokine expression. The Journal of biological
chemistry. 2010;285(46):35359-73. Epub 2010/09/14. doi: 10.1074/jbc.M110.138636. PubMed PMID:
20833713; PubMed Central PMCID: PMC2975160.
12.   Toth RK, Warfel NA. Strange Bedfellows: Nuclear Factor, Erythroid 2-Like 2 (Nrf2) and
Hypoxia-Inducible Factor 1 (HIF-1) in Tumor Hypoxia. Antioxidants (Basel). 2017;6(2). doi:
10.3390/antiox6020027. PubMed PMID: 28383481; PubMed Central PMCID: PMCPMC5488007.
13.   Budde MW, Roth MB. Hydrogen sulfide increases hypoxia-inducible factor-1 activity
independently of von Hippel-Lindau tumor suppressor-1 in C. elegans. Mol Biol Cell. 2010;21(1):212-7.
Epub 2009/11/06. doi: E09-03-0199 [pii]
10.1091/mbc.E09-03-0199. PubMed PMID: 19889840; PubMed Central PMCID: PMC2801715.
14.   Powell-Coffman JA. Hypoxia signaling and resistance in C. elegans. Trends Endocrinol Metab.
2010;21(7):435-40. Epub 2010/03/26. doi: S1043-2760(10)00041-X [pii]
10.1016/j.tem.2010.02.006. PubMed PMID: 20335046.
15.   Romanelli-Credrez L, Doitsidou M, Alkema MJ, Salinas G. HIF-1 Has a Central Role in
Caenorhabditis elegans Organismal Response to Selenium. Front Genet. 2020;11:63. doi:
10.3389/fgene.2020.00063. PubMed PMID: 32161616; PubMed Central PMCID: PMCPMC7052493.
16.   Saldanha JN, Parashar A, Pandey S, Powell-Coffman JA. Multiparameter behavioral analyses
provide insights to mechanisms of cyanide resistance in Caenorhabditis elegans. Toxicol Sci.
2013;135(1):156-68. doi: 10.1093/toxsci/kft138. PubMed PMID: 23805000; PubMed Central PMCID:
PMCPMC3748764.
17.   Shao Z, Zhang Y, Ye Q, Saldanha JN, Powell-Coffman JA. C. elegans SWAN-1 Binds to EGL-9
and regulates HIF-1-mediated resistance to the bacterial pathogen Pseudomonas aeruginosa PAO1. PLoS
Pathog. 2010;6(8):e1001075. doi: 10.1371/journal.ppat.1001075. PubMed PMID: 20865124; PubMed
Central PMCID: PMCPMC2928816.
18.   Kaelin WG, Jr., Ratcliffe PJ. Oxygen sensing by metazoans: the central role of the HIF
hydroxylase pathway. Mol Cell. 2008;30(4):393-402. Epub 2008/05/24. doi: S1097-2765(08)00292-X
[pii]
10.1016/j.molcel.2008.04.009. PubMed PMID: 18498744.
19.   Semenza GL. Oxygen sensing, homeostasis, and disease. N Engl J Med. 2011;365(6):537-47. doi:
10.1056/NEJMra1011165. PubMed PMID: 21830968.
20.   Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-
helix-PAS heterodimer regulated by cellular O2 tension. Proc Natl Acad Sci U S A. 1995;92(12):5510-4.
doi: 10.1073/pnas.92.12.5510. PubMed PMID: 7539918; PubMed Central PMCID: PMCPMC41725.
21.   Maxwell PH, Wiesener MS, Chang GW, Clifford SC, Vaux EC, Cockman ME, et al. The tumour
suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. Nature.
1999;399(6733):271-5. doi: 10.1038/20459. PubMed PMID: 10353251.

22.   Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, et al. Targeting of HIF-alpha
to the von Hippel-Lindau ubiquitylation complex by O2-regulated prolyl hydroxylation. Science.
2001;292(5516):468-72. doi: 10.1126/science.1059796. PubMed PMID: 11292861.
23.   Ivan M, Kondo K, Yang H, Kim W, Valiando J, Ohh M, et al. HIFalpha targeted for VHL-
mediated destruction by proline hydroxylation: implications for O2 sensing. Science.
2001;292(5516):464-8. doi: 10.1126/science.1059817. PubMed PMID: 11292862.
24.   Powell-Coffman JA, Bradfield CA, Wood WB. Caenorhabditis elegans orthologs of the aryl
hydrocarbon receptor and its heterodimerization partner the aryl hydrocarbon receptor nuclear
translocator. Proc Natl Acad Sci U S A. 1998;95(6):2844-9. Epub 1998/04/18. PubMed PMID: 9501178.
25.   Jiang H, Guo R, Powell-Coffman JA. The Caenorhabditis elegans hif-1 gene encodes a bHLH-
PAS protein that is required for adaptation to hypoxia. Proc Natl Acad Sci U S A. 2001;98(14):7916-21.
Epub 2001/06/28. doi: 10.1073/pnas.141234698
141234698 [pii]. PubMed PMID: 11427734; PubMed Central PMCID: PMC35443.
26.   Epstein AC, Gleadle JM, McNeill LA, Hewitson KS, O'Rourke J, Mole DR, et al. C. elegans
EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl
hydroxylation. Cell. 2001;107(1):43-54. Epub 2001/10/12. doi: S0092-8674(01)00507-4 [pii]. PubMed
PMID: 11595184.
27.   Bishop T, Lau KW, Epstein AC, Kim SK, Jiang M, O'Rourke D, et al. Genetic analysis of
pathways regulated by the von Hippel-Lindau tumor suppressor in Caenorhabditis elegans. PLoS Biol.
2004;2(10):e289. Epub 2004/09/14. doi: 10.1371/journal.pbio.0020289. PubMed PMID: 15361934.
28.   Shen C, Nettleton D, Jiang M, Kim SK, Powell-Coffman JA. Roles of the HIF-1 hypoxia-
inducible factor during hypoxia response in Caenorhabditis elegans. The Journal of biological chemistry.
2005;280(21):20580-8. Epub 2005/03/23. doi: M501894200 [pii]
10.1074/jbc.M501894200. PubMed PMID: 15781453.
29.   Shen C, Shao Z, Powell-Coffman JA. The Caenorhabditis elegans rhy-1 gene inhibits HIF-1
hypoxia-inducible factor activity in a negative feedback loop that does not include vhl-1. Genetics.
2006;174(3):1205-14. Epub 2006/09/19. doi: genetics.106.063594 [pii]
10.1534/genetics.106.063594. PubMed PMID: 16980385.
30.   Shao Z, Zhang Y, Powell-Coffman JA. Two distinct roles for EGL-9 in the regulation of HIF-1-
mediated gene expression in Caenorhabditis elegans. Genetics. 2009;183(3):821-9. Epub 2009/09/10. doi:
genetics.109.107284 [pii]
10.1534/genetics.109.107284. PubMed PMID: 19737748; PubMed Central PMCID: PMC2778979.
31.   Walker AK, See R, Batchelder C, Kophengnavong T, Gronniger JT, Shi Y, et al. A conserved
transcription motif suggesting functional parallels between Caenorhabditis elegans SKN-1 and
Cap'n'Collar-related basic leucine zipper proteins. The Journal of biological chemistry.
2000;275(29):22166-71. Epub 2000/04/15. doi: 10.1074/jbc.M001746200
M001746200 [pii]. PubMed PMID: 10764775.
32.   Oliveira RP, Abate JP, Dilks K, Landis J, Ashraf J, Murphy CT, et al. Condition-adapted stress
and longevity gene regulation by Caenorhabditis elegans SKN-1/Nrf. Aging Cell. 2009;8(5):524-41.
Epub 2009/07/07. doi: ACE501 [pii]
10.1111/j.1474-9726.2009.00501.x. PubMed PMID: 19575768.
33.   Park SK, Tedesco PM, Johnson TE. Oxidative stress and longevity in Caenorhabditis elegans as
mediated by SKN-1. Aging Cell. 2009;8(3):258-69. Epub 2009/07/25. doi: 10.1111/j.1474-
9726.2009.00473.x. PubMed PMID: 19627265; PubMed Central PMCID: PMC2762118.
34.   An JH, Blackwell TK. SKN-1 links C. elegans mesendodermal specification to a conserved
oxidative stress response. Genes Dev. 2003;17(15):1882-93. Epub 2003/07/19. doi:
10.1101/gad.1107803. PubMed PMID: 12869585.
35.   Horsman JW, Heinis FI, Miller DL. A Novel Mechanism To Prevent H2S Toxicity in
Caenorhabditis elegans. Genetics. 2019;213(2):481-90. doi: 10.1534/genetics.119.302326. PubMed
PMID: 31371406; PubMed Central PMCID: PMCPMC6781907.
36.   Li WH, Chang CH, Huang CW, Wei CC, Liao VH. Selenite enhances immune response against

Pseudomonas aeruginosa PA14 via SKN-1 in Caenorhabditis elegans. PLoS One. 2014;9(8):e105810.
doi: 10.1371/journal.pone.0105810. PubMed PMID: 25147937; PubMed Central PMCID:
PMCPMC4141825.
37.   Miller DL, Budde MW, Roth MB. HIF-1 and SKN-1 coordinate the transcriptional response to
hydrogen sulfide in Caenorhabditis elegans. PLoS One. 2011;6(9):e25476. Epub 2011/10/08. doi:
10.1371/journal.pone.0025476
PONE-D-11-09953 [pii]. PubMed PMID: 21980473; PubMed Central PMCID: PMC3183046.
38.   Kamath RS, Ahringer J. Genome-wide RNAi screening in Caenorhabditis elegans. Methods.
2003;30(4):313-21. Epub 2003/06/28. doi: S1046202303000501 [pii]. PubMed PMID: 12828945.
39.   Inoue H, Hisamoto N, An JH, Oliveira RP, Nishida E, Blackwell TK, et al. The C. elegans p38
MAPK pathway regulates nuclear localization of the transcription factor SKN-1 in oxidative stress
response. Genes Dev. 2005;19(19):2278-83. Epub 2005/09/17. doi: gad.1324805 [pii]
10.1101/gad.1324805. PubMed PMID: 16166371.
40.   Zhang Y, Shao Z, Zhai Z, Shen C, Powell-Coffman JA. The HIF-1 hypoxia-inducible factor
modulates lifespan in C. elegans. PLoS One. 2009;4(7):e6348. Epub 2009/07/28. doi:
10.1371/journal.pone.0006348. PubMed PMID: 19633713.
41.   Padilla PA, Nystul TG, Zager RA, Johnson AC, Roth MB. Dephosphorylation of cell cycle-
regulated proteins correlates with anoxia-induced suspended animation in Caenorhabditis elegans. Mol
Biol Cell. 2002;13(5):1473-83. Epub 2002/05/15. doi: 10.1091/mbc.01-12-0594. PubMed PMID:
12006646.
42.   Li X, Matilainen O, Jin C, Glover-Cutter KM, Holmberg CI, Blackwell TK. Specific SKN-1/Nrf
stress responses to perturbations in translation elongation and proteasome activity. PLoS Genet.
2011;7(6):e1002119. Epub 2011/06/23. doi: 10.1371/journal.pgen.1002119. PubMed PMID: 21695230;
PubMed Central PMCID: PMC3111486.
43.   Tullet JM, Hertweck M, An JH, Baker J, Hwang JY, Liu S, et al. Direct inhibition of the
longevity-promoting factor SKN-1 by insulin-like signaling in C. elegans. Cell. 2008;132(6):1025-38.
Epub 2008/03/25. doi: S0092-8674(08)00130-X [pii]
10.1016/j.cell.2008.01.030. PubMed PMID: 18358814.
44.   Wang J, Robida-Stubbs S, Tullet JM, Rual JF, Vidal M, Blackwell TK. RNAi screening
implicates a SKN-1-dependent transcriptional response in stress resistance and longevity deriving from
translation inhibition. PLoS Genet. 2010;6(8). Epub 2010/08/12. doi: 10.1371/journal.pgen.1001048.
PubMed PMID: 20700440; PubMed Central PMCID: PMC2916858.
45.   Darby C, Cosma CL, Thomas JH, Manoil C. Lethal paralysis of Caenorhabditis elegans by
Pseudomonas aeruginosa. Proc Natl Acad Sci U S A. 1999;96(26):15202-7. Epub 1999/12/28. PubMed
PMID: 10611362.
46.   Niu W, Lu ZJ, Zhong M, Sarov M, Murray JI, Brdlik CM, et al. Diverse transcription factor
binding features revealed by genome-wide ChIP-seq in C. elegans. Genome research. 2011;21(2):245-54.
Epub 2010/12/24. doi: 10.1101/gr.114587.110. PubMed PMID: 21177963; PubMed Central PMCID:
PMC3032928.
47.   Glover-Cutter KM, Lin S, Blackwell TK. Integration of the unfolded protein and oxidative stress
responses through SKN-1/Nrf. PLoS Genet. 2013;9(9):e1003701. doi: 10.1371/journal.pgen.1003701.
PubMed PMID: 24068940; PubMed Central PMCID: PMCPMC3772064.
48.   Paek J, Lo JY, Narasimhan SD, Nguyen TN, Glover-Cutter K, Robida-Stubbs S, et al.
Mitochondrial SKN-1/Nrf mediates a conserved starvation response. Cell Metab. 2012;16(4):526-37. doi:
10.1016/j.cmet.2012.09.007. PubMed PMID: 23040073; PubMed Central PMCID: PMCPMC3774140.
49.   Robida-Stubbs S, Glover-Cutter K, Lamming DW, Mizunuma M, Narasimhan SD, Neumann-
Haefelin E, et al. TOR signaling and rapamycin influence longevity by regulating SKN-1/Nrf and DAF-
16/FoxO. Cell Metab. 2012;15(5):713-24. doi: 10.1016/j.cmet.2012.04.007. PubMed PMID: 22560223;
PubMed Central PMCID: PMCPMC3348514.
50.   Bishop NA, Guarente L. Two neurons mediate diet-restriction-induced longevity in C. elegans.
Nature. 2007;447(7144):545-9. Epub 2007/06/01. doi: nature05904 [pii]

10.1038/nature05904. PubMed PMID: 17538612.

51.   Chen D, Thomas EL, Kapahi P. HIF-1 modulates dietary restriction-mediated lifespan extension

via IRE-1 in Caenorhabditis elegans. PLoS Genet. 2009;5(5):e1000486. Epub 2009/05/23. doi:

10.1371/journal.pgen.1000486. PubMed PMID: 19461873.

52.   Park EC, Ghose P, Shao Z, Ye Q, Kang L, Xu XZ, et al. Hypoxia regulates glutamate receptor

trafficking through an HIF-independent mechanism. The EMBO journal. 2012;31(6):1379-93. Epub

2012/01/19. doi: 10.1038/emboj.2011.499. PubMed PMID: 22252129; PubMed Central PMCID:

PMC3321172.

53.   Pocock R, Hobert O. Oxygen levels affect axon guidance and neuronal migration in

Caenorhabditis elegans. Nat Neurosci. 2008;11(8):894-900. Epub 2008/07/01. doi: nn.2152 [pii]

10.1038/nn.2152. PubMed PMID: 18587389.

54.   Chang AJ, Bargmann CI. Hypoxia and the HIF-1 transcriptional pathway reorganize a neuronal

circuit for oxygen-dependent behavior in Caenorhabditis elegans. Proc Natl Acad Sci U S A.

2008;105(20):7321-6. doi: 10.1073/pnas.0802164105. PubMed PMID: 18477695; PubMed Central

PMCID: PMCPMC2438248.

55.   Lin HC, Su SL, Lu CY, Lin AH, Lin WC, Liu CS, et al. Andrographolide inhibits hypoxia-

induced HIF-1alpha-driven endothelin 1 secretion by activating Nrf2/HO-1 and promoting the expression

of prolyl hydroxylases 2/3 in human endothelial cells. Environ Toxicol. 2017;32(3):918-30. doi:

10.1002/tox.22293. PubMed PMID: 27297870.

56.   Duarte TL, Talbot NP, Drakesmith H. NRF2 and Hypoxia-Inducible Factors: Key Players in the

Redox Control of Systemic Iron Homeostasis. Antioxid Redox Signal. 2020. doi: 10.1089/ars.2020.8148.

PubMed PMID: 32791852.

57.   Li L, Pan H, Wang H, Li X, Bu X, Wang Q, et al. Interplay between VEGF and Nrf2 regulates

angiogenesis due to intracranial venous hypertension. Sci Rep. 2016;6:37338. doi: 10.1038/srep37338.

PubMed PMID: 27869147; PubMed Central PMCID: PMCPMC5116754.

58.   Wong BW, Kuchnio A, Bruning U, Carmeliet P. Emerging novel functions of the oxygen-sensing

prolyl hydroxylase domain enzymes. Trends Biochem Sci. 2013;38(1):3-11. doi:

10.1016/j.tibs.2012.10.004. PubMed PMID: 23200187.

59.   Bellier A, Chen CS, Kao CY, Cinar HN, Aroian RV. Hypoxia and the hypoxic response pathway

protect against pore-forming toxins in C. elegans. PLoS Pathog. 2009;5(12):e1000689. Epub 2009/12/17.

doi: 10.1371/journal.ppat.1000689. PubMed PMID: 20011506.

60.   Lee SS, Lee RY, Fraser AG, Kamath RS, Ahringer J, Ruvkun G. A systematic RNAi screen

identifies a critical role for mitochondria in C. elegans longevity. Nat Genet. 2003;33(1):40-8. Epub

2002/11/26. doi: 10.1038/ng1056

ng1056 [pii]. PubMed PMID: 12447374.

61.   Hansen M, Hsu AL, Dillin A, Kenyon C. New genes tied to endocrine, metabolic, and dietary

regulation of lifespan from a Caenorhabditis elegans genomic RNAi screen. PLoS Genet. 2005;1(1):119-

28. Epub 2005/08/17. doi: 10.1371/journal.pgen.0010017. PubMed PMID: 16103914; PubMed Central

PMCID: PMC1183531.

62.   Hamilton B, Dong Y, Shindo M, Liu W, Odell I, Ruvkun G, et al. A systematic RNAi screen for

longevity genes in C. elegans. Genes Dev. 2005;19(13):1544-55. Epub 2005/07/07. doi: 19/13/1544 [pii]

10.1101/gad.1308205. PubMed PMID: 15998808.

63.   Curran SP, Ruvkun G. Lifespan regulation by evolutionarily conserved genes essential for

viability. PLoS Genet. 2007;3(4):e56. Epub 2007/04/07. doi: 07-PLGE-RA-0112R1 [pii]

10.1371/journal.pgen.0030056. PubMed PMID: 17411345.

64.   Kim Y, Sun H. Functional genomic approach to identify novel genes involved in the regulation of

oxidative stress resistance and animal lifespan. Aging Cell. 2007;6(4):489-503. Epub 2007/07/05. doi:

ACE302 [pii]

10.1111/j.1474-9726.2007.00302.x. PubMed PMID: 17608836.

65.   Sugimoto T, Mori C, Takanami T, Sasagawa Y, Saito R, Ichiishi E, et al. Caenorhabditis elegans

par2.1/mtssb-1 is essential for mitochondrial DNA replication and its defect causes comprehensive

transcriptional alterations including a hypoxia response. Experimental cell research. 2008;314(1):103-14.
Epub 2007/09/29. doi: 10.1016/j.yexcr.2007.08.015. PubMed PMID: 17900564.
66.   Lamitina T, Huang CG, Strange K. Genome-wide RNAi screening identifies protein damage as a
regulator of osmoprotective gene expression. Proc Natl Acad Sci U S A. 2006;103(32):12173-8. Epub
2006/08/02. doi: 0602987103 [pii]
10.1073/pnas.0602987103. PubMed PMID: 16880390.
67.   Nollen EA, Garcia SM, van Haaften G, Kim S, Chavez A, Morimoto RI, et al. Genome-wide
RNA interference screen identifies previously undescribed regulators of polyglutamine aggregation. Proc
Natl Acad Sci U S A. 2004;101(17):6403-8. Epub 2004/04/16. doi: 10.1073/pnas.0307697101
0307697101 [pii]. PubMed PMID: 15084750.
68.   Selak MA, Armour SM, MacKenzie ED, Boulahbel H, Watson DG, Mansfield KD, et al.
Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl hydroxylase.
Cancer Cell. 2005;7(1):77-85. Epub 2005/01/18. doi: S153561080400368X [pii]
10.1016/j.ccr.2004.11.022. PubMed PMID: 15652751.
69.   Koivunen P, Hirsila M, Remes AM, Hassinen IE, Kivirikko KI, Myllyharju J. Inhibition of
hypoxia-inducible factor (HIF) hydroxylases by citric acid cycle intermediates: possible links between
cell metabolism and stabilization of HIF. The Journal of biological chemistry. 2007;282(7):4524-32.
Epub 2006/12/22. doi: M610415200 [pii]
10.1074/jbc.M610415200. PubMed PMID: 17182618.
70.   Walker AK, Jacobs RL, Watts JL, Rottiers V, Jiang K, Finnegan DM, et al. A conserved SREBP-
1/phosphatidylcholine feedback circuit regulates lipogenesis in metazoans. Cell. 2011;147(4):840-52.
Epub 2011/11/01. doi: 10.1016/j.cell.2011.09.045. PubMed PMID: 22035958; PubMed Central PMCID:
PMC3384509.
71.   Simmer F, Moorman C, van der Linden AM, Kuijk E, van den Berghe PV, Kamath RS, et al.
Genome-wide RNAi of C. elegans using the hypersensitive rrf-3 strain reveals novel gene functions.
PLoS Biol. 2003;1(1):E12. Epub 2003/10/14. doi: 10.1371/journal.pbio.0000012. PubMed PMID:
14551910; PubMed Central PMCID: PMC212692.
72.   Blackwell TK, Bowerman B, Priess JR, Weintraub H. Formation of a monomeric DNA binding
domain by Skn-1 bZIP and homeodomain elements. Science. 1994;266(5185):621-8. Epub 1994/10/28.
PubMed PMID: 7939715.
73.   Mello CC, Kramer JM, Stinchcomb D, Ambros V. Efficient gene transfer in C.elegans:
extrachromosomal maintenance and integration of transforming sequences. The EMBO journal.
1991;10(12):3959-70. Epub 1991/12/01. PubMed PMID: 1935914.
74.   Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic
Acids Res. 2001;29(9):e45. Epub 2001/05/09. PubMed PMID: 11328886; PubMed Central PMCID:
PMC55695.
668

Supporting information

S1 Fig. *nhr-57* promoter HIF-1 chromatin immunoprecipitation experiments. (A) *nhr-57*

promoter sequence and positions of primers used for real-time quantitative PCR assays in HIF-1

chromatin immunoprecipitation experiments. (B) Primer sequences for real-time quantitative PCR assays

in *nhr-57* promoter HIF-1 chromatin immunoprecipitation experiments. (C) Chromatin co-

immunoprecipitation data. In these experiments the endogenous *hif-1* locus was disrupted by the

ia04 large deletion and HIF-1 function was restored by the *Phif-1:hif-1a:Myc:HA* transgene [40].

The relative amounts of *nhr-57* promoter regions that co-immunoprecipitated with HIF-

1:Myc:HA was determined by real-time quantitative PCR. The bars show the average enriched

fold from at least three independent replicates. *inf-1*, a gene not regulated by HIF-1, was used as

the reference gene.

S2 Fig. RNAi inactivation of *sams-1* or *sbp-1* increased *Pnhr-57:GFP* expression. (A) RNAi

for *sams-1* (S-adenosyl methionine synthetase) increased *Pnhr-57*:GFP expression more than 7-

fold in animals carrying the wild-type *hif-1* allele relative to control RNAi, and increased the

reporter 3-fold in animals carrying the *hif-1(ia04)* deletion. The difference in RNAi effect

between *hif-1(+)* and *hif-1(ia04)* strains is statistically significant (*$p < 0.05$, from six

independent experiments, by student *t*-test). (B) RNAi for the SREBP homolog *sbp-1* increased

expression of the reporter more than 3-fold in animals carrying the wild-type *hif-1* allele, but had

no effect on *Pnhr-57*:GFP expression in *hif-1(ia04)* mutants. The difference in RNAi effect

between *hif-1(+)* and *hif-1(ia04)* strains is statistically significant (*$p < 0.05$, from five

independent experiments, by student *t*-test). GFP levels were determined by protein blots, and

the control animals were fed on bacteria carrying the empty RNAi vector (L4440). The

experiments were conducted in RNAi-sensitive strains (*rrf-3(pk1426)*).

S1 Table. Genes increased *Pnhr-57:GFP* expression when knocked-down by RNAi.

S2 Table. Genes for which RNAi caused *hif-1*-dependent increase of *Pnhr-57:GFP*

expression.

A

RNAi inactivation of negative regulator of HIF-1 activity

B

- Metabolism: Electron transport chain
- Metabolism: Mitochondrial translation
- Metablism: Others
- Protein folding and turnover
- Vesicular transport
- Transporters and channels
- Signal transduction and cytoskeleton
- Transcription/translation/DNA/RNA processing
- Others

24%
13%
13%
18%
4%
6%
8%
5%
9%

A

egl-9	-365	CTTTGTCATCGTGTCTCTTTGATCCG -340
gcs-1	-122	CTTTATCATCATGA-GATTTAATGTT -98
med-1	-124	CTCTGTCATCATGATGATTTTTGGAG -99
med-2	-124	CTCTGTCATCATGATGATTTTTAGAG -99
		** * ***** ** ***

B

egl-9 mRNA level

control RNAi
skn-1 RNAi

Hypoxia treatment time (hours)

A

-1654 ——————————————————————●————————————— ▢
                                     -356

B C

D E