Yeast proteins reversibly aggregate like amphiphilic molecules

Pouria Dasmeh[1,3,4], Andreas Wagner[1,2,4]

[1]Institute for Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland. [2]The Santa Fe Institute, Santa Fe, New Mexico, [3]Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA USA 02139. [4]Swiss Institute of Bioinformatics (SIB).

Abstract:

More than a hundred proteins in yeast reversibly aggregate and phase-separate in response to various stressors, such as nutrient depletion and heat shock. We know little about the sequence and structural features behind this ability, which has not been characterized on a proteome-wide level. To identify the distinctive features of aggregation-prone regions, we apply machine learning algorithms to genome-scale limited proteolysis-mass spectrometry data from 96 yeast proteins that phase-separate upon heat shock. We find that the aggregation-prone regions (APRs) of our study proteins are significantly enriched in aliphatic residues and depleted in positively charged amino acids. Aggregator proteins with longer APRs show a greater propensity to aggregate, a relationship that can be explained by equilibrium statistical thermodynamics. Altogether, our observations suggest that proteome-wide reversible protein aggregation is mediated by sequence-encoded properties. Aggregating proteins resemble supra-molecular amphiphiles, where APRs are the hydrophobic parts, and non-APRs are the hydrophilic parts.

Body:

Proteins can aggregate reversibly or irreversibly. Irreversible aggregation is often pathological, indicates damaged cellular regulation[1-3], and is involved in multiple diseases such as Alzheimer's and Parkinson's diseases[4,5]. Reversible aggregation, however, can be beneficial and help cells survive stressors. For example, in yeast more than a hundred proteins in multiple subcellular compartments form reversible aggregates in response to nutrient starvation, heat shock, or chemical stress[6]. These aggregated proteins are not misfolded or tagged for degradation[7]. Instead, they help increase cells re-initiate growth during recovery from stress by protecting metabolic enzymes from degradation[6-9].

Reversible protein aggregation is a special case of a widespread phenomenon called protein phase-separation. In this process, a well-mixed solution of proteins de-mixes into two phases of high and low densities[10]. Proteins can phase separate with other proteins, but also with RNA or DNA molecules, into biomolecular condensates that regulate transcription[11], chromatin states[12,13], and RNA metabolism[14]. We know little about the sequence and structural features behind reversible phase separation, which has not been characterized on a proteome-wide level. In this work, we provide such a view and characterize sequence features that can help predict whether proteins will aggregate reversibly.

We took advantage of recent proteome-wide data from limited proteolysis-mass spectrometry (LiP-MS)[20,21]. This method permits the detection of both pronounced and subtle changes that protein and peptide structures experience in response to stressors such as heat and osmotic shock. Recently, Cappelletti *et. al.* characterized structural

changes within 96 proteins that reversibly aggregate upon heat shock in yeast[22]. These

proteins are also known as aggregators. They have diverse biological functions, such as

telomeric DNA-binding, RNA-helicase activity, or ribosome assembly. Because they are

functionally diverse, they constitute an excellent dataset to examine sequence features

that facilitate reversible protein aggregation. We studied regions within these proteins

whose proteolysis resistance changes significantly upon aggregation [22]. We refer to these

regions as aggregation-prone regions.

What are the characteristic sequence features of APRs? To answer this question,

we first performed a principal component analysis on the frequency of 20 amino acids in

all 270 APRs of yeast aggregators (Dataset S1). As shown in Figure 1A-B, the first and

the second principal components only accounted for ~12% and ~10% of the variation.

The PCA biplot for these components shows that no specific combinations of few amino

acids preferentially occur in APR sequences (Figure 1B), a further indication that APRs

do not show the kind of reduced sequence complexity characteristic of disordered

regions. We also looked for linear sequence motifs in APRs using DALEL[29], an algorithm

for the exhaustive identification of degenerate sequence motifs, but found no such

enriched motifs.

We then compared the frequencies of amino acids, dipeptides, and tripeptides in

APRs with the rest of the protein sequence in aggregators (Dataset S2). Here, we found

a significant depletion of the positively charged residues Arg and Lys, and their dipeptides

($p\sim10^{-16}$; Wilcoxon signed-rank test). We also observed that aliphatic residues Leu, Ile,

Ala, Val, were significantly more frequent in APRs compared to the rest of the protein

sequence ($p\sim10^{-16}$; Wilcoxon signed-rank test). The fraction of positively charged to

aliphatic residues was the strongest predictor of APRs compared to the fraction of all

other amino acids (Figure 1C; $p\sim10^{-21}$; Wilcoxon signed-rank test).

To find out whether these patterns are linked to differences in physicochemical

properties of amino acids, we used a random forest approach, a widely-used machine

learning technique for the classification of two or more data sets[30]. Specifically, we used

all 96 aggregators with experimentally characterized APRs[22], and subdivided each

protein into two sequence data sets, one comprising the APRs and one comprising the

non-APRs. For each of these datasets, we calculated a feature matrix that consisted of

500 amino acid properties (Dataset S3, and S4). We took these properties from the

AAindex database[31], which curates various physicochemical and biochemical properties

of amino acids. The classifier achieved an accuracy of ~93% in 100 independent runs,

with the data split into a training set (80% of the data) and a testing set (20%) (Figure 1D;

Dataset S5). Two amino acid properties were most important in this classification (Figure

1E). The first is a low positive charge of amino acids in APRs compared to non-APRs.

The second is a high amphiphilic propensity of these amino acids. The amphiphilic

propensity measures the preference of amino acids to occur at protein-solvent interfaces,

such as the end regions of transmembrane helices[32]. Amino acids like arginine and lysine

have the highest preference for these environments, and the largest value of the

amphiphilic index. In contrast, hydrophobic amino acids such as leucine, isoleucine,

valine, and alanine prefer internal protein environments and have the lowest amphiphilic

propensity[32].

Because of the statistically significant difference between the amphiphilic nature

of APRs and non-APRs, we argue that aggregator proteins resemble supramolecular

amphiphiles. These molecular structures have distinct hydrophobic and hydrophilic

parts[33]. In response to external stimuli such as temperature, pH, and ionic strength[33],

they can reversibly aggregate by non-covalent bonding forces, e.g., through electrostatic

and hydrophobic interactions.

We next investigated the relevance of amphiphilic aggregation to stress-induced

phase separation of our yeast aggregator proteins. An important feature of amphiphilic

molecules is that they are more likely to occur in an aggregated phase if their hydrophobic

chain is long[34]. We thus predicted that aggregator proteins with longer APRs will occur

preferentially in the aggregated phase. To test this prediction, we asked whether

aggregator proteins with long APRs preferentially occur in the pellet fraction of proteins

extracted from heat-stressed yeast cells[35]. Specifically, we used for this purpose the $\log_2$

ratio of the protein's abundance in the pellet fraction of yeast cells to that of supernatant

as a measure of protein's enrichment in the aggregated phase[35]. We found this

enrichment data for 31 aggregated proteins in our dataset. Indeed, aggregator proteins

with longer APRs preferentially accumulate in the aggregated pellet fraction (Figure 2A,

R = 0.53, p = 0.0021; Spearman's rank correlation). This association increases to R =

0.64 for aggregator proteins whose APRs have a low positive charge-to-aliphatic bias (p=

0.012; Spearman's rank correlation). To see whether this observation extends to other

stress-induced conditions, we also used the mass spectrometry dataset of nutrient-

starved yeast cells reported by Narayanasamy et al[36]. We identified 56 aggregator

proteins in this dataset and checked the association between APR lengths in these

proteins and the Z-score of proteins' enrichment in the pellet fraction. In nutrient-starved

cells too, the aggregator proteins with longer APRs preferentially accumulate in the

aggregated pellet fraction (Figures 2B, R = 0.47, p = 0.00014; Spearman's rank

correlation), and the association increased for the low positive charge-to-aliphatic bias (R

= 0.53, p = 0.0022; Spearman's rank correlation).

We then investigated the relationship between APR length in aggregator proteins

and their differential enrichment in the aggregated phase using equilibrium statistical

thermodynamics. We considered a two-state model where proteins exist either as a

soluble monomer or an insoluble aggregate. The fraction of proteins occurring in the

aggregate at the critical concentration (cc) for aggregation ($\mathcal{F}_{cc}$) is expressed as:

$$\mathcal{F}_{cc}(theory) = \frac{(P_{agg})_{cc}}{(P_{agg})_{cc}+(P_{mon})_{cc}} = \frac{1}{1+e^{-\beta\Delta G_{agg}}} \qquad \text{(Eq. 1)}$$

Here, P_{agg} and P_{mon} are the relative concentrations of proteins in the aggregated and the

monomer phase, $\beta = 1/k_B T$ where k_B is the Boltzmann's constant, and ΔG_{agg} is the free

energy change upon aggregation. Because the aggregation free energy increases in

proportion to the length of the hydrophobic chain[37], we rearranged Equation 1 to:

$$\mathcal{F}_{cc}(theory) = \frac{1}{1+e^{-\beta.B.L}} \qquad \text{(Eq. 2)}$$

Where B is the proportionality constant between the aggregation free energy and the APR

length. We then calculated the observed fraction of the aggregator proteins in the

aggregated phase from their reported $\log_2$ enrichment ratio in the pellet fraction of heat-

stressed yeast ($log_2 \frac{[Pellet]}{[Supernatant]}$) as:

$\mathcal{F}_{cc}(experiment) = (1 + 2^{-log_2 \frac{[Pellet]}{[Supernatant]}})^{-1}$ (Eq. 3)

Equation 2 predicted the observed relationship between APR length and a protein's

fraction in the aggregated phase well (Figure 2C, R=0.53, p=0.0021; Spearman's rank

correlation, see supplementary information for the fitted constants). Importantly, the

predictive power of Equation 2 was higher for proteins whose APRs had a low positive

charge-to-aliphatic bias (Figure 2C; R=0.64, p=0.012; Spearman's rank correlation).

In summary, our observations suggest that the aggregation-prone regions within

the aggregator proteins can readily unfold in response to stressors, form hydrophobic

long chains, and phase-separate in a manner akin to amphiphilic molecules. We also

observed that aggregator proteins with longer APRs are more prone to aggregation. This

relationship also suggests that aggregator proteins may leave the aggregated phase by

masking their APRs from self-assembly or from interaction with other proteins. Molecular

chaperones can facilitate this step: They have been co-purified with stress-induced

aggregated proteins and can solubilize them[39]. Altogether, our study demonstrates that

stress-induced phase separation is a sequence-encoded phenotype. It can be both

explained and predicted by the self-assembly and aggregation of amphiphilic molecules.

Methods:

We identified aggregation-prone regions in the 96 aggregator proteins from the dataset of Cappelletti et. al[22], and downloaded their sequences from the Uniprot database[40] (Dataset S6). We chose 500 amino acid properties from the AAindex database to distinguish the physicochemical properties of APRs from non-APR segments in the aggregator protein[31].

We used a binary classification to group regions within an aggregator protein into aggregation-prone regions (class A) and the remainder of the protein (class B). More specifically, we used a random forest classification, splitting aggregator proteins into a random subset comprising 80% of aggregator proteins for training and 20% for testing. To build features for the classification, we calculated the average value of 500 physicochemical properties for each sequence in the APR and non-APR dataset. This yielded two feature matrices for the APR and non-APR sequences. To apply random forest classification, we used the randomForest package in R[42], and evaluated the best number of trees (*nTree*) and the number of variables randomly sampled at each split (*mtry*), in the random forest algorithm. To do so, we systematically varied the *nTree* and *mtry*, and calculated the accuracy of classification with 10-fold cross-validation and 3 repeats. We defined accuracy as the percentage of correctly identified classes of proteins (APRs and non-APRs) out of all instances. The combination of *nTree*=5000 trees and *mtry*=10 variables achieved the highest accuracy of ~ 90%. We then used these parameters to perform 100 random forest clusterings, in which we randomly assigned proteins to the training and the testing datasets. To quantify the accuracy of classification

we counted the number of true positive and false positive predictions and calculated the

area under the curve (AUC). These values are shown as a receiver operating

characteristic curve (ROC) in Figure 2D. The most important physicochemical properties

were the ones whose Gini index in the classification decreased the most compared to all

other properties. This index is a widely-used measure of dispersion that reflects inequality

in the values of a frequency distribution. Within the random forest framework, this index

is calculated as the average probability that each of 500 physicochemical properties

wrongly classifies APRs and non-APRs. We performed all models and statistical analyses

using R. Scripts are available at: https://github.com/dasmeh/yeast_aggregators

References

1.   Aguzzi, A. & Calella, A.M. Prions: protein aggregation and infectious diseases.
Physiological reviews **89**, 1105-1152 (2009).
2.   Aguzzi, A. & O'connor, T. Protein aggregation diseases: pathogenicity and therapeutic
perspectives. *Nature reviews Drug discovery* **9**, 237-248 (2010).
3.   Ross, C.A. & Poirier, M.A. Protein aggregation and neurodegenerative disease. *Nature*
medicine **10**, S10-S17 (2004).
4.   Scheibel, T. & Buchner, J. Protein aggregation as a cause for disease. *Molecular*
chaperones in health and disease, 199-219 (2006).
5.   Shulman, J.M., De Jager, P.L. & Feany, M.B. Parkinson's disease: genetics and
pathogenesis. *Annual Review of Pathology: Mechanisms of Disease* **6**, 193-222 (2011).
6.   O'Connell, J.D. *et al.* A proteomic survey of widespread protein aggregation in yeast.
Molecular BioSystems **10**, 851-861 (2014).
7.   Wallace, E.W. *et al.* Reversible, specific, active aggregates of endogenous proteins
assemble upon heat stress. *Cell* **162**, 1286-1298 (2015).
8.   Saad, S. *et al.* Reversible protein aggregation is a protective mechanism to ensure cell
cycle restart after stress. *Nature cell biology* **19**, 1202-1213 (2017).
9.   Cereghetti, G., Saad, S., Dechant, R. & Peter, M. Reversible, functional amyloids: towards
an understanding of their regulation in yeast and humans. *Cell Cycle* **17**, 1545-1558
(2018).
10.   Hyman, A.A., Weber, C.A. & Jülicher, F. Liquid-liquid phase separation in biology. *Annual*
review of cell and developmental biology **30**, 39-58 (2014).

11. Hnisz, D., Shrinivas, K., Young, R.A., Chakraborty, A.K. & Sharp, P.A. A phase separation
model for transcriptional control. *Cell* **169**, 13-23 (2017).
12. Erdel, F. & Rippe, K. Formation of chromatin subcompartments by phase separation.
Biophysical journal **114**, 2262-2270 (2018).
13. Valsecchi, C.I.K. *et al.* RNA nucleation by MSL2 induces selective X chromosome
compartmentalization. *Nature* **589**, 137-142 (2021).
14. Youn, J.-Y. *et al.* Properties of Stress Granule and P-Body Proteomes. *Molecular Cell* **76**,
286-294 (2019).
15. Uversky, V.N. Intrinsically disordered proteins in overcrowded milieu: membrane-less
organelles, phase separation, and intrinsic disorder. *Current opinion in structural biology*
44, 18-30 (2017).
16. Posey, A.E., Holehouse, A.S. & Pappu, R.V. Phase separation of intrinsically disordered
proteins. in *Methods in Enzymology*, Vol. 611 1-30 (Elsevier, 2018).
17. Kuechler, E.R., Budzyńska, P.M., Bernardini, J.P., Gsponer, J. & Mayor, T. Distinct
features of stress granule proteins predict localization in Membraneless organelles.
Journal of molecular biology **432**, 2349-2368 (2020).
18. Riback, J.A. *et al.* Stress-triggered phase separation is an adaptive, evolutionarily tuned
response. *Cell* **168**, 1028-1040. e19 (2017).
19. Iserman, C. *et al.* Condensation of Ded1p Promotes a Translational Switch from
Housekeeping to Stress Protein Production. *Cell* (2020).
20. Gao, X. *et al.* High-throughput limited proteolysis/mass spectrometry for protein
domain elucidation. *Journal of structural and functional genomics* **6**, 129-134 (2005).
21. Schopper, S. *et al.* Measuring protein structural changes on a proteome-wide scale using
limited proteolysis-coupled mass spectrometry. *Nature protocols* **12**, 2391 (2017).
22. Cappelletti, V. *et al.* Dynamic 3D proteomes reveal protein functional alterations at high
resolution in situ. *Cell* **184**, 545-559. e22 (2021).
23. Dosztányi, Z. Prediction of protein disorder based on IUPred. *Protein Science* **27**, 331-
340 (2018).
24. Bonferroni, C.E., Bonferroni, C. & Bonferroni, C. Teoria statistica delle classi e calcolo
delle probabilita'. (1936).
25. Sugimoto, K., Matsumoto, K., Kornberg, R.D., Reed, S.I. & Wittenberg, C. Dosage
suppressors of the dominant G1 cyclin mutant CLN3-2: identification of a yeast gene
encoding a putative RNA/ssDNA binding protein. *Molecular and General Genetics MGG*
248, 712-718 (1995).
26. Altmeyer, M. *et al.* Liquid demixing of intrinsically disordered proteins is seeded by poly
(ADP-ribose). *Nature communications* **6**, 8088 (2015).
27. Jamieson, D.J., Rahe, B., Pringle, J. & Beggs, J.D. A suppressor of a yeast splicing
mutation (prp8-l) encodes a putative ATP-dependent RNA helicase. *Nature* **349**, 715-717
(1991).
28. Schäfer, I.B. *et al.* Molecular basis for poly (A) RNP architecture and recognition by the
Pan2-Pan3 deadenylase. *Cell* **177**, 1619-1631. e21 (2019).
29. Kelil, A., Dubreuil, B., Levy, E.D. & Michnick, S.W. Exhaustive search of linear information
encoding protein-peptide recognition. *PLoS computational biology* **13**, e1005499 (2017).

30. Ho, T.K. Random decision forests. in *Proceedings of 3rd international conference on*
document analysis and recognition Vol. 1 278-282 (IEEE, 1995).
31. Kawashima, S. & Kanehisa, M. AAindex: amino acid index database. *Nucleic acids*
research **28**, 374-374 (2000).
32. Mitaku, S., Hirokawa, T. & Tsuji, T. Amphiphilicity index of polar amino acids as an aid in
the characterization of amino acid preference at membrane–water interfaces.
Bioinformatics **18**, 608-616 (2002).
33. Zhang, X. & Wang, C. Supramolecular amphiphiles. *Chemical Society Reviews* **40**, 94-101
(2011).
34. Shaul, B. & Gelbart, W. Theory of chain packing in amphiphilic aggregates. *Annual*
Review of Physical Chemistry **36**, 179-211 (1985).
35. Zhu, M. *et al.* Proteomic analysis reveals the direct recruitment of intrinsically
disordered regions to stress granules in S. cerevisiae. *Journal of Cell Science* **133**(2020).
36. Narayanaswamy, R. *et al.* Widespread reorganization of metabolic enzymes into
reversible assemblies upon nutrient starvation. *Proceedings of the National Academy of*
Sciences **106**, 10147-10152 (2009).
37. Budin, I., Prywes, N., Zhang, N. & Szostak, J.W. Chain-length heterogeneity allows for the
assembly of fatty acid vesicles in dilute solutions. *Biophysical journal* **107**, 1582-1590
(2014).
38. Iserman, C. *et al.* Condensation of Ded1p promotes a translational switch from
housekeeping to stress protein production. *Cell* **181**, 818-831. e19 (2020).
39. Cherkasov, V. *et al.* Coordination of translational control and protein homeostasis during
severe heat stress. *Current biology* **23**, 2452-2462 (2013).
40. Consortium, U. UniProt: a hub for protein information. *Nucleic acids research* **43**, D204-
D212 (2015).
41. Kelley, L.A., Mezulis, S., Yates, C.M., Wass, M.N. & Sternberg, M.J. The Phyre2 web
portal for protein modeling, prediction and analysis. *Nature protocols* **10**, 845-858
(2015).
42. Liaw, A. & Wiener, M. Classification and regression by randomForest. *R news* **2**, 18-22
(2002).
43. Raudvere, U. *et al.* g: Profiler: a web server for functional enrichment analysis and
conversions of gene lists (2019 update). *Nucleic acids research* **47**, W191-W198 (2019).
293
294

295 **Figure captions:**
296
Figure 1. Sequence composition and physicochemical properties of aggregation-
prone regions in the yeast aggregators. A) The percentage of explained variance for
10 dimensions from a principal component analysis on the frequencies of 20 amino acids
in 270 aggregation-prone regions of 96 yeast proteins. B) Biplot for the first and the
second principal components. Grey circles, labeled by numbers from 1 to 270, represent
aggregation-prone regions. Red arrows represent the weight of different amino acid
frequencies. C) The amino acid frequencies in APRs (shown in red) versus the rest of the
protein sequence (shown in green) in 96 yeast aggregators. D) The receiver-operating
characteristic (ROC) curves for the clustering of APRs from non-APRs using random
forest classification. ROC curves in grey are from 100 random forest clustering and their
average is shown in black. E) Ranked importance of physicochemical vartiables
quantified as the average decrease in the Gini index for different physicochemical
properties.
310
Figure 2. Aggregator proteins reversibly aggregate in a manner akin to amphiphilic
molecules.
A) The log2 ratio of the protein's abundance in the pellet fraction to that of supernatant in
heat-stressed yeast cells (30 out of 96 aggregator proteins[35]) versus their APR length. B)
The aggregator proteins' enrichment in the pellet fraction upon nutrient starvation (59 out
of 96 proteins[35]) versus their APR length in log-scale. The red and the grey circles in both
panels A and B represent all aggregator proteins, and those where the positive charge-

to-aliphatic-bias in APRs was lower than its median value $\sim$ 0.25. C) The fraction of

aggregator proteins in the aggregated phase calculated from Equation 3 versus the length

of APRs. D) The fraction of aggregator proteins with a low positive charge-to-aliphatic-

bias in their APRs versus the length of APRs calculated from Equation 3. For both panels

C and D, the fitted curve to the observed fraction of proteins in the aggregated phase

using Equation 2 is shown in red. R values are the Spearman's rank correlation between

the expected (Equation 2) and observed (Equation 3) fractions of proteins in the

aggregated phase.

Figure 1

Figure 2

Figure 1

Figure 2