Systems analysis of gut microbiome influence on metabolic disease in HIV and high-risk populations

Abigail J.S. Armstrong[1,2,3], Kevin Quinn[4], Jennifer Fouquier[1], Sam X. Li[1], Jennifer M. Schneider[1], Nichole M. Nusbacher[1], Katrina A. Doenges[4], Suzanne Fiorillo[1], Tyson J. Marden[5], Janine Higgins[6], Nichole Reisdorph[4], Thomas B. Campbell[1], Brent E. Palmer[1#], Catherine A. Lozupone[1#]

Affiliations:

1. Department of Medicine, University of Colorado Denver, Anschutz Medical Campus, Aurora, CO, USA

2. Department of Immunology and Microbiology, University of Colorado Denver, Anschutz Medical Campus, Aurora, CO, USA

3. Center for Advanced Biotechnology and Medicine, Rutgers the State University, Piscataway, NJ, USA

4. Skaggs School of Pharmacy and Pharmaceutical Sciences, University of Colorado Anschutz Medical Campus, 80045 CO, Aurora, USA

5. Colorado Clinical and Translational Sciences Institute; University of Colorado Anschutz Medical Campus, Aurora, Colorado, 80045. USA

6. Department of Pediatrics, Section of Endocrinology, University of Colorado Anschutz Medical Campus. Aurora, CO

Corresponding authors

Brent E. Palmer: 303-724-7203; brent.palmer@cuanschutz.edu

Catherine A. Lozupone: 303-724-7942; catherine.lozupone@cuanschutz.edu

Abstract

Poor metabolic health, characterized by insulin resistance and dyslipidemia, is higher in people

living with HIV and has been linked with inflammation, anti-retroviral therapy (ART) drugs, and

ART-associated lipodystrophy (LD). Metabolic disease is associated with gut microbiome

composition outside the context of HIV but has not been deeply explored in HIV infection nor in

high-risk men who have sex with men (HR-MSM), who have a highly altered gut microbiome

composition. Furthermore, the contribution of increased bacterial translocation and associated

systemic inflammation that has been described in HIV-positive and HR-MSM individuals has not

been explored. We used a multi-omic approach to explore relationships between impaired

metabolic health, defined using fasting blood markers, gut microbes, immune phenotypes and

diet. Our cohort included ART-treated HIV positive MSM with and without LD, untreated HIV

positive MSM, and HR-MSM. For HIV positive MSM on ART, we further explored associations

with the plasma metabolome. We found that elevated plasma lipopolysaccharide binding protein

(LBP) was the most important predictor of impaired metabolic health and network analysis

showed that LBP formed a hub joining correlated microbial and immune predictors of metabolic

disease. Taken together, our results suggest the role of inflammatory processes linked with

bacterial translocation and interaction with the gut microbiome in metabolic disease among HIV

positive and negative MSM.

Importance Statement

The gut microbiome in people living with HIV (PLWH) is of interest as chronic infection often

results in long term comorbidities. Metabolic disease is prevalent in PLWH even in well-

controlled infection and has been linked with the gut microbiome in previous studies, but little

attention has been given to PLWH. Furthermore, integrated analyses that consider gut

microbiome together with diet, systemic immune activation, metabolites, and demographics

have been lacking. In a systems-level analysis of predictors of metabolic disease in PLWH and

men who are at high risk of acquiring HIV, we found that increased LBP, an inflammatory

marker indicative of compromised intestinal barrier function, was associated with worse

metabolic health. We also found impaired metabolic health associated with specific dietary

components, gut microbes, and host and microbial metabolites. This work lays the framework

for mechanistic studies aimed at targeting the microbiome to prevent or treat metabolic

endotoxemia in HIV-infected individuals.

Keywords

HIV; microbiome; men who have sex with men (MSM); metabolic disease; endotoxemia

Background

Poor metabolic health characterized by insulin resistance and dyslipidemia is frequent in people

living with HIV (PLWH) (1-3) and has been linked with chronic inflammation (4-7) and several

anti-retroviral therapy (ART) drugs (8). Metabolic disease is particularly prevalent in HIV-positive

individuals with lipodystrophy (LD), a disease linked with early ART drugs that is manifested by

lipoatrophy in the face, extremities, and buttocks with or without visceral fat accumulation.

Delineating pathophysiological mechanisms of impaired metabolic health is crucial for tailoring

strategies for prophylaxis and treatment to PLWH.

Metabolic disease has been linked with gut microbiome structure and function outside the

context of HIV infection (9-13), but this relationship has not been explored deeply in PLWH. We

and others have found an altered gut microbiome composition in both PLWH (14-16) and men

who have sex with men at high-risk of contracting HIV (HR-MSM) (15, 17). Furthermore, we

have demonstrated that the altered microbiome in HIV (14) and HR-MSM (14, 18) are pro-

inflammatory *in vitro* and/or in gnotobiotic mice (14, 18). This is of interest as peripheral

inflammatory signals have been implicated in both cardiovascular disease risk (7, 19) and

insulin sensitivity (4, 5, 20-22) in PLWH. A recent study conducted in PLWH found that HIV-

associated microbiome differences correlated with risk of metabolic syndrome, particularly in

individuals with a history of severe immunodeficiency (23).

One mechanism by which the gut microbiome may impact MS in PLWH is through effects on

intestinal barrier function. Impaired intestinal barrier, measured by increased bacterial products

such as lipopolysaccharide (LPS; endotoxin) in blood, has been linked with MS and particular

metabolic derangements (e.g., dyslipidemia and insulin resistance). This "metabolic

endotoxemia" has been described in chronic kidney disease/ hemodialysis patients(24) and

there are mixed data regarding a role in obesity(25-27). Murine studies have further supported

that an altered gut microbiome and translocation of LPS could trigger insulin resistance,

diabetes, and atherosclerosis(28-31). This is of interest for PLWH because bacterial

translocation, indicated by higher levels of LPS or LPS-binding protein (LBP) in blood, is well

known to occur in HIV infected populations, to be not completely ameliorated by antiretroviral

therapy (ART)(32, 33), and to positively correlate with HIV-associated gut microbiome

differences(23). Increased plasma LPS levels have also been observed in MSM and linked with

recent sexual behavior(34).

We hypothesized that PLWH and HR-MSM with poor metabolic health would harbor a distinct

gut microbial signature that was in turn also associated with elevated peripheral immune

activation. We evaluated this relationship while considering other factors known to influence the

microbiome, immunity and metabolic health. This analysis included typical diet; HIV, ART, and

LD status; and other demographic characteristics such as age and body mass index (BMI). For

HIV-positive individuals on ART with and without LD, we further explored associations with the

plasma metabolome (Figure 1). Our results suggest a central role of inflammatory processes

linked with

bacterial translocation as measured by LBP, and co-correlated intestinal microbes, dietary and

demographic attributes in metabolic disease risk.

Results

Study Population

This study examined a cohort of 113 men, including men who have sex with women (MSW;

n=22, 19.5%) and MSM (n=91, 80.5%) (Table 1). Of the MSM, 32 were HIV-negative (35.2%),

14 were HIV-positive and not on ART (15.4%), and 45 were HIV-positive and on ART (49.4%).

The HIV-positive, treated group included those with lipodystrophy (LD; n=25, 55.6%) and those

without (n=20, 44.4%). The HIV-negative MSM participated in activities that put them at high

risk of contracting HIV including: 1) a history of unprotected anal intercourse with one or more

male or male-to-female transgender partners; 2) anal intercourse with two or more male or

male-to-female transgender partners; or 3) being in a sexual relationship with a person who has

been diagnosed with HIV (35). In order to focus on HIV-associated metabolic disease, obese

individuals (BMI >30) were excluded. There was no significant difference in BMI between the

cohorts (Kruskal-Wallis test, p = 0.085, Supplemental Table 1). Individuals in the HIV-positive,

treated cohorts were significantly older than HIV-negative and HIV-positive, untreated MSM

(Kruskal-Wallis test, p < 0.001). Age matching across all cohorts was not feasible in part

because LD is associated with early-generation ART drugs and thus most common in older HIV-

positive individuals and HR-MSM behavior and new HIV infections are predominantly in younger

individuals. However, age is carefully considered in downstream analyses. All treated, HIV-

positive individuals were on successful ART with suppressed viral loads (Table 1).

<u>Metabolic Disease Score as a Marker for Metabolic Health</u>

We measured seven common clinical markers of metabolic health from fasting blood:

triglycerides, glucose, insulin, LDL, HDL, leptin, and adiponectin. Since these markers are often

correlated with each other, we used principal component analysis (PCA) to define a single

continuous measure of overall metabolic health as has been done previously (Figure 2A) (36,

37). Individuals with high values along the first principal component (PC1) generally had high

triglycerides and low HDL, indicating dyslipidemia, and higher levels of fasting blood glucose

and insulin, indicating insulin resistance (Figure 2A, Supplemental Figure 1). Values of PC1

were shifted to a minimum of one and log transformed to define the metabolic disease score,

which ranged from 0 as healthy and 2.5 as impaired. We used regression analysis to determine

how this score related to clinically defined cutoffs for normal levels of the markers

(Supplemental Figure 1). For example, triglycerides positively correlated with metabolic disease

score and almost all individuals with a score above 1.45 had triglyceride levels in the unhealthy

range of greater than 200 mg/dL. Similar patterns and cutoffs were true for HDL, LDL, and

glucose (Supplemental Figure 1). The intersect of the regression with these cutoffs were all

averaged to a single number of 1.4. Individuals below the cutoff were categorized as

metabolically normal and those above were categorized as metabolically impaired.

When comparing the metabolic disease score across cohorts, we found that ART-treated, HIV-

positive individuals with LD trended higher in both the average metabolic disease score and the

proportion of individuals with scores in the metabolically impaired group but intergroup

significance was lost after multiple test corrections (Figure 2B). Furthermore, because our HIV-

positive, treated cohorts were significantly older than our HIV-negative MSM and HIV-positive,

untreated MSM, we used a linear model to explore differences in the metabolic disease score

across cohorts while accounting for age (Figure 2C). This score was positively associated with

age only in HIV-negative MSM and HIV-positive, untreated MSM (Figure 2C; linear model; p <

0.001 and p = 0.036 respectively) and only HIV-negative MSM had significantly higher

metabolic disease score compared to HIV-negative MSW when accounting for age (linear

model; p < 0.001).

<u>Selection of Features that Predict the Metabolic Disease Score and Interactions Between</u>

<u>Selected Features</u>

We next explored the complex relationships of the gut microbiome, peripheral immune

activation, and diet to the metabolic disease score and to each other using only data from the

HIV positive and HR-MSM cohorts. We first selected features that were important predictors of

the metabolic disease score using the VSURF (Variable Selection Using Random Forest) tool

(38). VSURF is optimized for feature selection, returning all features that are highly predictive of

the response variable, even when a smaller subset of highly predictive variables with redundant

features removed could be just as accurate for prediction (38). We input the following features

into the VSURF tool: 1) 130 microbial features. 99% identity Operational Taxonomic Units

(OTUs) with highly co-correlated OTUs were binned into modules as described in the methods

(detailed in Supplemental Table 2). Only OTUs present in >20% of samples were included. 2)

21 immune features that were measured in plasma using multiplex ELISA (detailed in

Supplemental Table 2). These immune measures were selected based on a literature search for

those previously shown to be altered in HIV infection and/or metabolic disease. 3) 21

clinical/demographic features such as age, BMI, HIV infection and treatment status, and typical

gastrointestinal symptoms including constipation, diarrhea and bloating (detailed in

Supplemental Table 2). 4) 29 dietary features that were collected using a food frequency

questionnaire of typical dietary intake over the prior year as detailed in the methods. Highly co-

correlated diet features were also binned into modules (detailed in Supplemental Table 2).

From the initial 201 measures, VSURF identified 69 important variables (4 clinical data

measures, 6 diet measures, 14 immune measures, and 45 microbes) and a subset of ten highly

predictive variables (Supplemental Table 3). These 69 features could predict the metabolic

disease score using traditional random forest with an r^2 of 31.05%, which is significantly better

than a null model where the outcome was randomly permuted (p = 0.036, Supplemental Figure

2).

We found that 21 of the 69 selected variables were positively or negatively correlated with the

metabolic disease score (Spearman rank correlation, FDR p < 0.1, Supplemental Table 3).

Since random forest can detect non-linear relationships and/or features that are only important

when also considering another feature, it is not surprising that all features were not correlated

linearly with the metabolic disease score. All VSURF selected clinical measures were positively

correlated with metabolic disease score and included age, BMI, lipodystrophy, and bloating

(Supplemental Table 3). None of the six selected diet measures correlated with metabolic

disease score (Supplemental Table 3). VSURF selected immune markers that were positively

correlated with the metabolic disease score included LBP, intercellular adhesion molecule 1

(ICAM-1), interleukin (IL) 16, IL-12, and granulocyte-macrophage colony-stimulating factor (GM-

CSF) (Supplemental Table 3). The feature with the highest random forest importance score was

LBP.

Diet, the microbiome, and immune phenotypes can all influence each other and can also relate

to clinical/demographic factors such as BMI and age (Figure 1). For this reason, we also

investigated the relationship between the 69 important factors using pairwise Spearman rank

correlation and network visualization (Figure 3, Supplemental Figure 3, Supplemental Table 3).

The selected important microbes included many that were highly correlated with each other and

with dietary, clinical/demographic, and inflammatory phenotypes (Supplemental Figure 3A,

Supplemental Table 3). For example, a module of bacteria identified within the *Prevotella* genus

and the *Paraprevotellaceae* family, negatively associated with metabolic disease score and

positively associated with dietary fiber (Supplemental Figure 3B).

Because bacterial translocation is known to occur at increased levels in both HIV-positive

individuals (39) and HR-MSM (34), we were specifically interested in investigating which of the

selected features correlated with LBP. LBP was negatively correlated with several putative

butyrate producing bacteria/bacterial modules such as OTUs in the genera *Coprococcus* (40,

41)(Figure 3B). LBP was also positively correlated with *Dorea* species (Figure 3B; Supplemental

Table 3). In addition to correlations, we evaluated interactions between selected variables using

the tool iRF (iterative Random Forest)(42). These interactions represent variables that are in

adjacent nodes in a random forest tree in which the value of one influences the predictability of

the other. This analysis also identified that LBP interacted with age, BMI, an OTU in the

Lachnospiraceae family, and microbiome module 24 (*Coprococcus* sp. and *Blautia* sp.) in

predictions of the metabolic disease score (Figure 3C; Supplemental Table 3).

Because of the recognized role of inflammation in an increased risk of age-associated non-

AIDS-related morbidity and mortality in PLWH(43-46), we were also specifically interested in

investigating which of the selected features correlated with age. Age had the strongest positive

association with lipodystrophy and also positively correlated with several inflammatory markers

including LBP, CRP, IL-6, ICAM-1 and SAA (Supplemental Figure 3C, Supplemental Table 2).

Age was also correlated with several different gut microbes including a negative correlation with

OTUs assigned to *Bifidobacterium adolescentis*, *Eubacterium dolichum, Coprobacillus sp.*, and

Oscillospira sp. (Supplemental Figure 3C).

Random Forest analyses do not allow for gaps in data, and, as a result, HIV-specific variables

were omitted from these analyses. We thus performed Spearman's rank correlations to

investigate a relationship between variables only measured in our HIV positive cohorts and the

metabolic disease score and found no correlation with CD4 nadir (p=0.28) and CD4+ T-cell

count (p=0.076). We also found no significant correlation between viral load and metabolic

disease score among individuals with untreated HIV infection (p = 0.68).

Lastly, because many gut microbes were associated with metabolic disease score, we also

investigated associations with overall microbiome composition measures. Microbiome

evenness, as measured by Pielou's Evenness and Shannon Index were both weakly negatively

correlated (Spearman Rank Correlation, rho=-0.09; p = 0.047 and rho=-0.09; p = 0.045

respectively) and there was no significant relationship in measures of microbiome richness

(Faith's PD and Observed Features). Additionally, there was no significant relationship between

weighted and unweighted UniFrac distances and metabolic disease score differences between

individuals (Mantel Test, p = 0.7 and p = 0.4 respectively).

The Relationship Between The Plasma Metabolome and the Metabolic Disease Score in ART-

treated HIV-positive Individuals With and Without LD

To pursue a further mechanistic understanding of how the gut microbiome may influence the

metabolic disease score in PLWH, we performed untargeted metabolomics (LC/MS) on plasma

from our cohort of ART-treated, HIV-positive individuals with and without LD (n=44). Metabolite

identities were then validated using untargeted MS/MS. We used two approaches to determine

which plasma metabolites were either directly produced or indirectly influenced by the gut

microbiome. First, metabolites were run through the computational tool AMON (47), which uses

the KEGG database (48) and inferred metagenomes (calculated using PICRUSt2 (49)) to

determine which could have been produced by the microbiome. Second, LC/MS was run on

plasma from both germ free (GF) and humanized mice to determine metabolites that had

significantly altered levels upon colonization with human microbiomes. Specifically, GF mice

were gavaged using fecal samples from eight men from the study cohort (humanized mice)

while two mice were gavaged using PBS as control (Supplemental Table 4). Plasma was

collected before and after gavage. All mice were fed a high-fat western diet.

We found that 820 metabolites were different in abundance between GF and humanized mice

after multiple test corrections (Student's t-test, FDR p < 0.05), 493 of which were also present in

the human plasma samples (Figure 4). From the full set of 5,332 metabolites identified in the

human plasma, 416 could be annotated with KEGG IDs. These were further analyzed using

AMON. 146 microbiome-associated metabolites were identified that are putatively produced by

the gut microbiome; however, many of these could also be produced by the host. Twenty-six of

the 146 microbiome-associated metabolites identified by AMON also differed in colonized

versus GF mice (Figure 4, Supplemental Table 5).

Of the 5,332 total measured metabolites in the human samples, 150 correlated with the

metabolic disease score (Spearman rank correlation, FDR p < 0.05; Supplemental Table 5).

The correlated compounds were enriched in a number of different metabolic pathways with both

the Phospholipid and the Glycerolipid pathways of the Small Molecule Pathway Database

(SMDPH)(50) highly enriched (Supplemental Table 5). Consistent with the metabolic disease

score being defined in part by dyslipidemia, 17 of the significant compounds were annotated as

triglycerides. Seven of the significant compounds were associated with the microbiome as

determined by the process outlined above. We confirmed the identity of 5 of the 7 of these with

MS/MS (Supplemental Table 5). Of these seven microbiome-associated metabolites, two could

exclusively be explained by direct production by the microbiome. Specifically, dehydroalanine,

identified as a microbial product by AMON, negatively correlated with the metabolic disease

score and bacteriohopane-32,33,34,35-tetrol positively correlated with the metabolic disease

score (Supplemental Table 5). Two additional microbiome-associated compounds were

triglycerides (TG(54:6) and TG (16:0/18:2.20:4)) that were positively correlated with metabolic

disease score and elevated in humanized mice compared to GF mice. Another of these

metabolites, 1-Linleoyl-2-oleoyl-rac-glycerol, is a 1,2-diglyceride in the triglyceride biosynthesis

pathway. Finally, phosphatidylcholine (PC(17:0/18:2)) and phosphatidylethanolamine

(PE(20:3/18:0)) compounds were identified as microbiome associated and positively and

negatively correlated with the metabolic disease score respectively (51).

Since age was significantly associated with metabolic disease score in the full cohort, and thus

could present a potential confounding factor in the metabolite relationships, we constructed a

model that included age using a rank-based estimation for linear models using the R package

Rfit (52). Because of the statistical differences between linear regression and spearman

correlation, we looked at Rfit with and without age. Both analyses identified 60 metabolites

significant after FDR correction (Supplemental Table 5) and 45 of the 60 metabolites were

identified with both Rfit models, indicating that age was not very influential on the results

(Supplemental Figure 4).

Discussion

In this study, we identified gut microbes, dietary components, demographic and immune

measures that predicted impaired metabolic health in a cohort of MSM with and without HIV,

ART, and LD. Notably, we identified a strong relationship with circulating LBP which in turn

correlated with other markers of systemic inflammation, a loss of beneficial microbes such as

butyrate-producing bacteria, and higher BMI, indicating that diverse modifiable factors may

influence LPS/inflammation driven metabolic disease in this population.

There was a positive association between impaired metabolic health and age, as has been

reported previously for both HIV (53) and non-HIV populations (54), but linear modeling

suggested that this relationship was driven by an association in HIV-negative MSM and HIV-

positive untreated MSM in our study, revealing a possibly larger effect size than in our other

cohorts. Also, when controlling for age, HIV-negative MSM had the highest metabolic disease

score, even compared to HIV positive individuals on ART with LD, a population that has

previously been reported to have higher incidence (55). This result is intriguing given prior

research linking increased levels of LPS in blood with high-risk behavior in MSM (34). Larger

cohorts and more detailed behavior information are required, however, to make any definitive

claims on impaired metabolic health in aging in HR-MSM.

Our finding that age was a significant predictor of impaired metabolic health in this cohort is

consistent with previous studies showing an increased risk of age-associated non-AIDS-related

morbidity and mortality in PLWH (44-46, 56). Consistent with the previously reported role of

inflammation in this relationship, age was positively correlated with 5 of the 14 immune

measures that VSURF selected as important predictors of the metabolic disease score,

including LBP, CRP, IL-6, ICAM-1 and SAA. Consistent with age being associated with changes

in the microbiome previously (57-59), we also found age to be correlated with several different

predictive microbes (Supplemental Figure 3C, Supplemental Table 2). These included a

negative association with *B. adolescentis*, whose loss has previously been mechanistically

linked with health deficits that occur with aging(60-63). *B. adolescentis* has been shown to

prevent immunosenescence when fed probiotically to aged rats (62), and to have beneficial

effects on barrier function in a murine model(64).

Consistent with prior studies that have associated high BMI with dyslipidemia, insulin resistance,

and/or metabolic syndrome (65-67), BMI was a positive predictor of impaired metabolic health in

our cohort even though our study excluded obese individuals, but did include overweight. This

suggests the importance of weight management even among overweight, non-obese individuals

as a strategy for reducing metabolic health impairment in this population.

We did not find a positive association between ART status and the metabolic disease score, but

this may be because study participants were on a wide variety of drug combinations with the

potential to have varied/contrasting effects. For instance, both integrase stand transfer inhibitors

(ISTI) (68) and regimens including the nucleoside reverse transcriptase inhibitor (NRTI)

tenofovir have been shown to increase risk of weight gain (69). Conversely, the CCR5

antagonist, maraviroc, may confer a benefit to cardiovascular function and body weight

maintenance and evidence in mice suggests that these beneficial effects may be linked to gut

microbiome composition changes (20, 70). Thus, future studies will be required to understand

factors important in particular drug contexts. Other HIV-specific measures such as CD4 nadir,

viral load and CD4+ T cell count also did not have a significant association with metabolic

disease score. Low CD4+ T cell count (71) and CD4/CD8 T cell ratio (72) has previously been

linked with poor metabolic health in individuals on ART. Furthermore, low CD4 nadir has been

linked with gut microbiome dysbiosis in HIV-infected individuals (71) and an association

between HIV-associated gut microbiome dysbiosis and metabolic syndrome was significantly

stronger in individuals with past severe immunodeficiency compared to those without (23). Our

negative results may be a result of small sample size and also a lack of individuals included in

the study with severely compromised CD4+ T cell counts (Table 1).

Several dietary components that we identified have been previously associated with metabolic

health, including dietary carotenoid, lycopene, and fiber (73-77). Fiber's benefit in glucose

response has been linked with the activity of *Prevotella copri*. Specifically, individuals who had

improved glucose response after 3 days of high-fiber consumption had a greater increase in *P.*

copri, and these beneficial effects were confirmed in a mouse model (76). However, another

study found that *P. copri* actually promoted poor glucose response in the context of a western

diet low in fiber through the production of branched chain amino acids (BCAAs)(11). Interactions

between *Prevotella*, dietary fiber, and metabolic health are of particular interest in this cohort

since HIV positive and negative MSM have much higher *Prevotella*, including *P. copri*, than non-

MSM (15, 17). Additionally, our prior study using *in vitro* stimulations of human immune cells

with fecal bacteria of HIV positive and negative MSM indicated that the Prevotella-rich

microbiomes of MSM could drive systemic inflammation (14). Interestingly, in our data, a

module of three OTUs, two in the genus *Prevotella* and one in the family *Paraprevotellaceae,*

negatively associated with metabolic disease score and positively associated with dietary fiber

(Supplemental Figure 3B), supporting a relationship between *Prevotella* and dietary fiber in

improving metabolic health, and not supporting deleterious effects. Further work will be needed

to decompose the complex relationship between dietary fiber, particular *Prevotella* strains, and

metabolic health in HIV positive and negative MSM with unique Prevotella-dominated

communities.

LBP was the most important feature in the random forest analysis and also a highly interactive

measure in the iRF analysis. LBP binds to both microbial LPS and lipoteichoic acid (LTA) (78)

and the presence of elevated LBP in blood is indicative of increased intestinal barrier

permeability (79). LBP was correlated with age and BMI, a relationship that was previously

observed in a cohort of HIV-negative men of African ancestry with this trio being further

associated with adiposity and pre-diabetes (80). LBP levels were also correlated with other

inflammatory markers that have been linked with worse metabolic health suggesting a role as a

central mediator of metabolic-disease associated immune phenotypes. These included 1) I-

CAM 1, whose expression in adipose tissue has been associated with diet-induced obesity in

mice (81) and metabolic syndrome in humans (82) 2) IL-6, a pro-inflammatory cytokine that has

been shown to play a direct role in insulin resistance (83), and 3) SAA, which is regulated in part

by IL-6 and plays a role in cholesterol metabolism (82); SSA3 specifically has been shown to be

produced in response to gut bacteria in obesity in mice (84). We observed a positive association

between the metabolic disease score and frequency of abdominal bloating (Supplemental Table

3, Supplemental Figure 3A), further supporting a role of intestinal dysfunction in this population.

Taken together these associations suggest that inflammation originating from an impaired

intestinal barrier is promoting worse metabolic health.

An importance of microbiota-driven intestinal barrier dysfunction in HIV-associated metabolic

syndrome was suggested in a recent study of Gelpi *et al.,* which found that metabolic syndrome

in HIV-infected individuals was correlated with an HIV-associated gut microbiota (23). This

microbiota was characterized in part by a decrease in butyrate-producing bacteria, including

Coprococcus and *Butyrivibrio.* Consistent with this finding, we found that multiple *Coprococcus*

OTUs were selected as important predictors of the metabolic disease score and that they also

negatively correlated with LBP. Butyrate is well known to have beneficial effects on intestinal

barrier function (85-94), and low levels of intestinal butyrate producers has been previously

associated with microbial translocation and immune activation in PLWH (95). The prior study of

Gelpi *et al.* (23) also found increased levels of bacteria in the *Desulfovibrionaceae* family to be

linked with HIV-associated metabolic syndrome; These bacteria produce hydrogen sulfide, a

compound that can compromise the intestinal mucus layer by reducing disulfide bonds (96).

Further supporting that microbes that may compromise the mucus layer can impact barrier

function in this context, we found *Dorea sp*, which encodes the genes required to utilize the

canonical sialic acid Neu5Ac as a carbon source (97), to be an important predictor of the

metabolic disease score and to positively correlate with LBP. Hypo-sialylated intestinal glycans

have previously been linked with higher abundance of glycan-degrading species and higher

levels of microbial translocation in HIV-infected individuals (98). Taken together, these results

suggest that impaired metabolic health in HIV positive and negative MSM may be influenced by

impaired intestinal barrier linked with both protective and detrimental metabolic activity of

intestinal bacteria.

In our metabolomic analysis, we identified 150 metabolites in blood that correlated with the

metabolic disease score. To identify compounds whose prevalence may be related to the gut

microbiome we used two complimentary approaches. First, we used the bioinformatics tool

AMON (47), which allows us to specifically evaluate which compounds could have been directly

produced by the gut microbiome but is limited by a lack of KEGG annotations for many

compounds. Second, we measured which compounds changed in relative abundance in GF

versus mice colonized with feces from our study cohort, which can identify microbial influence in

unannotated compounds but cannot differentiate between direct production/consumption by

microbes versus indirect influence. These results will also be influenced by physiological

differences between mice and humans and incomplete colonization. Although these

weaknesses may have led us to underestimate which of the 150 metabolic disease associated

compounds may have been related to the microbiome, it still identified compounds that

supported a mechanistic link between gut microbes, metabolites, and metabolic disease in HIV-

infected individuals on ART.

First, we found a negative correlation between the microbially-produced non-canonical amino

acid, dehydroalanine, and metabolic disease score. Dehydroalanine is a component of

lantibiotics that are active against Gram-positive bacteria. Second, we found a positive

correlation with bacteriohopane-32,33,34,35-tetrol. This compound is a lipoxygenase inhibitor

that prevents the formation of hydroxyicosatetraenoic acid and various leukotrienes from

arachidonic acid (99), which have been linked with the development of cardiovascular disease

and metabolic syndrome (100). This association of a potentially protective metabolite with an

increase in metabolic impairment seems counterintuitive; however, it may be indicative of larger

systemic changes in arachidonic acid metabolism. Third, we identified a PC and a PE

associated with both the microbiome and metabolic disease score. Changes in PCs and/or PEs

have been previously implicated in atherosclerosis, insulin resistance and obesity (51). AMON

analysis indicated that both PCs and PEs can be synthesized by intestinal bacteria; however,

these compounds can also be synthesized in the host and may be found in the diet. In our

analysis, however, PE(18:1/20:1) levels were higher in colonized compared to GF mice

indicating that intestinal bacteria do influence overall levels despite diverse potential sources.

Finally, we observed increased levels of several plasma triglycerides in the humanized

compared to GF mice, including two plasma triglycerides that were significantly associated with

metabolic disease score. This confirms the influence of the gut microbiome on host plasma

triglycerides (101-103). However, we did not find any strong associations between these

triglycerides and specific microbes within our dataset, indicating a potential need for studies

conducted in larger cohorts or with shotgun metagenomics to look for functional correlates.

In conclusion, we observed a relationship between diet, gut microbiome, plasma metabolome,

and peripheral immune markers of inflammation and metabolic disease in HIV positive and HR-

MSM. Our results suggest a central role of inflammatory processes linked with bacterial

translocation and interaction with the gut microbiome, age and BMI in metabolic disease among

HIV positive and negative MSM. Our results also suggest contributions of low fiber, key

vitamins, and microbially produced metabolites. These results illuminate potential microbiome-

targeted therapies and personalized diet recommendation given an interacting set of gut

microbes and other host factors. Understanding these relationships further may provide novel

treatments to improve the metabolic disease and inflammatory outcomes of MSM living with

HIV.

Methods

<u>Subject Recruitment</u>

Participants were residents of the Denver, Colorado metropolitan area and the study was conducted at the Clinical Translational Research Center of the University of Colorado Hospital. The study was reviewed and approved by the Colorado Multiple Institutional Review Board and informed consent was obtained from all participants. For detailed criteria on recruitment of our five cohorts (HIV negative MSW; HIV negative MSM; HIV positive, ART naïve MSM; HIV positive ART-treated MSM with LD; and HIV positive ART-treated MSM without LD) see supplemental methods.

Feces, a fasting blood sample, and clinical surveys were collected from participants in order to obtain analytes for the study design outlined in Figure 3 (Supplemental Table 2). Additional information about relevant clinical measures such as probiotic use were collected via a questionnaire and study participants also filled out information on typical frequency of high-risk sexual practices and on typical levels of gastrointestinal issues such as bloating, constipation, nausea and diarrhea.

<u>Diet Data FFQ Collection</u>

Typical dietary consumption over the prior year was collected using the Diet History Questionnaire II (104). Diet composition was processed using the Diet*Calc software and the dhq2.database.092914 database (105). All reported values are based on USDA nutrition guidelines. Reported dietary levels were normalized per 1000 kcal. To reduce the number of comparisons within the diet survey data, we binned highly co-correlating groups of measures within the data types into modules (Supplemental Table 2). These modules were defined using the tool, SCNIC (106).

<u>Immune Data Collection</u>

Whole blood was collected in sodium heparin vacutainers and centrifuged at 1700rpm for 10

minutes for plasma collection. Plasma was aliquoted into 1mL microcentrifuge tubes and stored

at -80°C. For ELISA preparation, plasma was thawed, kept cold, and centrifuged at 2000xg for

20 minutes before ELISA plating. Markers for sCD14, sCD163, and FABP-2 were measured

from plasma using standard ELISA kits from R&D Systems (DC140, DC1630 &DFBP20).

Positive testing controls for each ELISA kit were also included (R&D Systems QC20, QC61, &

QC213). LBP was measured by standard ELISA using Hycult Biotech kit HK315-02. Markers for

IL-6, IL-10, TNF-α, MCP-1, and IL-22 were measured using Meso Scale Discovery's U-PLEX

Biomarker Group 1 multiplex kit K15067L-1. Markers for SAA, VCAM-1, ICAM-1, and CRP were

measured using Meso Scale Discovery's V-Plex Plus Vascular Injury Panel 2 multiplex kit

K15198G-1. Vascular Injury Control Pack 1 C4198-1 was utilized as a positive control for this

assay. Markers for GM-CSF, IL-7, IL-12/23p40, IL-15, IL-16, IL-17A, TNF-β, and VEGF were

measured using Meso Scale Discovery's V-Plex Plus Cytokine Panel 1 multiplex kit K151A0H-1.

Cytokine Panel 1 Control Pack C4050-1 was utilized as a positive control for this assay. Plasma

samples were diluted per manufacturer's recommendation for all assays. Standard ELISA kit

plates were measured using a Vmax® Kinetic Microplate Reader with Softmax® Pro Software

from Molecular Devices LLC. Multiplex ELISA kits from Meso Scale Discovery were measured

using the QuickPlex SQ 120 with Discovery Workbench 4.0 software.

<u>Gnotobiotic Mouse Protocols</u>

Germ-free C57/BL6 mice were purchased from Taconic and bred and maintained in flexible film

isolator bubbles, fed with standard mouse chow. Three days before they were gavaged, male

mice between 5-7 weeks of age were switched to a western high-fat diet and were fed this diet

for the remainder of the experiment. Diets were all obtained from Envigo (Indiana): Standard

chow - Teklad global soy protein-free extruded (item 2920X -

https://www.envigo.com/resources/data-sheets/2020x-datasheet-0915.pdf), Western Diet – New

Total Western Diet (item TD.110919). See Supplemental Table 4 for detailed diet composition.

Mice were gavaged with 200 µL of fecal solutions prepared from 1.5 g of donor feces mixed in 3

mL of anaerobic PBS (18). Mice were housed individually following gavage for three weeks in a

Tecniplast iso-positive caging system, with each cage having HEPA filters and positive

pressurization for bioexclusion. Feces were collected from mice at day 21 for 16S rRNA gene

sequencing. Mice were euthanized at 21 days post gavage using isoflurane overdose and all

efforts were made to minimize suffering. Blood from euthanized animals was collected using

cardiac puncture and cells were pelleted in K2-EDTA tubes; plasma was then aliquoted and

stored at -80° C.

<u>Metabolomics Methods</u>

Plasma Sample Preparation

A modified liquid-liquid extraction protocol was used to extract hydrophobic and hydrophilic

compounds from the plasma samples (107). Briefly, 50 µL of plasma spiked with internal

standards underwent a protein crash with 250 µL ice cold methanol. 750 µL methyl tert-butyl

ether (MTBE) and 650 µL 25% methanol in water were added to extract the hydrophobic and

hydrophilic compounds, respectively. 500 µL of the upper hydrophobic layer and 400 µL of the

lower hydrophilic layer were transferred to separate autosampler vials and dried under nitrogen.

The hydrophobic layer was reconstituted with 100 µL of methanol and the hydrophilic layer was

reconstituted with 50 µL 5% acetonitrile in water. Both fractions were stored at -80 °C until

LC/MS analysis.

Liquid Chromatography Mass Spectrometry

The hydrophobic fractions were analyzed using reverse phase chromatography on an Agilent

Technologies (Santa Clara, CA) 1290 ultra-high precision liquid chromatography (UHPLC)

system on an Agilent Zorbax Rapid Resolution HD SB-C18, 1.8um (2.1 x 100mm) analytical

column as previously described (107, 108). The hydrophilic fractions were analyzed using

hydrophilic interaction liquid chromatography (HILIC) on a 1290 UHPLC system using an Agilent

InfinityLab Poroshell 120 HILIC-Z (2.1 x 100mm) analytical column with gradient conditions as

previously described (109) with mass spectrometry modifications as follows: nebulizer pressure:

35psi, gas flow: 12L/min, sheath gas temperature: 275C, sheath gas flow: 12L/min, nozzle

voltage: 250V, Fragmentor: 100V. The hydrophobic and hydrophilic fractions were run on

Agilent Technologies (Santa Clara, CA) 6545 Quadrupole Time of Flight (QTOF) mass

spectrometer. Both fractions were run in positive electrospray ionization (ESI) mode.

Mass Spectrometry Data Processing

Compound data was extracted using Agilent Technologies (Santa Clara, CA) MassHunter

Profinder Version 10 software in combination with Agilent Technologies Mass Profiler

Professional Version 14.9 (MPP) as described previously (47). Briefly, Batch Molecular Feature

Extraction (BMFE) was used in Profinder to extract compound data from all samples and

sample preparation blanks. The following BMFE parameters were used to group individual

molecular features into compounds: charge state 1-2, with +H, +Na, +NH4 and/or +K charge

carriers. To reduce the presence of missing values, a theoretical mass and retention time

database was generated for compounds present in samples only from a compound exchange

format (.cef) file. This .cef file was then used to re-mine the raw sample data in Profinder using

Batch Targeted Feature Extraction.

An in-house database containing KEGG, METLIN, Lipid Maps, and HMDB spectral data was

used to putatively annotate metabolites based on accurate mass ($\leq$ 10 ppm), isotope ratios and

isotopic distribution. This corresponds to a Metabolomics Standards Initiative metabolite

identification level three (110). To improve compound identification, statistically significant

compounds underwent tandem MS using 10, 20, and 40V. Fragmentation patterns of identified

compounds were matched to either NIST14 and NIST17 MSMS libraries, or to the *in silico*

libraries, MetFrag (111) and Lipid Annotator 1.0 (Agilent) (112).

Microbiome-associated metabolites

Microbiome-associated metabolites were defined using metabolites identified as significantly

different in abundance between germ-free compared to humanized gnotobiotic mice and/or

metabolites identified as microbially produced by the tool AMON (47).

For the gnotobiotic mouse analysis aqueous and lipid metabolites were analyzed separately.

Metabolites that were present in <20% of samples were filtered out before analysis. Significant

difference was determined using a Student's t-test with FDR p-value correction. FDR-corrected

p values < 0.05 were deemed significant. Significant metabolites also present in the human

samples were retained for further analysis.

For the AMON-identified metabolites, the tool used an inferred metagenome, which was

calculated using the PICRUSt2 QIIME2 plugin (49) and default parameters; a list of all identified

KEGG IDs from the metabolite data (see metabolome methods); and KEGG flat files

(downloaded 2019/06/10). AMON determined metabolites observed that could be produced by

the given genes list. These metabolites were kept for analysis in addition to the gnotobiotic

mouse identified metabolites. Those without any putative classification were removed from

analysis.

<u>Microbiome Methods</u>

Sample Collection, Extraction, and Sequencing

Stool samples were collected by the patient within 24 hours of their clinic visit on sterile swabs

dipped into a full fecal sample deposited into a commode specimen collector. Samples were

kept cold or frozen at -20°C during transport prior to being stored at -80°C. DNA was extracted

using the standard DNeasy PowerSoil Kit protocol (Qiagen). Extracted DNA was PCR amplified

with barcoded primers targeting the V4 region of 16S rRNA gene according to the Earth

Microbiome Project 16S Illumina Amplicon protocol with the 515F:806R primer constructs (113).

Control sterile swab samples that had undergone the same DNA extraction and PCR

amplification procedures were also processed. Each PCR product was quantified using

PicoGreen (Invitrogen), and equal amounts (ng) of DNA from each sample were pooled and

cleaned using the UltraClean PCR Clean-Up Kit (MoBio). Sequences were generated on six

runs on a MiSeq sequencing platform (Illumina, San Diego, CA).

Microbiome Sequence Processing and Analysis

Microbiome processing was performed using QIIME2 version 2018.8.0 (114). Data was

sequenced across five sequencing runs. Each run was demultiplexed and denoised separately

using the DADA2 q2 plugin (115). Individual runs were then merged together and 99% *de novo*

OTUs were defined using vSEARCH (116). Features were classified using the skLearn classifier

in QIIME2 with a classifier that was pre-trained on GreenGenes13_8 (117). The phylogenetic

tree was building using the SEPP plugin (118). Features that did not classify at the phylum level

or were classified as mitochondria or chloroplast were filtered from the analysis. Samples were

rarefied at 19,986 reads. To reduce the number of comparisons within the microbiome, we

binned highly co-correlating groups of measures within the data types into modules

(Supplemental Table 2). These modules were defined using the tool, SCNIC (119). For

statistical analysis features present in <20% of samples were filtered out.

Bioinformatics

Module definition

Modules were called on microbiome and diet data. Modules were defined using the tool SCNIC

(119). The q2-SCNIC plugin was used with default parameters for the microbiome data and

standalone SCNIC was used for the diet data (https://github.com/shafferm/SCNIC). Specifically,

for each data type SCNIC was used to first identify pairwise correlations between all features.

Pearson correlation was used for diet and SparCC (120), which takes into account

compositionality, was used for microbiome data. Modules were then selected with a shared

minimum distance (SMD) algorithm. The SMD method defines modules by first applying

complete linkage hierarchical clustering to correlation coefficients to make a tree of features.

Modules are defined as subtrees where all pairwise correlations between all pairs of tips have

an R value greater than defined minimum. The diet modules were defined using a Pearson r^2

cutoff of 0.75. The microbiome modules were defined using a SparCC minimum r cutoff of 0.35.

To summarize modules SCNIC uses a simple summation of count data from all features in a

module. Application of SCNIC reduced the number of evaluated features from 6,913 to 6,818 for

microbiome and 59 to 29 for diet data.

Statistical Analysis

All statistics were performed in R. For non-parametric tests Spearman rank correlation and

Kruskal-Wallis test were used. For parametric tests linear models and Student's t-test were

used.

Data analysis tools

Metabolic disease score was calculated using PCA in R with prcomp. Data was scaled using

default method within the prcomp library. All random forest analysis tools were used in R.

Standard random forest was performed using the randomForest function. Variable selection was

performed in R using the tool VSURF (38). Interaction analysis was performed in R using the

tool iRF (42).

Data Availability

Microbiome data is available at EBI/ENA (https://www.ebi.ac.uk/ena) accession number:

ERP125300. Immune and diet data are available along with the microbiome data as associated

metadata. Metabolomics data will be available on Metabolomic Workbench

(https://www.metabolomicsworkbench.org) upon publishing. Until publicly available it is available

upon request.

Statements

Acknowledgements

We would like to thank our study participants for contributing their samples and time to this

study. We would also like to thank Christine Griesmer for her contributions to subject

recruitment and Brandi Wagner and Maggie Stanislawski for insights into statistical analysis

methods.

Funding sources

This work was funded by R01 DK104047 and R01 DK108366 with additional support from

NIH/NCATS Colorado CTSA Grant Number UL1 TR002535. High performance computing was

supported by a cluster at the University of Colorado Boulder funded by National Institutes of

Health 1S10OD012300. AJS Armstrong was supported by T32 AI052066. Contents are the

authors' sole responsibility and do not necessarily represent official NIH views.

Declarations of interests

The authors declare that they have no conflicts of interest.

Author contributions

AJSA analyzed and interpreted all data. AJSA and CAL wrote the manuscript. NR guided generation and interpretation of metabolomics data. KQ and KAD prepared, ran, and processed metabolomics. KQ ran metabolic pathway analysis. SXL prepared and conducted mouse experiments. JMS ran immunological assays. NMN prepared and ran sequencing and coordinated fecal sample and metadata collection from study subjects. SF recruited subjects, collected samples, and maintained regulatory compliance. TJM and JH collected and aided in interpretation and processing of diet data. CAL, BEP, and TC conceptualized and led the study.TC guided all clinical data collection and subject recruitment and provided clinical insight into study populations. BEP guided generation and interpretation of immune data. CAL guided microbiome data generation and multi'omic data analysis. All authors read and approved the final manuscript.

684 **Tables**

685 **Table 1. Description of full study cohort**

	HIV-negative MSW	HIV-negative MSM	HIV-positive MSM, untreated	HIV-positive MSM, treated	HIV-positive MSM, treated, with LD	P
n	22	32	14	20	25	
Age (years)	33 (27.3-38.5)	34 (29.8-44.5)	34 (26.5-40.3)	46 (42.8-50.5)	60 (54-64)	***
BMI (kg/m2)	25.2 (23.0-27.0)	25.5 (20.2-28.0)	21.4 (20.2-25.6)	23.9 (22.6-26.2)	25.8 (23.0-28.0)	NS
CD4 cell count	NA	NA	538 (406-732)	586 (420-878)	659 (550-908)	NS
CD4 nadir	NA	NA	496 (409-612)	256 (118-416)	152 (70-350)	***
Viral load	NA	NA	101,400 (20,300-292,514)	20 (0-20)	0 (0-20)	***
Cholesterol drugs/statins n (%)	2 (9.1%)	3 (9.4%)	1 (7.1%)	4 (20%)	14 (56%)	+++

Numbers reported are median (IQR); Kruskal-Wallis test * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$,
NS $p > 0.05$; Fisher's exact test +++ $p < 0.001$. See Supplemental Table 1 for pairwise
comparisons between groups.

Figures

Figure 1. Study design schematic. A. Measures were collected from four compartments: gut

microbiome, peripheral immune, diet questionnaire, and plasma metabolome. These separate

compartments can all influence each other and can be influenced by other clinical and

demographic characteristics such as HIV and treatment status. **B.** Analysis pipeline for the

study. First, a metabolic health outcome was determined. Second, informative variables were

selected using random forest analysis. Lastly, the relationships between these informative

variables and the metabolic health outcome were examined.

Figure 2. Calculation of the metabolic disease score. A. PCA of metabolic markers in fasting

blood of 164 men and women: 113 participants described in this paper along with 51 individuals

recruited at the same time and under the same exclusion criteria as study participants.

Metabolic disease score is calculated as the PC1 coordinates shifted to a minimum of one and

log transformed. **B.** Metabolic disease scores broken up by cohort. The percentages noted

above the groups are the percent of individuals with a score above our metabolic impairment

cutoff (Supplemental Figure 1). There is no significant difference between the proportions in

each group (Fischer's exact test, p = 0.11) or between mean ranks in each group (Kruskal-

Wallis test p = 0.13). **C.** Relationships between metabolic disease score and age stratified by

cohort. Statistical significance of slopes are indicated and were calculated with the linear model:

score ~ age + cohort + age*cohort. P-value annotations: ** < 0.01; * < 0.05.

Figure 3. Networks of selected measures reveal several strong associations with

metabolic disease score and between measures. Correlation sub-networks of **A.** all the non-

microbe selected measures. **B.** the nearest neighbors of LBP. All Spearman rank correlations

with an FDR p < 0.25 are shown. Subnetworks were pulled from a larger network of all VSURF

selected measures (Supplemental Figure 3, Supplemental Table 2). **C.** Network of interactions

between measures calculated using iRF. All edges represent an interaction (i.e., proximity in a

decision tree) that occurred in 30% or more of the decision trees.

Figure 4. Microbiome-associated metabolite workflow. Two-pronged approach of identifying

microbiome-associated metabolites. Numbers in bold are counts of metabolites.

Supplemental Tables

Supplemental Table 1. Pairwise test p-values for Table 1 demographics

Supplemental Table 2. Study measures by datatype and SCNIC-calculated modules for

microbiome and diet survey data

Supplemental Table 3. VSURF-selected features: correlation with metabolic disease score,

edge table for inter-variable correlations, and iRF analysis results

Supplemental Table 4. Gnotobiotic mouse experiment set-up and western diet

Supplemental Table 5. Microbiome-associated metabolites list with source of identification,

metabolites correlating with metabolic disease score and mBrole pathway analysis results

Supplemental Figures

Supplemental Figure 1. Metabolic disease score cutoff calculations using regressions between metabolic disease score and the metrics used in the PCA analysis. Triglycerides, fasting glucose, HDL and LDL have well-define clinical cut-offs for high values and were used to calculate the healthy-unhealth cutoff. Linear model was calculated modeling metabolic disease score by each marker. The high value intercept of the regression line is marked with a dotted line and value annotated on the plot. The solid line is the defined healthy-unhealthy cutoff calculated as the mean of the four cutoff values. P values are from the linear model.

Supplemental Figure 2. Histogram of percent variation explained in permuted VSURF. Metabolic disease score was permuted 1,000 times and passed through VSURF. The resulting variables were run through a standard random forest and the percent variation explained was calculated. The blue line represents the percent variation explained for the true VSURF. P value was calculated using a one tailed test.

Supplemental Figure 3. Correlation network of VSURF-selected variables. Correlation network of **A)** all VSURF-selected variables, **B)** neighboring nodes of dietary fiber, and **C)** neighboring nodes of age. All Spearman rank correlations with an FDR p < 0.25 are shown. See Supplemental Table 3 for the edge table and Supplemental Table 3 for the node table.

Supplemental Figure 4. Metabolic disease associated metabolites by model type with age accounting. Venn diagram summarizes all significant results (FDR-corrected p-value < 0.05) of each test and the metabolites that overlap between tests that do and do not account for age correction.

References

1.   Non LR, Escota GV, Powderly WG. 2016. HIV and its relationship to insulin resistance

and lipid abnormalities. Translational Research 183:41-56.

2.   Nix LM, Tien PC. 2014. Metabolic syndrome, diabetes, and cardiovascular risk in HIV.

Curr HIV/AIDS Rep 11:271-8.

3.   Lake JE, Li X, Palella FJ, Jr., Erlandson KM, Wiley D, Kingsley L, Jacobson LP, Brown

TT. 2018. Metabolic health across the BMI spectrum in HIV-infected and HIV-uninfected

men. AIDS 32:49-57.

4.   Willig AL, Overton ET. 2016. Metabolic Complications and Glucose Metabolism in HIV

Infection: A Review of the Evidence. Curr HIV/AIDS Rep 13:289-96.

5.   Monczor AN, Li X, Palella FJ, Jr., Erlandson KM, Wiley D, Kingsley LA, Post WS,

Jacobson LP, Brown TT, Lake JE. 2018. Systemic Inflammation Characterizes Lack of

Metabolic Health in Nonobese HIV-Infected Men. Mediators Inflamm 2018:5327361.

6.   Funderburg NT, Mehta NN. 2016. Lipid Abnormalities and Inflammation in HIV Inflection.

Curr HIV/AIDS Rep 13:218-25.

7.   Beltran LM, Rubio-Navarro A, Amaro-Villalobos JM, Egido J, Garcia-Puig J, Moreno JA.

2015. Influence of immune activation and inflammatory response on cardiovascular risk

associated with the human immunodeficiency virus. Vasc Health Risk Manag 11:35-48.

8.   Tsiodras S, Mantzoros C, Hammer S, Samore M. 2000. Effects of Protease Inhibitors on

Hyperglycemia, Hyperlipidemia, and Lipodystrophy A 5-Year Cohort Study. Arch Intern

Med 160:2050-2056.

9.   Tilg H, Kaser A. 2011. Gut microbiome, obesity, and metabolic dysfunction. The Journal

of Clinical Investigation 121:2126-2132.

10.   Sonnenburg JL, Bäckhed F. 2016. Diet–microbiota interactions as moderators of human

metabolism. Nature 535:56-64.

11. Pedersen HK, Gudmundsdottir V, Nielsen HB, Hyotylainen T, Nielsen T, Jensen BA,

Forslund K, Hildebrand F, Prifti E, Falony G, Le Chatelier E, Levenez F, Dore J, Mattila I,

Plichta DR, Poho P, Hellgren LI, Arumugam M, Sunagawa S, Vieira-Silva S, Jorgensen

T, Holm JB, Trost K, Meta HITC, Kristiansen K, Brix S, Raes J, Wang J, Hansen T, Bork

P, Brunak S, Oresic M, Ehrlich SD, Pedersen O. 2016. Human gut microbes impact host

serum metabolome and insulin sensitivity. Nature 535:376-81.

12. Matey-Hernandez ML, Williams FMK, Potter T, Valdes AM, Spector TD, Menni C. 2018.

Genetic and microbiome influence on lipid metabolism and dyslipidemia. Physiol

Genomics 50:117-126.

13. Karlsson F, Tremaroli V, Nielsen J, Bäckhed F. 2013. Assessing the human gut

microbiota in metabolic diseases. Diabetes 62:3341-9.

14. Neff CP, Krueger O, Xiong K, Arif S, Nusbacher N, Schneider JM, Cunningham AW,

Armstrong A, Li S, McCarter MD, Campbell TB, Lozupone CA, Palmer BE. 2018. Fecal

Microbiota Composition Drives Immune Activation in HIV-infected Individuals.

EBioMedicine 30:192-202.

15. Armstrong AJS, Shaffer M, Nusbacher NM, Griesmer C, Fiorillo S, Schneider JM,

Preston Neff C, Li SX, Fontenot AP, Campbell T, Palmer BE, Lozupone CA. 2018. An

exploration of Prevotella-rich microbiomes in HIV and men who have sex with men.

Microbiome 6:198.

16. Vujkovic-Cvijin I, Somsouk M. 2019. HIV and the Gut Microbiota: Composition,

Consequences, and Avenues for Amelioration. Curr HIV/AIDS Rep doi:10.1007/s11904-

019-00441-w.

17. Noguera-Julian M, Rocafort M, Guillén Y, Rivera J, Casadellà M, Nowak P, Hildebrand

F, Zeller G, Parera M, Bellido R, Rodríguez C, Carrillo J, Mothe B, Coll J, Bravo I,

Estany C, Herrero C, Saz J, Sirera G, Torrela A, Navarro J, Crespo M, Brander C,

Negredo E, Blanco J, Guarner F, Calle ML, Bork P, Sönnerborg A, Clotet B, Paredes R.

2016. Gut Microbiota Linked to Sexual Preference and HIV Infection. EBioMedicine

5:135-46.

18.   Li SX, Sen S, Schneider JM, Xiong KN, Nusbacher NM, Moreno-Huizar N, Shaffer M,

Armstrong AJS, Severs E, Kuhn K, Neff CP, McCarter M, Campbell T, Lozupone CA,

Palmer BE. 2019. Gut microbiota from high-risk men who have sex with men drive

immune activation in gnotobiotic mice and in vitro HIV infection. PLoS Pathog

15:e1007611.

19.   Cui HL, Ditiatkovski M, Kesani R, Bobryshev YV, Liu Y, Geyer M, Mukhamedova N,

Bukrinsky M, Sviridov D. 2014. HIV protein Nef causes dyslipidemia and formation of

foam cells in mouse models of atherosclerosis. FASEB Journal 28:2828-2839.

20.   Auclair M, Guenantin AC, Fellahi S, Garcia M, Capeau J. 2020. HIV antiretroviral drugs,

dolutegravir, maraviroc and ritonavir-boosted atazanavir use different pathways to affect

inflammation, senescence and insulin sensitivity in human coronary endothelial cells.

PLoS One 15:e0226924.

21.   Ahmed D, Roy D, Cassol E. 2018. Examining Relationships between Metabolism and

Persistent Inflammation in HIV Patients on Antiretroviral Therapy. Mediators Inflamm

2018:6238978.

22.   Reid M, Ma Y, Scherzer R, Price JC, French AL, Plankey MW, Grunfeld C, Tien PC.

2017. Higher CD163 levels are associated with insulin resistance in hepatitis C virus-

infected and HIV-infected adults. AIDS 31:385-393.

23.   Gelpi M, Vestad B, Hansen SH, Holm K, Drivsholm N, Goetz A, Kirkby NS, Lindegaard

B, Lebech AM, Hoel H, Michelsen AE, Ueland T, Gerstoft J, Lundgren J, Hov JR,

Nielsen SD, Troseid M. 2020. Impact of Hiv-Related Gut Microbiota Alterations on

Metabolic Comorbidities. Clin Infect Dis doi:10.1093/cid/ciz1235.

24.   Lim PS, Chang YK, Wu TK. 2019. Serum Lipopolysaccharide-Binding Protein is

833   Associated with Chronic Inflammation and Metabolic Syndrome in Hemodialysis

834   Patients. Blood Purif 47:28-36.

835   25.   Hersoug LG, Moller P, Loft S. 2018. Role of microbiota-derived lipopolysaccharide in

836   adipose tissue inflammation, adipocyte size and pyroptosis during obesity. Nutr Res Rev

837   31:153-163.

838   26.   Awoyemi A, Troseid M, Arnesen H, Solheim S, Seljeflot I. 2018. Markers of metabolic

839   endotoxemia as related to metabolic syndrome in an elderly male population at high

840   cardiovascular risk: a cross-sectional study. Diabetol Metab Syndr 10:59.

841   27.   Liu X, Lu L, Yao P, Ma Y, Wang F, Jin Q, Ye X, Li H, Hu FB, Sun L, Lin X. 2014.

842   Lipopolysaccharide binding protein, obesity status and incidence of metabolic syndrome:

843   a prospective study among middle-aged and older Chinese. Diabetologia 57:1834-41.

844   28.   Troseid M, Manner IW, Pedersen KK, Haissman JM, Kvale D, Nielsen SD. 2014.

845   Microbial translocation and cardiometabolic risk factors in HIV infection. AIDS Res Hum

846   Retroviruses 30:514-22.

847   29.   Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, Neyrinck AM, Fava F,

848   Tuohy KM, Chabo C, Waget A, Delmee E, Cousin B, Sulpice T, Chamontin B, Ferrieres

849   J, Tanti JF, Gibson GR, Casteilla L, Delzenne NM, Alessi MC, Burcelin R. 2007.

850   Metabolic endotoxemia initiates obesity and insulin resistance. Diabetes 56:1761-72.

851   30.   Lew WY, Bayna E, Molle ED, Dalton ND, Lai NC, Bhargava V, Mendiola V, Clopton P,

852   Tang T. 2013. Recurrent exposure to subclinical lipopolysaccharide increases mortality

853   and induces cardiac fibrosis in mice. PLoS One 8:e61057.

854   31.   Fei N, Zhao L. 2013. An opportunistic pathogen isolated from the gut of an obese human

855   causes obesity in germfree mice. ISME J 7:880-4.

32.   Somsouk M, Estes JD, Deleage C, Dunham RM, Albright R, Inadomi JM, Martin JN,

Deeks SG, McCune JM, Hunt PW. 2015. Gut epithelial barrier and systemic

inflammation during chronic HIV infection. AIDS (London, England) 29:43-51.

33.   Klase Z, Ortiz A, Deleage C, Mudd JC, Quiñones M, Schwartzman E, Klatt NR, Canary

L, Estes JD, Brenchley JM. 2015. Dysbiotic bacteria translocate in progressive SIV

infection. Mucosal Immunology 8:1009-1020.

34.   Palmer CD, Tomassilli J, Sirignano M, Romero-Tejeda M, Arnold KB, Che D,

Lauffenburger DA, Jost S, Allen T, Mayer KH, Altfeld M. 2014. Enhanced immune

activation linked to endotoxemia in HIV-1 seronegative MSM. AIDS 28:2162-6.

35.   Hammer SM, Sobieszczyk ME, Janes H, Karuna ST, Mulligan MJ, Grove D, Koblin BA,

Buchbinder SP, Keefer MC, Tomaras GD, Frahm N, Hural J, Anude C, Graham BS,

Enama ME, Adams E, DeJesus E, Novak RM, Frank I, Bentley C, Ramirez S, Fu R,

Koup RA, Mascola JR, Nabel GJ, Montefiori DC, Kublin J, McElrath MJ, Corey L, Gilbert

PB, Team HS. 2013. Efficacy trial of a DNA/rAd5 HIV-1 preventive vaccine. N Engl J

Med 369:2083-92.

36.   Wiley JF, Carrington MJ. 2016. A metabolic syndrome severity score: A tool to quantify

cardio-metabolic risk factors. Prev Med 88:189-95.

37.   Hillier TA, Rousseau A, Lange C, Lepinay P, Cailleau M, Novak M, Calliez E,

Ducimetiere P, Balkau B, Cohort DESIR. 2006. Practical way to assess metabolic

syndrome using a continuous score obtained from principal components analysis.

Diabetologia 49:1528-35.

38.   Genuer R, Poggi J-M, Tuleau-Malot C. 2015. VSURF: An R Package for Variable

Selection Using Random Forests. The R Journal, R Foundation for Statistical Computing

7:12-33.

39.   Marchetti G, Tincati C, Silvestri G. 2013. Microbial translocation in the pathogenesis of

HIV infection and AIDS. Clinical microbiology reviews 26:2-18.

40.   Duncan SH, Holtrop G, Lobley GE, Calder AG, Stewart CS, Flint HJ. 2004. Contribution

883   of acetate to butyrate formation by human faecal bacteria. Br J Nutr 91:915-23.

884   41.   Duncan SH, Barcenilla A, Stewart CS, Pryde SE, Flint HJ. 2002. Acetate utilization and

885   butyryl coenzyme A (CoA):acetate-CoA transferase in butyrate-producing bacteria from

886   the human large intestine. Appl Environ Microbiol 68:5186-90.

887   42.   Basu S, Kumbier K, Brown JB, Yu B. 2018. Iterative random forests to discover

888   predictive and stable high-order interactions. Proc Natl Acad Sci U S A 115:1943-1948.

889   43.   Aberg JA. 2020. Aging and HIV infection: focus on cardiovascular disease risk. Top

890   Antivir Med 27:102-105.

891   44.   Deeks SG. 2011. HIV infection, inflammation, immunosenescence, and aging. Annu Rev

892   Med 62:141-55.

893   45.   Leng SX, Margolick JB. 2015. Understanding frailty, aging, and inflammation in HIV

894   infection. Curr HIV/AIDS Rep 12:25-32.

895   46.   Pathai S, Bajillan H, Landay AL, High KP. 2014. Is HIV a model of accelerated or

896   accentuated aging? J Gerontol A Biol Sci Med Sci 69:833-42.

897   47.   Shaffer M, Thurimella K, Quinn K, Doenges K, Zhang X, Bokatzian S, Reisdorph N,

898   Lozupone CA. 2019. AMON: annotation of metabolite origins via networks to integrate

899   microbiome and metabolome data. BMC Bioinformatics 20:614.

900   48.   Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. 2017. KEGG: new

901   perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Research

902   45:D353-D361.

903   49.   Douglas GM, Maffei VJ, Zaneveld J, Yurgel SN, Brown JR, Taylor CM, Huttenhower C,

904   Langille MGI. 2020. PICRUSt2: An improved and extensible approach for metagenome

905   inference. doi:10.1101/672295.

906   50.   Frolkis A, Knox C, Lim E, Jewison T, Law V, Hau DD, Liu P, Gautam B, Ly S, Guo AC,

907   Xia J, Liang Y, Shrivastava S, Wishart DS. 2010. SMPDB: The Small Molecule Pathway

908   Database. Nucleic Acids Res 38:D480-7.

909   51.   van der Veen JN, Kennelly JP, Wan S, Vance JE, Vance DE, Jacobs RL. 2017. The

910   critical role of phosphatidylcholine and phosphatidylethanolamine metabolism in health

911   and disease. Biochim Biophys Acta Biomembr 1859:1558-1572.

912   52.   Kloke JD, McKean JW. 2012. Rfit: Rank-based Estimation for Linear Models. The R

913   Journal 4.

914   53.   Martin-Iguacel R, Negredo E, Peck R, Friis-Moller N. 2016. Hypertension Is a Key

915   Feature of the Metabolic Syndrome in Subjects Aging with HIV. Curr Hypertens Rep

916   18:46.

917   54.   Verhaegen P, Borchelt M, Smith J. 2003. Relation between cardiovascular and

918   metabolic disease and cognition in very old age: cross-sectional and longitudinal findings

919   from the berlin aging study. Health Psychol 22:559-69.

920   55.   Montessori V, Press N, Harris M, Akagi L, Montaner JSG. 2004. Adverse effects of

921   antiretroviral therapy for HIV infection. CMAJ : Canadian Medical Association journal =

922   journal de l'Association medicale canadienne 170:229-38.

923   56.   Aberg JA. 2012. Aging, inflammation, and HIV infection. Top Antivir Med 20:101-5.

924   57.   Zapata HJ, Quagliarello VJ. 2015. The microbiota and microbiome in aging: potential

925   implications in health and age-related diseases. J Am Geriatr Soc 63:776-81.

926   58.   Evans WJ, Cyr-Campbell D. 1997. Nutrition, Exercise, and Healthy Aging. Journal of the

927   American Dietetic Association 97:632-638.

928   59.   Nagpal R, Mainali R, Ahmadi S, Wang S, Singh R, Kavanagh K, Kitzman DW,

929   Kushugulova A, Marotta F, Yadav H. 2018. Gut microbiome and aging: Physiological

930   and mechanistic insights. Nutr Healthy Aging 4:267-285.

931   60.   Arboleya S, Watkins C, Stanton C, Ross RP. 2016. Gut Bifidobacteria Populations in

932   Human Health and Aging. Front Microbiol 7:1204.

933   61.   Roberts JL, Liu G, Darby TM, Fernandes LM, Diaz-Hernandez ME, Jones RM, Drissi H.

934   2020. Bifidobacterium adolescentis supplementation attenuates fracture-induced

935   systemic sequelae. Biomed Pharmacother 132:110831.

936   62.   El-Bakry HA, Zahran WM, Anter SA, Zahran AS. 2013. Role of some selected

937   Bifidobacterium strains in modulating immunosenescence of aged albino rats. The

938   Journal of Basic & Applied Zoology 66:255-262.

939   63.   Woodmansey EJ. 2007. Intestinal bacteria and ageing. J Appl Microbiol 102:1178-86.

940   64.   Li Y, Lv L, Ye J, Fang D, Shi D, Wu W, Wang Q, Wu J, Yang L, Bian X, Jiang X, Jiang H,

941   Yan R, Peng C, Li L. 2019. Bifidobacterium adolescentis CGMCC 15058 alleviates liver

942   injury, enhances the intestinal barrier and modifies the gut microbiota in D-

943   galactosamine-treated rats. Appl Microbiol Biotechnol 103:375-393.

944   65.   Alexander CM, Landsman PB, Teutsch SM, Haffner SM, Third National H, Nutrition

945   Examination S, National Cholesterol Education P. 2003. NCEP-defined metabolic

946   syndrome, diabetes, and prevalence of coronary heart disease among NHANES III

947   participants age 50 years and older. Diabetes 52:1210-4.

948   66.   Meigs JB. 2003. Epidemiology of the insulin resistance syndrome. Curr Diab Rep 3:73-9.

949   67.   Galescu O, Bhangoo A, Ten S. 2013. Insulin resistance, lipodystrophy and

950   cardiometabolic syndrome in HIV/AIDS. Reviews in Endocrine and Metabolic Disorders

951   14:133-140.

952   68.   Kerchberger AM, Sheth AN, Angert CD, Mehta CC, Summers NA, Ofotokun I, Gustafson

953   D, Weiser SD, Sharma A, Adimora AA, French AL, Augenbraun M, Cocohoba J,

954   Kassaye S, Bolivar H, Govindarajulu U, Konkle-Parker D, Golub ET, Lahiri CD. 2019.

955   Weight Gain Associated with Integrase Stand Transfer Inhibitor Use in Women. Clin

956   Infect Dis doi:10.1093/cid/ciz853.

69.   Venter WDF, Moorhouse M, Sokhela S, Fairlie L, Mashabane N, Masenya M, Serenata

C, Akpomiemie G, Qavi A, Chandiwana N, Norris S, Chersich M, Clayden P, Abrams E,

Arulappan N, Vos A, McCann K, Simmons B, Hill A. 2019. Dolutegravir plus Two

Different Prodrugs of Tenofovir to Treat HIV. N Engl J Med 381:803-815.

70.   Pérez-Matute P, Pérez-Martínez L, Aguilera-Lizarraga J, Blanco JR, Oteo JA. 2015.

Maraviroc modifies gut microbiota composition in a mouse model of obesity: a plausible

therapeutic option to prevent metabolic disorders in HIV-infected patients. Revista

española de quimioterapia : publicación oficial de la Sociedad Española de

Quimioterapia 28:200-6.

71.   Guillen Y, Noguera-Julian M, Rivera J, Casadella M, Zevin AS, Rocafort M, Parera M,

Rodriguez C, Arumi M, Carrillo J, Mothe B, Estany C, Coll J, Bravo I, Herrero C, Saz J,

Sirera G, Torrella A, Navarro J, Crespo M, Negredo E, Brander C, Blanco J, Calle ML,

Klatt NR, Clotet B, Paredes R. 2019. Low nadir CD4+ T-cell counts predict gut dysbiosis

in HIV-1 infection. Mucosal Immunol 12:232-246.

72.   Gojak R, Hadziosmanovic V, Baljic R, Zecevic L, Coric J, Mijailovic Z. 2019. CD4/CD8

Ratio as a Predictor for the Occurrence of Metabolic Syndrome in HIV / AIDS Patients

During 6 Months of cART Therapy. J Med Biochem 38:489-495.

73.   Han GM, Meza JL, Soliman GA, Islam KM, Watanabe-Galloway S. 2016. Higher levels

of serum lycopene are associated with reduced mortality in individuals with metabolic

syndrome. Nutr Res 36:402-7.

74.   Zeng YC, Peng LS, Zou L, Huang SF, Xie Y, Mu GP, Zeng XH, Zhou XL, Zeng YC.

2017. Protective effect and mechanism of lycopene on endothelial progenitor cells

(EPCs) from type 2 diabetes mellitus rats. Biomed Pharmacother 92:86-94.

75.   Sluijs I, Beulens JW, Grobbee DE, van der Schouw YT. 2009. Dietary carotenoid intake

is associated with lower prevalence of metabolic syndrome in middle-aged and elderly

men. J Nutr 139:987-92.

76.   Kovatcheva-Datchary P, Nilsson A, Akrami R, Lee Ying S, De Vadder F, Arora T, Hallen

A, Martens E, Björck I, Bäckhed F. 2015. Dietary Fiber-Induced Improvement in Glucose

Metabolism Is Associated with Increased Abundance of Prevotella. Cell Metabolism

22:971-982.

77.   O'Keefe SJD, Li JV, Lahti L, Ou J, Carbonero F, Mohammed K, Posma JM, Kinross J,

Wahl E, Ruder E, Vipperla K, Naidoo V, Mtshali L, Tims S, Puylaert PGB, DeLany J,

Krasinskas A, Benefiel AC, Kaseb HO, Newton K, Nicholson JK, de Vos WM, Gaskins

HR, Zoetendal EG. 2015. Fat, fibre and cancer risk in African Americans and rural

Africans. Nature communications 6:6342.

78.   Schröder NWJ, Schumann RR. 2016. Non-LPS targets and actions of LPS binding

protein (LBP). Journal of Endotoxin Research 11:237-242.

79.   Boulangé CL, Neves AL, Chilloux J, Nicholson JK, Dumas M-E. 2016. Impact of the gut

microbiota on inflammation, obesity, and metabolic disease. Genome Medicine 8:42.

80.   Tilves CM, Zmuda JM, Kuipers AL, Nestlerode CS, Evans RW, Bunker CH, Patrick AL,

Miljkovic I. 2016. Association of Lipopolysaccharide-Binding Protein With Aging-Related

Adiposity Change and Prediabetes Among African Ancestry Men. Diabetes Care 39:385-

91.

81.   Brake DK, Smith EO, Mersmann H, Smith CW, Robker RL. 2006. ICAM-1 expression in

adipose tissue: effects of diet-induced obesity in mice. Am J Physiol Cell Physiol

291:C1232-9.

82.   Hsu LA, Chang CJ, Wu S, Teng MS, Chou HH, Chang HH, Chang PY, Ko YL. 2010.

Association between functional variants of the ICAM1 and CRP genes and metabolic

syndrome in Taiwanese subjects. Metabolism 59:1710-6.

83.   Senn JJ, Klover PJ, Nowak IA, Mooney RA. 2002. Interleukin-6 induces cellular insulin

resistance in hepatocytes. Diabetes 51:3391-9.

84.　Reigstad CS, Lunden GO, Felin J, Backhed F. 2009. Regulation of serum amyloid A3

(SAA3) in mouse colonic epithelium and adipose tissue by the intestinal microbiota.

PLoS One 4:e5842.

85.　Peng L, He Z, Chen W, Holzman IR, Lin J. 2007. Effects of butyrate on intestinal barrier

function in a Caco-2 cell monolayer model of intestinal barrier. Pediatr Res 61:37-41.

86.　Wang HB, Wang PY, Wang X, Wan YL, Liu YC. 2012. Butyrate enhances intestinal

epithelial barrier function via up-regulation of tight junction protein Claudin-1

transcription. Dig Dis Sci 57:3126-35.

87.　Yan H, Ajuwon KM. 2017. Butyrate modifies intestinal barrier function in IPEC-J2 cells

through a selective upregulation of tight junction proteins and activation of the Akt

signaling pathway. PLoS One 12:e0179586.

88.　Peng L, Li ZR, Green RS, Holzman IR, Lin J. 2009. Butyrate enhances the intestinal

barrier by facilitating tight junction assembly via activation of AMP-activated protein

kinase in Caco-2 cell monolayers. J Nutr 139:1619-25.

89.　Xiaofeng Han HS, Yunlei Wang, Yingmo Sheng, Jie Chen. 2015. Sodium butyrate

protects the intestinal barrier function

in peritonitic mice. Int J Clin Exp Med 8:4000-7.

90.　Hu ED, Chen DZ, Wu JL, Lu FB, Chen L, Zheng MH, Li H, Huang Y, Li J, Jin XY, Gong

YW, Lin Z, Wang XD, Xu LM, Chen YP. 2018. High fiber dietary and sodium butyrate

attenuate experimental autoimmune hepatitis through regulation of immune regulatory

cells and intestinal barrier. Cell Immunol 328:24-32.

91.　Matheus VA, Monteiro L, Oliveira RB, Maschio DA, Collares-Buzato CB. 2017. Butyrate

reduces high-fat diet-induced metabolic alterations, hepatic steatosis and pancreatic

beta cell and intestinal barrier dysfunctions in prediabetic mice. Exp Biol Med (Maywood)

242:1214-1226.

92.   Kelly CJ, Zheng L, Campbell EL, Saeedi B, Scholz CC, Bayless AJ, Wilson KE, Glover

LE, Kominsky DJ, Magnuson A, Weir TL, Ehrentraut SF, Pickel C, Kuhn KA, Lanis JM,

Nguyen V, Taylor CT, Colgan SP. 2015. Crosstalk between Microbiota-Derived Short-

Chain Fatty Acids and Intestinal Epithelial HIF Augments Tissue Barrier Function. Cell

Host Microbe 17:662-71.

93.   Zheng L, Kelly CJ, Battista KD, Schaefer R, Lanis JM, Alexeev EE, Wang RX, Onyiah

JC, Kominsky DJ, Colgan SP. 2017. Microbial-Derived Butyrate Promotes Epithelial

Barrier Function through IL-10 Receptor-Dependent Repression of Claudin-2. J Immunol

199:2976-2984.

94.   Wang RX, Lee JS, Campbell EL, Colgan SP. 2020. Microbiota-derived butyrate

dynamically regulates intestinal homeostasis through regulation of actin-associated

protein synaptopodin. Proc Natl Acad Sci U S A 117:11648-11657.

95.   Dillon SM, Kibbie J, Lee EJ, Guo K, Santiago ML, Austin GL, Gianella S, Landay AL,

Donovan AM, Frank DN, Mc CM, Wilson CC. 2017. Low abundance of colonic butyrate-

producing bacteria in HIV infection is associated with microbial translocation and

immune activation. AIDS 31:511-521.

96.   Ijssennagger N, van der Meer R, van Mil SWC. 2016. Sulfide as a Mucus Barrier-

Breaker in Inflammatory Bowel Disease? Trends Mol Med 22:190-199.

97.   Haines-Menges BL, Whitaker WB, Lubin JB, Boyd EF. 2015. Host Sialic Acids: A

Delicacy for the Pathogen with Discerning Taste. Microbiol Spectr 3.

98.   Giron LB, Tanes CE, Schleimann MH, Engen PA, Mattei LM, Anzurez A, Damra M,

Zhang H, Bittinger K, Bushman F, Kossenkov A, Denton PW, Tateno H, Keshavarzian A,

Landay AL, Abdel-Mohsen M. 2020. Sialylation and fucosylation modulate

inflammasome-activating eIF2 Signaling and microbial translocation during HIV infection.

Mucosal Immunol 13:753-766.

99.   Moreau RA, Agnew J, Hicks KB, Powell MJ. 1997. Modulation of lipoxygenase activity

by bacterial hopanoids. J Nat Prod 60:397-8.

100. Tsai IJ, Croft KD, Mori TA, Falck JR, Beilin LJ, Puddey IB, Barden AE. 2009. 20-HETE

and F2-isoprostanes in the metabolic syndrome: the effect of weight reduction. Free

Radic Biol Med 46:263-70.

101. Raza GS, Putaala H, Hibberd AA, Alhoniemi E, Tiihonen K, Makela KA, Herzig KH.

2017. Polydextrose changes the gut microbiome and attenuates fasting triglyceride and

cholesterol levels in Western diet fed mice. Sci Rep 7:5294.

102. Rabot S, Membrez M, Bruneau A, Gérard P, Harach T, Moser M, Raymond F,

Mansourian R, Chou CJ. 2010. Germ-free C57BL/6J mice are resistant to high-fat-diet-

induced insulin resistance and have altered cholesterol metabolism. FASEB journal :

official publication of the Federation of American Societies for Experimental Biology

24:4948-59.

103. Velagapudi VR, Hezaveh R, Reigstad CS, Gopalacharyulu P, Yetukuri L, Islam S, Felin

J, Perkins R, Boren J, Oresic M, Backhed F. 2010. The gut microbiota modulates host

energy and lipid metabolism in mice. J Lipid Res 51:1101-12.

104. National Instiues of Health EaGRP, Nactional Cancer Institute. 2010. Diet History

Questionnaire, Verson 2.0.

105. National Cancer Institute EaGRP. 2012. Diet*Calc Analysis Program, Version 1.5.0. .

106. Shaffer M, Thurimella K, Lozupone CA. 2020. doi:10.1101/2020.11.13.380733.

107. Yang Y, Cruickshank C, Armstrong M, Mahaffey S, Reisdorph R, Reisdorph N. 2013.

New sample preparation approach for mass spectrometry-based profiling of plasma

results in improved coverage of metabolome. J Chromatogr A 1300:217-26.

108. Halper-Stromberg E, Gillenwater L, Cruickshank-Quinn C, O'Neal WK, Reisdorph N,

Petrache I, Zhuang Y, Labaki WW, Curtis JL, Wells J, Rennard S, Pratte KA, Woodruff

P, Stringer KA, Kechris K, Bowler RP. 2019. Bronchoalveolar Lavage Fluid from COPD

Patients Reveals More Compounds Associated with Disease than Matched Plasma.

Metabolites 9.

109.   Kennedy A, Bivens A. 2017. Methods for the Analysis of Underivatized Amino Acids by

LC/MS: For Food, Life Science, and Metabolomics Applications, Agilent Application

Note.

110.   Sumner LW, Amberg A, Barrett D, Beale MH, Beger R, Daykin CA, Fan TW, Fiehn O,

Goodacre R, Griffin JL, Hankemeier T, Hardy N, Harnly J, Higashi R, Kopka J, Lane AN,

Lindon JC, Marriott P, Nicholls AW, Reily MD, Thaden JJ, Viant MR. 2007. Proposed

minimum reporting standards for chemical analysis Chemical Analysis Working Group

(CAWG) Metabolomics Standards Initiative (MSI). Metabolomics 3:211-221.

111.   Ruttkies C, Schymanski EL, Wolf S, Hollender J, Neumann S. 2016. MetFrag

relaunched: incorporating strategies beyond in silico fragmentation. J Cheminform 8:3.

112.   Agilent. 2020. Lipidomics Analysis with Lipid Annotator and Mass Profiler Professional,

vol 5994-1111EN., Technical Overview.

113.   Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi

A, Gibbons SM, Ackermann G, Navas-Molina JA, Janssen S, Kopylova E, Vazquez-

Baeza Y, Gonzalez A, Morton JT, Mirarab S, Zech Xu Z, Jiang L, Haroon MF, Kanbar J,

Zhu Q, Jin Song S, Kosciolek T, Bokulich NA, Lefler J, Brislawn CJ, Humphrey G,

Owens SM, Hampton-Marcell J, Berg-Lyons D, McKenzie V, Fierer N, Fuhrman JA,

Clauset A, Stevens RL, Shade A, Pollard KS, Goodwin KD, Jansson JK, Gilbert JA,

Knight R, Earth Microbiome Project C. 2017. A communal catalogue reveals Earth's

multiscale microbial diversity. Nature 551:457-463.

114.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H,

Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ,

Brown CT, Callahan BJ, Caraballo-Rodriguez AM, Chase J, Cope EK, Da Silva R,

Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M,

Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo

J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch

AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, et al. 2019.

Reproducible, interactive, scalable and extensible microbiome data science using QIIME

2. Nat Biotechnol 37:852-857.

115.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. 2016.

DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods

13:581-3.

116.   Rognes T, Flouri T, Nichols B, Quince C, Mahe F. 2016. VSEARCH: a versatile open

source tool for metagenomics. PeerJ 4:e2584.

117.   McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL,

Knight R, Hugenholtz P. 2012. An improved Greengenes taxonomy with explicit ranks

for ecological and evolutionary analyses of bacteria and archaea. The ISME journal

6:610-8.

118.   Janssen S, McDonald D, Gonzalez A, Navas-Molina JA, Jiang L, Xu ZZ, Winker K, Kado

DM, Orwoll E, Manary M, Mirarab S, Knight R. 2018. Phylogenetic Placement of Exact

Amplicon Sequences Improves Associations with Clinical Information. mSystems 3.

119.   Shaffer M. SCNIC: Sparse Cooccurrence Network Investigation for Compositional data.

https://github.com/shafferm/SCNIC. Accessed

120.   Friedman J, Alm EJ. 2012. Inferring correlation networks from genomic survey data.

PLoS Comput Biol 8:e1002687.

1131