Systemic application of the TRPV4 antagonist GSK2193874 induces tail vasodilation in a

mouse model of thermoregulation

Fiona O'Brien, Caroline A Staunton & Richard Barrett-Jolley*

Department of Musculoskeletal Ageing, Faculty of Health and Life Sciences, University of

Liverpool, Liverpool, L7 8TX, U.K.

*Corresponding Author

Contact email: RBJ@liverpool.ac.uk

Abstract

15
In humans the skin is a primary thermoregulatory organ, with vasodilation leading to rapid
body cooling, whereas in Rodentia the tail performs an analogous function. Many
thermodetection mechanisms are likely to be involved including thermosensitive TRPV4 ion
channels. Previous studies have shown that TRPV4 can act as a vasodilator by local action in
blood vessels, and in this study we investigated whether there was a constitutive role for
TRPV4 in *mus muscularis* tail vascular tone and thermoregulation.

We measured tail blood flow and other cardiovascular parameters by pressure
plethysmography in lightly sedated CD-1 mice at a range of ambient temperatures, with and
without intraperitoneal administration of the blood brain barrier crossing TRPV4 antagonist
GSK2193874.

As expected for a thermoregulatory organ, we found that tail blood flow increased with
temperature. However, unexpectedly we found that the TRPV4 antagonist GSK2193874
increased tail blood flow.

Since arterial TRPV4 activation is known to cause vasodilation that would increase tail blood
flow, these data suggest that increases in tail blood flow resulting from the TRPV4
antagonist must arise from a site other than the blood vessels themselves, perhaps in
central cardiovascular control centres such as the hypothalamus.

33
34
35

Introduction

Thermoregulation is one of the key defining homeostatic processes common to mammals. Strict control of temperature is critical to a number of biological processes such as enzyme function and neuronal activity. Core body and brain temperatures are well maintained despite challenges such as changing ambient temperature and exercise to the degree that brain temperature rarely changes outside of a 3 °C range [1-3]. Typically, mammals detect temperatures at both central and peripheral sites and responses to changing temperatures can result both from local responses and central, hypothalamus-co-ordinated autonomic responses [4-6]. In turn, a temperature regulating effector system acts to return the core temperature (Tc) towards the set point [7], which changes with time of day [4, 8-10]. Typical thermogenic effector mechanisms include liver thermogenesis and skeletal muscle shivering whereas cooling mechanisms include counter-current mechanisms in some species, behavioural changes and redistribution of blood from core to peripheral vessels [4, 5, 7]. Brown adipose tissue (BAT) is recognised as the major site of non-shivering thermogenesis during cold exposure and is crucial for maintaining Tc and energy balance [11]. Rodents use basal metabolic rate (BMR) and non-shivering thermogenesis as their principle mechanisms for heat production, mainly because of their small size [12]. In terms of heat loss, transfer of excess heat to the environment is facilitated by so-called heat transfer zones, which are usually found at the body extremities and can be species specific, for example, humans, typically, acute heat loss is mediated by redistributing blood to cutaneous vascular beds [5]. The location of critical heat transfer zones are somewhat species specific however, so for example, the ear for elephants and rabbits [13, 14], head vasculature in large dinosaurs [15, 16], the feet [17] and the tail for rodents [18-20]. The tail of rodents is ideally suited to this role due to its glabrous nature, but unlike the similarly glabrous ear it has the necessary arterio-venous anastomoses [20]. Vasoconstriction and dilatation of the tail arteries are in turn controlled by increases and decreases (respectively) of tail sympathetic nerve activity (SNA) [21]. There is a strong correlation between each of abdominal temperature and tail SNA, skin temperature and tail SNA and brain temperature and tail SNA, but the latter has the steepest correlation slope [21].

In this work, we have investigated the role of *mus muscularis* TRPV4 in this homeostatic system using a potent and selective TRPV4 inhibitor, GSK2193874. TRPV4 is one of several temperature sensitive ion channels and expressed in both the hypothalamus and the

vasculature; in both smooth muscle and endothelial cells. TRPV4 is a relatively non-
selective Ca^{2+} channel (PCa/PNa 6-10) that was first characterised as mechanosensory [22,
23]. However, it is also activated by temperatures above approximately 30 ˚C, and so would
be expected to be constitutively active, under basal conditions [24]. Nevertheless,
activation of TRPV4 leads to vasodilation [25-27]. This is the opposite to what one might
expect for activation of a Ca^{2+}/non-specific cation channel in the vasculature, since they may
be expected to induce depolarisation which in turn induces contraction of vascular smooth
muscle [28]. The apparent anomaly can easily be explained as a relatively conserved
intracellular coupling between TRPV4 and Ca^{2+} activated potassium channels, seen in a wide
range of cells such as chondrocytes, neurones, endothelial and smooth muscle cells [25, 29-
32]. A small influx of Ca^{2+} that is insufficient to cause depolarisation but *is* sufficient to
activate nearby Ca^{2+}-activated potassium channels [33] leads to hyperpolarisation and a
reduction in over-all Ca^{2+} entry. Additionally, TRPV4 action on endothelial cells has been
proposed to induce both direct coupling to vascular smooth muscle cells (and dilatation)
and release of endothelially derived relaxant factors such as EDRF/EDHF [25, 26]. There is
therefore a consensus amongst several studies that TRPV4 activation is vasodilatory. The
logical extension of this body of evidence would be that transgenic elimination of TRPV4
(TRPV4-/- knock out) would increase blood pressure, but it does not [25]. Although counter-
intuitive data could be explained as resulting from potential compensatory changes of
expression in related ion channels, paradoxically, general linear model analyses of the
publicly available data from Sand *et al.,* 2015 [10] shows that over a 24hr period mean
arterial blood pressure is lower in TRPV4 -/- mice [34]. Furthermore, in terms of
temperature regulation, TRPV4 -/- mice do not have altered core temperature. On the
other hand, activation of TRPV4 with topological RN1747 decreases core temperature of
rattus norvegicus and increased tail vasodilatation [35]. The effects of a TRPV4 inhibitor
(HC067047) on body temperature, in the same study were mixed.
In this study we therefore had a straightforward question; does the small molecule TRPV4
inhibitor, GSK2193874, decrease tail vasodilation in response to elevated ambient
temperatures. As a surrogate for tail vasodilation we used tail blood flow measured by
volume plethysmography [36]. Surprisingly, we found the TRPV4 inhibitor had the opposite
effect, raising the possibility that the TRPV4 ion channels expressed outside of the

vasculature, for example in the central nervous system, may also be involved with rodent

thermoregulation.

Methods

Animals

Fourteen adult CD1-mice (Charles River, UK) were used, kept in our Animal Services unit with *ad libitum* access to food and water. All experimental procedures were performed under a UK Home Office licence (Number 70/8746, approved 09/11/2015) and complied with the UK Animals (Scientific Procedures) Act 1986 and were ethically approved by the University's Animal Welfare Committee (AWERB).

Volume pressure recording (VPR) plethysmography

To investigate the role of TRPV4 in the cardiovascular responses to temperature change, we used the CODA tail volume plethysmography (VPR) system (Kent Scientific, Torrington, CT, USA) on control CD1-mice and mice that had received the selective TRPV4 antagonist GSK2193874. Full details including figures are included in the supplementary materials.

Ambient temperature was increased slowly using heating pads under their housing and insulating the cage lids. Temperature was monitored by thermocouple during the experiment whilst the CODA device was continuously recording.

Statistical Analyses

Blood pressure (MAP), blood flow and heart rate (HR) statistical comparisons were made with the nlme package in R, which incorporates a repeated measures design (using R 4.1.0). For heart rate variability statistical comparions, we used MANOVA in Minitab (version19, Minitab Inc., PA,USA). $p < 0.05$ was taken as significant. The R code and data to repeat these analysis, together with the Minitab workbook are included in the Github repository for this work https://github.com/RichardBJ/TailCuff_Data_Analysis.

Drugs

Midazolam was supplied by our animal service unit, but GSK2193874 (300 µg/kg, *i.p.*) and DMSO were obtained from Sigma-Aldrich. GSK2193874 was dissolved in DMSO at 20 mg/ml stock then diluted 1:100 before i.p. injection (0.2 mg/ml). "Control" therefore includes 1% DMSO, and volume of injection was dependent upon the weight of the mouse.

Results

Using the VPR equipment we measured blood pressure (MAP), heart rate (HR) and blood

flow (Flow) in 14 animals with and without GSK2193874 over the ambient temperature

range of 31C to 36C. These are plotted in two-factor (treatment, temperature) format and

analysed with a repeated measures design. There was a statistically significant effect of

temperature on all parameters measured, MAP, HR and tail blood flow (Figure 1). I the

cases of HR and MAP there was no significant effect of treatment with the TRPV4 antagonist

(GSK2193874). However, with tail blood flow there was both a significant increase with

GSK2193874 treatment and a very highly significant interaction between temperature and

GSK2193874 treatment (Figure 1C).

Since we were able to derive beat-by-beat heart rate records for several seconds (for

example Supplementary Figure 1), we investigated whether heart rate variability (HRV)

could be captured over such a short period. To test whether this was feasible, we simulated

mouse heart rate interval records of decreasing length using a modified version of McSharry

et al., 2003 [37] and then measured HRV spectral powers by the Lomb-Scargle method [38,

39] for these over 3000 simulations. Figure 2A shows that just a few seconds of ECG are

sufficient to obtain a picture of the heart rate variability in a mouse, in so far as, increasing

the simulation duration beyond this does not greatly affect the HRV spectra. We therefore

measured HRV power in the 0.1 to 1.9Hz bands in our samples of control and GSK2193874

records (Figure 2B, C) and compared these statistically with a MANOVA model, over our full

range of temperatures. There was no over-all statistical difference with temperature,

however there was a statistically different set of spectra between control and GSK2193874

treated spectra. Furthermore, with univariate analyses, there was a significant difference

between treatment and control at each individual frequency except the 0.5Hz banding.

Discussion

In this work we investigate the role of TRPV4 in rat tail blood flow with a systemic inhibitor

of TRPV4, GSK2193874 using the CODA tail volume plethysmography (VPR) system.

Surprisingly, we find that tail blood flow is increased by GSK2193874. There was a

detectable effect also on HRV, but no significant change in blood pressure or heart rate

itself.

Using a potent drug inhibitor rather than constitutive transgenic knock down of TRPV4 has

the advantage that, realistically, in the time frame of an experiment there will be no

compensatory expression of functionally related proteins. The trade-off is selectivity. Whilst

GSK2193874 is considered highly selective for TRPV4 compared to the other 200+ proteins it

has been assayed against [40], there could still be unknown off target effects. GSK2193874

is a small lipid soluble inhibitor of TRPV4 [41] that crosses the blood-brain barrier well

(brain:plasma ratio = 0.6, personal communication with Dr David Behm of GSK) and so there

are several locations at which TRPV4 could potentially influence the control of blood flow in

response to elevated temperatures. A non-exhuastive list of possible sites of action could

include the vasculature, sensory neurones or cardiovascular control neurones; we will now

discuss each of these in turn.

The best studied of these potential sites of action is perhaps the vasculature itself. TRPV4 is

expressed in both vascular smooth muscle and the endothelial cell lining [42]. Activation of

these channels leads to clear vasodilatation. Whilst the mechanism of this vasodilatation is

complex, involving both endothelial cells and smooth muscle cells, potential release of local

transmitters and tight junctions, ultimately, small local increases of Ca^{2+} activate potassium

channels which hyperpolarize the muscle cells and allow relaxation/vasodilatation [25-27].

If there were to be no constitutive activity of TRPV4, an inhibitor would have little effect on

tail blood flow via this site of action, if however, there is constitutive activity, an inhibitor

would be expected to promote channel closure, vasoconstriction and decreased tail blood

flow. It therefore seems unlikely the increase in tail blood flow we report in this study

results from direct action on the vasculature. Furthermore, if the effect of GSK2193874

were simply on blood vessels to cause dilation, we would have expected to see an over-all

drop in blood pressure, and possibly then a reflex increase in heart rate. We saw no change

in blood pressure or heart rate, although multivariate analysis detected a small change in

short-range HRV analysis.

A second common site of action of TRPV4 channels is sensory neurones [43, 44] and have

been implicated in temperature nocifensive responses, but only in situations of heat

hyperalgesia [45]. Escape latencies (the time it takes before animals avoid a noxious stimuli)

at temperatures of 35-50°C are unaltered in TRPV4 -/- knock-out mice, but these were

delayed following induction of heat hyperalgesia. Together, these data offer no evidence

that the TRPV4 inhibitor vasodilating effect we see is likely due to modulation of sensory

neurones.

A third location of TRPV4 channels that may be of relevance is the central nervous system,

for example the hypothalamus [46]. It is known that other transient receptor potential

channels influence the cardiovascular system via changes in sympathetic activity [8, 47].

Our own work shows that TRPV4 channels are located on pre-autonomic neurones of the

hypothalamic paraventricular nucleus (PVN) and can influence cardiovascular control in

response to osmotic challenge [32, 48]. We have shown that intracerebroventricular (ICV)

injection of hypoosmotic solution decreased blood pressure but not heart rate, and this

effect was abolished with a TRPV4 inhibitor [32]. At the neuronal level, we have shown that

the action current frequency of parvocellular PVN neurones is dramatically reduced when

TRPV4 channels are blocked [48]. To date, there have been no studies that have explored

thermoregulatory roles for TRPV4 in central cardiovascular control neurones such as those

found within the PVN. The changes we observe here in flood flow, HRV and

thermoregulatory would be consistent with modulation of central control neurones.

This is the first recorded use of HRV analysis from such short segments of heart rate

recording and its measurement on non-invasive apparatus may prove a useful advance in

3Rs. We justify this in two ways; (i) since the average mouse heart rate is some 8 times that

of a human, (i) an 8s segment would be equivalent to the standard 1 minute of recording

necessary to detect higher frequency components of human ECG [49]. (ii) There is no clear

law that states how long a segment should be to faithfully recover a given frequency

component, since this is the *opposite* of the Nyquist theorem which sets the minimum

sample frequency. The landmark cardiac Electrophysiology Task-Force report stated that

"recording should last for at least 10 times the wavelength of the lower frequency" [49], but
without evidence.   Here, simulation shows that periodograms from very short segments of
ECG are similar to that of conventional 1-minute records (Figure 3A), and these data
themselves and this approach may be of field interest.

In conclusion, this whole animal study shows that a TRPV4 antagonist has a significant effect
on tail blood flow, in the context of thermoregulation, but its site of action, and the
mechanism of such modulation remain to be determined.   We also demonstrate non-
invasive measurement of frequency domain HRV Lomb-Scargle analysis from very short-
range data that may prove useful in future 3Rs friendly research.

Acknowledgements
This project received funding from BBSRC: BB/T002115/1, BB/T002115/1
BB/S008136/1 AND BB/N003020/1 awarded to RBJ.

References

[1] Fuller, A., Moss, D.G., Skinner, J.D., Jessen, P.T., Mitchell, G. & Mitchell, D. 1999 Brain, abdominal and arterial blood temperatures of free-ranging eland in their natural habitat. *Pflugers Arch* **438**, 671-680. (doi:10.1007/s004249900105).

[2] Fuller, A., Carter, R.N. & Mitchell, D. 1998 Brain and abdominal temperatures at fatigue in rats exercising in the heat. *Journal of applied physiology (Bethesda, Md. : 1985)* **84**, 877-883. (doi:10.1152/jappl.1998.84.3.877).

[3] Walters, T.J., Ryan, K.L., Tate, L.M. & Mason, P.A. 2000 Exercise in the heat is limited by a critical internal temperature. *Journal of applied physiology (Bethesda, Md. : 1985)* **89**, 799-806. (doi:10.1152/jappl.2000.89.2.799).

[4] Tan, C.L. & Knight, Z.A. 2018 Regulation of Body Temperature by the Nervous System. *Neuron* **98**, 31-48. (doi:10.1016/j.neuron.2018.02.022).

[5] Hemingway, A. & Price, W.M. 1968 The autonomic nervous system and regulation of body temperature. *Anesthesiology* **29**, 693-701. (doi:10.1097/00000542-196807000-00012).

[6] Liedtke, W.B. 2017 Deconstructing mammalian thermoregulation. *Proc Natl Acad Sci U S A* **114**, 1765-1767. (doi:10.1073/pnas.1620579114).

[7] Hammel, H.T. 1968 Regulation of internal body temperature. *Annu Rev Physiol* **30**, 641-710. (doi:10.1146/annurev.ph.30.030168.003233).

[8] Alawi, K.M., Aubdool, A.A., Liang, L., Wilde, E., Vepa, A., Psefteli, M.P., Brain, S.D. & Keeble, J.E. 2015 The sympathetic nervous system is controlled by transient receptor potential vanilloid 1 in the regulation of body temperature. *FASEB J* **29**, 4285-4298. (doi:10.1096/fj.15-272526).

[9] Lee, H., Iida, T., Mizuno, A., Suzuki, M. & Caterina, M.J. 2005 Altered thermal selection behavior in mice lacking transient receptor potential vanilloid 4. *J Neurosci* **25**, 1304-1310. (doi:10.1523/JNEUROSCI.4745.04.2005).

[10] Sand, C.A., Starr, A., Nandi, M. & Grant, A.D. 2015 Blockade or deletion of transient receptor potential vanilloid 4 (TRPV4) is not protective in a murine model of sepsis. *F1000Res* **4**, 93. (doi:10.12688/f1000research.6298.1).

[11] Mohler, F.S. & Heath, J.E. 1988 Comparison of IR thermography and thermocouple measurement of heat loss from rabbit pinna. *The American journal of physiology* **254**, R389-395. (doi:10.1152/ajpregu.1988.254.2.R389).

[12] Liu, J.-S., Wang, D.-H. & Sun, R.-Y. 2004 Metabolism and thermoregulation in three species of rodent from Northeastern China. *Journal of Thermal Biology* **29**, 177-183. (doi:https://doi.org/10.1016/j.jtherbio.2004.02.003).

[13] Phillips, P.K. & Heath, J.E. 1992 Heat exchange by the pinna of the african elephant (Loxodonta africana). *Comparative Biochemistry and Physiology Part A: Physiology* **101**, 693-699. (doi:https://doi.org/10.1016/0300-9629(92)90345-Q).

[14] Conley, K.E. & Porter, W.P. 1985 Heat loss regulation: role of appendages and torso in the deer mouse and the white rabbit. *Journal of comparative physiology. B, Biochemical, systemic, and environmental physiology* **155**, 423-431. (doi:10.1007/bf00684671).

[15] Porter, W.R. & Witmer, L.M. 2020 Vascular Patterns in the Heads of Dinosaurs: Evidence for Blood Vessels, Sites of Thermal Exchange, and Their Role in Physiological Thermoregulatory Strategies. *Anatomical record (Hoboken, N.J. : 2007)* **303**, 1075-1103. (doi:10.1002/ar.24234).

[16] Khamas, W.A., Smodlaka, H., Leach-Robinson, J. & Palmer, L. 2012 Skin histology and its
role in heat dissipation in three pinniped species. *Acta Veterinaria Scandinavica* **54**, 46.
(doi:10.1186/1751-0147-54-46).
[17] Hankenson, F.C., Marx, J.O., Gordon, C.J. & David, J.M. 2018 Effects of Rodent
Thermoregulation on Animal Models in the Research Environment. *Comp Med* **68**, 425-438.
(doi:10.30802/AALAS-CM-18-000049).
[18] Johansen, K. 1962 Heat exchange through the muskrat tail. Evidence for vasodilator
nerves to the skin. *Acta physiologica Scandinavica* **55**, 160-169. (doi:10.1111/j.1748-
1716.1962.tb02428.x).
[19] Gordon, C.J. 2017 The mouse thermoregulatory system: Its impact on translating
biomedical data to humans. *Physiol Behav* **179**, 55-66. (doi:10.1016/j.physbeh.2017.05.026).
[20] Grant, R.T. 1963 Vasodilation and body warming in the rat. *J Physiol* **167**, 311-317.
(doi:10.1113/jphysiol.1963.sp007151).
[21] Shafton, A.D., Kitchener, P., McKinley, M.J. & McAllen, R.M. 2014 Reflex control of rat
tail sympathetic nerve activity by abdominal temperature. *Temperature (Austin)* **1**, 37-41.
(doi:10.4161/temp.29597).
[22] Voets, T., Prenen, J., Vriens, J., Watanabe, H., Janssens, A., Wissenbach, U., Bodding,
M., Droogmans, G. & Nilius, B. 2002 Molecular determinants of permeation through the
cation channel TRPV4. *Journal of Biological Chemistry* **277**, 33704-33710.
(doi:10.1074/jbc.M204828200
M204828200 [pii]).
[23] Watanabe, H., Davis, J.B., Smart, D., Jerman, J.C., Smith, G.D., Hayes, P., Vriens, J.,
Cairns, W., Wissenbach, U., Prenen, J., et al. 2002 Activation of TRPV4 channels (hVRL-
2/mTRP12) by phorbol derivatives. *Journal of Biological Chemistry* **277**, 13569-13577.
(doi:10.1074/jbc.M200062200).
[24] Chung, M.K., Lee, H. & Caterina, M.J. 2003 Warm temperatures activate TRPV4 in
mouse 308 keratinocytes. *J Biol Chem* **278**, 32037-32046. (doi:10.1074/jbc.M303251200).
[25] Earley, S., Pauyo, T., Drapp, R., Tavares, M.J., Liedtke, W. & Brayden, J.E. 2009 TRPV4-
dependent dilation of peripheral resistance arteries influences arterial pressure. *Am J
Physiol Heart Circ Physiol* **297**, H1096-1102. (doi:10.1152/ajpheart.00241.2009).
[26] Campbell, W.B. & Fleming, I. 2010 Epoxyeicosatrienoic acids and endothelium-
dependent responses. *Pflugers Arch* **459**, 881-895. (doi:10.1007/s00424-010-0804-6).
[27] Filosa, J.A., Yao, X. & Rath, G. 2013 TRPV4 and the regulation of vascular tone. *J
Cardiovasc Pharmacol* **61**, 113-119. (doi:10.1097/FJC.0b013e318279ba42).
[28] Nelson, M.T. & Quayle, J.M. 1995 Physiological roles and properties of potassium
channels in arterial smooth muscle. *Am J Physiol* **268**, C799-822.
[29] Sonkusare, S.K., Bonev, A.D., Ledoux, J., Liedtke, W., Kotlikoff, M.I., Heppner, T.J., Hill-
Eubanks, D.C. & Nelson, M.T. 2012 Elementary Ca2+ signals through endothelial TRPV4
channels regulate vascular function. *Science* **336**, 597-601. (doi:10.1126/science.1216283).
[30] Nilius, B. & Droogmans, G. 2001 Ion channels and their functional role in vascular
endothelium. *Physiological Reviews* **81**, 1415-1459.
[31] Lewis, R., Asplin, K.E., Bruce, G., Dart, C., Mobasheri, A. & Barrett-Jolley, R. 2011 The
Role of the Membrane Potential in Chondrocyte Volume Regulation. *Journal of Cellular
Physiology* **226**, 2979-2986. (doi:10.1002/jcp.22646).
[32] Feetham, C.H., Nunn, N. & Barrett-Jolley, R. 2015 The depressor response to
intracerebroventricular hypotonic saline is sensitive to TRPV4 antagonist RN1734. *Front
Pharmacol* **6**, 83. (doi:10.3389/fphar.2015.00083).

[33] Fakler, B. & Adelman, J.P. 2008 Control of K(Ca) channels by calcium
nano/microdomains. *Neuron* **59**, 873-881. (doi:10.1016/j.neuron.2008.09.001).
[34] Sand, C.A., Starr, A., Nandi, M. & Grant, A.D. 2015 Blockade or deletion of transient
receptor potential vanilloid 4 (TRPV4) is not protective in a murine model of sepsis.
F1000Res **4**, 93-93. (doi:10.12688/f1000research.6298.1).
[35] Vizin, R.C., Scarpellini Cda, S., Ishikawa, D.T., Correa, G.M., de Souza, C.O., Gargaglioni,
L.H., Carrettiero, D.C., Bicego, K.C. & Almeida, M.C. 2015 TRPV4 activates autonomic and
behavioural warmth-defence responses in Wistar rats. *Acta Physiol (Oxf)* **214**, 275-289.
(doi:10.1111/apha.12477).
[36] Feng, M., Whitesall, S., Zhang, Y., Beibel, M., D'Alecy, L. & DiPetrillo, K. 2008 Validation
of volume-pressure recording tail-cuff blood pressure measurements. *Am J Hypertens* **21**,
1288-1291. (doi:10.1038/ajh.2008.301).
[37] McSharry, P.E., Clifford, G.D., Tarassenko, L. & Smith, L.A. 2003 A dynamical model for
generating synthetic electrocardiogram signals. *IEEE Transactions on Biomedical Engineering*
50, 289-294. (doi:10.1109/tbme.2003.808805).
[38] Lomb, N.R. 1976 Least-Squares Frequency Analysis Of Unequally Spaced Data.
Astrophysics and Space Science **39**, 447--446.
[39] Scargle, J.D. 1982 Studies in astronomical time series analysis. II. Statistical aspects of
spectral analysis of unevenly spaced data. *Astrophys J* **263**, 835-853.
[40] Thorneloe, K.S., Cheung, M., Bao, W., Alsaid, H., Lenhard, S., Jian, M.Y., Costell, M.,
Maniscalco-Hauk, K., Krawiec, J.A., Olzinski, A., et al. 2012 An orally active TRPV4 channel
blocker prevents and resolves pulmonary edema induced by heart failure. *Sci Transl Med* **4**,
159ra148. (doi:10.1126/scitranslmed.3004276).
[41] Cheung, M., Bao, W., Behm, D.J., Brooks, C.A., Bury, M.J., Dowdell, S.E., Eidam, H.S.,
Fox, R.M., Goodman, K.B., Holt, D.A., et al. 2017 Discovery of GSK2193874: An Orally Active,
Potent, and Selective Blocker of Transient Receptor Potential Vanilloid 4. *ACS Med Chem*
Lett **8**, 549-554. (doi:10.1021/acsmedchemlett.7b00094).
[42] White, J.P., Cibelli, M., Urban, L., Nilius, B., McGeown, J.G. & Nagy, I. 2016 TRPV4:
Molecular Conductor of a Diverse Orchestra. *Physiol Rev* **96**, 911-973.
(doi:10.1152/physrev.00016.2015).
[43] Kim, S., Barry, D.M., Liu, X.Y., Yin, S., Munanairi, A., Meng, Q.T., Cheng, W., Mo, P., Wan,
L., Liu, S.B., et al. 2016 Facilitation of TRPV4 by TRPV1 is required for itch transmission in
some sensory neuron populations. *Sci Signal* **9**, ra71. (doi:10.1126/scisignal.aaf1047).
[44] Brierley, S.M., Page, A.J., Hughes, P.A., Adam, B., Liebregts, T., Cooper, N.J., Holtmann,
G., Liedtke, W. & Blackshaw, L.A. 2008 Selective role for TRPV4 ion channels in visceral
sensory pathways. *Gastroenterology* **134**, 2059-2069. (doi:10.1053/j.gastro.2008.01.074).
[45] Todaka, H., Taniguchi, J., Satoh, J., Mizuno, A. & Suzuki, M. 2004 Warm temperature-
sensitive transient receptor potential vanilloid 4 (TRPV4) plays an essential role in thermal
hyperalgesia. *J Biol Chem* **279**, 35133-35138. (doi:10.1074/jbc.M406260200).
[46] Feetham, C.H., O'Brien, F. & Barrett-Jolley, R. 2018 Ion Channels in the Paraventricular
Hypothalamic Nucleus (PVN); Emerging Diversity and Functional Roles. *Frontiers in*
physiology **9**, 760. (doi:10.3389/fphys.2018.00760).
[47] Aubdool, A.A., Graepel, R., Kodji, X., Alawi, K.M., Bodkin, J.V., Srivastava, S., Gentry, C.,
Heads, R., Grant, A.D., Fernandes, E.S., et al. 2014 TRPA1 is essential for the vascular
response to environmental cold exposure. *Nature communications* **5**, 5732.
(doi:10.1038/ncomms6732).

[48] Feetham, C.H., Nunn, N., Lewis, R., Dart, C. & Barrett-Jolley, R. 2015 TRPV4 and KCa ion
channels functionally couple as osmosensors in the paraventricular nucleus. *Br J Pharmacol*
172, 1753-1768. (doi:10.1111/bph.13023).
[49] Malik, M. 1996 Heart rate variability: standards of measurement, physiological
interpretation and clinical use. Task Force of the European Society of Cardiology and the
North American Society of Pacing and Electrophysiology. *Circulation* **93**, 1043-1065.
376
377

Figure Legends

Figure 1. Effects of Pharmacological inhibition of TRPV4 on blood pressure, heart rate and
blood flow.

(A) Mean arterial pressure at a range of ambient temperatures in control and GSK2193874,
there was a significant association with temperature, but not drug. There was also no
significant association between temperature and drug (Mixed effects model: Temperature
F=5.34, p-val ≤ 0.05; drug F=0.38 p-val > 0.05, Drug x Temperature F=0.17, p>0.05). **(B)**
Mean heart rate across a range of ambient temperature in control and GSK2193874, over-
all there was a significant change of heart rate with temperature, but no significant
difference with drug and no significant association between drug and temperature (Mixed
effects model: Temperature F=7.37, p-val ≤ 0.05; drug F=0.68 p-val > 0.05, Drug x
Temperature F=0.23, p>0.05). **(C)** Tail blood flow across a range of ambient temperature in
control and GSK2193874. Overall, there was a statistically significant increase of blood flow
with temperature and with GSK2193874 and a significant interaction (Mixed effects model:
Temperature F=13.21, p-val ≤ 0.005; drug F=5.57, p-val ≤ 0.05, Drug x Temperature F=14.00,
p-val ≤ 0.0005). Overall, n = 14 animals.

Figure 2. Short range heart rate variability analysis. (A) Stacked Lomb periodograms for
3000 simulated ECG inter-beat interval datasets. Y-axis is the duration of the simulated ECG
record, X-axis is the frequency component of each Lomb-Scargle. The periodograms have
been normalized and scaled therefore the power (colour bar on the right) is in arbitrary
units. Below are periodogram surfaces recorded at different temperatures under control
(B), or after injection GSK2193874 **(C)**. Power given in the scale-bars. MANOVA (Minitab)
analyses show these two distributions are significantly different, Wilks lambda p<0.05.
Overall, n = 14 animals.

403
404
405
406

Figure 1

A

B   **C**

Figure 2