DNA methylation is not a driver of gene expression reprogramming

in young honey bee workers

Carlos A. M. Cardoso-Junior[1,2,*], Boris Yagound[2], Isobel Ronai[2], Emily J. Remnant[2], Klaus

Hartfelder[1], Benjamin P. Oldroyd[2,3]

[1] Departamento de Biologia Celular e Bioagentes Patogênicos, Faculdade de Medicina de

Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brasil.

[2] Behaviour, Ecology and Evolution (BEE) Laboratory, Ecology and Evolution, School of Life

and Environmental Sciences A12, University of Sydney, Sydney NSW 2006, Australia.

[3] Wissenschaftskolleg zu Berlin, Wallotstrasse 19, 14193 Berlin, Germany.

[*] **Corresponding author**: carloscardoso.bio@alumni.usp.br

Contact Information

Carlos A. M. Cardoso-Júnior – carloscardoso.bio@alumni.usp.br; Phone: +55 16 3315 3063

Boris Yagound - boris.yagound@sydney.edu.au

Isobel Ronai – isobel.ronai@sydney.edu.au

Emily Remnant - emily.remnant@sydney.edu.au

Klaus Hartfelder - klaus@fmrp.usp.br

Benjamin Oldroyd – benjamin.oldroyd@sydney.edu.au

Author contributions

CAMCJ conceived the study, performed the field work, carried out the dissections, laboratory
molecular work and bioinformatics analyses, and wrote a first draft of the manuscript and
revised the subsequent versions; BY contributed to the design of the study, helped with
bioinformatic analyses, performed field work and statistical analyses. ER participated in the
design of the study and performed bioinformatics analyses of amplicon sequencing. IR helped
with the dissections and participated in the gene expression analyses. KH conceived the study
and supervised the experiments. BPO participated in the design the study, performed the
artificial inseminations and field work, supervised the experiments and contributed to the
statistical analyses. All authors made critical contributions during the writing of the manuscript
and gave final approval for publication.

Disclosure declaration: There are no competing interests.

Running title: Social environment does not affect DNA methylation

Keywords: honeybees, whole genome bisulfite sequencing, amplicon sequencing, DNMT,

Apis mellifera

This PDF file includes:
Main Text
Figures 1 to 5

Abstract

Intragenic DNA methylation, also called gene body methylation, is an evolutionarily-conserved epigenetic mechanism in animals and plants. In social insects, gene body methylation is thought to contribute to behavioral plasticity, for example between foragers and nurse workers, by modulating gene expression. However, recent studies have suggested that the majority of DNA methylation is sequence-specific, and therefore cannot act as a flexible mediator between environmental cues and gene expression. To address this paradox, we examined whole-genome methylation patterns in the brains and ovaries of young honey bee workers that had been subjected to divergent social contexts: the presence or absence of the queen. Although these social contexts are known to bring about extreme changes in behavioral and reproductive traits through differential gene expression, we found no significant differences between the methylomes of workers from queenright and queenless colonies. In contrast, thousands of regions were differentially methylated between colonies, and these differences were not associated with differential gene expression in a subset of genes examined. Methylation patterns were highly similar between brain and ovary tissues and only differed in nine regions. These results strongly indicate that DNA methylation is not a driver of differential gene expression between tissues or behavioral morphs. Finally, despite the lack of difference in methylation patterns, queen presence affected the expression of all four DNA methyltransferase genes, suggesting that these enzymes have roles beyond DNA methylation. Therefore, the functional role of DNA methylation in social insect genomes remains an open question.

Introduction

DNA methylation is a reversible chemical modification of DNA whereby methyl groups are added to cytosines in CpG dinucleotides by enzymes of the DNA methyltransferase family (DNMTs). In mammals DNA methylation plays important roles in regulating gene expression, including inactivation of the X chromosome in females, transposon suppression and genomic imprinting, a mechanism by which parents influence gene expression in offspring (1). The role of DNA methylation in genomic regulation of invertebrates is less clear, in part because the two main model species, the fruit fly *Drosophila melanogaster* and the nematode *Caenorhabditis elegans,* either lack methylation (*C. elegans*) or have extremely low and transient levels of methylation (*D. melanogaster*) (2, 3). In contrast, the honey bee (*Apis mellifera*) genome project revealed a functional epigenetic system comparable to that of vertebrates (4, 5). Since this discovery the honey bee has emerged as a model species for epigenetics studies in invertebrates, as, in contrast to mammals, the methylation marks in honey bee genomes are sparse and mostly restricted to gene bodies (intragenic DNA methylation) (2, 6, 7). This sparseness brings a technical advantage to researchers, as it facilitates the study of gene body methylation and its potential roles in the regulation of gene expression without noise from methylated cytosines of other genomic compartments, such as promoters and transposons. Mechanistically, it has been proposed that gene body methylation modulates the affinity of cofactor binding in regulatory DNA methylation-dependent regions in order to regulate RNA polymerase II activity (8–11).

In honey bees and other social insects, gene body methylation has been associated with behavioral and phenotypic plasticity (reviewed in (11, 12)). For example, methylome differences have been found between queens and workers in honey bees (6, 13) and ants (7). Differences in gene body methylation were also associated with division of labor among ant (7, 14) and bee (15) workers. Furthermore, DNA methylation has been associated with several

biological processes, including aging, reproduction, aggressiveness, response to social stimuli,

memory and learning, and the haplodiploid sex determination system (15–25).

In support of the hypothesis that gene body methylation of honey bees modulates

transcription, the RNA interference-mediated knockdown of the *DNA methyltransferase 3*

(*Dnmt3*) gene, which codes for the enzyme responsible for *de novo* DNA methylation, affected

14% of the honey bee worker transcriptome (9). RNA splicing was also affected, corroborating

previous *in-silico* predictions that associated DNA methylation with alternative splicing (6, 7,

13, 15, 26–29). Even more spectacularly, the knockdown of *Dnmt3* in young female larvae

resulted in adults with a queen-like phenotype, mimicking the transcriptional program induced

by a royal jelly diet (16).

More recent studies, however, now cast doubt on the role of gene body DNA

methylation as a flexible regulator of gene expression in social insects, and an emerging

consensus is that DNA methylation is genotype-specific (30–41). For example, a reanalysis of

the *Dnmt3* knockdown data of Li-Byarlay et al. (2013) suggested that the original analysis had

overestimated the number of regulated genes, and that the *Dnmt3* knockdown had in fact only

a minor effect on the honey bee gene body methylation pattern, and hence, on gene expression

(37). Furthermore, recent studies have not provided support for the proposed association

between differential methylation and alternative splicing in honey bees (38) and other social

insects (34, 42–44).

Particularly compelling evidence that DNA methylation tends to be sequence-specific

rather than representing a flexible gene regulatory mechanism comes from the clonal raider ant

(*Ooceraea biroi*) (31). *O. biroi* reproduces asexually, and therefore allows experiments in a

uniform genetic background (45). Libbrecht et al. (2016) showed that DNA methylation is not

associated with reproductive and asexual stages in *O. biroi* workers, casting doubt on its role

in other ants and honey bees. These authors proposed that the previously reported differential

methylation patterns seen between experimental groups may have be an artefact arising from

combinations of colony-specific methylation patterns.

In this study, we test the hypothesis that differential methylation is associated with

differential gene expression in response to environmental change, versus the alternative

hypothesis that differentially methylated regions are a colony and/or individual-specific

character that does not vary in response to environmental change. To do this we compared the

gene expression and methylation patterns in the brains and ovaries of young honey bee workers

as they matured in the presence or absence of their queen. Honey bee queens, through their

mandibular gland pheromones, influence the behavioral maturation and reproductive capacity

of workers (46–48), and this process involves changes in the expression of hundreds of genes

in the brain and ovary (49–52), including the *Dnmt* genes (22, 53). Furthermore, instead of

generating data from whole body methylomes, we compared methylation patterns in the tissues

(brain and ovary) most likely to respond to the absence/presence of a queen. This allowed us

to further examine whether differential methylation is related to tissue function, social context

or to genotype.

Results

Social context does not affect the brain and ovary methylomes of young honey bee

workers

We sequenced and analyzed at single base-pair resolution the complete methylomes of brains

and ovaries of honey bee workers reared in queenright or queenless colonies. After removing

adaptors and reads of low quality, and aligning the methylomes to the honey bee reference

genome (54) we obtained high coverage in all sequenced samples (Supplemental Table S1).

The conversion rate of bisulfite treatment was above 99.4% (Supplemental Table S1),

indicating a low frequency of false-positive methylated CpGs. The observed frequency of

methylated CpGs (~1% - Supplemental Table S2) is consistent with previous studies on honey

bees and other holometabolous insects (6, 7, 15, 55).

With these data we were able to ask whether the methylomes of workers differ when

they are in a queenright (QR) or queenless (QL) social context, especially in the two tissues

that are known to respond transcriptionally to the presence of the queen: the brain and the ovary

(49–51). Comparing three pairs of QR colonies with three pairs of QL colonies we found 879

differently methylated regions (DMRs) for the brains of QL and QR workers (Supplemental

Table S3) and 376 DMRs for the ovaries (Supplemental Table S4). However, we found that

the number of DMRs shared by two or more colonies was very low for both tissues (Fig. 1A).

This indicates that the majority of candidate DMRs previously identified (e.g., Supplemental

Table S3 and S4) were driven by between-colony variability and do not reflect a reproductible

effect of 'social context'.

Hierarchical clustering of worker methylomes showed that samples from the same

colony cluster together, irrespective of social context (Fig. 1B). When comparing the number

of DMRs in the methylomes of the three colonies regardless of social context (i.e., A *vs.* B, A

vs. C, and B *vs.* C), we found over 10,000 significant DMRs in the pairwise colony comparisons

(Supplemental Fig. 1). Therefore, the effect of 'colony' on the worker methylomes is of much

greater magnitude than the effect of 'social environment', if there is such an effect.

For a more in-depth analysis, we next selected three top-ranked candidate DMRs for

both tissues from the QR *vs.* QL comparison for amplicon sequencing (Fig. 1C, Supplemental

Table S5). Bisulfite-treated DNA extracts from the same colonies as those used for WGBS

(colonies A-C) were PCR-amplified and sequenced in a high-throughput platform. To increase

sample size and check for data reproducibility we also added samples from three new pairs of

colonies (colonies D-F) to this analysis. The high-coverage obtained by amplicon sequencing

(average >10,000 reads/region, range 672-30,823, Supplemental Table S5) suggests that the

majority of cell types present in the brain and ovary tissues are represented in this dataset.

Amplicon sequencing revealed that the methylation differences (QR *vs.* QL) observed in the

original WGBS data (Fig. 1C – grey bars) did not reach the differential methylation threshold

set at 10%, despite the increased coverage for the original samples (Fig. 1C – blue bars,

Supplemental Fig. 2; Supplemental Table S5), or samples from the three new independent

colonies (Fig. 1C – yellow bars). Importantly, the differential methylation pattern for the

majority of regions (five out of six) from colonies D-F was in the opposite direction to that of

colonies A-C. Together, these comparisons provide strong support for the hypothesis that the

differences seen in the WGBS comparisons of the QR *vs.* QL workers were indeed a result of

strong differences in the methylomes of colonies A-C, and not a consistent consequence of

social context, either in the original colonies, nor in the three new colonies.

Differential methylation is not associated with differential gene expression when

comparing different colonies and social context

To further investigate whether DNA methylation-mediated transcriptional responses are

triggered by queen exposure, we determined the expression patterns of nine genes displaying

DMRs found in the contrast of the QR *vs.* QL methylomes (Supplemental Tables S3, S4). We

found that only three differentially methylated genes (*GB42836*, *GB49839* and *GB54664*),

were differentially expressed (Fig. 2, Supplemental Table S6), and there was no differential

expression for the other six genes in the respective tissue displaying differential methylation.

This lack of correlation between differential methylation and differential expression was also

statistically confirmed (Supplemental Fig. S3, Pearson correlation r = 0.37, p = 0.23, n = 12).

Taken together, although major transcriptional responses to the presence or absence of a queen

in a colony are regularly observed and reported for the brain and ovary of workers (49–52), our

data indicate that there is no direct association between the DNA methylation status of a gene

and its differential expression, even in the context of a very dramatic contrast in social context.

Given that the genome methylation differences appeared to be much more related to

colony genotype than to social context (Fig. 1 and Supplemental Fig. 1), we next asked whether

colony-specific methylation patterns are associated with differential gene expression between

colonies. To do so, we first estimated the "colony effect" by ascertaining how many regions

showed different methylation levels in the brain and ovary of workers, regardless of the social

context (Supplemental Fig. S1). We found large numbers of DMRs in both the brain (n = 3,603)

and ovary (n = 1,997) that differed in their methylation level by at least 10% when comparing

the same genomic region pairwise among the colonies (Fig. 3A). This indicates that there is a

core set of genomic regions that are hypervariable with respect to their methylation status.

Interestingly, these regions are not chromosome specific (Supplemental Fig. S4), suggesting a

colony-specific methylation fingerprint across all chromosomes. Also, almost 60% of the

hypervariable regions (n = 1,194) seen in the ovaries were the same as those found in the brains

(Fig. 3B), suggesting that these colony-specific signatures are independent of tissue.

Next, we asked whether changes in DNA methylation were associated with differential

gene expression between colonies. For these analyses we selected 11 genes that were either

hypermethylated or hypomethylated in one source colony, but methylated in the opposite

direction in the other two source colonies (e.g., hypermethylated in colony A but

hypomethylated in colonies B and C). We confirmed the methylation pattern of these regions

by amplicon sequencing of colonies A-C (Fig. 3C, Supplemental Table S5) and D-F

(Supplemental Fig. S5, Supplemental Table S5). We noticed that the methylation patterns of a

given genomic window were reasonably stable across different social contexts and tissues

(Supplemental Figs. S2, S6). These analyses revealed strong colony-specific methylation

patterns for all six analyzed colonies. Thus, if DNA methylation plays a role in regulating gene

expression, these genes would be strong candidates for differential gene expression, as their

differential methylation across colonies was much greater than the differences we observed for

social context.

Only one gene (*GB50283*) out of the 12 genes assessed, which was strongly methylated

in colony C compared to the other two colonies (Fig. 3C), turned out to be differently expressed

among colonies (Fig. 3D, $p < 0.01$, Supplemental Table S7). For the other genes assessed, we

found that even clear differences in DNA methylation profiles between colonies (Fig. 3C) did

not affect their expression (Fig. 3D). After all these comparisons between different colonies

and radically different social contexts, we conclude that there is little evidence in support of

the hypothesis that differences in DNA methylation drive alterations in gene expression in the

two tissues that are most likely to respond to social context. Hence, the brain and ovary

methylomes of young honey bee workers are primarily a manifestation of colony identity rather

than a mediator between social environmental changes and gene expression.

Brain and ovary methylomes are highly similar, despite the functional differences

between the two tissues

After showing that DNA methylation of young honey bee workers is colony specific rather

than influenced by the social context, we investigated whether there may be tissue-specific

differences between the brain and ovary methylomes. Comparing the methylomes of tissues

with completely distinct biological functions, we expected to identify hundreds of DMRs

associated with the transcriptomic differences between these two tissues. Surprisingly, we

found only nine DMRs across all three source colonies when comparing the methylomes of the

two tissues, irrespective of social context (Fig. 4A). These regions are associated with only

four genes: *GB47277*, *GB51802*, *GB55278*, and *GB50784*. The amplicon sequencing analysis

performed for two of these DMRs confirmed the differences originally seen in the WGBS data

for these two tissues (Fig. 4B and Supplemental Fig. S7).

When testing whether alterations in gene expression between the brain and ovaries were

associated with differences in DNA methylation levels we found that three of the four

differentially methylated genes were also differentially expressed (Fig. 4C, Supplemental

Table S6). However, differential expression between tissues were identified for 11 out of the

12 genes previously analyzed for colony specificity (Fig. 3D, Supplemental Table S7). Thus,

it is not yet clear whether the alterations in methylation level promotes causative alterations in

gene expression between the two tissues. From an overall perspective the methylomes of brains

and ovaries were strikingly similar, with significant differential methylation seen for only four

genes.

Expression of the four *Dnmt* genes is influenced by social context

As our data strongly indicate that the presence or absence of a queen does not drive significant

alterations in the gene body methylation patterns in the brain and ovaries of young honey bee

workers (Fig. 1) we wondered what the expression patterns of the genes encoding the four

honey bee DNA methyltransferases would look like, especially since previous studies have

used the expression of *Dnmt* genes as an indicator of epigenetic events in social insects (16,

18, 20, 23, 25, 56) . Furthermore, we have recently reported that the queen mandibular

pheromone upregulated the expression of three *Dnmt* genes (*Dnmt1b*, *Dnmt2* and *Dnmt3*) in

the brain of caged honey bee workers (53). Here we report that the expression of all four *Dnmt*

genes predicted in the *A. mellifera* genome is affected by the presence or absence of a queen in

the colony (Fig. 5, Supplemental Table S6), either in the brain or the ovary. Interestingly, the

same social cue (presence/absence of a queen) that upregulated two *Dnmt* genes (*Dnmt1b* and

Dnmt3) in the brain of young workers had an opposite effect on the expression of three *Dnmt*

genes (*Dnmt1a*, *Dnmt2* and *Dnmt3*) in the ovary. This suggests that expression of the *Dnmt*

genes responds to social cues in both tissues, even though their differential expression

apparently is not associated with alterations in the methylomes of honey bee workers.

266
267 **Discussion**
268
With the presence or absence of a queen being one of the strongest contrasts in the social

context of a colony, and with DNA methylation as a strong candidate for mediating adaptive

responses in the individual colony members, we had expected to see signatures of this social

context condition in the methylomes of young honey bee workers. However, all the evidence

we obtained suggests the opposite: that gene body methylation is not a major regulator of gene

expression reprogramming in young honey bees. First, the contrast between the presence *vs.*

absence of a queen was not reflected in significant alterations in the respective methylomes,

indicating that differential DNA methylation is not a mediator for the transcriptional

consequences of queen presence. Second, despite strong differences in gene expression

between brain and ovarian tissues (57), we found that their methylomes are very similar. Third,

thousands of DMRs were identified when comparing individuals from different colonies.

However, even when differences in methylation were greater than 50%, there was no

association between DNA methylation and gene expression, except for one gene (*GB50283*,

Fig. 3D). Therefore, these data provide strong support for the hypothesis that intragenic CpG

methylation is not a driver of gene expression reprograming in the brain and ovary of young

honey bees. This result is actually in line with and likely generalizable for other social insects

(31, 32, 34, 41, 43, 44, 58).

In contrast to our findings, two previous studies have suggested that honey bee workers

kept under different social conditions exhibit a low number of DMRs, and that these are

associated with differential gene expression (15, 17). A plausible explanation (32, 39) for these

contrasting results is that previously-observed differences in the methylomes of honey bee

workers in response to divergent social stimuli arose as an artifact of genotype-associated

methylation variants (33). Indeed, when the colony genetic background was standardized, no differences were found between the methylomes of newly-emerged queens and newly-emerged workers (15). Furthermore, ultra-deep analyses on the dynamics of brain DNA methylation of *Dynactin*, *Nadrin* and *PKCbp1* genes, indicated that their methylation content are not altered over time, and thereby not implicated in honey bee workers' aging and behavioral maturation (59). Thus, we conclude that the methylomes of honey bee workers do not respond dynamically to changes in the social milieu (e.g., loss of the queen), whereas gene expression levels do so (49–53).

Our results also show that DNMT-encoding genes are differentially expressed, and thus affected by the presence or absence of a queen. This finding is consistent with previous honey bee studies that showed an effect of queen mandibular pheromones on the expression of *Dnmt* genes (22, 53). However, these alterations seen in the expression of the *Dnmt* genes apparently does not promote a "real-time" epigenetic reprogramming process (this study; Harris et al. 2019), indicating that the expression levels of *Dnmt* genes should not be used as a proxy of possible differences in DNA methylation patterns or levels. This result further highlights the weak association between the expression of *Dnmt* genes, DNA methylation, and gene expression reprogramming seen in honey bees and other insects. For instance, even though the genome of the red flour beetle *Tribolium castaneum* is not methylated (60, 61), the knockdown of a *Dnmt1* ortholog resulted in offspring showing developmental arrest and high mortality (61). Similar developmental defects in response to a knockdown of *Dnmt1* were observed in the reproductive system of females of the milkweed bug *Oncopeltus fasciatus* (37, 62). These phenotypic alterations are not a consequence of transcriptomic changes, even though the *Dnmt1* knockdown successfully reduced the global levels of DNA methylation (37). Combined, these results imply that DNMTs have other, not yet understood functions, that go beyond DNA methylation and its maintenance.

After an unequivocal demonstration that colony-specific methylation patterns are major

signatures in the methylomes of honey bee workers (Fig. 3), an interesting question emerges:

why do individuals from different colonies exhibit such contrasting differences between their

methylomes? While with our current data we are not yet able to fully solve this puzzle, we

propose that colony-specific methylation patterns are both functional and genotype-associated;

with a minimum effect on transcription (37, 38). Our results show that colony-specific

methylation patters are predominantly chromosome and tissue independent (Fig. 3B, Fig. 4.

and Fig. S4). Therefore, the functional role of intragenic methylation in honey bees is likely to

be associated with basal processes necessary for cell viability, e.g., cell division, genome

integrity, and/or cell cycle progression (37, 61–63), rather than representing as a marker of

tissue identity (Fig. 4), or acting as a dynamic sensor of changes in the social environment (Fig.

1). The finding that the knockdown of the *Dnmt3* gene function, a key orchestrator of honey

bee development, is lethal at early development (16) suggests that DNA methylation is

essential to honey bee viability. Furthermore, the colony-specific methylation patterns are, at

least in part, sequence-specific (32, 39, 40). Ultra-deep amplicon sequencing analyses

identified that a honey bee brain displays only a small repertoire of methylation patterns. This

may reflect the small number of maternal and paternal epialleles predicted for polyandrous

insects (59). Interestingly, the colony-specific methylation patterns are inherited from drones

to their worker daughters (40), suggesting a role for the paternally inherited epialleles to the

composition of colony-specific worker methylomes.

In conclusion, all evidence generated in this study goes against the hypothesis that gene

body methylation is a driver of gene expression re-programming in adult honey bee workers.

With our experimental design, using several colonies of standardized within-colony genetic

backgrounds and high-throughput genomic sequencing approaches we had originally expected

to provide evidence in favor of this hypothesis and to see defined differentially-methylated

gene sets between queen states, with correlated expression patterns. As it has turned out, the

conclusion to be drawn is that gene body methylation patterns are essentially colony specific

and unaffected by even radical changes in social context. Nonetheless, it is possible that

intragenic DNA methylation plays a role in gene expression during other stages of a worker's

life cycle. If so, this is likely be restricted to a small set of genes under specific contexts (16,

35, 37, 38), including the *GB51802*, *GB55278* and *GB50784* genes that show both tissue-

specific methylation patterns and differential expression (Fig. 4C). Our results make it clear

that, whatever may be the specific function of gene body DNA methylation in the honey bee,

it is now undeniable that there is a strong genotype effect shaping the workers' methylomes,

and that future studies need to consider the possible effect of genotype variants (colony and

patriline genotypes) in their experimental design. In this sense, we believe that our work makes

a significant contribution to the understanding of the meaning of the enigmatic, but

evolutionarily-conserved intragenic DNA methylation in an important invertebrate epigenetic

model species, the honey bee. As it stands now, we must say that the functional role of gene

body methylation in social insects is still an unsolved issue.

Material and methods

Bees and manipulation of the social environment

We used workers from six source colonies. These colonies were headed by queens of standard

commercial stock (mostly *A. m. ligustica*) each of which had been instrumentally inseminated

with sperm from a single male (64). Hence, the workers within each colony pairs were all full

sisters, with a genetic relatedness of 0.75, thus minimizing genotypic heterogeneity within

colonies.

Queenright and queenless colonies were prepared by splitting each of the six host

colonies into two (n = 12), thus generating colony pairs of the same genetic background but

differing completely in terms of social context. Brood frames were equalized among the colony

pairs to ensure a similar condition in terms of brood presence, despite the presence or absence

of the queen. After splitting the host colonies into queenless and queenright halves at a remote

apiary, we moved the splits to the apiary at the University of Sydney, thus reducing the

tendency of the queenless workers moving back to the respective unit with the queen. The

experiments were conducted during the southern hemisphere summers of 2017 (source

colonies A-C) and 2018 (source colonies D-F).

Newly-emerged workers from source colonies A-F were obtained by placing sealed

brood frames overnight in separate boxes, in an incubator at 34 °C on the day of colony

splitting. The newly-emerged workers were divided into two groups (n = 200 bees per group),

paint marked with different colors, and then introduced into their respective pair of queenright

and queenless host colonies. After four days, the marked workers were collected, snap frozen

on dry ice, and stored at -80 °C. At this age workers are known to respond transcriptionally to

the presence or absence of a queen (49, 53). As expected, the queen was seen in all QR host

colonies at the time of sampling, but never in the QL host colonies. Samples from six of these

colonies (colonies queenright A-C and queenless A-C) were initially used for methylome

sequencing and subsequently also for the region-specific analyses, whereas the other six

colonies (colonies queenright D-F and queenless D-F) were used for the region-specific

analyses only, which served to validate and confirm the reproducibility of the methylome

results. The brains and ovaries of the workers were dissected as described elsewhere (53, 65).

Whole Genome Bisulfite Sequencing (WGBS) and Amplicon Sequencing

We first performed Whole Genome Bisulfite Sequencing (WGBS) to identify which genes

were differentially methylated in the samples from the first three queenright and queenless

colonies (colonies A-C). For each sample we pooled either eight brains or 20 pairs of non-

activated ovaries. Pooling was necessary to obtain a sufficient DNA yield for high-throughput

sequencing. This resulted in a total of 12 samples, represented by six brain and six ovary

samples for each of the two social contexts (three queenright and three queenless). DNA was

extracted with the DNeasy Blood and Tissue Kit[TM] (Qiagen) and quantified with a Qubit 2.0

Fluorometer system (Invitrogen). Before sequencing, 0.01% (w/w) of unmethylated Lambda

DNA (Promega) was added to each sample to be later used for calculating the bisulfite

conversation efficiency. For WGBS, the DNA samples were sent to the Beijing Genomics

Institute (China), where library construction, bisulfite treatment, and sequencing were

performed. Bisulfite treatment was done with the EZ-DNA methylation kit (Zymo Research),

and paired-end WGBS was performed on an Illumina NovaSeq platform. Each library was

sequenced twice in two separate lanes for high coverage. In the bioinformatics analysis, the

data from the two lanes were then merged, as we did not detect relevant differences between

the two runs. Data on coverage for each sample are given in Supplemental Table S1.

After processing the WGBS data and identification of DMRs, we performed a paired-

end ultra-deep amplicon sequencing analysis (59, 66) for 13 of the DMRs revealed by WGBS,

using a total of six colony pairs, the original three pairs (source colonies A-C) plus three new

colony pairs (source colonies D-F). DNA extractions, bisulfite conversion, and Lambda DNA

spiking was performed as described above. Columns were eluted with 20 μL of ultrapure water,

and 1 μL of the eluted solution was used as template in the PCR assays. Bisulfite PCR primers

were designed to amplify fragments between 140-300 bp of the forward strand of each

differentially methylated region and the control Lambda spike (Supplemental Table S8).

Bisulfite-treated DNA was amplified using the KAPA HiFi Uracil[+] Kit[TM] (Roche). PCR assays

were set up with 5 μL of Kapa Master Mix, 0.3 μL of each primer (forward and reverse – 3

pmol/reaction), 1 μL of bisulfite-treated DNA and 3.4 μL of water for a total reaction volume

of 10 μL, and performed with the annealing temperatures of the respective primers

(Supplemental Table S8). For multiplexing of the samples into two libraries, Nextera barcodes

were added to the 5' ends of all primers. Amplicons from different primers were pooled to

generate a separate library for each sample (24 samples in total: queenright and queenless

brains and ovaries from each of the source colonies A-F). Samples were purified, and Nextera

paired-end libraries were constructed at the Australian Genome Research Facility (AGRF).

Libraries were constructed in duplicate for each sample and sequenced (150 bp paired end) in

a single flow cell of an Illumina MiSeq platform. Data from the two libraries were merged for

downstream analyses.

Differential methylation analyses from WGBS data

Quality   of   the   raw   data   was   assessed   by   FastQC   0.11.8

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc) and reads with quality scores <

20   were   removed.   Trimming   was   performed   with   TrimGalore   0.5.0

(www.bioinformatics.babraham.ac.uk/projects/trim_galore/) with a stringency error of 2 bp.

Overall, we removed ~ 0.5% of the total sequencing reads after checking read quality and

trimming of the adaptors (Supplemental Table S1). The remaining reads were mapped onto the

honey bee reference genome assembly Amel_4.5 (54) using Bismark 0.16.1 (67) and Bowtie

2 2.3.5.1v (68). The honey bee genome assembly Amel_4.5 was the newest honey bee genome

version at the time the bioinformatic analyses were performed. Coverage varied between 27-

46 times across samples (Supplemental Table S1). Methylation calling was performed with

Bismark software. We used 10-times coverage as a threshold for adequate sequencing coverage

of each cytosine as in Herb et al. (2012), to make our data sets comparable. Methylation levels

were assessed by the C/T ratio of converted cytosines to unconverted bases (69). Significantly

methylated sites were identified using a binomial probability model that takes into account the

bisulfite conversion rate for each sample (Supplemental Table S1) as the probability of success,

followed by Bonferroni corrections at the 1% significance level using BWASP (34). We

removed methylated CpGs with > 500x coverage to avoid PCR-based bias in the analyses.

MethylKit (70) was used for differential methylation analysis. First, the honey bee

genome was partitioned into 200 bp sliding windows (step size = 100 bp). Only windows

containing at least four sufficiently covered CpGs (two in each strand) were analyzed. A

difference threshold of 10% in the methylation level between pairwise comparisons was

applied (Herb et al. 2012). A threshold >10 methylated cytosines (sum for all CpGs inside

given window) in at least one of the libraries was used to reduce methylome complexity. The

list of DMRs was then FDR-corrected and a q-value <0.01 was considered significant. A gene

containing at least one DMR was defined as a differentially methylated gene. Gene annotation

was performed with Homer 4.9.1 software (71). Analyses were performed in the *R* environment

(R Core Team 2018). Hierarchical clustering distances between methylome samples were

determined with the R package "pvclust" (73) .

Differential methylation analysis of amplicon sequencing data

Reads were checked for quality using FastQC 0.11.8

(www.bioinformatics.babraham.ac.uk/projects/fastqc), followed by trimming of adaptors and

removal of low quality reads (Phred score < 20) using Trimmomatic (74). Between 85-90% of

the reads were retained for each library. Bissulfite-converted DNA sequences of the 13

amplicon regions of interest and the Lambda control sequence were used as templates to

generate a Bowtie2 index prior to alignment. Data from each of the two duplicate libraries for

each sample were aligned with Bowtie2 2.3.5.1v (68) using paired-end default parameters and

then converted to BAM files with Samtools (75). BAM files were imported into Geneious

software 10.2.4 (76), and alignments for each amplicon were manually checked for each

sample. C-to-T variant frequencies were calculated using the 'Find variant' function for all CG

sites, with a minimum coverage of 50 and a minimum variant frequency of 2%. Additional

SNP variants that were still visible after bisulfite conversion, such as G-to-A polymorphisms,

were also recorded. The overall cytosine methylation frequency was determined for each of the

four treatment groups (QR brain, QL brain queenless, QR ovary, QL ovary) for all of the 13

amplicons by dividing the total amount of C (methylated cytosines) per the total amount of

C+T (total amount of methylated and unmethylated cytosines). The results are presented as

percentages, and when appropriate, regions were compared in relation to their colony of origin,

social context, and tissue type.

Gene expression analysis

Each sample consisted of four pairs of non-activated ovaries or one brain (n = 8 per source

colony and social context combination). Ovaries needed to be pooled to obtain a sufficient

amount of RNA. Brains and ovaries were macerated in TRIzol (Invitrogen). Total RNA was

extracted using the Direct-zol^{TM} RNA^{TM} Miniprep kit (Zymo Research) according to the

manufacturer's instructions. Samples were treated with Turbo DNase^{TM} (Thermo-Fisher

Scientific), and RNA concentrations were determined using a Qubit 2.0 Fluorometer system

(Invitrogen). RNA samples were diluted with ultrapure water to a final concentration of 40

ng/µL (brain) and 15 ng/µL (ovary). We used 142.5 ng of ovary RNA and 600 ng of brain

RNA to synthesize cDNA using the SuperScript^{TM} III Reverse Transcriptase Kit (Invitrogen)

with Oligo(dT) primer (Invitrogen). Ovary cDNA was diluted to 2 ng/µL due to its lower

concentration, while brain cDNA was diluted to 5 ng/µL in ultrapure water.

Relative expression was determined by RT-qPCR assays for total of 20 genes, 12 being

hypermethylated or hypomethylated in one source colony, four genes whose methylation

patterns differ between tissues and four *Dnmt* genes. Assays were set up with 2.5 µL

SsoAdvanced^{TM} Universal SYBR$^®$ Green Supermix (Bio Rad), 1.25 pmol of each primer, 1

µL diluted cDNA in a total volume of 5 µL using a CFX384 Real-Time System (Bio-Rad). For

each sample we conducted three technical replicates and used their mean as the data point.

Cycle conditions were as follows: 95 °C for 10 min followed by 40 cycles of 95 °C for 10 s,

annealing temperature (Supplemental Table 8) for 10 s and 72 °C for 15 s. At the end of the

PCR cycles a melting curve analysis was run to confirm a single amplification peak. Relative

gene expression analyses were performed using the a formula that accounts for primer's

efficiency: $E^{CqMin-CqSample}$ – where "E" (Table S3) is the efficiency of primers, "CqMin" is the

lowest Cq value for a given gene and "CqSample" is the Cq of that sample (65, 77, 78). Then,

two reference genes (*Rp49* [also known as *Rpl32*] and *Ef1α*) were used to normalize the

expression levels of the target genes. These control genes have been previously validated for

honey bee quantitative PCR analyses (79) and were stable in our analysis according to the

BestKeeper software (80). Primer efficiencies (Supplemental Table 8) were calculated based

on an amplification curve of 10 points obtained through serial dilution of mixed cDNA

samples. The list of primers used is given in Supplemental Table S8. Specificity of the

respective amplification products was validated by Sanger sequencing (Macrogen, South

Korea).

Statistical analysis

Gene expression levels were analyzed as the dependent variable using a Generalized Linear

Mixed Models (GLMM) with 'colony' as random effect, and 'social context' and 'tissue' as

fixed effects. We used a log link function to all genes to approximate gene expression data to

a Gaussian distribution, which was checked by analyzing the residuals' distribution.

Alternative link functions and data transformations were applied as necessary (see

Supplemental Table S6). Given that the 'social environment' (presence/absence of the queen)

might influence gene expression in opposite directions in different tissues, as seen for Dnmt's

expression, we performed Tukey's post-hoc tests for all gene/tissue combinations. To

investigate whether colony of origin influenced gene expression, 'colony' and 'tissue' were

treated as fixed effects and 'social environment' as a random effect. Tukey's post-hoc tests

were performed to identify differences in gene expression between individuals from different

colonies (Supplemental Table S7). The Pearson's correlation coefficient between gene

expression and differential methylation was calculated with a two-tailed test of significance.

Statistical testing was performed in *R* (R Core Team, 2018) using the packages lme4, car and

lsmeans, or in the GraphPad Prism 7 statistics package. For all analyses, a *p*-value < 0.05 was

considered significant.

Data access

All raw sequencing data generated in this study have been submitted to the NCBI Gene

Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession number (to

be filled soon).

Acknowledgments

This work was supported by grants from São Paulo Research Foundation (FAPESP

2016/15881-0 and 2017/09269-3 to CAM; and 2017/09128-0 to KH), the Brazilian National

Council for Scientific and Technological Development (CNPq 403646/20162 and

303401/2014-1 to KH) and Australian Research Council (DP180101696 to BO and A. Zayed).

References

1.   C. D. Allis, T. Jenuwein, The molecular hallmarks of epigenetic control. *Nat. Rev. Genet.* **17**, 487–500 (2016).

2.   G. L. Gabor Miklos, R. Maleszka, Epigenomic communication systems in humans and honey bees: from molecules to behavior. *Horm Behav* **59**, 399–406 (2011).

3.   S. Takayama, *et al.*, Genome methylation in *D. melanogaster* is found at specific short motifs and is independent of DNMT2 activity. *Genome Res* **24**, 821–830 (2014).

4.   Y. Wang, *et al.*, Functional CpG methylation system in a social insect. *Science* **314**, 645–647 (2006).

5.   C. Honeybee Genome Sequencing, Insights into social insects from the genome of the honeybee *Apis mellifera*. *Nature* **443**, 931–949 (2006).

6.   F. Lyko, *et al.*, The honey bee epigenomes: differential methylation of brain DNA in queens and workers. *PLoS Biol.* **8**, e1000506 (2010).

7.   R. Bonasio, *et al.*, Genome-wide and caste-specific DNA methylomes of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Curr. Biol.* **22**, 1755–1764 (2012).

8.   S. Shukla, *et al.*, CTCF-promoted RNA polymerase II pausing links DNA methylation to splicing. *Nature* **479**, 74–79 (2011).

9.   H. Li-Byarlay, *et al.*, RNA interference knockdown of *DNA methyl-transferase 3* affects gene alternative splicing in the honey bee. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 12750–12755 (2013).

10.   B. G. Hunt, K. M. Glastad, S. V. Yi, M. A. D. Goodisman, The function of intragenic DNA methylation: Insights from insect epigenomes. *Integr. Comp. Biol.* **53**, 319–328 (2013).

11.   H. Yan, *et al.*, DNA Methylation in social insects: how epigenetics can control behavior and longevity. *Annu. Rev. Entomol.* **60**, 435–452 (2015).

12.   H. Yan, *et al.*, Eusocial insects as emerging models for behavioural epigenetics. *Nat. Rev. Genet.* **15**, 677–688 (2014).

13.   S. Foret, *et al.*, DNA methylation dynamics, metabolic fluxes, gene splicing, and alternative phenotypes in honey bees. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 4968–4973 (2012).

14.   S. Alvarado, R. Rajakumar, E. Abouheif, M. Szyf, Epigenetic variation in the *Egfr* gene generates quantitative variation in a complex trait in ants. *Nat. Commun.* **6**, 6513 (2015).

15.   B. R. Herb, *et al.*, Reversible switching between epigenetic states in honeybee behavioral subcastes. *Nat. Neurosci.* **15**, 1371–1373 (2012).

16.   R. Kucharski, J. Maleszka, S. Foret, R. Maleszka, Nutritional control of reproductive status in honeybees via DNA methylation. *Science* **319**, 1827–1830 (2008).

17.   B. R. Herb, M. S. Shook, C. J. Fields, G. E. Robinson, Defense against territorial intrusion is associated with DNA methylation changes in the honey bee brain. *BMC Genomics* **19**, 216 (2018).

18.   G. A. Lockett, P. Helliwell, R. Maleszka, Involvement of DNA methylation in memory processing in the honey bee. *Neuroreport* **21**, 812–816 (2010).

19.   K. M. Glastad, B. G. Hunt, S. V. Yi, M. A. D. Goodisman, Epigenetic inheritance and genome regulation: Is DNA methylation linked to ploidy in haplodiploid insects? *Proc. R. Soc. B Biol. Sci.* **281**, 20140411 (2014).

20.   S. D. Biergans, C. Giovanni Galizia, J. Reinhard, C. Claudianos, *Dnmts* and *Tet* target memory-associated genes after appetitive olfactory training in honey bees. *Sci. Rep.* **5**, 16223 (2015).

21.   S. D. Biergans, C. Claudianos, J. Reinhard, C. G. Galizia, DNA methylation mediates neural processing after odor learning in the honeybee. *Sci. Rep.* **7**, 43635 (2017).

22. L. Holman, K. Trontti, H. Helantera, Queen pheromones modulate DNA
methyltransferase activity in bee and ant workers. *Biol Lett* **12**, 20151038 (2016).

23. S. Kay, D. Skowronski, B. G. Hunt, Developmental DNA methyltransferase expression
in the fire ant *Solenopsis invicta*. *Insect Sci.* **25**, 57–65 (2016).

24. C. A. M. Cardoso-Júnior, K. R. Guidugli-Lazzarini, K. Hartfelder, DNA methylation
affects the lifespan of honey bee (*Apis mellifera* L.) workers – Evidence for a regulatory
module that involves vitellogenin expression but is independent of juvenile hormone
function. *Insect Biochem. Mol. Biol.* **92**, 21–29 (2018).

25. C. A. M. Cardoso-Junior, M. Eyer, B. Dainat, K. Hartfelder, V. Dietemann, Social
context influences the expression of DNA methyltransferase genes in the honeybee. *Sci.*
Rep. **8**, 11076 (2018).

26. K. Flores, *et al.*, Genome-wide association between DNA methylation and alternative
splicing in an invertebrate. *BMC Genomics* **13**, 480 (2012).

27. P. Cingolani, *et al.*, Intronic non-CG DNA hydroxymethylation and alternative mRNA
splicing in honey bees. *BMC Genomics* **14**, 666 (2013).

28. K. M. Glastad, K. Gokhale, J. Liebig, M. A. D. Goodisman, The caste- and sex-specific
DNA methylome of the termite *Zootermopsis nevadensis*. *Sci. Rep.* **6**, 37110 (2016).

29. K. Kannan, M. Shook, Y. Li, G. E. Robinson, J. Ma, Comparative analysis of brain and
fat body gene splicing patterns in the honey bee, *Apis mellifera. G3 Genes, Genomes,*
Genet. **9**, 1055–1063 (2019).

30. E. J. Remnant, *et al.*, Parent-of-origin effects on genome-wide DNA methylation in the
Cape honey bee (*Apis mellifera capensis*) may be confounded by allele-specific
methylation. *BMC Genomics* **17** (2016).

31. R. Libbrecht, P. R. Oxley, L. Keller, D. J. C. Kronauer, Robust DNA methylation in the
clonal raider ant brain. *Curr. Biol.* **26**, 391–395 (2016).

32. X. Wu, *et al.*, Lineage and parent-of-origin effects in DNA methylation of honey bees
(*Apis mellifera*) revealed by reciprocal crosses and whole-genome bisulfite sequencing.
Genome Biol. Evol. **12**, 1482–1492 (2020).

33. B. P. Oldroyd, B. Yagound, The role of epigenetics in the evolution of caste in insect
societies. *Philos. Trans. R. Soc. B-Biological Sci.* **In press** (2021).

34. D. S. Standage, *et al.*, Genome, transcriptome and methylome sequencing of a
primitively eusocial wasp reveal a greatly reduced DNA methylation system in a social
insect. *Mol. Ecol.* **25**, 1769–1784 (2016).

35. L. Wedd, R. Kucharski, R. Maleszka, Differentially methylated obligatory epialleles
modulate context-dependent LAM gene expression in the honeybee *Apis mellifera.*
Epigenetics **11**, 1–10 (2016).

36. Z. Lonsdale, *et al.*, Allele specific expression and methylation in the bumblebee,
Bombus terrestris. *PeerJ* **5**, e3798 (2017).

37. A. J. Bewick, *et al.*, *Dnmt1* is essential for egg production and embryo viability in the
large milkweed bug, *Oncopeltus fasciatus*. *Epigenetics and Chromatin* **12**, 6 (2019).

38. K. D. Harris, J. P. B. Lloyd, K. Domb, D. Zilberman, A. Zemach, DNA methylation is
maintained with high fidelity in the honey bee germline and exhibits global non-
functional fluctuations during somatic development. *Epigenetics and Chromatin* **12**, 62
(2019).

39. B. Yagound, N. M. A. Smith, G. Buchmann, B. P. Oldroyd, E. J. Remnant, Unique DNA
methylation profiles are associated with cis-variation in honey bees. *Genome Biol. Evol.*
11, 2517–2530 (2019).

40. B. Yagound, E. J. Remnant, G. Buchmann, B. P. Oldroyd, Intergenerational transfer of
DNA methylation marks in the honey bee. *Proc. Natl. Acad. Sci. U. S. A.*, Ahead of print
(2020).

41. H. Marshall, A. R. C. Jones, Z. N. Lonsdale, E. B. Mallon, Bumblebee workers show
differences in allele-specific DNA methylation and allele-specific expression. *Genome*
Biol. Evol. **12**, 1471–1481 (2020).

42. X. Wang, *et al.*, Function and evolution of DNA methylation in *Nasonia vitripennis*.
PLoS Genet. **9**, e1003872 (2013).

43. S. V. Arsenault, B. G. Hunt, S. M. Rehan, The effect of maternal care on gene expression
and DNA methylation in a subsocial bee. *Nat. Commun.* **9**, 3468 (2018).

44. H. Marshall, Z. N. Lonsdale, E. B. Mallon, Methylation and gene expression differences
between reproductive and sterile bumblebee workers. *Evol. Lett.* **3**, 485–499 (2019).

45. P. R. Oxley, *et al.*, The genome of the clonal raider ant *Cerapachys Biroi*. *Curr. Biol.*
24, 451–458 (2014).

46. T. Pankiw, M. L. Winston, Z. Huang, G. E. Robinson, Queen mandibular gland
pheromone influences worker honey bee (*Apis mellifera* L.) foraging ontogeny and
juvenile hormone titers. *J. Insect Physiol.* **44**, 685–692 (1998).

47. M. L. Winston, *The biology of the honey bee.* (Harvard University Press, 1987).

48. S. E. Hoover, C. I. Keeling, M. L. Winston, K. N. Slessor, The effect of queen
pheromones on worker honey bee ovary development. *Naturwissenschaften* **90**, 477–
480 (2003).

49. C. M. Grozinger, N. M. Sharabash, C. W. Whitfield, G. E. Robinson, Pheromone-
mediated gene expression in the honey bee brain. *Proc. Natl. Acad. Sci. U. S. A.* **100**
Suppl, 14519–14525 (2003).

50. D. Niu, *et al.*, Transcriptome comparison between inactivated and activated ovaries of
the honey bee *Apis mellifera* L. *Insect Mol. Biol.* **23**, 668–681 (2014).

51. L. Holman, H. Helanterä, K. Trontti, A. S. Mikheyev, Comparative transcriptomics of
social insect queen pheromones. *Nat. Commun.* **10**, 1593 (2019).

52. E. J. Duncan, M. P. Leask, P. K. Dearden, Genome architecture facilitates phenotypic
plasticity in the honeybee (*Apis mellifera*). *Mol. Biol. Evol.* **37**, 1964–1978 (2020).

53. C. A. M. Cardoso-Júnior, I. Ronai, K. Hartfelder, B. P. Oldroyd, Queen pheromone
modulates the expression of epigenetic modifier genes in the brain of honey bee workers.
Biol. Lett. **16**, 20200440 (2020).

54. C. G. Elsik, *et al.*, Finding the missing honey bee genes: Lessons learned from a genome
upgrade. *BMC Genomics* **15**, 86 (2014).

55. R. A. Drewell, *et al.*, The dynamic DNA methylation cycle from egg to sperm in the
honey bee *Apis mellifera*. *Development* **141**, 2702–2711 (2014).

56. B. Li, *et al.*, Identification and caste-dependent expression patterns of DNA methylation
associated genes in *Bombus terrestris*. *Sci Rep* **8**, 2332 (2018).

57. Y. Wang, *et al.*, Regulation of behaviorally associated gene networks in worker honey
bee ovaries. *J. Exp. Biol.* **215**, 124–134 (2012).

58. R. Libbrecht, P. R. Oxley, D. J. C. Kronauer, Clonal raider ant brain transcriptomics
identifies candidate molecular mechanisms for reproductive division of labor. *BMC*
Biol. **16**, 89 (2018).

59. R. Kucharski, R. Maleszka, Exploring DNA methylation diversity in the honey bee brain
by ultra-deep amplicon sequencing. *Epigenomes* **4**, 10 (2020).

60. A. Zemach, I. E. McDaniel, P. Silva, D. Zilberman, Genome-wide evolutionary analysis
of eukaryotic DNA methylation. *Science* **328**, 916–919 (2010).

61. N. K. E. Schulz, *et al.*, *Dnmt1* has an essential function despite the absence of CpG DNA
methylation in the red flour beetle *Tribolium castaneum*. *Sci. Rep.* **8**, 16462 (2018).

62. J. T. Washington, *et al.*, The essential role of *Dnmt1* in gametogenesis in the large
milkweed bug *Oncopeltus fasciatus*. *BioRxiv Prepr.* (2020).

63. A. U. Amukamara, *et al.*, More than DNA methylation: Does pleiotropy drive the

complex pattern of evolution of *Dnmt1*? *Front. Ecol. Evol.* **8**, 4 (2020).

64.   J. Harbo, "Propagation and instrumental insemination." in *Rinderer T. E. (Ed.) Bee*
Genetics and Breeding, Rinderer T, (Academic Press, 1986), pp. 361–389.

65.   C. A. M. Cardoso-Júnior, B. Oldroyd, I. Ronai, Vitellogenin expression in the ovaries
of adult honeybee workers provides insights into the evolution of reproductive and social
traits. *Insect Mol Biol* **Epub** (2021).

66.   R. Kucharski, S. Foret, R. Maleszka, EGFR gene methylation is not involved in
Royalactin controlled phenotypic polymorphism in honey bees. *Sci. Rep.* **5**, 14070
(2015).

67.   F. Krueger, S. R. Andrews, Bismark: A flexible aligner and methylation caller for
Bisulfite-Seq applications. *Bioinformatics* **27**, 1571–1572 (2011).

68.   B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9, 357–359 (2012).

69.   K. M. Glastad, *et al.*, Variation in DNA methylation is not consistently reflected by
sociality in Hymenoptera. *Genome Biol. Evol.* **9**, 1687–1698 (2017).

70.   A. Akalin, *et al.*, MethylKit: a comprehensive R package for the analysis of genome-
wide DNA methylation profiles. *Genome Biol.* **13**, R87 (2012).

71.   S. Heinz, *et al.*, Simple combinations of lineage-determining transcription factors prime
cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**,
576–589 (2010).

72.   R. C. Team, R: A language and environment for statistical computing. R foundation for
statistical computing, Vienna, Austria. URL: http://www.R-project.org/ (2018).

73.   R. Suzuki, H. Shimodaira, Pvclust: An R package for assessing the uncertainty in
hierarchical clustering. *Bioinformatics* **22**, 1540–1542 (2006).

74.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

75.   H. Li, *et al.*, The sequence alignment/map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

76.   M. Kearse, *et al.*, Geneious basic: An integrated and extendable desktop software
platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–
1649 (2012).

77.   J. Hellemans, G. Mortier, A. De Paepe, F. Speleman, J. Vandesompele, qBase relative
quantification framework and software for management and automated analysis of real-
time quantitative PCR data. *Genome Biol.* **8**, R19 (2007).

78.   R. M. Brito, M. McHale, B. P. Oldroyd, Expression of genes related to reproduction and
pollen foraging in honey bees (*Apis mellifera*) narcotized with carbon dioxide. *Insect*
Mol. Biol. **19**, 451–461 (2010).

79.   A. P. Lourenço, A. Mackert, A. S. Cristino, Z. L. Simoes, Validation of reference genes
for gene expression studies in the honey bee, *Apis mellifera*, by quantitative real-time
RT-PCR. *Apidologie* **39**, 372–385 (2008).

80.   M. W. Pfaffl, A. Tichopad, C. Prgomet, T. P. Neuvians, Determination of stable
housekeeping genes, differentially regulated target genes and sample integrity:
BestKeeper - Excel-based tool using pair-wise correlations. *Biotechnol. Lett.* **26**, 509–
515 (2004).

Figures

Figure 1. Social context does not drive significant alterations in the methylomes of honey bee
workers. (*A*) Number of DMRs comparing QR and QL methylomes across all three colonies
assessed for WGBS. Intersections show the number of DMRs shared by more than one colony.
(*B*) Hierarchical clustering showing the correlation-based distances of methylation patterns
across all WGBS samples; the numbers show high support values for both approximately
unbiased (au) and bootstrap probability (bp) statistics. Samples are identified by their colony
origin (A, B or C), social context (QR or QL) and tissue (Br – Brain, Ov – Ovary), respectively.
(*C*) Methylation frequency (%QL - %QR) of three top-ranked DMRs identified in the brain
and ovaries for the six colonies. The gray bars represent the frequency of methylated sites
between QL and QR bees obtained from whole-genome bisulfite sequencing (WGBS), and the
blue bars for the amplicon bisulfite sequencing (AS) for the original colonies A-C. The yellow
bars show the frequency of methylated sites (%QL - %QR) obtained from amplicon sequencing
of samples from three additional colony pairs (colonies D-F). The dashed lines represent the
10% threshold for considering a region as differentially methylated. Data from individual
cytosines and coverage can be found in Supplemental Fig. S2 and Supplemental Table S5,
respectively.

Figure 2. Relative gene expression analysis of nine differentially methylated genes in the brain and ovaries of workers from queenright (QR) and queenless (QL) colonies. "Δ Meth" shows the tissue where DNA methylation was affected by social context (Supplemental Tables S3, S4). Note that only *GB42836* (brain), *GB49839* (ovary) and *GB54664* (both tissues) were both differently methylated and expressed in the QR *vs*. QL contrast. Each box shows the interquartile range (25^{th}-75^{th} percentiles) and the median (line), while whiskers represent the minimum and maximum values. Gray dots inside boxes represent individual samples (n = 3 colonies, 8 samples per colony, 24 in total). Statistical information: GLMM test of differences between means with Tukey correction for multiple pairwise comparisons, an asterisk (*) represents the genes with significant difference in the post-hoc test but non-significant for the main 'social context' effect, 'ns' indicates $p > 0.05$, see Supplemental Table S6 for further information.

Figure 3. Differences in the methylomes and gene expression levels between source colonies.
(*A*) Venn diagrams for the number of DMRs in the brain and ovaries comparing three different
colonies. Numbers inside the intersections represent the regions that show differential
methylation of at least 10% in one source colony compared to the other two source colonies.
For example, the intersect between blue and yellow region (e.g., 4798 DMRs for the brain or
2863 DMRs for the ovaries) represents the regions from colony A that differ in methylation
level by at least 10% from the same regions in colony B and C. The intersection of all three
comparisons, i.e. the central intersect, represents the regions with three degrees of DNA
methylation levels (for example, 0% methylation level for colony A, 20% for colony B and
70% for colony C). (*B*) Number of DMRs from central intersects displayed in Fig. 3A shared
by the brain and ovary tissues. (*C*) Heatmap showing colony-specific methylation validated by
high-throughput amplicon sequencing of 11 DMRs. For this analysis, we used the WGBS data
to select genes that had a specific pattern for one colony (hypermethylated or hypomethylated),
but differed from the patterns observed in the other two colonies. Methylation level for colonies
A-C are shown as follows: Br-QL – brain queenless; Br-QR – brain queenright; Ov-QL – ovary
queenless; Ov-QR – ovary queenright. Coverage of each region can be found in Supplemental
Table S5 and methylation level of individual CpG sites are displayed in Supplemental Figs.
S2, S6, S7. (*D*) Colony-specific gene expression of differentially methylated genes. Note that
only one gene, *GB50283* (red asterisk), had a methylation pattern (Fig. 3C) that seemed
correlated with its gene expression level. Note that the list of genes shown in Fig. 3D is the
same as in Fig. 3C, the exceptions being the gene *GB41147*, a differentially methylated gene
which relative expression was not examined, and the genes *GB40816* and *GB42770,* which
were differently methylated in the WGBS data but not validated by amplicon sequencing.
Different letters inside the heatmap boxes indicate statistical differences between colonies
(GLMM test, n = 16 samples per colony and tissue). Differential expression between tissues
was observed for all of the tested genes (Supplemental Table S7), with the exception of
GB54664.

Figure 4. Differences in the methylomes and gene expression levels between brain and ovary tissue. (*A*) Venn diagram shows the number of DMRs in the contrast between brain vs. ovary methylomes in all three colonies (A-C). These nine DMRs are associated with four genes. (*B*) Difference in methylation for two of the four genes seen in the WGBS data (gray bars) and validation by amplicon sequencing in colonies A-C (blue bars) and D-F (yellow bars). The dashed line represents the 10% threshold for considering a region as differentially methylated. Coverage of each region can be found in Supplemental Table S5 and methylation level of individual CpG sites in Supplemental Fig. S7. (*C*) Transcript levels of the four differentially methylated genes quantified by quantitative PCR in the brain and ovary of young honey bee workers. Note that *GB55278* was also found to be differentially methylated between the ovaries of QR and QL workers (Fig. 2 and Supplemental Table 4). Each box shows the interquartile range (25th-75th percentiles) and the median (line), while whiskers represent the minimum and maximum values. Sample size and statistical analysis are the same as in Fig. 2 and "ns" indicates $p > 0.05$.

Figure 5. Relative expression of DNMTs encoding genes in honey bee workers kept in two different social conditions, queen presence (QR) or queen absence (QL). Transcript levels were assessed by quantitative PCR in the brain and ovary of young honey bee workers. Each box shows the interquartile range (25^{th}-75^{th} percentiles) and the median (line), while whiskers represent the minimum and maximum values. Sample size and statistical analysis are the same as in Fig. 2; "*" represents the genes with a significant difference in the post-hoc test but non-significant for the main 'social context' effect, "ns" indicates $p > 0.05$.