Antibody evasion by the Brazilian P.1 strain of SARS-CoV-2

Wanwisa Dejnirattisai[#,1], Daming Zhou [#,2], Piyada Supasa[#,1], Chang Liu[#,1,3], Alexander J. Mentzer[#,1,4], Helen M. Ginn[5], Yuguang Zhao[2], Helen M.E. Duyvesteyn[2], Aekkachai Tuekprakhon[1], Rungtiwa Nutalai[1], Beibei Wang[1], Guido C. Paesen[2], César López-Camacho[1], Jose Slon-Campos[1], Thomas S. Walter[2], Donal Skelly[4,6,7], Sue Ann Costa Clemens[8], Felipe Gomes Naveca[9], Valdinete Nascimento[9], Fernanda Nascimento[9], Cristiano Fernandes da Costa[10], Paola C. Resende[11], Alex Pauvolid-Correa[11], Marilda M. Siqueira[11], Christina Dold[12,13], Robert Levin[14], Tao Dong[3,15,16], Andrew J. Pollard[12,13], Julian C. Knight[1,4], Derrick Crook[16], Teresa Lambe[17], Elizabeth Clutterbuck[12,13], Sagida Bibi[12,13], Amy Flaxman[17], Mustapha Bittaye[17], Sandra Belij-Rammerstorfer[17], Sarah Gilbert[17], Miles W. Carroll[1,18], Paul Klenerman[4,6,12,19], Eleanor Barnes[4,6,12,19], Susanna J. Dunachie[4,6,20,21], Neil G. Paterson[5], Mark A. Williams[5], David R. Hall[5], Ruben J. G. Hulswit[2], Thomas A. Bowden[2], Elizabeth E. Fry[2], Juthathip Mongkolsapaya[$,1,3,22], Jingshan Ren[$,2], David I. Stuart[$,1,2,3,5,23], Gavin R. Screaton[$,1,3]

1. Wellcome Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, Oxford, UK.
2. Division of Structural Biology, Nuffield Department of Medicine, University of Oxford, The Wellcome Centre for Human Genetics, Oxford, UK.
3. Chinese Academy of Medical Science (CAMS) Oxford Institute (COI), University of Oxford, Oxford, UK.
4. Oxford University Hospitals NHS Foundation Trust, Oxford, UK.
5. Diamond Light Source Ltd, Harwell Science & Innovation Campus, Didcot, UK.
6. Peter Medawar Building for Pathogen Research, Oxford, UK.
7. Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK
8. Institute of Global Health, University of Siena, Siena, Brazil; Department of Paediatrics, University of Oxford, Oxford, UK.
9. Laboratório de Ecologia de Doenças Transmissíveis na Amazônia, Instituto Leônidas e Maria Deane, Fiocruz, Manaus, Amazonas, Brazil.
10. Fundação de Vigilância em Saúde do Amazonas, Manaus, Amazonas, Brazil
11. Laboratorio de vírus respiratórios- IOC/FIOCRUZ, Rio de Janeiro, Brazil
12. NIHR Oxford Biomedical Research Centre, Oxford, UK.
13. Oxford Vaccine Group, Department of Paediatrics, University of Oxford, Oxford, UK.
14. Worthing Hospital, Worthing, UK.
15. MRC Human Immunology Unit, MRC Weatherall Institute of Molecular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford, UK.
16. Nuffield Department of Medicine, University of Oxford, Oxford, UK.
17. Jenner Institute, Nuffield Department of Medicine, University of Oxford, Oxford, UK.
18. National Infection Service, Public Health England (PHE), Porton Down, Salisbury, UK..
19. Translational Gastroenterology Unit, University of Oxford, Oxford, UK
20. Centre For Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, UK.
21. Mahidol-Oxford Tropical Medicine Research Unit, Bangkok, Thailand. Department of Medicine, University of Oxford, Oxford, UK.
22. Siriraj Center of Research Excellence in Dengue & Emerging Pathogens, Dean Office for Research, Faculty of Medicine Siriraj Hospital, Mahidol University, Thailand.
23. Instruct-ERIC, Oxford House, Parkway Court, John Smith Drive, Oxford, UK.

\# These authors contributed equally to this work.
$ corresponding authors

Summary

Terminating the SARS-CoV-2 pandemic relies upon pan-global vaccination. Current vaccines

elicit neutralizing antibody responses to the virus spike derived from early isolates. However,

new strains have emerged with multiple mutations: P.1 from Brazil, B.1.351 from South Africa

and B.1.1.7 from the UK (12, 10 and 9 changes in the spike respectively). All have mutations

in the ACE2 binding site with P.1 and B.1.351 having a virtually identical triplet: E484K,

K417N/T and N501Y, which we show confer similar increased affinity for ACE2. We show

that, surprisingly, P.1 is significantly less resistant to naturally acquired or vaccine induced

antibody responses than B.1.351 suggesting that changes outside the RBD impact

neutralisation. Monoclonal antibody 222 neutralises all three variants despite interacting with

two of the ACE2 binding site mutations, we explain this through structural analysis and use

the 222 light chain to largely restore neutralization potency to a major class of public

antibodies.

Introduction

For more than a year SARS-CoV-2 has caused enormous global dislocation, leading to more

than 2.5 million deaths (https://www.worldometers.info/coronavirus Accessed: 2021-03-01)

and leaving no country untouched. Successive waves of infection have led to the imposition of

draconian lock downs in many countries resulting in severe economic and societal disruption

(Donthu and Gustafsson, 2020).

Enormous investment has been made in vaccine development with hundreds of vaccine

candidates in different stages of development using an array of different platforms from RNA,

viral vectors, recombinant protein and inactivated (Krammer, 2020). Five vaccines have now been through large scale phase III trials and have demonstrated safety and efficacy (Polack et al., 2020;Voysey et al., 2020;Baden et al., 2020). Four of these, BNT162b2 (Pfizer-BioNTech; mRNA), mRNA-1273 (Moderna; mRNA), ChAdOx1 nCoV-19 (AZD1222) (Oxford-AstraZeneca; chimpanzee adenoviral vectored) and Ad26.COV2-S (Janssen; adenovirus serotype 26 vectored) have received emergency use authorization (EUA) in a variety of countries and are being rolled out at massive scale and NVX-CoV2373 (Novavax; recombinant protein) has also shown impressive efficacy and is likely to achieve EUA in the near future (https://www.medscape.com/viewarticle/944933 Accessed: 2021-03-01). All of these vaccines have been designed to raise antibodies (and T-cells) to spike protein (S) and because of the speed of development they all include S sequences derived from the first reported sequence from Wuhan in January 2020 (Lu et al., 2020).

SARS-CoV-2 like all RNA viruses has an error prone RNA polymerase, and despite some error correction, progressive accrual of mutational changes is inevitable. The massive scale of the pandemic which is largely uncontrolled leads to huge levels of viral replication, increasing the chances that adaptive mutations will occur. There are many possible ways whereby a mutation in SARS-CoV-2 may give the virus a selective advantage, however concentrating on mutation in S, there are two clear possibilities: increased efficiency of transmission and escape from neutralizing antibodies (Volz et al., 2021).

The S protein is a large Type-1 transmembrane glycoprotein which assembles into homo-trimers (Walls et al., 2020), which form most of the outer surface of coronaviruses. S is divided into two portions S1 and S2 which are cleaved by proteolysis. S1 is responsible for target cell engagement, whilst S2 completes membrane fusion allowing the viral RNA access to the host

cell cytoplasm where viral replication can begin. S1 contains an N terminal domain (NTD) and

receptor binding domain (RBD).

The RBD interacts with the cellular receptor angiotensin converting enzyme-2 (ACE2) which

is expressed on diverse cell types, including cells in the upper and lower respiratory tracts,

allowing SARS-CoV-2 to cause respiratory infection. The ACE2 interaction surface is a small

25 amino acid patch at the apex of spike, presented to ACE2 when the RBD swings upwards

(Hoffmann et al., 2020;Shang et al., 2020) and it is mutations in this region that are causing the

most concern. Three multiply mutated viral strains appeared independently at the end of 2020

in different regions where they rapidly expanded to become the dominant strains

(https://www.cogconsortium.uk/wp-content/uploads/2021/01/Report-2_COG-UK_SARS-

CoV-2-Mutations.pdf). It is not clear how these strains acquired so many changes without clear

intermediate variants. It has however been speculated, with some evidence, that they may have

evolved in immunosuppressed chronically infected patients (Kemp et al., 2021) who support

high levels of viral replication for months and may be treated with immune plasma or

monoclonal antibodies which may drive selection of variants displaying mutations that evade

antibody responses.

Of most concern are changes in the RBD. P.1 has three: K417T, E484K and N501Y, B.1.351

also has three: K417N, E484K and N501Y whereas, B.1.1.7 contains the single N501Y

mutation. All of these changes have the potential to modulate ACE2/RBD affinity potentially

leading to increased transmissibility, for which there is now good evidence in B.1.1.7. In

addition, these mutated residues also have the potential to modulate neutralization of SARS-

CoV-2 by naturally or vaccine induced antibody responses.

In this paper we examine an isolate of P.1 cultured from a throat swab taken from an infected patient in Manaus, Brazil in December 2020 and compare its interactions with serum and antibodies with those of three other viruses, an early isolate, B.1.1.7 and B.1.351. We test the ability of immune sera induced by infection with early strains of SARS-CoV-2 (Dejnirattisai et al., 2021), or by vaccination with the Oxford-AstraZenca or Pfizer-BioNTech vaccines to neutralize P.1 (Supasa et al., 2021;Zhou et al., 2021). We see a reduction in the neutralizing capacity of immune serum to P.1 similar to the reduction seen with B.1.1.7, but not as severe as that seen with B.1.351 (Zhou et al., 2021). We demonstrate an increased affinity of P.1 RBD for ACE2 and investigate the structural basis of this through crystallography. We also study neutralization by a panel of potent monoclonal antibodies which block RBD/ACE2 interaction and provide a crystallographic solution of how one potent antibody, mAb 222, of the panel (Dejnirattisai et al., 2021) which contacts both K417 and N501, is resistant to the 501Y and 417T/N mutations found in the P.1/B1.351 strains. We dissect the basis for this via a series of high resolution structures of RDB-Fab complexes and based on this restore neutralization of certain antibodies by swapping the light chain. Finally, we bring together data on P.1, B.1.351 and B.1.1.7 and attempt to interpret the different effects these have upon the neutralizing capacity of serum generated to early SARS-CoV-2 strains.

Results

The P.1 lineage

P.1 was first reported in December 2020 from Manaus in Amazonas province of Northern Brazil (Faria et al., 2021). A large first wave of infection was seen in Manaus in March-June 2020 and by October around 75% of individuals from the region are estimated to have been infected, representing a very high attack rate. A second large wave of infection began in

December 2020 leading to further hospitalizations. This second wave corresponded with the

rapid emergence of P.1, not seen before December when it was found in 52% of cases, rising

to 85% by January 2021 (**Figure S1**).

P.1 contains multiple changes compared to B.1.1.28 and P.2 which had been previously

circulating in Brazil (Faria et al., 2021). Compared to the Wuhan sequence P.1 contains the

following mutations: L18F, T20N, P26S, D138Y, R190S in the NTD, K417T, E484K, N501Y

in the RBD, D614G and H655Y at the C-terminus of S1 and T1027I, V1176F in S2. The

position of the changes seen in P.1 compared with those found in B.1.1.7 and B.1.351 together

with a representation on where they occur on the full spike protein and RBD are shown in

Figure 1. Mutations K417T, E484K, N501Y in the ACE2 interacting surface are of the greatest

concern because of their potential to promote escape from the neutralizing antibody response

which predominately targets this region (**Figure 1D)** (Dejnirattisai et al., 2021). We searched

the COVID-19 genomics UK (COG-UK) (Tatusov et al., 2000) and the global initiative on

sharing avian influenza data (GISAID) (https://www.gisaid.org) databases. A small number

of sequences including the K417T mutation, inclusive of the P.1 lineage, have been observed

in sequencing from Japan, France, Belgium, Italy, the Netherlands and Colombia (**Figure S1**).

It is noteworthy that P.1, B.1.1.7 and B.1.351 have accrued multiple mutations in the NTD, in

B.1.1.7 there are two deletions Δ69-70 and Δ144, in B.1.351 four amino acid changes and the

Δ242–244 deletion, while in P.1 there are 6 amino acid changes in the NTD but no deletions.

Of note, two of the NTD changes in P.1 introduce N-linked glycosylation sequons T20N (TRT

to NRT) and R190S (NLR to NLS, **Figure 1E)**. The NTD, in the absence of these changes,

reasonably well populated with glycosylation sites, indeed it has been suggested that a single

bare patch surrounded by N-linked glycans glycans attached at N17, N74, N122, and N149

defines a 'supersite' limiting where neutralizing antibodies can attach to the NTD (Cerutti et

al., 2021). Residue 188 is somewhat occluded whereas residue 20 is highly exposed, is close

to the site of attachment of neutralizing antibody 159 (Dejnirattisai et al., 2021) and impinges

on the proposed NTD supersite.

The effects of RBD mutations on ACE2 affinity.

We have previously measured the affinity of RBD/ACE2 interaction for Wuhan, B.1.1.7

(N501Y) and B.1.351 (K417N, E484K, N501Y) RBDs. N501Y increased affinity 7-fold and

the combination of 417, 484 and 501 mutations further increased affinity (19-fold compared to

Wuhan). Here we have expressed P.1 RBD (K417T, E484K, N501Y). The K_D for the

P.1/ACE2 interaction is 4.8 nM with Kon=1.08E5/Ms, Koff=5.18E-4/s (**Figure S2**, Methods),

showing that binding to P.1 is essentially indistinguishable from B.1.351 (4.0 nM).

To better understand RBD-ACE2 interactions we determined the crystal structure of the RBD-

ACE2 complex at 3.1 Å resolution (Methods, **Table S1**). As expected the mode of RBD-ACE2

engagement is essentially identical for P.1 and the original Wuhan RBD sequence (**Figure**

2A). The RMS deviation between the 791 Ca positions is 0.4 Å, similar to the experimental

error in the coordinates, and the local structure around each of the three mutations is conserved.

Nevertheless, calculation of the electrostatic potential of the contact surfaces reveals a marked

change, with much greater complementarity for the P.1 RBD consistent with higher affinity.

(**Figure 2B,C,D**).

Residue 417 lies at the back of the RBD neck (our RBD anatomy follows Dejnirattisai et al.,

2021) and in the original SARS-CoV-2 is a lysine residue which forms a salt-bridge with D30

of ACE2 (**Figure 2E**). The threonine of P.1 RBD no longer forms this interaction and the gap

created is open to solvent, so there is no obvious reason why the mutation would increase

affinity for ACE2, and this is consistent with directed evolution studies (Zahradník et al., 2021)

where this mutation was rarely selected in RBDs with increased affinity for ACE2.

Residue 484 lies atop the left shoulder of the RBD and neither the original Glu nor the Lys of

P1 make significant contact with ACE2, nevertheless the marked change in charge

substantially improves the electrostatic complementarity (**Figure 2F,G**), consistent with

increased affinity.

Residue 501 lies on the right shoulder of the RBD and the change from a relatively short Asn

sidechain to the large aromatic Tyr allows for favourable ring stacking interactions consistent

with increased affinity (**Figure 2H**).

Binding of P.1 RBD by potent human monoclonal antibodies

We have previously described a large panel of monoclonal antibodies generated from patients

infected with early strains of SARS-CoV-2, before the emergence of B.1.1.7 (Dejnirattisai et

al., 2021). From this panel we have selected 20 potent antibodies which have focus reduction

neutralization 50% (FRNT50) values <100ng/ml, 19 of these mAbs have an epitope on the

RBD and all of these block ACE2/RBD interaction, whilst mAb 159 binds the NTD. We used

biolayer interferometry (BLI) to measure the affinity of the RBD-binding antibodies and found

that compared to Victoria (SARS-CoV-2/human/AUS/VIC01/2020), an early isolate of SARS-

CoV-2, which has a single change S247R in S compared to the Wuhan strain (Seemann et al.,

2020; Caly et al., 2020) monoclonal antibody binding was significantly impacted with a

number showing complete knock-out of activity (**Figure 2I**). The results with P.1 showed a

greater impact compared to B.1.1.7 but similar to B.1.351 (Zhou et al., 2021), this is expected

since both contain mutation of the same 3 residues in the RBD, only differing at position 417,

K417N in B.1.351 and K417T in P.1. The localization of the impact on binding is shown in

Figure 2J and reflects direct interaction with mutated residues. Of note is mAb 222 which

maintains binding potency across all variants despite adjacency to mutated residues, as

discussed below.

Neutralization of P.1 by potent human monoclonal antibodies

Using the same set of 20 potent antibodies neutralization was measured by a focus reduction

neutralization test (FRNT) and compared with neutralization of Victoria and variants B.1.1.7

and B.1.351. Compared to Victoria neutralization by the monoclonal antibodies was

significantly impacted by P.1, with 12/20 showing >10-fold reduction in FRNT50 titre and a

number showing complete knock out of activity (**Figure 3A Table S2**). The results with P.1

showed a greater impact compared to B.1.1.7 but were as expected similar to those with

B.1.351 (Zhou et al., 2021). There is good correlation between the negative impact on

neutralization and on RBD-affinity (**Figure 2J**).

Reduced neutralization of P.1 by monoclonal antibodies being developed for clinical use.

A number of potent neutralizing antibodies are being developed for clinical use either

therapeutically of prophylactically (Ku et al., 2021;Baum et al., 2020;Kemp et al., 2021). We

performed neutralization assays against P.1 using antibodies S309 Vir (Pinto et al., 2020),

AZD8895, AZD1061 and AZ7442 (a combination of AZD8895 and AZD1061) AstraZeneca,

REGN10987 and REGN10933 Regeneron, LY-CoV555 and LY-CoV16 Lilly and ADG10,

ADG20 and ADG30 from Adagio (**Figure 3B**). The affinity of binding to P.1 RBD was also

investigated by BLI for the Regeneron and AstraZeneca antibodies and the results (**Figure 2I**)

parallel closely the neutralization results. Neutralization of both Lilly antibodies was severely

impacted with LY-CoV16 and LY-CoV555 showing almost complete loss of neutralization of

P.1 and B.1.351 while LY-CoV16 also showed marked reduction in neutralization of B.1.1.7.

There was also escape from neutralization of P.1 by REGN10933 and a modest reduction in

neutralization of P.1 by AZD8895, while AZD1061 and AZD 7442 showed equal

neutralization of all SARS-CoV-2 variants. The three Adagio antibodies neutralized all variants

with all reaching a plateau at 100% neutralization and interestingly, ADG30 showed a slight

increase of neutralization of P.1. S309 Vir was largely unaffected although for several viruses,

including P.1, the antibody failed to completely neutralize, conceivably reflecting incomplete

glycosylation at N343, since the sugar interaction is key to binding of this antibody N343 (Pinto

et al., 2020). The escape from REGN10933 and LY-CoV555 mirrors that of other potent

antibodies (including 316 and 384 in our set) which make strong interactions with residues

484-486 and are severely compromised by the marked change E484K, whereas LY-CoV016,

an IGHV3-53 mAb, is affected by changes at 417 and 501. The abrogation of the Lilly Ly-

CoV-16 and LyCoV-555 antibodies reflects the observation of Starr et al. (Starr et al.,

2021)(Greaney et al., 2021) that LY-CoV555 is sensitive to mutation at residue 384 and LY-

CoV16 is sensitive to changes at 417.

Reduced neutralization by an NTD-binding antibody

The neutralization titre of NTD-binding mAb159, was 133-fold reduced on P.1 compared to

Victoria with only 64% neutralization at 10μg/ml (**Figures 3A**). Although P.1 does not harbor

deletions in the NTD like B.1.1.7 (Δ69−70, Δ144) or B.1.351 (Δ242−244), it is clear that the 6

NTD mutations in P.1 (L18F, T20N, P26S, D138Y, R190S) disrupt the epitope for mAb159

(**Figure 4A**) (Dejnirattisai et al., 2021;Supasa et al., 2021). It is possible that the failure of this

antibody to achieve complete neutralization could be due to partial glycosylation at residue 20,

which is some 16 Å from bound Fab 159, however the L18F mutation is even closer and likely

to diminish affinity (**Figure 4A**). Since it has been proposed that there is a single supersite for

potent NTD binding antibodies we would expect the binding of many of these to be affected

(Cerutti et al., 2021).

Reduced neutralization by VH3-53 public antibodies

Five of the potent monoclonal antibodies used in this study (150, 158, 175, 222 and 269),

belong to the VH3-53 family and a further 2 (out of 5 of this family) belong to the almost

identical VH3-66, and the following discussion applies also to these antibodies. The binding

sites for these have been described (Dejnirattisai et al., 2021). The large majority of these

antibodies attach to the RBD in a very similar fashion. These motifs recur widely, VH3-53 are

the most prevalent deposited sequences and structures for SARS-CoV-2 neutralizing

antibodies. Their engagement with the RBD is dictated by CDR-H1 and CDR-H2 whilst the

CDR-H3 is characteristically short and makes rather few interactions (Yuan et al., 2020; Barnes

et al., 2020;Dejnirattisai et al., 2021). We have previously solved the structures of mAbs 150,

158 and 269 (**Figure 4B**) which show that whilst there are no contacts with residue 484, there

are interactions of CDR-H3 with K417 and CDR-L1 with N501, meaning that binding and

neutralization by VH3-53 antibodies would be predicted to be compromised by the N501Y

change in variant viruses B.1.1.7, B.1.351 and P.1, whilst the additional change at 417 in P.1

(K417T) and B.1.351 (K417N) might be expected to have an additive effect (Dejnirattisai et

al., 2021).

In practice, changes in the light chain and CDR-H3 between members of this family mean that

the story is more complex. Thus, neutralization of P.1 by 175 and 158 is severely impacted and

neutralization of P.1 by 269 is almost completely lost. However, for 150 P.1 neutralization is

less compromised than for B.1.351 (Zhou et al., 2021), whilst for 222 neutralization is

completely unaffected by the changes in P.1 and indeed all variants (**Figure 3A**).

We measured the affinity of 222 for both P.1 (KD = 1.92±0.01 nM) and Wuhan RBD (KD =

1.36±0.08 nM) showing no reduction in the strength of interaction despite the changes

occurring in the putative binding site for P.1 (**Table S2**).

To understand how 222 is able to still neutralize P.1 we solved the crystal structures of six

ternary complexes of 222 in complex with the RBDs for (i) the original virus, and bearing

mutations (ii) K417N; (iii) K417T; (iv) N501Y; the 417, 484 and 501 changes characteristic

of B.1.351 (v) and P.1 (vi). All crystals also contained a further Fab, EY6A as a crystallization

chaperone (Zhou et al., 2020), were isomorphous and the resolution of the structures ranged

from 1.95 to 2.67 Å, **Figure 4C,D**, Methods, **Table S1**. As expected, the structures are highly

similar with the binding pose of 222 being essentially identical in all structures (pairwise

RMSD in Cα atoms between pairs of structures are ~0.2-0.3 Å for all residues in the RBD and

Fv region of mAb 222, **Figure 4D**).

In the original virus residue 417 makes a weak salt bridge interaction with heavy chain CDR3

residue E99. Mutation to either Asn or Thr abolishes this and there is little direct interaction,

although there are weak (~3.5 Å) contacts to heavy chain Y52 and light chain Y92 (**Figure

4E**). However, a buffer molecule/ion rearranges to form bridging interactions and this may

mitigate the loss of the salt bridge, in addition the original salt bridge is weak and its

contribution to binding may be offset by the loss of entropy in the lysine sidechain. We note

that CDR-H3 of 222, at 13 residues is slightly longer than found in the majority of potent VH3-

53 antibodies, however this seems unlikely to be responsible for the resilience of 222, rather it

seems that there is little binding energy in general from the CDR3-H3, since most of the binding

energy contribution of the heavy chain comes from CDR-H1 and CDR-H2 which do not

interact with RBD residue 417, meaning that many VH3-53 antibodies are likely to be resilient

to the common N/T mutations (**Figure 4B**).

Residue 501 makes contact with CDR-L1 of mAb 222 (**Figure 4D,F**), however the interaction,

with P30 is probably slightly strengthened by the N501Y mutation which provides a stacking

interaction with the proline, conferring resilience. This is in contrast to the situation with most

other VH3.53 antibodies where direct contacts confer susceptibility to escape by mutation to

Tyr (**Figures 2I,J and 3A**) (Supasa et al., 2021;Zhou et al., 2021).

The 222 light chain can rescue neutralization by other VH3-53 mAbs

Reasoning that the relative robustness of mAb 222 to common variants (P.1, B.1.1.7 and

B.1.351) compared to other VH3-53 antibodies stems from the choice of light chain we

modelled the 222LC with the heavy chains of other VH3-53 antibodies to see if they might be

compatible **(Figure 4G)**. The result was striking, it appeared that there would likely be no

serious steric clashes. This contrasted with the numerous clashes seen when we docked the

light chains of other VH3-53 antibodies onto the heavy chain of 222 (**Figure 4G,H**). This

suggests that the 222 light chain might be an almost universal light chain for these 3-53

antibodies and could confer resilience to P.1, B.1.1.7 and B.1.351 variants. This led us to create

chimeric antibodies containing the 222LC combined with the HC of the other VH3-53 mAbs

150, 158, 175 and 269. In all cases, chimeric antibodies expressed well and we performed

neutralization assays against Victoria, B.1.1.7, B.1.351 and P.1 viruses **(Figure 5)**. For B.1.1.7

neutralization of 150HC/222LC, 158HC/222LC and 269HC/222LC was restored to near the

level seen on Victoria, whilst 175HC/222LC could not fully neutralize B.1.1.7. For B.1.351

and P.1 the activity of mAbs 150 and 158 was restored in chimeras containing the 222LC, with

the 150HC/222LC showing 50-fold greater potency against B.1.351 (7ng vs 350 ng/ml) and

13-fold greater potency against P.1 (3ng vs 40 ng/ml) than native 150. With an FRNT50 of

3ng/ml 150HC/222LC was the most potent antibody tested against P.1.

Neutralization of P.1 by convalescent plasma

We collected convalescent plasma samples from a cohort of volunteers who had suffered from

SARS-CoV-2 infection evidenced by a positive diagnostic PCR. Samples were collected

during the convalescent phase, 4-9 weeks following infection, all samples were taken during

the first wave of infection in the UK, prior June 2020 and well before the emergence of the

B.1.1.7 variant. We have also collected plasma from volunteers recently infected with B.1.1.7

as demonstrated by viral sequencing or S gene drop out from the diagnostic PCR (Dejnirattisai

et al., 2021;Supasa et al., 2021).

Neutralization of P.1 was assessed by FRNT on 34 convalescent samples **(Figure 6A Table**

S3A). P.1 neutralization curves are displayed alongside neutralization curves for Victoria,

together with B.1.1.7 and B.1.351. P.1 geometric mean neutralization titres were reduced 3.1-

fold compared to Victoria (p< 0.0001). This reduction was similar to B.1.1.7 (2.9-fold) and

considerably less than B.1.351 (13.3-fold) **(Figure 6C)**. When using plasma from individuals

infected with B.1.1.7 we saw only modest (1.8-fold p=0.0039) reductions in neutralization

comparing P.1 with Victoria **(Figure 6B and D Table S3B)**.

Neutralization of P.1 by vaccine serum

We next performed neutralization assays using serum collected from individuals who had

received either the BNT162b2 Pfizer-BioNTech or ChAdOx1 nCoV-19 Oxford-AstraZeneca

vaccine **Figure 7** (Supasa et al., 2021;Zhou et al., 2021). For the Pfizer BioNTech vaccine

serum was collected 4-14 days following the second dose of vaccine administered three weeks

after the first dose (n=25). For the Oxford-AstraZeneca vaccine serum was taken 14 or 28 days

following the second dose which was administered 8-14 weeks following the first dose (N=25).

Geometric mean neutralization titres against P.1 were reduced 2.6-fold (p<0.0001) relative to

the Victoria virus for the Pfizer-BioNTech vaccine serum **Figure 7A,C** and 2.9-fold

(P<0.0001) for the Oxford-AstraZeneca vaccine **Figure 7B,D Table S4.**

Neutralization titres against P.1 were similar to those against B.1.1.7 and only a minority of

samples failed to reach 100% neutralization at 1:20 dilution of serum, considerably better than

neutralization of B.1.351, where titres were reduced 7.6-fold and 9-fold for the BNT162b2

Pfizer and ChAdOx1 nCoV-19 AstraZeneca vaccines respectively.

Discussion

Large scale viral sequencing programmes have uncovered a spectrum of mutations containing

changes at many locations in the SARS-CoV-2 genome in correspondence with the concept of

viral quasispecies (Domingo and Perales, 2019). Mutations in S are of particular concern as S,

through the RBD, directs cellular tropism and in addition is the target for the neutralizing

antibody response. Mutations in S could therefore enhance viral fitness by increasing affinity

to ACE2 or provide escape from the antibody response induced by natural infection or

vaccination.

P.1 contains 12 individual changes spread throughout S with three changes in the RBD. In this

paper we demonstrate an increase in affinity of interaction for P.1 RBD with ACE2 to an

equivalent degree as that observed for B.1.351 with binding somewhat tighter than for B.1.1.7.

It seems conceivable that this increase in receptor affinity may drive increased virus

transmissibility, allowing the three variants to become dominant strains in the regions where

they emerged (Zhou et al., 2021;Supasa et al., 2021).

The ACE2 interacting surface of RBD is a small 25 amino acid patch at the apex of S and is

under extreme selection pressure as it not only mediates interaction with the cellular receptor

but is also the site of binding for a major class of neutralizing antibodies that block the

interaction of ACE2 with the RBD (Zost et al., 2020;Kreye et al., 2020;Wu et al., 2020;Yuan

et al., 2020). (Dejnirattisai et al., 2021). Recently, two elegant unbiased approaches have been

used to assess the influence of mutation on the ACE2/RBD binding affinity or the ability of

RBD mutations to evade the polyclonal antibody response. Firstly, a yeast display approach

was used to generate RBD mutants with enhanced ACE2 binding. Amongst a number of

mutations selected were the very same positions found in the recent variants of concern, namely

E484K and N501Y, and less frequently changes at residue 417 were also observed (Zahradník

et al., 2021). Multiple rounds of selection led to the emergence of mutant RBDs with 600-fold

higher affinity to ACE2, in the low picomolar range. In a second approach, polyclonal anti-

SARS-CoV-2 serum was used to select mutant RBD from a yeast display library which showed

reduced antibody binding (Greaney et al., 2021). This approach led to the identification of a

number of potential antibody escape mutants, amongst them E484 which is likely responsible

for a proportion of the escape from antibody neutralization we describe for P.1.

What is driving the emergence of the new strains is difficult to determine, the emergence of

B.1.1.7 occurred on the background of relatively low population immunity and may have been

primarily driven by increased transmissibility. The emergence of B.1.351 occurred on the

background of around 30% seropositivity in South Africa and P.1 on the background of an

estimated 75% seropositivity in Manaus Brazil (Faria et al., 2021). It seems possible that

selection of P.1 and B.1.351 may have been in part driven by immune escape, however until

methods are developed to screen at a population level for the frequency of reinfection, it is not

possible to determine this, especially as reinfection may lead to more mild or asymptomatic

disease.

Because P.1 and B.1.351 contain very similar changes in the RBD it might be assumed that

neutralization of both would be similarly affected. This was indeed the case for neutralization

by monoclonal antibodies directed at the RBD, where there was substantial escape from many

antibodies in our panel or from antibodies being developed for clinical use. However,

neutralization of P.1 was not compromised as severely as neutralization of B.1.351, when using

convalescent or vaccine serum induced by earlier SARS-CoV-2 strains(Zhou et al., 2021).

Using convalescent serum B.1.351 showed 13-fold reduction in neutralization compared to

Victoria whilst P.1 was only reduced 3.1-fold, comparable to the reduction seen with B.1.1.7,

which only harbours the single N501Y change in the RBD (Zhou et al., 2021;Supasa et al.,

2021). Similarly, neutralization of P.1 by vaccine serum was less impacted than neutralization

of B.1.351 meaning that vaccination with Wuhan S will likely provide some protection against

P.1. There is now clinical evidence that the ChAdOx1 nCoV-19 Oxford-AstraZeneca and

NVX-CoV2373 Novavax vaccines provide protection from B.1.1.7 (Emary et al.,

2021;Mahase, 2021). For B.1.351 both the Novavax

(https://www.webmd.com/vaccines/covid-19-vaccine/news/20210131/vaccine-not-as-

effective-against-south-african-variant) and Janssen vaccine

(https://www.reuters.com/article/us-health-coronavirus-vaccines-johnson-j-

idUSKBN29Z0F0) saw a marked decrease in efficacy against B.1.351, but still showed >50%

protection against moderate and severe disease, whilst in a Phase II trial, ChAdOx1 nCoV-19

efficacy against mild to moderate disease caused by B.1.351 was 10.4% (95% CI: –76.8; 43.8),

but efficacy against severe disease could not be assessed in this study (Madhi et al., 2021).

The reason for the differences in neutralization of B.1.351 and P.1 by immune serum are not

immediately clear, but presumably reflect the difference in the mutations introduced outside

the RBD. In addition to our mAb 159 a number of potent neutralizing mAbs have been reported

that map to the NTD (Cerutti et al., 2021), and this domain has multiple mutations in all three

major variant strains: B.1.1.7 has two deletions, B.1.351 has a deletion and four substitutions

and P.1 has 6 amino acid substitutions, including the creation of two N-linked glycan sequons

(**Figure 1 A-C**). Comparison of neutralization of pseudoviruses expressing only the three RBD

mutations (K417N E484K N501Y) of B.1.351 with pseudovirus expressing the full suite of

mutations in B.1.351 spike show that the non-RBD changes substantially increase escape from

neutralization (Wibmer et al., 2021;Dejnirattisai et al., 2021;Wang et al., 2021). The changes

in the NTD of the major variants are far less consistent than those found in the RBD, and there

are no strong trends in electrostatic properties (**Figure 1A-C**). It therefore remains unclear what

the drivers are for these changes, although one or more of immune escape, co-receptor binding,

and modulation of RBD dynamics affecting presentation of the receptor binding site are

plausible. Nonetheless, it seems likely that these changes are largely responsible for the non-

RBD component of neutralization variation between strains.

A number of public antibody responses have been reported for SARS-CoV-2, principal

amongst these being VH3-53/VH3-66 and VH1-58 (VH3-30 is also found but the antibodies

are not potent neutralizers) (Yuan et al., 2020;Barnes et al., 2020; Dejnirattisai et al., 2021).

We have previously shown that mixing heavy and light chains from antibodies within VH1-58

can increase the neutralization titre by 20-fold from the parent antibodies (chimera of 253HC

with 55LC or 165LC) (Dejnirattisai et al., 2021). Here we have shown that chimeras created

amongst the VH3-53 antibodies using the 222LC are able to confer broad neutralization to

antibodies which have reduced neutralization capacity against the viral variants. Furthermore,

the chimera of 150HC with 222LC achieved 13 and 3-fold increases in neutralization titre

compared to the parental 150 and 222 mAb respectively. Creation of such antibody chimeras

amongst other anti-SARS-CoV2 antibodies may similarly lead to the discovery of more

antibodies with enhanced activity. This also suggests that highly effective natural responses

against all three variants, and common cross-protective responses, will be found.

The recent emergence of a number of variants of concern has led to efforts to design new

vaccines which will be able to protect against the viral variants. Exactly which variants or

sequences should be selected is difficult to determine in what is likely to be an evolving

situation, as vaccine induced herd immunity increases the selection pressure for immune

escape. Based on the results reported here the South African B.1.351 is the variant of greatest

concern giving the largest reductions in neutralization titres and evidence of complete failure

to neutralize in some cases and we believe developing vaccine constructs to B.1.351 to be the

greatest priority.

In summary, we demonstrate that P.1 can escape neutralization by a number of monoclonal

antibodies including some being developed for prophylactic or therapeutic use, while other

antibodies with epitopes away from the mutated RBD residues retain broad neutralization.

Thus S309/AZD1061/REGN10987/ADG10/ADG20/ADG30 showed little to no reduction

(<4-fold) in neutralization activity across the three variants, consistent with their previously

described broadly neutralizing activities across clade I sarbecoviruses.

In contrast to B.1.351, neutralization of P.1 does not show such a substantial reduction by

polyclonal serum induced by natural infection or vaccination and there is no evidence of

widespread escape. Despite the reduction in neutralization titres it is hoped that immunization

with vaccines designed against parent/ancestral strains will provide protection from P.1.

Limitations of the study

The in vitro FRNT assays we report here do not measure the effect of complement or antibody

dependent cell mediated cytotoxicity which may enhance neutralization in vivo. The role that

T cells play in immunity to SARS-CoV-2 and in particular protection from severe disease is

unknown and worthy of investigation, but recent findings suggest that CD4 and CD8 T cell

responses raised to ancestral strains are minimally impacted by the variants (Alison Tarke et

al., 2021;Skelly et al., 2021). It will be interesting to determine the directionality of

neutralization between the different variant viruses and naturally acquired antibody responses

to them. For instance, there is some suggestion in this report that plasma induced by B.1.1.7 is

better able to neutralize B.1.351 and P.1. Measuring neutralization of viral variants by B.1.351

and P.1 serum will give a better idea of cross protection against the other strains.

Acknowledgements

This work was supported by the Chinese Academy of Medical Sciences (CAMS) Innovation

Fund for Medical Science (CIFMS), China (grant number: 2018-I2M-2-002) to D.I.S. and

G.R.S. H.M.E.D. and J.Ren are supported by the Wellcome Trust (101122/Z/13/Z), Y.Z. by

Cancer Research UK (C375/A17721), TAB and RJGH by the UKRI MRC (MR/S007555/1),

D.I.S. and E.E.F. by the UKRI MRC (MR/N00065X/1). D.I.S. is a Jenner Investigator. The

National Institute for Health Research Biomedical Research Centre Funding Scheme supports

G.R.S. We are also grateful for a Fast Grant from Fast Grants, Mercatus Center to support the

isolation of human monoclonal antibodies to SARS-CoV-2 and Schmidt Futures for support of

this work. G.R.S. is also supported as a Wellcome Trust Senior Investigator (grant

095541/A/11/Z). This is a contribution from the UK Instruct-ERIC Centre. The Wellcome

Centre for Human Genetics is supported by the Wellcome Trust (grant 090532/Z/09/Z). Virus

used for the neutralisation assays was isolated by Julian Druce, Doherty Centre, Melbourne,

Australia. Chanice Knight, Emily Chiplin, Ross Fothergill and Liz Penn contributed to assays.

We acknowledge Diamond Light Source for time on Beamline I03 under Proposal lb27009 for

COVID-19 Rapid Access. Huge thanks to the teams, especially at the Diamond Light Source

and Department of Structural Biology, Oxford University that have enabled work to continue

during the pandemic. The computational aspects of this research were supported by the

Wellcome Trust Core Award Grant Number 203141/Z/16/Z and the NIHR Oxford BRC. The

Oxford Vaccine work was supported by UK Research and Innovation, Coalition for Epidemic

Preparedness Innovations, National Institute for Health Research (NIHR), NIHR Oxford

Biomedical Research Centre, Thames Valley and South Midland's NIHR Clinical Research

Network. We thank the Oxford Protective T-cell Immunology for COVID-19 (OPTIC) Clinical

team for participant sample collection and the Oxford Immunology Network Covid-19

Response T cell Consortium for laboratory support. We acknowledge the rapid sharing of

Victoria, B.1.1.7 and B.1.351 which was isolated by scientists within the National Infection

Service at PHE Porton Down. We thank The Secretariat of National Surveillance, Ministry of

Health Brazil for assistance in obtaining P.1 samples. This work was supported by the UK

Department of Health and Social Care as part of the PITCH (Protective Immunity from T cells

to Covid-19 in Health workers) Consortium, the UK Coronavirus Immunology Consortium

(UK-CIC) and the Huo Family Foundation. EB and PK are NIHR Senior Investigators and PK

is funded by WT109965MA and NIH (U19 I082360). DS is an NIHR Academic Clinical

Fellow. The views expressed in this article are those of the authors and not necessarily those

of the National Health Service (NHS), the Department of Health and Social Care (DHSC), the

National Institutes for Health Research (NIHR), the Medical Research Council (MRC) or

Public Health, England.

Author Information

These authors contributed equally: WD, DZ, PS, CL, AJM.

Contributions

Contributions

D.Z. performed BLI interaction analyses. D.Z., J.R., N.G.P., M.A.W. and D.R.H. prepared the

crystals, enabled and performed X-ray data collection. J.R., E.E.F., H.M.E.D. and D.I.S.

analysed the structural results. G.R.S., J.M., P.S., Y.Z., D.Z., G.C.P, B.W., R.N., A.T., J.S-C.,

C.L-C. and C.L. prepared the Spike constructs, RBDs, ACE2 and antibodies and, W.D. and

P.S. performed neutralization assays. D.C. provided materials. H.M.G. wrote MABSCAPE and

performed mapping and cluster analysis, including sequence analyses. S.A.C.C., P. G. N.,

V.N., F. N., C. F. C., P.C.R., A.P-C., M.M.S., A.J.M., E.B., S.J.D., D.S., C.D., R.L., T.D.,

A.J.P., J.C.K., P.K., M.W.C., T.L., S.B., A.F., M.B., S.B-R., E.C. and S.G. assisted with patient

samples and vaccine trials. E.B., M.C., S.J.D., P.K. and D.S. conceived the study of vaccinated

healthcare workers and oversaw the OPTIC Healthcare Worker study and sample

collection/processing. G.R.S. and D.I.S. conceived the study and wrote the initial manuscript

draft with other authors providing editorial comments. R.J.G.H. and T.A.B. contributed to study

design. All authors read and approved the manuscript.

Figure Legends

Figure 1. Mutational landscape of P.1. Schematic showing the locations of amino acid

substitutions in (A) P.1, (B) B.1.351 and (C) B.1.1.7 relative to the Wuhan SARS-CoV-2

sequence. Under the structural cartoon is a linear representation of S with changes marked on.

Where there is a charge change introduced by mutations the change is coloured (red if the

change makes the mutant more acidic/less basic, blue more basic/less acidic). (D) Depiction of

the RBD as a grey surface with the location of the three mutations K417T, E484K and N501Y

(magenta) the ACE2 binding surface of RBD is coloured green. (E) locations of N-linked

glycan (red spheres) on the spike trimer shown in a pale blue surface representation, the two

new sequons found in P.1 are marked blue.

Figure 2. Comparison of WT RBD/ACE2 and P.1 RBD/ACE2 complexes. (A)

Comparison of P.1 RBD/ACE2 (grey and salmon) with WT RBD/ACE2 (blue and cyan)

(PDB ID 6LZG) by overlapping the RBDs. The mutations in the P.1 RBD are shown as

sticks. (B), (C) Open book view of electrostatic surface of the WT RBD/ACE2 complex, and

(C), (D) of the P.1 RBD/ACE2 complex. Note the charge difference between the WT and the

mutant RBDs. The charge range displayed is ±5 kJ/mol. (E) The K417 of the WT RBD forms

a salt bridge with D30 of ACE2. (F) and (G) Effect of E484K mutation on the electrostatic

surface (H) Y501 of the P.1 RBD makes a stacking interaction with Y41 of ACE2. (H) K_D of

RBD/mAb interaction measured by BLI for RBDs of Victoria, B.1.1.7, P.1 and B.1.351 (left

to right) (I) BLI data mapped onto the RBD using the method described (Dejnirattisai *et al.*,

2021). Front and back views of the RBD are shown. In the left pair the spheres represent the

antibody binding sites coloured according to the ratio (K_DP.1/K_DWuhan). For white the ratio

is 1, for red it is <0.1 (i.e. at least 10-fold reduction) black dots refer to mapped antibodies

not included in this analysis, dark green RBD ACE2 binding surface, yellow mutated K417T,

E484K, N501Y. For the right pair atoms are coloured according to the ratio of neutralisation

titres (IC_{50}B.1.351/IC_{50}Victoria), for white the ratio is 1, for red it is <0.01 (i.e. at least 100-

fold reduction). Note the strong agreement between K_D and $IC_{50.}$ 269 is very strongly

affected and is close to the IGHV3-53 and IGHV3-66 antibodies (*e.g.* 222).

Figure 3. Neutralization of P.1 by monoclonal antibodies. (A) Neutralization of P.1 by a

panel of 20 potent human monoclonal antibodies. Neutralization was measured by FRNT,

curves for P.1 are superimposed onto curves for Victoria, B.1.1.7 and B.1.351 as previously

reported (Supasa et al., 2021;Zhou et al., 2021). FRNT50 titres are reported in Table S2

Neutralization curves for monoclonal antibodies in different stages of development for

commercial use. (B) Shows equivalent plots for the Vir, Regeneron, AstraZeneca, Lilly and

Adagio antibodies therapeutic antibodies.

Figure 4. Structures of Fab 222 in complex with P.1 RBD. (A) Ribbon depiction of Fab

159/NTD complex with P1 mutations in the NTD highlighted as cyan spheres. (B) Front and

back surfaces of the RBD bound to a typical VH3-53. P1 mutations in the RBD are highlighted

in dark green and labelled. In this group, monoclonal antibody 222 has a slightly longer CDR3.

Sequences of VH3-53 CDR1-3 heavy and light chains are also shown. (C) Crystal structure of

P1 RBD, 222 Fab and EY6A Fab (Zhou et al., 2020). (D) Close up of 222 CDRs interacting

with the RBD (grey) mutations are highlighted in yellow on the green ACE2 interface. (E)

K417 interactions with Fab 222 (F) N501 interactions with Fab 222. (G), (H) Fab 222 chimera

models.

Figure 5. Neutralization curves of VH3-53 chimeric antibodies. Neutralization curves of

Victoria, B.1.1.7, B.1.351 and P.1. Left hand column; neutralization curves using the native

antibodies 222, 150, 158, 175 and 269. Right hand column; neutralization curves for chimeric

antibodies, the heavy chains of 150, 158, 175 and 269 are combined with the light chain of

222, native 222 is used as the control. FRNT50 titres are given in Table S2.

Figure 6 Neutralization of P.1 by convalescent plasma. Plasma (n=34) was collected from

volunteers 4-9 weeks following SARS-CoV-2 infection, all samples were collected before June

2020 and therefore represent infection before the emergence of B.1.1.7 in the UK. (A)

Neutralization of P.1 was measured by FRNT, comparison is made with neutralization curves

for Victoria, B.1.1.7 and B.1.351 that we have previously generated (Zhou et al., 2021;Supasa

et al., 2021). (B) Neutralization of P.1 by plasma taken from volunteers who had suffered

infection with B.1.1.7 as evidenced by sequencing or S-gene drop out by diagnostic PCR.

Samples were taken at varying times following infection. (C-D) Comparison of FRNT50 titres

between Victoria and P.1, data for B.1.1.7 and B.1.351 are included for comparison and, the

Wilcoxon matched-pairs signed rank test was used for the analysis and two-tailed P values

were calculated, geometric mean values are indicated above each column.

Figure 7 Neutralization of P.1 by vaccine serum. (A) Pfizer vaccine, serum (n=25) was taken

7-17 days following the second dose of the Pfizer-BioNTech vaccine. FRNT titration curves

are shown with Victoria, B.1.1.7 and B.1.351 as comparison (Supasa et al., 2021;Zhou et al.,

2021). (B) AstraZenca vaccine, serum was taken 14 or 28 days following the second dose of

the Oxford-AstraZeneca vaccine (n=25). (C-D) Comparison of FRNT50 titres for individual

samples for the Pfizer and AstraZeneca vaccine between Victoria, B.1.1.7, B.1.351 and P.1,

the Wilcoxon matched-pairs signed rank test was used for the analysis and two-tailed P values

were calculated, geometric mean values are indicated above each column.

STAR Methods

RESOURCE AVAILABILITY

Lead Contact

Resources, reagents and further information requirement should be forwarded to and will be

responded by the Lead Contact, David I Stuart (dave@strubi.ox.ac.uk).

Materials Availability

Reagents generated in this study are available from the Lead Contact with a completed

Materials Transfer Agreement.

Data and Code Availability

The coordinates and structure factors of the crystallographic complexes are available from the

PDB with accession codes (see Table S1). Mabscape is available from

https://github.com/helenginn/mabscape, https://snapcraft.io/mabscape. The data that support

the findings of this study are available from the corresponding authors on request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Viral stocks

SARS-CoV-2/human/AUS/VIC01/2020 (Caly et al., 2020), SARS-CoV-2/B.1.1.7 and SARS-CoV-2/B.1.351 were provided by Public Health England, P.1 from a throat swab from Brazil were grown in Vero (ATCC CCL-81) cells. Cells were infected with the SARS-CoV-2 virus using an MOI of 0.0001. Virus containing supernatant was harvested at 80% CPE and spun at 3000 rpm at 4 °C before storage at -80 °C. Viral titres were determined by a focus-forming assay on Vero cells. Victoria passage 5, B.1.1.7 passage 2 and B.1.351 passage 4 stocks were sequenced to verify that they contained the expected spike protein sequence and no changes to the furin cleavage sites. The P.1 virus used in these studies contained the following mutations: L18F, T20N, P26S, D138Y, R190S, K417T, E464K, N501Y, D614G, H655Y, T1027I, V1176F. Passage 1 P.1 virus was sequence confirmed and contained no changes to the furin cleavage site.

Bacterial Strains and Cell Culture

Vero (ATCC CCL-81) cells were cultured at 37 °C in Dulbecco's Modified Eagle medium (DMEM) high glucose (Sigma-Aldrich) supplemented with 10% fetal bovine serum (FBS), 2 mM GlutaMAX (Gibco, 35050061) and 100 U/ml of penicillin–streptomycin. Human mAbs were expressed in HEK293T cells cultured in UltraDOMA PF Protein-free Medium (Cat# 12-727F, LONZA) at 37 °C with 5% CO_2. *E.coli DH5α* bacteria were used for transformation of plasmids encoding wt and mutated RBD proteins. A single colony was picked and cultured in LB broth with 50 μg mL^{-1} Kanamycin at 37 °C at 200 rpm in a shaker overnight. HEK293T (ATCC CRL-11268) cells were cultured in DMEM high glucose (Sigma-Aldrich) supplemented with 10% FBS, 1% 100X Mem Neaa (Gibco) and 1% 100X L-Glutamine

(Gibco) at 37 °C with 5% CO_2. To express RBD, RBD K417T, E484K, N501Y, RBD K417N,

RBD K417T, RBD E484K and ACE2, HEK293T cells were cultured in DMEM high glucose

(Sigma) supplemented with 2% FBS, 1% 100X Mem Neaa and 1% 100X L-Glutamine at 37

°C for transfection.

Participants

Participants were recruited through three studies: Sepsis Immunomics [Oxford REC C,

reference:19/SC/0296]), ISARIC/WHO Clinical Characterisation Protocol for Severe

Emerging Infections [Oxford REC C, reference 13/SC/0149] and the Gastro-intestinal illness

in Oxford: COVID sub study [Sheffield REC, reference: 16/YH/0247]. Diagnosis was

confirmed through reporting of symptoms consistent with COVID-19 and a test positive for

SARS-CoV-2 using reverse transcriptase polymerase chain reaction (RT-PCR) from an upper

respiratory tract (nose/throat) swab tested in accredited laboratories. A blood sample was taken

following consent at least 14 days after symptom onset. Clinical information including severity

of disease (mild, severe or critical infection according to recommendations from the World

Health Organisation) and times between symptom onset and sampling and age of participant

was captured for all individuals at the time of sampling.

P.1 virus from throat swabs. The International Reference Laboratory for Coronavirus at

FIOCRUZ (WHO) as part of the national surveillance for coronavirus had the approval of the

FIOCRUZ ethical committee (CEP 4.128.241) to continuously receive and analyze samples of

COVID-19 suspected cases for virological surveillance. Clinical samples (throat swabs)

containing P.1 were shared with Oxford University, UK under the MTA IOC FIOCRUZ 21-

02.

Sera from Pfizer vaccinees

Pfizer vaccine serum was obtained 7-17 days following the second dose of the BNT162b2

vaccine. Vaccinees were Health Care Workers, based at Oxford University Hospitals NHS

Foundation Trust, not known to have prior infection with SARS-CoV-2 and were enrolled in

the OPTIC Study as part of the Oxford Translational Gastrointestinal Unit GI Biobank Study

16/YH/0247 [research ethics committee (REC) at Yorkshire & The Humber – Sheffield]. The

study was conducted according to the principles of the Declaration of Helsinki (2008) and the

International Conference on Harmonization (ICH) Good Clinical Practice (GCP) guidelines.

Written informed consent was obtained for all patients enrolled in the study. Each received two

doses of COVID-19 mRNA Vaccine BNT162b2, 30 micrograms, administered

intramuscularly after dilution as a series of two doses (0.3 mL each) 18-28 days apart. The

mean age of vaccines was 43 years (range 25-63), 11 male and 14 female.

AstraZeneca-Oxford vaccine study procedures and sample processing

Full details of the randomized controlled trial of ChAdOx1 nCoV-19 (AZD1222), were

previously published (PMID: 33220855/PMID: 32702298). These studies were registered at

ISRCTN (15281137 and 89951424) and ClinicalTrials.gov (NCT04324606 and

NCT04400838). Written informed consent was obtained from all participants, and the trial is

being done in accordance with the principles of the Declaration of Helsinki and Good Clinical

Practice. The studies were sponsored by the University of Oxford (Oxford, UK) and approval

obtained from a national ethics committee (South Central Berkshire Research Ethics

Committee, reference 20/SC/0145 and 20/SC/0179) and a regulatory agency in the United

Kingdom (the Medicines and Healthcare Products Regulatory Agency). An independent

DSMB reviewed all interim safety reports. A copy of the protocols was included in previous

publications (PMID: 33220855/PMID: 32702298).

Data from vaccinated volunteers who received two vaccinations are included in this paper. Vaccine doses were either 5×10^{10} viral particles (standard dose; SD/SD cohort n=21) or half dose as their first dose (low dose) and a standard dose as their second dose (LD/SD cohort n=4). The interval between first and second dose was in the range of 8-14 weeks. Blood samples were collected and serum separated on the day of vaccination and on pre-specified days after vaccination e.g. 14 and 28 days after boost.

METHOD DETAILS

Focus Reduction Neutralization Assay (FRNT)

The neutralization potential of Ab was measured using a Focus Reduction Neutralization Test (FRNT), where the reduction in the number of the infected foci is compared to a negative control well without antibody. Briefly, serially diluted Ab or plasma was mixed with SARS-CoV-2 strain Victoria or P.1 and incubated for 1 hr at 37 °C. The mixtures were then transferred to 96-well, cell culture-treated, flat-bottom microplates containing confluent Vero cell monolayers in duplicate and incubated for a further 2 hrs followed by the addition of 1.5% semi-solid carboxymethyl cellulose (CMC) overlay medium to each well to limit virus diffusion. A focus forming assay was then performed by staining Vero cells with human anti-NP mAb (mAb206) followed by peroxidase-conjugated goat anti-human IgG (A0170; Sigma). Finally, the foci (infected cells) approximately 100 per well in the absence of antibodies, were visualized by adding TrueBlue Peroxidase Substrate. Virus-infected cell foci were counted on the classic AID EliSpot reader using **AID ELISpot software.** The percentage of focus reduction was calculated and IC_{50} was determined using the probit program from the SPSS package.

Cloning of ACE2 and RBD proteins

The constructs of EY6A Fab, 222 Fab, ACE2, WT RBD, B.1.1.7 and B.1.351 mutant RBD are the same as previously described (Dejnirattisai et al. 2021, Zhou et al., 2021, Supasa et al., 2021). To clone RBD K417T and RBD K417N, primers of RBD K417T (forward primer 5'-GGGCAGACCGGCACGATCGCCGACTAC-3' and reverse primer 5'-GTAGTCGGCGATCGTGCCGGTCTGCCC) and primers of RBD K417N (forward primer 5'-CAGGGCAGACCGGCAATATCGCCGACTACAATTAC-3' and reverse primer 5'-GTAATTGTAGTCGGCGATATTGCCGGTCTGCCCTG-3') were used separately, together with two primers of pNEO vector (Forward primer 5'- CAGCTCCTGGGCAACGTGCT-3' and reverse primer 5'- CGTAAAAGGAGCAACATAG-3') to do PCR, with the plasmid of WT RBD as the template. To clone P.1 RBD, the construct of B.1.351 RBD was used as the template and the primers of RBD K417T and of pNEO vector mentioned above were used to do PCR. Amplified DNA fragments were digested with restriction enzymes AgeI and KpnI and then ligated with digested pNEO vector. All constructs were verified by sequencing.

Protein production

Protein production was as described in Zhou et al. (Zhou et al., 2020). Briefly, plasmids encoding proteins were transiently expressed in HEK293T (ATCC CRL-11268) cells. The conditioned medium was dialysed and purified with a 5-ml HisTrap nickel column (GE Healthcare) and further polished using a Superdex 75 HiLoad 16/60 gel filtration column (GE Healthcare).

Bio-Layer Interferometry

BLI experiments were run on an Octet Red 96e machine (Fortebio). To measure the binding

affinity of ACE2 with P.1 RBD and affinities of monoclonal antibodies and ACE2 with native

RBD and, RBD K417N, RBD K417T, RBD E484K and RBD K417T E484K N501Y, eachP.1

RBD, each RBD was immobilized onto an AR2G biosensor (Fortebio). Monoclonal antibodies

(Dejnirattisai et al., 2021) were used as analytes or serial dilutions of ACE2 were used as

analytes. All experiments were run at 30 °C. Data were recorded using software Data

Acquisition 11.1 (Fortebio) and Data Analysis HT 11.1 (Fortebio) with a 1:1 fitting model used

for the analysis.

Antibody production

AstraZeneca and Regeneron antibodies were provided by AstraZeneca, Vir, Lilly and Adagio

antibodies were provided by Adagio. For the chimeric antibodies heavy and light chains of the

indicated antibodies were transiently transfected into 293Y cells and antibody purified purified

from supernatant on protein A.

Crystallization

818
ACE2 was mixed with P.1 RBD in a 1:1 molar ratio to a final concentration of 12.5 mg ml^{-1}.

EY6A Fab, 222 Fab and WT or mutant RBD were mixed in a 1:1:1 molar ratio to a final

concentration of 7.0 mg ml^{-1}. All samples were incubated at room temperature for 30 min.

Most crystallization experiments was set up with a Cartesian Robot in Crystalquick 96-well X

plates (Greiner Bio-One) using the nanoliter sitting-drop vapor-diffusion method, with 100 nl

of protein plus 100 nl of reservoir in each drop, as previously described (Water et al., 2003).

Crystallization of B.1.1.7 RBD/EY6A/222 complex was set up by hand pipetting, with 500 nl

of protein plus 500 nl of reservoir in each drop. Good crystals of EY6A Fab and 222 Fab

complexed with WT, K417T, K417N, B.1.1.7, B.1.351 or P.1 RBD were all obtained from

Hampton Research PEGRx 2 screen, condition 35, containing 0.15 M Lithium sulfate, 0.1 M

Citric acid pH 3.5, 18% w/v PEG 6,000. Crystals of P.1 RBD/ACE2 complex were formed in

Hampton Research PEGRx 1 screen, condition 38, containing 0.1 M Imidazole pH 7.0 and

20% w/v Polyethylene glycol 6,000.

X-ray data collection, structure determination and refinement

Crystals of ternary complexes of WT and mutant RBD/EY6A and 222 Fabs and the P.1.

RBD/ACE2 were mounted in loops and dipped in solution containing 25% glycerol and 75%

mother liquor for a second before being frozen in liquid nitrogen prior to data collection.

Diffraction data were collected at 100 K at beamline I03 of Diamond Light Source, UK. All

data (except some of the WT RBD-EY6A-222 Fab complex images) were collected as part of

an automated queue system allowing unattended automated data collection

(https://www.diamond.ac.uk/Instruments/Mx/I03/I03-Manual/Unattended-Data-

Collections.html). Diffraction images of 0.1° rotation were recorded on an Eiger2 XE 16M

detector (exposure time of either 0.004 or 0.006 s per image, beam size 80×20 μm, 100% beam

transmission and wavelength of 0.9763 Å). Data were indexed, integrated and scaled with the

automated data processing program Xia2-dials (Winter, 2010;Winter *et al.*, 2018). A data set

of 1080° was collected from 3 positions of a frozen crystal for the WT RBD-EY6A-222 Fab

complex. 720° of data was collected for each of the B.1.1.7, P.1 and B.1.351 mutant

RBD/EY6A and 222 Fab complexes (each from 2 crystals), and 360° for each of the K417N

and K417T RBD with EY6A and 222 Fabs, and ACE2 complexes was collected from a single

crystal.

Structures of WT RBD-EY6A-222 and the P.1 RBD-ACE2 complexes were determined by

molecular replacement with PHASER (McCoy et al., 2007) using search models of SARS-

CoV-2 RBD-EY6A-H4 (PDB ID 6ZCZ) (Zhou et al., 2020) and RBD-158 (PDB ID, 7BEK)

(Dejnirattisai et al., 2021) complexes, and a RBD and ACE2 complex (PDB ID, 6LZG (Wang

et al., 2020)), respectively. Model rebuilding with COOT (Emsley and Cowtan, 2004) and

refinement with PHENIX (Liebschner et al., 2019) were done for all the structures. The ChCl

domains of EY6A are flexible and have poor electron density. Data collection and structure

refinement statistics are given in **Table S1**. Structural comparisons used SHP (Stuart et al.,

1979), residues forming the RBD/Fab interface were identified with PISA (Krissinel and

Henrick, 2007) and figures were prepared with PyMOL (The PyMOL Molecular Graphics

System, Version 1.2r3pre, Schrödinger, LLC).

862
Quantification and statistical analysis

Statistical analyses are reported in the results and figure legends. Neutralization was measured

by FRNT. The percentage of focus reduction was calculated and IC_{50} was determined using

the probit program from the SPSS package. The Wilcoxon matched-pairs signed rank test was

used for the analysis and two-tailed P values were calculated and geometric mean values. BLI

data were analysed using Data Analysis HT 11.1 (Fortebio) with a 1:1 fitting model.

References

Alison Tarke, A., Sidney, J., Methot, N., Zhang, Y., Dan, J.M., Goodwin, B., Rubiro, P.,

Sutherland, A., da Silva Antunes, R., Frazier, A., et al. Negligible impact of SARS-CoV-2

variants on CD4 + and CD8 + T cell reactivity in COVID-19 exposed donors and vaccinees.

BioRxiv. 2021; https://doi.org/10.1101/2021.02.27.433180.

Baden, L.R., El Sahly, H.M., Essink, B., Kotloff, K., Frey, S., Novak, R., Diemert, D., Spector,

S.A., Rouphael, N., Creech, C.B., et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2

Vaccine. *N. Engl. J. Med*. 2020; **384**: 403-416.

Barnes, C.O., West, A.P., Huey-Tubman, K.E., Hoffmann, M.A.G., Sharaf, N.G., Hoffman,

P.R., Koranda, N., Gristick, H.B., Gaebler, C., Muecksch, F., et al. Structures of Human

Antibodies Bound to SARS-CoV-2 Spike Reveal Common Epitopes and Recurrent Features

of Antibodies. *Cell* 2020; **182**: 828–842.

Baum, A., Fulton, B.O., Wloga, E., Copin, R., Pascal, K.E., Russo, V., Giordano, S., Lanza,

K., Negron, N., Ni, M., et al. Antibody cocktail to SARS-CoV-2 spike protein prevents rapid

mutational escape seen with individual antibodies. *Science.* 2020; *369*: 1014–1018.

Caly, L., Druce, J., Roberts, J., Bond, K., Tran, T., Kostecki, R., Yoga, Y., Naughton, W.,

Taiaroa, G., Seemann, T., et al. Isolation and rapid sharing of the 2019 novel coronavirus

(SARS-CoV-2) from the first patient diagnosed with COVID-19 in Australia. *Med. J. Aust.*

2020; *212*: 459–462.

Cerutti, G., Guo, Y., Zhou, T., Gorman, J., Lee, M., Rapp, M., Reddem, E.R., Yu, J., Bahna,

F., Bimela, J., et al.. Potent SARS-CoV-2 Neutralizing Antibodies Directed Against Spike N-

Terminal   Domain   Target   a   Single   Supersite.   *BioRxiv.*   2021:   doi:

https://doi.org/10.1101/2021.01.10.426120

Dejnirattisai, W., Zhou, D., Ginn, H.M., Duyvesteyn, H.M.E., Supasa, P., Case, J.B., Zhao, Y.,

Walter, T.S., Mentzer, A.J., Liu, C., et al. The antigenic anatomy of SARS-CoV-2 receptor

binding domain. *Cell* 2021; doi: https://doi.org/10.1016/j.cell.2021.02.032

Domingo, E., and Perales, C. Viral quasispecies. *PLoS Genet.* 2019; **15:** e1008271.

https://doi.org/10.1371/journal.pgen.1008271

Donthu, N., and Gustafsson, A. Effects of COVID-19 on business and research. *J. Bus. Res.*

2020; **117**: 284–289.

Emary, K.R.W., Golubchik, T., Aley, P.K., Ariani, C. V., Angus, B.J., Bibi, S., Blane, B.,

Bonsall, D., Cicconi, P., Charlton, S., et al. Efficacy of ChAdOx1 nCoV-19 (AZD1222)

Vaccine Against SARS-CoV-2 VOC 202012/01 (B.1.1.7). *SSRN*. 2021; doi: https://doi.org/10.2139/ssrn.3779160.

Emsley, P., and Cowtan, K. Coot: Model-building tools for molecular graphics. *Acta Crystallogr. Sect. D Biol. Crystallogr.* 2004; **60**: 2126–2132.

Faria, N.R., Claro, I.M., Candido, D., Moyses Franco, L.A., Andrade, P.S., Coletti, T.M., Silva, C.A.M., Sales, F.C., Manuli, E.R., Aguiar, R.S., et al. (2021). Genomic characterisation of an emergent SARS-CoV-2 lineage in Manaus: preliminary findings - SARS-CoV-2 coronavirus / nCoV-2019 Genomic Epidemiology - *Virological*, *virological.org*. Available at: https://virological.org/t/genomic-characterisation-of-an-emergent-sars-cov-2-lineage-in-manaus-preliminary-findings/586 (Accessed: 3 March 2021).

Greaney, A.J., Starr, T.N., Gilchuk, P., Zost, S.J., Binshtein, E., Loes, A.N., Hilton, S.K., Huddleston, J., Eguia, R., Crawford, K.H.D., et al. Complete Mapping of Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape Antibody Recognition. *Cell Host Microbe.* 2021; **29**: 44-57.e9.

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Mü, M.A., Drosten, C., and Pö, S. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*. 2020; *181*: 271-280.e8.

Kemp, S.A., Collier, D.A., Datir, R.P., Ferreira, I.A.T.M., Gayed, S., Jahun, A., Hosmillo, M., Rees-Spear, C., Mlcochova, P., Lumb, I.U., et al. SARS-CoV-2 evolution during treatment of chronic infection. *Nature*. 2021; https://doi.org/10.1038/s41586-021-03291-y

Krammer, F. SARS-CoV-2 vaccines in development. *Nature*. 2020; **586**: 516–527.

Kreye, J., Reincke, S.M., Kornau, H.C., Sánchez-Sendin, E., Corman, V.M., Liu, H., Yuan, M., Wu, N.C., Zhu, X., Lee, C.C.D., et al. A Therapeutic Non-self-reactive SARS-CoV-2 Antibody Protects from Lung Pathology in a COVID-19 Hamster Model. *Cell*. 2020; **183**: 1058–1069.

Krissinel, E., and Henrick, K. Inference of Macromolecular Assemblies from Crystalline State.

J. Mol. Biol. 2007; **372**: 774–797.

Ku, Z., Xie, X., Davidson, E., Ye, X., Su, H., Menachery, V.D., Li, Y., Yuan, Z., Zhang, X.,

Muruato, A.E., et al. Molecular determinants and mechanism for antibody cocktail preventing

SARS-CoV-2 escape. *Nat. Commun.* 2021; **12**: 469.

Liebschner, D., Afonine, P. V., Baker, M.L., Bunkoczi, G., Chen, V.B., Croll, T.I., Hintze, B.,

Hung, L.W., Jain, S., McCoy, A.J., et al. Macromolecular structure determination using X-

rays, neutrons and electrons: Recent developments in Phenix. *Acta Crystallogr. Sect. D Struct.*

Biol. 2019; **75**: 861–877.

Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., Wu, H., Wang, W., Song, H., Huang, B., Zhu, N., et

al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for

virus origins and receptor binding. *Lancet.* 2020; **395**: 565–574.

Madhi, S.A., Baillie, V., Cutland, C.L., Voysey, M., Koen, A.L., Fairlie, L., Padayachee, S.D.,

Dheda, K., Barnabas, S.L., Bhorat, Q.E., et al. Safety and efficacy of the ChAdOx1 nCoV-19

(AZD1222) cOVID-19 vaccine against the B.1.351 variant in South Africa. *MedRxiv.* 2021;

doi: https://doi.org/10.1101/2021.02.10.21251247.

Mahase, E. (2021). Covid-19: Novavax vaccine efficacy is 86% against UK variant and 60%

against South African variant. *BMJ* 2021;372:n296

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read,

R.J. Phaser crystallographic software. *J. Appl. Crystallogr. 2007; 40*: 658–674.

Pinto, D., Park, Y.J., Beltramello, M., Walls, A.C., Tortorici, M.A., Bianchi, S., Jaconi, S.,

Culap, K., Zatta, F., De Marco, A., et al. Cross-neutralization of SARS-CoV-2 by a human

monoclonal SARS-CoV antibody. *Nature* 2020; **583**: 290–295.

Polack, F.P., Thomas, S.J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J.L.,

Pérez Marc, G., Moreira, E.D., Zerbini, C., et al. Safety and Efficacy of the BNT162b2 mRNA

Covid-19 Vaccine. *N. Engl. J. Med*. 2020; **383**: 2603–2615.

Seemann, T., Lane, C.R., Sherry, N.L., Duchene, S., Gonçalves da Silva, A., Caly, L., Sait, M.,

Ballard, S.A., Horan, K., Schultz, M.B., et al. Tracking the COVID-19 pandemic in Australia

using genomics. *Nat. Commun*. 2020; **11**: 1–9.

Shang, J., Ye, G., Shi, K., Wan, Y., Luo, C., Aihara, H., Geng, Q., Auerbach, A., and Li, F.

Structural basis of receptor recognition by SARS-CoV-2. *Nature*. 2020; **581**: 221–224.

Skelly, D.T., Harding Sir William, A.C., Gilbert-Jaramillo Sir William, J., Knight Sir William,

M.L., Longet, S., Brown, A., Adele, S., Adland, E., Brown, H., and Stafford, L. (2021).

Vaccine-induced immunity provides more robust heterotypic immunity than natural infection

to   emerging   SARS-CoV-2   variants   of   concern.   Research   Square.   2021;

https://doi.org/10.21203/rs.3.rs-226857/v1.

Starr, T.N., Greaney, A.J., Dingens 1, A.S., and Bloom, J.D. (2021). Complete map of SARS-

CoV-2 RBD mutations that escape the monoclonal antibody LY-CoV555 and its cocktail with

LY-CoV016. B*ioRxiv*. 2021; https://doi.org/10.1101/2021.02.17.431683.

Stuart, D.I., Levine, M., Muirhead, H., and Stammers, D.K. Crystal structure of cat muscle

pyruvate kinase at a resolution of 2.6 Å. *J. Mol. Biol*. 1979; *134***:** 109–142.Supasa, P., Zhou,

D., Dejnirattisai, W., Liu, C., Mentzer, A.J., Ginn, H.M., Zhao, Y., Duyvesteyn, H.M.E.,

Nutalai, R., Tuekprakhon, A., et al. (2021). Reduced neutralization of SARS-CoV-2 B.1.1.7

variant   by   convalescent   and   vaccine   sera.   *Cell*

2021;doi:https://doi.org/10.1016/j.cell.2021.02.033.

Tatusov, R.L., Galperin, M.Y., Natale, D.A., and Koonin, E. V. The COG database: A tool for

genome-scale analysis of protein functions and evolution. *Nucleic Acids Res*. 2000; **28:** 33–36.

Volz, E., Hill, V., McCrone, J.T., Price, A., Jorgensen, D., O'Toole, Á., Southgate, J., Johnson,

R., Jackson, B., Nascimento, F.F., et al. Evaluating the Effects of SARS-CoV-2 Spike Mutation

D614G on Transmissibility and Pathogenicity. *Cell*. 2021; *184***:** 64-75.e11.

Voysey, M., Clemens, S.A.C., Madhi, S.A., Weckx, L.Y., Folegatti, P.M., Aley, P.K., Angus,

B., Baillie, V.L., Barnabas, S.L., Bhorat, Q.E., et al. Safety and efficacy of the ChAdOx1

nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised

controlled trials in Brazil, South Africa, and the UK. *Lancet.* 2020; **397**: 99-111.

Walls, A.C., Park, Y.J., Tortorici, M.A., Wall, A., McGuire, A.T., and Veesler, D. Structure,

Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell*. 2020; **181**: 281-

292.e6.

Wang, P., Wang, M., Yu, J., Cerutti, G., Nair, M.S., Huang, Y., Kwong, P.D., Shapiro, L., *and*

Ho, D.D. Increased Resistance of SARS-CoV-2 Variant P.1 to Antibody Neutralization.

bioRxiv 2021; doi: https://doi.org/10.1101/2021.03.01.433466.

Wang, Q., Zhang, Y., Wu, L., Niu, S., Song, C., Zhang, Z., Lu, G., Qiao, C., Hu, Y., Yuen,

K.Y., et al. Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2.

Cell. 2020; **181**: 894-904.e9.

Wibmer, C.K., Ayres, F., Hermanus, T., Madzivhandila, M., Kgagudi, P., Oosthuysen, B.,

Lambson, B.E., de Oliveira, T., Vermeulen, M., van der Berg, K., et al. SARS-CoV-2 501Y.V2

escapes neutralization by South African COVID-19 donor plasma.

bioRxiv. 2021; doi: https://doi.org/10.1101/2021.01.18.427166

Winter, G. Xia2: An expert system for macromolecular crystallography data reduction. *J. Appl.*

Crystallogr. 2010; **43**: 186–190.

Winter, G., Waterman, D.G., Parkhurst, J.M., Brewster, A.S., Gildea, R.J., Gerstel, M.,

Fuentes-Montero, L., Vollmar, M., Michels-Clark, T., Young, I.D., et al. *DIALS*:

implementation and evaluation of a new integration package. *Acta Crystallogr. Sect. D Struct.*

Biol. 2018; **74**: 85–97.

Wu, Y., Wang, F., Shen, C., Peng, W., Li, D., Zhao, C., Li, Z., Li, S., Bi, Y., Yang, Y., et al.

A noncompeting pair of human neutralizing antibodies block COVID-19 virus binding to its

receptor ACE2. *Science.* 2020; *368*: 1274–1278.

Yuan, M., Liu, H., Wu, N.C., Lee, C.C.D., Zhu, X., Zhao, F., Huang, D., Yu, W., Hua, Y.,

Tien, H., et al. (2020). Structural basis of a shared antibody response to SARS-CoV-2. Science.

2020; **369**: 1119–1123.

Zahradník, J., Marciano, S., Shemesh, M., Zoler, E., Chiaravalli, J., Meyer, B., Dym, O., Elad,

N., and Schreiber, G. SARS-CoV-2 RBD in vitro evolution follows contagious mutation

spread, yet generates an able infection inhibitor. *BioRxiv*. 2021;

https://doi.org/2021.01.06.425392.

Zhou, D., Duyvesteyn, H.M.E., Chen, C.P., Huang, C.G., Chen, T.H., Shih, S.R., Lin, Y.C.,

Cheng, C.Y., Cheng, S.H., Huang, Y.C., et al. Structural basis for the neutralization of SARS-

CoV-2 by an antibody from a convalescent patient. *Nat. Struct. Mol. Biol*. 2020; **27**: 950–958.

Zhou, D., Dejnirattisai, W., Supasa, P., Liu, C., Mentzer, A.J., Ginn, H.M., Zhao, Y.,

Duyvesteyn, H.M.E., Tuekprakhon, A., Nutalai, R., et al. (2021). Evidence of escape of SARS-

CoV-2 variant B.1.351 from natural and vaccine induced sera. *Cell*. 2021;

https://doi.org/10.1016/j.cell.2021.02.037.

Zost, S.J., Gilchuk, P., Case, J.B., Binshtein, E., Chen, R.E., Nkolola, J.P., Schäfer, A., Reidy,

J.X., Trivette, A., Nargi, R.S., et al. Potently neutralizing and protective human antibodies

against SARS-CoV-2. Nature 2020; **584**: 443–444.

1020

1023

1024

Figure2

Figure3

1025

1026

1027

1028

Figure4

1029

1030

Figure5

Figure7