Programmed mechano-chemical coupling in reaction-diffusion active matter

Anis Senoussi,* Jean-Christophe Galas,* and André Estevez-Torres*

Sorbonne Université, CNRS, Institut de Biologie Paris-Seine (IBPS), Laboratoire Jean Perrin (LJP), F-75005, Paris

E-mail: anis.senoussi@sorbonne-universite.fr; jean-christophe.galas@sorbonne-universite.fr; andre.estevez-torres@sorbonne-universite.fr

Abstract

Embryo morphogenesis involves a complex combination of pattern-forming mechanisms. However, classical *in vitro* patterning experiments explore only one mechanism at a time, thus missing coupling effects. Here, we conjugate two major pattern-forming mechanisms —reaction-diffusion and active matter— by integrating dissipative DNA/enzyme reaction networks within an active gel composed of cytoskeletal motors and filaments. We show that the strength of the flow generated by the active gel controls the mechano-chemical coupling between the two subsystems. We use this property to engineer the mechanical activation of chemical reaction networks both in time and space, thus mimicking key aspects of the polarization mechanism observed in *C. elegans* oocytes. We anticipate that reaction-diffusion active matter may be useful to investigate mechano-chemical transduction and to design new materials with life-like properties.

Keywords: DNA programming, active matter, morphogenesis, pattern formation, life-like material

Short title: Programmable reaction-diffusion active matter

Living embryos get their shape through a complex combination of chemical and physical processes that take place out of equilibrium. Basically, biochemical reaction networks process information at particular points of space and time, while active gels generate mechanical forces and flows.[1] These two generic processes are intertwined through diverse chemo-mechanical and mechano-chemical couplings and both continuously consume chemical energy. For instance, the pathway Rho GTPase brings about local contractions of the actomyosin cortex in *Xenopus*,[2] which regulates the cell cycle, while cytoplasmic flows of this same cortex trigger the PAR reaction network in *C. elegans*, inducing embryo polarization.[3]

The development of *in vitro* dissipative molecular systems is key to produce non-equilibrium materials and to investigate these processes in a controlled environment for testing theoretical predictions.[4,5] On the chemical side, *in vitro* out-of-equilibrium reaction networks produce spatio-temporal concentration patterns through reaction or reaction-diffusion instabilities.[6–14] On the mechanical side, active gels made of cytoskeletal motors and filaments reconstituted *in vitro*[4,15–24] convert chemical energy into mechanical work and generate static patterns and flows through hydrodynamic instabilities.[25] More recently, efforts have focused on coupling reaction networks with active fluids and gels, resulting in redox-controlled self-oscillating gels[26] and DNA-controlled passive[27] and active gels.[28–30] However, so far, only one of the two systems could be maintained out of equilibrium, either the reaction network,[26] or the gel,[27–31] and thus only one of them could exhibit spatio-temporal self-organization. This constraint suppresses the rich variety of mechano-chemical couplings that are essential for pattern generation during development.[1] It thus constitutes a major limitation for the design of self-shaping materials inspired from embryogenesis.[5]

To create a functional mechano-chemical dissipative material we assembled two dissipative molecular subsystems, a chemical and a mechanical one. The chemical subsystem is made of a network of DNA/enzyme reactions that produce single-stranded DNA (ssDNA) molecules[13] (Figure 1a). The mechanical subsystem is an active gel composed of bundles of protein filaments propulsed by molecular motors[15,17] (Figure 1b). Each subsystem is

Figure 1: In a reaction-diffusion active matter system, the coupling between the mechanical and the chemical subsystem is controlled by the strength of the active flow. (a) Scheme of the chemical subsystem involving the autocatalytic amplification of DNA strand **A** in the presence of enzymes and template strand T_{AA}. Harpoon-ended arrows denote ssDNA. (b) Cartoon of the mechanical subsystem: an active gel formed by microtubules bundled together by a depletion agent and clusters of kinesin-1 motors. (c) Sketch of the channel in which the front propagation and the active gel dynamics were observed by fluorescence microscopy. Initially, **A** (yellow) is present only on the left side and the microtubule bundles (light blue) are aligned along x. (d) A mechano-chemical coupling between the two subsystems is achieved by increasing the strength of the flows generated by the active gel, which induces four different microtubule structures (light blue) and two DNA patterns (yellow). The white arrows represent the hydrodynamic flows generated by the active gel. Fluorescence images of the microtubules are represented for each morphology. Scale bars are 0.5 mm.

maintained out of equilibrium *via* the hydrolysis of high-energy compounds, respectively

deoxynucleosidetriphosphates (dNTPs) and adenosinetriphosphate (ATP).

The chosen chemical subsystem has four advantages. Firstly, due to DNA hybridization

rules, it can be easily reprogrammed into a variety of dissipative dynamics such as oscilla-

tions,[32] bistability and excitability.[33] Secondly, it can be maintained out of equilibrium in a

closed reactor for days.[34] Thirdly, working in water at pH 7, it is *a priori* compatible with

other biochemical reactions.[35] Lastly, it generates a variety of reaction-diffusion patterns

such as traveling fronts,[36] waves[14] and stationary patterns.[37] In a first series of experiments,

the chemical subsystem encoded an autocatalytic loop that produces ssDNA species **A** —

the node— in the presence of ssDNA $\mathbf{T}_{AA}$ —the template—, a polymerase and a nicking

enzyme[13] (Figure 1a).

In the mechanical subsystem, the bundles are constituted of stabilized microtubule fil-

aments assembled together by attracting forces generated by the presence of a depletion

agent[17] (Figure 1b). The motors are clusters of kinesin-1 and thus can bind several micro-

tubules at once.† Such an active gel generates macroscopic flows that, depending on the

concentration of motors and filaments, produce a diversity of microtubule morphologies:[4,5]

local contractions,[15] corrugations,[24] chaotic flows,[17] and global contractions.[21] In the follow-

ing, we demonstrate that, when mixed together, the two subsystems retain their ability to

undergo, respectively, chemical and mechanical instabilities that generate spatio-temporal

patterns. We further show that the strength of the active flow generated by the mechani-

cal subsystem controls the mechanochemical coupling between the two subsystems. Finally,

we take advantage of this property to design materials that mimick crucial aspects of a

mechanochemical patterning mechanism observed in *C. elegans* embryo.

To check whether the two subsystems remained functional when combined in an opti-

mized buffer (Figure S1), we tested the propagation of a DNA front through and active gel

undergoing local contractions. To do so, a solution containing all the components of the

chemical and mechanical subsystems, except the strand **A**, were filled into a microchannel.*

An initial condition containing the same solution supplemented with **A** was injected on the

left side of the channel (Figure 1c). We recorded the spatiotemporal dynamics of each subsys-

tem by fluorescence microscopy thanks to the presence of a DNA intercalator that becomes

fluorescent upon binding to double-stranded DNA and of fluorescently-labeled microtubules

(SI Methods). In the chemical subsystem, we observed the propagation of a front of DNA

---

†Two types of clusters were used: biotinylated kinesin-1 (from *D. melanogaster*) assembled together by streptavidin,[15] or clusters made from SNAPtag modified kinesin-1 (from *R. norvegicus*) that spontaneously multimerize[24] (SI Methods).

*The experimental conditions for each figure are provided in Tables S2-S5.

fluorescence with constant velocity $v_c = 20$ μm/min across the whole length of the active gel,
i.e. > 1 cm (Figures 1d, 2a, S2 and Movie S1). Concomitantly, in the mechanical subsystem
the microtubules contracted locally with a characteristic time $\tau_m = 50$ min into aggregates
with a typical size of $100 - 500$ μm. The observation of a DNA front and microtubule ag-
gregates is in agreement with previous reports for each subsystem taken independently.[15,36]

Figure 2: A DNA/enzyme reaction-diffusion front propagates normally inside a locally-
contracting cytoskeletal active gel and the dynamics of each subsystem can be independently
tuned. (a) Time-lapse 2-color image of the fluorescence intensities associated to species **A**
(yellow) and to the microtubule network (light blue) (see also Movie S1). The dotted line
indicates the velocity of the chemical front, v_c. Plots of v_c (yellow disks) and contraction
time of the active gel, τ_m, (blue crosses) for different concentrations of DNA polymerase (b)
and motors (c). The lines are fits to the data with $v_c \sim [\mathrm{pol}]^{1/2}$ (dotted line), $\tau_m \sim [\mathrm{motor}]^{-1}$
(dashed line) and $\tau_m \sim [\mathrm{motor}]^{-2}$ (plain line). Error bars correspond to one standard
deviation from a triplicate experiment.

When the active gel produces local contractions, the dynamics of each subsystem can
be independently tuned. Increasing the polymerase concentration, [pol], increases the front
velocity, v_c, until reaching a plateau at 22 μm/min (Figure 2b). In these conditions, the
characteristic contraction time τ_m remained constant. We find a scaling $v_c \sim [\mathrm{pol}]^{1/2}$, in
agreement with previous results[36] and characteristic of Luther reaction-diffusion dynamics
where $v_c \sim r_c^{1/2}$, taking $r_c \sim [\mathrm{pol}]$ for the rate of the autocatalytic reaction as observed

in previous experiments.[36] In turn, when the motor concentration, [motor], increases, τ_m decreases until reaching a plateau at 100 min (Figure 2c) and the size of microtubule aggregates increases (Figure S3), while v_c remains constant. A hydrodynamic model of a contracting active gel[5] yields $\tau_m^{-1} \sim \zeta([\text{motor}])$, where ζ is the strength of the gel activity, which depends on the motor concentration, and two scalings are found in the literature[18,38] yielding $\tau_m \sim [\text{motor}]^{-\alpha}$, with $\alpha = 1, 2$ (Supplementary Text). Our data are compatible with both scalings (Figure 2c). In summary, Figure 2 shows that the two subsystems are both chemically and mechanically decoupled when the gel undergoes local contractions.

By varying the conditions, we can propagate the chemical front through active gels undergoing other spatial instabilities associated with different active flow strengths, as sketched in Figure 1d. When the dGTP concentration was reduced (Figures S4-S5), chaotic flows were observed in the mechanical subsystem during several hours before local contractions occurred (Movie S2). Such flows did not modify the velocity of the chemical front because transport remained dominated by Brownian diffusion.[*] When the length of microtubules was increased using taxol and the motor concentration reduced, the active gel formed corrugations reminiscent of those previously reported in the absence of the chemical subsystem,[24] again without perturbing the chemical front (Movie S3).

In contrast, a dramatic perturbation of the front propagation was observed when the active gel underwent a global contraction, associated with large hydrodynamic flows (Figures 3 and S6-S11 and Movie S4). Global contractions were observed for long microtubules and relatively high motor concentrations (Figure S7). When the gel contracted more rapidly than the front propagated, the front moved faster and we distinguished 4 phases (Figure 3c). During phase I, the active gel contracted rapidly towards the center of the channel, accelerating until reaching a maximum velocity $v_m = 400\ \mu\text{m/min}$ at the end of phase I and dragging DNA along, which formed a detached DNA islet ahead of the front. During phase II the gel

---

[*]The diffusivity of **A** due to the active flow, D_f, was estimated to be 10-fold smaller than the Brownian diffusivity of **A**, D_A, and thus $v_c \sim (D_A + D_f)^{1/2} \approx D_A^{1/2}$ corresponds to a purely reaction-diffusion front (Supplementary Text).

Figure 3: A globally-contracting active gel stretches and accelerates a reaction-diffusion front. (a) Time-lapse, 2-color fluorescence image with the DNA front in yellow and the microtubules in light blue. The inset shows the two fluorescence channels in separate images for the selected region. The white and yellow spots are dust particles concentrated by the contracting gel (see also Movie 4). (b) Position of the DNA front along x for different motor concentrations. (c) Time-lapse images of DNA fluorescence (color) at [motor] = 2.5 nM and 3 min per image. The extremities of the contracting gel are indicated with black markers. Roman numbers indicate the 4 phases described in the text. (d) Stroboscopic image averaged over 50 min showing the trajectories of fluorescent beads during gel contraction, the white arrows indicate the sense of the flow (see also Movie 5) and (e) plot of the bead velocity along x across the width of the channel for the beads in the red rectangle, 25 min after the beginning of the contraction. (f) Maximal front velocity during phase I (crosses) and steady-state velocity during phase IV (disks) for different motor concentrations. (g) Maximal front velocity during phase I $vs.$ maximal gel contraction velocity. Error bars correspond to one standard deviation from a triplicate experiment.

decelerated and the DNA islet was diluted in the xy plane resulting in a front with a skewed profile 5-fold wider than the initial one (Figure S6). Throughout phases I and II the Péclet number was greater than 1 (Figure S8), indicating that active convection predominated over diffusion. Finally, when the active gel stopped contracting, the DNA front slowly recovered

a sigmoidal shape (phase III, Figure S6) and eventually reached a steady state with constant velocity and width (phase IV).

In a control experiment with a passive dye, only phases I and II were observed, indicating that reaction was necessary for phases III and IV and that DNA islet formation was not related to the binding of DNA to the active gel (Figure S10). We confirmed the last interpretation by adding passive brownian beads to measure the hydrodynamic flow induced during contraction. We observed two counter-rotating fluid rolls, symmetric along the central axis of the channel, x, and producing water flows along x that reached $+150$ μm/min in the center of the channel and -100 μm/min at its borders (Figures 3d,e and S11 and Movie S5). Taken together, these results indicate that the stretching of the concentration profile of $\mathbf{A}$ leading to the formation of the DNA islet during phase I was a purely hydrodynamic process.

The active gel contraction velocity, v_m, was a sigmoidal function of the motor concentration (Figure S7). We quantified the two main regimes of front propagation with $\max(v_c^I)$ the maximal velocity during phase I and v_c^{IV} the velocity at steady state in phase IV. Figure 3f shows that the former strongly depended on the motor concentration while the latter was independent. Finally, the linear relationship between $\max(v_c^I)$ and v_m is consistent with the observation of a convection-dominated transport during phase I (Figure 3g). Taken together, these results show that when the flows generated by the active gel are sufficiently fast there is a mechano-chemical coupling between the gel and the reaction-diffusion front. This coupling happens through hydrodynamics and can be interpreted as a time-dependent Taylor dispersion.[39]

We have just seen that active flows can significantly modify heterogeneous concentration profiles present in the chemical subsystem. Can they induce an asymmetry in an initially homogeneous chemical subsystem? This is what happens in the *C. elegans* embryo, where the active flow generated by the actomyosin cortex breaks the symmetry of an initially homogeneous distribution of PAR-proteins. Later, this asymmetry is amplified by a PAR-dependent bistable reaction network, leading to embryo polarization.[3] The reaction-diffusion

active matter system developed here is a good candidate to mimick this process in a synthetic material. To do so, we first need to implement a mechanism that couples a variation in the microtubule concentration with a change in the concentration of a DNA species that is inatially homogeneously distributed and second to engineer a chemical network that amplifies this concentration change.

The first requirement was fulfilled by attaching DNA strands to $\sim 30~\mu$m diameter hydrogel beads, which were trapped by the microtubule mesh and concentrated during contraction. The second condition was satisfied by engineering a chemical subsystem whose kinetics depend on the concentration of DNA-bead conjugates, and thus on the contraction state of the gel. More precisely, the DNA autocatalytic loop was split into two nodes, $\mathbf{B}$ and $\mathbf{D}$, that cross-activate each other thanks to the templates $\mathbf{T}_{\mathrm{BD}}$ and $\mathbf{T}_{\mathrm{DB}}$ (Figure 4a). By attaching each of these templates to a set of hydrogel beads and supplementing the medium with an exonuclease that degrades $\mathbf{B}$ and $\mathbf{D}$ (Figure 4a) the cross-catalysis kinetics become diffusion-controlled[40] and thus should depend on bead density. As a result, the beads brought together in a contracted gel should activate faster, producing DNA that light them up in the presence of a DNA intercalator dye (Figure 4b). Indeed, when both types of beads where embedded in the active gel in the presence of a homogeneous, low concentration of $\mathbf{D}$, they first reached a high density as the gel contracted and later they became fluorescent (Figures 4c,d, S13 and Movie S6). In the absence of contraction, the bead fluorescence amplification was delayed (Figure S12) and its final amplitude reduced, both by a factor 2 (Figure 4d). As expected for diffusion-controlled kinetics, the mechano-chemical DNA amplification dynamics slowed down with increasing exonuclease concentration (Figure 4d).

To show that the activated beads may trigger downstream reactions in solution, the previous system was supplemented with freely-diffusing templates $\mathbf{T}_{\mathrm{BA}}$ and $\mathbf{T}_{\mathrm{AA}}$, that respectively convert $\mathbf{B}$ into $\mathbf{A}$ and sustain the autocatalytic reaction of $\mathbf{A}$ described earlier. In addition, to suppress the undesired self-activation of $\mathbf{T}_{\mathrm{AA}}$ a repressor strand $\mathbf{R}_{\mathrm{A}}$ was added[33] (Figure S15). In the presence of motors, the beads were all activated within 1 h at

Figure 4: A globally-contracting active gel triggers the activation of downstream reaction networks with temporal and spatial control. (a) Scheme of the cross-autocatalytic DNA/enzyme network (top). Plain and dotted arrows indicate activation and degradation reactions, respectively. Harpoon-ended arrows correspond to ssDNA and disks indicate hydrogel beads carrying templates T_{ij}. Detailed mechanism of bead activation where the diffusion of B and D between beads is indicated (bottom). (b) Sketch of the mechano-chemical activation of the reaction network in panel a through the contraction of the active gel (light blue) that brings the hydrogel beads (disks) close together, speeding up cross-catalysis. (c) Time-lapse images of DNA fluorescence from the template-bearing beads embedded in the active gel in the presence of motors. White dotted lines indicate the borders of the active gel and the channel walls are depicted in gray. (d) Average DNA fluorescence over the whole channel *vs.* time in the absence (dotted line) and in the presence (plain line) of motors for different exonuclease concentrations (colors). (e) Scheme of the bead-associated cross-autocatalytic network coupled to the autocatalysis of A in solution. Disks indicate templates linked to hydrogel beads. The blunt-ended arrow indicates repression. (f) Time-lapse images of DNA fluorescence in the channel for the network in panel e, in the presence (top) and in the absence (bottom) of motors. The bright spots are the beads, the arrows indicate the start of the fronts of A, in yellow.

the center of the channel, where the gel contracted, and they triggered a controlled front of **A** that propagated from the center of the channel to its extremities (Figures 4f and S14-S15 and Movie S7). In contrast, in the absence of gel contraction, the beads randomly activated over the course of 5h, which was followed by the uncontrolled amplification of **A**. Taken together, these results demonstrate that mechano-chemical coupling can be engineered to trigger either temporal or spatio-temporal chemical instabilities in a synthetic material.

The coupling of chemical and mechanical self-organization is a key ingredient of biological complexity, in particular during embryogenesis. We have demonstrated that it is possible to couple two archetypal examples of these mechanisms, reaction-diffusion and active matter, in a synthetic material. Our design is modular because it relies on two distinct subsystems with well-characterized and predictable spatiotemporal behaviors: DNA/enzyme reactions and kinesin/microtubule active gels. Considered independently, each subsystem reveals complex dynamics and macroscopic organizations which are subject to intense scrutiny.[14,15,17,19,21,24,35,37] When mixed-together, the coupling strength between the two subsystems is set by the magnitude of the flow generated by the active gel. As a result, this system may be useful for investigating self-organization when chemical and mechanical out-of-equilibrium processes are intertwined. Finally, reaction-diffusion active matter provides a framework for the rational engineering of functional out-of-equilibrium materials with life-like properties. On the one hand, it could be advantageously combined with the wide array of methods in DNA nanotechnology, such as nanostructure design,[41] logic gates,[11] analyte detection[42] or hydrogel swelling.[27] On the other hand, by using DNA-motor conjugates[28–30] or photosensitive motors[43] the system is extendable to chemo-mechanical as well as photo-mechanical couplings.

Acknowledgements

We thank H. Berthoumieux, V. Bormuth, M. Elez, A. Genot, G. Gines, N. Lobato-Dauzier, A. Maitra, L. Robert, Y. Rondelez and R. Voituriez for insightful discussions and K. Furuta

and Z. Gueroui for their kind gift of kinesin plasmids. This work has been funded by the European Research Council (ERC) under the European's Union Horizon 2020 programme (grant No 770940, A.E.-T.), by the Ville de Paris Emergences programme (Morphoart, A.E.-T.) and by MITI CNRS (J.-C. G.). The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

(1) Gross, P.; Kumar, K. V.; Grill, S. W. How Active Mechanics and Regulatory Biochemistry Combine to Form Patterns in Development. *Annual Review of Biophysics* **2017**, *46*, 337–356.

(2) Bement, W. M.; Leda, M.; Moe, A.; Kita, A.; Larson, M.; Golding, A.; Pfeuti, C.; Su, K.-C.; Miller, A.; Goryachev, A.; von Dassow, G. Activator–inhibitor coupling between Rho signalling and actin assembly makes the cell cortex an excitable medium. *Nature Cell Biology* **2015**, *17*, 1471–1483.

(3) Goehring, N. W.; Trong, P. K.; Bois, J. S.; Chowdhury, D.; Nicola, E. M.; Hyman, A. A.; Grill, S. W. Polarization of PAR proteins by advective triggering of a pattern-forming system. *Science* **2011**, *334*, 1137–1141.

(4) Needleman, D.; Dogic, Z. Active matter at the interface between materials science and cell biology. *Nature reviews materials* **2017**, *2*, 17048.

(5) Senoussi, A.; Vyborna, Y.; Berthoumieux, H.; Galas, J.-C.; Estevez-Torres, A. In *Out-of-Equilibrium Supramolecular Systems and Materials*; Giuseppone, N., Walther, A., Eds.; Wiley-VCH, in press.

(6) Epstein, I.; Pojman, J. A. *An introduction to nonlinear chemical reactions*; Oxford University Press: New York, 1998.

(7) Isalan, M.; Lemerle, C.; Serrano, L. Engineering gene networks to emulate Drosophila embryonic pattern formation. *PLoS biology* **2005**, *3*.

(8) Karzbrun, E.; Tayar, A. M.; Noireaux, V.; Bar-Ziv, R. H. Programmable on-chip DNA compartments as artificial cells. *Science* **2014**, *345*, 829–832.

(9) Nakajima, M.; Imai, K.; Ito, H.; Nishiwaki, T.; Murayama, Y.; Iwasaki, H.; Oyama, T.; Kondo, T. Reconstitution of Circadian Oscillation of Cyanobacterial KaiC Phosphorylation in Vitro. *Science* **2005**, *308*, 414–415.

(10) Loose, M.; Fischer-Friedrich, E.; Ries, J.; Kruse, K.; Schwille, P. Spatial regulators for bacterial cell division self-organize into surface waves in vitro. *Science* **2008**, *320*, 789–792.

(11) Zhang, D. Y.; Seelig, G. Dynamic DNA nanotechnology using strand-displacement reactions. *Nature chemistry* **2011**, *3*, 103.

(12) Chirieleison, S. M.; Allen, P. B.; Simpson, Z. B.; Ellington, A. D.; Chen, X. Pattern transformation with DNA circuits. *Nature chemistry* **2013**, *5*, 1000.

(13) Montagne, K.; Plasson, R.; Sakai, Y.; Fujii, T.; Rondelez, Y. Programming an in vitro DNA oscillator using a molecular networking strategy. *Molecular systems biology* **2011**, *7*.

(14) Padirac, A.; Fujii, T.; Estévez-Torres, A.; Rondelez, Y. Spatial waves in synthetic biochemical networks. *Journal of the American Chemical Society* **2013**, *135*, 14586–14592.

(15) Nedelec, F.; Surrey, T.; Maggs, A. C.; Leibler, S. Self-organization of microtubules and motors. *Nature* **1997**, *389*, 305.

(16) Bendix, P. M.; Koenderink, G. H.; Cuvelier, D.; Dogic, Z.; Koeleman, B. N.; Brieher, W. M.; Field, C. M.; Mahadevan, L.; Weitz, D. A. A quantitative analysis

of contractility in active cytoskeletal protein networks. *Biophysical journal* **2008**, *94*, 244 3126–3136.

(17) Sanchez, T.; Chen, D. T.; DeCamp, S. J.; Heymann, M.; Dogic, Z. Spontaneous motion 246 in hierarchically assembled active matter. *Nature* **2012**, *491*, 431.

(18) Foster, P. J.; Fürthauer, S.; Shelley, M. J.; Needleman, D. J. Active contraction of 248 microtubule networks. *Elife* **2015**, *4*, e10837.

(19) Wu, K.-T.; Hishamunda, J. B.; Chen, D. T.; DeCamp, S. J.; Chang, Y.-W.; Fernández-250 Nieves, A.; Fraden, S.; Dogic, Z. Transition from turbulent to coherent flows in confined 251 three-dimensional active fluids. *Science* **2017**, *355*, eaal1979.

(20) Kumar, N.; Zhang, R.; de Pablo, J. J.; Gardel, M. L. Tunable structure and dynamics 253 of active liquid crystals. *Science advances* **2018**, *4*, eaat7779.

(21) Torisawa, T.; Taniguchi, D.; Ishihara, S.; Oiwa, K. Spontaneous formation of a glob-255 ally connected contractile network in a microtubule-motor system. *Biophysical journal* 256 **2016**, *111*, 373–385.

(22) Ideses, Y.; Erukhimovitch, V.; Brand, R.; Jourdain, D.; Hernandez, J. S.; Gabinet, U.; 258 Safran, S.; Kruse, K.; Bernheim-Groswasser, A. Spontaneous buckling of contractile 259 poroelastic actomyosin sheets. *Nature communications* **2018**, *9*, 2461.

(23) Roostalu, J.; Rickman, J.; Thomas, C.; Nédélec, F.; Surrey, T. Determinants of polar 261 versus nematic organization in networks of dynamic microtubules and mitotic motors. 262 *Cell* **2018**, *175*, 796–808.

(24) Senoussi, A.; Kashida, S.; Voituriez, R.; Galas, J.-C.; Maitra, A.; Estevez-Torres, A. 264 Tunable corrugated patterns in an active nematic sheet. *Proceedings of the National* 265 *Academy of Sciences* **2019**, *116*, 22464–22470.

(25) Marchetti, M. C.; Joanny, J.-F.; Ramaswamy, S.; Liverpool, T. B.; Prost, J.; Rao, M.; Simha, R. A. Hydrodynamics of soft active matter. *Reviews of Modern Physics* **2013**, *85*, 1143.

(26) Yoshida, R.; Takahashi, T.; Yamaguchi, T.; Ichijo, H. Self-oscillating gel. *Journal of the American Chemical Society* **1996**, *118*, 5134–5135.

(27) Cangialosi, A.; Yoon, C.; Liu, J.; Huang, Q.; Guo, J.; Nguyen, T. D.; Gracias, D. H.; Schulman, R. DNA sequence–directed shape change of photopatterned hydrogels via high-degree swelling. *Science* **2017**, *357*, 1126–1130.

(28) Wollman, A. J.; Sanchez-Cano, C.; Carstairs, H. M.; Cross, R. A.; Turberfield, A. J. Transport and self-organization across different length scales powered by motor proteins and programmed by DNA. *Nature nanotechnology* **2014**, *9*, 44.

(29) Sato, Y.; Hiratsuka, Y.; Kawamata, I.; Murata, S.; Nomura, S.-i. M. Micrometer-sized molecular robot changes its shape in response to signal molecules. *Sci. Robot* **2017**, *2*.

(30) Keya, J. J.; Suzuki, R.; Kabir, A. M. R.; Inoue, D.; Asanuma, H.; Sada, K.; Hess, H.; Kuzuya, A.; Kakugo, A. DNA-assisted swarm control in a biomolecular motor system. *Nature communications* **2018**, *9*, 453.

(31) Bertrand, O. J. N.; Fygenson, D. K.; Saleh, O. A. Active, motor-driven mechanics in a DNA gel. *Proceedings of the National Academy of Sciences* **2012**, *10.1073/pnas.1208732109*.

(32) Fujii, T.; Rondelez, Y. Predator–prey molecular ecosystems. *ACS nano* **2012**, *7*, 27–34.

(33) Montagne, K.; Gines, G.; Fujii, T.; Rondelez, Y. Boosting functionality of synthetic DNA circuits with tailored deactivation. *Nature communications* **2016**, *7*, 13474.

(34) Urtel, G.; Van Der Hofstadt, M.; Galas, J.-C.; Estevez-Torres, A. rEXPAR: an isothermal amplification scheme that is robust to autocatalytic parasites. *Biochemistry* **2019**, *58*, 2675–2681.

(35) Van Der Hofstadt, M.; Galas, J.-C.; Estevez-Torres, A. Spatiotemporal Patterning of Living Cells with Extracellular DNA Programs. *ACS nano* **2020**,

(36) Zadorin, A. S.; Rondelez, Y.; Galas, J.-C.; Estevez-Torres, A. Synthesis of programmable reaction-diffusion fronts using DNA catalyzers. *Physical review letters* **2015**, *114*, 068301.

(37) Zadorin, A. S.; Rondelez, Y.; Gines, G.; Dilhas, V.; Urtel, G.; Zambrano, A.; Galas, J.-C.; Estevez-Torres, A. Synthesis and materialization of a reaction-diffusion French flag pattern. *Nature chemistry* **2017**, *9*, 990.

(38) Martínez-Prat, B.; Ignés-Mullol, J.; Casademunt, J.; Sagués, F. Selection mechanism at the onset of active turbulence. *Nature physics* **2019**, *15*, 362.

(39) Vedel, S.; Bruus, H. Transient Taylor–Aris dispersion for time-dependent flows in straight channels. *Journal of Fluid Mechanics* **2011**, *691*, 95–122.

(40) Gines, G.; Zadorin, A.; Galas, J.-C.; Fujii, T.; Estevez-Torres, A.; Rondelez, Y. Microscopic agents programmed by DNA circuits. *Nature nanotechnology* **2017**, *12*, 351.

(41) Jones, M. R.; Seeman, N. C.; Mirkin, C. A. Programmable materials and the nature of the DNA bond. *Science* **2015**, *347*, 1260901.

(42) Zhao, Y.; Chen, F.; Li, Q.; Wang, L.; Fan, C. Isothermal Amplification of Nucleic Acids. *Chemical Reviews* **2015**, *115*, 12491–12545.

(43) Ross, T. D.; Lee, H. J.; Qu, Z.; Banks, R. A.; Phillips, R.; Thomson, M. Controlling organization and forces in active matter through optically defined boundaries. *Nature* **2019**, *572*, 224–229.