2   Co-variation between glucocorticoids, behaviour and 3   immunity supports the pace-of-life syndrome hypothesis: 4   an experimental approach

Jeffrey Carbillet[1,2], Benjamin Rey[3], Rupert Palme[4], Chloé Monestier[5], Luca Börger[6],

Typhaine Lavabre[7,8], Marie-Line Maublanc[1], Nicolas Cebe[1], Jean-Luc Rames[1],

Guillaume Le Loc'h[9], Marine Wasniewski[10], Benoit Rannou[2], Emmanuelle Gilot-

Fromont[2,3], Hélène Verheyden[1]

1 CEFS, Université de Toulouse, INRAE, Castanet Tolosan, France

2 Université de Lyon, VetAgro Sup, Marcy-l'Etoile, France

3 Université de Lyon, Université Lyon 1, UMR CNRS 5558, Villeurbanne Cedex, France

4 Unit of Physiology, Pathophysiology, and Experimental Endocrinology, Department of 13   Biomedical Sciences, University of Veterinary Medicine, Vienna, 1210, Austria

5 Department of Evolution, Ecology and Behavior, University of Illinois, Urbana, IL, United 15   States

6 Department of Biosciences, Swansea University, Singleton Park, Swansea, United Kingdom

7 Equipe de Biologie médicale-Histologie, CREFRE, Inserm-UPS-ENVT, Toulouse, France

8 Inovie Vet, Laboratoire d'Analyses et Biologie Vétérinaires, Montpellier, France

9 Université de Toulouse, ENVT, INRAE, UMR IHAP, Toulouse, France.

10 French Agency for Food, Environmental and Occupational Health & Safety (ANSES), Nancy Laboratory for Rabies and Wildlife, European Union Reference Laboratory for Rabies, European Union Reference Laboratory for Rabies Serology, WHO Collaborating Centre for Research and Management in Zoonoses Control, OIE Reference Laboratory for Rabies, Technopôle agricole et vétérinaire de Pixérécourt, Malzéville, France

* Corresponding author: jeffrey.cm@live.fr

Summary statement

Experimental approach on roe deer highlighted a gradient of individual responses to stress and immune challenges, as revealed by co-variation between behavioural profiles and changes over time in glucocorticoids and immunity.

Abstract

The biomedical literature has consistently highlighted that long-term elevation of glucocorticoids might impair immune functions, while short-term elevation might enhance them. In contrast, in wild and free-ranging animals, patterns are less clear. In the present study, we explored the stress-immunity relationship in controlled conditions, taking into account the potential effects of behavioural profiles. Thirteen captive roe deer (*Capreolus capreolus*) were monitored over an eight-week period encompassing two capture events. We assessed how changes in baseline faecal cortisol metabolite (FCM) concentrations following a standardised capture protocol and vaccination affected changes in thirteen immune parameters of the innate and adaptive immunity, and whether behavioural profiles were linked to changes in baseline FCM levels and immune parameters. We found that individuals showing an increase in baseline FCM levels over time also exhibited an increase in immunity, for both the innate and adaptive immunity and were characterised by more reactive behavioural profiles (as indicated by low activity levels, docility to manipulation and neophilia). Consequently, our results suggest that immunity of large mammals may be influenced by their stress level, but also by their behavioural profiles, as it is predicted by the pace-of-life syndrome hypothesis. Our analysis therefore highlights the need to consider co-variations between behavioural profiles, immunity and stress hormones in order to gain a better understanding of the unexplained among-

62 individual variability in the glucocorticoids-immunity relationships observed in wildlife, as

63 they may be underpinned by different life-history strategies.

Introduction

The immune system is one of the most important mechanisms in vertebrates to improve their

survival. This complex system is composed of two complementary arms, the innate (relatively

fast and non-specific) and adaptive (slower at first encounter, but more long-lasting and

specific) immunity, each composed of numerous cells and effectors (Demas, 2004). This

system, however, is not cost-free (Lochmiller and Deerenberg, 2000; Demas 2004), and there

should be trade-offs between immune defences and other functions that use a common resource

and contributes to fitness (Sheldon and Verhulst, 1996; Rauw, 2012). Glucocorticoids (such as

cortisol and corticosterone) play a major role in regulating the use of energy (Landys et al.,

2006; Busch and Hayward, 2009) and may therefore underlie such trade-offs.

In addition to their role as metabolic hormones, glucocorticoids are also one of the main

mediators of the stress response. In response to external or internal stimuli, the activation of

behavioural and physiological responses allows an organism to cope with challenges

(Wingfield, 1998; Romero and Wingfield, 2015). In particular, the activation of the

hypothalamic-pituitary-adrenal (HPA) axis that results in the secretion of glucocorticoid

hormones helps organisms to cope with stressful situations by making stored energy available

(Sapolsky et al., 2000). However, when glucocorticoid hormones are secreted repeatedly or for

long periods of time, negative effects on other energetically demanding functions such as

reproduction (Boonstra, 2005) and immunity (Martin, 2009) may appear.

Over the past years, several studies have investigated the relationship between stress

and immunity, particularly in the biomedical domain where it has been shown that short-term

elevation of glucocorticoids (i.e., few minutes to few hours) enhances immune functions (Bilbo

et al., 2002; Viswanathan et al., 2005; Dhabhar, 2014) while chronically elevated glucocorticoid

levels are immunosuppressive (Dhabhar and McEwen, 1997; Sapolsky et al., 2000; Dhabhar,

2009). Focusing on long-term elevation of glucocorticoids (i.e., few days to few months),

studies on wild animals have shown mixed results, suggesting that chronically elevated

glucocorticoid levels may be associated with decrease, increase, or no changes in immune

functions (Matson et al., 2006; Delehanty and Boonstra, 2009; Merrill et al., 2012; Brooks and

Mateo, 2013). Evidence is also accumulating that glucocorticoid levels do not affect all aspects

of the immune system in the same manner, such that immunoglobulin production may be

impaired while other parameters (T-cell mediated, or constitutive immunity) might not be

affected (Bourgeon and Raclot, 2006; Stier et al., 2009). These contrasting patterns indicate that

our understanding of the nature of the stress-immunity relationship in the wild remains limited

and fragmentary.

Among the factors likely to modulate this relationship and to help to shade light on the

discrepancies, the consideration of behavioural profiles has been largely neglected. From a

theoretical point of view, close links between physiology and behaviour are expected due to the

underlying energetic basis of both traits (Careau et al., 2008). Accordingly, consistent among-

individual differences in behaviour, also called personality (Réale et al., 2007), are related to

numerous physiological traits including glucocorticoid secretion and immune function

(Koolhaas et al., 1999; Koolhaas et al., 2008; Réale et al., 2010). For instance, a study on wild

superb fairy-wrens (*Malurus cyaneus*) showed that individuals exhibiting proactive behavioural

traits (fast exploration of a novel environment) exhibited lower level of natural antibodies

(Jacques-Hamilton et al., 2017). In addition, previous works highlighted that slower explorer

or more reactive individuals tend to exhibit higher baseline and stress-induced glucocorticoid

levels compared to faster or more proactive ones (Koolhaas et al., 1999; Stöwe et al., 2010;

Montiglio et al., 2014). In addition, a recent study on laying hens (*Gallus gallus domesticus*)

highlighted that more reactive individuals exhibited greater stress and immunological (greater

swelling in response to phytohemagglutinin injection) responsiveness compared to more

proactive ones (Pusch and Navarra, 2018). Such co-variations between behavioural and

physiological traits can be interpreted within the pace-of-life syndrome hypothesis described

by Denis Réale and his collaborators (2010). This hypothesis posits that species, populations or

individuals experiencing different ecological conditions should differ in a suite of behavioural,

physiological and life-history traits that may have co-evolved according to the particular

ecological conditions encountered, leading to differences in life-history strategies. Within this

hypothesis, it is expected that individuals with slower life history strategies may have more

reactive behavioural profiles, higher glucocorticoid levels and higher investment in overall

immunity, while those with faster life history strategies may have more proactive behavioural

profiles, lower glucocorticoid levels and lower overall investment in immunity.

In the present study, we investigated the relationships between changes in baseline

glucocorticoid levels and changes in thirteen immune parameters, and how these were related

to behavioural profiles. To do so, we investigated the link between variations in baseline faecal

cortisol metabolite (FCM) levels and variations in adaptive and innate immune functions,

before and after a standardised capture stress protocol associated with an immune challenge

using anti-rabies vaccination, in a captive roe deer (*Capreolus capreolus*). In addition, we

evaluated how these changes may be related to behavioural profiles, as characterised by three

commonly used behavioural traits: docility, neophobia, and activity levels.

Based on the literature, we expected that 1) changes in baseline FCM levels between the

two observation periods (before/after capture) would co-vary with changes in immune

parameters and inflammatory markers; 2) changes in baseline FCM levels should be less related

to adaptive than innate immune parameters and inflammatory markers, due to the relatively low

cost of adaptive immunity (McDade et al., 2016). In addition, we expected 3) that baseline FCM levels as well as variations in baseline FCM levels should be related to individual behavioural profiles (Koolhaas et al., 1999; Montiglio et al., 2012), with higher baseline levels and higher increase in baseline levels in more docile, less active and more neophilic individuals (i.e., more reactive individuals). Finally, we predicted 4) the increase in immune parameters between the two observation periods should be more important for more reactive individuals, that are supposed to invest more in overall immunity (Réale et al. 2010; Pusch and Navarra, 2018).

Material and methods

Study site

This study was conducted on a captive population of roe deer living in the Gardouch research station, located in south-west of France, approximately 30 km south-east of Toulouse. The station is located on the slopes of a hill, approximately 230 m above sea level. The climate is oceanic with summer droughts. The station is owned and managed by the French Research Institute for Agriculture, Food and Environment (INRAE). It consists of 12 enclosures of 0.5 ha, each containing between one to six captive roe deer. Each enclosure is composed of meadow, trees and groves that provide shelter. Each enclosure contains a hut where individuals are supplemented with food pellets (600 g per individual per day) and can obtain a shelter. The current experiment included a total of 13 females, living in these enclosures since several years already, aged from 4 to 13 years old. All roe deer were raised at the station from their birth or their first year of life. In addition, they all had some degree of habituation to humans but were able to express normal behavioural responses (e.g., vigilance, escape) to stressful situations.

Experimental design

The experimental procedure was carried out between mid-September and mid-November 2018

and is summarised in Fig.1. To assess the baseline glucocorticoid levels of each individual, we

collected faeces every four days during four weeks to further determine faecal cortisol

metabolite concentrations (i.e., period 1). Faeces were collected immediately after defecation

was observed by experienced personnel and kept at +4 °C for a maximum of 1 h before being

stored at −20 °C until steroid analysis. At the end of period 1, each roe deer was subjected to a

standardised capture stress protocol involving restrained immobilisation (Monestier et al.,

2016). This procedure made it possible to collect faeces from the rectum, to draw blood

samples, to inject an anti-rabies vaccine, and to fit collars equipped with tri-axial accelerometers

(see Capture protocol and data collection for details). Collection of faecal samples was then

continued every four days for an additional four weeks, as describe above (i.e., period 2), in

order to evaluate the effect of capture on baseline glucocorticoid level for each individual.

Period 2 started two days after capture 1, in order to avoid measuring the acute increase in

glucocorticoid level due to capture (Dehnhard et al., 2001). At the end of period 2, roe deer

were recaptured following the same procedure (capture 2) and faeces and blood were again

collected.

Capture protocol and data collection

Roe deer were directed into their hut (one enclosure and one individual at a time) by slowly

approaching them. Each individual was then pushed through a trap door into a retention box.

Once in the box, animals were tranquilised with an intramuscular injection of acepromazine

(Calmivet®, Vetoquinol, France; targeted dose of 0.075 mg/kg), as recommended by Jordi

Montané and his colleagues (2003). Individuals were weighed with an electronic balance to the

nearest 100 g.

In addition, for each individual, we characterised three behavioural traits at three different moments. First, we recorded whether animals struggled during handling, and this measure was used as an index of docility, as previously used and described for roe deer (Gervais et al., 2020). A score of 0 was assigned to individuals that did not struggle (docile), while a score of 1 was attributed to those that did struggle (non-docile) during handling. This behavioural trait during handling has been shown to be repeatable over time (r=0.26) with a tendency to be heritable (h=0.17) in roe deer (Gervais et al., 2020). The second behavioural trait consisted in determining spontaneous daily activity (Réale et al., 2007) during four weeks with tri-axial accelerometery technology. To this end, we fitted collars equipped with tri axial accelerometers (Daily Diary tags, Wildbytes Ltd., Swansea University) recording at 20 Hz. Daily activity was measured using the Vectorial Dynamic Body Acceleration (VeDBA) metric (Wilson et al. 2020), a measure of dynamic body acceleration (and hence activity) derived from the 20 Hz tri-axial accelerometer channels – here, using a 2 second smoothing window and standard procedures (Wilson et al. 2020), as implemented in the DDMT software (Wildbytes Ltd., Swansea University). VeDBA values were summed for each individual, date, and hour of the day ('total VeDBA'), from which an average was taken for each individual as the mean daily activity through the four weeks between the two capture events (period 2). The third behavioural trait we considered was neophobia, defined as the avoidance of novel stimuli in the environment, and a widely studied dimension of animal personality (Réale et al., 2007; Greenberg and Mettke-Hofmann 2001). The neophilia-neophobia continuum was assed using the difference in feeding efficiency with and without the presence of a novel object, a measure previously used and validated for this roe deer population (Monestier et al., 2017). The complete protocol for evaluating feeding efficiency has been described in detail by Chloé Monestier and her colleagues (2017) for this roe deer population. Briefly, we calculated the ratio between the number of visits to the hut that resulted in a successful feeding bout

(numerator) and the total number of visits to the hut (denominator), with and without the

presence of a novel object in the hut, near the manger. Measurements were repeated for five

days for each condition (with and without novel object), and the difference in the ratio between

the two conditions was then evaluated. More neophobic individuals should be less inclined to

feed during a given visit when there was a novel object present, resulting in a higher score on

the neophilia-neophobia continuum.

Finally, we obtained a functional index of the adaptive immune system activity by

injecting a standardized dose of an anti-rabies vaccine (Rabisin®, Merial, France, 1ml)

administered subcutaneously during the first capture and by quantifying the circulating specific

antibodies at capture 1 and 2 (see below).

Immune parameters measurement

Blood samples were taken from each individual from the left jugular vein (up to 20 mL for a

20 kg roe deer) and divided in two. Part of blood was collected on EDTA, preserved at 4 °C

and served to measure total leukocyte concentration (white blood cell [WBC]). WBC were

determined at the Ecole Nationale Vétérinaire de Toulouse (ENVT) with an automat (Sysmex

2000iV, Sysmex). To obtain concentrations of each leukocyte type (neutrophil, basophil,

eosinophil, lymphocyte and monocyte), the total leukocyte count was multiplied by the

proportion of each cell type. The differential cell count was performed on the first 100 WBCs

on Wright-Giemsa-stained blood smears by trained technicians (Houwen, 2001). The remaining

blood was collected on a dry tube, centrifuged at 1500 g for 15 min, and the serum was stored

at -20°C. Serum was used to measure total proteins using a refractometer and to perform

electrophoresis on agarose gel that allowed differentiating albumin and alpha-1, alpha-2, beta,

and gamma globulins. Haptoglobin that belongs to the alpha-2-globulin fraction was also

measured by spectrophotometry (Konelab 30i PLC, Fisher Thermo Scientific) with a non-

species-specific method based on the ability of haptoglobin to bind to haemoglobin. Circulating

levels of natural antibodies (NAbs) was measured by a hemagglutination test (HA) that

measures NAbs ability to agglutinate exogenous cells, while the complement activity was

revealed by the ability of proteins to induce hemolysis (HL) (Matson et al., 2005). The HA/HL

protocol defined by Matson et al., (2005) in birds has been adapted and previously used in roe

deer (Gilot-Fromont et al., 2012). Finally, we quantified the level of anti-rabies antibody

following the method described by Florence Cliquet and her colleagues (1998). We therefore

measured 10 markers of innate immunity (neutrophils, basophils, monocytes, eosinophils,

hemagglutination and hemolysis titers) including 4 inflammatory markers (haptoglobin, alpha-

1, alpha-2 and beta-2 globulins), and 3 markers of adaptive immunity (lymphocytes, gamma-

globulins and anti-rabies antibodies).

Extraction and quantification of FCMs

FCMs were extracted following a methanol-based procedure and assayed using a group-specific

11-oxoaetiocholanolone enzyme immunoassay (EIA), as previously described (Möstl et al.,

2002) and validated for roe deer (Zbyryt et al., 2017). Briefly, each faecal sample was

homogenised and 0.5 ± 0.005 g of the homogenate was transferred to a glass tube containing

5 ml of methanol 80%. The suspended samples were vortexed at 1500 rpm for 30 min and

centrifuged at 2500 g for 15 min (Palme et al., 2013). An aliquot of the supernatant was further

diluted (1:10) with assay buffer prior to EIA analysis. Measurements were carried out in

duplicate (intra- and inter-assay coefficients of all samples were less than 10% and 15%,

respectively).

Ethical approval

All applicable institutional and/or national guidelines for the care and use of animals were

followed. The overall protocol, including captures and blood sampling of roe deer was approved

by the Ethical Committee 115 of Toulouse and was authorized by the French government

(APAFIS#14706).

Statistical analyses

Relationship between behaviour and changes in immunity and baseline FCMs

To analyse relationships between change in baseline FCM levels, change in immune parameters

and behavioural profiles, we first calculated the difference between the measures obtained

during the second capture event and the first one for immune parameters (Δ immunity).

Similarly, in order to obtain a measure of change in baseline FCM levels (Δ glucocorticoids),

we calculated the difference of averaged baseline FCM levels between period 2 and period 1

for each individual. In addition, behavioural scores from the first capture were used to index

docility since values did not differ between the two captures within-individual, except for one

individual for which the score passed from 1 to 0. We chose to use values from the first capture

rather the second one in order to avoid a potential habituation effect for the only individual that

exhibited a change in its docility score.

Then, to test our hypothesis, we used Partial Least Square Path Modelling (PLS-PM)

analysis. PLS-PM is a statistical method for studying complex multivariate datasets, in

particular when complex individual traits (here, immunity arms and behavioural profiles) have

been approached by several observed variables, that are generally correlated. In this case,

groups (blocks) of variables describing a character can be summarised by a latent variable

defined as a linear combination of observed variables, without imposing any restriction on

within-block covariance. Then relationships between blocks (summarised by latent variables)

are analysed taking into account previous knowledge on causes and consequences in the studied

system. This statistical analysis is particularly recommended when dealing with variables

showing high correlation in order to avoid redundancies and high type I error (Sanchez, 2013).

Here we built the following blocks, each being summarised by a latent variable: Δ

glucocorticoids (1 variable), Δ innate immunity (6 variables), Δ adaptive immunity (3

variables), Δ inflammatory markers (4 variables), and behavioural profile (3 variables, Table

1).

We then ran 3 PLS-PM analyses, each one evaluating the relationships between

behavioural profile, change in baseline glucocorticoids, and change in 1) innate immunity, 2)

adaptive immunity, and 3) inflammatory markers. For each of the three analyses, we built a

structural model (or inner model, i.e. model of relationships among latent variables) that

consisted of three latent variables: Δ glucocorticoids, behavioural profile, and Δ of one of the

three immune arms (innate, adaptive and inflammatory). The statement for the structural models

was as follows: change in immunity depends on both behavioural profile and change in baseline

glucocorticoids, which also depends on behavioural profile. Finally, in the measurement model

(or outer model, i.e. relationships between observed and latent variables), the observed variables

were considered as reflecting the corresponding latent variable (reflective mode), except for

innate immunity were observed variables were considered as constituting the latent variable

(formative mode). This option was chosen due to the high number of biomarkers used and the

complexity and diversity of the biological actions of the innate immune system (Demas et al.,

2011). This diversity is reflected in the moderate correlation among components of this latent

variable.

We then ran PLS-PM analysis to adjust both the structural (Fig. 2) and measurement

models, through multiple linear regressions. Lastly, we performed the diagnosis of each model

following the recommendations of Gaston Sanchez (2013). For blocks of reflective variables,

we checked for unidimentionality using Dillon-Goldstein's Rho and eigenvalues of the

correlation matrix, and also checked for cross-loadings, to determine the differentiation among

blocks. For all three models, we checked the measurement model using weights and loadings

of observed variables in the latent variable definition, as well as communality (squared

loadings, part of variance in common between observed and latent variable) and redundancy

(variance of indicators predicted from the corresponding latent variable) (presented in Table

S1). The structural models were checked using R^2, redundancy index (ability to predict) and

Goodness-of-fit index, a pseudo GoF measure that reflects the overall prediction power of the

model $(0 < Gof < 1)$.

Relationship between baseline FCMs throughout the experiment and behaviour

In order to test hypothesis 3 and evaluate the link between among-individual differences in

baseline glucocorticoid levels throughout the experiment (period 1 + period 2) and behavioural

profiles, while taking into account other factors affecting FCM levels, we analysed baseline

FCM levels by performing linear mixed-effects models (LMMs) on the 181 observations of 13

individuals (14 repetitions per individuals with 1 missing value for one individual). FCM values

were log transformed to achieve normality of model residuals. As we measured three different

behavioural traits, we analysed the overall correlation pattern between docility, neophobia and

activity using a normed Principal Component Analysis (PCA). Consequently, we used scores

from the first principal component (PC1) which indexed a gradient of behaviour ranging from

individuals showing more proactive behavioural profile (non-docile, active and neophobic) to

individuals showing a more reactive behavioural profile (docile, less active and neophilic)

(Table S2, Fig. S1). We then built a reference model that included all biologically relevant

variables to explain baseline glucocorticoids levels, and compared this model with all its sub-

models. The reference model included: PC1, age of individuals, and Julian date of sample

collection. Individual identity and enclosure identity were included as random effects to avoid

pseudo-replication issues (Hurlbert, 1984) and to control for unexplained variance due to

among-individual differences and among-enclosure variation.

To select the best models of variation in FCM levels we used a model selection

procedure based on the second-order Akaike Information Criterion (AICc, Burnham and

Anderson, 2003). Models with a difference in AICc (ΔAICc) > 2 units from the best model

were considered to have less support, following Burnham and Anderson (2003). In addition,

we removed models within two AICc units of the top model that differed from a higher-ranking

model by the addition of one or more parameters. These were rejected as uninformative, as

recommended by Todd Arnold (2010) and Shane Richards (2008). In addition, we calculated

AICc weights (AICcw) to measure the relative likelihood that a given model was the best

among the set of fitted models. Normality of the residuals for the selected models was tested

(Shapiro-Wilk test) and visually assessed. Goodness-of-fit was assessed by calculating

conditional (i.e. total variance explained by the best supported model) and marginal (i.e.

variance explained by fixed effects alone) R^2 values and standard residual plot techniques

(Nakagawa and Schielzeth, 2013).

All analyses were carried out with R version 3.6.0 (R Development Core Team 2016),

using the lmer function from the lme4 package (Bates et al., 2014) and the plspm function from

the plspm package (Sanchez, 2013).

Results

Co-variation between behaviour and changes in immunity and baseline FCMs

Among the 13 individuals considered in our study, 9 showed a decrease in baseline FCM levels

during period 2 compared to period 1 (ranging from -786 to -8 ng/g of wet faeces), while 4

showed an increase (ranging from 93 to 346 ng/g). In addition, for each individual, vaccination

increased the level of anti-rabies antibody, but large among-individual differences were

observed, with values ranging from 0.60 to 41.50 IU, with a median of 10.39. Regarding

behavioural profiles, we found, as predicted, that proactive individuals were characterised by

high daily activity levels, lack of docility and neophobia (Table S2). In addition, the 13

individuals appeared to be homogeneously distributed along the gradient ranging from

proactive to reactive behavioural profiles as showed by the PC1 axis scores ranging from -1.92

to 2.66, with a median value of 0.20 (Fig. S1).

Our analyses showed links between behavioural profiles and changes in the three

studied aspects of immunity (Table 2). Precisely, individuals that exhibited more proactive

behaviours, expressed by high daily activity levels, lack of docility and neophobia, showed an

overall strong decrease in innate ($r = -0.53$; $P < 0.05$; Fig. 2), adaptive ($r = -0.80$; $P < 0.001$;

Fig. 2), and inflammatory ($r = -0.77$; $P < 0.005$; Fig. 2) markers of immunity. However,

differences appeared among models as for the weights of observed variables in the definition

of latent variables. When analysing Δ adaptive immunity, gamma globulins, lymphocytes and

anti-rabies antibody contributed approximately equally to the latent variable (weights [w] of

0.37; 0.52 and 0.54 respectively), while behavioural profile was particularly represented by

docility and neophobia ($w = 0.65$ and 0.64 respectively) and very little by activity ($w = -0.04$).

On the opposite, for Δ inflammatory markers, behavioural profile was largely represented by

mean daily activity levels ($w = 1.05$) and weakly by docility ($w = -0.32$) and especially

neophobia ($w = 0.12$), while markers of inflammation contributed overall to the same

proportion to their latent variable (Table 3). Lastly, for Δ innate immunity, behavioural traits

contributed to the same extent to their latent variable while basophils and hemagglutination

contributed more strongly to their one (Table 3). In addition, it is important to note that among

the innate immune parameters, neutrophils were negatively correlated to other biomarkers of

innate immunity (see the negative loading in Table 3), meaning that the negative relationship

observed between behavioural profiles and innate immunity only occurred for other biomarkers

of innate immunity, while temporal changes in neutrophil concentrations were actually

positively linked to activity, neophobia and lack of docility.

Changes in baseline glucocorticoid levels also appeared to be associated with changes

in innate (r = 0.61; P < 0.05) and adaptive immunity (tendency: r = 0.37; P < 0.06), but not

inflammatory markers (Table 2; Fig. 2). Precisely, individuals that underwent an increase in

baseline FCM levels between periods also exhibited an increase in both innate and adaptive

immunity. However, as pointed out above, this positive relationship means that individuals

exhibiting an increase in baseline FCM levels actually had a decrease in neutrophil

concentration.

Finally, the relationship between behavioural profiles and changes in baseline FCM

levels was non-significant for all three models (Table 2).

Co-variation between baseline FCM levels throughout the experiment and

behaviour

According to the model selection procedure, the best model describing among-individual

differences in baseline FCMs throughout the experiment in relation to individual behavioural

profiles included PC1 score and period of the experimental protocol (Table S3). Specifically,

roe deer that exhibited a more reactive behavioural profile (as expressed by low daily activity

levels, docility and neophilia) also exhibited higher baseline FCM levels throughout the

experiment compared to roe deer exhibiting a more proactive behavioural profile (0.103; P <

0.005; Table 4; Fig. 3). In addition, baseline FCM levels decreased during the second part of

the experimental protocol compared to the first one (-0.1663; P < 0.05; Table 4), meaning that

the mean baseline FCM level of the studied roe deer population was lower during period 2 than

during period 1. Age, and Julian date did not increase the fit of the model.

Discussion

In this study, we first aimed to gain a better understanding on how changes in baseline glucocorticoid levels may affect simultaneous changes in immune parameters of the innate, adaptive and inflammatory markers of immunity, at the scale of few weeks (8 weeks). Our results demonstrated a correlation between baseline glucocorticoids and immunity, such that an increase in baseline FCM levels was related to an increase in immune parameters of the innate and adaptive arms, but not inflammation. Second, we tested whether behavioural profiles could underlie the co-variation between changes in immune parameters and baseline glucocorticoid levels. As predicted, behavioural profiles appeared to be strongly linked to changes in overall immunity, but also to baseline glucocorticoid levels throughout the experiment, while not related to changes in baseline glucocorticoids between the two periods of the study.

First, an increase in innate immunity was generally linked to an increase in baseline glucocorticoid levels between the two periods, except for neutrophil concentrations, which decreased as glucocorticoid levels increased. The observed negative relationship with neutrophils is consistent with the previous finding of an immunosuppressive effect of long-term elevation of glucocorticoids on immunity (Dhabhar, 2014; Martin 2009), while the overall increase in innate immunity may appear as more surprising, and opposite to our predictions. One of the assumptions underlying the hypothesis of a negative relationship between immunity and glucocorticoids is the energy cost of immunity, which should subsequently trade-off against other energy demanding functions (Lochmiller and Deerenberg 2000). However, it is important to note that the studied population, while living in large enclosures, is supplemented with food resources. Consequently, resources are not limiting and no unpredictable variations in food resources availability occurs. This particular context may lead to a disappearance of some trade-offs between energy demanding functions and may explain the absence of a negative

association between change in baseline glucocorticoids and immunity. Alternatively, to the

dominant energy cost hypothesis, it has also been proposed that, as the main function of the

stress response is to recover from stressors, a decrease in immunity should not necessarily occur

when glucocorticoids increase, as it could improve survival (Martin, 2009).

While the above hypothesis may partly explain the link between change in innate

immunity and change in baseline glucocorticoids, it does not explain the difference observed

between neutrophils and other biomarkers of innate immunity. Neutrophils are part of the

cellular immunity and reflect acute inflammatory response while monocytes reflect chronic

inflammatory response, and hemagglutination and hemolysis are both part of the humoral innate

response (Demas et al. 2011). In addition, eosinophils are known to be a specific component

that bridges innate and adaptive immunity (Demas et al. 2011). Considering these differences

in the functions of the different biomarkers we used, it is likely that they may not all be linked

in the same manner with glucocorticoids, which could explain the difference we observed.

Regarding basophils, which represent the major component of the latent variable "innate

immunity" alongside hemagglutination, they have been shown to be particularly secreted in

presence of ticks in order to develop an acquired resistance to this ectoparasite (Wada et al.,

2010; Karasuyama et al., 2018). Roe deer living in this experimental facility are particularly

exposed to ticks each year (unpublished data), and investment in production and maintenance

of basophils may therefore be important in order to decrease the costs associated with ticks'

parasitism and improve their fitness, which could be another explanation to the absence of a

trade-off and positive relationship between changes in glucocorticoid levels and innate

immunity.

With respect to a change in the adaptive arm of the immune system, our results did not

support the finding of the immunosuppressive effect of long-term elevation of glucocorticoids,

but instead showed that adaptive immune parameters increased in individuals that showed an

increase in baseline FCM levels. A first possible explanation of this result could be linked to

the transient increase in glucocorticoids that occurred during the first capture, where

vaccination was done. Previous work showed that a short-term elevation in glucocorticoids at

the time of vaccination reinforces the efficacy of vaccination (Dhabhar, 2014). A similar

scenario may have occurred in our study, and the stress of capture, resulting in a sudden increase

in circulating glucocorticoids, would have strengthened the immune response to anti-rabies

vaccine. This may partly explain our result if there is a correlation between the change observed

in baseline FCM levels and the stress-induced increase in glucocorticoids. Another possible

explanation is that the energetic cost of mounting an antibody response is relatively low

(McDade et al., 2016), and a trade-off between antibody levels and other functions was

therefore not necessarily expected. The positive relationship we observed in our study also

supports the pace-of-life syndrome hypothesis, where individuals exhibiting higher baseline

glucocorticoid levels should show stronger investment in immunity than those exhibiting lower

baseline glucocorticoid levels (Réale et al. 2010).

To summarise the immunity-glucocorticoids relationship found during our experiment,

individuals that experienced an increase in baseline glucocorticoid levels showed an investment

towards a strong immune system, while individuals that experienced a decrease in baseline

glucocorticoid levels showed a down-regulation of their immunity. Consequently, it appears

that in our study, individuals that underwent a decrease in baseline glucocorticoids may have

switched their investment from the immune system, possibly toward another unstudied

function, according to their changes in baseline stress levels. We suggest that this possible

switch could be underlain by a plastic response to the stressful event of capture, leading to an

adjustment of the individuals' life history strategies and change in investment between

functions supporting either long-term survival or current reproduction. This would be in

accordance with the pace-of-life syndrome hypothesis, where a positive association is expected

between glucocorticoid levels and immunity, with higher levels in individuals favouring their

long-term survival, while lower levels are expected in individuals favouring reproduction and

growth (Réale et al. 2010).

Regarding the association between behavioural profiles and change in immunity, we

found an overall similar association for the three components of the immune system assessed

in our study, namely, innate, adaptive and inflammatory immunity. Precisely, individuals

showing a propensity to be active, non-docile to manipulation and repulsed by novelty showed

a decreased investment in their immune system following the first capture event. This result is

in accordance with our predictions and supports the hypothesis that individuals with a fast pace-

of-life should exhibit proactive behavioural profile and should invest less in immune functions

than individuals with a slow pace-of-life, as they should prioritise investment in current

reproduction du to their relatively short longevity, and functions supporting aggressive

behaviours, high activity or elevated heart rate for instance, which characterised these

behavioural profiles (Réale et al. 2010). In this manner, the low investment in immunity showed

by fast-living individuals would support energy savings in order to invest in the functions cited

above rather than into mechanisms enhancing survival, such as immunity. On the opposite,

individuals showing more reactive behavioural profiles are supposed to have a slower pace-of-

life and are expected to favour functions enhancing survival and future reproduction (Réale et

al., 2010), hence a stronger investment in immunity as they are expected to frequently encounter

pathogens throughout their life, and most likely to face repeated encounter with the same

pathogens.

Finally, we investigated whether behavioural profiles could be associated with among-

individual variations in baseline glucocorticoid levels throughout the experiment, and also with

changes in baseline glucocorticoid levels over an 8-week period. Our results did not provide

any support for the latter, but instead supported the former. Specifically, individuals exhibiting

proactive behavioural profiles also showed lower baseline glucocorticoid levels throughout the

experiment compared to individuals exhibiting more reactive behavioural profiles. This result

supports previous studies on wild (Carbillet et al., 2019) and captive (Monestier et al., 2016)

roe deer. It is also in accordance with the coping style framework (Koolhaas et al., 1999) and

the pace-of-life syndrome hypothesis (Réale et al., 2010) which states that behavioural and

physiological responses to stressful situations are correlated. During our experiment, we

therefore observed a different relationship between behavioural profiles and baseline

glucocorticoids throughout the experiment, and between behavioural profiles and the change of

hormones between two specific periods. The difference we observed in our results may suggest

that activity (reflected by baseline levels) and reactivity (reflected by the changes following the

first capture event) of the HPA axis may not be associated in the same manner with among-

individual variations in behaviour (Koolhaas et al., 1999). Precisely, only activity of the HPA

axis may be associated with behavioural profiles, or at least more strongly associated with

behavioural profiles than reactivity of the HPA axis.

Overall, our results indicate that individual roe deer reacted differently to our protocol,

with part of the individuals showing an increase in baseline glucocorticoid levels, while others

showed a decrease. In addition, our results suggest that the increase in baseline glucocorticoid

levels may be associated with a re-allocation of energetic resources towards innate and adaptive

immunity, but also that this re-allocation towards immunity might be associated with a gradient

of behavioural profiles, such that individuals showing more reactive behaviours showed

increased investment in immunity and higher baseline glucocorticoid levels (see Fig. 4 for a

summary of the outcomes of the protocol). Immune functions are known to be costly in terms

of energy (Lochmiller and Deerenberg, 2000; McDade et al., 2016), and glucocorticoids are

metabolic hormones that regulate the use of energy (Palme, 2019). We therefore suggest that

the observed association between immunity and baseline glucocorticoid levels may be

associated with different life-history strategies and underpinned by energetic trade-offs between

functions enhancing survival, reproduction and growth, which would be congruent with the

pace-of-life syndrome hypothesis (Réale et al., 2010).

Finally, considering that all of our results tend to support a co-variation between stress

hormones, immunity and behaviour, it might seem important that future work goes one step

further and investigates how among-individual variations in behaviour could modulate the

variation of glucocorticoid levels, as well as the relationship between glucocorticoid hormones

and immunity. The extent to which co-variations between these traits are influenced by different

life history strategies in the wild also warrants further investigations as it may help to

understand the large and unexplained among-individual variability in the glucocorticoids-

immunity relationship observed in wildlife studies.

Acknowledgments
531

We thank Eric Bideau and Benoit Coracin for their help throughout this study regarding roe

deer manipulation and monitoring. We also thank Edith Klobetz-Rassam for EIA analysis,

veterinary students for their assistance during capture events, as well as Laura Gervais for

constructive discussions and comments that helped to improve this manuscript. We are also

very grateful to the ECODIV department of the INRAE for its financial support to the Gardouch

experimental station.

Author contributions
538

539 JC, BRey, EGF and HV conceived and designed the study. JC, HV, MLB, NC, JLR, GLL

540 undertook the fieldwork. LB performed the VeDBA analysis. RP performed the FCMs analysis.

541 MW quantified anti-rabies circulating antibodies. TL, BRannou and EGF performed the

542 immunological analysis. CM designed and performed the neophobia analysis. JC performed the

statical analysis, wrote the first draft of the paper and then received input from all other co-authors. All authors approved the final version of the manuscript and agree to be held accountable for the content therein.

Competing interests

The authors declare that they have no conflict of interest.

Funding

The study was funded by INRAE, VetAgro Sup and OFB, and was performed in the framework of the LABEX ECOFECT (ANR-11-LABX-0048) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007).

References

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823.*

Bilbo, S. D., Dhabhar, F. S., Viswanathan, K., Saul, A., Yellon, S. M., Nelson, R. J. (2002). Short day lengths augment stress-induced leukocyte trafficking and stress-induced enhancement of skin immune function. *Proc. Natl. Acad. Sci. U.S.A* 99, 4067-4072.

Boonstra, R. (2005). Equipped for life: the adaptive role of the stress axis in male mammals. *J. Mammal.* 86, 236-247.

Bourgeon, S., and Raclot, T. (2006). Corticosterone selectively decreases humoral immunity in female eiders during incubation. *J. Exp. Biol.* 209, 4957-4965.

Brooks, K. C., and Mateo, J. M. (2013). Chronically raised glucocorticoids reduce innate

immune function in Belding's ground squirrels (*Urocitellus beldingi*) after an immune

challenge. *Gen. Comp. Endocrinol.* 193, 149-157.

Burnham, K. P., and Anderson, D. R. (2002). Model selection and multimodel inference: a

practical information-theoretic approach. *Springer Science & Business Media*.

Busch, D. S., and Hayward, L. S. (2009). Stress in a conservation context: a discussion of

glucocorticoid actions and how levels change with conservation-relevant variables. *Biol.*

Conserv. 142, 2844-2853.

Carbillet, J., Rey, B., Lavabre, T., Chaval, Y., Merlet, J., Débias, F., Régis, C., Pardonnet, S.,

Duhayer, J., Gaillard, J.M., et al. (2019). The Neutrophil to Lymphocyte ratio and individual

coping styles across variable environments in the wild. *Behav. Ecol. Sociobiol.* 73, 1-13.

Careau, V., Thomas, D., Humphries, M. M., Réale, D. (2008). Energy metabolism and animal

personality. *Oikos* 117, 641-653.

Cliquet, F., Aubert, M., Sagne, L. (1998). Development of a fluorescent antibody virus

neutralisation test (FAVN test) for the quantitation of rabies-neutralising antibody. *J. Immunol.*

Methods 212, 79-87.

Dehnhard, M., Clauss, M., Lechner-Doll, M., Meyer, H. H. D., Palme, R. (2001). Noninvasive

monitoring of adrenocortical activity in roe deer (*Capreolus capreolus*) by measurement of

fecal cortisol metabolites. *Gen. Comp. Endocrinol.* 123, 111-120.

Delehanty, B., and Boonstra, R. (2009). Impact of live trapping on stress profiles of

Richardson's ground squirrel (*Spermophilus richardsonii*). *Gen. Comp. Endocrinol.* 160, 176-

182.

Demas, G. E. (2004). The energetics of immunity: a neuroendocrine link between energy

balance and immune function. *Horm. Behav.* 45, 173-180.

Demas, G. E., Zysling, D. A., Beechler, B. R., Muehlenbein, M. P., French, S. S. (2011).

Beyond phytohaemagglutinin: assessing vertebrate immune function across ecological

contexts. *J. Anim. Ecol.* 80, 710-730.

Dhabhar, F. S., and McEwen, B. S. (1997). Acute stress enhances while chronic stress

suppresses cell-mediated immunity in vivo: A potential role for leukocyte trafficking. *Brain*

Behav. Immun. 11, 286-306.

Dhabhar, F. S. (2009). Enhancing versus suppressive effects of stress on immune function:

implications for immunoprotection and immunopathology. *Neuroimmunomodulation* 16, 300-

317.

Dhabhar, F. S. (2014). Effects of stress on immune function: the good, the bad, and the

beautiful. *Immunol. Res.* 58, 193-210.

Gervais, L., Hewison, A. J., Morellet, N., Bernard, M., Merlet, J., Cargnelutti, B., Chaval, Y.,

Pujol, B., Quéméré, E. (2020). Pedigree-free quantitative genetic approach provides evidence

for heritability of movement tactics in wild roe deer. *J. Evol. Biol.* 33, 595-607.

Gilot-Fromont, E., Jégo, M., Bonenfant, C., Gibert, P., Rannou, B., Klein, F., Gaillard, J. M.

(2012). Immune phenotype and body condition in roe deer: individuals with high body

condition have different, not stronger immunity. *PLoS One* 7, e45576.

Greenberg, R., and Mettke-Hofmann, C. (2001). Ecological aspects of neophobia and neophilia

in birds. In Current ornithology (pp. 119-178). *Springer, Boston, MA*.

Houwen, B. (2001). The differential cell count. *Lab. Hematol.* 7, 89-100.

Hurlbert, S. H. (1984). Pseudoreplication and the design of ecological field experiments. *Ecol.*
Monogr. 54, 187-211.

Jacques-Hamilton, R., Hall, M. L., Buttemer, W. A., Matson, K. D., da Silva, A. G., Mulder,
R. A., Peters, A. (2017). Personality and innate immune defenses in a wild bird: Evidence for
the pace-of-life hypothesis. *Horm. Behav.* 88, 31-40.

Karasuyama, H., Tabakawa, Y., Ohta, T., Wada, T., Yoshikawa, S. (2018). Crucial Role for
Basophils in Acquired Protective Immunity to Tick Infestation. *Front. Physiol.* 9, 1-8.

Koolhaas, J. M., Korte, S. M., De Boer, S. F., Van Der Vegt, B. J., Van Reenen, C. G., Hopster,
H., et al. (1999). Coping styles in animals: current status in behavior and stress-
physiology. *Neurosci. Biobehav. Rev.* 23, 925-935.

Koolhaas, J. M. (2008). Coping style and immunity in animals: making sense of individual
variation. *Brain Behav. Immun.* 22, 662-667.

Landys, M. M., Ramenofsky, M., Wingfield, J. C. (2006). Actions of glucocorticoids at a
seasonal baseline as compared to stress-related levels in the regulation of periodic life
processes. *Gen. Comp. Endocrinol.* 148, 132-149.

Lochmiller, R. L., and Deerenberg, C. (2000). Trade-offs in evolutionary immunology: just
what is the cost of immunity?. *Oikos* 88, 87-98.

Matson, K. D., Tieleman, B. I., & Klasing, K. C. (2006). Capture stress and the bactericidal
competence of blood and plasma in five species of tropical birds. *Physiol. Biochem. Zool.* 79,
556-564.

McDade, T. W., Georgiev, A. V., & Kuzawa, C. W. (2016). Trade-offs between acquired and
innate immune defenses in humans. *Evol. Med. Public Health* 2016, 1-16.

Merrill, L., Angelier, F., O'Loghlen, A. L., Rothstein, S. I., & Wingfield, J. C. (2012). Sex-
specific variation in brown-headed cowbird immunity following acute stress: a mechanistic
approach. *Oecologia* 170, 25-38.

Monestier, C., Gilot-Fromont, E., Morellet, N., Debeffe, L., Cebe, N., Merlet, J., et al. (2016).
Individual variation in an acute stress response reflects divergent coping strategies in a large
herbivore. *Behav. Processes* 132, 22-28.

Monestier, C., Morellet, N., Verheyden, H., Gaillard, J. M., Bideau, E., Denailhac, A., Lourtet,
B., Cebe, N., Picot, D., Rames, J. M, Hewison, A. M. (2017). Neophobia is linked to
behavioural and haematological indicators of stress in captive roe deer. *Anim. Behav.* 126, 135-
143.

Montané, J., Marco, I., López-Olvera, J., Perpinan, D., Manteca, X., Lavin, S. (2003). Effects
of acepromazine on capture stress in roe deer (Capreolus capreolus). *J. Wild. Dis.* 39, 375-386.

Montiglio, P. O., Garant, D., Pelletier, F., & Réale, D. (2012). Personality differences are
related to long-term stress reactivity in a population of wild eastern chipmunks, *Tamias striatus*.
Anim. Behav. 84, 1071-1079.

Montiglio, P. O., Garant, D., Pelletier, F., & Réale, D. (2014). Intra-individual variability in
fecal cortisol metabolites varies with lifetime exploration and reproductive life history in
eastern chipmunks (*Tamias striatus*). *Behav. Ecol. Sociobiol.* 69, 1-11.

Möstl, E., Maggs, J. L., Schrötter, G., Besenfelder, U., & Palme, R. (2002). Measurement of
cortisol metabolites in faeces of ruminants. *Vet. Res. Commun.* 26, 127-139.

Palme, R., Touma, C., Arias, N., Dominchin, M. F., & Lepschy, M. (2013). Steroid extraction:
get the best out of faecal samples. *Wien Tierarztl. Monatsschr.* 100, 238-46.

Palme, R. (2019). Non-invasive measurement of glucocorticoids: advances and problems. *Physiol. Behav.* 199, 229-243.

Pusch, E. A., and Navara, K. J. (2018). Behavioral phenotype relates to physiological differences in immunological and stress responsiveness in reactive and proactive birds. *Gen. Comp. Endocrinol.* 261, 81-88.

R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. *URL http://www. R-project. org*.

Rauw, W. M. (2012). Immune response from a resource allocation perspective. *Front. Genet.* 3, 267.

Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemanse, N. J. (2007). Integrating animal temperament within ecology and evolution. *Biol. Rev.* 82, 291-318.

Réale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., & Montiglio, P. O. (2010). Personality and the emergence of the pace-of-life syndrome concept at the population level. *Philos. Trans. R. Soc. Lond, B Biol. Sci* 365, 4051-4063.

Romero, L. M., and Wingfield, J. C. (2015). Tempests, poxes, predators, and people: stress in wild animals and how they cope. Oxford University Press.

Sanchez, G. (2013). PLS path modeling with R. Berkeley: *Trowchez Editions* 383, 2013.

Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr. rev.* 21, 55-89.

Sheldon, B. C., and Verhulst, S. (1996). Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends Ecol. Evol.* 11, 317-321.

Sih, A., Bell, A., & Johnson, J. C. (2004). Behavioral syndromes: an ecological and evolutionary overview. *Trends Ecol. Evol.* 19, 372-378.

Stier, K. S., Almasi, B., Gasparini, J., Piault, R., Roulin, A., & Jenni, L. (2009). Effects of corticosterone on innate and humoral immune functions and oxidative stress in barn owl nestlings. *J. Exp. Biol.* 212, 2085-2091.

Stöwe, M., Rosivall, B., Drent, P. J., & Möstl, E. (2010). Selection for fast and slow exploration affects baseline and stress-induced corticosterone excretion in Great tit nestlings, *Parus major*. *Horm. Behav.* 58, 864-871.

Viswanathan, K., Daugherty, C., & Dhabhar, F. S. (2005). Stress as an endogenous adjuvant: augmentation of the immunization phase of cell-mediated immunity. *Int. Immunol.* 17, 1059-1069.

Wada, T., Ishiwata, K., Koseki, H., Ishikura, T., Ugajin, T., Ohnuma, N., Obata, K., Ishikawa, R., Yoshikawa, S., Mukai, K., et al. (2010). Selective ablation of basophils in mice reveals their nonredundant role in acquired immunity against ticks. *J. Clin. Investig.* 120, 2867-2875.

Wilson, R. P., Börger, L., Holton, M. D., et al. (2020) Estimates for energy expenditure in free-living animals using acceleration proxies: A reappraisal. *J. Anim. Ecol.* 89, 161– 172. https://doi.org/10.1111/1365-2656.13040

Wingfield, J. C., Maney, D. L., Breuner, C. W., Jacobs, J. D., Lynn, S., Ramenofsky, M., & Richardson, R. D. (1998). Ecological bases of hormone—behavior interactions: the "emergency life history stage". *Am. Zool.* 38, 191-206.

Wingfield J. C, Romero L. M. (2001) Adrenocortical responses to stress and their modulation in free-living vertebrates. In: McEwen BS, Goodman HM (eds) Handbook of Physiology;

694 Section 7: The Endocrine System; Volume IV: Coping with the Environment: Neural and

695 Endocrine Mechanisms. Oxford University Press, New York, pp 211-234.

696 Zbyryt, A., Bubnicki, J. W., Kuijper, D. P., Dehnhard, M., Churski, M., & Schmidt, K. (2017).

697 Do wild ungulates experience higher stress with humans than with large carnivores? *Behav.*

698 *Ecol.* 29, 19-30.

713 **Table 1. Summary of the latent variables used in the present study.** Observed variables are

714 presented in blocks reflecting their associated latent variable.

Latent variables	Observed variables	Average values (min ; max)
Δ baseline glucocorticoids	Δ FCMs (ng/g of wet faeces)	-146 (-786 ; 346)
Δ innate immunity	Δ Neutrophils (10^3 cells/mL)	-1.23 (-4.03 ; 1.5)
	Δ Eosinophils (10^3 cells/mL)	0.01 (-0.29 ; 0.38)
	Δ Basophils (10^3 cells/mL)	0.06 (-0.36 ; 0.46)
	Δ Monocytes (10^3 cells/mL)	-0.05 (-0.37 ; 0.09)
	Δ Hemagglutination (titer)	-0.04 (-1 ; 1.5)
	Δ Hemolysis (titer)	-0.07 (-1 ; 1.5)
Δ adaptive immunity	Δ Lymphocytes (10^3 cells/mL)	0.76 (-0.80 ; 2.23)
	Δ Anti-rabies antibody (IU)	12.30 (0.60 ; 41.5)
	Δ Gamma globulins (mg/mL)	0.10 (-4 ; 3.50)
Δ inflammatory markers	Δ Alpha-1 globulins (mg/mL)	0.10 (-0.20 ; 0.80)
	Δ Alpha-2 globulins (mg/mL)	-0.06 (-0.90 ; 2.90)
	Δ Beta-2 globulins (mg/mL)	0.13 (-0.40 ; 0.50)
	Δ Haptoglobin (mg/mL)	0.13 (-0.26 ; 1.19)
Behavioural profiles	Docility score	0.5 (0 ; 1)
	Neophobia score	0.20 (0.01 ; 0.40)
	Mean daily activity	228.8 (183.6 ; 259.9)

Table 2. Characteristics of the partial least squares path modelling analyses to explain the relationships between behavioural profiles, change in baseline glucocorticoid levels, and change in innate, adaptive and inflammatory markers of immunity. GoF indicates the goodness of fit of the model. SE stands for Standard Error. See text for definition of the observed variables that composed each latent variable. Variables in bold represent endogenous variables.

Parameter	Estimate	SE	t value	P value
Structural model for innate immunity biomarkers (GoF: 0.32)				
FCMs				
Intercept	2.4 e-17	0.301	7.98 e-17	1.00
Behavioural profile	0.055	0.301	0.183	0.86
Innate immunity				
Intercept	-4.56 e-17	0.197	-2.32 e-16	1.00
Behavioural profile	-0.53	0.197	-2.70	0.02
FCMs	0.61	0.197	3.08	0.01
Structural model for adaptive immunity biomarkers (GoF: 0.42)				
FCMs				
Intercept	1.15 e-17	0.299	3.84 e-17	1.00
Behavioural profile	0.127	0.299	0.424	0.68
Adaptive immunity				
Intercept	3.43 e-17	0.172	2.00 e-16	1.00
Behavioural profile	-0.801	0.173	-4.63	0.001
FCMs	0.373	0.173	2.16	0.057
Structural model for inflammatory biomarkers (GoF: 0.39)				
FCMs				
Intercept	1.41 e-16	0.294	4.79 e-16	1.00
Behavioural profile	0.226	0.294	0.183	0.46
Inflammatory markers				
Intercept	3.64 e-16	0.192	1.90 e-15	1.00
Behavioural profile	-0.770	0.197	-3.91	0.003
FCMs	0.089	0.197	0.454	0.660

Table 3. Characteristics of the observed variables that composed each latent variable in
the three partial least squares path modelling analyses to explain the relationships
between behavioural profiles, change in baseline glucocorticoid levels, and change in
innate, adaptive and inflammatory immunity. Weight represents the contribution of the
variable to the latent variable, and loadings indicate the direction of the correlation between the
observed variables and their latent variable. Communality indicates the amount of variability
in an observed variable that is captured by its latent variable, and particularly apply for
reflective latent variables. Redundancy indicates the ability to predict for a given observed
variable, and particularly apply for formative latent variables. See text for definition of the
observed variables composing each latent variable. Variables in bold represent endogenous
variables.

Parameter	Weight	Loading
Outer model for innate immunity biomarkers		
Behavioural profile		
Lack of docility	0.521	0.847
Neophobia	0.434	0.753
Activity	0.358	0.648
Glucocorticoids		
FCMs	1.00	1.00
Innate immunity		
Neutrophils	-0.324	-0.686
Eosinophils	0.396	0.495
Monocytes	-0.533	0.274
Basophils	0.843	0.416
Hemagglutination	0.662	0.435
Hemolysis	0.245	0.367
Outer model for adaptive immunity biomarkers		
Behavioural profile		
Lack of docility	0.551	0.824
Neophobia	0.637	0.877
Activity	-0.041	0.306
Glucocorticoids		
FCMs	1.00	1.00
Adaptive immunity		
Gamma globulins	0.365	0.581
Lymphocytes	0.523	0.754
Anti-rabies antibody	0.543	0.725
Outer model for inflammatory immunity biomarkers		
Behavioural profile		
Lack of docility	-0.319	0.115
Neophobia	0.115	0.215
Activity	1.051	0.962

Glucocorticoids		
FCMs	1.00	1.00
Inflammatory immunity		
Alpha-1 globulins	0.360	0.936
Alpha-2 globulins	0.279	0.618
Beta-globulins	0.217	0.456
Haptoglobin	0.456	0.859

Table 4. Characteristics of the selected linear mixed-effect models for explaining variation

in baseline FCM levels (log-transformed) in the roe deer population of Gardouch. The

effect of PC1 (behavioural profile ranging from proactive behavioural profiles to reactive

behavioural profiles), age of individuals, period of sample collection, and Julian date of sample

collection were fitted. Models included individual identity and enclosure number as random

effects. R^{2m} and R^{2c} are the marginal and conditional explained variance of the models,

respectively. SE stands for Standard Error. See text for definition of model sets.

Parameter	Estimate	SE	t-value	P-value
$(R^{2m}: 0.06 ; R^{2c}: 0.27)$				
Intercept	6.638	0.141	47.10	< 0.001
Behavioural profile (PC1)	0.103	0.036	2.849	< 0.005
Period (2)	-0.163	0.080	-2.045	< 0.05

 Fig. 1. Summary of the experimental design used in the present study. Data relative to the

780 assessment of neophobia scores were obtained prior to this protocol. Precisely, data were

781 collected in February 2015 for all the individuals, except for two that were assessed for

782 neophobia in February 2018, following the same protocol (see details in Monestier et al. 2017).

Fig. 2. Structural models of relationships among behavioural profiles, change in baseline

glucocorticoid levels and change in immunity as determined by the partial least squares

path modelling analyses. Arrows indicates the direction of effect, and the shape of arrows

indicate the strength of the observed correlation among the latent variables. Stars indicate the

degree of significance (see Table 2 for details), and absence of star indicate the relationship was

not significant.

Fig. 3. Relationship between baseline FCMs level (log-transformed) and behavioural

profiles. Behavioural profiles' scores correspond to the score for the first axis (PC1) of the PCA

conducted using docility, activity and neophilia as co-variables. The three variables were all

negatively correlated with PC1. Thus, this axis represents a gradient of behavioural profiles,

with negative values indicating proactive behavioural profiles (high activity levels, neophobia,

and lack of docility), and positive values indicating reactive behavioural profiles (low activity

levels, neophilia, and docility). Points represent observed values, lines represent model

predictions and dashed lines represent the 95% confidence interval.

Fig. 4. Summary of the observed outcomes and relationships between the latent variables

considered in our study. Our protocol (see details in the main text) resulted in different

outcomes, with part of the individuals showing an increase in baseline cortisol between the two

study periods (indicated by an increase in FCMs), while others showed a decrease. Individuals

that showed more reactive behavioural profiles (indicated by low activity levels, docility to

manipulation, and neophilia) also exhibited an increase in baseline cortisol levels, and an

increase in immunity (both innate and adaptive immunity), while the opposite occurred for

individuals that showed more proactive behavioural profile.

826