Gene duplications and phylogenomic conflict underlie major pulses of phenotypic evolution in gymnosperms

Gregory W. Stull[1,2,†], Xiao-Jian Qu[3,†], Caroline Parins-Fukuchi[4], Ying-Ying Yang[1], Jun-Bo Yang[2], Zhi-Yun Yang[2], Yi Hu[5], Hong Ma[5], Pamela S. Soltis[6], Douglas E. Soltis[6,7], De-Zhu Li[1,2,*], Stephen A. Smith[8,*], Ting-Shuang Yi[1,2,*].

[1]Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan, China.

[2]CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan, China.

[3]Shandong Provincial Key Laboratory of Plant Stress Research, College of Life Sciences, Shandong Normal University, Jinan, Shandong, China.

[4]Department of Geophysical Sciences, University of Chicago, Chicago, IL, USA.

[5]Department of Biology, Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA, USA.

[6]Florida Museum of Natural History, University of Florida, Gainesville, FL, USA.

[7]Department of Biology, University of Florida, Gainesville, FL, USA.

[8]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA.

[†]Co-first author.

*Correspondence to: dzl@mail.kib.ac.cn; eebsmith@umich.edu; tingshuangyi@mail.kib.ac.cn.

Abstract

Inferring the intrinsic and extrinsic drivers of species diversification and phenotypic disparity

across the Tree of Life is a major challenge in evolutionary biology. In green plants, polyploidy

(or whole-genome duplication, WGD) is known to play a major role in microevolution and

speciation[1], but the extent to which WGD has shaped macroevolutionary patterns of

diversification and phenotypic innovation across plant phylogeny remains an open question.

Here we examine the relationship of various facets of genomic evolution—including gene and

genome duplication, genome size, and chromosome number—with macroevolutionary patterns

of phenotypic innovation, species diversification, and climatic occupancy in gymnosperms. We

show that genomic changes, such as WGD and genome-size shifts, underlie the origins of most

major extant gymnosperm clades, and notably our results support an ancestral WGD in the

gymnosperm lineage. Spikes of gene duplication typically coincide with major spikes of

phenotypic innovation, while increased rates of phenotypic evolution are typically found at

nodes with high gene-tree conflict, representing historic population-level dynamics during

speciation. Most shifts in gymnosperm diversification since the rise of angiosperms are

decoupled from putative WGDs and instead are associated with increased rates of climatic

occupancy evolution, particularly in cooler and/or more arid climatic conditions, suggesting that

ecological opportunity, especially in the later Cenozoic, and environmental heterogeneity have

driven a resurgence of gymnosperm diversification. Our study provides critical insight on the

processes underlying diversification and phenotypic evolution in gymnosperms, with important

broader implications for the major drivers of both micro- and macroevolution in plants.

Main Text

Connecting microevolutionary processes with emergent macroevolutionary patterns of

phenotypic disparity and species diversification is one of the grand challenges of evolutionary

biology. In green plants, polyploidy is a common phenomenon in many lineages[1,2] that has long

been considered a major evolutionary force[3,4], unique in its ability to generate manifold

phenotypic novelties at the population level and in its potential to drive long-term evolutionary

trends. However, despite its prevalence and clear microevolutionary importance, its

macroevolutionary significance remains unclear and, in major clades such as gymnosperms,

understudied. For example, analyses of the relationship between WGD and diversification have

produced conflicting results[5,6,7,8], and no studies to our knowledge have examined whether spikes

in gene duplication tend to directly coincide with bursts of phenotypic novelty on a

macroevolutionary scale, although clearly gene duplications have facilitated the origins of

numerous key traits in plants[9,10,11]. It is possible that other processes—e.g., major fluctuations in

population size during rapid speciation events, in concert with extrinsic factors such as climate

change or ecological opportunity—may contribute more fundamentally to major phases of

morphological innovation and diversification[12,13]. More realistically, however, these various

phenomena may interact in complex ways to shape macroevolutionary patterns in plants and,

more broadly, across the Tree of Life, and detailed case studies are needed to disentangle their

relative contributions to the origins of phenotypic and taxonomic diversity.

Here we examine potential drivers of phenotypic innovation and species diversification in

seed plants (=*Spermatophyta*[14]), with a focus on *Acrogymnospermae*[14], the most ancient subclade

of living spermatophytes, including all extant gymnosperms (conifers, cycads, Gnetales, and

Ginkgo biloba) as well now-extinct groups such as Cordaitales[15]. For the past ~300 million years,

seed plants have occupied center stage in terrestrial ecosystems[16], constituting the most dominant

plant clade in terms of both biomass and diversity and forming myriad ecological relationships

with other organisms across the Tree of Life, such as bacteria, fungi, insects, and vertebrates[17].

Reconstructing the diversification of seed plants is therefore essential for a better understanding

of the history of terrestrial ecosystems across the latter half of the Phanerozoic. Many aspects of

seed plant phylogeny remain poorly known or contentious[18], including the phylogenetic

placements of numerous extinct lineages (e.g., Bennettitales and Caytoniales) with respect to

living gymnosperms (*Acrogymnospermae*) and flowering plants (*Angiospermae*[14]). Nevertheless,

based on the extensive fossil record of seed plants[19] and phylogenetic studies of living

representatives *Acrogymnospermae*[20,21] (which comprise ca. 1100 species), extant gymnosperm

lineages clearly exhibit a complex history of ancient radiations, major extinctions, extraordinary

stasis, and recent diversification, but the correlates and causes of major phases of gymnosperm

evolution remain understudied. Given that recent studies[22–25] have challenged the traditional

notion[26,27] that WGD is rare or insignificant in gymnosperms, a focused investigation on the

prevalence and macroevolutionary significance of polyploidy in gymnosperms is timely.

We synthesize phylogenomic analyses, trait reconstructions, and examinations of

extensive comparative data on a comprehensive phylogeny of gymnosperms to understand major

processes that have shaped phenotypic innovation, species diversification, and climatic evolution

in this clade. Specifically, we analyze a novel transcriptome dataset including representatives of

nearly all extant genera of gymnosperms to identify hotspots of genomic conflict and gene

duplication across gymnosperm phylogeny, which we relate to reconstructed bursts of

phenotypic innovation and major shifts in species diversification and climatic evolution.

Gymnosperm phylogenomics, polyploidy, and genome evolution

Based on phylogenomic analysis of our transcriptome dataset (including 790 inferred orthologs

across 121 ingroup species, 77 of which were newly sequenced for this study; Supplementary

Table 1), we recovered a well-resolved and highly supported species tree for gymnosperms (Fig.

1). We then took a multifaceted approach for WGD inference to circumvent the limitations of

some methods and datatypes[28]. Using the species-tree phylogeny and the inferred orthogroups,

we performed gene-tree reconciliation to identify hotspots of gene duplication supporting WGD

(Fig. 1; Methods), and compared these results against synonymous substitution (*Ks*) plots

(Extended Data Figs. 1 and 2; Supplementary Fig. 1; data deposit), as well as reconstructions of

chromosome number (Supplementary Fig. 2) and genome size (Extended Data Fig. 3) across a

broader sampling of gymnosperm species from our supermatrix phylogeny (Methods). Particular

attention was paid to branches with previously inferred (and debated) instances of WGD in

gymnosperms, such as the gymnosperm root, Pinaceae, and Cupressaceae[22,29–32].

We inferred elevated levels of gene duplication (Fig. 1) in the branches subtending seed

plants (56.2 % = percentage of orthogroups showing a duplication), gymnosperms (41.7 %),

Coniferae[14] (including Gnetales, 18.8 %), Pinaceae (12.1 %), *Cupressophyta*[14] (i.e.,

Araucariaceae + Podocarpaceae + Sciadopityaceae + Taxaceae + Cupressaceae; 13.9 %),

Podocarpaceae (11.3 %), and Araucariaceae (10.1 %). Most of these branches also show at least

one of the following features: shared Ks peaks, inferred changes in chromosome number, or

extreme shifts in genome size (Fig. 1; Extended Data Figs. 1–3; Supplementary Fig. 1): seed

plants (shared Ks peak), *Acrogymospermae* (shared Ks peak), *Cupressophyta* ($n = 7$ to $n = 11$),

Pinaceae (shared Ks peak; $n = 7$ to $n = 12$), Podocarpaceae ($n = 11$ to $n = 10$; major C-value

decrease), and Araucariaceae ($n = 11$ to $n = 13$). Cycads and *Ginkgo* show clear Ks spikes around

0.8, previously thought to represent a potential WGD event unique to this clade[30], but these

paralog Ks peaks are deeper than the ortholog Ks peak for gymnosperms as a whole (Extended

Data Fig. 1), suggesting instead that these peaks represent a gymnosperm-wide WGD event. This

putative gymnosperm peak is also evident in Ks plots of conifers, but less prominently and at a

Ks range of around 1.0 to 1.5. We suggest these Ks differences between 'ginkads' and conifers

might be due to differences in rates of molecular evolution or differences in the timing of

diploidization in each lineage if they diverged prior to the diploidization of the polyploid

gymnosperm ancestor.

Ancestral chromosomal changes were also observed in Cycadaceae ($n = 7$ to $n = 11$),

Zamiaceae ($n = 7$ to $n = 8$), and Gnetaceae ($n = 7$ to $n = 12$), as well as within Cupressaceae,

Podocarpaceae, and the genera of Gnetales, due to more recent paleopolyploidy or

neopolyploidy[24,25,33] (Supplementary Fig. 2). Regarding genome size, rate shifts (inferred using

BAMM) and/or major 'jumps' (i.e., extreme changes in reconstructed ancestor to descendant C-

values) in genome size evolution (Methods) were inferred at the origins of several major clades,

including abrupt genome downsizing in Podocarpaceae (as noted above), Gnetales, and

Cycadaceae and rate shifts in Zamiaceae (excluding *Dioon*), Taxaceae + Cupressaceae, and

Ephedraceae; jumps or rate shifts were also inferred in numerous more recent clades (Extended

Data Fig. 3). See Supplementary Information for additional results and discussion of

gymnosperm genome evolution, including more detailed comparisons of our results with those of

previous studies on WGD in gymnosperms[22,29,30,32].

Genomic correlates of diversification and phenotypic evolution

Our diversification analysis (Supplementary Fig. 3), based our dated supermatrix phylogeny of

extant gymnosperms (including 82% species representation), recovered 17 major shifts, only two

of which correspond directly to branches with possible WGDs: *Cupressophyta* and Pinaceae

(Fig. 1). Seven other diversification shifts corresponded to major jumps or rate shifts in genome

size evolution or changes in chromosome number (Fig. 1, Extended Data Fig. 3, and

Supplementary Information). Although most diversification shifts (15/17) were not associated

with putative WGDs, it is possible that major extinctions have obscured instances of

diversification shifts associated with WGD in deep time. On the other hand, extinction of

depauperate branches separating a WGD event and a diversification shift could also lead to their

erroneous association. With these considerations in mind, we argue at least that WGD has played

a minimal role in driving the more geologically recent diversification shifts detected here. Our

results also align with recent studies showing an inconsistent relationship between WGD and

diversification[5,6,7]. We suggest that in cases where WGD is associated with diversification, it is a

product of WGD mediating changes in traits (such as key innovations[9]) or life history that

influence population structuring, with diversification only being a secondary consequence[7]. This

is exemplified by Pinaceae, where a possible ancestral WGD is associated with not only a shift in

diversification (Fig. 1) but also a major spike in morphological innovation (Fig. 2) and a rate

shift in climatic evolution (see below).

To examine the general relationship of gene duplications and genomic (gene-tree)

conflict with levels of phenotypic innovation and rates of phenotypic evolution, we analyzed a

newly compiled trait dataset for gymnosperms (available in the data deposit) comprising 148

phenotypic characters, with sampling matched to the transcriptome phylogeny (Supplementary

Information). Sub- and neo-functionalization of duplicate genes are considered the primary

means by which polyploidy can lead to phenotypic novelty[34,35], and therefore an examination of

the direct correspondence of gene duplication numbers to levels of phenotypic novelty at nodes

across gymnosperm phylogeny serves as a means of assessing the potential contribution of large-

scale (WGD) and small-scale genomic events to macroevolutionary patterns of phenotypic

evolution. Genomic conflict, on the other hand, records information about population-level

processes during speciation events—e.g., rapid population expansions or fragmentation, resulting

in incomplete lineage sorting (ILS), or introgression—that might also play a role in rapid

acquisition of phenotypic novelties[36]. Thus, these represent two different, albeit not mutually

exclusive, processes that might shape macroevolutionary patterns of morphological evolution in

plants. We reconstructed levels of phenotypic innovation or novelty (defined here simply as state

changes along a branch) and rates (state changes/time) for each branch in the phylogeny[13] and

plotted these against corresponding levels of gene duplication and gene-tree conflict (Figs. 2 and

3). Linear regression and permutation tests were also performed to evaluate the statistical

strength of these relationships (Methods).

We find a positive relationship between levels of phenotypic innovation and levels of

gene duplication (linear regression: $r^2 = 0.2038$, $p = 0.000$; permutation test: $p = 0.038$;

Supplementary Figs. 4 and 5), suggesting that gene duplications may have played a vital role in

the generation of phenotypic novelty in gymnosperms evident at a macroevolutionary scale (Fig.

2). Major examples include Araucariaceae, Cupressaceae, *Pinus*, Pinaceae, Podocarpaceae, and

Zamiaceae, which all show corresponding, elevated levels of gene duplication and phenotypic

innovation. There are also evident temporal patterns, with, for example, a noticeable increase in

duplications and phenotypic innovation near the present (Fig. 2). While causation between gene

duplication and phenotypic innovation cannot be demonstrated on the basis of these observed

patterns, this connection is plausible in cases of complex innovations and clearly deserves future

study given that gene duplications have long been considered to play a pivotal role in the origin

of evolutionary novelties[34]. It has been suggested that small-scale and large-scale (e.g., WGD)

duplication events might tend to have different immediate consequences and evolutionary

trajectories[35]. Some of the putative WGD events examined here seem most salient in terms of

morphological innovation, but the general correspondence (Supplementary Figs. 4 and 5)

suggests that perhaps both large- and small-scale duplication events have played an important

role in the generation of novel traits in gymnosperms. It is possible that some of the

phylogenetically deeper spikes in phenotypic innovation are exaggerated due to prevalent

extinction of stem lineages with intermediate morphologies. However, the sudden appearance of

major lineages with distinct morphologies is a common phenomenon across the Tree of Life[13],

and neofunctionalization of duplicate genes (and/or other results of gene or genome duplication)

may serve as an important driver of instances of radical and rapid phenotypic change. A better

integration of fossil and phylogenomic data might offer important insight into this question.

Although we observed a significant relationship between gene duplication and

phenotypic novelty, there are notable discrepancies, the most salient being Gnetales (Fig. 2).

However, this is perhaps not surprising given that Gnetales have experienced significant life

history changes and stands out among gymnosperms in having relatively small genomes (despite

at least recent WGD events[22,25]) and remarkably fast ("angiosperm-like") rates of molecular

evolution[37,38]. *Coniferae* (including Gnetales) also represents a major discrepancy, and the

exclusion of Gnetales from the phenotypic reconstructions results in an increase in the number of

inferred apomorphies for conifers (Supplementary Fig. 6). However, another explanation for the

low levels of innovation inferred for the *Coniferae* node might be the prevalence of traits shared

between conifers and *Ginkgo*. Although our species tree places *Ginkgo* sister to cycads (Fig. 1),

morphological phylogenetic analyses of seed plants[18] have typically placed *Ginkgo* with extinct

and extant conifers (rather than cycads), and among our gene trees (Fig. 1) the dominant

alternative topology for *Ginkgo* is sister to *Coniferae* (with 81 gene trees showing this alternative

topology with strong support). This finding suggests that the proposed gymnosperm-wide WGD

event might have involved reticulate evolution in the early history of *Acrogymnospermae*,

providing a partial explanation for this discrepancy. It is also noteworthy that several well-

documented gymnosperm polyploids (e.g., *Sequoia sempervirens*, *Fitzroya cupressoides*[25]) do

not show major phenotypic changes. However, these species both show polysomic inheritance[33]

with gene copies still functioning as alleles, thus constraining the ability of duplicate genes to

generate novel phenotypes via neofunctionalization or other processes.

In contrast to cumulative levels of phenotypic innovation, rates of phenotypic evolution

show a strong relationship with levels of gene-tree conflict (Fig. 3; linear regression: $r^2 = 0.2308$,

$p = 0.000$; Supplementary Fig. 7), with extreme values of conflict and rate in particular tending

to co-occur (permutation test: $p = 0.001$; Supplementary Fig. 8). This finding is in line with a

recent study[13] documenting a similar correspondence in multiple clades across the Tree of Life.

Figure 3 clearly shows that the branches with the highest rates of morphological evolution also

have extreme levels of gene-tree conflict. A noteworthy example is the Podocarpaceae subclade

including *Phyllocladus* and *Parasitaxus* (Fig. 3), which is characterized by extensive conflict as

well as remarkable shifts in phenotype (e.g., the acquisition of phylloclades) and life history

(e.g., parasitism) in several of the constituent lineages.

If gene-tree conflict reflects a signature of historical population-level processes such as

ILS[39], the observed relationship between conflict and phenotypic rates suggests that, at

macroevolutionary scales, many instances of rapid morphological evolution in gymnosperms are

a product of episodic and dramatic population perturbations associated with rapid speciation

events. Given the life history characteristics of (most) gymnosperms—including large genome

sizes, long time to reproduction, and, in some species, exceptional longevity[40,41]—instances of

population reduction or isolation should theoretically be essential for instances of rapid

phenotypic evolution. Thus, areas of gymnosperm phylogeny with elevated conflict and

morphological rates perhaps reflect the rapid fragmentation of broad, variable ancestral

populations. As discussed below, this type of process might explain the prevalence of young

conifer and cycad species with highly restricted distributions today. We also note that the

processes outlined above are not mutually exclusive putative drivers of phenotypic evolution,

given that gene duplications might serve as a source of novelties sorted by population-level

processes and that WGD can theoretically result in the rapid establishment of small,

reproductively isolated populations[9,42], which could result in the rapid fixation of polymorphisms

via genetic drift.

Temporal patterns of climatic occupancy evolution and species diversification

Reconstructions of climatic variables on the dated supermatrix phylogeny—focusing on mean

annual temperature (Bio 1), mean annual precipitation (Bio 12), and principal components one

and two of the 19 WorldClim bioclimatic layers—revealed that rate shifts and jumps in climatic

occupancy are concentrated in the last ~70 Myr (and particularly the last ~25 Myr;

Supplementary Figs. 9–14). Additionally, many of the branches (10/17) with a diversification

shift also included a shift in the rate of climatic evolution (Fig. 4), with four additional

diversification shifts separated from a climatic shift by only one or two nodes (i.e., 14/17

diversification shifts are directly or strongly associated with climatic shifts; Fig. 4). Except for

two of the concerted shifts (in the branches subtending the cycad genera *Ceratozamia* and

Zamia), all others involved shifts associated with cooler and/or drier climatic conditions (Fig. 4

and Supplementary Figs. 9–14). Undoubtedly, there have been many deeper bursts of

diversification and shifts in climatic occupancy in gymnosperms that are not evident in our

phylogeny due to extinction[19]; this renders the inference of diversification dynamics across the

depth of gymnosperm history a major, if not insurmountable, challenge without extensive data

from the fossil record[43]. Nevertheless, considering an extensive body of research concerning

gymnosperm vs. angiosperm functional morphology and competitive dynamics today[44] and

patterns in the fossil record[45], we might predict many gymnosperms (and especially conifers) to

thrive with the expansion of cooler and more arid conditions after late Eocene climatic

deterioration and particularly after the mid-Miocene[46].

The global ecological ascendance of angiosperms, from the early Cretaceous to early

Cenozoic, was a complex and relatively protracted process, with gymnosperms showing

heterogeneous patterns of decline, in terms of abundance and diversity, across different lineages,

ecological guilds, geographic areas, and points in time[45,47]. It was not until the early Cenozoic,

perhaps in part facilitated by major ecological upheaval at the K-Pg boundary, that angiosperms

came to dominate (or co-dominate) nearly all geographic areas and vegetation types[47]. Numerous

attributes of angiosperms may underpin their macroevolutionary success, including their

reproductive innovations and ecological associations[48], proclivity for polyploidy[32,49,50], and faster

rates of molecular evolution[37]. However, their functional morphology and correspondingly

elevated growth rates have been highlighted as particularly important in the context of

competition with gymnosperms[44]. The early Cenozoic was a time of warm, equable climate—

conditions well suited to rapid growth strategies of many angiosperms, facilitated by broad

leaves and a more efficient vascular system including vessels. Gymnosperms, with narrow

tracheids and (generally) smaller, more poorly vascularized evergreen leaves, may have

struggled to compete with angiosperms across much of the landscape of the early Cenozoic given

the growth-rate limitations imposed by their less efficient vascular systems[44]. Exceptions that

prove the rule include *Gnetum* and some members of Podocarpaceae that independently evolved

angiosperm-like vasculature and/or broader leaves, allowing them to compete in wet tropical

conditions where gymnosperms (especially conifers) are otherwise largely excluded[51].

However, the same basic traits (and consequent slow-growth strategies) that exclude most

gymnosperms from habitats that are more favorable to angiosperm plant growth allow many to

persist in more extreme, stressful habitats[52]. The expansion of cool and arid conditions after the

Eocene and especially after the mid-Miocene therefore may have created many new

opportunities for gymnosperm expansion and diversification. This hypothesis is consistent with

our results showing many gymnosperm lineages shifting into cooler and/or more arid conditions

after the late Eocene (Supplementary Figs. 9–14). Diversification shifts associated with cool/arid

transitions that occurred prior to the late Eocene (e.g., *Pinus* subsection *Ponderosae*, ~56 Ma; the

Juniperus-Cupressus clade, ~70 Ma; the *Podocarpus-Retrophyllum* clade, ~68 Ma) might

represent competitive displacement by angiosperms and/or expansion into upland or montane

habitats[53]. Future biogeographic analyses integrating fossil evidence might shed light on the

geographic context of these coordinated climatic-diversification shifts. In general, our results

challenge suggestions that recent climatic cooling precipitated macroevolutionary declines in

conifers[54] (Supplementary Information). The Northern and Southern Hemispheres, however,

clearly do show distinct patterns with respect to the extinction and turnover of conifer diversity

during the Cenozoic[21], and numerous extant gymnosperm species (especially cycads) are at risk

of extinction due to pressures including habitat destruction[55,56].

While climatic shifts have played a significant role in the Cenozoic history of

gymnosperms, many major climatic shifts are not associated with changes in diversification (Fig.

4 and Supplementary Figs. 9–14), suggesting that climatic evolution might be an important but

insufficient vehicle for diversification. Ecological factors (e.g., environmental heterogeneity

related to elevation or soil type) and/or traits (e.g., dispersal strategies) that more directly control

population size and spatial structuring are likely necessary concomitant factors for driving

diversification patterns in gymnosperms, and more broadly in green plants. This finding is

consistent with the observation that many recent gymnosperm radiations (e.g., multiples lineages

of cycads, Pinaceae, and Cupressaceae) include numerous species with insular/restricted

distributions[57]. The small distributions of many gymnosperm species have typically been

interpreted as relictual[40], but clearly many of these species stem from recent, Neogene radiations

given their remarkably short molecular branch lengths[20,21,58]. Aside from these numerous recent

radiations, however, extensive data from the fossil record show that many extant

gymnosperms—e.g., *Ginkgo biloba*, members of Taxaceae (including *Cephalotaxus*), and the

"taxodioid" Cupressaceae—are clearly relics of historically more diverse and broadly distributed

groups.

Macroevolutionary synthesis

Our analyses provide an unprecedented and multifaceted view of genome evolution in

gymnosperms, with broad implications for how different attributes of genome evolution might

both drive and signal major phases of phenotypic evolution in plants. Using multiple lines of

evidence, we show that polyploidy and dynamic shifts in chromosome number and genome size

are common occurrences across the phylogenetic breadth of gymnosperms. We also provide

compelling evidence for a WGD event in the direct ancestor of gymnosperms, a possibility that

has been suggested previously but with limited and conflicting evidence[59]. Furthermore, we

provide one of the first direct documentations of a general relationship between levels of gene

duplication and phenotypic innovation at a macroevolutionary scale. Gene duplications have

long been emphasized as a fundamental source of novelty, but their evolutionary outcomes have

typically been studied at a microevolutionary scale or with respect to particular traits or nodes of

interest[9,10,11] (but see Walden et al.[8]). The broader patterns documented here provide a framework

of promising areas of gymnosperm phylogeny in which to investigate connections between

duplications or gene-family expansions and phenotypic novelties.

Contrary to the association of duplications and innovation, we do not observe a consistent

relationship between polyploidy and diversification—consistent with other recent studies also

showing a sporadic association[5,6,7]. This implies that, across the euphyllophyte radiation, WGD is

more fundamentally tied to phenotypic innovation, and that subsequent diversification is

contingent on the nature of the traits acquired and the presence of ecological opportunity[9]. In

gymnosperms, recent phases of diversification appear closely tied with shifts in climatic

occupancy, suggesting that the expansion of cool and arid climates in the later Cenozoic created

new ecological opportunities suited to the basic functional morphology of this clade. Finally, we

show that elevated rates of phenotypic evolution appear most closely tied to phylogenetic regions

with extensive gene-tree conflict—a signature of dynamic population processes during rapid

speciation events. Collectively these threads create a nuanced picture of how different

processes—including gene and genome duplication, population dynamics such as ILS and

introgression, and ecological opportunity in the face of climate change—may interact to drive the

emergence of novel morphologies and major radiations in plants.

1. Otto, S. P. & Whitton, J. Polyploid incidence and evolution. *Ann. Rev. Genet.* **34**, 401–437

(2000).

2. Wood, T. E. et al. The frequency of polyploid speciation in vascular plants. *Proc. Natl. Acad.*

Sci. U.S.A. **106**, 13875–13879 (2009).

3. Stebbins, G. L. *Variation and evolution in plants* (Columbia Univ. Press, 1950).

4. Levin, D. A. Polyploidy and novelty in flowering plants. *Am. Nat.* **122**, 1–25 (1983).

5. Tank, D. C. et al. Nested radiations and the pulse of angiosperm diversification: increased
diversification rates often follow whole genome duplications. *New Phytol.* **207**, 454–467
(2015).

6. Landis, J. B. et al. Impact of whole-genome duplication events on diversification rates in
angiosperms. *Am. J. Bot.* **105**, 348–363 (2018).

7. Smith, S. A. et al. Disparity, diversity, and duplications in the Caryophyllales. *New Phytol.*
217, 836–854 (2017).

8. Walden, N. et al. Nested whole-genome duplications coincide with diversification and high
morphological disparity in Brassicaceae. *Nature Comm.* **11**, 3795 (2020).

9. Soltis, P.S. & Soltis, D. E. Ancient WGD events as drivers of key innovations in angiosperms.
Curr. Opin. Genet. Dev. **30**, 159–165 (2016).

10. Guo, J., et al. Phylotranscriptomics in Cucurbitaceae reveal multiple whole-genome
duplications and key morphological and molecular innovations. *Mol. Plant* **13**, 1117–1133
(2020).

11. Sheehan, H. et al. Evolution of L-DOPA 4,5-dioxygenase activity allows for recurrent
specialization to betalain pigmentation in Caryophyllales. *New Phytol.* **227**, 914–929 (2020).

12. Stroud, J. T. & Losos, J. B. Ecological opportunity and adaptive radiation. *Annu. Rev. Ecol.*
Evol. Syst. **47**, 507–532 (2016).

13. Parins-Fukuchi, Stull, G. W. & Smith, S. A. Phylogenomic conflict coincides with rapid
morphological innovation. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2023058118 (2021).

14. Cantino, et al. Towards a phylogenetic nomenclature of *Tracheophyta*. *Taxon* **56**, E1–E44.

15. Doyle, J. A. Phylogenetic analyses and morphological innovations in land plants. *Ann. Plant*
Rev. **45**, 1–50 (2013).

16. DiMichele, W. A., Pfefferkorn, H. W. & Gastaldo, R. A. Response of Late Carboniferous
and Early Permian plant communities to climate change. *Annu. Rev. Earth Pl. Sci.* **29**, 461–
487 (2001).

17. Behrensmeyer, A. K., et al. *Terrestrial ecosystems through time: evolutionary paleoecology*
of terrestrial plants and animals (Univ. Chicago Press, 1992).

18. Crane, P. R. Phylogenetic analysis of seed plants and the origin of angiosperms. *Ann. Mo.*
Bot. Gard. **72**, 716–793 (1985).

19. Taylor, E. L., Taylor, T. N. & Krings, M. *Paleobotany: the biology and evolution of fossil*
plants (Academic Press, 2009).

20. Nagalingum, N. S. et al. Recent synchronous radiation of a living fossil. *Science* **334**, 796–
799 (2011).

21. Leslie, A. B. et al. Hemisphere-scale differences in conifer evolutionary dynamics. *Proc.*
Natl. Acad. Sci. U.S.A. **109**, 16217–16221 (2012).

22. Li, Z. et al. Early genome duplications in conifers and other seed plants. *Sci. Adv.* **1**,
e1501084 (2015).

23. Guan, R. et al. Draft genome of the living fossil *Ginkgo biloba*. *GigaScience* **5**, 49 (2016).

24. Farhat, P. et al. Polyploidy in the conifer genus *Juniperus*: an unexpectedly high rate. *Front.*
Plant Sci. **10**, 676 (2019).

25. Ickert-Bond, S. M. et al. Polyploidy in gymnosperms—Insights into the genomic and
evolutionary consequences of polyploidy in *Ephedra*. *Mol. Phylogenet. Evol.* **147**, 106786
(2020).

26. Khoshoo, T. N. Polyploidy in gymnosperms. *Evolution* **13**, 24–39 (1959).

27. Lewis, W. H. *Polyploidy—Biological Relevance* (Plenum Press, 1980).

28. Yang, Y. et al. Improved transcriptome sampling pinpoints 26 ancient and more recent
polyploidy events in Caryophyllales, including two allopolyploidy events. *New Phytol.*, **217**,
855–870 (2018).

29. Jiao, Y. et al. Ancestral polyploidy in seed plants and angiosperms. *Nature* **473**, 97–100
(2011).

30. Roodt, D. et al. Evidence for an ancient whole genome duplication in the cycad lineage.
PLoS One **12**, e0184454 (2017).

31. Ruprecht, C. et al. Revisiting ancestral polyploidy in plants. *Science Adv.* **3**, e1603195
(2017).

32. Leebens-Mack, J. H. et al. One thousand plant transcriptomes and the phylogenomics of
green plants. *Nature* **574**, 679–685 (2019).

33. Scott, A. D., Stenz, N. W. M., Ingvarsson, P. K. & Baum, D. A. Whole genome duplication
in coast redwood (*Sequoia sempervirens*) and its implications for explaining the rarity of
polyploidy in conifers. *New Phytol.* **211**, 186–193 (2016).

34. Ohno, S. *Evolution by Gene Duplication* (Springer Verlag, 1970).

35. Rensing, S. A. Gene duplication as a driver of plant morphogenetic evolution. *Curr. Opin.*
Plant Biol. **14**, 43–48 (2014).

36. Marques, D. A., Meier, J. I. & Seehausen, O. A combinatorial view on speciation and
adaptive radiation. *Trends Ecol. Evol.* **34**, 531–544 (2019).

37. De La Torre, A. R., Li, Z., Van de Peer, Y. & Ingvarsson, P. K. Contrasting rates of
molecular evolution and patterns of selection among gymnosperms and flowering plants. *Mol.*
Biol. Evol. **34**, 1363–1377 (2017).

38. Ran, J.-H., Shen, T.-T., Wang, M.-M. & Wang, X.-Q. Phylogenomics resolves the deep
phylogeny of seed plants and indicates partial convergent or homoplastic evolution between
Gnetales and angiosperms. *Proc. R. Soc. B* **285**, 20181012 (2018).

39. Oliver, J. C. Microevolutionary processes generate phylogenomic discordance at ancient
divergences. *Evolution* **67**, 1823–1839 (2013).

40. Farjon, A. *A natural history of conifers* (Timber Press, 2008).

41. Pellicer, J., Hidalgo, O., Dodsworth, S. & Leitch, I. J. 2018. Genome size diversity and its
impact on land plants. *Genes* **9**, 88 (2018).

42. Otto, S. P. 2007. The evolutionary consequences of polyploidy. *Cell* **131**, 452–462.

43. Louca, S. & Pennell, M. W. Extant timetrees are consistent with a myriad of diversification
histories. *Nature* **580**, 502–505 (2020).

44. Bond, W. J. The tortoise and the hare: ecology of angiosperm dominance and gymnosperm
persistence. *Biol. J. Linn. Soc.* **36**, 227–249 (1989).

45. Lupia, R., Ligard, S. & Crane, P. R. Comparing palynological abundance and diversity:
implications for biotic replacement during the Cretaceous angiosperm radiation. *Paleobiology*
25, 305–340. (1999).

46. Zachos, J., Pagani, M., Sloan, L., Thomas, E. & Billups, K. Trends, rhythms, and aberrations
in global climate 65 Ma to present. *Science* *292*, 686–693 (2001).

47. Wing, S. L., & Boucher, L. D. Ecological aspects of the Cretaceous flowering plant
radiation. *Annu Rev. Earth Plant. Sci.* **26**, 379–421 (1998).

48. Tiffney, B. H. Seed size, dispersal syndromes, and the rise of the angiosperms: evidence and
hypothesis. *Ann. Missouri Bot.* **71**, 551–576 (1984).

49. Rice, A., et al. The Chromosome Counts Database (CCBD)—a community resource of plant
chromosome numbers. *New Phyt.* **206**, 19–26 (2015).

50. Pellicer, J., & Leitch, I. J. The Plant DNA C-values database (release 7.1): an updated online

repository of plant genome size data for comparative studies. *New Phyt.* **226**, 301–305 (2020).

51. Biffin, E. et al. Leaf evolution in Southern Hemisphere conifers tracks the angiosperm

ecological radiation. *Proc. R. Soc. B* **279**, 341–348 (2012).

52. Brodribb, T. J., McAdam, S. A., Jordan, G. J. & Martins, S. C. Conifer species adapt to low-

rainfall climates by following one of two divergent pathways. *Proc. Natl. Acad. Sci.*

U.S.A. **111**, 14489–14493 (2014).

53. English, J. M. & Johnson, S. T. The Laramide orogeny: what were the driving forces? *Int.*

Geol. Rev. **46**, 83–838 (2004)

54. Condamine, F. L., Silvestro, D., Koppelhus, E. B. & Antonelli, A. The rise of angiosperms

pushed conifers to decline during global cooling. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 28867–

28875 (2020).

55. Fragnière, Y., Bétrisey, S., Cardinaux, L., Stoffel, M. & Kozlowski, G. Fighting their last

stand? A global analysis of the distribution and conservation status of gymnosperms. *J.*

Biogeogr. **42**, 809–820 (2015).

56. Forest, F. et al. Gymnosperms on the EDGE. *Sci. Rep.* **8,** 6053 (2018).

57. Rundel, P. W. A Neogene Heritage: conifer distributions and endemism in Mediterranean-

climate ecosystems. *Front. Ecol. Evol.* **7**, 364 (2019).

58. Jin, W.-T., et al. Phylogenomic and ecological analyses reveal the spatiotemporal evolution

of global pines. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2022302118 (2021).

59. Zwaenepoel, A., Van de Peer, Y. Inference of ancient whole-genome duplications and the

evolution of gene duplication and loss rates. *Mol. Biol. Evol.* **36**, 1384–1404 (2019).

Acknowledgements. We thank the Germplasm Bank of Wild Species at the Kunming Institute

of Botany (KIB) for facilitating this study. We thank the curators and staff of the Kunming

Botanical Garden of the Kunming Institute of Botany, the University of California Botanical

Garden at Berkeley, the Arnold Arboretum of Harvard University, the Missouri Botanical

Garden, the Royal Botanic Garden Edinburgh, and the Royal Botanical Gardens Kew for

providing fresh and silica-dried leaves and DNA samples. We also thank Andrew Leslie and an

anonymous reviewer for their thoughtful comments and suggestions, which greatly improved the

manuscript. This work was funded by the Strategic Priority Research Program of the Chinese

Academy of Sciences (CAS) (grant No. XDB31000000 to D.-Z.L. and T.-S.Y.), CAS' large-

scale scientific facilities (grant No. 2017-LSF-GBOWS-02 to D.-Z.L.), the National Natural

Science Foundation of China [key international (regional) cooperative research project No.

31720103903 to T.-S.Y.], the Yunling International High-end Experts Program of Yunnan

Province (grant No. YNQR-GDWG-2017-002 to P.S.S. and YNQR-GDWG-2018-012 to

D.E.S.), and the Natural Science Foundation of Shandong province (ZR2020QC022 to X.-J.Q.).

G.W.S. acknowledges support from the Chinese Academy of Sciences (CAS) President's

International Fellowship Initiative (No. 2020PB0009) and the China Postdoctoral Science

Foundation (CPSF) International Postdoctoral Exchange Program.

Author contributions. G.W.S., D.-Z.L., S.A.S., and T.-S.Y. conceived the study; X.-J.Q., Y.-

Y.Y., Y.H., J.-B.Y., Z.-Y.Y., and H.M. and collected and prepared samples for transcriptome

and plastome sequencing; G.W.S. generated the trait dataset, and compiled publically available

data for the supermatrix and comparative analyses; G.W.S. conducted analyses with help from

C.P.-F., S.A.S., and X.-J.Q; G.W.S, C.P.-F., P.S.S., D.E.S., S.A.S., and T.-S.Y. interpreted the

results; G.W.S. wrote the manuscript, with contributions from C.P.-F., P.S.S., D.E.S., S.A.S., and

T.-S.Y. All authors approved the manuscript.

Figures

Figure 1. **Transcriptome species tree showing major genomic events**. Species tree inferred

from the 790 gene trees using ASTRAL, scaled to time by fixing the nodes to the ages inferred in

the supermatrix dating analysis. Pie charts at each node show the proportion of gene trees in

concordance (blue), in conflict (chartreuse = dominant alternative; magenta = other conflicting

topologies), or without information for that particular bipartition (grey). Nodes with the highest

percentages of duplicate genes (indicative of WGD) are highlighted in the inset plot in the top

left corner; the percentages of duplicate genes inferred for these outlier nodes are also displayed

on the tree. Orange stars indicated WGD events supported by *Ks* plots; uncertainty regarding the

phylogenetic placement(s) of such events is indicated with a question mark. Changes in the

haploid chromosome number (based on the ChromEvol analyses; Methods) are indicated on the

phylogeny. Black circles indicate diversification rate shifts (inferred using BAMM; Methods).

Plant silhouettes were obtained from PhyloPic (http://phylopic.org); credit is acknowledged for

the silhouettes of *Araucaria angustifolia* (LeonardoG and T. Michael Keesey); *Zamites* (Robert

Gay), which closely resembles the architecture of cycads; and *Ginkgo* (Joe Schneid and T.

Michael Keesey). The *Taxus* silhouette was generated from a line drawing of *Taxus brevifolia*

obtained from the website of Natural Resources Canada, Canadian Forest Service

(https://tidcf.nrcan.gc.ca/en/trees/factsheet/288).

Figure 2. **Gene duplications vs. phenotypic innovation**. Plots juxtaposing levels of gene

duplication and phenotypic innovation (i.e., the number of state changes) at corresponding

branches across gymnosperm phylogeny (*n* = 119). The x-axis represents time in millions of

years. Nodes with an inferred diversification shift are marked with an asterisk.

Figure 3. **Gene-tree conflict vs. phenotypic rates**. Plots juxtaposing rates of phenotypic

evolution and levels of gene-tree conflict at corresponding branches across gymnosperm

phylogeny (*n* = 119). Note that the conflict node highlighted as corresponding with the *Juniperus*

rate node is its immediate parent, comprising *Cupressus sempervirens*, *Juniperus*. The x-axis

represents time in millions of years.

Figure 4. **Climatic and diversification shifts across gymnosperms**. Ancestral reconstruction of

annual precipitation (Bio 12) on the supermatrix phylogeny, showing diversification rate shifts

(black circles) and shifts in rates of climatic evolution (Bio 1, mean annual temperature = red

circle; Bio 12, annual precipitation = cyan circle) inferred using BAMM. The diversification

shifts marked with asterisks (corresponding to *Macrozamia* and *Zamia*) show concerted rate

shifts with PC2 of the climatic variables, bringing the number of directly concerted

diversification and climate shifts to ten. See Supplementary Fig. 9 for species tip labels.

Extended Data Figure 1. **Plots of synonymous substitutions per site (*Ks*) for within-taxon paralog pairs (black lines) and between-taxon ortholog pairs (blue lines).** The relative positions of ortholog vs. paralog *Ks* spikes help clarify the phylogenetic positions of possible WGD events. The seed plant WGD event (GINK-α) is labeled. *Ks* peaks corresponding to an inferred WGD for gymnosperms are highlighted with an asterisk. The taxa compared capture the root nodes of (A, B) seed plants and gymnosperms, (C) the 'ginkad' clade, (D) conifers, (E) the cupressophyte clade, and (F) the Taxaceae-Cupressaceae clade. Ortholog and paralog *Ks* plots were generated using the pipelines of Walker et al.[83] and Yang et al.[28], respectively.

Extended Data Figure 2. **Plots of synonymous substitutions per site (*Ks*) for within-taxon**

paralog pairs of representatives of the 'ginkad' clade. The seed plant WGD event (GINK-α)

is labeled. *Ks* peaks corresponding to an inferred WGD for gymnosperms are highlighted with an

asterisk.

Extended Data Figure 3. **Genome size evolution in gymnosperms**. Ancestral reconstruction of

genome size (C-value) on the pruned supermatrix phylogeny, showing BAMM rate shifts (red

circles) and jumps (i.e., extreme differences in ancestor-descendent values; white circles) in

genome size evolution, as well as BAMM diversification shifts (larger black circles).

Supplementary Fig. 15 shows this figure with species tip labels.

Methods

Sampling overview

We generated two major phylogenetic datasets. The first is a transcriptomic and genomic dataset comprising 144 species (including 121 ingroup species, representing 83 of the 85 gymnosperm genera recognized by the Kew checklist: https://wcsp.science.kew.org/home.do); this was used to resolve broad-scale phylogenetic relationships, characterize patterns of phylogenomic conflict, and infer hotspots of gene duplication. Representatives of angiosperms, monilophytes, and lycophytes were used as outgroups. In all, 77 transcriptomes were newly generated for this study; the remaining transcriptomes were obtained from the One Thousand Plants (1KP) Consortium (https://sites.google.com/a/ualberta.ca/onekp) and several genomes were obtained from the Phytozome database v9.1[60]. See Supplementary Table 1 for transcriptomic and genomic sampling details. The two missing gymnosperm genera, *Pectinopitys* and *Pherosphaera*, represent genera nested within Podocarpaceae and thus their absence should not impact the resolution of major relationships. The newly generated transcriptomes were deposited in the Sequence Read Archive (BioProject PRJNA726756). We also generated a phenotypic dataset of 148 traits for the seed plant species in the transcriptomic dataset to permit examination of the correspondence of different aspects of genomic and phenotypic evolution; the trait dataset is described below.

The other phylogenetic dataset is a plastid supermatrix, based on publicly available plastomes and plastid genes in NCBI (data deposit), as well as 32 plastid genomes newly generated through genome skimming (Supplementary Table 2). In total, this dataset includes 78 plastid genes across 1055 accessions, including 995 ingroup accessions. The ingroup names were checked against the Kew gymnosperm checklist (https://wcsp.science.kew.org/home.do) to resolve issues with synonymy. In total, 890 gymnosperm species are represented, comprising ca. 82% of the 1090 recognized gymnosperm species[56]; our sampling also includes numerous infraspecific taxa (i.e., varieties and subspecies; https://wcsp.science.kew.org/home.do).

The supermatrix dataset was used to generate a comprehensive dated phylogeny for gymnosperms for reconstructions of diversification shifts, climatic evolution, genome size evolution, and chromosome evolution. We used exclusively plastid data for two reasons: (1) plastid genes are more widely sequenced across gymnosperms, allowing for greater species representation; and (2) conflict within phylogenetic datasets can greatly (and negatively) impact

branch length estimation and downstream analyses such as divergence-time estimation[61]. While

studies show that the plastome is not without conflict[62], it clearly shows more minimal levels of

conflict compared to nuclear genes. Plastid relationships of gymnosperms, as shown by previous

studies[56,63] and our results here, largely agree with those of the nuclear genome, and thus a plastid

supermatrix tree should accurately capture major macroevolutionary patterns.

Transcriptome and plastome sequencing and assembly

To generate the new transcriptomes, total RNAs were isolated from young buds or leaves using

the Spectrum™ Plant Total RNA Kit (SIGMA-ALDRICH). Approximately 5 ug of total RNAs

were used to construct cDNA libraries (NEBNext Ultra Directional RNA Library Prep Kit for

Illumina, Illumina, San Diego, CA, USA). Transcriptome sequencing was performed using the

Illumina Hi-Seq 2000 platform. Paired-end reads of 150 bp were *de novo* assembled with Trinity

v2.4.0 (Grabherr et al., 2011) using default parameters. Transcriptomes from the 1KP project

were previously assembled (Wickett 2014; Leebens-Mack et al. 2019) using SOAPdenovo v1 (Li

et al. 2010), and amino acid sequences for three gymnosperm species derived from genome

annotations were downloaded from the Phytozome database v9.1[60]. All transcripts were

translated using TransDecoder v5.3.0[64], and then cd-hit-est[65] (-c 0.99 –n 5) was used to reduce

sequence redundancy in all CDS.

Plastid genomes were sequenced and assembled as follows. Total genomic DNA was

extracted from fresh leaves using the CTAB method[66]. Approximately one microgram of total

genomic DNA from each sample was used for ~400-bp library construction with the NEBNext

DNA Library Prep Kit (New England Biolabs, Ipswich, MA, USA) and Covaris-based

fragmentation. The Illumina MiSeq instrument (Illumina, San Diego, CA, USA) was used to

carry out 150-bp, paired-end sequencing. We assembled the plastomes using both the

GetOrganelle pipline[67] and SPAdes v3.13.0[68], the latter with the "careful"-option and kmers of

61, 81, 101, and 121. In order to validate the plastome assembly, we mapped all paired-end reads

to the consensus sequence of the assembled plastomes with the local-sensitive option of Bowtie

v2.3.2[69]. Plastid Genome Annotator[70] was used to perform plastome annotation, coupled with

manual correction in Geneious v8.0.2 (https://www.geneious.com/).

Phylogenomic workflow

Homology and orthology inference. We followed two major procedures for homology and

orthology inference in the transcriptomic dataset. First, we employed the hierarchical homology

inference approach of Walker et al.[71]. Given the size of our transcriptomic dataset (including 144

transcriptomes), performing an all-by-all BLAST search across all samples for initial homology

assessment would have been computationally prohibitive. The Walker et al.[71] approach

overcomes this problem by conducting all-by-all BLAST searches on subsets of the data (*tip*

clustering), which are then combined using a post-order (tip to root) tree traversal method (*node*

clustering). We divided our samples into 11 subgroups for tip clustering: (**1**) monilophytes and

lycophytes (7 samples), (**2**) angiosperms (16 samples), (**3**) cycads and *Ginkgo* (14 samples), (**4**)

Gnetales (5 samples), (**5**) Pinaceae (22 samples), (**6**) Podocarpaceae (23 samples), (**7**)

Araucariaceae (6 samples), (**8**) *Sciadopitys* and Taxaceae (12 samples), (**9**) 'basal' Cupressaceae

(11 samples), (**10**) cupressoid Cupressaceae (17 samples), and (**11**) callitroid Cupressaceae (11

samples); see Supplementary Table 1 for lists of species included in each grouping. For tip

clustering, for each subset, we conducted an all-by-all BLASTP search[72] with an e-value cutoff

of 10. The tip clusters were then sequentially combined in a post-order fashion (node-clustering)

using the scripts developed by Walker et al.[71] (https://github.com/jfwalker/Clustering).

For homolog tree and orthology inference from the combined clusters, we followed the

Yang and Smith[73] pipeline (scripts available at

https://bitbucket.org/yangya/phylogenomic_dataset_construction/src/master/), which we briefly

outline here. Fasta files were written from each cluster. The clusters were then aligned using

MAFFT v7[74] and cleaned using the phyx[75] program 'pxclsq' (retaining alignment columns with

at least 10% occupancy), after which homolog trees were built using FastTree v2 with the WAG

model[76]. The trees were then pruned of extreme long branches, monophyletic tips were masked

(i.e., only one sequence was retained of species/sample-specific alleles/paralogs/duplications),

and then long internal branches (representing deep paralogs) were pruned. Fasta files were then

written again from the resulting subtrees, and two more iterations of alignment, homolog tree

construction, and homolog tree pruning were conducted. A final stage of homolog tree

construction was then conducted using RAxML v8.2.12[77] to generate more robust homolog trees.

The resulting 1,992 homolog trees were then used for orthology inference and gene-duplication

mapping. For orthology inference, we used the 'RT' method of Yang and Smith[73], which extracts

ingroup clades and then cuts paralogs from root to tip. For this procedure, the three lycophyte

samples were designated as outgroups (and thus these were pruned and excluded from

subsequent analyses), and the minimum ingroup sampling was set at 100. This approach resulted

in a set of 790 orthologs. Nucleotide fasta files were then written from each extracted ortholog,

re-aligned using MAFFT, and then cleaned by removing alignment columns with under 25%

occupancy using 'pxclsq' in phyx.

Coalescent species-tree inference. Gene trees (from the 790 orthologs inferred above) were

constructed using RAxML, using the GTRGAMMA model and 100 bootstrap replicates. The

gene trees were then used for species-tree analysis in ASTRAL-III[78]. Prior to the ASTRAL

analysis, low supported branches (<10 BS) in the gene trees were collapsed using Newick

utilities[79], as this has been shown to increase accuracy[78]. ASTRAL was then run under default

settings, producing a 'species tree' tree with local posterior probability (LPP) support values for

the bipartitions and branch lengths in coalescent units (Supplementary Fig. 16). The resulting

phylogeny was used as a framework for various subsequent analyses, including conflict analysis,

analysis of gene duplications, inference of whole-genome duplications, morphological trait

reconstructions, and comparisons of phenotypic innovation and rates of phenotypic evolution

with levels of genomic conflict and gene duplications.

Concatenation species-tree inference. The individual gene/ortholog alignments were

concatenated using the phyx program 'pxcat'. We then inferred phylogenies from the

concatenated nucleotide alignment (comprising 1,444,037 bp) using IQ-TREE[80] with the

GTRGAMMA model and ultrafast bootstrapping (1000 replicates). Two analyses were

conducted: one unpartitioned, and the other with separate model parameters estimated for each

gene region[81] (Supplementary Fig. 17).

Conflict analysis. To examine patterns of gene-tree conflict across gymnosperm phylogeny, we

used the program PhyParts[82], which maps gene trees onto an input species tree and summarizes

the number of gene trees in concordance or conflict with each bipartition. When gene trees have

inadequate species sampling or bootstrap support (if a support threshold is specified) to speak to

a particular bipartition, those are considered uninformative. We use the python script

'phypartspiecharts.py'

(https://github.com/mossmatters/phyloscripts/tree/master/phypartspiecharts) to visualize the

PhyParts results, which are shown in Fig. 1. The levels of conflict calculated for each bipartition

were also plotted against corresponding levels of morphological innovation (i.e., the number of

new states appearing at each node; see below). We also used the program 'minority_report.py'

(https://github.com/mossmatters/phyloscripts/tree/master/minorityreport) to examine the number

of gene trees supporting alternative placements for particular lineages (namely, *Ginkgo*).

Inference of whole-genome duplication

We used a multi-faceted approach to examine genomic changes and infer putative WGDs across

gymnosperm phylogeny, employing gene-duplication mapping, plots of synonymous

substitutions per site (*Ks*), and reconstructions of chromosome number and genome size (C-

value). Gene duplications were mapped onto the species tree topology using the scripts

(https://bitbucket.org/blackrim/clustering/src/master/) and method of Yang et al.[28] This involved

extracting rooted orthogroups from the homolog trees (with lycophytes designated as the

outgroup), which were then mapped onto the species tree topology. We used two methods of

mapping: one maps duplications to the most recent common ancestor (MRCA) in the species

tree, with a bootstrap filter of 50% (Supplementary Fig. 18); the other imposes a topology filter

such that duplications are only mapped when the sister clade of the gene duplication node

contains at least a subset of the taxa present in that clade in the species tree[28] (Supplementary

Fig. 19).

Ks plots of within-taxon paralog pairs were generated for the newly sequenced

transcriptomes. First, CD-HIT (-c 0.99 –n 5) was used to reduce highly similar sequences, after

which an all-by-all BLASTP search was carried out within each taxon (e-value cutoff = 10). The

results were then filtered such that sequences with ten or more hits were removed (to avoid

overrepresentation of large gene families), as were hits with pident/nident less than 20/50%. *Ks*

values were then calculated from the CDS of the remaining paralog pairs, using the pipeline

https://github.com/tanghaibao/bio-pipeline/tree/master/synonymous_calculation, and plotted

using scripts from Yang et al.[28] We used between-taxon ortholog pairs to generate ortholog *Ks*

plots following the method of Walker et al.[83]; relative comparisons of ortholog and paralog *Ks*

peaks are useful for determining whether observed peaks are shared (occurred prior to

ortholog/lineage divergence) or more species- or clade-specific (occurred after ortholog/lineage

divergence). The ortholog and paralog *Ks* plots (Extended Data Fig. 1 and 2; data deposit) were

visually inspected to identify clade-specific shared peaks, which were compared against the

results from gene-duplication mapping, as well as the reconstructions of chromosome number

and genome size (outlined below). We also generated within-species paralog *Ks* plots using the

pipeline of Scott et al.[33] ([https://github.com/nstenz/plot-ks)](https://github.com/nstenz/plot-ks) and these are also provided in the

data deposit.

Supermatrix tree construction and dating

To assemble a comprehensive species-level phylogeny of gymnosperms, we mined plastid gene

and genome sequences available on NCBI and combined these with genes extracted from the 32

newly sequenced and assembled plastomes. While the plastomes from NCBI were directly

downloaded, individual plastid genes were mined from NCBI using the program PyPhlawd[84].

Large-scale plastid gene clusters obtained using PyPhlawd were then combined with the

corresponding individual gene sequences derived from the plastomes.

The plastid genes were aligned individually using MAFFT v7[74], after which alignment

columns with under 25% occupancy were removed using the phyx function 'pxclsq'. The

alignments were then concatenated using the phyx function 'pxcat'. The final concatenated

matrix included 20.35% matrix occupancy (i.e., 79.65% missing data). Prior to phylogenetic

analysis, PartitionFinder v2[85] was used to determine the optimal model partitioning strategy for

the concatenated alignment. Phylogenetic analyses were conducted using RAxML-ng[86] with a

topological constraint enforcing major relationships to match the transcriptomic species tree

topology: ((*Ginkgo*,cycads),((Gnetales,Pinaceae),((Araucariaceae,Podocarpaceae),

(Sciadopityaceae, (Taxaceae,Cupressaceae))))). When unconstrained, the plastome tree only

differed from the constraint in the placement of Gnetales. Using RAxML-ng, we conducted 25

independent searches for the maximum likelihood tree ("raxml-ng --search1"), from which the

tree with the best likelihood score was selected for downstream analyses of diversification,

climate evolution, and genome evolution (chromosome number and cval), although all 25 trees

were dated in treePL[87] to obtain confidence intervals on node ages (see below). We also inferred

100 bootstrap trees to evaluate topological support for the supermatrix phylogeny.

Divergence dating. Given the scale of the supermatrix tree (1055 terminals), typical methods of

Bayesian divergence dating were unfeasible. Therefore, we used treePL[87] to generate a dated

phylogeny. We used extensive fossil evidence to inform our calibration scheme (see

Supplementary Information for detailed discussion of the fossils used, their justification, and

references). We conducted independent analyses on the 25 inferred ML trees using the same set

of calibrations as a means of obtaining confidence intervals for inferred node ages

(Supplementary Fig. 20). Cross-validation was performed for each analysis to determine the

optimal smoothing parameter. Node ages from the dated supermatrix tree were then applied as

fixed ages (also using treePL) to the corresponding nodes in the transcriptomic species tree. This

allowed us to place subsequent analyses involving the transcriptomic phylogeny (i.e.,

examinations of conflict and rates of morphological evolution) in a temporal context.

Given uncertainty in the placements of two calibrations (for the *Acmopyle* stem and the

Keteleeria stem; see Supplementary Methods), we also conducted a dating analysis on the best

ML tree excluding these calibrations to assess their impact on the ages inferred for major

relevant clades (i.e., Pinaceae, Podocarpaceae). The resulting ages for these clades fell within the

confidence intervals defined by the analyses of the 25 ML trees noted above (Supplementary Fig.

20), and therefore we proceeded with our initial analyses including these calibrations.

Chromosome number and genome size reconstructions

Chromosome counts were obtained from the Chromosome Count Database[49]

(http://ccdb.tau.ac.il; accessed 9 December 2019), and the parsed (haploid) counts were used for

subsequent analysis. In cases where several different counts were recorded for a given species,

the most frequently occurring count was used. Measurements of genome size (i.e., C-values, the

amount of DNA in the haploid genome) were obtained from the Plant DNA C-values Database[50]

(https://cvalues.science.kew.org; accessed 9 December 2019). The dated supermatrix phylogeny

was pruned to match the sampling in each dataset (chromosome counts and C-values), and these

two trees were used in the analyses outlined below. The chromosome count tree/dataset included

478 species, and the cval tree/dataset include 373 species. The chromosome count and C-value

datasets examined are available in the data deposit.

Reconstructions of ancestral chromosome number were implemented in ChromEvol

v2.0[88]. We examined the data under eight different available models, each including different

combinations of the rate parameters *gain* (ascending dysploidy), *loss* (descending dysploidy),

duplication (polyploidization), and *demi-duplication* (multiplication of the chromosome number

by a factor of 1.5), with the rates for each treated either as constant or linear. The model with the

lowest Akaike information criterion (AIC) score (i.e., 'LINEAR RATE DEMI EST') was used in

a more thorough analysis involving default settings and 100,000 simulations. We then visualized

changes in chromosome number across the tree using FigTree (Supplementary Fig. 2) and

compared changes against reconstructions of genome size and gene duplication.

Ancestral reconstructions of genome size (using C-values) were implemented in R[89] (R

Core Team 2020) using the 'fastAnc' function in phytools[90]. The C-value data were not

transformed prior to analysis as they conform to a more-or-less normal distribution. To identify

the most extreme changes in genome size (which we call C-value 'jumps'), we then calculated

the difference between each node's reconstructed value and that of its immediate parent node,

and selected branches in the 95th percentile of the distribution of parent-child C-value

differences. These are shown in Extended Data Fig. 3. We also examined rates of genome size

evolution using BAMM[91,92], using the same tree as above. The priors for the BAMM analysis

were determined using the function 'setBAMMpriors' in the BAMMtools package[93]. The

analysis was run for one billion generations using the phenotypic evolution model, with

convergence assessed using BAMMtools. Rate shifts from the maximum a posteriori probability

(MAP) shift configuration are shown in Extended Data Fig. 3.

Species diversification analysis

We used BAMM to detect diversification rate shifts across gymnosperm phylogeny, using the

dated supermatrix tree as input. The analysis employed the speciation-extinction model, with the

priors determined using the function 'setBAMMpriors' in BAMMtools. We also specified

sampling fractions specified for each family of *Acrogymnospermae*. This analysis was initially

run for 1 billion generations, but then was re-run for 2 billion generations in order to achieve

convergence. Rate shifts from the MAP shift configuration are shown in Fig. 1, 4, and

Supplementary Fig. 3.

Reconstructions of climatic occupancy evolution

Geographic occurrences of gymnosperms were pulled from GBIF (https://www.gbif.org;

accessed 15 May 2020), and then compared against occurrences from iDigBio

(https://www.idigbio.org/portal/search) to check for available data of the species missing from

GBIF. Occurrences were then cleaned using the R package CoordinateCleaner[94], removing

duplicate records as well as those with problematic coordinates (e.g., non-terrestrial, identical

lat/lon, located within major cities or biodiversity institutions). Synonymy was reconciled using

data from the Kew checklist (https://wcsp.science.kew.org/home.do). The occurrences were then

used to extract climatic values from the 19 WorldClim bioclimatic variables, and means were

calculated for each species for each climatic variable. We dropped tips from the dated

supermatrix tree that were absent from the climatic dataset sampling, resulting in a matched

sampling of 908 species for subsequent climatic analyses. We then conducted a principal

component analysis in R, and principal components (PC) 1 and 2 were used in subsequent

analyses of climatic evolution. We also conducted analyses of mean annual temperature (Bio1)

and annual precipitation (Bio12), using the mean values for each species.

As with the genome size analyses, for the climate analysis we reconstructed major jumps

and rate shifts in climatic evolution, involving separate analyses for PC1, PC2, Bio1, and Bio12.

Jumps were determined by reconstructing ancestral climatic values (using the 'fastAnc' function

in phytools), calculating the difference between each node and its immediate parent, and

identifying those branches in the 95th percentile. Extreme climatic jumps for PC1, PC2, Bio1,

and Bio12 are shown in Supplementary Figs. 13 and 14. BAMM analysis of rates of climatic

occupancy evolution were also conducted on PC1, PC2, Bio1, and Bio12, with each run for 1

billion generations, employing the phenotypic evolution model, with priors determined using

BAMMtools. Rate shifts from the MAP shift configuration for Bio1 and Bio 12 are shown in

Fig. 4 and Supplementary Figs. 9 and 10; those of PC1 and PC2 are shown in Supplementary

Figs. 11 and 12.

Although analysis of historical climatic information would be ideal for understanding the

climatic context of previous diversification shifts in greater detail, this absence of reliable

estimates spanning the depth of gymnosperm evolutionary history makes this, at the present, not

feasible given the scope of our study. Thus, we only used current global climate data for our

reconstructions. However, the use of historical climate data to investigate particular Cenozoic

radiations of gymnosperms would be a promising avenue for future research.

Morphological reconstructions and genomic comparisons

A phenotypic dataset comprising 148 traits across the 137 seed plant species included in the

transcriptomic dataset was assembled to examine the correspondence of phenotypic and genomic

evolution (matrix available in the data deposit). Many of the traits examined were also included

in previous morphological datasets. However, the taxonomic sampling in some of these previous

datasets was limited; furthermore, many traits from clade-specific datasets needed to be modified

and reconciled with similar traits in other datasets in order to establish characters applicable

across all sampled taxa. The traits examined (and their corresponding states) are listed in

Supplementary Information, and we indicate how these relate to traits in previously published

datasets. The primary source datasets were those of Hart[95], Hilton and Bateman[96], Mao et al.[97],

Escapa and Catalano[98], Coiro and Pott[99], Herrera et al.[100,101], Gernandt et al.[102], and Andruchow-

Colombo et al.[103] See Supplementary Information for a complete list of references from which

phenotypic data were obtained. We note that while determining homology for character

delimitation in seed plants is a major challenge[104], we do not believe that instances of

misidentified homology (i.e., treating independently evolved structures as a single trait or

expression of a trait) should necessarily undermine our goals here, given that we are not using

this trait set to reconstruct phylogeny. Our goal is to identify state changes along branches, and

even in instances of conflated homology, e.g., with two distantly related lineages scored with the

same state for a particular trait despite independent origins of the relevant structures, a state

change should nevertheless be 'correctly' reconstructed and recorded for each branch.

Morphological reconstructions on the species-tree were performed using Maximum

Parsimony, following the procedure of Parins-Fukuchi et al.[13], briefly outlined here. Each trait

was reconstructed on the phylogeny, and then the number of state changes along each branch

was tallied—this is considered the degree or level of phenotypic 'innovation' or 'novelty' along

each branch. To determine rates of phenotypic evolution for each branch, we divided the number

of state changes by the duration of branch (in million years). We then plotted levels of

morphological innovation and rates of phenotypic evolution for each branch/node (with the x-

axis representing time or node age), and compared these against plots of the same nodes showing

their levels of gene duplication and gene-tree conflict (Figs. 2 and 3), allowing for visual

inspection of these patterns and their correspondence through time. Because Gnetales was

detected as a major outlier, with extreme levels of morphological innovation perhaps driven by

life history shifts, we also conducted morphological reconstructions excluding Gnetales to ensure

that the inclusion/placement of Gnetales was not significantly biasing reconstructions for other

near-by nodes (Supplementary Fig. 6).

On the basis of the visual correspondences observed from the plots noted above, we then

explicitly tested the relationships of (1) phenotypic innovation vs. gene duplication and (2)

phenotypic rates vs. gene tree conflict by conducting linear regression and permutation analyses

of each pair of variables. Linear regression was performed using the 'lm' function in R, treating

phenotypic innovation and rate as the dependent variables in the respective analyses. For the

examination of innovation vs. duplication, we conducted an analysis including all nodes

(Supplementary Fig. 21), as well as one excluding two major outlier nodes: Gnetales and

conifers (Supplementary Fig. 6). As noted in the main text, we believe exclusion of these outliers

is warranted given that (1) Gnetales has experienced major changes in genome size (i.e.,

reduction) and life history with generally elevated rates of molecular evolution that might erase

signatures of a relationship between gene duplication and morphological innovation, and (2) trait

reconstruction for conifers is complicated by its inclusion of Gnetales (a highly derived lineage)

and the prevalence of features shared among conifers and *Ginkgo*, which may have been

involved in a reticulation event early in the evolution of the conifer lineage based on patterns of

gene-tree conflict.

The permutation test, developed by Parins-Fukuchi et al.[13], identifies nodes with

significantly elevated values for a particular variable by resampling (with replacement) observed

values 1000 times and using the means from each replicate to create an empirical null

distribution. We executed this procedure for each of the four variables considered above

(duplication levels, conflict levels, innovation levels, and phenotypic rate), with nodes/values

falling in the 95th percentile deemed significant. Then, to examine the strength of the association

between (1) innovations and duplications and (2) rates and conflict, we determined whether the

number of nodes with significantly high values for both of the variables in question (e.g.,

innovations and duplications) departed from the expectation if the same number of nodes was

randomly sampled (Supplementary Figs. 5 and 8). This permutation test was designed

specifically to examine whether extreme/significant values (in, for example, conflict and rates)

tend to coincide, even if there is not a strict positive monotonic relationship between the two

variables across the spectrum of values. Finally, we also used this test to determine if

diversification shifts tend to co-occur with significant/extreme levels of phenotypic innovation

(Supplementary Fig. 22).

Data availability

The newly generated raw sequence data is available at the NCBI Sequence Read Archive

(https://www.ncbi.nlm.nih.gov/sra) under BioProject PRJNA726756 (transcriptomic samples)

and PRJNA726638 (genome skimming samples). The newly assembled plastid genomes are also

available at NCBI (https://www.ncbi.nlm.nih.gov); see Supplementary Table 2 for sample

accession numbers. Sequence alignments, phylogenies, *Ks* plots, phenotypic trait data, and other

data analyzed (chromosome counts, c-values) are available on figshare

(10.6084/m9.figshare.14547354).
956

60. Goodstein, D. M. et al. Phytozome: a comparative platform for green plant genomics.

Nucleic Acids Res. **40**, 1178–1186 (2012).

61. Mendes, F. K. & Hahn, M. W. Gene tree discordance causes apparent substitution rate

variation. *Syst. Biol.* **65**, 711–721 (2016).

62. Walker, J. F., Walker-Hale, N., Vargas, O. M., Larson, D. A. & Stull, G. W. Characterizing

gene tree conflict in plastome-inferred phylogenies. *PeerJ* **7**, e7747 (2019).

63. Zhong, B., Yonezawa, T., Zhong, Y., Hasegawa, M. The position of Gnetales among seed

plants: overcoming pitfalls of chloroplast phylogenomics. *Mol. Biol. Evol.* **27**, 2855–2863

(2010).

64. Haas, B.J. et al. De novo transcript sequence reconstruction from RNA-seq using the Trinity

platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494–1512 (2013).

65. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W., 2012. CD-HIT: accelerated for clustering the next-

generation sequencing data. *Bioinformatics* **28**, 3150-3152 (2012).

66. Doyle, J. J. & Doyle, J. L. A rapid DNA isolation procedure for small quantities for fresh leaf

tissue. *Phytochemical Bulletin* **19**, 11–15 (1987).

67. Jin, J-J. et al. GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of

organelle genomes. *Genome Biol.* **21**, 241 (2020).

68. Bankevich, A. et al. SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

69. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357 (2012).

70. Qu, X. J., Moore, M. J., Li, D. Z. & Yi, T. S. PGA: a software package for rapid, accurate,
and flexible batch annotation of plastomes. *Plant Methods* **15**, 50 (2019).

71. Walker, J. F. et al. From cacti to carnivores: improved phylotranscriptomic sampling and
hierarchical homology inference provide further insights into the evolution of Caryophyllales.
Am. J. Bot. **105**, 446–462 (2018).

72. Altschul, S. F. et al. Gapped BLAST and PSI-BLAST: a new generation of protein database
search programs. *Nucleic Acids Res.* **25**, 3389–3402 (1997).

73. Yang, Y. & Smith, S. A. Orthology inference in nonmodel organisms using transcriptomes
and low-coverage genomes: improving accuracy and matrix occupancy for
phylogenomics. *Mol. Biol. Evol.* **31**, 3081–3092 (2014).

74. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

75. Brown, J. W., Walker, J. F. & Smith, S. A. Phyx: phylogenetic tools for
unix. *Bioinformatics* **33**, 1886–1888 (2017).

76. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately Maximum-Likelihood
Trees for Large Alignments. *PLoS One* **5**, e9490 (2010).

77. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

78. Zhang, C. et al. ASTRAL-III: polynomial time species tree reconstruction from partially
resolved gene trees. *BMC Bioinformatics* **19**, 153 (2018).

79. Junier, T. & Zdobnov, E. M. The Newick utilities: high-throughput phylogenetic tree
processing in the UNIX shell. *Bioinformatics* **26**, 1669–1670 (2010).

80. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum likelihood phylogenies. *Mol. Biol.*
Evol. **32**, 268–274 (2015).

81. Chernomor, O., von Haeseler, A. & Minh, B. Q. Terrace aware data structure for
phylogenomic inference from supermatrices. *Syst. Biol.* **65**, 997–1008 (2016).

82. Smith, S. A., Moore, M. J., Brown, J. W. & Yang, Y. Analysis of phylogenomic datasets
reveals conflict, concordance, and gene duplications with examples from animals and
plants. *BMC Evol. Biol.* **15**, 150 (2015).
83. Walker, J. F., et al. Widespread paleopolyploidy, gene tree conflict, and recalcitrant
relationships among the carnivorous Caryophyllales. *Am. J. Bot.* **104**, 858–867 (2017).
84. Smith, S. A. & Walker, J. F. Pyphlawd: a python tool for phylogenetic dataset
construction. *Methods Ecol. Evol.* **10**, 104–108 (2019).
85. Lanfear, R., Frandsen, P.B., Wright, A.M., Senfeld, T. & Calcott, B. PartitionFinder 2: new
methods for selecting partitioned models of evolution for molecular and morphological
phylogenetic analyses. *Mol. Biol. Evol.* **34**, 772–773 (2017).
86. Kozlov, A.M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: a fast,
scalable and user-friendly tool for maximum likelihood phylogenetic
inference. *Bioinformatics* **35**, 4453–4455 (2019).
87. Smith, S. A. & O'Meara, B. C. treePL: divergence time estimation using penalized likelihood
for large phylogenies. *Bioinformatics* **28**, 2689–2690 (2012).
88. Glick, L. & Mayrose, I. ChromEvol: assessing the pattern of chromosome number Evolution
and the inference of polyploidy along a phylogeny. *Mol. Biol. Evol.* **31**, 1914–1922 (2014).
89. R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org (2020).
90. Revell, L. J. Phytools: an R package for phylogenetic comparative biology (and other
things). *Methods Ecol. Evol.* **3**, 217–223 (2012).
91. Rabosky, D. L. et al. Rates of speciation and morphological evolution are correlated across
the largest vertebrate radiation. *Nat. Commun.* **4**, 1–8 (2013).
92. Rabosky, D. L. Automatic detection of key innovations, rate shifts, and diversity-dependence
on phylogenetic trees. *PLoS One* **9**, e89543 (2014).
93. Rabosky, D. L. et al. BAMM tools: an R package for the analysis of evolutionary dynamics
on phylogenetic trees. *Methods Ecol. Evol.* **5**, 701–707 (2014).
94. Zizka, A. et al. CoordinateCleaner: standardized cleaning of occurrence records from
biological collection databases. *Methods Ecol. Evol.* **10**, 744–751 (2019).
95. Hart, J. A. A cladistics analysis of conifers: preliminary results. *J. Arnold Arb.* **68**, 269–307
(1987).

96. Hilton, J. & Bateman, R. M. Pteridosperms are the backbone of seed-plant phylogeny. *J.*

Torrey Bot. Soc. **133**, 119–168 (2006).

97. Mao, K. et al. Distribution of living Cupressaceae reflects the breakup of Pangea. *Proc. Natl*

Acad. Sci. USA **109**, 7793–7798 (2012).

98. Escapa, I. H. & Catalano, S. A. Phylogenetic analysis of Araucariaceae: integrating

molecules, morphology, and fossils. *Int. J. Plant Sci.* **174**, 1153–1170 (2013).

99. Coiro, M. & Pott, C. *Eobowenia* gen. nov. from the Early Cretaceous of Patagonia: indication

for an early divergence of Bowenia? *BMC Evol. Biol.* **17**, 97 (2017).

100. Herrera, F. et al. Cupressaceae conifers from the Early Cretaceous of Mongolia. *Int. J. Plant*

Sci. **178**, 19–41 (2017).

101. Herrera, F. et al. Reconstructing *Krassilovia mongolica* supports recognition of a new and

unusual group of Mesozoic conifers. *PLoS One* **15**, e0226779 (2020).

102. Gernandt, D. S. et al. Incorporating fossils into the Pinaceae tree of life. *Am. J. Bot.* **105**,

1329–1344 (2018).

103. Andruchow-Colombo, A., Wilf, P. & Escapa, I. H. A South American fossil relative of

Phyllocladus: *Huncocladus laubenfelsii* gen. et sp. nov. (Podocarpaceae), from the early

Eocene of Laguna del Hunco, Patagonia, Argentina. *Aust. Syst. Bot.* **32**, 290–309 (2019).

104. Nixon, K. C., Crepet, W. L., Stevenson, D. & Friis, E. M. A reevaluation of seed plant

phylogeny. *Ann. Missouri Bot. Gard.* **81**: 484–533 (1994).