Ecological correlates of gene family size: the draft genome of the redheaded pine sawfly *Neodiprion lecontei*

Kim L. Vertacnik[1,2], Danielle K. Herrig[1], R. Keating Godfrey[3], Tom Hill[4,5], Scott M. Geib[6], Robert L. Unckless[4], David R. Nelson[7], and Catherine R. Linnen[1]

[1]Department of Biology, University of Kentucky, Lexington, KY 40506, USA
[2]Columbia River Inter-Tribal Fish Commission, Hagerman, ID 83332, USA (current address)
[3]Department of Neuroscience, University of Arizona, Tucson, AZ 85721, USA
[4]Department of Molecular Biosciences, University of Kansas, Lawrence, Kansas 66045, USA
[5]NIH Frederick National Laboratory for Cancer Research, Frederick, MD 21702, USA
[6]Tropical Crop and Commodity Protection Research Unit, United States Department of Agriculture: Agriculture Research Service Pacific Basin Agricultural Research Center, Hilo, Hawaii 96720, USA
[7]Department of Microbiology, Immunology and Biochemistry, University of Tennessee Health Science Center, Memphis, TN 38163, USA

Correspondence: kvertacnik@critfc.org

Keywords: evolutionary predictability, adaptation, chemosensation, detoxification, immunity, Hymenoptera

Abstract

A central goal in evolutionary biology is to determine the predictability of adaptive genetic changes. Despite many documented cases of convergent evolution at individual loci, little is known about the repeatability of gene family expansions and contractions. To address this void, we examined gene family evolution in the redheaded pine sawfly *Neodiprion lecontei*, a non-eusocial hymenopteran and exemplar of a pine-specialized lineage evolved from angiosperm-feeding ancestors. After assembling and annotating a draft genome, we manually annotated multiple gene families with chemosensory, detoxification, or immunity functions and characterized their genomic distributions and evolutionary history. Our results suggest that expansions of bitter gustatory receptor (GR), clan 3 cytochrome P450 (CYP3), and antimicrobial peptide (AMP) subfamilies may have contributed to pine adaptation. By contrast, there was no evidence of recent gene family contraction via pseudogenization. Next, we compared the number of genes in these same families across insect taxa that vary in diet, dietary specialization, and social behavior. In Hymenoptera, herbivory was associated with large GR and small olfactory receptor (OR) families, eusociality was associated with large OR and small AMP families, and—unlike investigations among more closely related taxa—ecological specialization was not related to gene family size. Overall, our results suggest that gene families that mediate ecological interactions may expand and contract predictably in response to particular selection pressures, however, the ecological drivers and temporal pace of gene gain and loss likely varies considerably across gene families.

Introduction

Changes in gene family size are a potentially important source of evolutionary innovation. When gene families grow via duplication, for example, reduced functional constraints may facilitate the development of phenotypic novelty (Ohno 1970; Demuth and Hahn 2009). Reductions in gene family size can also give rise to novel traits. For example, rampant pseudogenization in particular gene families is linked to the colonization of highly specialized niches like oligotrophic caves (Protas et al. 2006; Gross et al. 2009; Yang et al. 2016) and toxic host plants (Matsuo et al. 2007; McBride 2007; Good et al. 2014). Together, these observations suggest that gene families predictably expand or contract in response to specific selection pressures. Yet compared to the rich and growing literature on genetic convergence at individual loci (Martin and Orgogozo 2013), the repeatability and predictability of gene family evolution remains understudied.

The evolution of many gene families, defined here as groups of genes that share sequence and functional similarity from common ancestry (Dayhoff 1976; Demuth and Hahn 2009), is consistent with a birth-death model in which genes arise via duplication (gene gain) and are lost via deletion or pseudogenization (gene loss) (Hughes and Nei 1992; Nei and Rooney 2005). When the frequencies of gene duplications and deletions evolve primarily via genetic drift, gene families contract and expand over time via a process dubbed genomic drift (Nei 2007; Nozawa et al. 2007). Overall, this stochastic birth-death process (which differs from Nei's conceptual birth-death model of gene family evolution (Hahn et al. 2005)) sufficiently explains most gene family size distributions within genomes (Karev et al. 2002) and between species (Hahn et al. 2007).

But during an ecological shift, natural selection can influence birth-death dynamics by promoting the expansion or contraction of specific gene families. For lineages that adapt to a novel niche,

their genomes may exhibit evidence of selective maintenance of duplications or deletions in
certain gene families. For example, if selection favors gene gain, novel gene duplicates will tend
to persist in the population, resulting in subfamilies of recently diverged paralogs. If the
mutational mechanism that generates new duplicate genes is unequal crossing over during
meiosis, these recently diverged paralogs will be found in tandem arrays across the genome
(Zhang 2003). Moreover, if gene duplicates experience positive selection for novel functions,
this can result in elevated amino acid substitution rates among paralogs. Conversely, some
genetic functions may become obsolete or even deleterious in the novel habitat. When this
occurs, positive or relaxed purifying selection will cause some gene families to accumulate loss-
of-function mutations at an accelerated rate.
81
Sometime after an ecological shift, impacted gene families will reach a new equilibrium state in
which gene number evolves primarily via negative selection and genomic drift. Likewise,
pseudogenes will fade into the genomic background (Petrov et al. 1996; Petrov and Hartl 1997,
1998) and tandem array lengths will reflect local recombination rates (Akhunov et al. 2003;
Zhang and Gaut 2003; Rizzon et al. 2006; Thomas 2006). Thus, within-genome signatures of
adaptive changes in gene family size are likely ephemeral and best detected in lineages that
recently shifted to a novel niche. Nevertheless, if natural selection consistently favors the
expansion or contraction of specific gene families in specific environments, among-taxon
correlations between gene family size and ecology will be maintained. Currently, the extent to
which different taxa exhibit convergence at the level of gene family size changes is largely
unknown.
93
Arguably, the genes most likely to expand and contract convergently in response to similar
selection pressures are those that mediate organismal interactions with their biotic and abiotic
environments. These "environmentally responsive genes" include chemosensory (e.g., olfactory
and gustatory receptors), detoxifying (e.g., cytochrome P450), and immunity (e.g., MHC and
immunoglobulin) genes. To cope with constantly changing pressures, environmentally
responsive genes tend to be characterized by elevated sequence diversity, duplication rates,
substitution rates, and genomic clustering, as well as tissue- or temporal-specific expression
(Berenbaum 2002) and limited pleiotropy (Arguello et al. 2016). Importantly, causal links
between changes in environmentally responsive genes and adaptation to novel niches have been
established for multiple taxa (Després et al. 2007; Matsuo et al. 2007; Dobler et al. 2012; Zhen et
al. 2012; Sezutsu et al. 2013).

With their exceptionally diverse ecologies and ever-increasing availability of annotated genomes
(Consortium 2013; Poelchau et al. 2015), insects are a powerful system for investigating the
predictability of size changes in environmentally responsive gene families. To date, at least two
ecological transitions are hypothesized to have a predictable impact on gene family size in insect
lineages. In plant-feeding insects, the evolution of increased dietary specialization (i.e., smaller
diet breadth) is associated with reduced chemosensory and detoxifying gene family sizes
(McBride 2007; McBride and Arguello 2007; Good et al. 2014; Goldman-Huertas et al. 2015;
Calla et al. 2017; Comeault et al. 2017) but see (Gardiner et al. 2008). In hymenopteran insects,
eusociality is associated with expansions of the olfactory-receptor family and contractions of the
gustatory-receptor family (Robertson and Wanner 2006; Zhou et al. 2015; McKenzie et al. 2016;
Brand and Ramírez 2017) but see (Fischman et al. 2011; Johnson et al. 2018). Most of these
studies, however, considered a single ecological characteristic or gene family (but see Robertson

and Wanner 2006) which is potentially problematic in that changes in social behavior may often
be accompanied by changes in other ecological characteristics and vice versa (Faulkes et al.
1997; Duffy and Macdonald 2010; Ross et al. 2013). A better understanding of ecology and gene
family size relationships will require simultaneous consideration of multiple ecological
characteristics and diverse gene families.

Here, we characterize multiple environmentally responsive gene families in the genome of the
redheaded pine sawfly, *Neodiprion lecontei* (Order: Hymenoptera; Family: Diprionidae). This
species provides an opportunity to examine both within-genome signatures of adaptive gene
family contractions and expansions, and among-lineage correlations between ecology and gene
family size. First, for within-genome signatures, *N. lecontei* is an exemplar of an herbivorous
hymenopteran lineage (Diprionidae) that underwent a drastic host shift: sometime within the last
60 million years, this lineage transitioned from angiosperms to coniferous host plants in the
family Pinaceae (Boevé et al. 2013; Peters et al. 2017). To defend against herbivores and
pathogens, Pinaceae produce viscous oleoresin secretions that are sticky and have unique
antimicrobial properties (Trapp and Croteau 2001; Gershenzon and Dudareva 2007). To manage
these toxic and extraordinarily sticky resins, *N. lecontei* and related diprionids evolved
specialized feeding and egg-laying traits (Figure 1). Beyond these traits, we hypothesize that
pine specialization likely resulted in pronounced changes to the selection pressures acting on
multiple gene families, especially those involved in chemosensation, detoxification, and immune
function. Second, with respect to among-lineage correlations between ecology and gene-family
size, *N. lecontei* is an herbivorous, non-eusocial insect from the Eusymphyta, a massively
understudied hymenopteran clade (Peters et al. 2017). Although many assembled and annotated
hymenopteran genomes are currently available, almost all have come from apocritans (bees,
wasps, and ants, but see (Robertson et al. 2018)). Thus, *N. lecontei* increases the ecological,
behavioral, and taxonomic diversity of hymenopteran genomes for evaluating ecological
correlates of gene family size among taxa.
143
To evaluate the predictability of gene family evolution, we assembled a draft genome for *N.*
lecontei and manually annotated genes for five environmentally responsive gene families:
olfactory receptor (OR), gustatory receptor (GR), odorant binding protein (OBP), cytochrome
P450 (CYP), and antimicrobial peptide (AMP). For gene families that underwent a size change
related to pine adaptation, we expected one or more of the following patterns: (1) clusters of
recently diverged paralogs in gene-family trees, (2) a high proportion of genes in tandem arrays,
(3) signatures of positive selection among paralogs, and (4) elevated rates of pseudogenization.
Then, for the same five gene families, we asked whether gene-family size correlated with
ecology among distantly related insect taxa. To do so, we compiled published gene annotations
and ecological variables (diet type, degree of ecological specialization, presence/absence of
eusociality) for hymenopteran taxa. Together, these analyses identify possible candidate gene
families underlying pine specialization and reveal that relationships between gene family size
and ecology differ among environmentally responsive gene families.
157
158

Results
Genome assembly and annotation
Sequencing and assembly

We sequenced one mate-pair and two small-insert Illumina libraries made from haploid male
siblings (see Methods). After read processing, we retained 268 billion PE100 reads with a
combined read depth of 112x (Table S1). ALLPATHS-LG (v47417) (Gnerre et al. 2011)
produced a 239-Mbp assembly consisting of 4523 scaffolds, with a scaffold N50 of 243 kbp
(Table S2). Prior studies identified seven chromosomes in *N. lecontei* (Smith 1941; Maxwell
1958; Sohi and Ennis 1981; Linnen et al. 2018). With an estimated genome size (1C) of 331 ±9.6
Mbp, our assembly captured 72% of the genome. Overall, these metrics are comparable to other
hymenopteran assemblies (Table S2).
170
To measure assembly completeness and artificial sequence duplication, we used CEGMA (Parra
et al. 2007) and BUSCO (Simão et al. 2015). Both search the assembly for a set of single-copy,
conserved genes, however, the CEGMA software has been deprecated
(http://korflab.ucdavis.edu/Datasets/cegma). Of the 248 CEGMA core eukaryotic genes, 90%
aligned as complete, single copies and 8% aligned complete but duplicated. For BUSCO, we
used the OrthoDB arthropod dataset, and out of 2675 groups 77% were complete, single copies
and 3% were complete but duplicated. These metrics indicate the presence of artificial duplicate
sequences, but otherwise the assembly was reasonably complete and suitable for annotation.
179
About 15.8% of the assembly consisted of repetitive elements, including 122 unknown
transposable elements that were mostly unique to *N. lecontei* (Table S3), and 212 other families
of transposable elements and simple repeats. This 15.8% corresponds to 11.4% of the actual 331-
Mb genome, of which we predict 27.6% is repetitive, suggesting that ~16.1% of the missing
~28% of the genome is repetitive content (Table S3). For de novo gene prediction, we
included the *N. lecontei* transcriptome and protein-coding genes from *Atta cephalotes*
(OGSv1.2), *Acromyrmex echinatior* (OGSv3.8), *Apis melifera* (OGSv3.2), *Athalia rosae*
(OGSv1.0), and *Nasonia vitripennis* (OGSv1.0) to guide annotation. The official gene set
(OGSv1) had 12,980 gene models while the transcriptome had an average of 26,000 transcripts
per tissue (Table S4).
190
Olfactory receptor
The OR gene family had 56 genes total, including the co-receptor *Orco*; one gene contained stop
codons, three were partial annotations, and 52 genes were intact (Table 1). In *D. melanogaster*
most olfactory sensory neurons (OSNs) express a single OR (along with the coreceptor, *Orco*),
and OSNs expressing a particular OR converge on a single glomerulus in the antennal lobe (Gao
et al. 2000; Vosshall et al. 2000; Couto et al. 2005) but see (Fishilevich and Vosshall 2005). This
anatomy results in a general one-to-one correspondence between the number of ORs and the
number of glomeruli, a correspondence also observed in the hymenopteran European honey bee
(*Apis mellifera*, (Robertson and Wanner 2006)). Based on these studies and examination of the
antennal lobes of adult male and adult female *N. lecontei*, we expected to find a minimum of 49
functional ORs (Table S5, Figure 2). The close correspondence between our gene annotations
and glomeruli counts suggests that we have located all or most *N. lecontei* OR genes.
203
59% of ORs were in genomic clusters of two or more genes (Figure 3), a low proportion
compared to many other hymenopteran OR families (Robertson and Wanner 2006; Zhou et al.
2015; McKenzie et al. 2016; Brand and Ramírez 2017). A phylogenetic analysis of OR protein
sequences from *Neodiprion,* six other hymenopterans, and *D. melanogaster* identified three

Neodiprion-specific clades with at least five genes (Figure S1a). These same three clades were
also recovered in a phylogenetic analysis of *Neodiprion* OR cDNA sequences (Figure S1b). For
each *Neodiprion*-specific OR clade (and *Neodiprion*-specific clades in other gene families, see
below), we used the *Neodiprion* cDNA tree, the codeml program in the PAML package (Yang
2007), and likelihood-ratio tests to ask: (1) whether the ratio of non-synonymous to synonymous
substitution rates (dN/dS or ω) for the focal OR clade differed from the rest of the *Neodiprion*
OR gene family and, if so, whether they exhibited evidence of positive selection (ω>1) (branch
tests); and (2) whether ω differed among sites across members of *Neodiprion*-specific clades
and, if so, which sites exhibited evidence of positive selection (site tests). For only one OR clade
(OR clade 1) did we detect evidence of branch-specific positive selection (i.e., rejection of both
1-ratio and fixed-ω models), but this clade lacked evidence of site-specific positive selection
(Table 2).
220
Gustatory receptor
The GR gene family had 44 genes total; two genes contained stop codons, two were partial
annotations (one annotation was both partial and pseudogenized), and 41 were intact (Table 1).
76% of the GRs that could be placed on chromosomes were in genomic clusters (Figure 3) with
three *Neodiprion*-specific clades of at least five genes (Figures S2a and S2b). Only one clade
(GR clade 3) had evidence of branch-specific positive selection (Table 2). This clade also had
evidence of positive selection at some amino acid positions among paralogs (Table 2; sites with
evidence of positive selection include: 77E, 79S, 146N, 275S, 301S). Notably, GR Clade 3 is an
expansion of six paralogs orthologous to *DmGR66a*, a bitter receptor specifically for caffeine
(Moon et al. 2006). However, *N. lecontei* orthologs were not found for *DmGR93a* (Lee et al.
2009) and *DmGr33a* (Moon et al. 2009), coreceptors possibly required for caffeine detection.
Together, these data suggest that caffeine-like GR receptors have been coopted for novel
functions in *N. lecontei*.
234
The GR family also had orthologs for sugar receptors *DmGR5a* (trehalose) (Dahanukar et al.
2001), *DmGR43a* (fructose) (Miyamoto et al. 2012), and *DmGR64a-f* (multiple sugars) (Slone et
al. 2007) as well as carbon dioxide receptors *DmGR21a* and *DmGR63a* (Jones et al. 2007)
(Figure S2a). Orthologs to these carbon dioxide receptors have not been found in Apocrita but
seem to be preserved in Symphyta, like *N. lecontei* (Robertson and Kent 2009; Robertson et al.
2018).
241
Odorant binding protein
The OBP gene family had 13 genes total; none were pseudogenized or partial annotations (Table
1). In this family, 38% of genes were in genomic clusters, including a cluster of five genes on
chromosome 6 (Figure 3). *Neodiprion*-specific OBP clades were not found, even for the
chromosome 6 cluster. We note, however, that the OBP phylogenies had low bootstrap support
(Figure S3a,b), making it difficult to infer relationships among paralogs.
248
Cytochrome P450
The CYP gene family had 107 genes total; twelve genes contained stop codons, two were partial
annotations, and 93 were intact (Table 1). In insects, CYPs belong to four major clades, which
are referred to as clans (Feyereisen 2012). When we split the CYP gene family by clan, the
CYP2 clan had nine intact genes; the CYP3 clan had 47 intact genes and eight pseudogenes; the

CYP4 clan had 27 intact genes, four pseudogenes, and two partial genes; and the mitochondrial
CYP clan had 10 intact genes (Table 1). Across all CYPs, 66% were in genomic clusters (Figure
3). Looking at the four major clans separately, the percentage of clustered genes were: 33% for
CYP2, 81% for CYP3, 55% for CYP4, and 50% for mitochondrial CYP.
258
The CYP gene family had five *Neodiprion*-specific clades with at least five genes (Figure S4a,b),
four of which were in the CYP3 clan. Of these, two clades that were both within the CYP3 clan
(CYP clades 3 and 5) had evidence of branch-specific, but not site-specific, positive selection
(Table 2). CYP clade 3 contained members of the CYP6 subfamily, and the CYP clade 5
contained members of the CYP336 subfamily. Several studies to date suggest that members of
the CYP3 clan—and the CYP6 subfamily in particular—play an important role in detoxifying
pesticides and host-plant allelochemicals (Feyereisen 2012).
266
Orthologs were found for all the Halloween genes (which include genes from both the CYP2 and
mitochondrial CYP clans) of the 20-hydroxy ecdysone biosynthesis pathway: *CYP302A1*
(disembodied), *CYP306A1* (phantom), *CYP307A2* (spookier), *CYP307B1* (spookiest),
CYP314A1 (shade), *CYP315A1* (shadow), and *CYP18A1* which turns over 20-hydroxy ecdysone
(Rewitz et al. 2007; Feyereisen 2011; Guittard et al. 2011; Qu et al. 2015). The juvenile hormone
biosynthesis gene *CYP15A1* was present as well (Helvig et al. 2004). Finally, *N. lecontei* had
orthologs for the two CYP4G enzymes that synthesize the cuticular hydrocarbons used as
external waterproof coating (Qiu et al. 2012).
275
Immunity
Antimicrobial peptides (AMPs) are expressed upon infection to kill or inhibit microbes. Based
on hymenopteran sequences, the *N. lecontei* AMP gene family had 21 genes (Table 1; Table S6),
including single copies of *Hymenoptaecin*, *Abaecin*, and *Tachystatin*, but no clear *Defensin*
ortholog. Over 18 *Hisnavicin* genes were identified, including a *Neodiprion*-specific expansion
of eight histidine-rich paralogs orthologous to *Hisnavicin-4*, which has been characterized as a
larval cuticle protein and AMP, but not functionally tested (Tian et al. 2010). The *N. lecontei*
Hisnavicins had a conserved 62 amino acid motif that appeared up to 19 times in a single
protein; the purpose of this amplification is unknown. 95% of the AMPs were in genomic
clusters (Figure 3). Due to low bootstrap support on many of the branches in our Hisnavicin
protein tree, we could not identify unambiguous *Neodiprion*-specific clades (Figure S5a).
However, our *Neodiprion* cDNA tree (Figure S5b) did reveal strong support for the monophyly
of a cluster of 15 *Hisnavicins* on linkage group 5 (Figure 3), and this cluster had some evidence
of positive selection (Table 2).
290
Outside of the AMP family, most immune pathways had direct orthologs between *N. lecontei*
and *D. melanogaster* (Figure S6, Table S7). The basic viral siRNA response pathway was
completely conserved between species. The immune deficiency (IMD) pathway was missing an
ortholog for the peptidoglycan recognition receptor *PGRP-LC*, but it is likely that another *PGRP*
replaced *PGRP-LC* in *N. lecontei*; assigning PGRP orthology was also difficult in ants (Gupta et
al. 2015). Also missing is the *Drosophila* mitogen activated protein kinase kinase kinase, TGF-β
activated kinase 1 (*Tak1*), but *N. lecontei* had a similar TGF-β activated kinase that is a close
ortholog to several *Tak1-like D. melanogaster* proteins possibly involved in immune deficiency
signaling. The encapsulation/melanization pathway was missing one of the two *Drosophila*

GTPases (*Rak2*). The *N. lecontei Rak1* ortholog may be playing both roles, but again this is
likely due to the difficulty of assigning one-to-one orthologs. The Duox pathway was missing the
top G-protein coupled receptor, but this is unknown in *D. melanogaster* and unidentified in other
Hymenoptera (Evans et al. 2006). Interestingly, *N. lecontei* had two copies of Dual Oxidase
(*Duox*), which regulates commensal gut microbiota and infectious microbes (Ha et al. 2005; Lee
et al. 2015); *Apis mellifera* had one copy. Finally, the Toll pathway *NF-kappaB* transcription
factor, *Dorsal-related immunity factor (Dif)* does not have a one-to-one ortholog in *N. lecontei*,
but two copies of its paralog, *Dorsal*, were present.
308
309   **Within-genome signatures of adaptive expansions and contractions**
310
Evidence of selection in Neodiprion-*specific gene family clades*
Massive gene family expansions with dozens of genes were not found in *N. lecontei* (in contrast
to (Smadja et al. 2009; Zhou et al. 2015)). Instead, the largest *Neodiprion*-specific clade had 22
genes (CYP gene family) and the rest had fewer than 10 genes. Nevertheless, we did identify 11
Neodiprion-specific clades containing at least 5 closely related paralogs and a monophyletic
clade of 15 AMPs with ambiguous ancestry (Table 1). Of these 12 clades, four had significant
branch positive selection (OR clade 1, GR clade 3, CYP clade 3, and CYP clade 5) (Table 2). Of
these four clades, only one also had significant site-specific positive selection (GR clade 3)
(Table 2).
320
Clustering
Our five focal gene families varied in the proportion of genes that were found in clusters of two
or more genes (Fisher's exact test, $P = 0.002$; Table 1). Post-hoc tests revealed that much of this
variation was due to differences between the highly clustered AMP family and all other families
except GR (AMP vs. OR: $P = 0.0091$; AMP vs. OBP: $P = 0.0053$; AMP vs. CYP: $P = 0.024$,
AMP vs. GR: $P = 0.12$; all p-values are FDR-corrected). The only other difference in clustering
that we detected was between the GR and singleton-heavy OBP families (FDR-corrected $P =$
0.045).
329
Differences in clustering were even more pronounced when we separated the CYP family by
clan (Fisher's exact test, $P < 0.0001$; Table 1). In addition to the pairwise differences described
above, we also found that the proportion of CYP3 genes found in clusters differed significantly
from ORs, CYP2s, and CYP4s (all FDR-corrected $P < 0.05$), but not AMPs, GRs, and
mitochondrial CYPs. Additionally, AMP clustering differed from CYP2, CYP4, and
mitochondrial CYP, while GR differed from CYP2 (all FDR-corrected $P < 0.05$). Together, these
analyses identified AMP, GR, and CYP3 as having an unusually high proportion of genes found
in clusters compared to other environmentally responsive gene families.
338
Pseudogenization
Overall, we found very few pseudogenes, and the proportion of pseudogenized genes did not
differ significantly among gene families (Fisher's exact test, $P = 0.12$; Table 1). The
chemoreceptors had one pseudogene each while CYP had 12, which is about 10% of the family,
but this was also the largest gene family. Although CYP3 had more pseudogenes than other CYP
clans, the proportion of pseudogenized genes still did not differ when we compared CYP clans

(Fisher's exact test, $P = 0.10$). Given these low rates of pseudogenization, it is unlikely that *N.*
lecontei gene families underwent substantial, recent contractions.
347
Ecological correlates of gene-family size across insects
We first examined broad-scale variation in the sizes of our five focal gene families and four CYP
clans among different insect orders (Figure S7). Not surprisingly, sample sizes were highly
variable across gene families and insect orders. Despite this variation, we observed some
intriguing differences among gene families and taxa. We detected significant differences in gene
family size among orders for OR (Kruskal-Wallas chisq = 48.2, df = 12, $P < 1 \times 10^{-5}$), GR (K-
W chisq = 25.5, df = 9, $P = 0.0025$), and OBP (K-W chisq = 37.6, df = 9, $P < 1 \times 10^{-4}$), but not
CYP (K-W chisq = 10.3, df = 7, $P = 0.17$) or AMP (K-W chisq = 7.93, df = 5, $P = 0.16$). We
note, however, that the AMP sample size was considerably smaller than the other gene families.
When we looked at CYP clans individually, we found differences among orders for CYP4 (K-W
chisq = 19.0, df = 7, $P = 0.0083$) and mitochondrial CYP (K-W chisq = 16.3, df =7, $P = 0.022$),
but not CYP2 (K-W chisq = 9.19, df = 7, $P = 0.24$) or CYP3 (K-W chisq = 8.76, df = 7, $P =$
0.27).
361
For the OR family, among-group differences in gene number were mostly attributable to an
unusually large number of OR genes in Hymenoptera (significant post-hoc tests include Diptera
vs. Hymenoptera: $P = 0.0018$; Hemiptera vs. Hymenoptera: $P = 0.00014$; and Odonata vs.
Hymenoptera: $P = 0.011$; all p-values are FDR-corrected). By contrast, the size of the OBP
family was larger in Diptera than other orders (significant post-hoc tests include Diptera vs.
Hymenoptera: $P = 0.00037$; Diptera vs. Hemiptera: $P = 0.00092$; all p-values are FDR-
corrected). Although none of the post-hoc tests were significant for GR family size, the Blattodea
appear to have more GRs on average than other insect orders (Figure S7). For CYP clans,
posthoc tests revealed that hymenopterans have fewer CYP4s than dipterans (FDR-corrected $P =$
0.010) and fewer mitochondrial CYPs than both dipterans and lepidopterans (FDR-corrected $P =$
0.024 and 0.023, respectively).
373
We next examined how gene family size correlated with ecology within the hymenopteran clade
(Figures 4 and 5). Once again, we observed differences among gene families. We found that the
number of ORs differed significantly among hymenopteran species that differed in diet (Kruskal-
Wallas chisq = 15.8, df = 3, $P = 0.0012$) and sociality (Wilcoxon rank-sum test W = 115; $P =$
0.00094). For diet, we found that herbivores had fewer ORs than all other diet types (fungivores
vs. herbivores: $P = 0.015$; omnivores vs. herbivores: $P = 0.015$; insectivores vs. herbivores: $P =$
0.048; all p-values are FDR-corrected). We observed an even more striking difference between
eusocial and non-eusocial hymenopterans, with the former having larger OR families, on
average. By contrast, GR family size was related to diet (Kruskal-Wallas chisq = 11.8, df = 3, P
= 0.0082), but not sociality (Wilcoxon rank sum test W = 30; $P = 0.65$). And CYP family size
was related to sociality (W = 2; $P = 0.045$), but not diet ($P = 0.38$). Finally, specialists and
generalists did not differ significantly in gene family size in any of the gene families and ecology
was unrelated to gene family size for OBP and CYP (total CYP number and individual CYP
clans). Although these analyses have several limitations (see discussion), these results are
consistent with the hypothesis that environmentally responsive gene families may contract or
expand predictably in response to particular selection pressures.
390

Discussion

The predictability of gene family expansion or contraction in response to specific selection pressures remains an open question. Here, we evaluated genomic signatures of adaptive gene family size changes in five environmentally responsive gene families within the *N. lecontei* draft genome, a hymenopteran exemplar of a pine-specialized lineage. Although we saw minimal evidence of recent gene loss via pseudogenization, at least three gene families (AMP, GR, and CYP3) had genomic distributions consistent with selective maintenance of novel gene duplicates, and two of these families also exhibited evidence of positive selection within *Neodiprion*-specific clades (GR and CYP3). Next, we examined these same gene families in other hymenopterans to see if family size correlated with diet, ecological specialization, or eusocial behavior. Among Hymenoptera, we found that OR family size was correlated with both eusociality and diet type, but not dietary specialization; GR family size was correlated with diet type; and AMP family size was associated with eusociality. These results suggest that ecology can have a predictable impact on gene family size and that different selection pressures impact different gene families. Below, we discuss both the implications and limitations of our analyses and suggest priorities for future comparative work on gene family size evolution.

Within-genome signatures of gene-family size change

During a niche shift, new selective pressures can leave footprints in the genomes of evolving lineages; such signatures of positive selection are well described for individual loci (Nielsen et al. 2005; Vitti et al. 2013). Similarly, strong selection for increases or decreases in the size of a particular gene family should also leave characteristic genomic footprints. We argue that these footprints include monophyletic groups of closely related paralogs in gene-family trees (from the selective maintenance of novel duplicates), genomic clustering (when novel genes arise via unequal crossing over), evidence of positive selection among paralogs (if there is selection for sub- or neofunctionalization), and high rates of pseudogenization (from the selective maintenance of loss-of-function mutations). Of the environmentally responsive gene families we evaluated, none exhibited patterns consistent with selection for a decrease in gene family size. By contrast, at least three families had characteristics consistent with selection for an increase in gene family size. Two of these families, GR and CYP3, were highly clustered in the genome and exhibited evidence of positive selection, making these especially promising candidates for expansions related to a novel coniferous host. Additionally, although the AMP family lacked evidence of positive selection, its unusually clustered distribution in the *Neodiprion* genome could be related to selection for increased dosage of a conserved protein function (Perry et al. 2007). Below we discuss the functions of these three candidate families in more detail.

Shifts to pine feeding likely involved changes in the detection of and response to pine-specific cues. Intriguingly, the one GR clade with evidence of positive selection—GR clade 3—is an expansion of six paralogs (one is pseudogenized) orthologous to *DmGR66a*, a bitter receptor specifically for caffeine (Moon et al. 2006). However, orthologs were not found for *DmGR93a* (Lee et al. 2009) and *DmGr33a* (Moon et al. 2009), coreceptors possibly required for caffeine detection. Nevertheless, honeybees, which also lack clear orthologs to these putative coreceptors (Wanner and Robertson 2008), can detect and even prefer low concentrations of caffeine and nicotine (Singaravelan et al. 2005, but see de Brito Sanchez 2011). Although pines do not

contain caffeine, they do synthesize alkaloids that could confer some bitterness (Mumm and
Hilker 2006). Thus, despite lacking caffeine coreceptor orthologs, members of GR clade 3 may
still be involved in the detection of pine-specific bitter compounds. Duplications of putative
bitter GRs are documented in other host-specialized insects, such as *Heliconius, Danaus,* and
Bombyx butterflies (Wanner and Robertson 2008; Briscoe et al. 2013). Our sawfly-specific GR
expansion, coupled with the finding that GR family size is associated with diet (see below), lends
support to the hypothesis that expansions of GR bitter receptors repeatedly contribute to changes
in oviposition and feeding behaviors in plant-feeding insects.
444
Because pines contain toxic components like terpenoids and phenolics, detoxifying gene families
are also promising candidates for pine adaptation. The mountain pine beetle (*Dendroctonus*
ponderosae), feeds on pine bark and wood and has gene "blooms" (species-specific gene gains)
in the CYP3 and CYP4 clans (Keeling et al. 2013). Similarly, in *N. lecontei*, the CYP family had
five blooms (Figure S4a): four CYP3 and one CYP4. CYP3 blooms are also found in wood-
feeding insects that do not use pine, such as the emerald ash borer (*Agrilus planipennis*) (David
Nelson, unpublished data) and the Asian longhorned beetle (*Anoplophora glabripennis*)
(McKenna et al. 2016). Notably, *N. lecontei* larvae frequently ingest pine bark in addition to pine
needles (Wilson 1992), suggesting that CYP3 may expand predictably in wood feeders.
Additionally, one of the two *Neodiprion*-specific CYP3 clades with evidence of positive
selection (Table 3) belongs to the CYP6 subfamily, which is linked to host plant adaptation in
several insect taxa (Li et al. 2003; Li et al. 2007; Feyereisen 2012; Mittapelly et al. 2019).
457
Because pine resin has antimicrobial (Himejima et al. 1992; Cowan 1999; Gershenzon and
Dudareva 2007) and fungicidal properties (Grayer and Harborne 1994), we hypothesized that *N.*
lecontei co-opted these compounds for its own defense, leading to relaxed selection on genes
involved in immunity and a reduced innate immune response. In other Hymenoptera, honeybees
(*Apis mellifera*) exposed to plant resin have reduced expression of immune-related genes
(Simone et al. 2009) and wood ants (*Formica paralugubris*) that use conifer resin as building
material have slightly reduced inducible immune system activity and nests with lower bacterial
and fungal loads (Castella et al. 2008). In Diptera, AMP loss is associated with herbivorous
lineages that live within host tissue, a more sterile habitat than is experienced by most dipterans
(Hanson et al. 2019). Unexpectedly, we found a large species-specific clade of *Hisnavicin*-like
AMPs in *Neodiprion*. Although additional data are needed to confirm that *Hisnavicin* orthologs
act as AMPs in *N. lecontei*, one possible explanation for this putatively adaptive expansion that
lacked an accompanying change in non-synonymous substitution rate is that having large
numbers of *Hisnavicin*-like AMPs confers protection against pathogens unique to pine trees.
That said, our data do not rule out adaptive AMP loss. For example, *N. lecontei* lacks a clear
Defensin ortholog, a gene present in all dipterans tested to date (Hanson et al. 2019).
474
Limitations of within-genome analyses
One benefit to studying adaptive expansions/contractions within a single taxon is that gene
families have likely experienced similar demographic histories, which can also impact gene birth
and death rates. That said, each of our within-genome signatures of selection has limitations that
should be revisited with additional data. First, our analysis of genomic clustering does not
account for local recombination rate variation, which correlates with tandem array size in several
taxa (Gaut et al. 2007). A fine-scale recombination rate map, coupled with clustering analyses

for many additional gene families, would more rigorously test the extent to which individual
gene family clustering deviates from the genome-wide relationship between recombination rate
and tandem array size.
485
Second, a lack of comparable data from other Eusymphyta meant that our gene family
phylogenies lacked orthologues from closely related sawfly taxa. Thus, the "*Neodiprion*-
specific" clades may not be unique to pine-feeding sawflies. If these paralogs were present prior
to the shift to pine hosts, this would not support a scenario in which new duplicates were
selectively maintained in the novel niche. Signatures of positive selection may still be related to
pine adaptation but would indicate selection on preexisting loci rather than selection favoring
gene family expansion.
493
Third, signatures of adaptive gene family expansions and contractions may be ephemeral, and
the shift to pine use could have occurred too long ago to detect these signatures in *N. lecontei*.
For example, in *Drosophila*, pseudogenes have an estimated half-life of ~14.3 million years
(Petrov et al. 1996; Petrov and Hartl 1997, 1998). If the rate of gene decay is similar in
Neodiprion, then pseudogenes that formed after a shift to pine (up to 60 mya) may no longer be
detectable in the genomes of extant sawflies. Likewise, gene clustering patterns are likely to
change over time from chromosomal rearrangements and additional gene duplications and
deletions. To investigate how the number and position of genes in these focal families has
changed over time, high quality gene annotations for diprionids and many additional sawfly
outgroups are needed. Fortunately, even if footprints of recent gene family size changes are too
ephemeral to be detected in most taxa, consistent relationships with ecology should still be
detectable given sufficient sampling of taxa differing in ecological traits of interest.
506
Ecological correlates of gene family size among hymenopteran taxa
The largest insect OR gene families are in eusocial Hymenoptera, leading to the hypothesis that
OR family size expansions were favored in these lineages because they facilitate complex
chemical communication (Robertson and Wanner 2006; LeBoeuf et al. 2013; Zhou et al. 2015).
To date, evidence in support of this hypothesis has been mixed (e.g., (Roux et al. 2014; Brand
and Ramírez 2017). Consistent with the OR-eusociality hypothesis, we found that, on average,
eusocial hymenopterans had larger OR families than non-eusocial hymenopterans. However, it is
likely that eusocial taxa differ from non-eusocial taxa in many other aspects of their ecology that
should also impact OR evolution. Indeed, we found that herbivorous hymenopterans tended to
have fewer OR genes than non-herbivores.
517
Whereas all eusocial hymenopterans had relatively large OR families, some eusocial
hymenopterans had relatively small GR families (Figures 4, 5; (Zhou et al. 2015)). To explain
the strikingly small set of GR genes in honeybee, Wanner and Robinson (2006) proposed that a
stable hive environment and a mutualistic relationship with flowering plants resulted in a lack of
selection for GR expansions. Intriguingly, our data indicate that among hymenopterans, GR
family size is associated with diet, but not eusociality. Like the ORs, GR gene family size tends
to be smaller in herbivores than in non-herbivorous taxa, regardless of social behavior. The
directionality of this change, however, is unclear: do shifts to plant diets favor reductions in GR
families, do shifts to non-plant diets favor GR expansions, or is it both? Answering this question
will require characterizing GR families across many independent transitions to and from

herbivory, as well as polarizing directions of change (i.e., distinguishing GR gains from GR
losses). Fortunately, there are many such diet transitions across diverse clades of insects (Wiens
et al. 2015).
531
Unlike sociality and diet, ecological specialization was not associated with gene family size in
any of the five gene families we evaluated. This result was unexpected because specialization-
associated reductions in gene family size are documented in diverse taxa and multiple gene
families, including the families examined here (McBride 2007; Smadja et al. 2009; Cao et al.
2014; Goldman-Huertas et al. 2015; Suzuki et al. 2018). One explanation for the lack of
association between gene family sizes and specialization in our data is that our "generalist" and
"specialist" categories are not meaningful across diverse diets (Forister et al. 2012).
Additionally, within a particular diet, the degree of specialization may be highly labile, with
rapid fluctuations that are not captured in our broad, order-wide comparison. Indeed, previous
studies that reported correlations between gene family size and ecological specialization focused
on closely related species. Thus, to fully understand how changes in ecology shape gene family
evolution, it will be necessary to evaluate ecological correlates of gene family size at multiple
time scales of taxonomic divergence.
545
Compared to ORs and GRs, our other focal gene families had far less manual annotation data
available for analysis. This may explain, in part, why we did not detect strong ecological
correlates for the other gene families. It is also possible that by focusing on the sizes of entire
gene families, we missed relevant signals in particular subfamilies (Hahn et al. 2007). For
example, as noted above, expansions of CYP3 and CYP4 subfamilies are associated with wood-
feeding insects and CYP3 clan subfamilies were also linked to detoxification in honey bee
(Berenbaum and Johnson 2015; Johnson et al. 2018). However, we did not detect any
correlations between ecology and CYP clan sizes. Despite these limitations, we did uncover hints
that AMP gene family size may be larger in non-eusocial lineages. If eusocial taxa tend to
inhabit more sterile environments (nests and hives) than non-eusocial taxa, this finding is
consistent with associations between habitat and AMP loss reported in dipterans (Hanson et al.
2019). Given that AMPs were also implicated in our within-genome analysis, immune-related
genes are especially promising candidates for future manual annotation projects.
559
Limitations of among-taxon analyses
Comparative analysis is a powerful approach for evaluating the repeatability and predictability of
evolutionary outcomes. Although our comparison of candidate gene family sizes among
ecologically diverse hymenopterans hints at intriguing relationships between ecology and gene
family size, it also had several limitations that should be revisited in future work. First, because
several taxa in our manual annotation dataset are missing from published hymenopteran
phylogenies (Peters et al. 2017), we were unable to correct for phylogenetic non-independence
and polarize gene gain/loss (e.g., as in (Hahn et al. 2005; Han et al. 2013) without losing
unacceptable amounts of data. Without accounting for similarity in ecology and gene family size
due to recent common ancestry, our Type I error rate is likely inflated and p-values should be
interpreted with caution. Nevertheless, variation in patterns of association among ecological
traits and gene families suggest that phylogeny and ecology are, to some extent, decoupled.
572

The gene annotation and ecological datasets also had limitations. For example, across studies
that included manual annotations, we observed a lack of consistency in the methods and criteria
for manually curated gene family datasets. The most problematic inconsistency was in the
criteria for delineating intact, partial, and pseudogenized gene annotations. "Intact" could mean
an exon-by-exon check against closely related orthologs, a minimum amino acid length, or
merely the presence of an expected domain. Meanwhile, in reference publications, the number of
pseudogenized and partial annotations were not always reported or were conflated. This is in
addition to variation in the methods used to search for genes. Inconsistency in annotation
methods and criteria across studies may introduce taxon-specific biases that have nothing to do
with ecology. Regarding ecology, categorizations are somewhat subjective. For example, this
study and Rane et al. (2016) classified bees as generalists since they collect nectar and pollen
from multiple plant families (we defined specialization as the use of a single taxonomic family).
But Johnson et al. (2018) classified bees as specialists as their diet consists of only nectar and
pollen.
587
Finally, our attempts to correlate the size of different gene families with ecology suffered from
sampling biases in which species had genome assemblies and which gene families were
manually annotated. Species skewed heavily towards *Drosophila* and apocritan Hymenoptera,
and annotations toward the OR and CYP families (Table S8). To evaluate ecological correlates
of gene family expansions and contractions, it is essential to expand both the taxonomic breadth
and depth of annotation sampling. Taxa that capture independent ecological transitions (e.g.,
between herbivory and other diets) would be especially useful, as would replicated groups of
closely related species that vary in ecological axes of interest (e.g., specialization or social
behavior). By systematically sampling different ranges of divergence times, we can evaluate the
extent to which the tempo of gene family size change varies across different gene families. To do
so, however, will require high quality, manually curated datasets produced using consistent
methods and standards for many different environmentally responsive gene families.
600
601
Conclusions
Gene families that mediate ecological interactions may expand and contract predictably in
response to changing selection pressures. These adaptive changes in gene family size should
leave detectable footprints in the genomes of recent niche colonists, and across taxa with
convergent niche shifts. Consistent with these predictions, our analysis of gene family evolution
in a derived pine feeder suggests that expansions of GRs, CYP3s, and AMPs may have
accompanied pine adaptation, and our comparison among ecologically diverse hymenopterans
links two of these same two gene families to variation in diet (GR) and eusociality (AMP). In the
order Hymenoptera, the OR gene family was associated with ecology (eusociality), while other
ecological axes of variation (specialization/generalization) were not linked to gene family size
(but have been in comparisons of other closely related species (McBride 2007)). Together, these
results suggest that environmentally responsive gene families vary in both temporal dynamics
(shallow vs. deep divergence times) and in ecological drivers of size change. Teasing apart these
relationships will require high quality annotation data across diverse gene families, ecologies,
and divergence times. For hymenopterans, increased effort in understudied symphytan,
parasitoid, and herbivorous taxa would be especially useful for disentangling different axes of
ecological variation contributing to gene family size change.

619
Materials and methods
Biological material
To minimize the confounding effects heterozygosity has on genome assembly, we sequenced
haploid siblings. Like all Hymenoptera, sawflies have haplodiploid sex determination in which
males (haploid genomes) emerge from unfertilized eggs and females (diploid genomes) from
fertilized eggs. A virgin female will bear a clutch of all-male offspring with haploid
recombinants of the maternal genome. But the individual genomes are not identical, so an
assembly derived from a single clutch is akin to a diploid assembly made from a single
individual.

All insects were reared in custom, climate-controlled environmental chambers (18:6 light cycle,
22°C, 70% RH) on jack pine (*Pinus banksiana*) foliage. Our laboratory line of *N. lecontei* was
established from multiple larval colonies collected from a mugo pine (*P. mugo*) in Lexington,
Kentucky, USA (37°59'01.6"N 84°30'38.8"W; population ID: RB017). For the transcriptome,
adults and larvae were collected from the first lab-reared generation; both were stored at -80°C.
For the genome assembly, the founding population was propagated in the lab for two
generations, followed by brother-sister matings for an additional two generations. At this point, a
single, virgin, adult female (I2G2-V, 4th generation in the lab) was allowed to lay unfertilized
eggs onto jack pine seedlings. The offspring (haploid male brothers from an inbred mother) were
reared until the eonymph (prepupal) life stage, at which point they were isolated without food for
24 hours prior to preservation in absolute ethanol at -20°C. Although eonymphs are non-feeding,
they were starved to ensure the gut contents were completely voided.

Sample preparation and sequencing
Genomic DNA
Whole eonymph bodies were individually frozen inside microcentrifuge tubes with liquid
nitrogen and ground with pestles made from 1-mL micropipette tips; the resulting powder was
incubated in CTAB buffer supplemented with proteinase K and RNase A. After PCI extraction
and ethanol precipitation, the precipitate was dried overnight before being resuspended in TE
buffer. DNA integrity was assessed with 0.7% agarose gel, purity was measured with the
260/280 ratio, and concentration was measured with a Quant-iT dsDNA High-Sensitivity
fluorescence assay (Thermo Fisher Scientific).

The HudsonAlpha Genomic Services Lab (Huntsville, AL, USA) prepared and sequenced the
DNA libraries. Two small-insert, barcoded libraries with average fragment sizes of 337 bp and
864 bp were made from a single individual. A 4.6-kbp mate-pair, barcoded library was made
from 25 pooled individuals. All individuals were brothers from the same I2G2-V mother. The
libraries were sequenced on Illumina HiSeq 2000 with paired-end, 100 bp (PE100) reads: the
small-insert libraries each had ¼ of a flow cell lane and the mate-pair library had an entire lane.

mRNA
The RNeasy Mini extraction kit (Qiagen) was used to collect total RNA from adult female body,
adult female head, adult male body, adult male head, eonymph body, feeding larval body, and
feeding larval head. RNA from eonymph head was extracted but not sequenced due to
insufficient yield. Each tissue was represented with one replicate that had equal RNA

665　contributions from eight individuals, except for eonymph body which was comprised of three
666　individuals. RNA integrity and concentration were measured with a 2100 Bioanalyzer (Agilent).
667
668　The HudsonAlpha Genomic Services Lab (Huntsville, AL, USA) handled library preparation and
669　sequencing. Non-stranded, barcoded libraries were made for each of the seven tissue samples; on
670　average, mRNA was sheared to 200 bp. The libraries were combined and sequenced on an entire
671　flow cell of Illumina MiSeq with PE250 reads in addition to one lane of Illumina HiSeq 2000
672　with PE50 reads.
673
Read processing and assembly
De novo genome assembly
Sequencing reads were chastity-filtered and adaptor-trimmed with fastq-mcf (ea-utils v1.04.803)
(Aronesty 2011), and quality-filtered with fastq_quality_filter (FASTX Toolkit v0.0.13.2)
(Gordon and Hannon 2019). The 337-bp small-insert reads and the 4.6-kbp mate-pair reads were
quality-filtered to retain reads where at least 80% of the bases had a quality score of 20 or higher
(parameters: `-q 20 -p 80`). Due to sequencing quality, the 864-bp small-insert reads were filtered
to retain reads where at least 70% of the bases had a quality score of 20 or higher (R1) or 60%
(R2) (parameters: `-q 20 -p 60/70`). In situations where only one end of the paired-end reads
passed filtering, the passed reads were kept and treated as single-end data. Kmer counting was
used to measure read depth before and after filtering (Jellyfish v1.1.11) (Marçais and Kingsford
2011). Finally, reads were screened for sequencing contamination by mapping the reads (BWA
v0.7.12-r1039) (Li and Durbin 2009) to reference genomes for *Escherichia coli* (K12 substr.
DH10B uid58979), human (v37), loblolly pine (*Pinus taeda*, v0.8), and *Wolbachia*
(endosymbiont of Dmel uid57851).
689
The genome was assembled with ALLPATHS-LG (v47417) (Gnerre et al. 2011) using default
settings, including a minimum scaffold size of 1000 bp. The error-correction module was run on
the reads prior to assembly. After assembly, GapFiller (v1.11) (Boetzer and Pirovano 2012) was
used to help close intra-scaffold gaps. Spurious scaffolds were identified with SOAP.coverage
(v2.7.7) (Li et al. 2009): reads were mapped to the assembly scaffolds and scaffolds with a read
depth < 15 and nucleotide percentage < 40 were removed. The completeness of the final
assembly was measured with CEGMA (v2.5) (Parra et al. 2007) and BUSCO (v1.22) (Simão et
al. 2015) benchmarks. BUSCO was run with the arthropoda-25oct16 database (parameters: `--`
`long`).
699
De novo transcriptome assembly
For both the PE250 MiSeq and the PE50 HiSeq reads, fastq-mcf (ea-utils v.1.04.803) (Aronesty
2011) was used for chastity filtering and Trimmomatic (v0.32) (Bolger et al. 2014) was used to
adaptor clip, trim, and quality-filter. The PE250 MiSeq reads were processed with the
Trimmomatic parameters `ILLUMINACLIP: 2:15:5, HEADCROP: 10, CROP: 60, MINLEN: 60,`
`AVGQUAL: 25` whereas the PE50 HiSeq reads were processed with `ILLUMINACLIP: 2:15:5,`
`HEADCROP: 15, MINLEN: 35, AVGQUAL: 25`. Because the mRNA libraries had an average insert
size of 200 bp, the MiSeq reads required extensive adaptor trimming. Reads were screened for
contamination as described in *De novo genome assembly*.
709

710 For each tissue, transcriptomes were assembled with Trinity (r2013_08_14) (Grabherr et al.
711 2011; Haas et al. 2013) using default settings and the --jaccard_clip option. Spurious sequences
712 were identified by mapping the sequencing reads to the assembled transcripts with RSEM
713 (v1.2.18) (Li and Dewey 2011); transcripts with either FPKM or TPM values < 1 were removed.
714 After filtering, the transcriptomes were combined, and duplicate sequences were removed.
715
Genome size estimation
Flow cytometry was performed as described in (Harper et al. 2016). For this analysis, we used
adult males and females from a lab line of *N. lecontei* established from a colony collected in
Auburn, GA (33°59'22.4" N, 83°47'44.6"W; population ID: RB027). Briefly, cell nuclei were
collected from the heads of 7 individuals (4 female, 3 male) and stained with propidium iodide.
Mean fluorescence for each sample was measured with a BD FACSCalibur flow cytometer (BD
Biosciences) and compared to two external standards: *Drosophila melanogaster* (adult female
heads, 1C = 175 Mbp) and *Gallus gallus domesticus* (CEN singlets from BioSure, Grass Valley,
CA, 1C = 1222.5 Mbp). To correct for ploidy differences between haploid males and diploid
standards, we multiplied the *N. lecontei* male estimates by 2. To obtain a single size estimate for
each *N. lecontei* sample, we averaged values obtained for the two standards.
727
728 **Repeat annotation**
The *N. lecontei* genome assembly was masked with a custom repeat library. A lineage-specific
de novo repeat library was made with RepeatModeler (v1.0.7) (Smit and Hubley 2008-2015) and
combined with the hymenopteran repetitive element database (Nov. 2013) from Repbase (Jurka
et al. 2005). The custom library was used by RepeatMasker (v4.0.3) (Smit et al. 2013-2015)
(parameters: -cutoff 250 -s -pa 15 -gc 40 -a -poly) to identify and mask repetitive elements
in the genome, including low-complexity DNA and simple repeats.
734
Transposable element (TE) family consensus sequences were identified by rerunning
RepeatModeler (Smit and Hubley 2008-2015) on the genome assembly using the "ncbi" search
engine. The resulting sequences were provided to RepeatMasker (Smit et al. 2013-2015) as a
custom library to locate associated TE copies in the genome (parameters: -gc 40 -cutoff 250 -
gff -gccalc -norna -nolow -no_is -poly). TE families with at least 10 fragments longer than
100 bp were extracted for further analysis.
741
The sequencing reads were mapped to a concatenation of the masked genome and the consensus
TE sequences (BWA MEM (parameters: -M) (Li and Durbin 2009)). Families that had at least 1x
the median coverage to the reference genome for at least 80% of their sequence (to support at
least one full insertion found by RepeatModeler) and at least 2x the maximum coverage of the
reference genome (to support multiple insertions of the family) were extracted with
genomeCoverageBed (BEDtools (Quinlan and Hall 2010)). We attempted to identify the
consensus sequences with BLASTN and BLASTX (Altschul et al. 1990) searches against a
database of repeat elements, but the only hits were to the lineage-specific elements identified by
RepeatModeler. Sequences were also filtered for BLAST hits to rRNA or mitochondrial
sequences.
752
753 We also used dnaPipeTE (Goubert et al. 2015) to identify what proportion of our short reads was
754 composed of repetitive content, we used a random subset of reads corresponding to 1-fold
755 coverage of the genome (331Mb) and took the total for three separate random samplings of reads

756   (parameters: genome size = 331000000 genome coverage = 1 samples number = 3). We then
757   compared this annotation to the RepeatModeler annotation.
758
759

Gene and functional annotation
Automated gene annotation
RNA-Seq data for *N. lecontei* was used to generate training models for gene prediction along
with utilization of peptide sequences from other species. PASA (r20130425beta) was used to
build a comprehensive transcriptome set from Trinity assembled transcripts along with RNA-Seq
read mapping predictions generated from the Tuxedo pipeline. To improve annotation quality, in
addition to this *N. lecontei* transcriptome, annotated proteins from *Atta cephalotes* (OGSv1.2),
Acromyrmex echinatior (OGSv3.8), *Apis melifera* (OGSv3.2), *Athalia rosae* (OGSv1.0), and
Nasonia vitripennis (OGSv1.0) were provided to Maker (2.09) (Cantarel et al. 2008) as evidence
for structural gene prediction. Prior to annotation, the genome was masked using a custom repeat
database built using RepeatModeler (v1.0.8) and the annotation was run using the *ab initio* gene
predictors Augusts, Genemark-ES and snap in addition to the evidence provided. The functions
of the predicted protein-coding genes were putatively established with BLASTP alignments
(Altschul et al. 1990) to the Swiss-Prot database (accessed 20 Apr 12) (Apweiler et al. 2004). In
cases of multiple matches, the top-ranked alignment was assigned to the gene annotation. Protein
motifs and functional domains within the annotations were also identified with an InterProScan
(v5.3.46.0) (Jones et al. 2014) search against the InterPro database with gene ontology and IPR
lookup (Finn et al. 2016). For the official gene set (OGS), the Maker annotations were filtered by
hits to the reference databases and/or a minimum eAED score of 0.1. A second set of gene
annotations was generated with the NCBI GNOMON pipeline (annotation release 100 on
Nlec1.0 assembly, GCF_001263575.1) (Souvorov et al. 2010).
781
782   Because the genome was annotated prior to submission to NCBI, we encountered a problem
783   when the NCBI contamination software flagged vector/adaptor sequences for removal; this
784   would disrupt the coordinates provided by Maker. We used a modified version of GAG (Hall et
785   al. 2014) that could accept the flagged coordinates from NCBI to edit the assembly and update
786   annotation coordinates accordingly.
787

Chemoreceptor genes
The olfactory (OR) and gustatory (GR) receptor genes were manually curated following
Robertson et al. (2003, 2006). Amino acid sequences of manually curated chemoreceptor genes
from *Apis mellifera* (Robertson and Wanner 2006; Smith et al. 2011), *Bombus terrestris* (Sadd et
al. 2015) and *Cephus cinctus* (Robertson et al. 2018), *Drosophila melanogaster* (Flybase release
FB2017_04), and *Nasonia vitripennis* (Robertson et al. 2010) were used as queries in TBLASTN
(v2.2.19) (Altschul et al. 1990) searches against the *N. lecontei* draft genome (parameters: –e
`100000 –F F`). Gene models were manually built in TextWrangler (v5.5) (Bare Bones Software),
using protein alignment to identify exons and refine the gene structures; alignments were
visualized with Clustal X (v2.1) (Larkin et al. 2007). The Neural Network Splice Predictor
program from the Berkeley *Drosophila* Genome Project was used to help identify intron splice
sites (http://www.fruitfly.org/seq_tools/splice.html). New gene models were added to subsequent
TBLASTN searches and this process continued iteratively until new chemoreceptors were no
longer found. The gene models were checked against RNAseq reads from tissue-specific

transcriptomes (adult antennae, mouthparts, heads, legs, genitalia, and larval heads (Herrig et al.
2019)) and against orthologs in the *N. pinetum* draft genome assembly (NCBI accession
GCA_004916985.1).
805
806

Odorant binding proteins

Custom scripts were used to identify Maker gene annotations (see *Automated gene annotation*)
that contained the classic/6C, Plus-C, Minus-C, or atypical odorant binding protein (OBP) motif
(Xu et al. 2009). These as well as OBPs from *Apis mellifera* and *Nasonia vitripennis* were used
as queries for TBLASTN searches against the *N. lecontei* genome; searches did not yield any
new OBPs. All genomic regions identified as potential OBPs were manually curated as described
for chemoreceptor genes. After manual annotation, duplicate annotations or genes that lacked
OBP motifs were removed.
815

Cytochrome P450 genes

A broad set of 52 insect CYP genes (covering the diversity of insect CYP families) were
searched against the *N. lecontei* genome assembly (E-value cutoff 1e3). Scaffolds with hits were
then searched against 8782 known insect CYPs. The top 10 hits were returned (later increased to
15 to recover more sequences) and filtered for duplicates. An alternative search of the NCBI
GNOMON predictions ("Neodiprion lecontei[orgn] AND P450 NOT reductase") was also
performed and new sequences were added to the dataset. This approach found all the loci
identified by the initial search, indicating that the GNOMON annotation tool was able to
comprehensively search for CYP sequences. Finally, the candidate *N. lecontei* CYP sequences
were manually curated based on comparison to the best BLAST hits.
826

Immune-related genes

Because of the relative completeness of its immune annotation, *Drosophila melanogaster*
immunity genes were used to guide annotation. Reference immune genes from *D. melanogaster*
tagged with the gene ontology term "GO:0002376 – Immune system process" were compiled
from Flybase (release 6.13). Orthology with *N. lecontei* proteins was assigned initially with
reciprocal BLASTP (Altschul et al. 1990) searches (E-value cutoff 1e-10). Reference *D.*
melanogaster genes without obvious one-to-one orthologs in *N. lecontei* were examined
individually to determine whether closely related paralogs in one or both species interfered with
the inference of orthology. If not, they were searched against the *N. lecontei* genome assembly
using TBLASTN (Altschul et al. 1990) in an attempt to identify unannotated orthologs.
837
Since antimicrobial peptides (AMP) are unlikely to be conserved between *D. melanogaster* and
N. lecontei, AMPs from three representative hymenopterans *Apis mellifera* (Danihlík et al.
2015), *Nasonia vitripennis* (Tian et al. 2010), and *Camponotus floridanus* (Ratzka et al. 2012;
Zhang and Zhu 2012; Gupta et al. 2015) were used for BLAST queries. Furthermore, since AMP
copy number is fast evolving, we attempted to find all the *N. lecontei* orthologs of each
hymenopteran AMP instead of focusing on one-to-one orthology. Once again, BLASTP searches
were performed against the annotated proteins and TBLASTN searches were performed against
the assembled genome; the TBLASTN search did not reveal additional AMPs. Putative *N.*
lecontei orthologs were reciprocally blasted against the appropriate hymenopteran proteome to
assure that the best hits were indeed AMPs.

Amino acid and cDNA sequences for chemoreceptor genes, CYP genes, and AMP genes are
available in File S1.

Glomeruli counts

Antennal lobe histology
Whole heads of adult *N. lecontei* of both sexes were fixed in 2% paraformaldehyde, 2%
glutaraldehyde in PBS for 5 days. Heads were rinsed for 40 minutes three times and the brains
dissected out in cold PBS. Following blocking with goat serum, brains were permeabilized with
1% Triton X-100 in PBS (Electron Microscopy Supply, Fort Washington, PA; PBS-TX), rinsed
with 0.1% PBS-TX, and incubated on a shaker at 25°C for three nights in primary antibody
(1:500 in 2% goat serum in 0.2% PBS-TX). Monoclonal *Drosophila* synapsin I antibody
(SYNORF1, AB_2315426) from the Developmental Studies Hybridoma Bank (catalog 3C11)
was used to label synapsin. Subsequently, brains were washed in 0.1% PBS-TX and incubated
for two nights in Alexa Fluor 568 (ThermoFisher) goat anti-mouse secondary antibody (1:100 in
PBS) in the dark at room temperature on a shaker. After secondary incubation, brains were
rinsed with distilled water, dehydrated in increasing concentrations of ethanol, and mounted in
custom-made aluminum well slides. Brains were cleared by removing ethanol and replacing it
with methyl salicylate. Brains were imaged on an inverted Zeiss 880 Laser Scanning Confocal
Microscope with a 20X plan-Apochromat 20x 0.8 aperture objective and optically sectioned in
the horizontal plane at 3-micron intervals.

Glomeruli segmentation
Whole-brain images of one female and one male were manually segmented using the TrakEM2
software package in ImageJ (Cardona et al. 2012; Schindelin et al. 2012). Individual glomeruli
were traced in both brain hemispheres. Glomeruli near the center of the antennal lobe can be
difficult to distinguish, meaning counts are biased toward fewer glomeruli. Therefore, the largest
number of glomeruli confidently detected represents a minimum of the number of expected
glomeruli.   Male *Neodiprion* have a collection of smaller synaptic clusters in their antennal lobe
(Dacks and Nighorn 2011), but the functional significance of this anatomy is not known. There
are more than 50 of these smaller synaptic clusters and we suspect they do not represent the
traditional one-to-one OR-to-glomerulus organization. Therefore, these structures were not
included in counts. Male glomeruli number may be lower if particular OSNs contribute to these
clusters instead of forming traditional glomeruli.

Within-genome signatures of adaptive expansions and contractions
Clustering and pseudogene analyses
To evaluate the extent to which members of our five focal gene families were located in tandem
arrays, we placed our annotated genes on a linkage-map anchored version of the *N. lecontei*
genome assembly described in Linnen et al. 2018. We considered members of a gene family to
be clustered if they were located within a genomic region of 20(n - 1) kilobases, where n is the
number of genes in the cluster under consideration. This criterion was chosen based on average
gene densities in *Nasonia* (Niehuis et al. 2010) and clustering criteria described *Drosophila*
(Vieira et al. 2007). For scaffolds that could not be placed on linkage groups, we evaluated
clustering only if genes were more than 20 kb from either end of the scaffold.

894
To evaluate whether the five focal gene families differed in (1) the proportion of genes that were
found in clusters of two or more genes or (2) the proportion of genes that were pseudogenized,
we performed Fisher's exact tests in R v3.5.0 ("fisher.test" function) (R-Core-Team 2018). For
significant Fisher's exact tests, we performed additional posthoc tests using the
"fisher.multcomp" function (from R package RVAideMemoire v. 0.9-72) with FDR correction
(Benjamini-Hochberg method) for multiple comparisons.
901
Identification of* Neodiprion-*specific clades and tests of positive selection
First, we identified clades unique to *N. lecontei*. For each gene family, a multi-species, amino
acid phylogeny was constructed with manually curated annotations from *N. lecontei*, select
Hymenoptera, and *D. melanogaster*. Sequences were size filtered (350≥ for GR, OR, CYP; 100≥
for histnavicin and OBP), but pseudogenes and partial annotations that met the length
requirement were retained. MAFFT alignments (v7.305b) (Katoh et al. 2002) (parameters: --
`maxiterate 1000 –localpair`) were visually inspected to remove sequences with large alignment
gaps, and sites with more than 20% gaps were removed with trimAl (v1.4.rev15 build[2013-12-
17]) (Capella-Gutiérrez et al. 2009) (parameters: `–gapthreshold 0.8`). Maximum likelihood
phylogenies were made in RAxML (v8.2.4) (Stamatakis 2014) (parameters: `–f a –x 12345 –p`
`12345 –# autoMRE`) using protein substitution models chosen from ProtTest3 (v3.4.2) (Abascal et
al. 2010; Darriba et al. 2011).
914
Neodiprion-specific clades were defined as those with at least five *N. lecontei* genes (not
including partial and pseudogenes) and a bootstrap score ≥70 (Engsontia et al. 2015). Second,
the clades were confirmed with cDNA phylogenies for each *N. lecontei* gene family. Amino acid
sequences were aligned as above, however, after alignment TranslatorX (Abascal et al. 2010)
was used to map cDNA sequences to the amino acid alignment. After trimming, the cDNA
alignments were passed to RAxML to construct maximum likelihood gene family trees with the
nucleotide substitution model `–m GTRGAMMA`.
922
Site tests were conducted with codeml (part of the PAML package (PAML v4.9e) (Yang 2007))
using the cDNA phylogenies and sequences as inputs. For each *Neodiprion*-specific clade, the
gene family cDNA phylogeny was pruned to remove all branches except for that clade. Codeml
models M7, M8, and M8a were fitted to the cDNA sequence and phylogeny data. Likelihood-
ratio tests were performed for the nested models M7-M8 (null model M7 that equally distributes
amino acid sites across 10 classes of ω parameter values (p, q) against alternative model M8 that
has an 11[th] class for positively selected sites) and M8-M8a (null model M8a that has 11 classes
and does not allow positive selection against alternative model M8). Bonferroni correction was
applied to the likelihood-ratio test probability values; each value was multiplied by two since two
tests that used M8 as the alternative model were performed on each clade. For clades with
significant likelihood-ratio tests, sites under selection were identified by looking at the Bayes
Empirical Bayes analysis within the alternative models.
935
For branch tests, the cDNA phylogenies for each *N. lecontei* gene family were used to compare
the lineage-specific clade to the rest of the gene family. To determine if the dN/dS of the
foreground branch (i.e., the branch with the species-specific expansion) was significantly
different from the background (i.e., the rest of the gene family), in codeml we ran a two-ratio

model (Model=2, fix_omega=0) and a one-ratio model (Model=0, fix_omega=0) for that clade
and performed a likelihood-ratio test comparing the two models. To determine if the foreground
branch is under positive selection (dN/dS>1), we performed a likelihood-ratio test comparing the
two-ratio model to a neutral model (fix_omega=1).
944
945
Ecological correlates of gene family size among insects
All the insect genome assembly projects we could find (published and unpublished) were
searched for manually curated chemoreceptor, odorant binding protein, cytochrome P450, and
antimicrobial peptide gene annotations. If fasta sequence files were available, the number of
intact, partial, and pseudogenized genes was determined based on gene names (e.g., labels with
"pse" or "partial"); values were compared to those reported in the publication. Otherwise, we
relied on reported values. If gene family size was reported but not broken down into intact,
partial, and pseudogenized, and sequence files were unavailable, we assumed that the reported
number referred to intact genes. Splice variants were not included in the gene count. It is
important to note that different authors likely used different criteria for these categories.
956
For comparisons, only the number of putatively functional (intact) genes was used. Species were
classified according to taxonomic order, diet type, dietary specialization, and sociality. An order
needed at least two species to be included. Specialization was defined as the use of a single
taxonomic family and only referred to the realized diet niche, ignoring reports of feeding under
laboratory conditions. If a species had a preferred host or both specialist and generalist life
stages, it was classified as specialist. Gene family size comparisons were made in R (v3.5.0)
where species were grouped by the different classifications.
964
Because both gene family size and ecology are likely to correlate with phylogeny, the ideal
approach to identifying ecological correlates of gene family evolution is to use statistical
methods that account for phylogenetic relationships (Hahn et al. 2005; De Bie et al. 2006; Han et
al. 2013). Unfortunately, a lack of overlap between species with manual annotations for our focal
gene families and species included in published hymenopteran genomes precluded us from such
an analysis without a substantial loss of sample size. Therefore, as a first step to evaluating
ecological correlates of gene family size, we used non-parametric tests to determine whether
gene family size differed among taxa. For sociality and specialization, we used two-tailed
Wilcoxon rank-sum tests ("wilcox.exact" function in the exactRankTests v0.8-30 package). For
taxonomic order and diet, both of which have more than two categories, we used Kruskal-Wallis
tests ("kruskal.test" function) followed by Dunn's post-hoc tests of multiple comparisons
("dunnTest" function in the FSA v0.8.23 package).
977

Acknowledgements
We thank Linnen lab members for insect collection, insect reading, and reading earlier
manuscripts; Jeramiah Smith and Erin Scully for advice. We are especially grateful to Hugh
Robertson for his guidance on annotating ORs and GRs. This work was supported by
the University of Kentucky Center for Computational Sciences and the Lipscomb High
Performance Computing Cluster, the United States Department of Agriculture National Institute
of Food and Agriculture (2016-67014-2475; CRL), the Kentucky Science and Engineering

Foundation (KSEF-3492-RDE-019; CRL), and the University of Kentucky (Lyman T. Johnson
Fellowship; KV).
987
988   **Data availability**
989   The genome assembly, official gene set (OGS), and transcriptome described in this paper (v1
990   versions) can be found at https://i5k.nal.usda.gov/neodiprion-lecontei
991
On GenBank (NCBI), the genome assembly is labeled whole genome shotgun sequencing
project accession PRJNA28045 and the genomic sequencing reads are RefSeq accession
PRJNA312506. The transcriptome is transcriptome shotgun assembly accession
GEDM00000000; this is a combined transcriptome file with sequences from all seven tissue
types. The mRNA sequencing reads for each tissue type was submitted separately under
BioSample and short read archive accessions SAMN04302192 (adult female head),
SAMN04302193 (adult female body), SAMN04302194 (adult male head), SAMN04302195
(adult male body), SAMN04302196 (feeding larval head), SAMN04302197 (feeding larval
body), and SAMN04302198 (eonymph body). The predicted gene annotations on NCBI are from
Gnomon, the NCBI annotation pipeline, and were not described in this paper. Finally, the
clustering analysis was based on a linkage-map anchored version of the genome assembly
described in Linnen et al. 2018. This anchored assembly is denoted as v1.1 in NCBI and the *N.*
lecontei i5k Workspace@NAL (USDA).
1005
1006

References

1007
1008
Abascal F, Zardoya R, Telford MJ. 2010. TranslatorX: multiple alignment of nucleotide
sequences guided by amino acid translations. Nucleic Acids Res 38(suppl 2):W7-W13.
Akhunov ED, Goodyear AW, Geng S, Qi L-L, Echalier B, Gill BS, Gustafson JP, et al. 2003.
The organization and rate of evolution of wheat genomes are correlated with recombination
rates along chromosome arms. Genome Res 13(5):753-763.
Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool.
J Mol Biol 215(3):403-410.
Apweiler R, Bairoch A, Wu CH, Barker WC, Boeckmann B, Ferro S, Gasteiger E, et al. 2004.
UniProt: the universal protein knowledgebase. Nucleic Acids Res 32(suppl 1):D115-D119.
Arguello JR, Cardoso-Moreira M, Grenier JK, Gottipati S, Clark AG, Benton R. 2016.
Extensive local adaptation within the chemosensory system following Drosophila
melanogaster's global expansion. Nat Commun 7(1):1-12.
Aronesty E. 2011. ea-utils : "Command-line tools for processing biological sequencing data".
Berenbaum MR. 2002. Postgenomic chemical ecology: from genetic code to ecological
interactions. J Chem Ecol 28(5):873-896.
Berenbaum MR, Johnson RM. 2015. Xenobiotic detoxification pathways in honey bees. Curr
Opin Insect Sci 10:51-58.
Boetzer M, Pirovano W. 2012. Toward almost closed genomes with GapFiller. Genome Biol
13(6):R56.
Boevé J-L, Blank SM, Meijer G, Nyman T. 2013. Invertebrate and avian predators as drivers of
chemical defensive strategies in tenthredinid sawflies. BMC Evol Biol 13(1):198.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics 30(15):2114-2120.
Brand P, Ramírez SR. 2017. The evolutionary dynamics of the odorant receptor gene family in
corbiculate bees. Genome Biol Evol 9(8):2023-2036.
Briscoe AD, Macias-Munoz A, Kozak KM, Walters JR, Yuan F, Jamie GA, Martin SH, et al.
2013. Female behaviour drives expression and evolution of gustatory receptors in butterflies.
PLoS Genet 9(7):e1003620.
Calla B, Noble K, Johnson RM, Walden KKO, Schuler MA, Robertson HM, Berenbaum MR.
2017. Cytochrome P450 diversification and hostplant utilization patterns in specialist and
generalist moths: Birth, death and adaptation. Mol Ecol 26(21):6021-6035.
Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, Holt C, Alvarado AS, Yandell M.
2008. MAKER: an easy-to-use annotation pipeline designed for emerging model organism
genomes. Genome Res 18(1):188-196.
Cao D, Liu Y, Walker WB, Li J, Wang G. 2014. Molecular characterization of the Aphis
gossypii olfactory receptor gene families. PLoS One 9(6):e101187.
Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25(15):1972-1973.
Cardona A, Saalfeld S, Schindelin J, Arganda-Carreras I, Preibisch S, Longair M, Tomancak P,
Hartenstein V, Douglas RJ. 2012. TrakEM2 software for neural circuit reconstruction. PLoS
One 7(6):e38011.
Castella G, Chapuisat M, Moret Y, Christe P. 2008. The presence of conifer resin decreases the
use of the immune system in wood ants. Ecological Entomology 33(3):408-412.
Comeault AA, Serrato-Capuchina A, Turissini DA, McLaughlin PJ, David JR, Matute DR.
2017. A nonrandom subset of olfactory genes is associated with host preference in the fruit
fly Drosophila orena. Evolution Letters 1(2):73-85.
Consortium iK. 2013. The i5K Initiative: advancing arthropod genomics for knowledge, human
health, agriculture, and the environment. J Hered, 104(5):595-600.
Couto A, Alenius M, Dickson BJ. 2005. Molecular, anatomical, and functional organization of
the Drosophila olfactory system. Curr Biol 15(17):1535-1547.
Cowan MM. 1999. Plant products as antimicrobial agents. Clin Microbiol Rev 12(4):564-582.
Dacks AM, Nighorn AJ. 2011. The organization of the antennal lobe correlates not only with
phylogenetic relationship, but also life history: a basal hymenopteran as exemplar. Chem
Senses 36(2):209-220.
Dahanukar A, Foster K, Carlson JR. 2001. A Gr receptor is required for response to the sugar
trehalose in taste neurons of Drosophila. Nat Neurosci 4(12):1182-1186.
Danihlík J, Aronstein K, Petřivalský M. 2015. Antimicrobial peptides: a key component of
honey bee innate immunity: Physiology, biochemistry, and chemical ecology. Journal of
Apicultural Research 54(2):123-136.
Darriba D, Taboada GL, Doallo R, Posada D. 2011. ProtTest 3: fast selection of best-fit models
of protein evolution. Bioinformatics 27(8):1164-1165.
Dayhoff MO. 1976. The origin and evolution of protein superfamilies Federation Proceedings
35(10):2132-2138.
De Bie T, Cristianini N, Demuth JP, Hahn MW. 2006. CAFE: a computational tool for the
study of gene family evolution. Bioinformatics 22(10):1269-1271.
de Brito Sanchez MG. 2011. Taste perception in honey bees. Chem Senses 36(8): 675-692.

Demuth JP, Hahn MW. 2009. The life and death of gene families. Bioessays 31(1):29-39.
Després L, David J-P, Gallet C. 2007. The evolutionary ecology of insect resistance to plant
chemicals. Trends Ecol Evol 22(6):298-307.
Dobler S, Dalla S, Wagschal V, Agrawal AA. 2012. Community-wide convergent evolution in
insect adaptation to toxic cardenolides by substitutions in the Na, K-ATPase. Proc Natl
Acad Sci USA 109(32):13040-13045.
Duffy JE, Macdonald KS. 2010. Kin structure, ecology and the evolution of social organization
in shrimp: a comparative analysis. Proc Royal Soc B 277(1681):575-584.
Engsontia P, Sangket U, Robertson HM, Satasook C. 2015. Diversification of the ant odorant
receptor gene family and positive selection on candidate cuticular hydrocarbon receptors.
BMC Res Notes 8(1):380.
Evans JD, Aronstein K, Chen YP, Hetru C, Imler JL, Jiang H, Kanost M, Thompson GJ, Zou
Z, Hultmark D. 2006. Immune pathways and defence mechanisms in honey bees Apis
mellifera. Insect Mol Biol 15(5):645-656.
Faulkes CG, Bennett NC, Bruford MW, O'Brien HP, Aguilar GH, Jarvis JU. 1997. Ecological
constraints drive social evolution in the African mole–rats. Proc Royal Soc B
264(1388):1619-1627.
Feyereisen R. 2011. Arthropod CYPomes illustrate the tempo and mode in P450 evolution.
Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics 1814(1):19-28.
Feyereisen R. 2012. Insect CYP genes and P450 enzymes. In. Insect Mol Biol and Biochemistry:
Academic Press. p. 236-316.
Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, et al. 2016.
The Pfam protein families database: towards a more sustainable future. Nucleic Acids Res
44(D1):D279-D285.
Fischman BJ, Woodard SH, Robinson GE. 2011. Molecular evolutionary analyses of insect
societies. Proc Natl Acad Sci USA 108(suppl 2):10847-10854.
Fishilevich E, Vosshall LB. 2005. Genetic and functional subdivision of the Drosophila antennal
lobe. Curr Biol 15(17):1548-1553.
Forister ML, Dyer LA, Singer MS, Stireman III JO, Lill JT. 2012. Revisiting the evolution of
ecological specialization, with emphasis on insect–plant interactions. Ecology 93(5):981-991.
Gao Q, Yuan B, Chess A. 2000. Convergent projections of Drosophila olfactory neurons to
specific glomeruli in the antennal lobe. Nat Neurosci 3(8):780-785.
Gardiner A, Barker D, Butlin RK, Jordan WC, Ritchie MG. 2008. Drosophila chemoreceptor
gene evolution: selection, specialization and genome size. Mol Ecol 17(7):1648-1657.
Gaut BS, Wright SI, Rizzon C, Dvorak J, Anderson LK. 2007. Recombination: an
underappreciated factor in the evolution of plant genomes. Nat Rev Genet 8(1):77-84.
Gershenzon J, Dudareva N. 2007. The function of terpene natural products in the natural world.
Nat Chem Biol 3(7):408-414.
Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, et al. 2011.
High-quality draft assemblies of mammalian genomes from massively parallel sequence data.
Proc Natl Acad Sci USA 108(4):1513-1518.
Goldman-Huertas B, Mitchell RF, Lapoint RT, Faucher CP, Hildebrand JG, Whiteman NK.
2015. Evolution of herbivory in Drosophilidae linked to loss of behaviors, antennal responses,
odorant receptors, and ancestral diet. Proc Natl Acad Sci USA 112(10):3026-3031.

Good RT, Gramzow L, Battlay P, Sztal T, Batterham P, Robin C. 2014. The molecular
evolution of cytochrome P450 genes within and between Drosophila species. Genome Biol
Evol 6(5):1118-1134.

Gordon A, Hannon GJ. 2019. "Fastx-toolkit" FASTQ/A short-reads preprocessing tools
(unpublished).

Goubert C, Modolo L, Vieira C, Claire ValienteMoro, Mavingui P, Boulesteix M. 2015. De novo
assembly and annotation of the Asian tiger mosquito (Aedes albopictus) repeatome with
dnaPipeTE from raw genomic reads and comparative analysis with the yellow fever
mosquito (Aedes aegypti). Genome Biol Evol 7(4):1192-1205.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, et al. 2011.
Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat
Biotechnol 29(7):644-652.

Grayer RJ, Harborne JB. 1994. A survey of antifungal compounds from higher plants, 1982–
1993. Phytochemistry 37(1):19-42.

Gross JB, Borowsky R, Tabin CJ. 2009. A novel role for Mc1r in the parallel evolution of
depigmentation in independent populations of the cavefish Astyanax mexicanus. PLoS Genet
5(1):e1000326.

Guittard E, Blais C, Maria A, Parvy J-P, Pasricha S, Lumb C, Lafont R, Daborn PJ, Dauphin-
Villemant C. 2011. CYP18A1, a key enzyme of Drosophila steroid hormone inactivation, is
essential for metamorphosis. Dev Bio l349(1):35-45.

Gupta SK, Kupper M, Ratzka C, Feldhaar H, Vilcinskas A, Gross R, Dandekar T, Förster F.
2015. Scrutinizing the immune defence inventory of Camponotus floridanus applying total
transcriptome sequencing. BMC Genom 16(1):1-21.

Ha E-M, Oh C-T, Bae YS, Lee W-J. 2005. A direct role for dual oxidase in Drosophila gut
immunity. Science 310(5749):847-850.

Haas BJ, Papanicolaou A, Yassour M, Grabherr MG, Blood PD, Bowden J, Couger MB, et al.
2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform
for reference generation and analysis. Nat Protoc 8(8):1494-1512.

Hahn MW, de Bie T, Stajich JE, Nguyen C, Cristianini N. 2005. Estimating the tempo and
mode of gene family evolution from comparative genomic data. Genome Res 15(8):1153-
1160.

Hahn MW, Han MV, Han S-G. 2007. Gene family evolution across 12 Drosophila genomes.
PLoS Genet 3(11):e197.

Hall B, DeRego T, Geib SM. 2014. GAG: the genome annotation generator (version 1.0).

Han MV, Thomas GW, Lugo-Martinez J, Hahn MW. 2013. Estimating gene gain and loss rates
in the presence of error in genome assembly and annotation using CAFE 3. Mol Biol Evol
30(8):1987-1997.

Hanson MA, Lemaitre B, Unckless RL. 2019. Dynamic evolution of antimicrobial peptides
underscores trade-offs between immunity and ecological fitness. Front Immunol 10:2620.

Harper KE, Bagley RK, Thompson KL, Linnen CR. 2016. Complementary sex determination,
inbreeding depression and inbreeding avoidance in a gregarious sawfly. Heredity 117(5):326-
335.

Helvig C, Koener JF, Unnithan GC, Feyereisen R. 2004. CYP15A1, the cytochrome P450 that
catalyzes epoxidation of methyl farnesoate to juvenile hormone III in cockroach corpora
allata. Proc Natl Acad Sci USA 101(12):4024-4029.

Herrig DK, Vertacnik KL, Linnen CR. 2019. Testing the adaptive decoupling hypothesis in a
hypermetamorphic and sexually dimorphic insect. bioRxiv 2019.12.20.882803.
Himejima M, Hobson KR, Otsuka T, Wood DL, Kubo I. 1992. Antimicrobial terpenes from
oleoresin of ponderosa pine tree Pinus ponderosa: A defense mechanism against microbial
invasion. J Chem Ecol 18(10):1809-1818.
Hughes AL, Nei M. 1992. Maintenance of MHC polymorphism. Nature 335:402–403.
Johnson RM, Harpur BA, Dogantzis KA, Amro Z, Berenbaum MR. 2018. Genomic footprint of
evolution of eusociality in bees: floral food use and CYPome "blooms". Insectes Soc
65(3):445-454.
Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, et al. 2014.
InterProScan 5: genome-scale protein function classification. Bioinformatics 30(9):1236-1240.
Jones WD, Cayirlioglu P, Kadow IG, Vosshall LB. 2007. Two chemosensory receptors together
mediate carbon dioxide detection in Drosophila. Nature 445(7123):86-90.
Jurka J, Kapitonov VV, Pavlicek A, Klonowski P, Kohany O, Walichiewicz J. 2005. Repbase
Update, a database of eukaryotic repetitive elements. Cytogenet Genome Res 110(1-4):462-
467.
Karev GP, Wolf YI, Rzhetsky AY, Berezovskaya FS, Koonin EV. 2002. Birth and death of
protein domains: a simple model of evolution explains power law behavior. BMC Evol Biol
2(1):18.
Keeling CI, Yuen MMS, Liao NY, Docking TR, Chan SK, Taylor GA, Palmquist DL, et al.
2013. Draft genome of the mountain pine beetle, Dendroctonus ponderosae Hopkins, a major
forest pest. Genome Biol 14(3):R27.
Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F,
et al. 2007. Clustal W and Clustal X version 2.0. Bioinformatics 23(21):2947-2948.
LeBoeuf AC, Benton R, Keller L. 2013. The molecular basis of social behavior: models, methods
and advances. Curr Opin Neurobiol 23(1):3-10.
Lee K-A, Kim B, Bhin J, Kim DH, You H, Kim E-K, Kim S-H, Ryu J-H, Hwang D, Lee W-J.
2015. Bacterial uracil modulates Drosophila DUOX-dependent gut immunity via Hedgehog-
induced signaling endosomes. Cell Host Microbe 17(2):191-204.
Lee Y, Moon SJ, Montell C. 2009. Multiple gustatory receptors required for the caffeine response
in Drosophila. Proc Natl Acad Sci USA 106(11):4495-4500.
Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinform 12(1):323.
Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics 25(14):1754-1760.
Li R, Yu C, Li Y, Lam T-W, Yiu S-M, Kristiansen K, Wang J. 2009. SOAP2: an improved
ultrafast tool for short read alignment. Bioinformatics 25(15):1966-1967.
Li W, Schuler MA, Berenbaum MR. 2003. Diversification of furanocoumarin-metabolizing
cytochrome P450 monooxygenases in two papilionids: specificity and substrate encounter
rate. Proc Natl Acad Sci USA 100(suppl 2):14593-14598.
Li X, Schuler MA, Berenbaum MR. 2007. Molecular mechanisms of metabolic resistance to
synthetic and natural xenobiotics. Annu Rev Entomol 52:231-253.
Linnen CR, O'Quin CT, Shackleford T, Sears CR, Lindstedt C. 2018. Genetic basis of body
color and spotting pattern in redheaded pine sawfly larvae (Neodiprion lecontei). Genetics
209(1):291-305.

Marçais G, Kingsford C. 2011. A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. Bioinformatics 27(6):764-770.
Martin A, Orgogozo V. 2013. The loci of repeated evolution: a catalog of genetic hotspots of
phenotypic variation. Evolution 67(5):1235-1250.
Matsuo T, Sugaya S, Yasukawa J, Aigaki T, Fuyama Y. 2007. Odorant-binding proteins
OBP57d and OBP57e affect taste perception and host-plant preference in Drosophila
sechellia. PLoS Biology 5(5):e118.
Maxwell DE editor. Proceedings of the 10th International Congress of Entomology. 1958
Montreal, Canada.
McBride CS. 2007. Rapid evolution of smell and taste receptor genes during host specialization
in Drosophila sechellia. Proc Natl Acad Sci USA 104(12):4996-5001.
McBride CS, Arguello RJ. 2007. Five Drosophila genomes reveal nonneutral evolution and the
signature of host specialization in the chemoreceptor superfamily. Genetics 177(3):1395-1416.
McKenna DD, Scully ED, Pauchet Y, Hoover K, Kirsch R, Geib SM, Mitchell RF, et al. 2016.
Genome of the Asian longhorned beetle (Anoplophora glabripennis), a globally significant
invasive species, reveals key functional and evolutionary innovations at the beetle–plant
interface. Genome Biol 17:227
McKenzie SK, Fetter-Pruneda I, Ruta V, Kronauer DJC. 2016. Transcriptomics and
neuroanatomy of the clonal raider ant implicate an expanded clade of odorant receptors in
chemical communication. Proc Natl Acad Sci USA 113(49):14091-14096.
Mittapelly P, Bansal R, Michel A. 2019. Differential expression of cytochrome P450 CYP6 genes
in the brown marmorated stink bug, Halyomorpha halys (Hemiptera: Pentatomidae). J Econ
Entomol 112(3):1403-1410.
Miyamoto T, Slone J, Song X, Amrein H. 2012. A fructose receptor functions as a nutrient
sensor in the Drosophila brain. Cell 151(5):1113-1125.
Moon SJ, Köttgen M, Jiao Y, Xu H, Montell C. 2006. A taste receptor required for the caffeine
response in vivo. Curr Biol 16(18):1812-1817.
Moon SJ, Lee Y, Jiao Y, Montell C. 2009. A Drosophila gustatory receptor essential for aversive
taste and inhibiting male-to-male courtship. Curr Biol 19(19):1623-1627.
Mumm R, Hilker M. 2006. Direct and indirect chemical defence of pine against folivorous
insects. Trends Plant Sci 11(7):351-358.
Nei M. 2007. The new mutation theory of phenotypic evolution. Proc Natl Acad Sci USA
104(30):12235-12242.
Nei M, Rooney AP. 2005. Concerted and birth-and-death evolution of multigene families. Annu
Rev Genet 39:121-152.
Niehuis O, Gibson JD, Rosenberg MS, Pannebakker BA, Koevoets T, Judson AK, Desjardins
CA, et al. 2010. Recombination and its impact on the genome of the haplodiploid parasitoid
wasp Nasonia. PLoS One 5(1):e8597.
Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C. 2005. Genomic scans
for selective sweeps using SNP data. Genome Res 15(11):1566-1575.
Nozawa M, Kawahara Y, Nei M. 2007. Genomic drift and copy number variation of sensory
receptor genes in humans. Proc Natl Acad Sci USA 104(51):20421-20426.
Ohno S. 1970. The enormous diversity in genome sizes of fish as a reflection of nature's
extensive experiments with gene duplication. Trans Am Fish Soc 99(1):120-130.

Parra G, Bradnam K, Korf I. 2007. CEGMA: a pipeline to accurately annotate core genes in
eukaryotic genomes. Bioinformatics 23(9):1061-1067.
Perry GH, Dominy NJ, Claw KG, Lee AS, Fiegler H, Redon R, Werner J, et al. 2007. Diet and
the evolution of human amylase gene copy number variation. Nat Genet 39(10):1256-1260.
Peters RS, Krogmann L, Mayer C, Donath A, Gunkel S, Meusemann K, Kozlov A, et al. 2017.
Evolutionary history of the Hymenoptera. Curr Biol 27(7):1013-1018.
Petrov DA, Hartl DL. 1997. Trash DNA is what gets thrown away: high rate of DNA loss in
Drosophila. Gene 205(1-2):279-289.
Petrov DA, Hartl DL. 1998. High rate of DNA loss in the Drosophila melanogaster and
Drosophila virilis species groups. Mol Biol Evol 15(3):293-302.
Petrov DA, Lozovskaya ER, Hartl DL. 1996. High intrinsic rate of DNA loss in Drosophila.
Nature 384(6607):346-349.
Poelchau M, Childers C, Moore G, Tsavatapalli V, Evans J, Lee C-Y, Lin H, Lin J-W, Hacket
K. 2015. The i5k Workspace@ NAL—enabling genomic data access, visualization and
curation of arthropod genomes. Nucleic Acids Res 43(D1):D714-D719.
Protas ME, Hersey C, Kochanek D, Zhou Y, Wilkens H, Jeffery WR, Zon LI, Borowsky R,
Tabin CJ. 2006. Genetic analysis of cavefish reveals molecular convergence in the evolution
of albinism. Nat Genet 38(1):107-111.
Qiu Y, Tittiger C, Wicker-Thomas C, Le Goff G, Young S, Wajnberg E, Fricaux T, Taquet N,
Blomquist GJ, Feyereisen R. 2012. An insect-specific P450 oxidative decarbonylase for
cuticular hydrocarbon biosynthesis. Proc Natl Acad Sci USA 109(37):14858-14863.
Qu Z, Kenny NJ, Lam HM, Chan TF, Chu KH, Bendena WG, Tobe SS, Hui JHL. 2015. How
did arthropod sesquiterpenoids and ecdysteroids arise? Comparison of hormonal pathway
genes in noninsect arthropod genomes. Genome Biol Evol 7(7):1951-1959.
Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26(6):841-842.
R-Core-Team. 2018. R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing.
Ratzka C, Förster F, Liang C, Kupper M, Dandekar T, Feldhaar H, Gross R. 2012. Molecular
characterization of antimicrobial peptide genes of the carpenter ant Camponotus floridanus.
PLoS One 7(8):e43036.
Rewitz KF, O'Connor MB, Gilbert LI. 2007. Molecular evolution of the insect Halloween family
of cytochrome P450s: phylogeny, gene organization and functional conservation. Insect
Biochem Mol Biol 37(8):741-753.
Rizzon C, Ponger L, Gaut BS. 2006. Striking similarities in the genomic distribution of
tandemly arrayed genes in Arabidopsis and rice. PLoS Comput Biol 2(9):e115.
Robertson HM, Gadau J, Wanner KW. 2010. The insect chemoreceptor superfamily of the
parasitoid jewel wasp Nasonia vitripennis. Insect Mol Biol 19:121-136.
Robertson HM, Kent LB. 2009. Evolution of the gene lineage encoding the carbon dioxide
receptor in insects. J Insect Sci 9(1):19.
Robertson HM, Wanner KW. 2006. The chemoreceptor superfamily in the honey bee, Apis
mellifera: expansion of the odorant, but not gustatory, receptor family. Genome Res
16(11):1395-1403.
Robertson HM, Waterhouse RM, Walden KKO, Ruzzante L, Reijnders MJMF, Coates BS,
Legeai F, et al. 2018. Genome sequence of the wheat stem sawfly, Cephus cinctus,

representing an early-branching lineage of the Hymenoptera, illuminates evolution of
hymenopteran chemoreceptors. Genome Biol Evol 10(11):2997-3011.
Ross L, Gardner A, Hardy N, West SA. 2013. Ecology, not the genetics of sex determination,
determines who helps in eusocial populations. Curr Biol 23(23):2383-2387.
Roux J, Privman E, Moretti S, Daub JT, Robinson-Rechavi M, Keller L. 2014. Patterns of
positive selection in seven ant genomes. Mol Biol Evol 31(7):1661-1685.
Sadd BM, Barribeau SM, Bloch G, de Graaf DC, Dearden P, Elsik CG, Gadau J, et al. 2015.
The genomes of two key bumblebee species with primitive eusocial organization. Genome
Biol 16(1):1-32.
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, et al.
2012. Fiji: an open-source platform for biological-image analysis. Nature Methods 9(7):676-
682.
Sezutsu H, Le Goff G, Feyereisen R. 2013. Origins of P450 diversity. Philos Trans R Soc Lond B
Biol Sci 368:20120428.
Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics 31(19):3210-3212.
Simone M, Evans JD, Spivak M. 2009. Resin collection and social immunity in honey bees.
Evolution 63(11):3016-3022.
Singaravelan N, Nee'man G, Inbar M, Izhaki I. 2005. Feeding responses of free-flying honeybees
to secondary compounds mimicking floral nectars. J Chem Ecol 31(12):2791-2804.
Slone J, Daniels J, Amrein H. 2007. Sugar receptors in Drosophila. Curr Biol 17(20):1809-1816.
Smadja C, Shi P, Butlin RK, Robertson HM. 2009. Large gene family expansions and adaptive
evolution for odorant and gustatory receptors in the pea aphid, Acyrthosiphon pisum. Mol
Biol Evol 26(9):2073-2086.
Smit AFA, Hubley R. 2008-2015. RepeatModeler Open-1.0.
Smit AFA, Hubley R, Green P. 2013-2015. RepeatMasker Open-4.0.
Smith CR, Smith CD, Robertson HM, Helmkampf M, Zimin A, Yandell M, Holt C, et al. 2011.
Draft genome of the red harvester ant Pogonomyrmex barbatus. Proc Natl Acad Sci USA
108(14):5667-5672.
Smith SG. 1941. A new form of spruce sawfly identified by means of its cytology and
parthenogenesis. J Agric Sci 21(5):245-305.
Sohi SS, Ennis TJ. 1981. Chromosomal characterization of cell lines of Neodiprion lecontei
(Hymenoptera: Diprionidae). Proc Entomol Soc Ont 112:45–48.
Souvorov A, Kapustin Y, Kiryutin B, Chetvernin V, Tatusova T, Lipman DJ. 2010. Gnomon–
NCBI eukaryotic gene prediction tool. National Center for Biotechnology Information.
Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30(9):1312-1313.
Suzuki HC, Ozaki K, Makino T, Uchiyama H, Yajima S, Kawata M. 2018. Evolution of
gustatory receptor gene family provides insights into adaptation to diverse host plants in
nymphalid butterflies. Genome Biol Evol 10(6):1351-1362.
Thomas JH. 2006. Analysis of homologous gene clusters in Caenorhabditis elegans reveals
striking regional cluster domains. Genetics 172(1):127-143.
Tian C, Gao B, Fang Q, Ye G, Zhu S. 2010. Antimicrobial peptide-like genes in Nasonia
vitripennis: a genomic perspective. BMC Genom 11(1):187.

Trapp S, Croteau R. 2001. Defensive resin biosynthesis in conifers. Annu Rev Plant Biol
52(1):689-724.
Vieira FG, Sánchez-Gracia A, Rozas J. 2007. Comparative genomic analysis of the odorant-
binding protein family in 12 Drosophila genomes: purifying selection and birth-and-death
evolution. Genome Biol 8(11):R235.
Vitti JJ, Grossman SR, Sabeti PC. 2013. Detecting natural selection in genomic data. Annu Rev
Genet 4797-120.
Vosshall LB, Wong AM, Axel R. 2000. An olfactory sensory map in the fly brain. Cell
102(2):147-159.
Wanner KW, Robertson HM. 2008. The gustatory receptor family in the silkworm moth
Bombyx mori is characterized by a large expansion of a single lineage of putative bitter
receptors. Insect Mol Biol 17(6):621-629.
Wiens JJ, Lapoint RT, Whiteman NK. 2015. Herbivory increases diversification across insect
clades. Nat Commun 6(1):1-7.
Wilson LF, Wilkinson RC, C AR. 1992. Redheaded pine sawfly: its ecology and management.
In: Service USDoAF, editor. Agriculture Handbook. Washington, DC.
Xu Y-L, He P, Zhang L, Fang S-Q, Dong S-L, Zhang Y-J, Li F. 2009. Large-scale identification
of odorant-binding proteins and chemosensory proteins from expressed sequence tags in
insects. BMC Genom 10(1):632.
Yang J, Chen X, Bai J, Fang D, Qiu Y, Jiang W, Yuan H, et al. 2016. The Sinocyclocheilus
cavefish genome provides insights into cave adaptation. BMC Biol 14(1):1-13.
Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol
24(8):1586-1591.
Zhang J. 2003. Evolution by gene duplication: an update. Trends Ecol Evol 18(6):292-298.
Zhang L, Gaut BS. 2003. Does recombination shape the distribution and evolution of tandemly
arrayed genes (TAGs) in the Arabidopsis thaliana genome? Genome Res 13(12):2533-2540.
Zhang Z, Zhu S. 2012. Comparative genomics analysis of five families of antimicrobial peptide-
like genes in seven ant species. Dev Comp Immunol 38(2):262-274.
Zhen Y, Aardema ML, Medina EM, Schumer M, Andolfatto P. 2012. Parallel molecular
evolution in an herbivore community. Science 337(6102):1634-1637.
Zhou X, Rokas A, Berger SL, Liebig J, Ray A, Zwiebel LJ. 2015. Chemoreceptor evolution in
hymenoptera and its implications for the evolution of eusociality. Genome Biol Evol
7(8):2407-2416.

1376

Figure Legends

1378

Figure 1. Like other diprionids, *N. lecontei* has multiple morphological and behavioral
adaptations to *Pinus* foliage. A. An egg-laying *N. lecontei* female demonstrating several
adaptations for dealing with thick, resinous pine needles, including: a robust saw-like ovipositor
(visible within the needle), a tendency to lay many closely spaced eggs per needles, and a
tendency to cut resin-draining slits on egg-bearing needles (circled). **B.** Prior to hatching, *N.*
lecontei eggs absorb water from the host, causing the eggs to swell and the pockets to open.
Throughout development, embryos are in close contact with living host tissue. **C.** Early-instar
larvae have skeletonizing feeding behavior in which only the outer needle tissue is consumed,
leaving the resinous interior intact. This strategy prevents small larvae from being overwhelmed

by sticky resin. **D.** Mid- and late-instar larvae consume the entire pine needle. Larvae sequester
pine resin in specialized pouches for use in self-defense (All photos by R.K. Bagley).
1390
1391   **Figure 2. Optical sections through the antennal lobes of adult female (left) and male (right)**
1392   *N. lecontei.* White arrows indicate regions of male-specific synaptic clusters. Scale bars = 500
1393   µm.
1394
Figure 3. Position of genes belonging to five environmentally responsive gene families along
seven *N. lecontei* linkage groups. Linkage groups (LG) are drawn to scale and ordered as in the
linkage-group anchored assembly described in Linnen et al. 2018 (GenBank accession numbers
are as follows: LG1 = CM009916.1; LG2 = CM009917.1; LG3 = CM009918.1; LG4 =
CM009919.1; LG5 = CM009920.1; LG6 = CM009921.1; LG7 = CM009922.1). Gene family
abbreviations: OR (olfactory receptor), GR (gustatory receptor), OBP (odorant binding protein),
CYP (cytochrome P450), AMP (antimicrobial protein). Each gene family is represented by a
different color. Horizontal lines indicate the approximate locations of genes within LG; diagonal
lines that connect to horizontal lines are used to highlight groups of genes that met our clustering
criteria. Genes that were found on scaffolds that have not been placed on linkage groups are
indicated on the bottom left, with abbreviated scaffold names given in parentheses (e.g., S-210 =
scaffold_210 = LGIB01000210.1 in the assemblies available on NCBI).
1407
1408   **Figure 4. Number of intact genes in hymenopteran genomes for each of five**
1409   **environmentally responsive gene families.** Phylogenetic relationships are as in Moreau et al.
1410   (2006); Hedtke et al. (2013); Roux et al. (2014); Brand et al. (2017); Branstetter et al. (2017);
1411   Peters et al. (2017). Branch lengths are arbitrary. Gene family abbreviations are as in Figure 3.
1412
Figure 5. Ecological correlates of gene family size in Hymenoptera. Each point represents the
number of intact genes for a hymenopteran species for which both manually curated gene
annotations and ecological data are available. Asterisks indicate that gene number varies
significantly among the ecological categories under consideration; for significant categories with
>2 groups, letters indicate significance in post-hoc tests (groups that do not share a letter are
significantly different). Gene number and ecological data for all taxa are provided in Table S8.
1419

1420 **Table 1. Summary of within-genome signatures of adaptive expansions and contractions**
1421 **for five environmentally responsive gene families.**

Gene family[*]	Gene family size					Genomic Clustering		Molecular evolution		
	Intact genes	Partial	Pseudo	Total genes	Prop. pseudo	Prop. in clusters[†]	Largest cluster	*Neodiprion*-specific clades[‡]	Significant branch tests[§]	Significant site tests[**]
OR	52	3	1	56	0.02	0.59	8	3	1	0
GR	41	2	2	44[††]	0.05	0.76	10	3	1	1
OBP	13	0	0	13	0	0.38	3	0	n/a	n/a
CYP (all)	93	2	12	107	0.11	0.66	16	5	2	0
CYP2 clan	9	0	0	9	0	0.33	2	0	0	0
CYP3 clan	47	0	8	55	0.15	0.81	16	4	2	0
CYP4 clan	27	2	4	33	0.12	0.55	3	1	0	0
mito CYP clan	10	0	0	10	0	0.50	3	0	0	0
AMP	21	0	0	21	0	0.95	15	?[‡‡]	0	0

1422

[*] Abbreviations: OR = olfactory receptor genes; GR = gustatory receptor genes; OBP = odorant binding protein genes; CYP = cytochrome P450 genes ("clans" refer to four major clades of CYPs present in insects); AMP = antimicrobial peptide genes.
[†] Calculated as: (number of genes in clusters of 2 or more)/(genes for which clustering could be evaluated).
[‡] Defined as monophyletic clusters of 5 or more *Neodiprion* paralogs with a bootstrap support $\geq$ 70% in an amino acid phylogeny constructed with gene annotations from *Neodiprion*, select Hymenoptera, and *Drosophila melanogaster*.
[§] To be counted, clades had to reject both 1-ratio and fixed-ratio models in dN/dS branch tests (see Table 2).
[**] To be counted, clades had to reject both M7 and M8a models in dN/dS site tests (see Table 2).
[††] One gene was both a partial annotation and a pseudogene.
[‡‡] Low bootstrap support precluded the identification of *Neodiprion*-specific clades.

Table 2. Likelihood-ratio tests (LRTs) of positive selection on *Neodiprion*-specific clades (branch models) and on amino acid sites within these clades (site models).

Clade Names[*]	n[†]	Model comparison[‡]	LRT statistic[§]	df	P-value[**]
Olfactory Receptor					
Clade 1	6	M8 vs M7	2.932	2	0.231
		M8 vs M8a	0.748	1	0.387
		2 ratio vs 1 ratio	5.408	1	**0.020**
		2 ratio vs neutral	7.800	1	**0.005**
Clade 2	5	M8 vs M7	3.941	2	0.139
		M8 vs M8a	1.525	1	0.217
		2 ratio vs 1 ratio	1.426	1	0.232
		2 ratio vs neutral	0.050	1	0.822
Clade 3	5	M8 vs M7	0	2	1
		M8 vs M8a	0	1	1
		2 ratio vs 1 ratio	2.395	1	0.122
		2 ratio vs neutral	6.371	1	**0.012**
Gustatory Receptor					
Clade 1	7	M8 vs M7	0.809	2	0.667
		M8 vs M8a	0.379	1	0.538
		2 ratio vs 1 ratio	0.003	1	0.954
		2 ratio vs neutral	0.645	1	0.422
Clade 2	8	M8 vs M7	6.049	2	**0.049**
		M8 vs M8a	2.654	1	0.103
		2 ratio vs 1 ratio	0.003	1	0.959
		2 ratio vs neutral	0.781	1	0.377
Clade 3	5	M8 vs M7	39.328	2	**2.884e-09**
		M8 vs M8a	35.167	1	**3.026e-09**
		2 ratio vs 1 ratio	14.789	1	**1.202e-04**
		2 ratio vs 2 ratio neutral	27.810	1	**1.338e-07**
Cytochrome P450					
Clade 1 (CYP4 clan)	8	M8 vs M7	0.615	2	0.735
		M8 vs M8a	0.866	1	0.352
		2 ratio vs 1 ratio	0.658	1	0.417
		2 ratio vs 2 ratio neutral	0.089	1	0.766
Clade 2 (CYP3 clan)	19	M8 vs M7	0	2	1
		M8 vs M8a	0	1	1

[*] Clade names are as in Figures S1a,S1b,S2a,S2b,S3a,S3b,S4a,S4b, and S5a.

[†] Putatively functional genes. Pseudogenes and partial annotations were excluded from analysis.

[‡] Site models unshaded; neutral M7 and M8a do not allow for positive selection. Branch models shaded; 1 ratio estimates a single ω value for all branches, 2 ratio estimates a separate ω value for the foreground branch, 2 ratio neutral fixes $\omega=1$ for all branches.

[§] Likelihood ratio test statistic, calculated twice the difference in model log likelihoods.

[**] Bolded values are significant at critical value 0.05.

		2 ratio vs 1 ratio	2.077	1	0.149
		2 ratio vs 2 ratio neutral	0.076	1	0.783
Clade 3 (CYP3 clan)	6	M8 vs M7	7.152	2	**0.028**
		M8 vs M8a	0.649	1	0.421
		2 ratio vs 1 ratio	6.325	1	**0.012**
		2 ratio vs 2 ratio neutral	14.261	1	**1.59 e-04**
Clade 4 (CYP3 clan)	6	M8 vs M7	0	2	1
		M8 vs M8a	0.151	1	0.697
		2 ratio vs 1 ratio	0.002	1	0.964
		2 ratio vs 2 ratio neutral	0.936	1	0.333
Clade 5 (CYP3 clan)	5	M8 vs M7	0	2	1
		M8 vs M8a	0	1	1
		2 ratio vs 1 ratio	5.327	1	**0.021**
		2 ratio vs 2 ratio neutral	12.286	1	**4.56 e-04**
Hisnavicin (Antimicrobial Peptide)					
Clade 1[††]	15	M8 vs M7	2.388	2	0.665
		M8 vs M8a	0	1	1
		2 ratio vs 1 ratio	7.908	1	**0.010**
		2 ratio vs 2 ratio neutral	0.999	1	0.635

1429
1430

[††] Although this clade did not meet the bootstrap criteria for species-specific clades (>70), it was included in this analysis because it contained almost all *N. lecontei* hisnavicin paralogs.

1431
1432
Figure 1. Like other diprionids, *N. lecontei* has multiple morphological and behavioral
adaptations to *Pinus* foliage. A. An egg-laying *N. lecontei* female demonstrating several
adaptations for dealing with thick, resinous pine needles, including: a robust saw-like ovipositor
(visible within the needle), a tendency to lay many closely spaced eggs per needles, and a
tendency to cut resin-draining slits on egg-bearing needles (circled). **B.** Prior to hatching, *N.*
lecontei eggs absorb water from the host, causing the eggs to swell and the pockets to open.
Throughout development, embryos are in close contact with living host tissue. **C.** Early-instar
larvae have skeletonizing feeding behavior in which only the outer needle tissue is consumed,
leaving the resinous interior intact. This strategy prevents small larvae from being overwhelmed
by sticky resin. **D.** Mid- and late-instar larvae consume the entire pine needle. Larvae sequester
pine resin in specialized pouches for use in self-defense (All photos by R.K. Bagley).

1449
1450
1451 **Figure 2. Optical sections through the antennal lobes of adult female (left) and male (right)**
1452 ***N. lecontei.*** White arrows indicate regions of male-specific synaptic clusters. Scale bars = 500
1453 μm.
1454

1455
Figure 3. Position of genes belonging to five environmentally responsive gene families along seven *N. lecontei*
linkage groups. Linkage groups (LG) are drawn to scale and ordered as in the linkage-group anchored assembly
described in Linnen et al. 2018 (GenBank accession numbers are as follows: LG1 = CM009916.1; LG2 =
CM009917.1; LG3 = CM009918.1; LG4 = CM009919.1; LG5 = CM009920.1; LG6 = CM009921.1; LG7 =
CM009922.1). Gene family abbreviations: OR (olfactory receptor), GR (gustatory receptor), OBP (odorant binding
protein), CYP (cytochrome P450), AMP (antimicrobial protein). Each gene family is represented by a different
color. Horizontal lines indicate the approximate locations of genes within LG; diagonal lines that connect to
horizontal lines are used to highlight groups of genes that met our clustering criteria. Genes that were found on
scaffolds that have not been placed on linkage groups are indicated on the bottom left, with abbreviated scaffold
names given in parentheses (e.g., S-210 = scaffold_210 = LGIB01000210.1 in the assemblies available on NCBI).

	OR	GR	OBP	CYP	AMP
Bee					
Bombus terrestris	151	21	16	44	
Melipona quadrifasciata	142	10	6		
Apis mellifera	169	10	21	46	6
Apis cerana	119	10		41	
Apis dorsata				42	
Apis florea	159			44	
Habropoda laboriosa	100			38	
Euglossa dilemma	123	13	15		
Eufriesea mexicana	111	16	13	45	
Megachile rotundata				49	
Osmia bicornis bicornis				47	
Lasioglossum albipes	158	23			
Dufourea novaeangliae	77			45	
Ant					
Solenopsis invicta	333	219	18		7
Cardiocondyla obscurior	232	34			
Monomorium pharaonis	240	159			
Pogonomyrmex barbatus	274	58	16	72	10
Acromyrmex echinatior	375	116		73	
Atta cephalotes	341	89		52	
Camponotus floridanus	352	46	13	128	6
Linepithema humile	301	93	13	111	6
Cerapachys biroi	256	20	15	69	6
Harpegnathos saltator	347	17	13	95	8
Wasp					
Nasonia vitripennis	217	47	82	92	44
Ceratosolen solmsi	56	5	7	34	8
Microplitis demolitor	203	79			
Sawfly					
Neodiprion lecontei	52	41	13	94	11

Figure 4. Number of intact genes in hymenopteran genomes for each of five environmentally responsive gene families. Phylogenetic relationships are as in Moreau et al. (2006); Hedtke et al. (2013); Roux et al. (2014); Brand et al. (2017); Branstetter et al. (2017); Peters et al. (2017). Branch lengths are arbitrary. Gene family abbreviations are as in Figure 3.

1473
Figure 5. Ecological correlates of gene family size in Hymenoptera. Each point represents the number of intact genes for a hymenopteran species for which both manually curated gene annotations and ecological data are available. Asterisks indicate that gene number varies significantly among the ecological categories under consideration; for significant categories with >2 groups, letters indicate significance in post-hoc tests (groups that do not share a letter are significantly different). Gene number and ecological data for all taxa are provided in Table S8.
1480