The virucidal effects of 405 nm visible light on SARS-CoV-2 and influenza A virus

Raveen Rathnasinghe[1,2,3], Sonia Jangra[1,2], Lisa Miorin[1,2], Michael Schotsasert[1,2], Clifford

Yahnke[6#], Adolfo García-Sastre[1,2,4,5#1]

[1]Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
[2]Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
[3]Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
[4]Department of Medicine, Division of Infectious Diseases, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
[5]The Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
[6] Kenall Manufacturing, Kenosha, WI 53144
#Correspondence: cliff.yahnke@kenall.com , adolfo.garcia-sastre@mssm.edu

Abstract

Germicidal potential of specific wavelengths within the electromagnetic spectrum is an

area of growing interest. While ultra-violet (UV) based technologies have shown

satisfactory virucidal potential, the photo-toxicity in humans coupled with UV associated

polymer degradation limit its use in occupied spaces. Alternatively, longer wavelengths

with less irradiation energy such as visible light (405 nm) have largely been explored in

the context of bactericidal and fungicidal applications. Such studies indicated that 405

nm mediated inactivation is caused by the absorbance of porphyrins within the

organism creating reactive oxygen species which result in free radical damage to its

DNA and disruption of cellular functions. The virucidal potential of visible-light based

technologies has been largely unexplored and speculated to be ineffective given the

lack of porphyrins in viruses. The current study demonstrated increased susceptibility of

lipid-enveloped respiratory pathogens of importance such as SARS-CoV-2 (causative

agent of COVID-19) as well as the influenza A virus to 405nm, visible light in the

absence of exogenous photosensitizers indicating a potential porphyrin-independent alternative mechanism of visible light mediated viral inactivation. These results were obtained using less than expected irradiance levels which are generally safe for humans and commercially achievable. Our results support further exploration of the use of visible light technology for the application of continuous decontamination in occupied areas within hospitals and/or infectious disease laboratories, specifically for the inactivation of respiratory pathogens such as SARS-CoV-2 and Influenza A.

Key words – Visible light, 405nm, Virucidal, SARS-CoV-2, Influenza, inactivation

Introduction

The severe-acute respiratory syndrome corona virus 2 (SARS-CoV-2), the causative agent of the COVID-19 pandemic, is a member of the beta-coronavirus family and it emerged at the end of 2019 in the Hubei province in Wuhan China[1]. By late February 2021, more than 112 million cases had been reported while accounting for approximately 2.5 million deaths, underscoring the rapid dissemination of the virus on a global scale[2]. As a complement to standard precautions such as handwashing, masking, surface disinfection, and social distancing, other enhancements to enclosed spaces such as improved ventilation and whole-room disinfection are being considered by segments beyond acute healthcare such as retail, dining, and transportation[3].

Initial guidance from health authorities such as the CDC and WHO on environmental transmission focused on contaminated surfaces as fomites[4]. Data pertaining to the survival of SARS-CoV-2 and other related coronaviruses to date has indicated that virions are able to persist on fomites composed of plastic[5], wood[6], paper[5], metal[7] and glass[8] potentially up to nine days. Recent studies have suggested that SARS-CoV-2 may also remain viable approximately at least three days in such surfaces and another two studies showed that at room temperature (20-25°C), a 14-day time-period was required to see a 4.5-5 Log_{10} of the virus[9, 10].

Since the start of the pandemic, transmission of the virus by respiratory droplets and aerosols has become an accepted method of transmission although the relative impact of each mode of transmission is the subject of much debate. Nevertheless, enclosed spaces with groups of people exercising or singing have been associated with

increased transmission. The half-life survival of SARS-CoV-2 in this type of environment

has been estimated between 1-2 hours[6, 11, 12].

Taking this information into consideration, several methods have been evaluated to

effectively inactivate SARS-CoV-2. Chemical methods, which focus on surface

disinfection, utilize 70% alcohol and bleach and their benefits are well established.

These methods are also episodic (or non-continuous) meaning that in-between

applications, the environment is not being treated[13].

In addition to chemicals, one of the most utilized methods for whole-room disinfection is

germicidal ultra-violet C (UVC; ~254 nm)[14]. This technology is well established[15] and

has been shown to inactivate a range of pathogens including bacteria[16], fungi[17] and

viruses[18]. The mechanism of action of UVC is photodimerization of genetic material

such as RNA (relevant for SARS-CoV2 and IAV) and DNA (relevant for DNA viruses

and bacterial pathogens, among others)[19]. Unfortunately, this effect has been

associated with deleterious effects in exposed humans such as photokeratoconunctivitis

in eyes and photodermatitis in skin[20]. For these reasons, UVC irradiation requires safety

precautions and cannot be used to decontaminate fomites and high contact areas in the

presence of humans[21].

Germicidal properties of violet-blue visible light (380-500 nm), especially within the

range of 405 to 450 nm wavelengths have been appreciated as an alternative to UVC

irradiation in whole-room disinfection scenarios where it has shown reduction of

bacteria[22, 23] in occupied rooms and reductions in surgical site infections[24]. Although 405

nm or closely related wavelengths have been shown to be less germicidal than UVC, its

inactivation potential has been assessed in pathogenic bacteria such as *Listeria* spp

and *Clostridium* spp[24, 25], and in fungal species such as *Saccharomyces* spp and

Candida spp[26] . It is thought that the underlying mechanism of blue-light mediated

inactivation is associated with absorption of light via photosensitizers such as

porphyrins which results in the release of reactive oxygen species (ROS) [27, 28]. The

emergence of ROS is associated with direct damage to biomolecules such as proteins,

lipids and nucleic acids which are essential constituents of bacteria, fungi and viruses.

Further studies have shown that ROS can also lead to the loss of cell membrane

permeability mediated by lipid oxidation[29]. Given the lack of endogenous

photosensitizers such as porphyrins in virions, efficient decontamination of viruses (both

enveloped and non-enveloped) may require the addition of exogenous

photosensitizers[23]. With the use of media suspensions containing both endogenous

and/or exogenous photosensitizers, inactivation of viruses such as feline calcivirus

(FCV)[30], viral hemorrhagic septicemia virus (VHSV)[31] and murine norovirus-1[32] has

demonstrated the virucidal potency of 405 nm visible light using porphyrin rich media

such as saliva, blood, etc. (to create the ROS required for inactivation) using unsafe,

commercially impractical irradiance levels. This highlights the importance of answering

the basic scientific questions related to viral inactivation within the context of the applied

science required for clinical application. Our study specifically focused on three

questions: 1) does the same wavelength of light (405nm) that inactivates bacteria also

inactivate enveloped viruses, 2) is a porphyrin rich medium required for this inactivation,

and 3) can this inactivation be achieved using safe, commercially practical irradiance

levels?

In the current study, we show the impact of 405 nm irradiation on inactivation of SARS-

CoV-2 and influenza A H1N1 viruses without the use of photosensitizers making it

directly relevant to the clinical environment. We show this using a commercially

available visible light disinfection system ensuring that the irradiance used is both safe

and practically achievable in a clinical setting supporting the possible use of 405 nm

irradiation as a tool to confer continuous decontamination of respiratory pathogens such

as SARS-CoV-2 and influenza A viruses. We further show the increased susceptibility

of lipid-enveloped viruses for irradiation in comparison to non-enveloped viruses, further

characterizing the virucidal effects of visible light.

Materials and methods.

405 nm Exposure System

The visible light disinfection product used in this study was a commercially available 6"

LED downlight (Indigo-Clean, Kenall, Kenosha, WI) to allow for use within a BSL-3 level

containment hood. Within the hood, the distance between the face of the fixture and the

sample was 10"- much less than the normal 1.5 m used in normal, whole-room

disinfection applications. The output of the fixture was modified electronically during its

manufacture to match this difference and ensure that the measurements would

represent the performance of the device in actual use. This test setup is shown in a

BSL-2 hood in Figure 1 below. Note the spectroradiometer and the bottom portion of the

rig (~6") used for calibration were removed during the actual study.

Figure 1. Test setup shown with spectroradiometer and extension used for

calibration.

Controls were placed outside the test rig but within the BSL hood as shown in Figure 2.

Note that this picture contains the bottom portion of the test rig to highlight the position

of the radiometer

Figure 2. Test setup showing the placement of the control and sample for irradiation. Note that the bottom portion of the test rig was removed during the actual experiment. This ensures the 10" distance used in the study.

For the range of output used in this study, multiple discrete levels were created using pulse width modulation within the LED driver itself. These levels were made to be individually selectable using a simple knob on the attached control module.

As expected, the amount of visible light within the 400nm-420nm bandwidth, measured in $mWcm^{-2}$, is a measurement of the "dose" delivered to the target organism and is used to quantify this relationship similar to that used in UV disinfection applications.

To fully examine this effect, a range of irradiance values were used representing actual

product deployment conditions in occupied rooms. The lowest value (0.035 mWcm^{-2})

represents a single-mode, lower wattage used in general lighting applications while the

highest value (0.6 mWcm^{-2}) represents a dual-mode, higher wattage used in critical care

applications such as an operating room.

The device was placed in a rig to ensure a consistent distance (10") between the fixture

and the samples. The output of the fixture in the test rig was measured using a Stellar-

RAD Radiometer from StellarNet configured to make wavelength and irradiance

measurements from 350nm-1100nm with < 1nm spectral bandwidth using a NIST

traceable calibration. To ensure that the regular white light portion of the illumination

(which is non-disinfecting) was not measured, the measurement was electronically

limited to a 1nm bandwidth over the 400nm-420nm range. The normalized spectral

profile is shown in Fig. 3 below. The absolute value of the measurement was

determined using a NIST traceable calibration as previously described.

Figure 3. Normalized spectral power distribution showing peak irradiance at

405nm.

Samples were irradiated with the range of wavelengths depicted in Figure 3. This was

deliberately done for two reasons: 1) prior work had shown that visible light disinfection

was primarily active at 405nm[33] and 2) to emphasize the applied science associated

with actual clinical use where a virus in the environment could be exposed to both

405nm and regular white light in an occupied room.

To isolate the contribution of 405nm light, the control samples were placed outside the

field of irradiation created by the disinfection product but within the biosafety hood.

Accordingly, these samples were exposed to the overhead lights within the room which

contained virtually no 405nm light (< 0.001 mWcm^{-2}).

In any assessment of viral inactivation, thermal denaturing of the organism is a concern.
Older lighting technologies such as incandescent sources heat a resistive element and
were widely used in a variety of applications. This creates heat at both the source and
to objects within its field. Fortunately, the disinfecting light (sample) and the overhead
lights in the room (control) did not use this technology and therefore contain no infrared
emissions (> 800 nm) a commonly known benefit associated with LED lighting. As a
confirmation, the temperature beneath the disinfecting light was measured using a
commercially available thermocouple during a 24-hour period. During this time, the
temperature at the sample position was constant at 20°C +/- 0.5°C. even at the highest
disinfecting power (0.6 mWcm^{-2}).

For reference, two spectral profiles, one for traditional fluorescent lighting and the other
for standard LED lighting are provided in Figure 4 below. While both types of lighting
are commonly used in the overhead lights within buildings, the lights in the BSL-3
laboratory were traditional fluorescent. As shown in Figure 2, the controls were
exposed only to traditional fluorescent lighting with a negligible amount of disinfecting
light (< 0.001 mWcm^{-2}) between 400nm and 420nm. Due to the inherent differences
between fluorescent and LED lighting, the standard LED spectra has a small, but
measurable amount of disinfecting light (0.006 mWcm-2) between 400nm and 420nm.
As will be later shown, this amount of light can have a measurable disinfecting effect.

Figure 4. Normalized spectral power distribution for the fluorescent control light

(non-disinfecting) and standard LED light (without 405nm) used in the study.

Each spectrum is normalized relative to its own peak value.

Cells and viruses

Vero-E6 cells (ATCC® CRL-1586™, clone E6) were maintained in Dulbecco's Modified

Eagle Medium (DMEM) complemented with 10% heat-inactivated Fetal Bovine Serum

(HI-FBS; PEAK serum), penicillin-streptomycin (Gibco; 15140-122), HEPES buffer

(Gibco; 15630-080) and MEM non-essential amino-acids (Gibco; 25025CL) at 37°C with

5% CO_2. Vero-CCL81 (ATCC® CRL-81™) cells and MDCK cells (ATCC® CCL-34)

were cultured in DMEM supplemented with 10% HI-FBS and penicillin/streptomycin at -

37ºC with 5% CO_2. All experiments involving SARS-CoV2 (USA-WA1/202, BEI

resource – NR52281) were conducted within a biosafety-level 3 (BSL3) containment

facility at Icahn school of medicine at Mount Sinai by trained workers upon authorization

of protocols by a biosafety committee. Amplification of SARS-CoV-2 viral stocks was

done in Vero-E6 cell confluent monolayers by using an infection medium composed of

DMEM supplemented with 2% HI-FBS, Non-essential amino acids (NEAA), Hepes and

penicillin-streptomycin at 37°C with 5% CO_2 for 72 hours. Influenza A virus used here

was generated using plasmid based reverse genetics system as previously described[34].

The backbone used in the study was A/Puerto Rico/8/34/Mount Sinai(H1N1) under the

GenBank accession number AF389122. IAV-PR8 virus was grown and titrated in MDCK

as previously described[34]. As a non-enveloped virus, the cell culture adapted murine

Encephalomyocarditis virus (EMCV; ATCC® VR-12B) was propagated and titrated in

Vero-CCL81 cells with DMEM and 2% HI-FBS and penicillin-streptomycin at 37°C with

5% CO_2 for 48 hours[35].

405nm inactivation of viruses

The SARS-CoV-2 virus was exclusively handled at the Icahn school of Medicine BSL-3

and studies involving IAV and EMCV were handled in BSL-2 conditions. Indicated PFU

amounts were mixed with sterile 1X PBS and were irradiated in 96 well format cell

culture plates in triplicates. In these studies, A starting dose of $5x10^5$ PFU for SARS-

CoV-2 and starting doses of $1x10^5$ PFU for IAV and EMCV were used. The final

volumes for inactivation were 250 µl per replicate. The untreated samples were

prepared the same way and were left inside the biosafety cabinet isolated from the

inactivation device at room temperature. The plates were sealed with qPCR plate

transparent seal and an approximate 10% reduction of the intensity was observed due

to the sealing film. The distance from the lamp and the samples was measured to be

10". All samples were extracted at indicated times and were frozen at -80°C and were

thawed together for titration via plaque assays.

Plaque assays

For SARS-CoV-2 studies, confluent monolayers of Vero-E6 cells in 12-well plate format

were infected (with an inoculum volume 150μl) with 10-fold serially diluted samples in

1X phosphate-buffered saline (PBS) supplemented with bovine serum albumin (BSA)

and penicillin-streptomycin for an hour while gently shaking the plates every 15 minutes.

Afterwards, the inoculum was removed, and the cells were incubated with an overlay

composed of MEM with 2% FBS and 0.05% Oxoid agar for 72 hours at 37°C with 5%

CO_2. The plates were subsequently fixed using 10% formaldehyde overnight and the

formaldehyde was removed along with the overlay. Fixed monolayers were blocked with

5% milk in Tris-buffered saline with 0.1% tween-20 (TBS-T) for an hour. Afterwards,

plates were immunostained using a monoclonal antibody against SARS-CoV2

nucleoprotein (Creative-Biolabs; NP1C7C7) at a dilution of 1:1000 followed by 1:5000

anti-mouse IgG monoclonal antibody and was developed using KPL TrueBlue

peroxidase substrate for 10 minutes (Seracare; 5510-0030). After washing the plates

with distilled water, the number of a plaques were counted. Plaque assays for IAV and

EMCV were done in a similar fashion. For IAV, confluent monolayers of MDCK cells

supplemented with MEM-based overlay with TPCK-treated trypsin was used and was

incubated for 48 hours at 37°C with 5% CO_2. For EMCV, Vero-CCL81 cells were used

to do plaque assays in 6 well plate format with an inoculum volume of 200μl and was

incubated for 48 hours at 37°C with 5% CO_2. Plaques for IAV and EMCV were

visualized using crystal violet. Data shown here is derived from three independent

experimental setups.

Results.

Dose and time dependent inactivation of SARS-CoV-2 in the absence of

photosensitizers.

The lowest irradiation dose of 0.035 mWcm^{-2} was applied for SARS-CoV-2 and when

compared to the T$_{4H}$ untreated control, a reduction of 0.3288 log$_{10}$ was seen as early as

4 hours and after 24 hours of irradiation, an inactivation of 1.0325 log$_{10}$ (approximately

10 times reduction in infectivity) was observed for SARS-CoV-2 via plaque assays

(Figure 5A). A slightly higher dose of 0.076mWcm^{-2} yielded 0.4123 log$_{10}$, 0.6118 log$_{10}$

and 1.5393 log$_{10}$ reduction by 4, 8 and 12 hours post irradiation when compared to the

respective untreated controls (Figure 5B). Subsequent increase of the irradiation dose

to 0.150 mWcm^{-2} resulted in a reduction of 0.4771 log$_{10}$ after 4 hours which then had a

1.1206 log$_{10}$ after 12 hours. Irradiation for 24 hours at 0.150 mWcm^{-2} suggested a total

reduction of 2.0056 log$_{10}$ (256 times) for SARS-CoV-2 and (Figure 5C). As a final

experiment, a high irradiation dose of 0.6 mWcm^{-2} was used to assess the inactivation

potential within a much shorter time frame. Irradiation for one hour resulted in a

reduction of 0.4150 log$_{10}$ which reached 1.2943 log$_{10}$ reduction by four hours and 2.309

log$_{10}$ (385 times) after 8 hours in comparison to untreated controls samples at the

corresponding times. (Figure 5 D and E). All experimental conditions demonstrated the

stability of untreated SARS-CoV-2 which was left at room temperature in PBS, as

shown by the marginal reduction of viral titer over time.

Figure 5. Dose and time dependent inactivation of SARS-CoV-2 virus in PBS by 405 nm irradiation. A. A dose of 0.035 mWcm^{-2} or **B.** a dose of 0.076 mWcm^{-2} or **C.** a dose of 0.150 mWcm^{-2} or **D.** a dose of 0.6 mWcm^{-2} was applied to irradiate samples at 405 nm over a course of 24 while sampling at 4, 8, 12 and 24 hours (for A, B and C) or over a course of 8 hours while sampling at 1, 2, 4 and 8 hours (D) was done in independent triplicates. Blue bars indicate treated samples and red bars correspond to the untreated equivalent that was left at the biosafety cabinet under the same conditions while not subjecting to disinfecting irradiation. Data shown as PFUml^{-1} in triplicate assessed by plaque assay. **E.** Plaque phenotype comparison from one independent experiment at an irradiation dose of 0.6 mWcm^{-2}. Fixed and blocked plaques were immunostained using anti-SARS-CoV-2/NP antibody before developing using TrueBlue reagent. Data show in here are from three independent replicates (Mean+SEM).

Influenza A virus is susceptible to 405 nm inactivation in the absence of photosensitizers.

Given the observations derived from SARS-CoV-2, a separate inactivation study using a different lipid-enveloped RNA virus was conducted by using influenza A Puerto Rico (A/H1N1/PR8-Mount Sinai) virus strain. Irradiation with a high dose of 0.6 mWcm^{-2}

suggested a time dependent reduction of infectious titers as calculated by the 0.1619

$\log_{10}$, 0.5609 $\log_{10}$, and 1.6115 $\log_{10}$ (66 times) reductions at 1, 2, 4 and 8 hours

respectively (Figure 6A). And the reduction of plaques was apparent (figure 6B).

Figure 6 Inactivation of Influenza A virus in PBS by 405 nm irradiation. A. A dose of 0.6 mWcm^{-2} was applied to irradiate

samples at 405 nm over a course 8 hours while sampling at 1, 2, 4 and 8 hours (done in independent triplicates). Blue bars indicate

treated samples and red bars correspond to the untreated equivalent that was left at the biosafety cabinet under the same

conditions while not subjected to disinfecting irradiation. Data shown as PFUml^{-1} in triplicate assessed by plaque assay. **B.** Plaque

phenotype comparison from one independent experiment at an irradiation dose of 0.6 mWcm^{-2}. Fixed and blocked plaques were

stained using crystal violet. Data show in here are from three independent replicates (Mean+SEM).

The stability of IAV virus at room temperature for a period of 8 hours was found to be

the negligible in untreated IAV spiked PBS samples (Figure 6A).

Encephalomyocarditis virus (EMCV) as a model non-enveloped virus indicates

reduced susceptibility to 405 nm inactivation in the absence of photosensitizers.

In order to better understand the effect of the lipid-envelope in viral inactivation by 405

nm irradiation, we used a non-lipid enveloped RNA virus derived from the

Picornaviridae family. EMCV virus was irradiated at a high dose of 0.6 mWcm^{-2} similar

to SARS-CoV-2 and IAV.

Figure 7. Encephalomyocarditis virus (EMCV) in PBS shows reduced susceptibility to 405 nm irradiation. A. A dose of 0.6 mWcm^{-2} was applied to irradiate samples at 405 nm over a course 8 hours while sampling at 1, 2, 4 and 8 hours (done in independent triplicates). Blue bars indicate treated samples and red bars correspond to the untreated equivalent that was left at the biosafety cabinet under the same conditions while not subjected to disinfecting irradiation. Data shown as PFUml^{-1} in triplicate assessed by plaque assay. **B.** Plaque phenotype comparison from one independent experiment at an irradiation dose of 0.6 mWcm^{-2}. Fixed and blocked plaques were stained using crystal violet. Data show in here are from three independent replicates (Mean+SEM).

In this case however, a total reduction of 0.0969 (approximately 2 times) in comparison to the untreated after 8 hours of irradiation was observed (Fig 7A and 7B) indicating a lower rate of inactivation (despite identical dosing) in contrast to the lipid-enveloped RNA viruses tested in this study. The plaque reduction at 8 hours did not indicate the same dramatic reduction as observed with the latter studies.

As shown in Figure 4, standard LED lighting (without the specific addition of 405nm light) has a small but measurable amount of disinfecting light (0.006 mWcm-2) in the 400nm to 420nm range. To quantify the disinfecting effect of this light, irradiations were performed for 4h, 8h, 12h, and 24 h. A reduction was observed. {Raveen to add detail}. These results at 24h were largely consistent with other irradiation levels and reinforce the general observation of a 405nm dose-dependent effect. The intermediate irradiation levels (4h, 8h, and 12h) were generally larger than expected even when accounting for experimental uncertainty.

Discussion

The ongoing SARS-CoV-2 pandemic has affected the day-to-day functions in the entire world, raising concerns not only with regards to therapeutics but also in the context of virus survivorship and decontamination[36]. Taking into consideration the rapid spread of SARS-CoV-2 from person to person by droplets, aerosols, and fomites, whole-room disinfection systems can be viewed as a supplement to best practices for interrupting transmission of the virus.

Given the ongoing COVID-19 pandemic, we wanted to explore the impact of 405 nm enriched visible light technology on inactivation of respiratory pathogens such as SARS-CoV-2 and influenza A virus.

Without the use of exogenous photosensitizers, we were able to show that irradiation with low intensity (0.035 mWcm^{-2}) visible light yielded a reduction of $\log_{10}$ 0.3288 inactivation after four hours (0.5 Jcm^{-2}) and a total $\log_{10}$ 1.0325 inactivation of SARS-CoV-2 after 24 hours (3.02 Jcm^{-2}). A slightly higher dose (0.076 mWcm^{-2}) resulted in $\log_{10}$ 1.5393 reduction after 24 hours (6.56 Jcm^{-2}) while an irradiation dose of 0.150 mWcm^{-2} showed a reduction $\log_{10}$ 2.0056 after 24 hours (12.96 Jcm^{-2}) of irradiation. Finally, increasing the dose to 0.6 mWcm^{-2} yielded $\log_{10}$ 2.3010 reduction only after eight hours (12.96 Jcm^{-2}), indicating a both time and dose dependent inactivation of infectious viruses.

The irradiations using standard LED lighting raise some interesting questions for further discussion. With nearly 6x the amount of disinfecting light as compared to traditional fluorescent lighting but 1/6th the lowest amount of 405nm used in the study; it is conceivable that a reduction could be observed. The magnitude of this effect at less

than 24h was larger than expected based on other irradiations performed in this study. A similar effect was observed by Bache, et. al {REF} in whole-room bacterial reduction studies. They saw a disinfecting effect for irradiance values as low as 0.023 mWcm-2. This effect was shown to be uncorrelated to the level of irradiance used suggesting that the mere abundance of 405nm light can initiate the oxidizing chemical reaction. Nevertheless, our results clearly show a dose-dependent effect. One possible explanation for this observation is that the lower irradiance levels expose different responses by specific organisms within a population based on the individual biology of that organism. Clearly, these results suggest that additional factors may be at work and warrant further investigation.

We selected conventional plaque assays as the read out to specifically estimate infectious virus titers upon disinfection. Methods based in the quantification of viral RNA via PCR based techniques might be misleading as they detect viral RNA from both infectious and noninfectious virions.

SARS-CoV-2 is a lipid-enveloped virus composed of a ssRNA genome and our data indicates its susceptibility to visible light mediated inactivation. To further confirm these observations, we used influenza A virus. which is another human respiratory virus with a lipid envelop and an segmented-RNA genome. Upon irradiating for 1 hour at 0.6 mWcm^{-2} (2.16 Jcm^{-2}), we observed a total reduction of $\log_{10}$ 0.1619 for the influenza A virus compared to the reduction of $\log_{10}$ 0.4150 for SARS-CoV-2 under the same conditions. While both viruses have lipid envelopes, there is clearly a difference here that will require further study. One possible explanation is the difference in the virion size creating a physically smaller cross-section for absorption. (IAV ~120 nm and

SARS-CoV-2 ~200 nm)[37, 38]. Nevertheless, both viruses were largely inactivated after

eight hours achieving more 1.5 $\log_{10}$ reduction. Intriguingly, it was observed that both

RNA viruses were able to remain stable at room temperature for at least 24 hours,

indicating minimal decay which is consistent with previous studies[36, 39]. We next

irradiated a non-enveloped RNA virus, EMCV. Previous results for visible light against

non-enveloped viruses demonstrated the need for external photosensitizers such as

artificial saliva, blood, feces, etc[30, 36]. Without a porphyrin containing medium, we

expected little to no inactivation when this virus was irradiated with visible light. For

these measurements, we used the highest available irradiance of 0.6 $mWcm^{-2}$. As

anticipated, we observed only a $\log_{10}$ 0.0969 reduction after eight hours, however, this

appears to be with the statistical precision of the measurement based on the results

obtained from shorter irradiations (1, 2, and 4 hours). For comparison, a study involving

the M13-bacteriophage virus (a non-enveloped virus) showed a 3-Log reduction using

an irradiance of $50mWcm^{-2}$ (almost 100 times greater than the highest irradiance used

in this study) for 10 hours at 425 nm further supporting the idea that non-enveloped

viruses may require higher doses of visible light[40].

Our study was conducted using a neutral liquid media composed of PBS without any

photosensitizers and we were able to show that visible light can indeed inactivate lipid-

enveloped viruses, differing from the theory that states that photosensitizers are a

requirement for inactivation. While these results provide insight into the basic science

involved, they were performed within the context of the applied science needed to show

the potential impact of this technology upon the current COVID-19 pandemic. By using

safe, commercially practical irradiance levels, our results are more directly translatable

to occupied rooms in the clinical environment.

Other studies which used visible light-based irradiation have shown similar results in the

absence of photosensitizers, indicating the possibility of an alternative inactivation

mechanism[23, 25, 30]. Studies have proposed two theories for this observation primarily

due to non-405nm wavelengths emitted by the source: 1) some amount of 420-430 nm

emitted from the source is contributing to the viral inactivation [41], and 2) the presence of

UV-A (390 nm) within the source. This wavelength is known to create oxidative stress

upon viral capsids[42].

Longer wavelengths, such as 420-430nm, have shown inactivation of the murine

leukemia virus (MRV-A) [41]. While this is an intriguing study, it used a broad-spectrum

lamp with optical filters to selectively identify the spectrum primarily responsible with

their results. Unfortunately, they did not quantify the amount of light (using radiometric

units) within the spectrum of interest used to irradiate the virus. While transmission

profile of the filters used were provided, it does not consider the spectral composition of

the source itself making any direct quantitative comparison between our studies

impossible. It is interesting to note that they did observe viral inactivation in their

controls from wavelengths less than 420nm confirming the qualitative findings of our

study without confirming the specific use of 405nm. This suggests that the viral

inactivation is a likely a broad response (> 20nm) with relative contributions unique to

the chemistry of each organism. They also considered much longer exposures (~7

days) and much higher illuminance (> 200 lux) than that used in our study although this

is again difficult to compare given the lack of radiometric quantification of their light

source. It is important to note that the control samples used in our study were exposed

to the same overhead (non-405nm) lights as the irradiated samples and our results are

the observed difference between the two demonstrating the contribution from 405nm

over and above that potentially from 420-430nm. Future experiments can further

quantify the potential effect.

The other theory, potential UV-A irradiation, was historically applied to lamp-based

sources with broad spectral (> 100nm) outputs. Again, the use of LED technology

addresses this question as the peak irradiance at 390nm of the device used in this

study was < 1% of its peak irradiance at 405nm without the need for any additional

filtration. Future experiments can further quantify the potential effect.

Another consideration to be addressed is thermal heating of the virus by the LED

source. Tsen and Achilefu used a pulsed laser method at 425nm[43] with ~100 $mWcm^{-2}$

average power density for < 2h while simultaneously measuring the sample temperature

with a thermocouple. They detected less than a 2°C demonstrating minimal temperature

impact even under a power density nearly 9 orders of magnitude larger than that used

in this study. This was confirmed by our thermocouple measurements as stated earlier.

Nazari, *et al* used an 805nm source with an average power density of > 0.3 Wcm^{-2} for

10s, nearly 1000 times that used in this study[44]. While the total energy delivered was

more comparable to that used in our study, they did not make explicit temperature

measurements, their analysis ruled out any potential thermal effects.

One possible explanation for the observed differences between the enveloped and non-

enveloped organisms is absorption of the 405nm light by the lipid envelope itself. This

could, in turn, lead to the creation of reactive oxygen species (causing an oxidative

effect) or simply destruction of the envelope leading to a denaturing of the organism.

This question could serve as the basis for a range of future studies.

The results obtained suggest that the performance of visible light against SARS-CoV-2

is similar to other organisms commonly found in the environment such as *S. aureus.*

Previous studies have shown that the visible light irradiance levels used in this study

(0.035 mWcm^{-2} to 0.6 mWcm^{-2}) reduce bacteria levels in occupied rooms and improve

outcomes for surgical procedures. It is therefore reasonable to conclude that visible light

might be an effective disinfectant against SARS-CoV-2. More importantly, this

disinfection can operate continuously as it is safe for humans based upon the exposure

guidelines in IEC 62471[45]. This means that once it has been in use for a period of time,

the environment will be cleaner and safer the next time it is occupied by humans.

One limitation of this study is that the inactivation assays were performed in static liquid

media as opposed to aerosolized droplets. While the use of visible light in air

disinfection has been briefly studied where it was shown that its effectiveness increased

approximately 4-fold[46], further studies involving dynamic aerosolization are needed to

better understand the true potential of visible light mediated viral inactivation.

In any case, our study shows the increased susceptibility of enveloped respiratory viral

pathogens to 405 nm mediated inactivation in the absence of photosensitizers. The

irradiances used in this study are very low and might be easily applied to disinfect

occupied areas safely and continuously within hospitals, schools, restaurants, offices

and other locations. Of particular interest is the potential for standard LED lighting to

play a role in reducing the presence of SARS-CoV-2 in the environment.

Conclusions

We have demonstrated the basic science of inactivation of enveloped viruses such as

SARS-CoV-2 and Influenza-A using 405nm visible light within the context of the applied

science required for this technology to have an impact upon the current COVID-19

pandemic. Without the need for exogenous photosensitizers and by using safe,

commercially practical irradiance levels, our results can be easily translated to the

clinical environment.

Future Efforts

Future work should focus on explaining the difference between the enveloped and non-

enveloped results. This may include transmission electron microscopy (TEM),

hemagglutination assay (HA), or other methods focusing on the potential role that a

mediated reaction (due to the envelope itself) might play. The size of the virion particle

may play a role in photoelectric absorption and could be studied for different viral

species. We acknowledge that while unlikely, other wavelengths of visible light, beyond

400-420nm may play a role in the inactivation process and future studies should explore

this possibility as well. Finally, the inactivation kinetics of low irradiances could add

valuable insight into clinical applications of this technology.

Acknowledgments

We thank Kenall Manufacturing for supplying the M4DLIC6 fixture and test rig used in

this study. We thank Randy Albrecht for support with the BSL3 facility and procedures

at the ISMMS, and Richard Cadagan for technical assistance. This research was partly

funded by CRIP (Center for Research for Influenza Pathogenesis), a NIAID supported

Center of Excellence for Influenza Research and Surveillance (CEIRS, contract #

HHSN272201400008C); by the generous support of the JPB Foundation, the Open

Philanthropy Project (research grant 2020-215611 (5384)) and anonymous donors to

AG-S, and by a research contract from Kenall Manufacturing to the AG-S lab.

Conflicts of interest

511   References

1. Andersen, K. G., Rambaut, A., Lipkin, W. I., Holmes, E. C. & Garry, R. F. The
proximal origin of SARS-CoV-2. *Nat. Med.* **26**, 450-452 (2020).

2. Worldometer, D. COVID-19 coronavirus pandemic. *World Health Organization,*
www.worldometers.info (2020).

3. Buitrago-Garcia, D. *et al.* Occurrence and transmission potential of asymptomatic and
presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis.
PLoS medicine **17**, e1003346 (2020).

4. https://www.who.int/news-room/commentaries/detail/modes-of-transmission-of-virus-
causing-covid-19-implications-for-ipc-precaution-recommendations.

5. Dehbandi, R. & Zazouli, M. A. Stability of SARS-CoV-2 in different environmental
conditions. *The Lancet Microbe* **1**, e145 (2020).

6. Van Doremalen, N. *et al.* Aerosol and surface stability of SARS-CoV-2 as compared
with SARS-CoV-1. *N. Engl. J. Med.* **382**, 1564-1567 (2020).

7. Behzadinasab, S., Chin, A., Hosseini, M., Poon, L. & Ducker, W. A. A surface coating
that rapidly inactivates SARS-CoV-2. *ACS applied materials & interfaces* **12**, 34723-
34727 (2020).

8. Chan, K. *et al.* Factors affecting stability and infectivity of SARS-CoV-2. *J. Hosp.*
Infect. **106**, 226-231 (2020).

9. Biryukov, J. *et al.* Increasing Temperature and Relative Humidity Accelerates
Inactivation of SARS-CoV-2 on Surfaces. *mSphere* **5**, 10.1128/mSphere.00441-20
(2020).

10. Aboubakr, H. A., Sharafeldin, T. A. & Goyal, S. M. Stability of SARS-CoV-2 and
other coronaviruses in the environment and on common touch surfaces and the
influence of climatic conditions: A review. *Transboundary and emerging diseases*
(2020).

11. Smither, S. J., Eastaugh, L. S., Findlay, J. S. & Lever, M. S. Experimental aerosol
survival of SARS-CoV-2 in artificial saliva and tissue culture media at medium and high
humidity. *Emerging microbes & infections* **9**, 1415-1417 (2020).

12. Schuit, M. *et al.* Airborne SARS-CoV-2 is rapidly inactivated by simulated sunlight.
J. Infect. Dis. **222**, 564-571 (2020).

13. Kampf, G., Todt, D., Pfaender, S. & Steinmann, E. Persistence of coronaviruses on
inanimate surfaces and their inactivation with biocidal agents. *J. Hosp. Infect.* **104**, 246-
251 (2020).

14. Rutala, W. A. & Weber, D. J. Disinfection and sterilization in health care facilities:
what clinicians need to know. *Clinical infectious diseases* **39**, 702-709 (2004).

15. Rathnasinghe, R. *et al.* Scalable, effective, and rapid decontamination of SARS-
CoV-2 contaminated N95 respirators using germicidal ultra-violet C (UVC) irradiation
device. *medRxiv* (2020).

16. Escombe, A. R. *et al.* Upper-room ultraviolet light and negative air ionization to
prevent tuberculosis transmission. *PLoS Med* **6**, e1000043 (2009).

17. Nakpan, W., Yermakov, M., Indugula, R., Reponen, T. & Grinshpun, S. A.
Inactivation of bacterial and fungal spores by UV irradiation and gaseous iodine
treatment applied to air handling filters. *Sci. Total Environ.* **671**, 59-65 (2019).

18. Tseng, C. & Li, C. Inactivation of viruses on surfaces by ultraviolet germicidal
irradiation. *Journal of occupational and environmental hygiene* **4**, 400-405 (2007).

19. Kowalski, W. in *Ultraviolet germicidal irradiation handbook: UVGI for air and surface*
disinfection (Springer science & business media, 2010).

20. Zaffina, S. *et al.* Accidental exposure to UV radiation produced by germicidal lamp:
case report and risk assessment. *Photochem. Photobiol.* **88**, 1001-1004 (2012).

21. Leung, K. C. P. & Ko, T. C. S. Improper Use of the Germicidal Range Ultraviolet
Lamp for Household Disinfection Leading to Phototoxicity in COVID-19 Suspects.
Cornea **40**, 121-122 (2021).

22. Maclean, M. *et al*. Environmental decontamination of a hospital isolation room using
high-intensity narrow-spectrum light. *J. Hosp. Infect.* **76**, 247-251 (2010).

23. Maclean, M., McKenzie, K., Anderson, J. G., Gettinby, G. & MacGregor, S. J. 405
nm light technology for the inactivation of pathogens and its potential role for
environmental disinfection and infection control. *J. Hosp. Infect.* **88**, 1-11 (2014).

24. Murrell, L. J., Hamilton, E. K., Johnson, H. B. & Spencer, M. Influence of a visible-
light continuous environmental disinfection system on microbial contamination and
surgical site infections in an orthopedic operating room. *Am. J. Infect. Control* **47**, 804-
810 (2019).

25. Maclean, M., Murdoch, L. E., MacGregor, S. J. & Anderson, J. G. Sporicidal effects
of high-intensity 405 nm visible light on endospore-forming bacteria. *Photochem.*
Photobiol. **89**, 120-126 (2013).

26. Murdoch, L., McKenzie, K., Maclean, M., Macgregor, S. & Anderson, J. Lethal
effects of high-intensity violet 405-nm light on Saccharomyces cerevisiae, Candida
albicans, and on dormant and germinating spores of Aspergillus niger. *Fungal Biology*
117, 519-527 (2013).

27. Dai, T. *et al*. Blue light for infectious diseases: Propionibacterium acnes,
Helicobacter pylori, and beyond? *Drug Resistance Updates* **15**, 223-236 (2012).

28. Bumah, V. V. *et al*. Spectrally resolved infrared microscopy and chemometric tools
to reveal the interaction between blue light (470 nm) and methicillin-resistant
Staphylococcus aureus. *Journal of Photochemistry and Photobiology B: Biology* **167**,
150-157 (2017).

29. Hadi, J., Dunowska, M., Wu, S. & Brightwell, G. Control Measures for SARS-CoV-2:
A Review on Light-Based Inactivation of Single-Stranded RNA Viruses. *Pathogens* **9**,
737 (2020).

30. Tomb, R. M. *et al*. New proof-of-concept in viral inactivation: virucidal efficacy of 405
nm light against feline calicivirus as a model for norovirus decontamination. *Food and*
environmental virology **9**, 159-167 (2017).

31. Ho, D. T. *et al*. Effect of blue light emitting diode on viral hemorrhagic septicemia in
olive flounder (Paralichthys olivaceus). *Aquaculture* **521**, 735019 (2020).

32. Wu, J. *et al.* Virucidal efficacy of treatment with photodynamically activated
curcumin on murine norovirus bio-accumulated in oysters. *Photodiagnosis and*
photodynamic therapy **12**, 385-392 (2015).

33. Maclean, M., MacGregor, S. J., Anderson, J. G. & Woolsey, G. High-intensity
narrow-spectrum light inactivation and wavelength sensitivity of Staphylococcus aureus.
FEMS Microbiol. Lett. **285**, 227-232 (2008).

34. Martinez-Sobrido, L. & Garcia-Sastre, A. Generation of recombinant influenza virus
from plasmid DNA. *J. Vis. Exp.* **(42). pii: 2057. doi**, 10.3791/2057 (2010).

35. Carocci, M. & Bakkali-Kassimi, L. The encephalomyocarditis virus. *Virulence* **3**, 351-
367 (2012).

36. Derraik, J. G., Anderson, W. A., Connelly, E. A. & Anderson, Y. C. Rapid evidence
summary on SARS-CoV-2 survivorship and disinfection, and a reusable PPE protocol
using a double-hit process. *MedRxiv* (2020).

37. Bouvier, N. M. & Palese, P. The biology of influenza viruses. *Vaccine* **26**, D49-D53
(2008).

38. Bar-On, Y. M., Flamholz, A., Phillips, R. & Milo, R. Science Forum: SARS-CoV-2
(COVID-19) by the numbers. *Elife* **9**, e57309 (2020).

39. Wang, X., Zoueva, O., Zhao, J., Ye, Z. & Hewlett, I. Stability and infectivity of novel
pandemic influenza A (H1N1) virus in blood-derived matrices under different storage
conditions. *BMC infectious diseases* **11**, 1-6 (2011).

40. Tomb, R. M. *et al.* Inactivation of Streptomyces phage φC31 by 405 nm light:
Requirement for exogenous photosensitizers? *Bacteriophage* **4**, e32129 (2014).

41. Richardson, T. B. & Porter, C. D. Inactivation of murine leukaemia virus by exposure
to visible light. *Virology* **341**, 321-329 (2005).

42. Girard, P. et al. *UVA-induced damage to DNA and proteins: direct versus indirect*
photochemical processes (Journal of Physics: Conference Series Ser. 261, IOP
Publishing, 2011).

43. Tsen, S. D. *et al.* Studies of inactivation mechanism of non-enveloped icosahedral
virus by a visible ultrashort pulsed laser. *Virology journal* **11**, 1-9 (2014).

44. Nazari, M. *et al.* Plasmonic enhancement of selective photonic virus inactivation.
Scientific reports **7**, 1-10 (2017).

45. IEC 62471:. Photobiological safety of lamps and lamp systems. (2006).

46. Dougall, L. R., Anderson, J. G., Timoshkin, I. V., MacGregor, S. J. & Maclean, M.
Efficacy of antimicrobial 405 nm blue-light for inactivation of airborne bacteria (Light-
Based Diagnosis and Treatment of Infectious Diseases Ser. 10479, International
Society for Optics and Photonics, 2018).

630
631
632   Ref3 = Nazari M, Xi M, Lerch S, Alizadeh MH, Ettinger C, Akiyama H, Gillespie C,
633   Gummuluru S, Erramilli S, Reinhard BM, *Plasmonic Enhancement of Selective Photonic*
634   *Virus Inactivation.* Scientific Reports, **7**, 11951 (2017).
635
636

637