Chromosome counting in the mouse zygote using low-invasive super-resolution live-cell imaging

Yu Hatano[1], Daisuke Mashiko[1], Mikiko Tokoro[1,2], Tatsuma Yao[1,3], Kazuo Yamagata[1]

[1]Faculty of Biology-Oriented Science and Technology, KINDAI University, Kinokawa, Wakayama, Japan;

[2]Asada Institute for Reproductive Medicine, Asada Ladies Clinic, Aichi, Japan;

[3]Research and Development Center, Fuso Pharmaceutical Industries, Ltd., Osaka, Japan

Correspondence should be addressed to: yamagata@waka.kindai.ac.jp (KY)

Impact Statement

Low-toxic super-resolution observation enables chromosome counting in preimplantation

embryos without cell collection.

Abstract

In preimplantation embryos, an abnormal chromosome number causes developmental failure

and a reduction in the pregnancy rate. Conventional chromosome testing methods requiring

biopsy reduce the risk of associated genetic diseases; nevertheless, the reduction in cell number

also reduces the pregnancy rate. Therefore, we attempted to count the chromosomes in mouse

(Slc:ICR) embryos using super-resolution live-cell imaging as a new method of chromosome

counting that does not reduce the cell number or viability. We counted the forty chromosomes

at the first mitosis by injecting embryos with histone H2B-mCherry mRNA under conditions

by which pups could be obtained; however, the results were often an underestimation of

chromosome number and varied by embryo and time point. Therefore, we developed a method

to count the chromosomes via CRISPR/dCas-mediated live-cell fluorescence *in situ*

hybridization targeting the sequence of the centromere region, enabling us to count the

chromosomes more accurately in mouse embryos. The methodology presented here may

provide useful information for assisted reproductive technologies, such as those used in

livestock animals/humans, as a technique for assessing the chromosomal integrity of embryos

prior to transfer.

Introduction

In preimplantation embryos, the number of chromosomes is correlated with the pregnancy rate in assisted reproductive technology (ART) and animal breeding (*Magli et al., 2000, Sandalinas et al., 2001, Rubio et al., 2007, Mantikou., 2012, Yao et al., 2018*). In addition, even if embryos with abnormal chromosome numbers achieve full-term development, they may exhibit severe genetic disease (e.g., trisomy 21) (*Lejeune et al., 1959*). Therefore, evaluating chromosome numbers prior to embryo transplantation is important to mitigate these risks. Conventional chromosomal testing methods such as collecting and disrupting cells, subsequent spreading of the chromosomes, or scanning the genome using microarrays or next-generation sequencing can reduce the risk of genetic disease (Hens et al., 2013). Meanwhile, an associated risk of decreased pregnancy rate consequent to cell harvesting from chromosomally normal embryos has been reported (*Mastenbroek et al., 2007*). As approximately 90% of the causes of trisomy 21 are derived from meiosis (*Antonarakis et al., 1991, Freeman et al., 2007*), determining the number of chromosomes immediately following fertilization may facilitate the application of strategies to reduce the risk of genetic disease by excluding embryos with abnormal chromosome numbers from transplantation. Nevertheless, the number of chromosomes in early-stage embryos cannot be evaluated using these techniques since they are conventionally performed using blastocysts, which contain a sufficient number of cells to support cell collection. Therefore, the development of technology that enables real-time testing of chromosomes immediately after fertilization without reducing the number of pre-implantation germ cells will represent a breakthrough that will bring new insights into ART.

Live-cell imaging constitutes a technology for observing the interior of cells in their native environment. Through low-toxic, long-term live-cell imaging of mouse early embryos using fluorescence microscopy (*Yamagata 2009a*), we previously demonstrated that the early division of preimplantation embryos affects subsequent ontogeny. For example, abnormal chromosome segregation during early division affects development up to the blastocyst stage (*Mashiko et al., 2020*), resulting in extremely low birth rates following transplantation of 2-cell stage embryos (*Yamagata et al., 2009b*). However, although obvious abnormalities leading to the formation of micronuclei could be detected, it was not possible to count the number of chromosomes owing to limited resolution (*Mashiko et al., 2020*).

In recent years, super-resolution microscopes based on various principles have been proposed and used for imaging cells (i.e., stimulated emission depletion microscopy (STED): *Hell et al., 1994; Hell et al., 1995*; photoactivated localization microscopy (PALM): *Betzig et al., 2006*; fluorescence photoactivation localization microscopy (FPALM): *Hess et al., 2006*;

stochastic optical reconstruction microscopy (STORM): *Rust et al., 2006*; structured illumination microscopy (SIM): *Gustafsson et al., 2000*; and Airyscan: *Huff, 2015*). Nevertheless, long-term observation using a super-resolution microscope may be considered as highly invasive to observed cells. In particular, during the observation of preimplantation embryos, acquisition of three-dimensional images and repeated irradiation by the laser for time-lapse imaging causes embryo damage. Thus, fluorescence observation under conditions that allow the development of embryos to term without prior arrest is important to ensure that the observed phenomenon is not an artifact caused by dying cells. To this end, we attempted to identify appropriate long-term live-cell imaging conditions using a disk confocal type fluorescence microscope to observe early embryogenesis and predict prognosis (*Yamagata et al., 2009a*). In general, confocal microscopy scans a sample at a single plane, thus requiring substantial time to obtain an image. In addition, application of high-intensity laser energy may readily discolor the sample or have cytotoxic effects. In comparison, spinning disk confocal microscopy can overcome these limitations by scanning multiple points through rotation of the disk and is therefore suitable for live-cell imaging of early embryos. Moreover, super-resolution has recently been realized using a disk confocal, optical photon reassignment microscopy (OPRA)-type microscope by optically reducing individual focal points projected on pinholes using microlenses (super-resolution via optical re-assignment (SoRa): *Azuma and Kei, 2015*).

The objective of this study is therefore to develop a technology capable of performing chromosome counts in preimplantation embryos without requiring cell collection to identify embryos with aberrant chromosome numbers. Using mice, which have successfully been used as model animals for the study of mammalian embryos and the establishment of associated techniques, we applied the SoRa system to count chromosomes of early-stage embryos under conditions that allowed the embryo to maintain full-term developmental potential following the super-resolution observation.

Results

Construction of the SoRa live-cell imaging system and optimization of imaging conditions with minimal phototoxicity

We constructed a disk confocal type super-resolution live-cell imaging system to observe mouse fertilized eggs (*Fig. 1*). As the system has a switching mechanism between the spinning disk with (SoRa disk) and without (Normal disk) microlenses, we can observe the same specimen under these two different modes. To cover the broad viewing field of the system, a scientific complementary metal–oxide–semiconductor (sCMOS) image sensor was used and denoise algorithms (*Boulanger et al., 2009*) were activated at every acquisition to compensate for the lower sensitivity compared to electron multiplying charge-coupled devices (EMCCD) sensor. For observation of the first mitosis (approximately 17 h from the pronuclear stage to the end of the first mitosis), an incubator and gas chamber were set up on the stage for culturing embryos. A constant room temperature was maintained at 30 °C so that the embryos would not be affected by outside air.

To evaluate the phototoxicity resulting from observation using the constructed SoRa live-cell imaging system, we focused on the decay of the brightness (fading) of H2B-mCherry during the observation of early embryos. We injected mRNA encoding histone H2B-mCherry into fertilized eggs. One plane of the z-axis of the nucleus of the mouse 2-cell stage embryo was observed for 100 s continuously in streaming mode. Despite observation under the same conditions, such as excitation laser power, exposure time, and camera sensitivity, a decrease in signal was observed with SoRa compared to that observed using the conventional disk confocal microscopy system W1 (time constant 82.4 vs 137 ms, respectively) (*Fig. 2A, B*). This result suggested that the SoRa system requires more stringent conditions to suppress the phototoxicity than the conventional W1 system, which likely arises because the SoRa system has a higher light density (power of laser per unit area) than the conventional W1 system.

We next searched for conditions that did not affect embryogenesis following observation with a super-resolution microscope (*Table. 1*). We examined the wavelength, laser power, and time intervals. At an excitation wavelength of 561 nm, laser power of 0.1 mW (emitted from tip of objective), and 5 min intervals, the number of embryos that reached the 2-cell stage was 19/20 (95%) and 14/20 (70%) reached the blastocyst stage. At an excitation wavelength of 561 nm, laser power of 0.1 mW, and 10 min intervals, 18/20 (90%), embryos reached 2 cells and 16/20 (80%) reached the blastocyst stage. The blastocyst arrival rate under these conditions did not differ from that in the non-observed group (52/65 (80%), $P = 0.89$, $P = 1$, prop-test). Laser

power of 0.2 mW affected the growth up to the blastocyst stage more than 0.1 mW (0.2 mW,

5 min: 0/10 (0%); 0.2 mW, 10 min: 3/19 (15.8%), prop-test P = 0.016, 0.013, respectively).

These results suggested that a laser power of 0.1 mW is suitable for an excitation wavelength

of 561 nm. However, at an excitation wavelength of 488 nm and laser power of 0.1 mW, no

embryos reached the blastocyst stage regardless of the time interval, indicating that an

excitation wavelength of 488 nm is more toxic. We therefore utilized the conditions supportive

of normal embryo development following super-resolution observation using a wavelength of

561 nm in subsequent experiments.

To examine whether super-resolution live-cell imaging allowed subsequent full-term

embryo development, 2-cell embryos observed at an excitation wavelength of 561 nm, laser

power of 0.1 mW, and interval of 5 or 10 min were transplanted into the oviduct of

pseudopregnant mice. Notably, pups were obtained under both intervals (5 min (1/10): P =

0.57, 10 min (7/15): P = 0.66, respectively vs. imaging (−) prop-test) (***Table. 2; Fig S1***). We

therefore adopted a 10 min interval as the preferred observation condition.

Under the condition that the pups could be obtained (i.e., excitation wavelength 561 nm,

laser power 0.1 mW, 10 min interval), time-lapse observation was then performed for 17 h until

the completion of the first mitosis (***Fig. 3A, B; Movie. 1***).

SoRa system allows observation of the grooves between the sister chromatids

Subsequently, the resolution of the SoRa system was compared with that of the W1 system

using movies captured by each system. Chromosomes could be seen as two separating sister

chromatids by observation using SoRa but not the conventional W1 system. Furthermore, by

performing deconvolution processing on the images acquired with SoRa, the grooves of sister

chromosomes could be separated more clearly (peak-to-peak length: 618.7 $\pm$ 137.5 nm) (***Fig.***

4). When interphase nuclei were observed, the contrast of the chromatin in the nuclei could be

observed in more detail by imaging using the SoRa system than when using the conventional

W1 system (***Fig. 4***).

Chromosome segmentation and auto-counting

For counting the number of chromosomes, we attempted the segmentation of M-phase

chromosomes in living cells. Deconvoluted images were acquired using the Tikhonov

regularization algorithm (***Sage et al., 2017***). Images were noise-processed using the Top Hat

filter (***Legland et al., 2016***) and binarized with Otsu's method (***Otsu, 1979***). As a result of

automatically counting binarized objects (*Fig. 5A; Movie. 2*), we were able to count 40 objects

(*Fig. 5B*). However, a risk existed of recognizing two objects as one when the chromosomes

were close; consequently, the number of chromosomes detected were varied among the

embryos and time points (*Fig. 5C*). Although the chromosomes were clearly separated

compared to the results from conventional W1 system observation (*Fig. 5; Fig. S2*), the

inconsistent counting of individual chromosomes represented a limitation of this technique.

Chromosome counting by centromere labeling with live-fluorescence in situ

hybridization (live-FISH) technology

To overcome the issue of miscounting at sites of overlapping chromosome signals, we sought

to visualize the centromere, a more restricted DNA region in which the distal ends of all

chromosomes are located (*Fig. 6A*). Toward this end, CRISPR-mediated live imaging

technology (*Wang et al., 2019*) was performed by targeting the minor satellite sequence, a

repeat sequence constituting the centromere (*Fig. 6B*). First, we designed the crRNA based on

the target sequence used in a previous report of ES cells (*Anton et al., 2014*). As counter

staining, major satellite sequence locating pericentromere, a region on a chromosome that is

larger and existing inward than the centromere (*Fig. 6A*), was also labeled using TALE

technology (*Miyanari et al, 2013*). As a result, the signals of both minor and major satellites

were observed at the distal ends of metaphase and anaphase chromosomes (*Fig. 6C*).

Importantly, when compared to major satellites, the dot-like signals of minor satellites were

more restricted, and overlapping satellites were not detected (*Fig. 6C*).

Finally, the number of chromosomes at metaphase were counted by combining centromeres

labeled by live-FISH and the SoRa system. Consequently, it was possible to count 40 pairs of

dot-shaped signals (*Fig. 7A, B; Movies. 3, 4*). Notably, the number of counted dot-like signals

were 40 in all three embryos analyzed, which did not change with time (*Fig. 7C; Fig. S3*). In

contrast, conventional W1 system did not consistently detect the dot-like signal (*Fig. 7; Fig.*

S4). Moreover, through multicolor live-cell imaging using histone H2B-mCherry and

ATTO647N‐labeled gRNA for the minor satellite, we found that embryos would reach the

blastocyst stage (3/6, *Fig. 7; Fig. S5, Movie. 5*).

Discussion

The objective of this study was to establish a method to count the number of chromosomes in the one-cell zygote without the need for cell collection and fixation. We continuously observed the first cleavage of fertilized mouse eggs using an OPRA-type disk confocal super-resolution microscope and identified the observation conditions under which full-term embryos could be obtained. Depending on the image quality and timing, we succeeded in counting M-phase chromosomes using these conditions. Moreover, the CRISPR-mediated live-cell imaging of the minor-satellite region dramatically improved the accuracy of chromosome counting.

Implications of phototoxicity assessment and full-term development

When observing a thick sample such as an oocyte, the image becomes darker and the resolution in the Z-direction becomes lower than that of a thinner sample, such as most somatic cells. Although these challenges can be overcome by increasing the laser intensity or the number of images obtained in the Z-axis direction, such measures tend to decrease the embryo viability. Moreover, the effect of super-resolution imaging on the viability of preimplantation embryos has not been reported in the few prior studies that utilized such systems (e.g., Airyscan, SIM, and STED) to observe mouse embryos (*Zielinska et al., 2019*). In the present study, because observations using the SoRa system evinced faster fading of H2B-mCherry, which indicated higher phototoxicity than that in conventional systems (*Fig. 2A, B*), we searched for observation conditions that did not affect the prognosis of the embryo. *Squirrell et al. 1999* reported that embryo phototoxicity could be evaluated by determining the rate of full-term development following in-utero transplantation of hamster embryos observed using two-photon excitation microscopy. This method is convincing since it can confirm that the observed phenomenon is not caused by dying cells. In the same manner, phototoxicity consequent to observation using a wide-field microscope (*Yamagata et al., 2005*) and a disk confocal system (*Yamagata et al., 2009a*) has been evaluated. In these studies, EGFP-α-tubulin and histone H2B-mRFP1 were injected into fertilized eggs and observed in two colors (excitation wavelengths: 488 nm/561 nm). In comparison, in the present study, using super-resolution microscopy, we found that 2-cell embryos could not grow into blastocysts when observed with an excitation wavelength of 488 nm (*Table. 1*). Conversely, following observation at an excitation wavelength of 561 nm, the cells successfully developed into blastocysts and full-term embryos (*Table. 2*). These findings suggest that the phenomenon observed at the excitation wavelength of 561 nm did not reflect the abnormal behavior of cells dying from

226 phototoxicity. Furthermore, as we identified conditions under which embryo development
227 proceeded to full-term following super-resolution observation, it should be possible in future
228 experiments to associate phenomena observed using the SoRa system with the prognosis of
229 embryo development and the acquisition of offspring.

230
231

Possibilities afforded by super-resolution live-cell imaging of the first mitosis or chromatin

In the present study, we observed the first division of the mouse embryo (*Fig. 3A, B*). Failure
of chromosomal segregation during the first mitosis constitutes a major factor in the
termination of embryonic development (*Yamagata et al., 2009b*, *Mashiko et al., 2020*); thus,
detailed observation of the first mitosis using super-resolution live-cell imaging will allow for
the prediction of developmental failure. We have previously identified a mitotic pattern that
causes aneuploidy using a combination of next generation sequencing and low-magnification
live-cell imaging (*Mashiko et al., 2020*). By repeatedly associating super-resolution
observation of the first mitosis with genomic testing by next generation sequencing methods,
it may be possible to clarify the behavior of chromosome segregation, which is prone to
producing chromosome abnormalities.

Notably, analysis of the image data obtained in this study revealed that the SoRa system
provided sufficiently high resolution to observe chromatin density at the 2-cell stage (*Fig. 4*),
which suggests that this system offers considerable promise as a research tool. Since we have
established a low-invasive super-resolution observation method, it has become possible to link
the images with chromatin information in combination with conventional methods for
examining chromatin. For example, the combination of a method for detailed observation of
chromatin dynamics that does not affect embryogenesis (*Fig. 4*) with the profiles of chromatin
fold structure as observed by chromosome conformation capture (3C)/Hi-C (*Ke et al., 2017*)
and transcription as determined by live-imaging (*Bertrand et al., 1998; Park et al., 2014*) will
be expected to accelerate chromatin research.

Application of chromosome counting to zootechnical science and ART

Counting the chromosomes of mouse zygotes from the obtained super-resolution imaging data
(*Fig. 5A, B*) successfully allowed the detection of 40 objects (i.e., 40 pairs of sister chromatids)
with a risk of miscounting (*Fig. 5C*). The concept that chromosome counts could be obtained
using super-resolution microscopy had previously been tested on U2OS cells (*Gao et al.,*

2012); however, the effects on cell viability or risk of miscounting were not reported. In the
present study, by further applying the CRISPR-mediated live-FISH technique to our super-
resolution live-cell imaging system (*Fig. 6*), we obtained 40-pairs (mice: *Fig. 7*) of dot signals
without measurement error over time.

This suggests that if the chromosomes of livestock/human fertilized eggs could be
segmented in the same manner as mouse embryos and the number of chromosomes can be
measured, the risk of transplanting abnormal ploidy embryos may be reduced. Since the nuclear
envelope breakdown (NEBD) timing is different for each embryo, live-cell imaging is useful
to find the different timings more easily. Moreover, counting at multiple time points prevents
overlooking or overcounting the number of chromosomes.

Although chromosome counting has been attempted by observing the centromere and
kinetochore (*Chiang and Lampson, 2013*), to our knowledge the number of chromosomes has
not yet been counted in living cells. Notably, the human alpha-satellite sequence (*Willard,*
1985) is homologous to the mouse minor satellite sequence used in the present study; cattle are
also recognized to carry centromere satDNA (*Macaya et al., 1978; Modi et al., 1993; Modi et*
al., 1996). Thus, human and bovine embryo chromosomes could also be counted using a gRNA
that targets the alpha-satellite. Taken together, our methodology presented in this report, the
combination of SoRa system and live-FISH technology, has the potential for use in many
animal species and shows considerable promise for widespread application in ART and
breeding livestock animals.

Material and Methods

Animals

This study conformed to the requirements of the Guide for the Care and Use of Laboratory Animals. All animal experiments were approved by the Animal Care and Use Committee at the Research Institute for Kindai University (permit number: KABT-28-001). ICR strain mice (15-week-old) were obtained from Japan SLC, Inc. (Shizuoka, Japan), and were bred under a specific pathogen-free environment. Room conditions were standardized with temperature maintained at 23 °C, relative humidity at 50%, and 12 h/dark: 12 h light cycle. Animals had free access to water and commercial food pellets. Mice used for the experiments were euthanized by cervical dislocation.

In vitro **fertilization**

To obtain the unfertilized egg, superovulation was induced in female ICR mice (15 weeks old) by intraperitoneal injections of 10 IU pregnant mare serum gonadotropin (ASKA Animal Health Co., Ltd., Tokyo, Japan) and 10 IU human chorionic gonadotropin (hCG) (ASKA Animal Health) at 48 h intervals. Cumulus-intact oocytes were recovered at euthanasia 15–17 h following hCG injection; these oocytes were arrested at metaphase II of the meiotic cell cycle. Spermatozoa were collected from the cauda epididymis of male ICR mice (11–16 weeks old) in 0.2 mL droplets of TYH medium (*Toyoda et al., 1971*) and capacitated by incubation for 1.5 h at 37 °C under 6% CO_2 in humidified air. Cumulus-intact oocytes were collected in 0.2 mL of TYH medium and inseminated with spermatozoa (final concentration 75 sperm/μL). Upon insemination, oocytes arrested at metaphase II restarted the meiotic cell cycle and proceeded to interphase in approximately 3 h. After 1.5 h incubation at 37 °C under 6% CO_2 in humidified air, the cumulus cells were dispersed by brief treatment with hyaluronidase (Type I-S, 120–300 units/mL; Sigma-Aldrich, St Louis, MO, USA). A total of 30 female and 6 male mice were used, with three experimental replications (female 10). The obtained fertilized eggs were frozen after mRNA injection. In this process, the eggs of each group used in the experiment were randomized, which was expected to mask individual differences in mice. Frozen eggs were used as follows: 20 for the examination of fading, 57 for the examination of developmental capacity, 45 for the examination of birth rate, 10 for observation of 1st mitosis, 10 for W1 and SoRa comparison, 10 for chromosome counting using H2B-mCherry, 10 for chromosome counting using CRISPR-mediated live-FISH, and 20 for measuring the distance between sister

316   chromosomes.

Preparation of mRNAs to express proteins of interest

After linearization of the template plasmids (pcDNA3.1-polyA83: *Yamagata et al., 2005*,

pTALYM3 (Addgene #47874): *Miyanari et al., 2013*) at the XhoI (histone H2B-mCherry,

histone H2B-EGFP) and ApaI (pTALYM3) site, mRNA was synthesized using RiboMAX^{TM}

Large Scale RNA Production Systems-T7 (Promega, Madison, WI, USA). For efficient

translation of the fusion proteins in embryos, the 5′-end of each mRNA was capped using the

Ribo m7G Cap Analog (Promega), according to the manufacturer's protocol. To circumvent

the integration of template DNA into the embryonic genome, the reaction mixtures for in vitro

transcription were treated with RQ-1 RNase-free DNase I (Promega). Synthesized mRNAs

were treated with phenol-chloroform to remove protein components. The mRNAs were further

purified by filtration using MicroSpin^{TM} S-200 HR columns (GE Healthcare, Chicago, IL,

USA) to remove unreacted substrates (RNA reaction intermediates) and then stored at −80 °C

until use.

Preparation of the dCas/gRNA complex

The target sequence of minor satellites of mouse was 5′-ACACTGAAAAACACATTCGT-3′

(*Anton et al., 2014*) crRNA and tracrRNA-ATTO550/ATTO647N (Integrated DNA

Technologies, Redwood City, CA) hybridized using a T100 thermal cycler (Bio-Rad

Laboratories, Hercules, CA, USA) (94 °C: 5 min, 60 °C: 5 min) were mixed with dCas protein

(Integrated DNA Technologies, Redwood City, CA). The final concentrations of gRNA and

dCas for mice embryos were 20 and 200 ng/µL, respectively.

Microinjection of mRNA or dCas/gRNA complex

Probe injection into fertilized eggs was performed as described previously (*Yamagata et al.,*

2005). Briefly, mRNAs were diluted to 10 ng/µL using ultrapure water (Thermo Fisher

Scientific Barnstead Smart2Pure; Waltham, MA, USA) and an aliquot was placed in a

micromanipulation chamber. Fertilized eggs (approximately 4–6 h following insemination)

were transferred to HEPES-buffered Chatot-Ziomek-Bavister (CZB) (*Chatot et al., 1989*)

medium in the chamber and injected with mRNA using a piezo-driven manipulator with a

narrow glass pipette (1 µm diameter). Once the mRNA solution had been aspirated into the

pipette, piezo pulses were applied to the fertilized eggs to break the zona pellucida and plasma

membrane. A few picoliters of the solution were introduced into the cytoplasm and the pipette

was removed gently. The mRNA-injected fertilized eggs were incubated at 37 °C under 6%
CO_2 in air for at least 2 h prior to injection to allow time for protein production. After these
procedures, the embryos were frozen.
353
Imaging
The motor-control (Mac5000; Ludl Electronic Products, Hawthorne, NY, USA) and piezo
control (P-725xDD, PIFOC high dynamics piezo scanner) methods were used to control the Z-
axis direction. However, as the images obtained using the piezo-control method were blurred,
we adopted the motor-control method for subsequent experiments. The fertilized eggs were
transferred to 5 µL droplets of potassium simplex optimization medium with amino acids
(KSOMaa) containing 0.00025% polyvinyl alcohol (P8136-250G; Sigma-Aldrich) and 100
µM ethylenediaminetetraacetic acid on a glass-bottomed dish and placed in an incubation
chamber (Tokai Hit, Shizuoka, Japan) set at 37°C on the microscope stage. A gas mixture of
5% O_2, 6% CO_2, and 89% N_2 was introduced into the chamber (138 mL/min). An inverted IX-
73 microscope (Olympus, Tokyo, Japan) was used with an attached Nipkow disk confocal
microscope (CSU-W1 SoRa; Yokogawa Electric, Tokyo, Japan), a Prime95B scientific
complementary metal oxide semiconductor (sCMOS) camera (Teledyne Photometrics, Tuscon,
AZ, USA), and z-motor (Mac5000; Ludl Electronic Products, Hawthorne, NY, USA). As our
imaging device contained an attached auto x-y stage (Sigma Koki, Tokyo, Japan), multiple
embryos could be monitored simultaneously. The SoRa system can be switched to the
traditional nipkow disk without micro lenses (CSU-W1 mode), allowing parallel observation
of the same sample. A set of imaging systems was placed in a dark room with the room
temperature maintained at 30 °C. Device control was performed using µ-Manager microscopy
software (**https://micro-manager.org**). When observing in the Z-direction, 101 images were
taken at 0.5 µm intervals, 25 µm above and below the equatorial plane of the fertilized egg
(total 50 µm). The observed embryos were subsequently moved to an incubator to examine
their developmental potential. The power of laser emitted from tip of the objective was
measured by optical power meter (TB200; Yokogawa Electric, Tokyo, Japan)
378
379 **Image analysis**
380 The 2D images were constructed using MetaMorph software ver. 7.7.10 (Molecular Devices,
381 San Jose, CA, USA), µ-Manager microscopy software, and the ImageJ/Fiji image analysis
382 platform (**https://imagej.net/Fiji**). The measurements of fluorescence intensity were
383 performed by ImageJ/Fiji. The time constant of the SoRa system was obtained by measuring

the time until H2B-mCherry decayed to 36.8% (derived from the definition of the time constant in exponential decay and from the reciprocal of the Napier number), and the time constant of the W1 system was derived from the time until H2B-mCherry decayed to 36.8% after fitting the exponential curve. Centromere images obtained using the W1 and SoRa systems were binarized using the yen-method, denoised using median-filter, and the region of interest (ROI) was detected using Icy (**http://icy.bioimageanalysis.org/**).

Chromosome counting/dot signal counting

The FIJI distribution of ImageJ (*Schindelin et al., 2012*) was used for image processing. Deconvolved images were acquired using the Tikhonov regularization algorithm in the DeconvolutionLab2 plugin (*Sage et al., 2017*) with a theoretical point spread function calculated using the Diffraction PSF 3D plugin. Following deconvolution, these images were applied to the Top Hat filter in the MorphoLibJ plugin (*Legland et al., 2016*) for noise removal and binarized using Otsu's algorithm. Proximate objects in the image were separated using the Transform Watershed 3D algorithm in the MorphoLibJ plugin. The dot signals of the minor satellite were counted using FIJI and FluoRender (*Wan et al., 2012*; **https://www.sci.utah.edu/software**).

Embryo transfer

Embryo transfer has been described previously (*Yamagata et al., 2009b*). Two-cell mice embryos were transferred into the oviducts of day 0.5 pseudopregnant mice. At 18 days following the transfer, cesarean section was performed.

Sample-size estimation

The blastocyst formation rate in mice is approximately 70–90% and is expected to decrease to 0% to 20% when problems such as environmental or embryo damage occur. We determined that a total of n = 5.08 embryos were required to test the hypothesis that blastocyst formation rate is reduced by super-resolution observation by assuming power = 0.8, sig.level = 0.05, blastocyst ratio of control group = 0.8, blastocyst ratio of observed group = 0.1, one-sided test and prop-test. In turn, the birth rate of mice is approximately 40–60% and is expected to be reduced to 0–10% owing to embryonic damage. We calculated that n = 11.08 embryos would be required to test the hypothesis that super-resolution observation reduced the birth rate by assuming the power was 0.8, sig.level = 0.05, term development ratio of the control group = 0.5, term ratio of the observed group = 0.05, one-sided test and prop-test. However, the lack of

significant difference under these conditions does not necessarily suggest the lack of effect. Thus, in the present study, we confirmed whether any abnormalities in blastocyst formation were present following observation of the embryos under these conditions and considered whether live offspring could be obtained. Power analysis was performed using R.

Experimental replication

With super-resolution observation, the number of eggs that could be observed in one experiment was approximately 10. Thus, the number of eggs used in one experiment was approximately 5 in each of the two groups (biological replication n = 5/once). Technical replication was N = 1 in the examination of fading, N = 7 for blastocyst formation rate, N = 5 for term ratio, N = 3 for first mitosis, N = 3 in the comparison between conventional W1 and SoRa systems, N = 3 each for chromosome counting by H2B-mCherry mRNA and by CRISPR-mediated live-FISH, and N = 2 for measuring the distance between sister chromosomes. The technical replication of experiments related to qualitative conclusions and calculation of the time constant was set to 1. By setting a control group for each experimental replication, we confirmed that inconsistent observation results were not obtained for each experimental replication control group.

Statistical analysis

One-sided proportion test was performed using the prop.test function in R. Statistics on independent repeats were not possible due to the small number of embryos that could be observed each time in the case of super-resolution live-cell imaging system, so we performed statistics on the total number of embryos.

Competing interests

The authors declare no conflict of interest.

Author contributions

Yu Hatano: Methodology, Validation, Formal analysis, Investigation, Resources, Visualization, manuscript review & editing; Daisuke Mashiko: Data curation, Formal analysis, original manuscript draft preparation; Mikiko Tokoro: Formal analysis, Investigation, Resources; Tatsuma Yao: Formal analysis, Software; Kazuo Yamagata: Conceptualization, Supervision, manuscript review & editing, Funding acquisition.

Acknowledgements

We would like to thank Editage for English language editing. We would like to thank Takuya Azuma for producing the image of the SoRa W1 switching system.

Funding

This work was supported by JSPS KAKENHI Grant Numbers JP25712035, JP25116005, JP18H05528, and JP18H02357 to KY. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

References

Anton T, Bultman S, Leonhardt H, Markaki Y. 2014. Visualization of specific DNA sequences in living mouse embryonic stem cells with a programmable fluorescent CRISPR/Cas system. *Nucleus* **5**:163–172. DOI: https://doi.org/10.4161/nucl.28488

Antonarakis SE. 1991. Parental origin of the extra chromosome in trisomy 21 as indicated by analysis of DNA polymorphisms. Down Syndrome Collaborative Group. *New England Journal of Medicine* **324**:872–876. DOI: https://doi.org/10.1056/NEJM199103283241302

Azuma T, Kei T. 2015. Super-resolution spinning-disk confocal microscopy using optical photon reassignment. *Optics Express* **23**:15003–15011. DOI: https://doi.org/10.1364/OE.23.015003

Bertrand E, Chartrand P, Schaefer M, Shenoy SM, Singer RH, Long RM. 1998. Localization of ASH1 mRNA particles in living yeast. *Molecular Cell* **2**:437–445. DOI: https://doi.org/10.1016/S1097-2765(00)80143-4

Betzig E, Patterson GH, Sougrat R, Lindwasser OW, Olenych S, Bonifacino JS, Davidson MW, Lippincott-Schwartz J, Hess HF. 2006. Imaging intracellular fluorescent proteins at nanometer resolution. *Science*, **313**:1642–1645. DOI: https://doi.org/10.1126/science.1127344

Boulanger J, Kervrann C, Bouthemy P, Elbau P, Sibarita J B, Salamero J. 2009. Patch-based nonlocal functional for denoising fluorescence microscopy image sequences. *IEEE transactions on medical imaging*, *29*(2), 442-454. DOI: https://doi.org/10.1109/TMI.2009.2033991

Chatot CL, Ziomek CA, Bavister BD, Lewis JL, Torres I. 1989. An improved culture medium supports development of random-bred 1-cell mouse embryos in vitro. *Journal of Reproduction and Fertility* **86**:679–688. DOI: https://doi.org/10.1530/jrf.0.0860679

Chiang T, Lampson MA. 2013. Counting chromosomes in intact eggs. Methods in Molecular Biology **957**:249–253. DOI: https://doi.org/10.1007/978-1-62703-191-2_17

Freeman SB, Allen EG, Oxford-Wright CL, Tinker SW, Druschel C, Hobbs CA, O'Leary LA, Romitti PA, Royle MJ, Torfs CP, Sherman SL. 2007. The National Down Syndrome Project: design and implementation. *Public Health Reports* 122:62–72. DOI: https://doi.org/10.1177/003335490712200109

Gao L, Shao L, Higgins CD, Poulton JS, Peifer M, Davidson MW, Wu X, Goldstein B, Betzig E. 2012. Noninvasive imaging beyond the diffraction limit of 3D dynamics in thickly fluorescent specimens. *Cell* **151**:1370–1385. DOI: https://doi.org/10.1016/j.cell.2012.10.008

Gustafsson MG. 2000. Surpassing the lateral resolution limit by a factor of two using structured illumination microscopy. *Journal of Microscopy* **198**:82–87. DOI: https://doi.org/10.1046/j.1365-2818.2000.00710.x

Hell SW, Wichmann J. 1994. Breaking the diffraction resolution limit by stimulated emission: stimulated-emission-depletion fluorescence microscopy. *Optics Letters* **19**:780–782. DOI: https://doi.org/10.1364/OL.19.000780

Hell SW, Kroug M. 1995. Ground-state-depletion fluorescence microscopy: A concept for breaking the diffraction resolution limit. *Applied Physics B* **60**:495–497. DOI: https://doi.org/10.1007/BF01081333

Hens K, Dondorp W, Handyside A H, Harper J, Newson A J, Pennings G, ... & de Wert G. 2013. Dynamics and ethics of comprehensive preimplantation genetic testing: a review of the challenges. *Human Reproduction Update*, *19*(4), 366-375. DOI: https://doi.org/10.1093/humupd/dmt009

Hess ST, Girirajan TP, Mason MD. 2006. Ultra-high resolution imaging by fluorescence photoactivation localization microscopy. *Biophysical Journal* **91**:4258–4272. DOI: https://doi.org/10.1529/biophysj.106.091116

Huff J. 2015. The Airyscan detector from ZEISS: confocal imaging with improved signal-to-noise ratio and super-resolution. *Nature Methods* **12**: i–ii. DOI: https://doi.org/10.1038/nmeth.f.388

Ke Y, Xu Y, Chen X, Feng S, Liu Z, Sun Y, Yao X, Li F, Zhu W, Gao L, Chen H, Du Z, Xie W, Xu X, Huang X, Jiang L.. 2017. 3D chromatin structures of mature gametes and structural reprogramming during mammalian embryogenesis. *Cell* **170**:367–381. DOI: https://doi.org/10.1016/j.cell.2017.06.029

Legland D, Arganda-Carreras I, Andrey P. 2016. MorphoLibJ: integrated library and plugins for mathematical morphology with ImageJ. *Bioinformatics* **32**:3532–3534. DOI: https://doi.org/10.1093/bioinformatics/btw413

Lejeune J, Gauthier M, Turpin R. 1959. [Human chromosomes in tissue culture]. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* **248**:602–603.

Macaya G, Cortadas J, Bernardi G. 1978. An analysis of the bovine genome by density-gradient centrifugation. Preparation of the dG+dC-rich DNA components. *European Journal of Biochemistry* **84**:179–188 DOI: https://doi.org/10.1111/j.1432-1033.1977.tb11565.x

Magli MC, Jones GM, Gras L, Gianaroli L, Dorman I, Trounson AO. 2000. Chromosome mosaicism in day 3 aneuploid embryos that develop to morphologically normal blastocysts

in vitro. *Human Reproduction* **15**:1781–1786. DOI:
https://doi.org/10.1093/humrep/15.8.1781

Mantikou E, Wong KM, Repping S, Mastenbroek S. 2012. Molecular origin of mitotic
aneuploidies in preimplantation embryos. *Biochimica Biophysica Acta* **1822**:1921–1930.
DOI: https://doi.org/10.1016/j.bbadis.2012.06.013

Mashiko D, Ikeda Z, Yao T, Tokoro M, Fukunaga N, Asada Y, Yamagata K. 2020.
Chromosome segregation error during early cleavage in mouse pre-implantation embryo
does not necessarily cause developmental failure after blastocyst stage. *Scientific Reports*
10:854. DOI: https://doi.org/10.1038/s41598-020-57817-x

Mastenbroek S, Twisk M, van Echten-Arends J, Sikkema-Raddatz B, Korevaar JC, Verhoeve
HR, Vogel NEA, Arts EGJM, de Vries JWA, Bossuyt PM, Buys CHCM, Henieman MJ,
Repping S, van der Veen F. 2007. In vitro fertilization with preimplantation genetic
screening. *New England Journal of Medicine* **357**:9–17. DOI:
https://doi.org/10.1056/NEJMoa067744

Miyanari Y, Ziegler-Birling C, Torres-Padilla ME. 2013. Live visualization of chromatin
dynamics with fluorescent TALEs. *Nature Structural & Molecular Biology* **20:**1321–1324.
DOI: https://doi.org/10.1038/nsmb.2680.

Modi WS, Gallagher DS, Womack JE. 1993. Molecular organization and chromosomal
localization of six highly repeated DNA families in the bovine genome. *Animal*
Biotechnology **4**:143–161. DOI: https://doi.org/10.1080/10495399309525793

Modi WS, Gallagher DS, Womack JE. 1996. Evolutionary histories of highly repeated DNA
families among the Artiodactyla (Mammalia). *Journal of Molecular Evolution* **42**:337–349.
DOI: https://doi.org/10.1007/BF02337544

Otsu N, 1979. A Threshold Selection Method from Gray-Level Histograms. in *IEEE*
Transactions on Systems, Man, and Cybernetics **9**:62–66. DOI:
https://doi.org/10.1109/TSMC.1979.4310076.

Park HY, Lim H, Yoon YJ, Follenzi A, Nwokafor C, Lopez-Jones M, Meng X, Singer RH.
2014. Visualization of dynamics of single endogenous mRNA labeled in live mouse.
Science **343**:422–424. DOI: https://doi.org/10.1126/science.1239200

Rubio C, Rodrigo L, Mercader A, Mateu E, Buendía P, Pehlivan T, viloria T, De los Santos,
MJ, Simón C, Remohí J, Pellicer A. 2007. Impact of chromosomal abnormalities on
preimplantation embryo development. *Prenatal Diagnosis* **27**:748–756. DOI:
https://doi.org/10.1002/pd.1773

Rust MJ, Bates M, Zhuang X. 2006. Sub-diffraction-limit imaging by stochastic optical

reconstruction microscopy (STORM). *Nature Methods* **3**:793–795. DOI:
https://doi.org/10.1038/nmeth929

Sage D, Donati L, Soulez F, Fortun D, Schmit G, Seitz A, Guiet R, Vonesch C, Unser M. 2017.
DeconvolutionLab2: An open-source software fordeconvolution microscopy. *Methods*
115:28–41. DOI: https://doi.org/10.1016/j.ymeth.2016.12.015

Sandalinas M, Sadowy S, Akikani M, Calderon G, Cohen , Munné S. 2001. Developmental
ability of chromosomally abnormal human embryos to develop to the blastocyst stage.
Human Reproduction **16**:1954–1958. DOI: https://doi.org/10.1093/humrep/16.9.1954

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietsch T Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis.
Nature Methods **9**:676–82. DOI: https://doi.org/10.1038/nmeth.2019

Squirrell JM, Wokosin DL, White JG, Bavister BD. 1999. Long-term two-photon
fluorescence imaging of mammalian embryos without compromising viability. *Nature*
Biotechnology **17**:763–767. DOI: https://doi.org/10.1038/11698

Toyoda Y, Yokoyama M, Hoshi T. 1971. Studies on the fertilization of mouse eggs *in*
vitro. *Japanese Journal of Animal Reproduction* **16**, 147–151. DOI:
https://doi.org/10.1262/jrd1955.16.147

Wan Y, Otsuna H, Chien CB, Hansen C. 2012. FluoRender: an application of 2D image space
methods for 3D and 4D confocal microscopy data visualization in neurobiology research.
In *2012 IEEE Pacific Visualization Symposium* (pp. 201–208). IEEE, New York. DOI:
https://doi.org/10.1109/PacificVis.2012.6183592

Wang H, Nakamura M, Abbott TR, Zhao D, Luo K, Yu C, Nguyen CM, Lo A, Daley TP, La
Russa M, Liu Y, Qi LS. 2019. CRISPR-mediated live imaging of genome editing and
transcription. *Science* **365**.6459:1301–1305. DOI: https://doi.org/10.1126/science.aax7852

Willard HF. 1985. Chromosome-specific organization of human alpha satellite DNA.
American Journal of Human Genetics **37**:524–532 DOI: https://doi.org/10.1016/0888-
7543(87)90103-0

Yamagata K, Suetsugu R, Wakayama T. 2009a. Long-term, six-dimensional live-cell imaging
for the mouse preimplantation embryo that does not affect full-term development. *Journal*
of Reproduction and Development **55**:343–350. DOI: https://doi.org/10.1262/jrd.20166

Yamagata K, Suetsugu R, & Wakayama T. 2009b. Assessment of chromosomal integrity using
a novel live-cell imaging technique in mouse embryos produced by intracytoplasmic sperm
injection. *Human reproduction*, *24*(10), 2490-2499. DOI:

https://doi.org/10.1093/humrep/dep236

Yamagata K, Yamazaki T, Yamashita M, Hara Y, Ogonuki N, Ogura A. 2005. Noninvasive

visualization of molecular events in the mammalian zygote. *Genesis* **43**:71–79. DOI:

https://doi.org/10.1002/gene.20158

Yao T, Suzuki R, Furuta N, Suzuki Y, Kabe K, Tokoro M, Sugawara A, Yajima A, Nagasawa

T, Matoba S, Yamagata K, Sugimura S. 2018. Live-cell imaging of nuclear–chromosomal

dynamics in bovine in vitro fertilised embryos. *Scientific Reports* **8**:7460. DOI:

https://doi.org/10.1038/s41598-018-25698-w

Zielinska AP, Bellou E, Sharma N, Frombach AS, Seres KB, Gruhn JR, Blayner M, Eckel H,

Moltrecht r, Elder K, Hoffman ER, Schuh M. 2019. Meiotic kinetochores fragment into

multiple lobes upon cohesin loss in aging eggs. *Current Biology* **29**:3749–3765.e7. DOI:

https://doi.org/10.1016/j.cub.2019.09.006

Figure legends

Figure 1. SoRa live-cell imaging system. Photograph (left) and list (right) of the equipments for imaging. A conventional inverted microscope was attached to a Nipkow disk confocal unit, sCMOS camera, Z-motor, and XY auto stage. All devices were controlled using a micro-manager. Embryos were cultured in a CO_2 incubator at this stage. The Nipkow disk confocal unit can switch between a conventional unit without microlenses and a unit with microlenses (lower left).

Figure 2. Assessment of the phototoxicity of super-resolution imaging. (**A**) Photographs of 2-cell nuclei obtained using the W1 (upper panels) and SoRa (below panels) systems. bar: 1 µm. (**B**) Graph of the brightness change.

Figure 3. Super-resolution live-cell imaging of the first mitosis using chromosome counting by histone H2B-mCherry mRNA was the SoRa system. (**A**) Schematic diagram of super-resolution imaging of the first mitosis. (**B**) Super-resolution imaging of the first mitosis in mouse embryos. The upper panels show the bright-field images. Middle panels show the histone H2B-mCherry images of the x–y plane, and the panels below show that of the x–z plane. See also Supplementary movie 1.

Figure 4. The SoRa system enables higher resolution imaging than the conventional W1 system. (**A**) Left six panels show images of the metaphase of an embryo. From left to right, the image was obtained using W1 (x–y and x–z), SoRa (x–y and x–z), and SoRa (x–y and x–z; deconvoluted). The right three panels show enlargements of the rectangular area (red). The right three graphs show the relative intensity on the yellow dashed line. (**B**) The left six panels show images of the interphase of a 2-cell embryo. The right three panels show enlargements of the rectangular area (red).

Figure 5. Chromosome counting using histone H2B-mCherry mRNA. (**A**) The left panel shows the histone H2B-mCherry signal. The right image shows the processed image. (**B**) Each panel shows the recognized object. The white arrowheads indicate a processed chromosome with a risk of miscount. (**C**) Relationship between the number of recognized objects and time from nuclear envelope breakdown (NEBD). See also Supplementary Movie 2 and Supplementary Figure 2.

Figure 6. Chromosome counting by CRISPR-mediated live-FISH. (**A**) The left panel shows

the illustrated chromosome; the grey and black areas show the mouse centromere and

pericentromere regions, respectively. The below shows the repeated sequence of minor

satellite; the underlined sequence shows the sequence targeted by gRNA. (**B**) Illustrated

scheme of CRISPR-mediated live-cell imaging. The ATTO550 or 647N probe was added to

tracrRNA. (**C**) Snapshot of Histone H2B-mCherry, TALMaj-mClover3, and minor satellite

targeted gRNA using the SoRa system.

Figure 7. CRISPR-mediated live-cell imaging of the centromere region in the mouse

preimplantation embryo. (**A**) Left panel shows the centromere region and right panel shows the

segmented image. (**B**) Segmented 40 centromere pairs. (**C**) The number of pairs of dot signals

was measured at three points prior to the start of anaphase (t = 0). See also Supplementary

Figures 3 and 4.

Table captions

Table 1. Developmental capacity of the preimplantation embryo following super-resolution

imaging.

From left to right column: injected mRNA, excitation wavelength of the laser (nm), optical

intensity (mW), time interval (min), imaging period (h), number of embryos examined, number

of 2-cell embryos, and number of blastocysts.

Table 2. Full-term development of transferred embryos following super-resolution imaging.

From left to right column: injected mRNA, excitation wavelength of the laser (nm), optical

intensity (mW), time interval (min), imaging period (h), number of embryos examined, number

of 2-cell stage embryos, number of recipients (transferred mice), number of transferred

embryos, and number of pups. See also Supplementary figure 1.

Additional files

Movie 1 (Related to Figure 3). Super-resolution observation of the first mitosis of a mouse

embryo. The left panel shows the movie from bright field (BF) imaging while the right shows

the movie of the histone H2B-mCherry signal.

Movie 2 (Related to Figure 5). Three-dimensional (3D) construction of segmented

chromosomes. Rotated 3D images of the histone H2B-mCherry signal (left) and processed

image (right) are shown.

Movie 3 (Related to Figure 7). Live-cell imaging using minor satellite targeted gRNA.

Imaging of the minor satellite region for approximately 17 h from the one-cell to two-cell stage.

Movie 4 (Related to Figure 7). Three-dimensional (3D) construction of the segmented

centromere region. Rotated 3D images of CRISPR-mediated minor satellite signal (left) and

the segmented centromere region (right).

Movie 5 (Related to Figure 7). Multi color live-cell imaging of a mouse embryo using histone

H2B-mCherry and minor satellite-targeted gRNA. The left panel shows the bright-field movie

while the right shows the merged movies of histone H2B-mCherry and CRISPR-mediated

minor satellite signals.

Table 1. Developmental capacity of the preimplantation embryo following super-resolution imaging

Injected mRNA	Optical wavelength (nm)	Optical intensity (mW)	Time intervals (min)	Imaging period (hour)	No. of embryos examined	No. (%) of 2-cell embryos	No. (%) of blastocyst
Histone H2B-mCherry	-	-	-	-	65	63 (96.9)	52 (80)
	561	0.1	5	17	20	19 (95)	14 (70)
	561	0.1	10	17	20	18 (90)	16 (80)
	561	0.2	5	17	10	4 (40)	0 (0)
	561	0.2	10	17	19	15 (78.9)	3 (15.8)
Histone H2B-EGFP	-	-	-	-	7	7 (100)	7 (100)
	488	0.1	5	17	5	1 (20)	0 (0)
	488	0.1	10	17	5	5 (100)	0 (0)

Table 2. Full-term development of transferred embryos following super-resolution imaging

Injected mRNA	Optical wavelength (nm)	Optical intensity (mW)	Time intervals (min)	Imaging period (hour)	No. of embryos examined	No. (%) of 2-cell embryos	No. of recipients	No. (%) of Pups
Histone H2B-mCherry	-	-	-	-	27	26 (96.3)	6	8 (29.6)
	561	0.1	5	17	13	10 (76.9)	3	1 (7.7)
	561	0.1	10	17	15	15 (100)	3	7 (46.7)

No.	Devices	Manufacturer	Model
1	Inverted microscope	Olympus	IX-73
2	Nipkow disk Confocal unit	Yokogawa electric	CSU-W1 SoRa
3	sCMOS camera	Photometrics	Prime 95B
4	Z motor	Ludl	Mac5000
5	XY auto stage	Sigmakoki	BIOS-206T
6	CO_2 incubator	Tokai hit	MI-IBC
7	Software	https://micro-manager.org	Micro-Manager
8	Lasers	Coherent	405 nm 488 nm 561 nm 640 nm
9	Laser combiner	Sigmakoki	-
10	Vibration isolation table	Nippon boushin industry	SK-AS-1875T

Figure 1.

Figure 2.

A

IVF mRNA injection Live-cell imaging

B

Figure 3.

A

| | W1 (Digital zoom 1.6x) | SoRa | SoRa Deconvoluted | Enlarged W1 | SoRa | SoRa Deconvoluted |

Metaphase

B

Interphase (2cell)

Figure 4.

Figure 5.

Figure 6.

Figure 7.