dGPredictor: Automated fragmentation method for metabolic reaction free energy

prediction and *de novo* pathway design

Lin Wang[1,¶], Vikas Upadhyay[1,¶], Costas D. Maranas[1*]

Department of Chemical Engineering, The Pennsylvania State University, University Park, PA 16803, USA.

1 Department of Chemical Engineering, Pennsylvania State University, University Park, Pennsylvania

* Corresponding author

E-mail: costas@psu.edu

[¶] These authors contributed equally to this work

Abstract

Group contribution (GC) methods are conventionally used in thermodynamics analysis of metabolic pathways to estimate the standard Gibbs free energy change ($\Delta_r G'^o$) of enzymatic reactions from limited experimental measurements. However, these methods are limited by their dependence on manually curated groups and inability to capture stereochemical information, leading to low reaction coverage. Herein, we introduce an automated molecular fingerprint-based thermodynamic analysis tool called dGPredictor that enables the consideration of stereochemistry within metabolite structures and thus increases reaction coverage. dGPredictor has a higher prediction accuracy as compared to existing GC methods and can capture free energy changes for isomerase and transferase reactions, which exhibit no overall group changes. We also demonstrate dGPredictor's ability to predict the Gibbs free energy change for novel reactions and seamless integration within *de novo* metabolic pathway design tools such as novoStoic. This enables performing a thermodynamic analysis for synthetic pathways, thus safeguarding against the inclusion of reaction steps with infeasible directionalities. To facilitate easy access to dGPredictor, we developed a graphical user interface to predict the standard Gibbs free energy change for reactions at various pH and ionic strengths. The tool allows customized user input of known metabolites as KEGG IDs and novel metabolites as InChI strings (https://github.com/maranasgroup/dGPredictor).

Author summary

The genome-scale metabolic networks consist of a large number of biochemical reactions interconnected in a complex system. The standard Gibbs free energy change is commonly used to check for the feasibility of enzyme-catalyzed reactions as thermodynamics plays a crucial role in pathway design for biochemical synthesis. The group contribution methods using expert-defined functional groups have been extensively used for estimating standard Gibbs free energy change with limited experimental measurements. However, current methods using functional groups have major issues that limit its ability to cover all the metabolites and reactions as well as the inability to consider stereochemistry leads to erroneous estimation of free energy that undergoes only stereochemical change such as isomerases. Here,

we introduce a molecular fingerprint-based thermodynamic tool dGPredictor that enables stereochemistry

in metabolites and thus improves the reaction coverage with higher prediction accuracy compared to

current GC methods. It also allows the ability to predict free energy change for novel reactions which can

aid the *de novo* metabolic pathway design tool to ensure the reaction feasibility. We apply and test our

method on reactions in the KEGG database and isobutanol synthesis pathway. In addition, we provide an

open-source user-friendly web interface to facilitate easy access for standard Gibbs free energy change of

reactions at different physiological states.

Introduction

Thermodynamics determines both the directionality and efficiency of enzymatic reactions, thereby

dictating metabolic phenotypes for product synthesis in microbial engineering platforms. For example, the

beta-oxidation pathway can be reversed because the overall standard Gibbs free energy change in the

reverse direction becomes negative when utilizing ferredoxin as the reducing equivalent [1]. This

engineered pathway enables the production of higher-chain linear alcohols and fatty acids with greater

efficiency. The biological methanogenic and acetogenic reduction pathways are highly efficient in

converting CO_2 to CH_4 due to lower thermodynamic barriers compared to the corresponding geochemical

pathways [2]. Thus, thermodynamic analysis is an important tool for assessing and selecting feasible

heterologous metabolic pathways [3] and quantifying the thermodynamic driving force for biosynthesis

in different production hosts using intracellular metabolomic data [4]. However, direct experimental

measurements of standard Gibbs free energy change of reactions ($\Delta_r G'^o$) are still limited to ~600

enzymatic reactions cataloged in the Thermodynamics of Enzyme-catalyzed Reactions Database

(TECRDB) [3]. Emerging isotopic labeling experiments such as deuterium-labeled studies [5] can

directly quantify $\Delta_r G'^o$ of enzymatic reactions but have so far been limited to central carbon metabolism

thus necessitating the use of predictive computational frameworks.

In order to expand the prediction of $\Delta_r G'^\circ$ for reactions lacking experimental measurements, group

contribution (GC) methods were developed. They assume that the $\Delta_r G'^\circ$ can be expressed as the sum of

contributions from all functional groups ($\Delta_g G^\circ$) (based on a predefined list of substructures [6]), which

in turn were fitted from experimental data. Multiple linear regression is typically applied to determine

each group's contribution in a reaction by minimizing the mean squared error between the observed and

estimated $\Delta_r G^\circ$. Various group contribution methods were developed to improve the prediction accuracy

[7]. Table 1 contains a brief description of existing methods for $\Delta_r G'^\circ$ estimation.

Table 1. Existing methods for the prediction of standard Gibbs free energy of biochemical reactions

Algorithm/Method	Citation	Major contributions
Group contribution	Mavrovouniotis et al. (1990)	• Estimate free energy of formation by decomposing compounds into functional groups based on biochemical knowledge
Group contribution	Jankowski et al. (2008)	• Free energy estimation for biochemical reactions in *E. coli* metabolic network and improved previous method by introducing additional groups and interaction factors
Group contribution	Noor et al. (2012)	• Improved last method by integrating known thermodynamic data for molecules when available • Considered pseudoisomers of molecules at different protonation levels to accounts for the effect of pH and ionic strength
eQuilibrator	Flamholz et al. (2012)	• User-friendly web-based platform based on previous group contribution method by Noor et al.
Component contribution	Noor et al. (2013)	• Prioritize on reactant contribution over group contribution, which directly uses the formation energy when available
Updated group-contribution	Du et al. (2018)	• Improved previous methods using entropy and enthalpy information • Considered influence of magnesium binding with metabolites and temperature

Despite the extensive efforts to improve group contribution predictions, the approach itself suffers from

many inherent limitations [7]. The expert-defined groups provide incomplete coverage leading to (i)

metabolites that cannot be decomposed, and thus their $\Delta_f G^o$ cannot be estimated, (ii) an assignment of

zero for $\Delta_r G^o$ of reactions with no group changes despite experimental values being non-zero (such as

isomerase reactions), and (iii) large uncertainties in $\Delta_r G^o$ for reactions that involve groups occurring

sparingly in the training dataset.

Moving beyond simple molecular groups, more than 11,145 descriptors have been developed for

extracting chemical information for describing Quantitative Structure-Activity Relationships (QSAR)

[13]. Molecular descriptors such as molecular [14] and circular fingerprints [15] use fragments/moieties

to represent substructure information as vectors, matrices, or other mathematical representations such as

encoding group information in GC methods. They were applied extensively, alongside statistical and

machine learning algorithms, to predict physical and (bio-)chemical properties (such as dissociation

constant, viscosity, and toxicity) for drug design [16]. However, their application for $\Delta_r G'^o$ estimation of

enzymatic reactions has been limited to only a few studies [17,18]. The computation tool IGERS [17]

predicts $\Delta_{\square} \square'^{\square}$ of a new reaction based on its similarity (calculated using 2D molecular descriptors) to

reference reactions with experimental measurements. More recently, a machine learning algorithm using

chemical fingerprints as a feature [18] showed improved prediction accuracy.

Chemical substructures have been used extensively to encode molecular information in many

computational *de novo* pathway design tools [19]. Substructure changes in enzymatic reactions can be

generalized as rules, thereby codifying *de novo* reactions to fill in missing chemical conversion steps.

However, current *de novo* pathway design tools allow only *a posteriori* evaluation of thermodynamic

feasibility of a proposed metabolic pathway as novel reaction steps are generally treated as being

reversible [20]. Therefore, significant computational resources may be expended in generating pathways

with one or more steps operating in a thermodynamically unfavorable direction. Even though tilting the

relative reactant/product concentrations can maintain feasibility of the designed pathway, the imposed

concentration ranges may not be physiologically viable. Furthermore, operating near thermodynamic

equilibrium dramatically increases the required enzyme level, incurring a significant metabolic burden

[10,21]. Thus, an automated approach for estimating the $\Delta_r G'^o$ of all novel steps is required, which can

also be easily embedded within pathway design tools such as novoStoic [22]. This would help refine *de*

novo predictions by constraining the reaction steps/rules in only the thermodynamically feasible direction.

Herein, we developed a moiety-based automated fragmentation tool called dGPredictor for $\Delta_r G'^o$

estimation of enzymatic reactions. Moieties are descriptors of the bonding environment of all non-

hydrogen atoms in a chemical structure. They differ from functional groups, which are a set of connected

atoms responsible for a shared characteristic such as chemical nomenclature, reactivity, and bond types

(single, double, etc.) [23]. Multiple models were tested within dGPredictor using moiety descriptions of

different spans and using both explainable linear regression and neural network-based nonlinear

formalisms. Resultant prediction accuracies and overfitting potential were calculated as the mean squared

error (MSE) over training data and median of mean absolute error (MAE) of leave-one-out-cross-

validation (LOOCV) results, respectively, which were used as two metrics to compare the prediction

accuracies for $\Delta_r G'^o$ estimates and facilitate the direct comparison with widely used GC method

(specifically, component contribution (CC) [10]). dGPredictor improved the accuracy of prediction over

the widely used CC method [10] by 78.76% (i.e., MSE over training data from TECRDB database (see

Table 2)). It also led to an increase in the coverage of $\Delta_f G'^o$ and $\Delta_r G'^o$ estimation for metabolites and

reactions present in the KEGG database by 17.23% and 102%, respectively, over GC by allowing for

stereochemical considerations not captured in previous expert-defined chemical groups. Examining the

sensitivity of model predictions to moiety definition revealed that moieties spanning a bonding distance

of two are more prone to overfitting as compared to a one due to combinatorial explosion of unique

moiety types (e.g., cross-validation MAE of 5.83 vs. 15.46 for bonding distance one and two,

respectively). Moieties spanning distances one and two in a combined linear model were found to be less

prone to overfitting than the corresponding non-linear variants (cross-validation MAE 5.48 vs 7.27 for the

best performing non-linear NN model). This is likely due to the relatively small size (i.e., 4,001 reactions

and 673 metabolites) of the training dataset that does not seem to benefit from using a more extensive set

of descriptors embedded in a non-linear modeling framework.

Next, we employed dGPredictor to aid metabolic pathway design by improving upon the current practice

of considering *de novo* reactions as being always reversible, thereby necessitating additional scrutiny to

ensure thermodynamic feasibility. We show that dGPredictor can estimate the $\Delta_r G'^o$ of *de novo* reactions

(i.e., involving novel metabolites) by utilizing their chemical structure information in the IUPAC

International Chemical Identifier (InChI) strings [24]. Furthermore, pathway design tools can use the

moiety changes in dGPredictor as reaction rules to ensure the thermodynamic feasibility.

Results

Automated fragmentation of metabolites

In contrast to the widely used GC methods that rely on expert-defined groups, we apply an automated

fragmentation approach similar to Carbonell et al. [14] to encode novel molecules. dGPredictor classifies

every (non-hydrogen) atom in a structure by assessing its bonding environment at a distance of one (M1

moieties) or two bonds (M2 moieties). We do not consider bonding distance zero (M0 moieties) and more

than two (M2+ moieties), as M0 only encode the element associated with each atom and to avoid

overfitting with M2+ moieties as the available regression dataset was relatively limited in size. We use

(+)-epi-isozizaene as an example to demonstrate the chemical moieties-based metabolite descriptors used

in dGPredictor. Fig. 1A shows the seven chemical moieties that are generated from (+)-epi-isozizaene

when stereochemistry is considered. The occurrence of the seven moieties is counted and used as the

group/moiety incidence matrix $G_{i,g}$ (shown in blue, where rows denote moieties, and the columns count

the occurrence of each moiety in the chemical structure and corresponding Simplified molecular-input

line-entry system (SMILES) representation). The moiety shown in orange was obtained from the

SMILES annotation C[C@@H](C)C, where the chiral specification (i.e., "C@@H") indicates the carbon

atom is the tetrahedral center. The two different symbols, "@" and "@@" indicate anticlockwise and

clockwise configurations, respectively. Without stereochemistry, (+)-epi-isozizaene can be decomposed

into only five moieties (see Fig. 1B). The moiety shown in green is now represented by the SMILES

string CC(C)C without chiral specifications. Thus, including stereochemistry-based decomposition within

dGPredictor allows incorporating moiety changes for reactants and products that differ only in

stereochemistry (e.g., isomerases). Fig. 1A and B highlight the possible loss of stereochemical

information that occurs when not considering the two possible configurations around an asymmetric

tetrahedral atom, which results in the same moiety description for both configurations.

It is worth noting that the recent GC method, component contribution [10] cannot decompose (+)-epi-

isozizaene due to the incomplete coverage of all substructures in CC. As shown in Fig. 1, dGPredictor can

encode metabolites with both very complex structures as well as very small chemicals such as

hydroxylamine (NH_2OH). We applied dGPredictor to construct the group/moiety incidence matrix for the

673 metabolites with experimental measurements in TECRDB (see Fig. 1C). For moieties spanning a

bonding distance of one and two, we generated 263 and 1,380 members, respectively, using the RDkit

python package (See Methods section 'Automated fragmentation of metabolites').

Figure 1. The substructures/chemical moieties of (+)-epi-isozizaene decomposed by dGPredictor (A)
with stereochemistry consideration and (B) without stereochemistry consideration, and (C) group/moiety
incidence matrix created by dGPredictor for all metabolites in TECRDB.

Improved goodness of fit of   by dGPredictor using an automated moiety classification

method

We apply Bayesian ridge regression [25,26] and feedforward neural networks [27,28] to estimate the

standard Gibbs free energy contributions of the moieties. Moiety decomposition and the regression

models employed are described in detail in the Method section. Note that as experimental data was

collected at different pH and ionic strength, a mixture of pseudoisomers (i.e., multiple structures with

different protonation states) exists for each metabolite. We used the Inverse Legendre Transform (see

details in Methods section 'Pseudoisomer') to standardize the experimental values of   at pH 7 and

ionic strength 0.25 M [29]. Inverse Legendre Transform reduces the ensemble of pseudoisomers to a

single pseudoisomer (i.e., the most abundant pseudoisomer at pH 7) and was used as the reference for

regression analysis in this section. The difference between $\Delta_r G^o$ calculated for a single pseudoisomer and

the transformed Gibbs energy $\Delta_r G'^o$ (i.e., Gibb's energy of the ensemble) at a specific cellular condition

is corrected as a function of the dissociation constant *pKa* for each pseudoisomer, pH, and the ionic

strength when making predictions (see details in Methods section 'Pseudoisomers').

First, we applied Bayesian ridge regression to estimate the individual contribution of each moiety from

available experimental measurements of $\Delta_r G^o$. We chose to assess moieties associated with a bonding

distance one and two to select the model with better prediction accuracy. In the remainder of the paper,

we refer to the models of distance one and two as M1-linear (263 unique moieties for 4,001 reactions) and

M2-linear (1,380 moieties for 4,001 reactions), respectively. In addition, we examined the efficacy of

using moieties of both distances simultaneously within a single model referred to as the M1,2-linear

(1,643 moieties for 4,001 reactions). Model prediction accuracy is defined as the mean squared error

(MSE) over the entire training dataset. For the M2-linear model, MSE improved by 35.77% over the M1-

linear (see Table 2), which is expected as the number of moieties increases >5-fold from 263 to 1,380. We

found that combining moieties of radius one and two in model M1,2-linear further lowered the MSE by

60.97% compared to the M2-linear. Model overfitting potential in the cross-validation analysis is defined

as the mean absolute error (MAE) $|\Delta_r G^o_{obs} - \Delta_r G^o_{est}|$ between the experimentally observed Gibbs free

energy change of a reaction (i.e., $\Delta_r G^o_{obs}$ in the validation dataset) and the predicted value from the linear

regression model trained without the experimental data of that particular reaction (i.e., $\Delta_r G^o_{est}$ in the

training dataset). This metric can be used to flag model overfitting by comparing MAE values between

models – a larger MAE value on the validation data alongside a lower MSE value on the training data

indicates overfitting. However, a leave-one-out cross-validation (LOOCV) analysis of all models revealed

over-fitting by the M2-linear as compared to the other two linear models. We found a significant increase

in MAE for M2-linear compared to M1-linear and M1,2-linear models, with a median MAE of 15.46,

5.83, and 5.48, respectively (see Table 2). The M1,2-linear model provides the lowest MSE on the

training dataset with a value of 9.60 compared to 38.30 for M1-linear and 24.60 for M2-linear and a

significantly lower overfitting potential. Therefore, we choose the M1,2-linear as the best among the three

linear models. Do note that this should not be interpreted as a result that would hold universally. It is

likely due to the small training set size (i.e., 4,001 reactions) compared to the total number of possible

moieties.

Table 2. Details of prediction accuracy and cross-validation scores for different regression models

Model	Mean squared error	cross-validation: Mean absolute error
M1-linear	38.30	5.83
M2-linear	24.60	15.46
M1,2-linear	**9.60**	**5.48**
M1-nonlinear	20.81	6.64
M2-nonlinear	6.92	14.69
M1,2-nonlinear	6.95	7.27
GC [10]	**45.20**	**5.32**

Introducing neural network-based nonlinear moiety contribution leads to a marginal

increase in prediction performance

We next explored whether the inherent linearity of moiety contributions in the developed models limits

their predictive capability. We used a feed-forward multi-layer neural network to allow for a nonlinear

description of moiety contribution for predicting $\Delta_r G^o$. Similar to linear models, three models with

varying moiety spanning distance (same as linear regression models in the above section) were generated

to determine the effect of moiety description on prediction accuracy - M1-nonlinear, M2-nonlinear, and

M1,2-nonlinear. Similar to the linear models, $\Delta_r G^o$ prediction accuracy (i.e., a lower MSE value)

improved with increasing bonding distance (see Table 2). Models M2-nonlinear and M1,2-nonlinear were

nearly identical in their prediction accuracy within the training datasets (MSE 6.92 vs. 6.95) and

outperformed the M1-linear model (>3-fold higher MSE 20.81). The LOOCV MAE was lower for the

M1,2-nonlinear model than the M2-nonlinear (7.27 vs. 14.69, see Table 2), indicating that the former is

the best-performing nonlinear model.

A comparison among the six models (three linear and three nonlinear) suggests MSE values for all three

nonlinear models were somewhat lower than their linear counterparts. However, a correspondingly higher

LOOCV MAE revealed that the increased accuracy was likely due to overfitting. M1,2-nonlinear has a

slightly lower MSE compared to the M1,2-linear (6.95 vs. 9.60). However, cross-validation scores

showed that the M1,2-linear is significantly less susceptible to overfitting with MAE 5.48 vs. 7.27. This

implies that the extra complexity and lack of interpretability associated with the M1,2-nonlinear model do

not come with any appreciable increase in prediction performance. Since the MSE of both models is

comparable, we chose the model that is least prone to overfitting as our overall best model, i.e., the M1,2-

linear, and all subsequent results were computed using this model.

Finally, we assessed model performance between the automated decomposition scheme using moieties

proposed herein and expert-based groups. We applied the same set of thermodynamics data compiled by

Noor et al. [10] (available at https://github.com/eladnoor/component-contribution/) on dGPredictor and

the GC method to enable direct comparison. dGPredictor using the M1,2-linear model outperformed GC,

with a lower MSE (9.60 vs 45.20) and a higher R^2 score (0.9998 vs 0.9989) (Fig. 2A) estimated over the

entire training dataset. This can be attributed to the underlying energy contribution of groups/moieties

estimated by the two competing methods. As evaluated by Du et al. [7], the linear regression-based GC

model often predicts unrealistically large $|\Delta_g G^o|$ for functional groups with limited representation in the

training data which was observed in GC that predicted a $|\Delta_g G^o|$ value greater than 1,500 kJ/mol for two

groups (Fig. 2B), while the largest value in dGPredictor is ~200 kJ/mol. dGPredictor mitigates this issue

by applying regularization in the regression analysis. It must be noted that the median MAE from GC and

dGPredictor is 5.32 kJ/mol and 5.48 kJ/mol respectively (Fig. 2C) indicating similar overfitting potential.

(A) Regression analysis

(B) Group information

(C) Cross validation

Figure 2. Comparison between group contribution method by Noor et al. [10] and dGPredictor (best model: M1,2-linear). (A) Comparison of regression analysis for the best dGPredictor model (M1,2-linear model) and previous group contribution method. It shows the improvement using two metrics mean squared error (MSE improved from 45.20 to 9.60) and R-squared (increased from 0.9989 to 0.9998). (B) Histogram denoting the estimated contribution of groups. dGpredictor group contribution estimates do not predict unrealistically high values, unlike group contribution. (C) leave-one-out cross-validation results, which indicates very close performance with the previous method with an improved MSE value.

Increased coverage of metabolites and reactions by dGpredictor

A rigorous test of any predictive model lies in its ability to generalize over unseen data. Thus, we used the KEGG database [31] to test dGPredictor's scope and prediction capability for reactions that are not included in the training dataset. This would also help evaluate the model's ability to provide genome-scale coverage and estimate $\Delta_r G^o$ for novel reactions when designing *de novo* biochemical pathways. The KEGG database consists of 15,278 metabolites and 7,053 reactions with chemical structure information defined by InChI strings. We first compared the ability of the GC method and the automated fragmentation method of dGPredictor to describe all metabolites present in the database as chemical groups/moieties. The GC method could describe 85.3% of the 15,278 metabolites, while dGPredictor succeeded in covering all metabolites (see Table 3). The GC method missed 14.7% metabolites due to them containing unique substructures which are not included in its expert-defined 163 groups. Fig. 3A illustrates a few of these unique moieties and the corresponding dGPredictor decomposition. Most of the substructures that were not covered in GC methods decomposition involved bonds with N, P, and S atoms. Next, the $\Delta_r G^o$ of reactions in the KEGG database was calculated using dGPredictor and GC. Metabolites from the TECRDB database [32] were decomposed using expert-defined GC groups and the automated moiety-based framework in dGPredictor and used to calculate $\Delta_r G^o$. GC could successfully describe 33.8% of the database reactions, while dGPredictor could parameterize twice that number (i.e., 69.3%), indicating that the developed formalism can successfully generalize to provide larger reaction coverage.

Table 3. Statistics of 15,278 metabolites and 7,053 reactions from KEGG that can be decomposed into groups/moieties.

	GC	dGPredictor
Coverage of KEGG metabolites	85.3% (13,032/15,278)	100% (15,278/15,278)
Coverage of KEGG reactions	33.8% (2,385/7,053)	69.3% (4,887/7,053) (using moieties from TECRDB training data) 100% (using moieties from KEGG metabolites)

Figure 3. (A) Unique substructures in the 14.7% metabolites that are covered by dGPredictor. Here the count indicates the occurrence of such substructures in reactions (B) Example of a reaction with groups (i.e., group "C-N-N") missing from the 263 groups from TECRDB.

Even though dGPredictor improved the coverage of KEGG metabolites and their corresponding reactions,

there are still metabolites with associated moieties absent from the TECRDB dataset, leading to

incomplete moiety coverage during model training. For example, the reaction shown in Fig 3B represents

six group changes. However, group "C-N-N" is not in the list of the 1,643 moieties generated from

metabolites in the TECRDB training dataset (for both radius one and two). To warrant the lack of

information in GC, Noor et al. [10] used a standard deviation of 10^{10} J/mol to indicate that it cannot

estimate $\Delta_r G^o$ with any level of certainty. Contrary to the wide confidence interval used in GC,

dGPredictor uses Bayesian ridge regression to provide a narrower $\Delta_r G^o$ range. Briefly, it assumes an

isotropic Gaussian distribution with precision parameter α and identity matrix I for the standard Gibbs

free energy contributions of the groups ($\Delta_g G^o$):

$$p(\Delta_g G^o | \alpha) = N(0, \alpha^{-1} I)$$

After the hyperparameter optimization during model training (see Methods section 'Bayesian ridge

regression to determine the Gibbs free energy of groups $\Delta_g G^o$'), maximizing the log of the posterior

distribution in Bayesian ridge regression yields the optimized precision parameters α as $9.023 \cdot 10^{-4}$.

The standard deviation of $\Delta_g G^o$ in dGPredictor can thus be computed as $\sigma = \sqrt{\alpha^{-1}} = 33.29 \, J/mol$. As

shown in Figure 2B, $\Delta_g G^o$ of 1,643 moieties are well-determined by the normal distribution because the

values for 99% of the moieties fall within three standard deviations of the mean (i.e., $\Delta_g G^o \in$

$[-99.87, 99.87]$). Therefore, Bayesian ridge regression provides a much narrower confidence interval for

all moieties with no prior thermodynamic information. The model predictions can thus increase to 100%

reaction coverage with Bayesian regression, although with a wider confidence interval in $\Delta_r G'^o$ for

reactions that contain moieties without experimental data. Hence, the use of an automated moiety-based

description with Bayesian regression allows expanding the set of Gibbs free energy predictions while

narrowing the overall confidence interval as compared to the previous GC method.

dGPredictor enhances prediction scope by accounting for reactions with no GC-defined group changes

A limitation of previous GC methods lies in their inability to predict $\Delta_r G'^o$ of reactions with no group changes, which are ultimately assigned a $\Delta_r G'^o$ value of zero. These reactions mainly belong to classes such as isomerase and transaminase, which are known to have a $\Delta_r G'^o$ far from zero [7]. One such example is the enzymatic reaction catalyzed by GDP-D-mannose 3,5-epimerase (EC number: 5.1.3.18), with an experimentally measured equilibrium constant K' = 1.94, implying a $\Delta_r G'^o$ of 1.7 kJ/mol [33]. The reactant and product, GDP-mannose and GDP-L-galactose, respectively, are structural isomers (Fig. 4B), due to which GC-based methods are unable to capture a group change in the reaction. dGPredictor accounts for the inherent stereochemical changes by capturing the clockwise and anticlockwise configurations of the two chiral centers (Fig 4B). The predicted $\Delta_r G'^o$ is 1.74 kJ/mol, which is quite close to the experimental measurement of 1.7 kJ/mol.

Figure 4. (A) The number of reactions that have no group changes defined by GC (blue) and resolved by dGPredictor (orange), (B) GDP-mannose 3,5-epimerase (R00889) reaction that can be resolved to give group change using stereochemistry information, and examples of reactions in EC 2 (C) and EC 5 (D) that cannot be resolved by dGPredictor using only bonding distance one.

319 reactions in the KEGG database are associated with no group changes, as defined by GC-based

methods (Fig. 4A). Most of them are transferases (EC 2) and isomerases (EC 5), which cannot be

captured due to no group changes from the 163 expert-defined groups. dGPredictor described 86.83% of

the reactions with no group changes (i.e., 277/319) and, in particular, 39.71% isomerases (i.e., 110/277).

This is because the simultaneous inclusion of moieties spanning distances one and two allows us to

consider additional details of the localized bonds and atoms, thus alleviating problematic cases of no

moiety changes being registered when considering single bonding distance. For example, the moiety

description of an aminotransferase reaction (EC 2) (Figure 4C) generates identical moieties for both

substrates and products, leading to an empty group change vector when considering bonding distance one

moieties. However, allowing an additional bonding distance of two helps dGPredictor generate unique

moieties, thus resolving the issue of zero change and in turn leading to a non-zero $\Delta_r G'^o$ value.

Nevertheless, 42 reactions could still not be resolved because both M1 and M2 descriptions used by

dGPredictor were identical between reactants and products. This is because the RDkit tool [34] used for

computing moieties stores molecules with implicit hydrogen (i.e., not explicitly present in the molecular

graph) and ignores the "C-H" bond when generating the SMILES string. Thus, cannot identify the chiral

center when the "C-H" bond is responsible for tetrahedral stereochemistry which leads to same SMILES

string for different moieties and a loss of stereochemical information for reactions using M1 moiety

change. An example is shown in Fig. 4D, where KEGG reaction R10764 that converts α-L-Fucopyranose

to β-L-Fructopyranose is expected to have non-empty group changes because the moieties have different

stereo configurations (i.e., [C@H] or [C@@H]) for bonding distance one. In this case, RDkit returning an

empty group can be alleviated by combining both distances to capture a non-zero moiety change. This

flexible moiety consideration is the primary reason behind dGPredictor providing ~87% more coverage

for reactions with no group/moiety change compared to previous GC methods and the remaining 13%

could be tackled using more advanced molecular descriptors such as Neural Graph Fingerprints [15] that

account for higher-order interactions instead of local atom/bond information.

348 **User-friendly interface for Gibbs free energy calculation of novel reactions and metabolites**

349 **beyond KEGG**

350 Advances in computational pathway design have expanded the range of microbial product synthesis to

351 non-natural synthetic molecules and drug precursors, leveraging broad-substrate range enzymes [35,36]

352 promiscuous activity [37], and even *de novo* enzyme design [38,39]. However, tools such as novoStoic

353 [40] generally treat novel transformations as being reversible, necessitating additional scrutiny to ensure

354 the thermodynamic feasibility of the designed pathway. The ability of dGPredictor to characterize unseen

355 metabolites, and by extension, chemical transformations, accompanied by an automated moiety-based

356 framework enables its use in pathway design tools to eliminate thermodynamically infeasible predictions.

357 In this section, we demonstrate how dGPredictor can predict $\Delta_r G'^o$ for novel reactions which do not span

358 known biochemical reactions using chemical moieties, and a GUI that can help users query the same.

359 The dGPredictor input format uses KEGG IDs to indicate the substrates and products of a reaction. For

360 example, dGPredictor can recognize 'C00096 <=> C02280' as the reaction "GDP-mannose <=> GDP-L-

361 galactose" (discussed in above section). However, KEGG is only one of many databases consolidating

362 biochemical reactions, has a lower metabolite content [41], and does not capture novel molecules that

363 often show up in the reactions of retrosynthetic metabolic pathways [42]. Therefore, we designed

364 dGPredictor to allow for user-defined chemical structures as an additional input to estimate $\Delta_r G'^o$ for any

365 novel   reaction.   A   user-friendly   interface   has   been   developed

366 (https://github.com/maranasgroup/dGPredictor) to facilitate thermodynamic analysis for reactions.

367 Metabolites with known chemical structures can be entered using KEGG IDs and InChI strings are

368 required for molecules not present in the KEGG database. For example, in the *de novo* pathway found by

369 novoStoic for pinosylvin (C01745) degradation, a deoxygenase enzyme catalyzes the first step and

370 produces an aromatic product (see Fig. 5). KEGG IDs identify all metabolites except the reduced product

371 'N00001' in the reaction   "C01745 + C00004 <=> N00001 + C00003 + C00001". Here, we use id

"N00001" to represent the novel metabolite 3-Phenethyl-phenol, where N refers to novel. The user can

indicate that the reaction involves a novel metabolite (i.e., absent in the KEGG database) by clicking a

checkbox and filling in the InChI string placeholder under the stoichiometry of the reaction to describe

the atom composition and bonds in metabolite N00001 (see Fig. 5). Similar to eQuilibrator [11], we allow

the customized input of intracellular conditions (i.e., pH and ionic strength) via two sliders. With all the

information properly defined, on clicking the "search" button, dGPredictor first show the chemical

structures in the reaction and then calculate the standard transformed Gibbs energy $\Delta_r G'^\circ$ of the reaction

at a particular pH and ionic strength. The output information also displays the standard deviation of

predictions from Bayesian ridge regression and the moiety changes as defined by dGPredictor. Note that

the standard transformed Gibbs energy $\Delta_r G'^\circ$ is not a function of the actual metabolite concentrations.

The measured concentrations of metabolites (c_i) can be used to calculate the actual Gibbs free energy

for a reaction j ($\Delta_r G'_j$) via the equation $\Delta_r G'_j = \Delta_r G'^\circ_j + \ln(\prod_i [c_i]^{S_{ji}})$ where, S_{ji} is the stoichiometry

of metabolite i in reaction j. We envision the developed GUI to facilitate easy adoption of dGPredictor to

the broader metabolic engineering and synthetic biology community.

Figure 5. The graphical user interface of dGPredictor. It allows the input of metabolites using their KEGG IDs (for known compounds) and also InChI strings (for novel compounds) in a chemical reaction. Intracellular conditions of pH and ionic strength can also be adjusted using sliders. The final output shows the standard transformed Gibbs energy   of the reaction at a particular pH and ionic strength.

Assigning thermodynamics-derived directionality to reaction rules in *de novo* pathway design

As dGPredictor generates automated molecular descriptions, it can be used in conjunction with pathway design tools to ensure that individual reaction, and in turn, the overall pathway is thermodynamically feasible. Pathway design tools such as novoStoic, RetroPath, and RetroPath2.0 [40,43,44] can be

integrated with the group change vector of dGPredictor as reaction rules to design *de novo* pathways. We

used the novoStoic tool to illustrate this integration and deployed dGPredictor on the 3,603 unique

reaction rules generated from KEGG database. We found that based on the predicted standard free energy

of change, a number of novel reactions can reliably be flagged as irreversible (i.e., $\left| \Delta_r G'^\circ \right| > 20$

kJ/mol). This enables the elimination of intermediate steps with a high free energy barrier, thereby

significantly reducing the number of candidate pathways to be explored.

Similar to the definition of reaction rules in novoStoic [45], which captures changes in the topology of

molecular graphs for a substrate to product conversion [46], dGPredictor assumes that reactions with the

same rule undergo same substrate to product change, thereby conforming to identical moiety change. We

estimated $\Delta_r G'^\circ$ for all 3,603 reaction rules and use ± 3 standard deviations to obtain confidence intervals

(i.e., 99.7% probability that the free energy estimate is within the calculated interval) (Fig. 6A and B). We

identified 331 reaction rules with predicted free energy confidence intervals that span only positive values

(see Fig. 6B) implying that they can only be deployed in the reverse direction. As an additional safeguard,

we only applied the irreversibility restriction if the absolute value of the predicted free energy of change

$\left| \Delta_r G'^\circ \right|$ exceeds 20 kJ/mol, to account for varying metabolite concentrations that may ultimately tilt

reaction directionality [47]. This additional constraint reduces the number of reaction rules that can

confidently be treated as irreversible from 331 to 325 (see Fig. 6C for EC classification for irreversible

reaction rules). Therefore, the dGPredictor framework can help identify the direction *a priori* for

irreversible reaction rules and eliminate the thermodynamically infeasible intermediate steps. This allows

novoStoic to consider not only the overall thermodynamic feasibility of the pathway but also evaluate

every single step.

To assess the extent of reaction rules' directionality affecting pathway design predictions, we applied the

set of reaction rules to search for isobutanol production pathways (a well-studied product from the Ehrlich

pathway [48–51]) using 2-ketoisovalerate as the precursor. Two distinct production pathways were

identified by novoStoic (Fig. 6D). Pathway A (shown in blue) is the engineered pathway in *E. coli* and *C.*

thermocellum with a demonstrated thermodynamic feasibility *in vivo*. Pathway B (shown in orange),

however, is thermodynamically infeasible. The first step converts 2-ketoisovalerate to isobutyric acid

using a novel reaction R1, which has the same group change as the native reaction of indole-3-pyruvate

monooxygenase. However, the second step is a novel reaction R2 similar to the reaction catalyzed by

nucleoside oxidase that favors the reverse direction based on dGPredictor thermodynamics analysis. Thus,

instead of relying on manual post-processing, direct inclusion of thermodynamic constraints from

dGPredictor within the novoStoic pathway design tool can help eliminate infeasible pathways.

Figure 6. Group change vectors can be used as reaction rules to design *de novo* pathways. (A) $\Delta_r G'^o$ for all the 3,603 reaction rules, (B) 331 reaction rules found to be irreversible from the uncertainty analysis using ± 3 standard deviations (see supplementary table 1 for the KEGG Id of reactions on the x-axis), (C) The EC classification for irreversible reaction rules, and (D) an example of thermodynamically infeasible pathways eliminated by using irreversible reaction rules.

Discussion

This paper presents an automated thermodynamic analysis tool, dGPredictor, based on molecular fingerprints and chemical moieties to estimate the $\Delta_r G'^o$ of biochemical reactions. Compared with previous group contribution methods, dGPredictor expands thermodynamic calculation coverage to more molecules and reactions with improved accuracy. This is primarily due to the automated fragmentation method that allows incorporating stereochemical information while generating chemical moieties/groups, which was lacking in the group decomposition scheme in previous methods [6,8–10]. The versatility of the fragmentation method is further illustrated by extending predictions to *de novo* reactions, which enables determining the thermodynamical feasibility of synthetic pathways and thus aid strain design algorithms.

First, we showed that the moiety-based automated fragmentation using the SMILES notations can account for stereochemistry in metabolite descriptions. Moieties spanning bonding distances of one and two were generated and used in a linear and non-linear regression framework to ascertain the best performing model. We found that employing a non-linear framework and moieties spanning a bonding distance of two leads to a slight increase in prediction accuracy while being significantly more prone to overfitting. Therefore, an explainable linear model comprising moieties spanning bonding distances of one and two (i.e., M1,2-linear) was determined to be the best performing model and the default for dGPredictor. Notably, the cross-validation MAE (representing overfitting) was comparable to the current state-of-the-art CC method and the MSE (representing prediction accuracy) improved by 78.76% over same training (i.e., the TECRDB database) and cross-validation (i.e., the KEGG database) datasets for a direct comparison of model predictions. We found that the proposed automated fragmentation approach

can be implemented on all metabolites in the TECRDB database, unlike the CC method, which uses

expert-defined functional groups to decompose metabolites and can only cover 85.3% of metabolites.

Molecular descriptions thus obtained were used to estimate $\Delta_r G'^o$ for reactions in the KEGG database,

leading to an increase in (69.3% vs. 33.8%). However, there remains scope for improvement in the

dGPredictor prediction pipeline, which can be aided by increasing the experimental coverage of reactions

with experimental $\Delta_r G'^o$ estimates, in turn, increasing the number of characterized moieties. dGPredictor

can be used to prioritize these experimental targets for $\Delta_r G'^o$ estimations by focusing on reactions that

comprise unique moieties frequently occur in the uncovered reactions. In addition, quantum chemical

calculation [52,53] that *de novo* estimate $\Delta_f G'^o$ for metabolites with no experimental measurements

and/or novel metabolites generated by retrosynthetic pathway design algorithms can be used to

supplement the available experimental data.

A graphical user interface was built for the dGPredictor tool, similar to the group contribution-based web

interface eQuilibrator. It can estimate $\Delta_r G'^o$ at different intracellular conditions, namely pH and ionic

strength. dGPredictor improves upon eQuilibrator by not only estimating the $\Delta_r G'^o$ for reactions with

compounds in the KEGG database and allowing the user to input novel metabolites using their InChI

strings. Thus, our tool broadens the capability of $\Delta_r G'^o$ predictions by incorporating novel reactions. This

prediction ability is extended to predict the $\Delta_r G'^o$ of designed pathways by restricting the directionality to

new reaction rules in the *de novo* pathway design tool novoStoic [40]. dGPredictor can help eliminate

unnecessary solutions in novoStoic that are not thermodynamically feasible. The moiety change vectors in

the dGPredictor can be directly used as allowed reaction rules in novoStoic [45]. dGPredictor thus

facilitates an effective search for thermodynamically feasible metabolic pathways by discarding on

average 10% of the reaction rules as infeasible, thereby reducing the search space and computational

expense for pathway design. dGPredictor can be easily generalized to allow a detailed thermodynamic

analysis of large-scale metabolic networks to understand cell metabolism better.

However, it must be noted that there still exists scope for improvement in the molecular descriptors proposed herein. dGPredictor cannot fully resolve reactions with no moiety changes in isomerase reactions due to limitations in the fingerprints defined by RDkit. Therefore, applying advanced 3D molecular descriptors to capture the exact stereochemistry might help capture such reactions. 2D molecular descriptors only contain information for the localized atoms/bonds, whereas a 3D descriptor allows precise capture of molecular shape and interactions [13]. In a recent study, the Quantitative Structure-Activity Relationship (QSAR) method based on Smooth Overlap of atomic position (SOAP) descriptors was shown to successfully capture the 3D atomic environment from conformers [54]. Therefore, utilizing a 3D molecular descriptor for molecule decomposition in the free energy prediction has great potential to account for even more accurate estimation.

Methods

Automated fragmentation of metabolites

The algorithm first decomposes the chemical structures of a metabolite into groups of adjacent atoms within a specified bonding distance (i.e., moieties). The bonding distance defines the bond's proximity to consider from an atom to be described as chemical moieties. We use the InChI string of metabolites as input. Next, we represent each fragment/group with canonical SMILES string. Finally, a group incidence matrix $G_{i,g}$ is used to represent the count of each moiety as group g for every metabolite i. We use the automated fragmentation algorithm and represent each group by a unique SMILES string utilizing the Cheminformatic tool RDkit[34] accessed through python.

Bayesian ridge regression to determine the Gibbs free energy of groups $\Delta_g G^o$.

Using the molecular descriptions thus obtained, we determine the standard Gibbs free energy contributions of groups $\Delta_g G^o$ that allows the best fit of experimental data from TECRDB.

Parameters

S: stoichiometric matrix ($R^{i \times j}$)

G: group incidence matrix that represents group decomposition ($R^{i \times g}$)

$\Delta_r G^o_{obs,j}$: Gibbs free energy change of reaction j observed in enzyme thermodynamic database TECRDB.

Variables

$\Delta_g G^o$: standard Gibbs free energy contributions of groups ($R^{g \times 1}$)

$\Delta_r G^o_{est}$: estimated standard Gibbs free energy change of reactions ($R^{n \times 1}$)

The optimization formulation of multi-linear regression with L2 regularization (i.e., ridge regression) to

estimate $\Delta_g G^o$ by minimizing the sum of the squared estimate of errors (SSE) and the regularization term

is shown as follows:

$$\min \sum_j^n \left(\Delta_r G^o_{obs,j} - \Delta_r G^o_{est,j} \right)^2 + \lambda \left(\Delta_g G^o \right)^T \Delta_g G^o \tag{1}$$

s.t.

$$\Delta_r G^o_{est} = S^T G \cdot \Delta_g G^o \tag{2}$$

Because the multi-linear regression is prone to overfitting, especially since more groups are defined in our

method than previous GC-based methods, L2 regularization ($\lambda \left(\Delta_g G^o \right)^T \Delta_g G^o$) is often applied in ridge

regression to reduce cross-validation errors.

Confidence interval analysis from strongly biased models (e.g., ridge regression) relies on bootstrap-

based methods. It often produces inaccurate uncertainty estimations [55]. Credible intervals used by

Bayesian inference provide an alternative metric to confidence intervals. They have been shown to

provide more reliable uncertainty estimation in 13C metabolic flux analysis [56]. Herein, we apply

Bayesian ridge regression and define the prior of the $\Delta_g G^o$ as the isotropic Gaussian distributions with

precision parameter α:

$$p(\Delta_g G^o | \alpha) \ = \ N(\Delta_g G^o | 0, \alpha^{-1} I) \tag{3}$$

530 The likelihood function for the $\Delta_r G^o_{est}$ is defined as a Gaussian distribution with a precision parameter β:

$$\square(\Delta_r G^o_{est} | \Delta_g G^o, \beta) = N(\Delta_r G^o_{est} | S^T G \cdot \tag{4}$$

$$\Delta_g G^o, \beta^{-1})$$

531 Based on the prior and likelihood function, $\Delta_g G^o$ is then estimated by the method of maximum a posteriori

532 estimation (MAP) of the log of the posterior distribution:

$$\max_{\Delta_g G^o} ln(p(\Delta_r G^o_{est} | \Delta_g G^o, \beta) \cdot p(\Delta_g G^o | \alpha) \tag{5}$$

533 Note that the maximization of the log of the posterior distribution is equivalent to the multi-linear

534 regression with L2 regularization defined in equations (1) and $\lambda = \square / \square$ as shown in [25]:

$$ln(p(\Delta_r G^o_{est} | \Delta_g G^o, \beta) \cdot p(\Delta_g G^o | \alpha) = -\frac{\beta}{2} \sum_j^n \left(\Delta_r G^o_{obs,j} - \Delta_r G^o_{est,j} \right)^2 - \frac{\alpha}{2} \left(\Delta_g G^o \right)^T \Delta_g G^o \tag{5}$$

536 The prediction interval of $\Delta_r G^o_{est,j}$ for a new reaction j with a group change vector $\square_\square$ from Bayesian

537 ridge regression can then be calculated as:

$$p(\Delta_r G^o_{est,j} | x_j, \Delta_g G^o, \alpha, \beta) = N(x_j \cdot \Delta_g G^o, \sigma^2(x_j)) \tag{7}$$

538 where the variance of the prediction interval can be calculated following the derivation in [34] as:

539
$$\sigma^2(x_j) = 1/\beta + x_j^T S_N x_j$$

$$S_N = (\alpha + \beta (S^T G)^T S^T G)^{-1}$$

540 We apply the Bayesian ridge regression function from the scikit-learn python package to train the

541 Bayesian ridge regression model. Scikit-learn optimizes the precision parameters α and β by iterative re-

542 estimation based on an estimate for "how well-determined" the corresponding $\Delta_g G^o$ is by the training

543 data [26]. We found the same parameters and mean squared error in the fitted data upon Bayesian ridge

544 regression to 50 different initial guesses of precision parameters. We infer that the iterative method

545 applied in Bayesian ridge regression in scikit-learn can produce unique, optimized parameters α and β.

546 Then, the Bayesian ridge regression model can estimate the mean ($\square_\square \cdot \Delta_g G^o$) and standard deviation (σ)

547 of $\Delta_r G^o_{est,j}$.

548 **Neural networks models for estimating Gibbs free energy of groups $\Delta_g G^o$**

549 We choose a feed-forward multi-layer perceptron neural network for nonlinear regression. These

550 networks have neurons that are ordered in layers. The model starts with an input layer (moiety incidence

551 vector), followed by a hidden layer, and ends with an output layer. Fig. 7 shows the architecture for a

552 feed-forward neural network. The multi-layer perceptron model maps the moiety incidence matrix of the

553 training data with the output layer using nonlinear functions. The output of the hidden layers is

554 determined by using a rectified linear unit transfer function [57].

Figure 7. The architecture for multi-layer perceptron neural network with an input layer of moiety incidence matrix (j x g matrix), a hidden layer with n neurons, and an output layer of estimates with j neurons. Here *j* represents the reactions, g represents the groups/moieties, and n represents the neurons in the hidden layer.

We applied the multi-layer perceptron regression from the scikit-learn python package to train the neural network model. LBFGS (Limited memory Broyden-Fletcher-Goldfarb-Shanno) backpropagation algorithm [27] is used to minimize the mean squared error and update the weights of the hidden layers based on the estimates in the output layer. The LBFGS is a faster technique in the family of quasi-newton methods commonly used for parameter estimation in machine learning [27]. Our model has three layers: input, hidden, and output. We considered the scikit-learn package default single hidden layer with 100 neurons to build the neural network model. Studies suggest that a single layer can adequately approximate any function which maps one finite space to another [28]. Scikit-learn package automatically estimates the number of neurons based on the cross-validation results for an accurate fit.

At last, we build three different models for bonding distance one, two, and combining both distances. The

M1,2-nonlinear model inputs the moiety incidence matrix for both the radiuses in a single input matrix.

The radius one and two have 263 and 1380 unique moieties for 673 metabolites in 4001 reactions.

Therefore, the combined-NN model considers a total of 1643 moieties to generate moiety incidence. The

same leave-one-out cross-validation was performed as linear regression models for a direct comparison of

the model performance.

Pseudoisomers

The exact structure of the metabolite inside a cell is typically unknown because it exists as a mixture of

pseudoisomers with different protonation states. Pseudoisomers are incorporated into dGPredictor by

assuming that the intracellular mixture of pseudoisomers follows the Boltzmann distribution. We use

Inverse Legendre Transform [9] to calculate the difference between the standard Gibbs free energy of

formation of a compound $\Delta_f G^o$ of the major pseduoisomer at pH 7 and the transformed Gibbs free energy

of formation of the mixture $\Delta_f G'^o$:

$$\Delta_f G'^o = \Delta_f G^o + RT \ln(10) \cdot m \cdot pH - RT \ln \sum_{n=0}^{N_H} \exp \left[\ln(10) \sum_{i=m+1}^{n} (pKa(i) - pH) + \right.$$

$$\left. \frac{2.91482(z_n^2 - n)\sqrt{I}}{RT(1 + 1.6\sqrt{I})} \right] \quad (8)$$

where the acid-base dissociation constant *pKa(i)* of pseudoisomer *i* is calculated using ChemAxon

Marvin, $\square_\square$ is the maximum number protonated hydrogens within a molecule (i.e., the number of *pKas)*,

m is the number of hydrogens of the reference pseudoisomer, $\square_\square$ is the total charge the pseudoisomer *n*,

I is the ionic strength, R is the gas constant, and T is the temperature.

The experimental measurement in TECRDB for a reaction for the apparent equilibrium constants $\square'$,

which is used to calculate the standard transformed Gibbs energy of reaction *j*:

$$\Delta_r G_j'^o = -RT \ln(K') = \sum_i S_{ij} \Delta_f G'^o \quad (9)$$

Replacing $\Delta_f G'^o$ in equation (9) with equation (8), the resulting linear equations can be used to calculate

$\Delta_f G^o$ using Gaussian elimination. When predicting $\Delta_f G'^o$ of a reaction, the difference between $\Delta_f G'^o$

and $\Delta_r G^o$ (i.e., $\Delta\Delta G = \Delta_f G'^o - \Delta_f G^o$) of all the metabolites in the reaction calculated from equation (8)

can be added to the $\Delta_r G^o$ calculated from regression analysis in above sections to correct the

contribution of the pseudoisomer mixture. Both the calculation of $\Delta_f G^o$ using Inverse Legendre

Transform and $\Delta\Delta G$ are implemented using the functions from the component contribution package

(https://github.com/eladnoor/component-contribution).

References

1. Dellomonaco C, Clomburg JM, Miller EN, Gonzalez R. Engineered reversal of the β-oxidation cycle for the synthesis of fuels and chemicals. Nature. 2011;476: 355–359. doi:10.1038/nature10333

2. Yung Y, Russell M, Parkinson C. The Search for Life on Mars. J Cosmol. 2010;5: 1121–1130. Available: https://authors.library.caltech.edu/48859/

3. Dash S, Olson DG, Joshua Chan SH, Amador-Noguez D, Lynd LR, Maranas CD. Thermodynamic analysis of the pathway for ethanol production from cellobiose in Clostridium thermocellum. Metab Eng. 2019;55: 161–169. doi:10.1016/j.ymben.2019.06.006

4. McCloskey D, Xu J, Schrübbers L, Christensen HB, Herrgård MJ. RapidRIP quantifies the intracellular metabolome of 7 industrial strains of E. coli. Metab Eng. 2018;47: 383–392. doi:10.1016/j.ymben.2018.04.009

5. Xu J, Martien J, Gilbertson C, Ma J, Amador-Noguez D, Park JO. Metabolic flux analysis and fluxomics-driven determination of reaction free energy using multiple isotopes. Current Opinion in Biotechnology. Elsevier Ltd; 2020. pp. 151–160. doi:10.1016/j.copbio.2020.02.018

6. Mavrovouniotis ML. Group contributions for estimating standard gibbs energies of formation of biochemical compounds in aqueous solution. Biotechnol Bioeng. 1990;36: 1070–1082. doi:10.1002/bit.260361013

7. Du B, Zielinski DC, Palsson BO. Estimating Metabolic Equilibrium Constants: Progress and Future Challenges. Trends in Biochemical Sciences. Elsevier Ltd; 2018. pp. 960–969. doi:10.1016/j.tibs.2018.09.009

8. Jankowski MD, Henry CS, Broadbelt LJ, Hatzimanikatis V. Group contribution method for thermodynamic analysis of complex metabolic networks. Biophys J. 2008;95: 1487–1499. doi:10.1529/biophysj.107.124784

9. Noor E, Bar-Even A, Flamholz A, Lubling Y, Davidi D, Milo R. An integrated open framework for thermodynamics of reactions that combines accuracy and coverage. Bioinformatics. 2012;28: 2037–2044. doi:10.1093/bioinformatics/bts317

627   10.   Noor E, Haraldsdó ttir HS, Milo R, T Fleming RM. Consistent Estimation of Gibbs Energy Using
628   Component Contributions. PLoS Comput Biol. 2013;9: 1003098.
629   doi:10.1371/journal.pcbi.1003098

630   11.   Flamholz A, Noor E, Bar-Even A, Milo R. EQuilibrator - The biochemical thermodynamics
631   calculator. Nucleic Acids Res. 2012;40. doi:10.1093/nar/gkr874

632   12.   Du B, Zhang Z, Grubner S, Yurkovich JT, Palsson BO, Zielinski DC. Temperature-Dependent
633   Estimation of Gibbs Energies Using an Updated Group-Contribution Method. Biophys J.
634   2018;114: 2691–2702. doi:10.1016/j.bpj.2018.04.030

635   13.   Masand VH, Rastija V. PyDescriptor: A new PyMOL plugin for calculating thousands of easily
636   understandable molecular descriptors. Chemom Intell Lab Syst. 2017;169: 12–18.
637   doi:10.1016/j.chemolab.2017.08.003

638   14.   Carbonell P, Planson AG, Fichera D, Faulon JL. A retrosynthetic biology approach to metabolic
639   pathway design for therapeutic production. BMC Syst Biol. 2011;5: 122. doi:10.1186/1752-0509-
640   5-122

641   15.   Duvenaud D, Maclaurin D, Aguilera-Iparraguirre J, Gómez-Bombarelli R, Hirzel T, Aspuru-Guzik
642   A, et al. Convolutional Networks on Graphs for Learning Molecular Fingerprints. Adv Neural Inf
643   Process Syst. 2015;2015-January: 2224–2232. Available: http://arxiv.org/abs/1509.09292

644   16.   Piir G, Kahn I, García-Sosa AT, Sild S, Ahte P, Maran U. Best practices for QSAR model
645   reporting: Physical and chemical properties, ecotoxicity, environmental fate, human health, and
646   toxicokinetics endpoints. Environmental Health Perspectives. Public Health Services, US Dept of
647   Health and Human Services; 2018. doi:10.1289/EHP3264

648   17.   Rother K, Hoffmann S, Bulik S, Hoppe A, Gasteiger J, Holzhütter HG. IGERS: Inferring gibbs
649   energy changes of biochemical reactions from reaction similarities. Biophys J. 2010;98: 2478–
650   2486. doi:10.1016/j.bpj.2010.02.052

651   18.   Alazmi M, Kuwahara H, Soufan O, Ding L, Gao X. Systematic selection of chemical fingerprint
652   features improves the Gibbs energy prediction of biochemical reactions. Valencia A, editor.
653   Bioinformatics. 2019;35: 2634–2643. doi:10.1093/bioinformatics/bty1035

19.   Wang L, Dash S, Ng CY, Maranas CD. A review of computational tools for design and

reconstruction of metabolic pathways. Synthetic and Systems Biotechnology. KeAi
Communications Co.; 2017. pp. 243–252. doi:10.1016/j.synbio.2017.11.002

20.   Duigou T, Du Lac M, Carbonell P, Faulon JL. Retrorules: A database of reaction rules for
engineering biology. Nucleic Acids Res. 2019;47: D1229–D1235. doi:10.1093/nar/gky940

21.   Ng CY, Wang L, Chowdhury A, Maranas CD. Pareto Optimality Explanation of the Glycolytic
Alternatives in Nature. Sci Rep. 2019;9: 1–15. doi:10.1038/s41598-019-38836-9

22.   Wang L, Ng CY, Dash S, Maranas CD. Exploring the combinatorial space of complete pathways
to chemicals. Biochemical Society Transactions. Portland Press Ltd; 2018. pp. 513–522.
doi:10.1042/BST20170272

23.   Ertl P. An algorithm to identify functional groups in organic molecules. J Cheminform. 2017;9:
36. doi:10.1186/s13321-017-0225-z

24.   Heller SR, McNaught A, Pletnev I, Stein S, Tchekhovskoi D. InChI, the IUPAC International
Chemical Identifier. J Cheminform. 2015;7: 23. doi:10.1186/s13321-015-0068-4

25.   Bishop C. Pattern Recognition and Machine Learning. springer; 2006. Available:
https://www.springer.com/gp/book/9780387310732

26.   Tipping M. SparseBayesianLearningandtheRelevanceVector Machine.

27.   Liu DC, Nocedal J. On the limited memory BFGS method for large scale optimization. Math
Program. 1989;45: 503–528. doi:10.1007/BF01589116

28.   Heaton J. Introduction to Neural Networks for Java, 2nd Edition. 2nd ed. Heaton Research, Inc.;
2008.

29.   Alberty RA. Inverse legendre transform in biochemical thermodynamics: Illustrated with the last
five reactions of glycolysis. J Phys Chem B. 2002;106: 6594–6599. doi:10.1021/jp020764w

30.   Noor E, Haraldsdóttir HS, Milo R, Fleming RMT. Consistent Estimation of Gibbs Energy Using
Component Contributions. PLoS Comput Biol. 2013;9: 1003098.
doi:10.1371/journal.pcbi.1003098

31.   Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new perspectives on

genomes, pathways, diseases and drugs. Nucleic Acids Res. 2016;45: 353–361.

doi:10.1093/nar/gkw1092

32.   Goldberg RN, Tewari YB, Bhat TN. Thermodynamics of enzyme-catalyzed reactions--a database

for quantitative biochemistry. Bioinformatics. 2004;20: 2874–2877.

doi:10.1093/bioinformatics/bth314

33.   Barber GA, Hebda PA. GDP-D-Mannose: GDP-L-Galactose Epimerase from Chlorella

pyrenoidosa. Methods Enzymol. 1982;83: 522–525. doi:10.1016/0076-6879(82)83046-2

34.   Landrum G. RDKit: Open-source cheminformatics. 2006.

35.   Rodriguez GM, Tashiro Y, Atsumi S. Expanding ester biosynthesis in Escherichia coli. Nat Chem

Biol. 2014;10: 259–265. doi:10.1038/nchembio.1476

36.   Atsumi S, Hanai T, Liao JC. Non-fermentative pathways for synthesis of branched-chain higher

alcohols as biofuels. Nature. 2008;451: 86–89. doi:10.1038/nature06450

37.   Khersonsky O, Roodveldt C, Tawfik DS. Enzyme promiscuity: evolutionary and mechanistic

aspects. Current Opinion in Chemical Biology. Curr Opin Chem Biol; 2006. pp. 498–508.

doi:10.1016/j.cbpa.2006.08.011

38.   Siegel JB, Smith AL, Poust S, Wargacki AJ, Bar-Even A, Louw C, et al. Computational protein

design enables a novel one-carbon assimilation pathway. Proc Natl Acad Sci U S A. 2015;112:

3704–3709. doi:10.1073/pnas.1500545112

39.   Khersonsky O, Kiss G, Röthlisberger D, Dym O, Albeck S, Houk KN, et al. Bridging the gaps in

design methodologies by evolutionary optimization of the stability and proficiency of designed

Kemp eliminase KE59. Proc Natl Acad Sci U S A. 2012;109: 10358–10363.

doi:10.1073/pnas.1121063109

40.   Kumar A, Wang L, Ng CY, Maranas CD. Pathway design using de novo steps through uncharted

biochemical spaces. Nat Commun. 2018;9: 1–15. doi:10.1038/s41467-017-02362-x

41.   Ebastien Moretti S´, Martin O, Van T, Tran D, Bridge A, Morgat A, et al. MetaNetX/MNXref-

reconciliation of metabolites and biochemical reactions to bring together genome-scale metabolic

networks. Nucleic Acids Res. 2015;44: 523–526. doi:10.1093/nar/gkv1117

42.   Lin GM, Warden-Rothman R, Voigt CA. Retrosynthetic design of metabolic pathways to
chemicals not found in nature. Current Opinion in Systems Biology. Elsevier Ltd; 2019. pp. 82–
107. doi:10.1016/j.coisb.2019.04.004

43.   Carbonell P, Parutto P, Baudier C, Junot C, Faulon JL. Retropath: Automated pipeline for
embedded metabolic circuits. ACS Synth Biol. 2014;3: 565–577. doi:10.1021/sb4001273

44.   Delépine B, Duigou T, Carbonell P, Faulon JL. RetroPath2.0: A retrosynthesis workflow for
metabolic engineers. Metab Eng. 2018;45: 158–170. doi:10.1016/j.ymben.2017.12.002

45.   Kumar A, Wang L, Ng CY, Maranas CD. Pathway design using de novo steps through uncharted
biochemical spaces. Nat Commun. 2018;9: 1–15. doi:10.1038/s41467-017-02362-x

46.   Kumar A, Maranas CD. CLCA: Maximum Common Molecular Substructure Queries within the
MetRxn Database. 2014 [cited 26 May 2020]. doi:10.1021/ci5003922

47.   Chowdhury A, Maranas CD. Designing overall stoichiometric conversions and intervening
metabolic reactions. Sci Rep. 2015;5: 16009. doi:10.1038/srep16009

48.   Hazelwood LA, Daran JM, Van Maris AJA, Pronk JT, Dickinson JR. The Ehrlich pathway for
fusel alcohol production: A century of research on Saccharomyces cerevisiae metabolism. Applied
and Environmental Microbiology. American Society for Microbiology (ASM); 2008. pp. 2259–
2266. doi:10.1128/AEM.02625-07

49.   Wang L, Ng CY, Dash S, Maranas CD. Exploring the combinatorial space of complete pathways
to chemicals. Biochemical Society Transactions. Portland Press Ltd; 2018. pp. 513–522.
doi:10.1042/BST20170272

50.   Wess J, Brinek M, Boles E. Improving isobutanol production with the yeast Saccharomyces
cerevisiae by successively blocking competing metabolic pathways as well as ethanol and glycerol
formation. Biotechnol Biofuels. 2019;12: 173. doi:10.1186/s13068-019-1486-8

51.   Matsuda F, Ishii J, Kondo T, Ida K, Tezuka H, Kondo A. Increased isobutanol production in
Saccharomyces cerevisiae by eliminating competing pathways and resolving cofactor imbalance.
Microb Cell Fact. 2013;12: 119. doi:10.1186/1475-2859-12-119

52.   Aprà E, Bylaska EJ, De Jong WA, Govind N, Kowalski K, Straatsma TP, et al. NWChem: Past,
present, and future. J Chem Phys. 2020;152: 184102. doi:10.1063/5.0004997

53.   Jinich A, Rappoport D, Dunn I, Sanchez-Lengeling B, Olivares-Amaya R, Noor E, et al. Quantum
737   chemical approach to estimating the thermodynamics of metabolic reactions. Sci Rep. 2014;4.
738   doi:10.1038/srep07022

739   54.   Mccorkindale W, Poelking C, Lee AA. Investigating 3D Atomic Environments for Enhanced
740   QSAR. 2020.

741   55.   Goeman JJ. L1 penalized estimation in the Cox proportional hazards model. Biometrical J.
742   2010;52: 70–84. doi:10.1002/bimj.200900028

743   56.   Theorell A, Leweke S, Wiechert W, Nöh K. To be certain about the uncertainty: Bayesian
744   statistics for 13C metabolic flux analysis. Biotechnol Bioeng. 2017;114: 2668–2684.
745   doi:10.1002/bit.26379

746   57.   Murtagh F. Multilayer perceptrons for classification and regression. Neurocomputing. 1991;2:
747   183–197. doi:10.1016/0925-2312(91)90023-5
748