Histone H3 clipping is a novel signature of human neutrophil
extracellular traps

Authors

Dorothea O Tilley[1], Ulrike Abuabed[2], Ursula Zimny Arndt[3], Monika Schmid[3], Stefan Florian[4], Peter
R. Jungblut[3], Volker Brinkmann[2], Alf Herzig*[1], Arturo Zychlinsky*[1]

Author affiliations

[1]Department of Cellular Microbiology, Core facilities for [2]Microscopy and [3]Protein Analysis, Max
Planck Institute for Infection Biology, Charitéplatz 1, 10117 Berlin.
[4]Institut für Pathologie Charité - Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin.
*shared last authorship

Abstract

Neutrophils are critical to host defence, executing diverse strategies to perform their antimicrobial
and regulatory functions. One tactic is the production of neutrophil extracellular traps (NETs). In
response to certain stimuli neutrophils decondense their lobulated nucleus and release chromatin
into the extracellular space through a process called NETosis. However, NETosis, and the
subsequent degradation of NETs, can become dysregulated. NETs are proposed to play a role in
infectious as well as many non-infection related diseases including cancer, thrombosis,
autoimmunity and neurological disease. Consequently, there is a need to develop specific tools for
the study of these structures in disease contexts. In this study, we identified a NET-specific histone
H3 cleavage event and harnessed this to develop a cleavage site-specific antibody for the detection
of human NETs. By microscopy, this antibody distinguishes NETs from chromatin in purified and

mixed cell samples. It also detects NETs in tissue sections. We propose this antibody as a new

tool to detect and quantify NETs.

Introduction

Neutrophil extracellular traps (NETs) are extracellular structures consisting of chromatin

components, including DNA and histones, and neutrophil proteins (Brinkmann et al., 2004, Urban

et al., 2009). NETs were first described as an antimicrobial response to infection, facilitating

trapping and killing of microbes (Brinkmann et al., 2004). They are found in diverse human tissues

and secretions where inflammation is evident (recently reviewed by Sollberger et al., 2018). NETs

are produced in response to a wide-range of stimuli; bacteria (Brinkmann et al., 2004); fungi (Urban

et al., 2006); viruses (Saitoh et al., 2012; Schonrich et al., 2015); crystals (Schauer et al., 2014);

and mitogens (Amulic et al., 2017). Both NADPH oxidase (NOX) dependent and NOX independent

mechanisms lead to NET formation (Bianchi et al., 2009; Hakkim et al., 2011; Kenny et al., 2017;

Neeli and Radic, 2013). NETs are also observed in sterile disease, including multiple types of

thrombotic disease (recently reviewed by Jimenez-Alcazar et al., 2017) and even neurological

disease (Zenaro et al., 2015). NETs, or their components, are implicated in the development and

exacerbation of autoimmune diseases including psoriasis, vasculitis and systemic lupus

erythematosus (recently reviewed by Papayannopoulos, 2018) as well as cancer and cancer

metastasis (Albrengues et al., 2018; Cools-Lartigue et al., 2013; Demers et al., 2016).

Consequently, there is an urgency across multiple fields to establish the pathological contribution

of NETs to disease. However, the detection of NETs in affected tissues remains a challenge.

NETs are histologically defined as areas of decondensed DNA and histones that colocalise with

neutrophil granular or cytoplasmic proteins. Thus, reliable detection of NETs requires a combination

of anti-neutrophil and anti-chromatin antibodies as well as DNA stains. Immunofluorescent

microscopy is a useful method to detect NETs in tissue sections and *in vitro* experiments. However,

this can be challenging since NET components are distributed across the large decondensed

structure resulting in a weak signal. For example, the signal of antibodies to neutrophil elastase

(NE) is significantly dimmer in NETs than in the granules of resting cells where this protease is

highly concentrated. Conversely, anti-histone antibodies stain NETs strongly but not nuclei of naïve

neutrophils, where the chromatin is compact and less accessible. This differential histone staining

property can be exploited for the detection and quantification of NETs (Brinkmann et al., 2012).

However, sample preparation and the subsequent image analysis make results between different

labs difficult to compare. Thus, there is a need to identify antibodies against NET antigens.

NETs can also be detected through post-translational modifications (PTMs) that occur during

NETosis. Histone 3 (H3) is deaminated in arginine residues - the conversion to citrulline

(citrullination) - by protein arginine deiminase 4 (PAD4) (Wang et al., 2009). Citrullinated H3 (H3cit)

is widely used as a surrogate marker of NETs in both *in vitro* and *in vivo* experiments (Gavillet et

al., 2015; Pertiwi et al., 2018; Wang et al., 2009; Yoo et al., 2014; Yoshida et al., 2013). Cleavage

of histones by granular derived neutrophil serine proteases (NSPs) also contributes to NETosis

(Papayannopoulos et al., 2010). Histone cleavage, or clipping, by cysteine or serine proteases is a

bona fide histone PTM that facilitates the gross removal of multiple, subtler, PTMs in the histone

tail and is conserved from yeast to mammals (Dhaenens et al., 2015). Until now, histone clipping

has not been exploited for the detection of NETs but recent work by our group showed that histone

H3 cleavage is a conserved response to diverse NET stimuli, including *Candida albicans* and Group

B Streptococcus (Kenny 2017). Thus, in this study we map the site(s) of histone H3 cleavage during

NETosis. We developed a new monoclonal antibody against cleaved H3 that detects human NETs

in vitro and in histological samples. This antibody also facilitates easier NET quantification.

Results

Serine protease dependent cleavage of Histone H3 N-terminal tails during NET formation

Histones are processed in response to phorbol 12 myristate 13 acetate - PMA (Papayannopoulos et al., 2010; Urban et al., 2009) and other NET stimuli (Kenny et al., 2017). Indeed, the cleavage products of H3 were consistent between stimuli (Kenny et al., 2017). This suggests that H3 proteolysis occurs at specific sites during NETosis. In a time course experiment of human neutrophils incubated with PMA, we detected a H3 cleavage product of ~14 kDa as early as 30 min post-stimulation (Fig.1A). Further cleavages occurred between 60 and 90 min, yielding products of approximately 13 kDa and 10 kDa, respectively. The histone N-terminal tails protrude from the nucleosome core and are a major PTM target (Bannister and Kouzarides, 2011). A C-terminal, but not an N-terminal, histone antibody detected the cleavage products of H3 (Fig. 1A). These results indicate that the N-terminus is cleaved in truncated H3.

Neutrophil azurophilic granules are rich in serine proteases that can degrade histones *in vitro* (Papayannopoulos et al., 2010). We tested the contribution of these proteases to H3 cleavage during NET formation. Preincubation with the serine protease inhibitor, AEBSF (4- [2-aminoethyl] benzensulfonylfluoride) (Fig. 1A), but not with the cysteine protease inhibitor E64 (S.Fig 1A), inhibited H3 cleavage upon PMA stimulation. AEBSF also inhibited NET formation and nuclear decondensation as shown by immunofluorescent microscopy (Fig. 1B). PMA induced NETosis requires NADPH oxidase activity and, at high concentrations, AEBSF can inhibit activation of NAPDH oxidase (Diatchuk et al., 1997). To rule out this upstream effect, we showed that AEBSF did not inhibit ROS production at the concentrations used in our assay (S.Fig 1B [i]). Similarly, we verified that at these concentrations AEBSF was not cytotoxic, as shown by limited LDH release (S.Fig 1B [ii]). This data shows that serine proteases cleave the N-terminus of H3 early during NET formation.

A

B

Fig 1. PMA induced Histone H3 cleavage occurs in the N-terminal domain & is prevented by serine protease inhibition.

(**A**) Neutrophils were preincubated with the serine protease inhibitor AEBSF for 30 min in microcentrifuge tubes and then stimulated with PMA as indicated in the figure. Lysates were resolved by SDS-PAGE and immunoblotted with N- and C-terminal antibodies to H3. GAPDH was used as a loading control. (**B**) Immunofluorescent confocal microscopy of NET formation. Neutrophils were seeded on coverslips and preincubated with AEBSF before stimulation with PMA (50 nM) as indicated in the figure. At 150 min the cells were fixed and stained for neutrophil elastase (NE), chromatin (using a H2A-H2B-DNA antibody PL2.3) and DNA (DAPI [4',6-diamidino-2-phenylindole]). NS: non-stimulated. Images were taken at 63X and the scale bar is 20 µm. Blots and images are representative of 3 independent experiments.

Histone H3 is cleaved at a novel site in the globular domain

To identify the precise H3 cleavage sites we prepared histone enriched extracts from primary

neutrophils stimulated with PMA for 90 min and then purified H3 by RP-HPLC as previously

described (Shechter et al., 2007). A schematic summary of this is presented in S.Fig. 2A. We

identified the fractions containing H3 and its cleaved products by Western blot with anti-H3 C-

terminal antibodies (Fig.2A). As expected, H3 was the last core histone to elute (at 45-46 min).

We further separated H3 and its truncated forms by two-dimensional electrophoresis (2-DE) and

confirmed their identity by mass spectrometry (Fig. 2B and S.Fig. 2B i-ii). The sequence coverage

did not include residues that allowed the differentiation of H3 variants. The N-terminals of the

separated H3 fragments were not covered by MS and therefore sequenced by Edman degradation

from 2 independent experiments (Fig. 2C). The N-terminal sequence of the largest molecular

weight H3 spot matched that of intact H3 (Fig 2B, spot 1). We did not obtain reliable sequencing of

spots 2 and 3 but the cleavage sites of spots 4 and 5 were identified. The most truncated H3

fragment (spot 5) was cleaved between L48 and R49, in the globular domain of the protein, within

the nucleosome core structure (Fig. 2D). This is a previously unidentified cleavage site in H3 and,

thus, we selected cleavage at H3R49 as a candidate marker of NETs.

A

C

Table 1: N-terminal histone H3 cleavage sites identified by Edman degradation sequencing

Spot #	Name	Round 1	Round 2
1	Intact H3	ARTKQ	ARTKQ
2	CS1	*NA*	T
3	CS2	AARKS	AAAAS TKRRR
4	CS3	TGGV	TGGVK* AAA
5	CS4	REIRR	REIRR* AL - EI

*confirmation of first round identification
NA: not analysable

Fig 2. Identification of histone H3 cleavage sites in NET formation.
(**A**) Representative RP-HPLC chromatogram of acid extracted histones from NETs and corresponding 1D-SDS-PAGE and immunoblots to identify H3 and H4 containing fractions. Histone enriched supernatants were prepared from neutrophils stimulated with PMA for 90 min. (**B**) Representative Coomassie stained blot of pooled H3 fractions separated by 2-DE. Inset is a zoom of all spots (1-5) identified as histone H3 by mass spectrometry. Other proteins identified are listed in the supplementary figures. (**C**) Summary of Edman degradation sequencing results of the H3 spots in two independent experiments. In the second experiment overlapping sequences were detected. However, the detected amino acids for spot 4 and 5 confirmed the initial sequence identification from round 1*. (**D**) Schematic representation of the cleavage site of the truncated H3 product in both the linear sequence of H3 and in the nucleosomal context. H3 is represented in blue and the pink tail region and partial alpha helix represent the part of H3 that is removed. The nucleosome structure is adapted from PDB 2F8N (Chakravarthy, 2005).

Generation of a histone H3 cleavage site monoclonal antibody

We adopted a similar strategy to Duncan and colleagues (2008) to raise antibodies against the cleaved site. We designed a lysine branched immunogen containing the 5 amino acids at the carboxylic side of the H3R49 cleavage site (outlined in S.Fig 3A) and used it to immunize mice. After preliminary screening by ELISA against the immunogen and control peptides, we selected sera, and later hybridoma clones, that detected cleaved H3 in immunoblots of PMA stimulated cell

lysates. We excluded sera and clones that detected full length histone H3 in addition to cleaved H3

(Fig 3A). We selected sera and clones that detected NETs but not resting chromatin of naive

neutrophils by immunofluorescence microscopy. Of the 6 mice immunised, we obtained one stable

clone, 3D9, that functioned in both Western blot and microscopy – other clones performed only in

Western blot (data not shown). 3D9 recognised a protein of ~10 kDa in neutrophils stimulated with

PMA for 120 min and longer but did not detect any protein in resting or early stimulated cells (Fig

3A). This band corresponded in size with the smallest H3 fragment detected by the H3 C-terminal

antibody. Interestingly, by microscopy, 3D9 exclusively recognised neutrophils undergoing

NETosis – with decondensing chromatin (Fig 3B). To validate the specificity of the antibody for the

de novo N-terminal H3 epitope of NETs, we performed competition experiments with the

immunising peptide and demonstrated that it could block 3D9 binding to NETs as shown by

immunofluorescent microscopy (S.Fig 3B).

3D9 binds specifically to cleaved H3. This antibody binds to the immunizing peptide and to isolated

NETs, but not to equal concentrations of chromatin, recombinant H3 or purified calf thymus DNA,

by direct ELISA (Fig 3C). Furthermore, in immunoprecipitation experiments, 3D9, but not an isotype

control, pulled down intact H3 in lysates of naïve and activated neutrophils, but only the cleaved

fragment from activated cells (S.Fig 3C). We detected these pull downs both by Coomassie and

silver stained gels. In a similar experiment, we immunoblotted the immunoprecipitate with the C-

terminal antibody as well as 3D9 (S.Fig 3C [ii]), and showed that only our monoclonal antibody

recognizes cleaved H3. Together this data shows that 3D9 is selective for cleaved H3 under the

denaturing conditions of SDS-PAGE but also recognizes full length H3 under the more native

conditions of IP.

A

H3 C-terminal   3D9

B

C

Fig 3. Screening and detection of cleaved H3 & NETs by 3D9
(**A**) Immunoblots of lysates prepared from neutrophils stimulated with PMA (50 nM) for the times indicated in
the figure. H3 C-terminal antibody was used as a control to detect all H3 forms while a single band (cleaved
H3) was detected by the newly generated monoclonal antibody, 3D9. (**B**) Immunofluorescent microscopy of
neutrophils stimulated with PMA and fixed at the indicated times. Samples were stained with Hoechst (DNA
- blue) and 3D9 (with Alexafluor-568 conjugated secondary antibody - red). NS: non-stimulated. Images were

taken on an upright fluorescent microscope at 20X. Scale bars – 50 µm (full field) and 10 µm (inset) (**C**) Direct
ELISA for cleaved H3 in NETs, chromatin (A549 lung epithelial cells), recombinant histone H3 and DNA.
Samples were serially diluted and immobilized on a high affinity ELISA plate according to DNA content (for
NETs, chromatin and DNA) or protein content (for recombinant histone H3) as determined by PicoGreen and
bicinchoninic acid assays respectively. Starting concentration was 1 µg/ml DNA or protein. Cleaved H3 was
detected using 3D9 (2 µg/ml) and HRP conjugated anti-mouse secondary antibody and reactions were
developed using TMB (3,3',5,5'-tetramethylbenzidine) as a substrate. Data is presented for dilution
200 ng/ml. REIRR peptide control was coated at 20 ng/ml. Data represents mean ± SD of 3 experiments
using independent NET donors.

Epitope mapping & sequencing of antibody clone 3D9

To determine the binding site of 3D9 in histone H3, we tested the antibody by ELISA with

overlapping linear peptide arrays and helical peptide mimic arrays (S.file 1) based on a sequence

(residues 30-70; PATGGVKKPHRYRPGTVALREIRRYQKSTELLIRKLPFQRL) around the H3R49

cleavage site (S.Fig 4A & 4B). In such assays acetylation is often used to neutralise the contribution

of the amino terminal charge. However, to mimic any potential charge created at the newly revealed

N-terminus, R49, we also included arrays of unmodified peptides. Based on overlapping peptides,

the putative core epitope in the linear array was (R)EIRR. The peptides ending in REIRR were in

all cases in the top 2 of each peptide mimic (S.Table 3). Interestingly, peptides extended at the N-

terminus were still recognized. Moreover, acetylation at the N-terminus of the peptide ending in the

REIRR sequence did not affect the binding, suggesting that a free N-terminus may not be

recognized by the antibody. We further refined the epitope mapping by amino acid replacement

analysis of linear peptides and helical peptide mimetics ending in REIRR (S.Fig 4C [i, ii]). Mutations

in Glu51, Ile52, and Arg54 negatively impacted the signal, indicating these residues are critical for

epitope recognition. A schematic of the antibody epitope mapped onto H3 is presented in S.Fig 4C

(iii).

To preserve the primary hybridoma sequence integrity and to confirm monoclonality, we sequenced

the full antibody gene sequence. The heavy and light chain constant and variable gene sequences

are presented in the supplemental results (S.Table 4).

Automatic quantification of in vitro generated NETs by microscopy

We tested how 3D9 stained NETs in immunofluorescence in samples that were robustly

permeabilized (Triton X-100, 0.5% for 10 min) to facilitate the distribution of the antibody throughout

the sample. Fig 4A shows that 3D9 detects decondensed chromatin almost exclusively. In contrast,

the anti-chromatin antibody (PL2.3) - directed against a H2A-H2B-DNA epitope (Losman et al.,

1992) – stains NETs in addition to condensed nuclei. We compared the staining characteristics of

3D9 and PL2.3 during NET formation. We determined the nuclear area and signal intensity at the

indicated time points, from multiple fields of view (Fig. 4B). Both antibodies detect the increase in

nuclear area characteristic of NETosis between 15 and 180 min after simulation. At later time

points, the intensity of PL2.3 staining decreased and failed to discriminate between resting cell

nuclei and NETs. In contrast, 3D9 stained nuclei undergoing NETosis more brightly than nuclei of

non-activated cells.

Publicly available software (ImageJ) can be used to quantify *in vitro* NETosis. We compared 3D9

versus the anti-chromatin antibody with our previously published semi-automatic image analysis

(Brinkmann et al., 2012) and with a modified automatic method (S.Fig 5A). Both methods use

automatic thresholding of the DNA channel (Hoechst) to count total cells/objects. The historical

semi-automatic method exploits the differential staining by chromatin antibodies of decondensed

chromatin (high signal) over compact chromatin (weak signal) to count cells in NETosis. This

method uses a manual thresholding and segmentation procedure (denoted manual in the figure).

This manual thresholding step is subject to observer bias. In contrast, the modified method uses

automatic thresholding at both stages; total cell and NET counts. Both methods use a size

exclusion particle analysis step so that only structures larger than a resting nucleus are counted.

Both 3D9 and PL2.3 antibodies effectively quantified NETs using previously published method (Fig

4C - manual). However, PL2.3 failed to accurately quantify the number of NETs with the fully

automatic method, specifically at early time points after stimulation (15 min). In this case, the weakly

staining lobulated nucleus can extend over a larger surface area during cell activation and adhesion

resulting in these cells being wrongly categorised as NETs by the algorithm. Together, these data

indicate that the automatic method using 3D9 staining can reduce experimental bias.

Fig 4. Comparison of NET quantification using an anti-chromatin antibody versus 3D9
(**A**) Confocal immunofluorescent microscopy of neutrophils stimulated with PMA (180 min) and stained with
Hoechst and anti-chromatin antibody (PL2.3) or 3D9. Insets represent selected cells examined at higher
magnification (63X) and presented as split channels in grayscale or merged as per the total field of view (20X
magnification). White scale bar - 50 μm, cyan scale bar - 10 μm. Images are representative of 3 experiments

(**B**) Comparison of the fluorescent distribution of PL2.3 versus 3D9 staining of PMA stimulated cells over time
(6h). Staining intensities were normalized over all images of the respective time course. NS:360: non-
stimulated at 360 min. Analysis is performed on one data set that is representative of 3-4 independent time
course experiments. (**C**) Comparison of NET quantification using manual or automatic thresholding and
segmentation procedures for chromatin antibody (PL2.3) and cleaved H3 antibody (or 3D9). Manual
thresholding excludes cells/NETs with a weak signal whereas automatic thresholding includes all objects
irrespective of signal. Images for analysis were taken using a fluorescence microscope. Graph represents
the mean ± standard deviation, where n=3-4.
273
274

3D9 detects NETs induced by multiple stimuli

Histone H3 cleavage is a feature of the neutrophil response to multiple NET stimuli (Kenny et al.,

2017). 3D9 detects NETs induced by the bacterial toxin nigericin, which induces NETs

independently of NADPH oxidase activation (Kenny et al., 2017), by heme in TNF primed

neutrophils (Knackstedt et al., 2019) and by the fungal pathogen *Candida albicans* (Fig 5).

Interestingly, in *C. albicans* infections, we detected both 3D9 positive and negative NETs.

Fig 5. Detection by 3D9 of NETs from diverse stimuli.
Immunofluorescence microscopy of neutrophils left unstimulated (NS), stimulated with PMA (50 nM, 2.5h), nigericin (15 µM, 2.5 h), TNF primed and then stimulated with heme (20 µM, 6h), and neutrophils co-cultured with *Candida albicans* hyphae (MOI 5) for 4h. Samples were stained with Hoechst, anti-neutrophil elastase (NE) and 3D9. Scale bar – 50 µm. Images were taken on a confocal microscope at 20X and are representative of 3 experiments with independent donors. Scale bar - 20 µm.

3D9 distinguishes NETs in mixed cell samples

Histone H3 clipping, albeit at other sites in the N-terminal tail, was observed in mast cells (Melo et al., 2014) and unstimulated PBMC fractions (Howe and Gamble, 2015). Of note, PBMC fractions often contain contaminating neutrophils (Hacbarth and Kajdacsy-Balla, 1986). To test if 3D9 specifically stained neutrophils treated with NET stimuli, we incubated PBMCs with PMA or nigericin (Fig. 6). Importantly, 3D9 detected only nuclei that appeared decondensed in cells that were positive for NE, a specific neutrophil marker. In contrast, the chromatin antibody stained both

neutrophils in NETosis and nuclei of other cells. This shows that 3D9 detects NETs specifically

even in the presence of other blood cells.

3D9 distinguishes NETosis from other forms of cell death in neutrophils

Neutrophils can commit to other cell death pathways (recently reviewed by Dabrowska et al., 2019) besides NETosis. Naïve neutrophils undergo apoptosis after overnight incubation (Kobayashi et al., 2005) and necroptosis upon TNFα stimulation in the presence of a SMAC (second mitochondria-derived activator of caspase) mimetic and if caspases are inhibited (Galluzzi et al.,

311 2012). Interestingly, the anti-chromatin antibody (S.Fig7), but not 3D9, stained the condensed

312 nuclei of cells undergoing apoptosis (Fig. 7) and neither of the antibodies stained cells during

313 necrosis induced by the staphylococcal toxin α-haemolysin nor after stimulation with necroptosis

314 inducers.

Fig 7. Comparison of 3D9 detection in response to apoptotic, necroptotic & necrotic cell death stimuli.
Confocal immunofluorescent microscopy of neutrophils stimulated with different cell death stimuli and
subsequently stained with Hoechst, anti-neutrophil elastase (NE) and 3D9. NETs were induced with PMA
(100 nM, 3h). Apoptosis was induced in resting neutrophils by incubation for 24h without stimulation
overnight. Neutrophils were stimulated with Z-VAD-FMK (50 µM) plus SMAC mimetic (100 nM) plus TNF (50
ng/ml) for 6h to induce necroptosis. Necrosis was induced with the pore forming toxin α-haemolysin (25
µg/ml). Images were taken at 20X and are representative of 3 experiments. Scalebar 20 µm. A comparison
was made with parallel samples stained with the chromatin antibody PL2.3 and are presented in S.Fig 7.

3D9 labels NETs in human tissue sections

NETs are found in inflamed tissues based on the juxtaposition of chromatin and granular makers as well as the detection of citrullinated H3. 3D9 stains areas of decondensed DNA (Hoechst) that colocalise with NE in both inflamed human tonsil (Fig. 8A) and human kidney (Fig. 8B). This indicates that 3D9 labels NETs in histological samples - hematoxylin and eosin (HE) tissue overviews are provided in the supplemental figures (S.Fig 8-10). Indeed, in kidney (Fig. 8C), inflamed appendix (Fig. 9) and gallbladder (Fig. 10), 3D9 labelled DNA in the same cluster as anti-H3cit or anti-H2B. Interestingly, 3D9 stained decondensed, more NET-like structures, while anti-H3cit or anti-H2B antibodies stained relatively compact chromatin. Furthermore, colocalization analysis of 3D9 with H2B or with H3cit revealed that 3D9 was more commonly colocalised with H2B as compared to H3cit; overlap coefficients 0.463 (3D9-H2B) v 0.125 (3D9-H3cit), and 0.533 (3D9-H2B) v 0.122 (3D9-H3cit) for S.Fig 9 and 10 respectively.

Fig 8. Detection of clipped histone 3 & NETs in human tissues
Paraffin embedded sections were stained with Hoechst, anti-NE and 3D9 or H3cit antibodies and examined
by confocal microscopy. Scale bar - 10 μm (**A**) inflamed human tonsil (**B** & **C**) Inflamed human kidney.

Fig. 9. Comparison of Clipped H3, H3cit & H2B staining in the gallbladder
Paraffin embedded sections were stained with Hoechst, anti-NE and histone antibodies 3D9, H3cit and H2B and examined by confocal microscopy. Scale bar – 20 μm.

Fig. 10. Comparison of Clipped H3, H3cit & H2B staining in the appendix
Paraffin embedded sections were stained with Hoechst, anti-NE and 3D9 or H3cit antibodies and examined
by confocal microscopy. Scale bar - 20 μm.

Discussion

Decondensed chromatin is a defining feature of NETs. It occurs through PTMs that partially

neutralise the histone positive charge and thus the affinity of histones for negatively charged DNA

(Papayannopoulos et al., 2010; Wang et al., 2009). One way to achieve this is through proteolytic

removal of the lysine and arginine rich histone tails. Using a biochemical and proteomic approach,

we determined that H3 is cleaved within its globular domain during NETosis. We exploited the

specificity of this event to produce a mouse monoclonal antibody to the *de novo* histone H3 epitope,

the new N-terminal beginning at R49. This antibody, 3D9, recognises human NETs induced by both

microbial and host derived physiological stimuli and distinguishes netotic neutrophils from

neutrophils that die via alternative pathways. It also discriminates between NETs and other cells in

mixed blood cell fractions and, importantly, NETs in human histological samples.

Until now, histone citrullination is the only PTM that has been used for antibody-based detection of

NETs. In this paper, we propose histone cleavage at H3R49 as a new histone PTM that can be

used for broad detection of NETs from human samples. The use of H3cit for the detection of NETs

is not without controversy. Not all NETs are citrullinated and NET formation can occur in the

absence or inhibition of citrullinating enzymes (Kenny et al., 2017; Konig and Andrade, 2016). By

identifying the precise histone cleavage site, we shed further light on this. The most commonly

used H3cit antibody detects citrullination of R2, R8 and R17. However, histone cleavage at R49

would remove the H3cit epitopes, rendering these NET defining PTMs mutually exclusive on a

single histone level. Indeed, co-staining by anti-H3cit and 3D9 in inflamed kidney, gallbladder and

appendix paraffin sections revealed extensive mutual exclusion of the two marks and more

abundant staining of decondensed chromatin by 3D9. Thus, we propose that 3D9 will allow broad

detection of NETs but may display a preference for more mature or proteolytically processed NETs.

In contrast to the present study, histone cleavage was reported as discriminating between different

pathways of NET formation (Pieterse et al., 2018). Using a sandwich ELISA approach, Pieterse *et*

al concluded that, generally, the N-terminal histone tails are removed in NOX dependent but not

NOX independent NET formation. They used a suite of N-terminal directed antibodies for H2B, H3

and H4, and all H3 epitopes were located N-terminal to the cleavage site H3R49. However, in the

final biological sample testing the authors did not examine H3. Interestingly, the authors observed

that, by immunofluorescent microscopy, all histone N-terminal antibodies failed to stain NETs at

time points after cell lysis. Therefore, to us, this data suggests that at later stages in NETosis the

histone N-terminal tails, at least for H3, are removed irrespective of the pathway of activation. This

is in line with our observations that both NOX dependent (PMA, heme) and NOX independent

(nigericin) stimuli result in NETs that are recognised by 3D9 and supports our finding that histone

H3 cleavage at R49 is a general feature of human NET formation.

In this study we detect NETs in fixed or denatured human samples from *in vitro* experiments and

histological samples. While an ELISA with 3D9 revealed a preference for NETs over isolated

chromatin or recombinant H3, immunoprecipitation of naïve cell lysates by 3D9 also pulled down

full length H3 as confirmed by immunoblot. It is not yet clear if this is due to co-immunoprecipitation

due to the presence of low levels of clipped histone or if this represents true recognition of intact

H3 by 3D9. Thus, care should be taken when detecting cleaved H3 or NETs under native and mild

detergent conditions and all sample types e.g. serum samples, need careful validation for cross

reactivity.

More broadly, and applying to the general principles of NET detection, it is not yet possible to prove

conclusively that the detected decondensed chromatin originates from the same cell source as the

neutrophil proteins which decorate it. For example, in an infected necrotic wound to which to high

numbers of neutrophils are recruited. Here, activated neutrophils might release both proteases

(Borregaard et al., 1993) and citrullinating enzymes (Spengler et al., 2015; Zhou et al., 2017) that

bind to and modify extracellular chromatin generating a NET - according to the histological

definition. This remains a conundrum that requires further exploration.

N-terminal histone cleavage at H3R49 is a novel and so far undescribed cleavage site in any

eukaryotic organism. Unusually, it is located in the globular rather than the unstructured tail region

of H3. H3R49 is one of 6 key residues important for the regulation of H3K36me^3 and forms part of

the structured nucleosome surface (Endo et al., 2012). Thus, we speculate that removal of the N-

terminal tail, in its entirety, could lead to removal of higher order structure interactions and facilitate

chromatin decondensation e.g. removal of H3K9me and its associated heterochromatin protein 1

interactions (Jacobs and Khorasanizadeh, 2002). To determine the contribution of histone cleavage

at H3R49 to the process of chromatin decondensation, future work will focus on establishing the

protease(s) responsible and the sequence of proteolytic events leading to this final truncation of

H3 and NET formation. Given the specificity of this event and its restriction to NETotic forms of cell

death, we propose that N-terminal cleavage at H3R49 is an example of histone 'clipping' in

neutrophils – a term proposed by the histone/histone proteolysis field for specific histone cleavage

sites for which a biological function has been demonstrated (Dhaenens et al., 2015).

In conclusion, this study represents the first identification of a distinctive and exclusive marker of

NETs and describes the development and characterisation of a complementary antibody to

facilitate easier detection of human NETs. Analogous to finding a smoking gun at a crime scene,

the monoclonal antibody 3D9 detects evidence of the proteolytic events that occur in NETosis –

the proteolytic signature, histone cleavage at H3R49. In doing so, 3D9 discriminates NETs from

chromatin of other cells and chromatin of neutrophils that die via alternative mechanisms. This

added layer of specificity will simplify the detection of NETs in tissue samples and facilitate

comparison of quantitative studies between labs. This will be an important step in assessing the

contribution of extracellular chromatin and NETs to disease pathology.

428 Materials and methods

429 *Reagents*

430 All reagents were purchased from common vendors of laboratory reagents e.g. Sigma Aldrich or

431 VWR Deutschland unless otherwise stated.

Blood collection and ethical approval

Venous blood was collected from healthy donors who had provided informed consent according to the Declaration of Helsinki. Ethical approval was provided by the ethics committee of Charité-Universitätsmedizin Berlin and blood was donated anonymously at Charité Hospital Berlin.

Purification and culture of human peripheral blood neutrophils

Neutrophils were isolated as described by Amulic et al (2017). Briefly, venous whole blood was collected in EDTA and separated by layering over equal volume Histopaque 1119 and centrifugation at 800 g (20 min). The pinkish neutrophil rich fraction was collected and washed once by the addition of 3 volumes of wash buffer (PBS, without Mg^{2+} or Ca^{2+} [Gibco] supplemented with 0.5% [w/v] human serum albumin [HSA, Grifols]) and centrifugation at 300 g (10 min). The neutrophil fraction was further purified by density gradient centrifugation - Percoll (Pharmacia) gradient from 85%-65% (v/v). Purified cells were collected from the 80-70% fractions and washed once before being resuspended in wash buffer. Cells were counted using a CASY cell counter.

For all experiments, unless indicated, neutrophils were cultured RPMI (GIBCO 32404014) supplemented with 10 mM HEPES and 0.1% (w/v) HSA, which had been preequilibrated in CO_2 conditions for 1 h. For some stimuli, the HSA content was reduced to 0.05% or 0% HSA as indicated in the figure legends. Cells were routinely cultured at 37°C, 5% CO_2 unless indicated. For all experiments, stimuli were added to cell reactions as 10X working stock solutions freshly diluted in RPMI. For inhibition experiments, a 10X inhibitor stock and appropriate vehicle controls, were added to the cells and preincubated for the times stated in the figure legends.

Neutrophil and NET lysate preparation

To analyse proteins, lysates were prepared from stimulated or resting neutrophils. Cells were seeded in culture medium in 1.5 ml microcentrifuge tubes at 1×10^7 cells/ml with 5×10^6 cells per time point. After addition of the inhibitor or agonist, cells were gently mixed and incubated at 37 °C with gentle rotation. At the specified time points, protease inhibitors - 1 mM AEBSF, 20 µM Cathepsin

G inhibitor I (Calbiochem), 20 µM neutrophil elastase inhibitor GW311616A (Biomol), 2X Halt

protease inhibitor cocktail (PIC, Thermofisher Scientific), 10 mM EDTA, 2 mM EGTA - were added

directly to the cell suspension. Cells were gently mixed and centrifuged at 1000 g (30 s) to collect

all residual liquid. Boiling 5X sample loading buffer (50 mM Tris-HCl pH 6.8, 2% [w/v] SDS, 10%

glycerol, 0.1% [w/v] bromophenol blue, 100 mM DTT) was added to samples which were then

briefly vortexed and boiled (98 °C) for 10 min and flash frozen in liquid nitrogen for storage at -

80 °C.

464 *1D SDS-PAGE and immunoblot blot*

For routine protein analysis, samples were analysed by 1D SDS-PAGE and immunoblotted.

Samples were thawed on ice, boiled at 98 °C (10 min) and sonicated to reduce viscosity (Braun

sonicator, 10 s, cycle 7, power 70%). Proteins were applied to NuPAGE 12% gels (Invitrogen,

Thermofisher) and run at 150 V in MES buffer (Thermofisher Scientific). Proteins were transferred

by western blot onto PVDF (0.2 µm pore size, Amersham GE Healthcare) using the BioRad wet

transfer system (buffer: 25 mM Tris, 192 mM glycine, 20% methanol, protocol: 30 min at 100 mA,

120 min at 400 mA). Blotting efficacy was assessed by Ponceau S staining. Blots were blocked

with TBST (TBS pH 7.5, 0.1% [v/v] Tween-20) with 5% [w/v] skimmed milk, for 1 h at RT. Blots

were then incubated with the following primary antibodies overnight at 4 °C or for 2 h at RT: rabbit

anti-histone H3 C-terminal pAb, 1:15000 (Active motif #61277); rat anti-histone H3 N-terminal mAb,

1:1000 (Active Motif #61647, aa 1-19); Histone H4 C-terminal, 1:5000 (Abcam 10158 – aa 50 to C

terminal); rabbit anti-histone H4 N-terminal mAb, 1:30,000 (Upstate, Millipore #05-858, aa17-28);

rabbit GAPDH mAb, 1:5000 (Cell Signalling Technology, #2118); mouse 3D9 1ug/ml (produced in

this study) – all diluted in TBST with 3% (w/v) skimmed milk. After washing with TBST (3 x 5 min),

blots were blocked for 15 min as before and then probed with secondary HRP conjugated

antibodies (Jackson ImmunoResearch -diluted 1:20000 in 5% skimmed milk TBST) for 1 h at RT.

Blots were washed in TBST (3 x 5 min) and developed using SuperSignal™ West Dura Extended

Duration Substrate (ThermoFisher Scientific) and an ImageQuant Gel imager (GE Healthcare).

Immunofluorescent staining of in vitro samples

For immunofluorescent imaging of purified cells, neutrophils/PBMCs were seeded in 24 well dishes containing glass coverslips, with 1×10^5 cells per well and incubated at 37 °C for 1 h to allow to adhere to the coverslip. At this stage, inhibitors and priming factors were included as indicated. Reactions were stopped by the addition of paraformaldehyde (2% [w/v]) for 20 min at RT or 4°C overnight. After fixation, cells were washed and stained as previously described (Brinkmann et al 2012). Briefly, all steps were performed by floating inverted coverslips on drops of buffer on laboratory parafilm. Cells were permeabilised with PBS, pH 7.5, 0.5% (v/v) Triton X-100 for 3 min. For screening and quantification experiments, this permeabilization step was extended to 10 min. Samples were washed (3 x 5 min) with PBS, 0.05% (v/v) Tween 20 and incubated with blocking buffer - PBS pH 7.5, 0.05% (v/v) Tween 20, 3% (v/v) normal goat serum, 3% (w/v) freshwater fish gelatin, 1% (w/v) BSA - for 20 min at RT and then probed with primary antibodies diluted in blocking buffer and incubated overnight at 4 °C. Primary antibodies: anti-chromatin (H2A-H2B-DNA complex) mouse mAb, 1 µg/ml (Losman 1992); neutrophil elastase, rabbit pAb 1:500 (Calbiochem); mouse serum for screening, 1:100; hybridoma supernatants, neat; 3D9 mouse mAb, 1 µg/ml. Samples were then washed as before. Alexa labelled secondary antibodies (Invitrogen) were diluted 1/500 in blocking buffer and incubated for 2 h at RT. DNA was stained with Hoescht 33342 (Invitrogen, Molecular Probes) 1 ug/ml, incubated with the secondary antibody step. Samples were washed in PBS followed by water and mounted in Mowiol mounting medium.

Histone extraction from neutrophils

Histone enriched fractions were prepared from resting neutrophils and NETs according to a method modified from Shechter et al (2007). Neutrophils (4-8 $\times 10^7$) were resuspended in 13 ml of RPMI (without HSA) in a 15 ml polypropylene tube and incubated on a roller at 37 °C with PMA 50 nM for 90 min. After stimulation, 1 mM AEBSF was added to inhibit further degradation by NSPs and cells were cooled on ice for 10 min. All subsequent steps were performed on ice or 4 °C where possible. Cells and NETs were pelleted by centrifugation at 1000 *g*, 10 min. Samples were

resuspended in ice-cold hypotonic lysis buffer (10 mM Tris-HCl pH 8.0, 1 mM KCL, 1.5 mM MgCl$_2$,

1 mM DTT supplemented with protease inhibitors just before use – 1 mM AEBSF, 20 µM NEi, 20

µM CGi, 2X PIC, 10 mM EDTA) using 1 ml of buffer/5x10^6 cells. Cells were then incubated at 4 °C

on a rotator for 30 min before being passed through a syringe to aid lysis and shearing of intact

cells. Nuclei and NETs were collected by centrifugation at 10,000 g, 10 min, discarding the

supernatant. To disrupt nuclei, samples were resuspended in dH$_2$O (1 ml/1x10^7 cells)

supplemented with protease inhibitors, as before, and incubated on ice for 5 min with intermittent

vortexing. NP40 (0.2% [v/v]) was added to help lysis and disruption of NETs and samples were

sonicated briefly (10 s, mode 7, power 70%). To extract histones, H$_2$SO$_4$ (0.4 N) was added to

samples, and vortexed briefly. Samples were then incubated, rotating, for 2-3h. Histone enriched

fractions were collected by aliquoting samples into multiple 1.5 ml microcentrifuge tubes and

centrifuging at 16,000 g for 10 min, followed by collection of the supernatants. To minimise further

processing of histones, proteins were immediately precipitated overnight by dropwise addition of

trichloroacetic acid to a final concentration of 33% followed by mixing. The next day precipitated

proteins were pelleted by centrifugation at 16,000 g, 10 min. The supernatants were discarded and

waxy pellets were washed once with equal volume ice-cold acetone with 0.2% (v/v) HCl and 5

times with ice-cold acetone alone. Pellets were allowed to air dry for 5 min before being

resuspended with 1 ml (per 5x10^6 cells) of dH$_2$O plus 1 mM AEBSF. For difficult to resuspend

pellets the mixture was vigorously shaken at 4°C overnight before samples were centrifuged, as

before, to remove undissolved protein. Pooled supernatants for each sample were lyophilised and

stored at -80°C until histone fractionation.

Purification of histone H3

Histones were fractionated by RP-HPLC according to the method described by Shechter et al

(2007). After lyophilisation samples were resuspended in 300 µl Buffer A (5% acetonitrile, 0.1%

trifluoroacetic acid) and centrifuged at 14,000 g to remove particulate matter. 150 µl of clarified

sample was mixed with 40 µl Buffer A before being applied to a C18 column (#218TP53, Grave

Vydac) and subjected to RP-HPLC (Waters 626 LC System, MA, US) as described by Schecter et

al (2007). The flow rate was set to 1 ml min^{-1} and fractions were collected at 30 s intervals from

minute 30 to 55. All fractions were lyophilised and stored at -80 °C until analysis. To determine

which fractions contained H3 and cleaved species, each fraction was dissolved in 50 µl dH$_2$O and

5 µl was subjected to SDS-PAGE and either stained with Coomassie blue stain or transferred to

PVDF and immunoblotted for H3 and H4 as described already.

2D electrophoresis (2-DE) of purified histones

To determine the cleavage sites, H3 containing fractions were pooled and subjected to a small gel

2-DE procedure (Jungblut and Seifert, 1990). Briefly, pooled fractions were denatured in 9 M urea,

70 mM DTT, 2% Servalyte 2-4 and 2 % Chaps. Samples (30 µl) were applied to 1.5 mm thick

isoelectric focusing (IEF) gels using ampholytes 7-9 and a shortened IEF protocol was used: 20

min 100 V, 20 min 200 V, 20 min 400 V, 15 min 600 V, 5 min 800 V, and 3 min 1000 V, (a total of

83 min and 500 Vh) in 8 cm long IEF tube gels. Separation in the second dimension was performed

in 6.5 cm x 8.5 cm x 1.5 mm SDS-PAGE gels. Duplicate gels were prepared; one stained with

Coomassie Brilliant Blue R250 for excision of spots for mass spectrometry identification; and the

second transferred to PVDF as follows. Proteins were blotted onto PVDF blotting membranes (0.2

µm) with a semidry blotting procedure (Jungblut et al., 1990) in a blotting buffer of 100 mM borate.

Spots were stained by Coomassie Brilliant Blue R250 and analysed by N-terminal Edman

degradation sequencing (Proteome Factory, Berlin, Germany).

Antibody generation

Immunising and screening peptides are outlined in S.Table 2 (S.Fig 3A) and were synthesized by

Eurogentec (Belgium). A portion was further conjugated to Key Lymphocyte Haemoglutinin (KLH)

for immunization. Immunisation of mice, preliminary ELISA screening and production of

hybridomas were performed by Genscript as follows. Six mice (3x Balb/c and 3x C57/BL6) were

immunized with branched peptides. Mice were bled and effective immunization was assessed using

a direct ELISA. The ELISA and subsequent inhouse immunoblot and immunofluorescent

microscopy screening strategy are outline in S.Fig 3A. Following selection of effectively immunized

animals, a further boost injection of the immunogen was given before isolation of spleen cells for

hybridoma production. The resulting hybridoma supernatants were screened similarly. Large-scale

culture of supernatants and purification of antibodies was performed by Genscript.

ELISA

To assess 3D9 specificity for NETs, 3D9 was used in an indirect ELISA to detect cleaved H3 in

purified NETs, chromatin, recombinant H3 and DNA. NETs were prepared by seeding 3×10^6

neutrophils in a 6 well dish and incubating for 3-6 h with 100 nM PMA. NETs were gently washed

3 times with equal volume PBS, before being collected in 300 µl PBS. Clumped NETs were

disrupted by sonicating briefly (3 s, mode 7, power 70% - Braun Sonicator) and then snap frozen

and stored at -80°C. Chromatin was prepared from lung epithelial cells (A549) as previously

described by Shechter *et al* (2007). The final nuclear pellet was resuspended in dH_2O and

sonicated as before and stored at -80 °C. The DNA content of NETs and chromatin was assessed

by PicoGreen assay according to the manufacturer's instructions (Thermofisher Scientific).

Beginning at 1 µg/ml (of DNA content), serial dilutions of NETs, chromatin and calf thymus DNA

(Invitrogen) were prepared in lo-DNA bind Eppendorf microcentrifuge tubes. A similar dilution series

of recombinant histone H3 (New England Biolabs) was prepared starting at 1 µg/ml protein. All

dilutions were performed in PBS. One hundred microliters of each sample, in duplicate, at dilutions

1 ug/ml to 1 ng/ml, was aliquoted in a Nunc Maxisorb 96 well dish and immobilised overnight at

4°C, 250 rpm. The immunising peptide, REIRR (10ng/ml) was used as a positive control. The

following day all wells were washed 6 times with wash buffer (PBS, 0.05% Tween 20) and then

blocked with 200 µl of blocking solution (1% BSA in wash buffer) for 2 h (RT). Wells were washed

once with wash buffer and incubated with 100 µl of 3D9 (2 µg/ml, in blocking solution) at RT (2 h)

with gentle shaking (250 rpm). Wells were washed 6 times as before and then incubated with 100

µl of secondary HRP conjugated anti-mouse (Jackson laboratories) at 1:10,0000 dilution in blocking

solution and incubated as before. Finally, wells were washed 6 times as before and HRP activity

was detected using TMB (3,3', 5,5' tetramethylbenzidine) reagent (BD OptEIA™) according to the

manufacturer's instructions (incubating for 15 -30 min). The assay was stopped by the addition of

100 µl H_2SO_4 (0.16 M) and absorbance (450 nm) was measured on a 96 well plate reader

(VERSAmax, Molecular Devices, CA, US).

Quantification of staining characteristics by Image J and R

In order to assess staining characteristics of antibodies during NET formation we developed a

bundle of Image J and R scripts. These scripts allow for an automated workflow starting from 2-

channel microscopic images of an experimental series (DNA stain, antibody stain), to a graphical

representation and classification of individual cells and eventually to mapping these classifications

back to the original images as a graphical overlay. In the first step nuclei are segmented based on

intensity thresholding (either programmatic or manual), including options for lower and upper size

selection limits. The same threshold is applied to the entire experimental series and the upper size

limit is used to exclude fused structures that cannot be assigned individual cells. A quality score is

assigned to every image based on the fraction of the total DNA stained area that can be assigned

to individual cells (or NETs). This score along with all other parameters of the analysis is exported

as report file and can be used to automatically exclude images from the analysis. In addition, this

part of the script generates a result file that includes the area, circumference (as x,y coordinates)

and cumulative intensities for each channel for every detected nucleus along with information such

as time point or stimulus that can be assigned programmatically. In order to analyse these data

sets we implemented a series of functions in R. These functions include import of Image J result

files, classification of cells based on nuclear area and staining intensity, various plot and data export

functions, as well mapping functions that allow to display the classification of nuclei as color coded

circumferences overlaid on the original images. The scripts are available for download at

https://github.com/tulduro/NETalyser

Quantification of NETs by Image J

NETs were quantified by the semi-automatic method described by Brinkmann et al (2012) and via a second modified method that allowed automatic quantification. All microscopy image datasets were processed by both methods to allow comparison. Hoechst was used to stain total DNA and NETs were additionally stained with anti-chromatin (PL2.3) or 3D9 antibodies (1 μg/ml) and Alexa-568 coupled secondary antibody according to the previous section. Images were acquired with a Leica DMR upright fluorescence microscope equipped with a Jenoptic B/W digital microscope camera and 10x or 20x objective lens. For each experiment, the same exposure settings were used for all samples and a minimum of 3 random fields of view (FOV) were collected. Images were analysed using ImageJ/FIJI software. As described by Brinkmann et al (2012), each channel was imported as an image sequence and converted into a stack. To count total cells/NETs per FOV, the Hoechst channel stack was imported and segmented using the automatic thresholding function (Bernsen method) with radius 15 and parameter 1 set to 35 to produce a black and white thresholded image. Particle analysis was then performed to count all objects the size of a cell nucleus or bigger (10x objective: particle size 25-infinity; 20x objective: particle size 100-infinity) and to exclude background staining artefacts. Total NETs were then counted using the anti-chromatin (PL2.3) or 3D9 channels accordingly. For the method published in 2012, anti-chromatin stains were segmented using manually adjusted thresholding so that the less intense staining of resting cell nuclei was excluded. Particle analysis was then performed to count all objects larger than a resting cell nucleus (10x objective: particle size 75-infinity; 20x objective: particle size 250-infinity). This was also performed for 3D9. In contrast, in the second analysis, the workflow was modified so that the automatic Bernsen thresholding and segmentation were used for both Hoechst and NET channels, total cells and NETs respectively. For each method percentage NETs were calculated as (NETs/Total cells) x100. Results per FOV were then averaged according to sample. A schematic of the different workflows is presented in S.Fig 5.

Immunofluorescent staining of histological samples from tissue sections

Paraffin sections (2 µm thick) were deparaffinized in two changes of 100% xylene for 5 min each and then rehydrated in two changes of 100% ethanol for 5 min each and followed by 90% and 70% ethanol for 5 min each. Sections were washed with 3 changes of deionized water and incubation in TBS (Tris buffered saline). For antigen retrieval, Target Retrieval Solution (TRS pH9) (Dako S2367) was used to incubate the slides in a steam cooker (Braun) for 20 min. After cooling down to room temperature in antigen retrieval buffer, slides were rinsed 3x in deionized water and incubated in TBS until further processing. Slides were blocked with blocking buffer (1% BSA, 5% normal donkey serum, 5% cold water fish gelatin and 0.05% Tween20 in TBS, pH7.4) for 30 min. Blocking buffer was removed, and sections were incubated with primary antibodies at appropriate dilution in blocking buffer (containing 0.05% Triton-X100) overnight at RT. Sections were rinsed in TBS and then incubated with secondary antibody at an appropriate concentration (1:100) for 45 min in the dark at RT and then rinsed three times in TBS for 5 min followed by rinsing with deionized water. Slides were incubated with DNA stain Hoechst 33342 (1:5000) for 5 min, rinsed with water before mounting with Mowiol. Primary antibodies for detection of NETs were as follows: mouse anti-cleaved histone 3 clone 3D9 (2 µg/ml); rabbit anti-histone H3 antibody (citrulline R2 + R8 + R17; ab5103, Abcam); chicken anti-Histone H2B ab134211 Abcam (1:400); sheep anti-ELANE (NE) LS-Bio LSB 4244 Lot 75251. Secondary antibodies were the following: donkey anti-rabbit immunoglobulin G (IgG) heavy and light chain (H&L) Alexa Fluor 488 (Jackson 711-225-152); and donkey anti-mouse IgG H&L Cy3 (Jackson 715-165-151); donkey anti-sheep (IgG) H&L Alexa Fluor 647 (Jackson 713-605-147). An upright widefield microscope (Leica DMR) equipped with a JENOPTIK B/W digital microscope camera or a Leica confocal microscope SP8 were used for fluorescent imaging. Z-stack images were collected at 63× magnification. Where stated colocalisation analysis was performed on confocal images using Volocity 6.5.1 software. Human tonsil and kidney paraffin tissue blocks were purchased from AMSbio. Inflamed tissue from a gallbladder and an appendix was obtained from archived leftover paraffin embedded diagnostic

663 samples and used in an anonymised way after approval through the Charité Ethics Committee
664 (Project EA4/124/19, July 24, 2019). Informed consent from patients for use of biomaterials for
665 research was obtained as part of the institutional treatment contract at Charité.

Contributions

DOT- Conceptualization, methodology, investigation, data curation, figure preparation, writing
original draft, review and editing; UZA - 2-DE, blotting and sample preparation for protein
sequencing; UA -preparation of human pathology tissue samples and microscopy of tissue
samples, image curation; MS - MALDI mass spectrometry; SF - provision and description of human
pathology tissues and advice on histology figure presentation; PRJ - proteomic and 2-DE analysis;
VB - colocalisation analysis, image curation, reviewing and critical feedback on manuscript; AH -
conceptualization, data curation for R analysis, R scripting; AZ- conceptualization, reviewing and
editing of manuscript; AZ and AH provided critical feedback to DOT and helped shape the research,
analysis and manuscript.

Acknowledgements

677 The authors would like to give special thanks to Borko Amulic and Gerben Marsman for their critical
678 feedback at the early stages of manuscript preparation.

Competing interests

680 DOT, AH and AZ and the Max Planck Society have submitted a patent application concerning the
681 contents of this manuscript.

List of supplemental figures

S.Fig 1A. Cysteine protease inhibition by E64 does not inhibit histone H3 cleavage.
S.Fig 1B. AEBSF does not inhibit ROS production & is not cytotoxic.
S.Fig 2A. Schematic summary of extraction & identification of histone H3 cleavage sites in NETs.
S.Fig 2B. Separation of histone H3 by two dimensional electrophoresis & protein identification by
mass spectrometry (contains **S.Table 1**)

S.Fig 3A. Outline of immunisation & screening strategy for antibody production (contains **S.Table**
2)

S.Fig 3B. Peptide inhibition of 3D9 binding to NETs

S.Fig 3C. 3D9 immunoprecipitation

S.Fig 4A. Binding profiles recorded for 3D9 on the linear peptide array.

S.Fig 4B. Binding profiles recorded for 3D9 on helical peptide mimics arrays

S.Table 3. Summary of identified 3D9 binding regions in the peptide array.

S.Fig 4C. Fine epitope mapping by replacement analysis & visualisation of the 3D9 epitope in the
nucleosome core complex.

S.Table 4. 3D9 complete antibody gene sequencing & IMGT V(D)J junction analysis.

S.Fig 5. Workflow of NET analysis methods

S.Fig 7. Comparison of PL2.3 detection in response to apoptotic, necroptotic & necrotic cell death
stimuli

S.Fig 8. Hematoxylin & eosin (HE) stain of kidney section

S.Fig 9. Hematoxylin & eosin (HE) stain of gallbladder & colocalization analysis

S.Fig 10. Hematoxylin & eosin stain of inflamed appendix & colocalization analysis.

Supplemental files

S.file 1. Peptides for epitope mapping
S.file 2. Supplemental Methods
S.file 3. Supplemental Results
709

References

Albrengues, J., Shields, M.A., Ng, D., Park, C.G., Ambrico, A., Poindexter, M.E., Upadhyay, P.,
Uyeminami, D.L., Pommier, A., Kuttner, V.*, et al.* (2018). Neutrophil extracellular traps produced
during inflammation awaken dormant cancer cells in mice. Science *361*.
10.1126/science.aao4227

Amulic, B., Knackstedt, S.L., Abu Abed, U., Deigendesch, N., Harbort, C.J., Caffrey, B.E.,
Brinkmann, V., Heppner, F.L., Hinds, P.W., and Zychlinsky, A. (2017). Cell-Cycle Proteins Control
Production of Neutrophil Extracellular Traps. Dev Cell *43*, 449-462 e445.
10.1016/j.devcel.2017.10.013

Bannister, A.J., and Kouzarides, T. (2011). Regulation of chromatin by histone modifications. Cell
Res *21*, 381-395. 10.1038/cr.2011.22

Bianchi, M., Hakkim, A., Brinkmann, V., Siler, U., Seger, R.A., Zychlinsky, A., and Reichenbach,
J. (2009). Restoration of NET formation by gene therapy in CGD controls aspergillosis. Blood
114, 2619-2622. 10.1182/blood-2009-05-221606.

Borregaard, N., Lollike, K., Kjeldsen, L., Sengelov, H., Bastholm, L., Nielsen, M.H., and Bainton,
D.F. (1993). Human neutrophil granules and secretory vesicles. Eur J Haematol *51*, 187-198.
10.1111/j.1600-0609.1993.tb00629.x.

Brinkmann, V., Goosmann, C., Kuhn, L.I., and Zychlinsky, A. (2012). Automatic quantification of
in vitro NET formation. Frontiers in immunology *3*, 413. 10.3389/fimmu.2012.00413.

Brinkmann, V., Reichard, U., Goosmann, C., Fauler, B., Uhlemann, Y., Weiss, D.S., Weinrauch,
Y., and Zychlinsky, A. (2004). Neutrophil extracellular traps kill bacteria. Science *303*, 1532-1535.
10.1126/science.1092385

Chakravarthy, S., Luger, K. (2005). Nucleosomes containing the histone domain of macroH2A: In
vitro possibilities. 10.2210/pdb2f8n/pdb

Cools-Lartigue, J., Spicer, J., McDonald, B., Gowing, S., Chow, S., Giannias, B., Bourdeau, F.,
Kubes, P., and Ferri, L. (2013). Neutrophil extracellular traps sequester circulating tumor cells
and promote metastasis. The Journal of clinical investigation. 10.1172/JCI67484

Dabrowska, D., Jablonska, E., Iwaniuk, A., and Garley, M. (2019). Many Ways-One Destination:
Different Types of Neutrophils Death. Int Rev Immunol *38*, 18-32.
10.1080/08830185.2018.1540616

Demers, M., Wong, S.L., Martinod, K., Gallant, M., Cabral, J.E., Wang, Y., and Wagner, D.D.
(2016). Priming of neutrophils toward NETosis promotes tumor growth. Oncoimmunology *5*,
e1134073. 10.1080/2162402X.2015.1134073

Dhaenens, M., Glibert, P., Meert, P., Vossaert, L., and Deforce, D. (2015). Histone proteolysis: a
proposal for categorization into 'clipping' and 'degradation'. Bioessays *37*, 70-79.
10.1002/bies.201400118

Diatchuk, V., Lotan, O., Koshkin, V., Wikstroem, P., and Pick, E. (1997). Inhibition of NADPH
oxidase activation by 4-(2-aminoethyl)-benzenesulfonyl fluoride and related compounds. The
Journal of biological chemistry *272*, 13292-13301. 10.1074/jbc.272.20.13292.

Duncan, E.M., Muratore-Schroeder, T.L., Cook, R.G., Garcia, B.A., Shabanowitz, J., Hunt, D.F.,

and Allis, C.D. (2008). Cathepsin L proteolytically processes histone H3 during mouse embryonic

stem cell differentiation. Cell *135*, 284-294. 10.1016/j.cell.2008.09.055

Endo, H., Nakabayashi, Y., Kawashima, S., Enomoto, T., Seki, M., and Horikoshi, M. (2012).

Nucleosome surface containing nucleosomal DNA entry/exit site regulates H3-K36me3 via

association with RNA polymerase II and Set2. Genes Cells *17*, 65-81. 10.1111/j.1365-

2443.2011.01573.x

Galluzzi, L., Vitale, I., Abrams, J.M., Alnemri, E.S., Baehrecke, E.H., Blagosklonny, M.V.,

Dawson, T.M., Dawson, V.L., El-Deiry, W.S., Fulda, S.*, et al.* (2012). Molecular definitions of cell

death subroutines: recommendations of the Nomenclature Committee on Cell Death 2012. Cell

Death Differ *19*, 107-120. 10.1038/cdd.2011.96

Gavillet, M., Martinod, K., Renella, R., Harris, C., Shapiro, N.I., Wagner, D.D., and Williams, D.A.

(2015). Flow cytometric assay for direct quantification of neutrophil extracellular traps in blood

samples. Am J Hematol *90*, 1155-1158. 10.1002/ajh.24185

Hacbarth, E., and Kajdacsy-Balla, A. (1986). Low density neutrophils in patients with systemic

lupus erythematosus, rheumatoid arthritis, and acute rheumatic fever. Arthritis and rheumatism

29, 1334-1342. 10.1002/art.1780291105.

Hakkim, A., Fuchs, T.A., Martinez, N.E., Hess, S., Prinz, H., Zychlinsky, A., and Waldmann, H.

(2011). Activation of the Raf-MEK-ERK pathway is required for neutrophil extracellular trap

formation. Nat Chem Biol *7*, 75-77. 10.1038/nchembio.496.

Howe, C.G., and Gamble, M.V. (2015). Enzymatic cleavage of histone H3: a new consideration

when measuring histone modifications in human samples. Clin Epigenetics *7*, 7. 10.1186/s13148-

014-0041-5

Jacobs, S.A., and Khorasanizadeh, S. (2002). Structure of HP1 chromodomain bound to a lysine

9-methylated histone H3 tail. Science *295*, 2080-2083. 10.1126/science.1069473

Jimenez-Alcazar, M., Kim, N., and Fuchs, T.A. (2017). Circulating Extracellular DNA: Cause or

Consequence of Thrombosis? Seminars in thrombosis and hemostasis. 10.1055/s-0036-1597284

Jungblut, P., Eckerskorn, C., Lottspeich, F., and Klose, J. (1990). Blotting efficiency investigated

by using two-dimensional electrophoresis, hydrophobic membranes and proteins from different

sources. Electrophoresis *11*, 581-588. 10.1002/elps.1150110709

Jungblut, P.R., and Seifert, R. (1990). Analysis by high-resolution two-dimensional

electrophoresis of differentiation-dependent alterations in cytosolic protein pattern of HL-60

leukemic cells. J Biochem Biophys Methods *21*, 47-58. 10.1016/0165-022x(90)90044-d.

Kenny, E.F., Herzig, A., Kruger, R., Muth, A., Mondal, S., Thompson, P.R., Brinkmann, V.,

Bernuth, H.V., and Zychlinsky, A. (2017). Diverse stimuli engage different neutrophil extracellular

trap pathways. Elife *6*. 10.7554/eLife.24437.

Knackstedt, S.L., Georgiadou, A., Apel, F., Abu-Abed, U., Moxon, C.A., Cunnington, A.J.,

Raupach, B., Cunningham, D., Langhorne, J., Kruger, R*., et al.* (2019). Neutrophil extracellular

traps drive inflammatory pathogenesis in malaria. Science immunology *4*.

10.1126/sciimmunol.aaw0336

Kobayashi, S.D., Voyich, J.M., Whitney, A.R., and DeLeo, F.R. (2005). Spontaneous neutrophil

apoptosis and regulation of cell survival by granulocyte macrophage-colony stimulating factor.

Journal of leukocyte biology *78*, 1408-1418. 10.1189/jlb.0605289

Konig, M.F., and Andrade, F. (2016). A Critical Reappraisal of Neutrophil Extracellular Traps and

NETosis Mimics Based on Differential Requirements for Protein Citrullination. Frontiers in

immunology *7*, 461. 10.3389/fimmu.2016.00461

Losman, M.J., Fasy, T.M., Novick, K.E., and Monestier, M. (1992). Monoclonal autoantibodies to

subnucleosomes from a MRL/Mp(-)+/+ mouse. Oligoclonality of the antibody response and

recognition of a determinant composed of histones H2A, H2B, and DNA. Journal of immunology

(Baltimore, Md : 1950) *148*, 1561-1569. (*DOI not assigned*)

Melo, F.R., Vita, F., Berent-Maoz, B., Levi-Schaffer, F., Zabucchi, G., and Pejler, G. (2014).

Proteolytic histone modification by mast cell tryptase, a serglycin proteoglycan-dependent

secretory granule protease. The Journal of biological chemistry *289*, 7682-7690.

10.1074/jbc.M113.546895

Neeli, I., and Radic, M. (2013). Opposition between PKC isoforms regulates histone deimination

and neutrophil extracellular chromatin release. Frontiers in immunology *4*, 38.

10.3389/fimmu.2013.00038

Papayannopoulos, V. (2018). Neutrophil extracellular traps in immunity and disease. Nat Rev

Immunol *18*, 134-147. 10.1038/nri.2017.105

Papayannopoulos, V., Metzler, K.D., Hakkim, A., and Zychlinsky, A. (2010). Neutrophil elastase

and myeloperoxidase regulate the formation of neutrophil extracellular traps. J Cell Biol *191*, 677-

691. 10.1083/jcb.201006052

Pertiwi, K.R., de Boer, O.J., Mackaaij, C., Pabittei, D.R., de Winter, R.J., Li, X., and van der Wal,

A.C. (2018). Extracellular traps derived from macrophages, mast cells, eosinophils and

neutrophils are generated in a time-dependent manner during atherothrombosis. The Journal of

pathology. 10.1002/path.5212

Pieterse, E., Rother, N., Yanginlar, C., Gerretsen, J., Boeltz, S., Munoz, L.E., Herrmann, M.,

Pickkers, P., Hilbrands, L.B., and van der Vlag, J. (2018). Cleaved N-terminal histone tails

distinguish between NADPH oxidase (NOX)-dependent and NOX-independent pathways of

neutrophil extracellular trap formation. Annals of the rheumatic diseases *77*, 1790-1798.

10.1136/annrheumdis-2018-213223

Posthumus, W.P., Lenstra, J.A., Schaaper, W.M., van Nieuwstadt, A.P., Enjuanes, L., and

Meloen, R.H. (1990). Analysis and simulation of a neutralizing epitope of transmissible

gastroenteritis virus. J Virol *64*, 3304-3309. 10.1128/JVI.64.7.3304-3309.1990.

Saitoh, T., Komano, J., Saitoh, Y., Misawa, T., Takahama, M., Kozaki, T., Uehata, T., Iwasaki, H.,

Omori, H., Yamaoka, S.*, et al.* (2012). Neutrophil extracellular traps mediate a host defense

response to human immunodeficiency virus-1. Cell Host Microbe *12*, 109-116.

10.1016/j.chom.2012.05.015

Schauer, C., Janko, C., Munoz, L.E., Zhao, Y., Kienhofer, D., Frey, B., Lell, M., Manger, B., Rech,

J., Naschberger, E.*, et al.* (2014). Aggregated neutrophil extracellular traps limit inflammation by

degrading cytokines and chemokines. Nature medicine *20*, 511-517. 10.1038/nm.3547

Schonrich, G., Kruger, D.H., and Raftery, M.J. (2015). Hantavirus-induced disruption of the

endothelial barrier: neutrophils are on the payroll. Frontiers in microbiology *6*, 222.

10.3389/fmicb.2015.00222

Shechter, D., Dormann, H.L., Allis, C.D., and Hake, S.B. (2007). Extraction, purification and

analysis of histones. Nat Protoc *2*, 1445-1457. 10.1038/nprot.2007.202

Sollberger, G., Tilley, D.O., and Zychlinsky, A. (2018). Neutrophil Extracellular Traps: The Biology

of Chromatin Externalization. Dev Cell *44*, 542-553. 10.1016/j.devcel.2018.01.019

Spengler, J., Lugonja, B., Ytterberg, A.J., Zubarev, R.A., Creese, A.J., Pearson, M.J., Grant,

M.M., Milward, M., Lundberg, K., Buckley, C.D.*, et al.* (2015). Release of Active Peptidyl Arginine

Deiminases by Neutrophils Can Explain Production of Extracellular Citrullinated Autoantigens in
Rheumatoid Arthritis Synovial Fluid. Arthritis & rheumatology (Hoboken, NJ) *67*, 3135-3145. doi:
10.1002/art.39313

Urban, C.F., Ermert, D., Schmid, M., Abu-Abed, U., Goosmann, C., Nacken, W., Brinkmann, V.,
Jungblut, P.R., and Zychlinsky, A. (2009). Neutrophil extracellular traps contain calprotectin, a
cytosolic protein complex involved in host defense against Candida albicans. PLoS Pathog *5*,
e1000639. 10.1371/journal.ppat.1000639

Urban, C.F., Reichard, U., Brinkmann, V., and Zychlinsky, A. (2006). Neutrophil extracellular
traps capture and kill Candida albicans yeast and hyphal forms. Cell Microbiol *8*, 668-676.

Wang, Y., Li, M., Stadler, S., Correll, S., Li, P., Wang, D., Hayama, R., Leonelli, L., Han, H.,
Grigoryev, S.A.*, et al.* (2009). Histone hypercitrullination mediates chromatin decondensation and
neutrophil extracellular trap formation. J Cell Biol *184*, 205-213. 10.1083/jcb.200806072

Yoo, D.G., Winn, M., Pang, L., Moskowitz, S.M., Malech, H.L., Leto, T.L., and Rada, B. (2014).
Release of cystic fibrosis airway inflammatory markers from Pseudomonas aeruginosa-stimulated
human neutrophils involves NADPH oxidase-dependent extracellular DNA trap formation. Journal
of immunology (Baltimore, Md : 1950) *192*, 4728-4738. 10.4049/jimmunol.1301589

Yoshida, M., Sasaki, M., Sugisaki, K., Yamaguchi, Y., and Yamada, M. (2013). Neutrophil
extracellular trap components in fibrinoid necrosis of the kidney with myeloperoxidase-ANCA-
associated vasculitis. Clin Kidney J *6*, 308-312. 10.1093/ckj/sft048

Zenaro, E., Pietronigro, E., Della Bianca, V., Piacentino, G., Marongiu, L., Budui, S., Turano, E.,
Rossi, B., Angiari, S., Dusi, S.*, et al.* (2015). Neutrophils promote Alzheimer's disease-like
pathology and cognitive decline via LFA-1 integrin. Nature medicine *21*, 880-886.
10.1038/nm.3913

Zhou, Y., Chen, B., Mittereder, N., Chaerkady, R., Strain, M., An, L.L., Rahman, S., Ma, W., Low,
C.P., Chan, D.*, et al.* (2017). Spontaneous Secretion of the Citrullination Enzyme PAD2 and Cell
Surface Exposure of PAD4 by Neutrophils. Frontiers in immunology *8*, 1200.
10.3389/fimmu.2017.01200