**Structural basis for late maturation steps of the human mitoribosomal large subunit**

**Authors:** Miriam Cipullo[1,2,†], Genís Valentín Gesé[3,†], Anas Khawaja[1,2],

B. Martin Hällberg[3,4,*], Joanna Rorbach[1,2,*]

**Affiliations:**

[1] Department of Medical Biochemistry and Biophysics, Division of Molecular Metabolism,

Karolinska Institutet, Solnavägen 9, 171 65 Solna, Sweden

[2] Max Planck Institute Biology of Ageing - Karolinska Institutet Laboratory, Karolinska

Institutet, Stockholm, Sweden

[3] Department of Cell and Molecular Biology, Karolinska Institutet, Solnavägen 9, 171 65

Solna, Sweden

[4] Centre for Structural Systems Biology (CSSB) and Karolinska Institutet VR-RÅC,

Notkestrasse 85, 22607 Hamburg, Germany.

†Joint First Authors

*Correspondence to:

B. Martin Hällberg

Address: Department of Cell and Molecular Biology, Karolinska Institutet, Solnavägen 9, 171

65 Solna, Sweden

Email: martin.hallberg@ki.se

Joanna Rorbach

Department of Medical Biochemistry and Biophysics, Research Division of Molecular

Metabolism, Karolinska Institutet, Solnavägen 9, 171 65 Solna, Sweden

Email: joanna.rorbach@ki.se

**Abstract**

Mitochondrial ribosomes (mitoribosomes) synthezise a critical set of proteins essential for oxidative phosphorylation. Therefore, their function is vital to cellular energy supply and mitoribosomal defects give rise to a large and diverse group of human diseases [1]. The architecture of mitoribosomes is strikingly different from that of their bacterial and eukaryotic cytosolic counterparts and display high divergence between species [2–6]. Mitoribosome biogenesis follows distinct molecular pathways that remain poorly understood. Here, we determined the cryo-EM structures of mitoribosomes isolated from human cell lines with either depleted or overexpressed mitoribosome assembly factor GTPBP5. This allowed us to capture consecutive steps during mitoribosomal large subunit (mt-LSU) biogenesis that involve normally short-lived assembly intermediates. Our structures provide important insights into the last steps of 16S rRNA folding, methylation and peptidyl transferase centre (PTC) completion, which require the coordinated action of nine assembly factors. We show that mammalian-specific MTERF4 contributes to the folding of 16S rRNA, allowing 16S rRNA methylation by MRM2, while GTPBP5 and NSUN4 promote fine-tuning rRNA rearrangments leading to PTC formation. Moreover, our data reveal an unexpected role for the elongation factor mtEF-Tu in mt-LSU assembly, in which mt-EF-Tu interacts with GTPBP5 in a manner similar to its interaction with tRNA during translational elongation. Together, our approaches provide detailed understanding of the last stages of mt-LSU biogenesis that are unique to mammalian mitochondria.

## Main

Mammalian mitoribosomes assemble in a multi-step process that includes the maturation of two ribosomal RNAs (rRNAs; 12S and 16S), a structural tRNA, and incorporation of 82 mitoribosomal proteins (MRPs) [7]. Multiple assembly factors, many specific to mammalian mitochondria, assist in the mitoribosome assembly process. A growing number of studies in recent years have shown that a family of GTP-binding proteins (GTPBPs) is crucial for mammalian mitoribosome assembly [8–13]. Among these GTPBPs, GTPBP5 participates in the late steps of large subunit (mt-LSU) maturation, and its deletion leads to severe translational defects [8,11].

To understand the molecular basis for the late stages of human mitochondrial mt-LSU assembly, we used single-particle electron cryomicroscopy (cryo-EM) to determine the structure of an mt-LSU intermediate isolated from GTPBP5-deficient cells (GTPBP5$^{KO}$). In addition, we determined the cryo-EM structure of a GTPBP5-bound mt-LSU intermediate immunoprecipitated from cells expressing a tagged variant of GTPBP5 (GTPBP5$^{IP}$).

## Composition of the GTPBP5$^{KO}$ and GTPBP5$^{IP}$ mt-LSU assembly intermediates

Both the GTPBP5$^{KO}$ and GTPBP5$^{IP}$ mt-LSU assembly-intermediates reveal several trapped assembly factors: the MTERF4-NSUN4 complex, MRM2, MTG1 and the MALSU1:L0R8F8:mt-ACP module (Fig. 1a,b, Extended Data Fig. 1 and 2). Furthermore, the GTPBP5$^{IP}$ mt-LSU structure features GTPBP5 and the mitochondrial elongation factor mtEF-Tu (Fig. 1b, Extended Data Fig. 2). Comparing the GTPBP5$^{KO}$ and the GTPBP5$^{IP}$ mt-LSU intermediates with the mature mt-LSU [14] reveals two crucial differences in the 16S rRNA conformation (Fig. 1c,d). First, in both GTPBP5$^{KO}$ and GTPBP5$^{IP}$ intermediates, MTERF4 in the MTERF4-NSUN4 complex, binds an immaturely folded region of the 16S domain IV. This region (C2548-G2631) – corresponding to helices H68, H69, and H71 of the mature mt-LSU – is folded into a novel intermediate rRNA helical structure, hereafter denoted helix-X (Fig. 1). The helix-X occupies a different position on the mt-LSU than H68-71 in the mature mt-LSU, where helices H68-71 and H89-90 jointly form the peptidyl-transferase centre (PTC) (Fig. 1d). MTERF4s binding of helix-X partly orders the disordered rRNA in the mt-LSU assemblies' subunit interface side (Extended Data Fig. 1) and thereby enables MRM2 to bind (Fig. 1d). Second, in the GTPBP5$^{KO}$ – but not in the GTPBP5$^{IP}$ – the junction between H89 and H90 of domain V is significantly different compared to the mature mitoribosome (Fig. 1c, d). Specifically, at the base of H89, one helical turn remains unfolded and instead forms a flexible loop in the GTPBP5$^{KO}$ (Fig. 1d).

**MTERF4-NSUN4 complex steers the final steps of 16S rRNA folding and allows for**

**MRM2 binding**

The MTERF4-NSUN4 complex, previously shown to be essential for monosome assembly [16,17], binds at the intersubunit interface in our GTPBP5[KO] and GTPBP5[IP] structures (Fig. 1a, b). The C-terminal part of MTERF4 binds to NSUN4 close to the NSUN4 N-terminus in a mixed hydrophobic-polar binding interface similar to earlier crystal structures of the isolated complex [18,19] (Fig. 2a). NSUN4 was previously shown to $m^5C$-methylate the C1488 carbon 5 in 12S mt-rRNA [17]. In our structures, the active site of NSUN4 is turned towards the mt-LSU core (Extended Data Fig. 3a), impeding methylation of the 12S mt-rRNA. Although the methyl-donor S-adenosyl-methionine (SAM) is observed in the NSUN4 active site, no RNA substrate is present. Furthermore, in the GTPBP5[KO] and GTPBP5[IP] structures, the MTERF4-NSUN4 complex is bound and bent from two sides by uL2m. Specifically, a uL2m C-terminal extension penetrates in between NSUN4 and MTERF4 to further stabilize the MTERF4-NSUN4 binding interface and decreases the curvature of the MTERF4 solenoid relative to the crystal structures (Fig. 2a). This reforming of the MTERF4 solenoid is necessary to bind the helix-X rRNA region in the strongly positively charged concave side of MTERF4 (Extended Data Fig. 3b). Here, MTERF4 forms an extensive network of contacts with helix-X that stabilizes the association and promotes helix-X folding (Fig. 2b). The mature H71 base-pairing is already formed within helix-X. Thus, by binding to helix-X, MTERF4 intiates the folding of this 16S mt-rRNA region. Furthermore, it also exposes the A-loop, which is obstructed by H68, H69, and H71 in the mature mt-LSU (Fig. 1d), thereby allowing MRM2 binding.

Similarly to a previously determined mt-LSU assembly intermediate [20] there is a MALSU1-module positioned adjacent to uL14m in both the GTPBP5[KO] and GTPBP5[IP] (Fig.1a,b). Furthermore, MTG1 (GTPBP7), which assists in late-stage mt-LSU maturation [21], is bound in the vicinity of helix-X (Fig. 1a,b). MTG1 contacts the C-terminus of MALSU1 (Extended Data Fig. 4a) and the region encompassing A2554-U2602 of helix-X. This region could not modelled due to the lower local resolution, but the contact is visible in the electron density map (Extended Data Fig. 4b). Interestingly, the position of human MTG1 in our structures differs significantly from its bacterial and trypanosomal counterparts ([3,22], Extended Data Fig. 4c). Specifically, while in other systems MTG1 homologs contact the rRNA, reaching out towards the PTC (Extended Data Fig. 4c), in the trapped intermediates described here MTG1 is unlikely to induce pronounced conformational changes of the PTC or participate in the recruitment/dissociation of assembly factors.

MRM2 2′-O-methylates U3039 in the 16S A-loop during mt-LSU assembly [23,24] and

in our GTPBP5[KO] and GTPBP5[IP] structures, MRM2 binds in the mt-LSU intersubunit

interface (Fig. 1a,b). It features two N-terminal $\alpha$-helices followed by a canonical S-adenosyl-

L-methionine-dependent methyltransferase domain (SAM MTase) (Fig. 2c). In GTPBP5[KO],

but not in GTPBP5[IP], the two N-terminal $\alpha$-helices extend from MRM2 and insert into the

rRNA core to thereby displace and retrieve the A-loop (16S mt-rRNA domain V) through a

complex interaction network (Fig. 2d,e). This places the 2′-hydroxyl of U3039 close to the

ideal methyl-acceptor position in the MRM2 active site (Fig. 2d). However, there is no

density for either SAM or S-adenosyl homocysteine (SAH) in the MRM2 active site and there

is no apparent density for a 2′-O-methyl on U3039 (Extended Data Fig. 5). Interestingly,

G3040 that is 2′-O-methylated by MRM3 [23,24], is methylated in our structures (Extended Data

Fig. 5). Hence, 2′-O-methylation by MRM3 takes place prior to MRM2 methylation in human

mitoribosome biogenesis.

**GTPBP5 promotes remodelling of the PTC**

GTPBP5 consists of a glycine-rich N-terminal domain (Obg-domain) and a C-terminal

GTPase domain (G-domain) (Fig. 3a). In our GTPBP5[IP] structure, the G-domain has GTP in

its active site and is wedged between the L7/L12 stalk and the MALSU1 module (Fig. 1b and

Extended Data Fig. 6a). The Obg-domain protrudes into the PTC (Fig. 3b), thereby displacing

the A-loop from the MRM2 active site and expelling the MRM2 N-terminal $\alpha$-helices from

the rRNA core (Fig. 3c), while the A-loop folds into the fully mature position (Fig. 3b3).

The protruding Obg-domain is positioned between H89 and H93 and occupies the space that

accommodates the acceptor arm of the A-site tRNA during translation (Extended Data Fig.

6b). Hereby, GTPBP5 adopts a tRNA mimicry strategy, similar to ObgE of *E. coli* [25]. The

Obg-domain contains six glycine-rich sequence motifs that form antiparallel polyproline-II

helices (helices a–f) (Fig. 3b). Helices c and d bind the A-loop, while the loop between

helices e and f inserts into the major groove of H93. The loop between a and b inserts at the

triple-junction formed between H89-H90-H93 (Fig. 3b).

Comparison of the GTPBP5[KO] and the GTPBP5[IP] structures with the mature mt-LSU reveals

extensive maturation of the PTC upon GTPBP5 binding. The partly unfolded H89 in the

GTPBP5[KO] is folded in the GTPBP5[IP] (Fig. 3 b1). This folding is coordinated by the joint

action of GTPBP5 and NSUN4: in the presence of GTPBP5, the extended N-terminal region

of NSUN4 inserts into the rRNA core and temporarily displaces the P-loop (Fig. 3 b2),

thereby breaking the P-loop interaction with H89 (Fig. 3d). As a consequence, H89 is given

the space necessary to fold into a structure similar to its mature form (Fig. 3 b1, lower).

The GTPBP5[IP] structure shows a rotation of the L7/L12 stalk in comparison to the GTPBP5[KO]

structure (Extended Data Fig. 6c). Here, the rRNA in the L54/L11 region of the stalk forms π-

stacking interactions with two residues of the GTP-ase switch I element of GTPBP5

(Extended Data Fig. 6a,c). Thereby, the L7/L12 stalk stabilizes the "state 2" conformation of

the switch I. In this way, the rotated L7/L12 stalk stabilizes the GTP-state of GTPBP5[26] and

consequently a GTP is bound in our structure (Extended Data Fig. 6a,c). The requirement for

GTPBP5 to be in a GTP-bound state is supported by the inability of a GTPBP5 Walker A

mutant (GTPBP5-S238A) to bind mt-LSU intermediates [8]. A back-rotation of the L7/12 stalk,

presumably by binding of another maturation factor to the mt-LSU assembly intermediate,

would lead to a release of the switch I and the activation of GTP hydrolysis, followed by

release of GTPBP5 from the mt-LSU assembly intermediate. Taken together, GTPBP5 plays

a direct and active role in rRNA remodeling and, together with the NSUN4 N-terminus,

orchestrates the maturation of mitoribosomal PTC.

**Translation elongation factor mtEF-Tu is involved in mitoribosome assembly**

mtEF-Tu consists of a GTPase domain (G-domain; domain I) and two structurally

similar β-stranded domains (domains II and III) (Fig. 4a). It was recently shown that during

translation, mtEF-Tu·GTP delivers aminoacylated-tRNA to the mitoribosome in a manner

similar to its bacterial EF-Tu counterparts (Extended Data Fig. 7a,[27]). In contrast, the binding

of an EF-Tu·GTP·aa-tRNA complex is sterically hindered by the MALSU1 module bound in

our mt-LSU intermediates (Extended Data Fig. 7a). Unexpectedly, mtEF-Tu binds to the

mitoribosome in a unique manner in our GTPBP5[IP] structure (Fig. 4b). Here, domains II and

III establish extensive interactions with GTPBP5, the sarcin-ricin loop (SRL) and the

MALSU1 stalk (Fig. 4b,c). In addition, the G-domain switch I element, in its "state 1"/GDP

conformation (Extended Data Fig. 7b), extends and binds MALSU1. Thereby, mtEF-Tu,

together with the SRL and MALSU1, forms a platform for GTPBP5 binding (Fig. 4b). These

structural conclusions are supported by earlier mass-spectrometry data on isolated GTPBP5[IP]

assembly intermediates and protein-proximity interactome analysis [8,28].

The G-domain of mtEF-Tu does not contact the SRL as in mtEF-Tu's canonical role in

translation but instead binds to the C-terminal region of a bL12m that also contacts uL10m at

the stalk base (Fig. 4b). In bacteria, homologs to bL12m and uL10m, recruit and activate

translational GTPases such as EF-Tu via the bL12m C-terminal domain[29,30] and stimulate

GTP hydrolysis 1000-fold[31]. Taken together, this suggests that mtEF-Tu hydrolysis –

stimulated by bL12m and uL10m – is used to accommodate GTPBP5 on the maturating mt-

LSU in analogy to the canonical EF-Tu role in translation, in which aminoacylated-tRNA is

accommodated on the translating ribosome (Extended Data Fig. 7a).

208
**Concluding remarks**

Our analyses shed new light into mammalian mitoribosome maturation and explain

the essential roles of several assembly factors that together promote fine RNA rearrangments

and lead to the mt-LSU completion. Thanks to our approaches that combine biochemical tools

with structural determination, we were able to uncovered several features unique to

mammalian mitochondria. Based on these data, we propose a model of the late-stage mt-

LSU assembly that requires the interplay of nine auxiliary factors (Fig. 4d).

Lastly, as defects in mitoribosome biogenesis – resulting from, for example, mutations

in MRM2 and GTPBP5 – are increasingly implicated in mitochondrial disease [32,33], the

current work does not only describes fundamental cellular processes but may also further new

diagnostic and therapeutic approaches to mitochondrial diseases.

220
221   **Methods**

222
**Generation of GTPBP5 knock-out cell line**

The knock-out cell line (GTPBP5[KO]) was generated in the Flp-In T-Rex human embryonic

kidney 293 (HEK293T) cell line (Invitrogen) using CRISPR/Cas9 technology targeted on

exon 1 of *MTG2* gene, which encodes for GTPBP5, as described [8]. In short, two pairs of

gRNAs were designed and cloned into the pSpCas9(BB)-2A-Puro (pX459) V2.0 vector to

generate out-of-frame deletions. Transfection of HEK293T cell line with the pX459 variants

was performed using Lipofectamine 3000 following manufacturer's instructions. Selection of

transfected cells was done using puromycin treatment at a final concentration of 1.5 mg/ml

for 48 hours. Subsequently, cells were single-cell diluted and transferred into a 96-well plate.

Selected clones were screened via Sanger sequencing and Western blotting.

233
**Purification of the mt-LSU from GTPBP5[KO] cell line via sucrose gradient centrifugation**

Isolation of mitochondria was performed from GTPBP5[KO] cell line as described in Rorbach *et*

*al.* [24], with some modifications. Crude mitochondria were further purified via differential

centrifugation by being loaded onto a sucrose gradient (1 M and 1.5 M sucrose, 20 mM
Tris-HCl pH 7.5, 1 mM EDTA) and centrifuged at 25000 rpm for 1 hour at 4°C (Beckman
Coulter SW41-Ti rotor). Mitochondria forming a band at the interphase between the 1 M and
1.5 M sucrose were collected and resuspended in 10 mM Tris-HCl pH = 7.5 in 1:1 ratio. After
centrifugation, the final mitochondrial pellet was resuspended in mitochondrial freezing
buffer (200 mM trehalose, 10 mM Tris-HCl pH 7.5, 10 mM KCl, 0.1% BSA, 1 mM
EDTA), snap frozen in liquid nitrogen and stored at -80°C.
The mt-LSU was purified from the GTPBP5[KO] cell line via a sucrose gradient centrifugation
experiment. Mitochondria were lysed at 4°C for 20 minutes (25 mM HEPES-KOH pH = 7.5,
20 mM Mg(OAc)$_2$, 100 mM KCl, 2% (v/v) Triton X-100, 2 mM dithiothreitol (DTT), 1x
cOmplete EDTA-free protease inhibitor cocktail (Roche), 40 U/µl RNase inhibitor
(Invitrogen)) and later centrifuged at 13000 rpm for 5 minutes at 4°C. For mitoribosome
purification, the mitolysate was subjected to sucrose cushion ultracentrifugation method (0.6
M sucrose, 25 mM HEPES-KOH pH = 7.5, 10 mM Mg(OAc)$_2$, 50 mM KCl, 0.5% (v/v)
Triton X-100, 2 mM DTT) by being centrifuged at 73000 rpm for 45 minutes at 4°C
(Beckman Coulter TL120.2 rotor). The mitoribosomal pellet was subsequently resuspended in
ribosome resuspension buffer (25 mM HEPES-KOH pH = 7.5, 10 mM Mg(OAc)$_2$, 50 mM
KCl, 0.05% DDM, 2 mM DTT) and centrifuged at 13000 rpm for 10 minutes at 4°C. The
obtained supernatant was then loaded onto a linear sucrose gradient (15-30% (w/v)) in 1x
gradient buffer (25 mM HEPES-KOH pH = 7.5, 10 mM Mg(OAc)$_2$, 50 mM KCl, 0.05%
DDM, 2 mM DTT) and centrifuged for 2 hours and 15 minutes at 39000 rpm at 4°C
(Beckman Coulter TLS55 rotor). Fractions corresponding to the large mitochondrial subunit
were collected and  subjected to buffer exchange (25 mM HEPES-KOH pH = 7.5, 10 mM
Mg(OAc)$_2$, 50 mM KCl) using Vivaspin 500 centrifugal concentrators.
261
**Generation of a mammalian cell line expressing GTPBP5**
A stable mammalian cell line overexpressing C-terminal FLAG-tagged GTPBP5
(GTPBP5::FLAG) in a doxycycline-inducible dose-dependent manner was generated as
described in Cipullo et al. [8]. The GTPBP5 cDNA (hORFeome Database; Internal ID: 12579)
was cloned into pcDNA5/FRT/TO. Flp-In T-Rex human embryonic kidney 293 (HEK293T,
Invitrogen) cells were cultured in DMEM (Dulbecco's modified eagle medium) containing
10% (v/v) tetracycline-free fetal bovine serum (FBS), 2 mM Glutamax (Gibco), 1x
Penicillin/Streptomycin (Gibco), 50 µmg/ml uridine, 10 µg/ml Zeocin (Invitrogen) and 100
µg/ml blasticidin (Gibco) at 37 °C under 5% $CO_2$ atmosphere. Cells were seeded in a 6-well

plate, grown in medium without antibiotics and co-transfected with pcDNA5/FRT/TO-GTPBP5::FLAG and pOG44 using Lipofectamine 3000 according to manufacturer's recommendations. After 48 hours, selection of cells was promoted by addition of hygromycin (100 µg/ml, Invitrogen) and blasticidin (100 µg/ml) to culture media. After two to three weeks post-transfection, single colonies were picked and GTPBP5 overexpression was tested via Western Blot analysis 48 hours after induction with 50 ng/ml doxycycline.

**Immunoprecipitation experiment**

Isolation and purification of mitochondria from GTPBP5::FLAG overexpressing cell line was performed as described in the above paragraph "Purification of the mt-LSU from GTPBP5[KO] cell line via sucrose gradient centrifugation". The mt-LSU bound with GTPBP5 was isolated via FLAG-immunoprecipitation analysis (IP). Pelleted mitochondria were lysed at 4°C for 20 minutes (25 mM HEPES-KOH pH = 7.5, 20 mM $Mg(OAc)_2$, 100 mM KCl, 2% (v/v) Triton X-100, 0.2 mM DTT, 1x cOmplete EDTA-free protease inhibitor cocktail (Roche), 40 U/□l RNase inhibitor (Invitrogen)) and centrifuged at 5000g for 5 minutes at 4°C. The supernatant was then added to ANTI-FLAG M2-Agarose Affinity Gel (Sigma-Aldrich) previously equilibrated (25 mM HEPES-KOH pH = 7.5, 5 mM $Mg(OAc)_2$, 100 mM KCl, 0.05% DDM) and incubated for 3 hours at 4°C. After incubation, the sample was centrifuged at 5000g for 1 minute at 4°C, the supernatant was removed and the gel was washed three times with wash buffer. Elution (25 mM HEPES-KOH pH = 7.5, 5 mM $Mg(OAc)_2$, 100 mM KCl, 0.05% DDM, 2 mM DTT) was performed using 3x FLAG Peptide (Sigma-Aldrich) for about 40 minutes at 4°C.

**Cryo-EM data acquisition and image processing**

Prior to cryo-EM grid preparation, grids were glow-discharged with 20 mA for 30 seconds using a PELCO easiGlow glow-discharge unit. Quantifoil Cu 300 mesh (R 2/2 geometry; Quantifoil Micro Tools GMBH) covered with a thin layer of 3 nm carbon were used for the for the GTPBP5[KO] sample. Carbon lacey films (400 mesh Cu grid; Agar Scientific) mounted with ultrathin carbon support film were used for the GTPBP5[IP] sample. Three □l aliquots of sample were applied to the grids, which were then vitrified in a Vitrobot Mk IV (Thermo Fisher Scientific) at 4°C and 100% humidity (blot 10 s, blot force 3, 595 filter paper (Ted Pella Inc.)). Cryo-EM data collection (Extended Data Table 1) was performed with EPU (Thermo Fisher Scientific) using a Krios G3i transmission-electron microscope (Thermo Fisher Scientific) operated at 300 kV in the Karolinska Institutet's 3D-EM facility. Images

were acquired in nanoprobe EFTEM SA mode with a slit width of 10 eV using a K3

Bioquantum during 1 second during which 60 movie frames were collected with a flux of

0.82 e$^-$/Å$^2$ per frame. Motion correction, CTF-estimation, Fourier binning (to 1.02 Å/px),

picking and extraction in 600 pixel boxes (size threshold 300 Å, distance threshold 20 Å,

using the pretrained BoxNet2Mask_20180918 model) were performed on the fly using Warp

[34]. Only particles from micrographs with an estimated resolution of 3.6 Å and underfocus

between 0.2 and 3 □m were retained for further processing.

For the GTPBP5$^{KO}$ dataset, 704720 particles were picked from 37307 micrographs (Extended

Data Fig. 1). The particles were imported into CryoSPARC 2.15 [35] for further processing.

After 2D classification, 130289 particles were selected for an ab-initio reconstruction. This

reconstruction, in addition to two "bad" reconstructions created from bad 2D class-averages,

were used for heterogeneous refinement of the complete particle set resulting in one of the

three classes yielding a large-subunit reconstruction with high resolution features (196318

particles). After homogeneous refinement of these particles, the PDB model of a

mitochondrial LSU assembly intermediate (PDB: 5OOL [20]) was fitted in the density. The

reconstruction contained the MALSU1 module and also featured weak unexplained densities

for several additional components in the intersubunit interface. A 3D variability analysis was

performed with a mask on the intersubunit interface and a low pass resolution of 10 Å, and

subsequently used for clustering into six particle classes representing different assembly

intermediates. Two of the classes (43057 and 41619 particles) lacked the density for the A-

and P-loops, H89, helices 68-71 and the L7/12 stalk. The A- and P-loops become visible in

the third class (28001 particles). The fourth class revealed a number of biogenesis factors:

MRM2, MTERF4-NSUN4, MTG1 and the structured H67-H71 rRNA region (helix-X)

(48646 particles) as well as H89. All the biogenesis factors are absent in the fifth class, in

which helices 68 and 71 move to the mature position (26678 particles). H69 is nevertheless

not visible. The last class contains the small subunit (8317 particles). Non-uniform refinement

of the fourth particle set yielded a reconstruction at 2.64. Å, which was used for model

building and refinement. As the density for MTG1 was weaker than for the other factors, 3D

variability analysis was performed with a mask on the MTG1 region and a 10 Å low-pass

filter to select particles containing MTG1 (19254 particles, which was subsequently subjected

to homogeneous refinement yielding a reconstruction at 2.90 Å).

For the GTPBP5$^{IP}$ dataset, 283598 particles were picked from 112076 micrographs using

WARP and imported into CryoSPARC 2.15 (Extended Data Fig. 2) [35]. The complete particle

set was used in heterogeneous refinement against the same three references derived from the

GTPBP5[KO] dataset. One of the classes (78306 particles) yielded a high-resolution

reconstruction of the mt-LSU assembly intermediate. After homogeneous refinement,

additional density for GTPBP5 was visible in the intersubunit interface. 3D variability

analysis was performed with a mask on the GTPBP5 region and a low pass resolution of 10

Å. Subsequent clustering into two particle clusters revealed a particle subset containing

GTPBP5 (71834 particles), which was used for model building and refinement. This

reconstruction also features densities for MRM2 and MTERF4-NSUN4. In addition, a weak

density was present for mtEF-TU, the bL12m C-terminal domain and MTG1. The refined

particles were subject to 2D classification and the bad classes were removed. The remaining

particles were polished and refined in Relion 3.1 and re-imported into CryoSPARC for further

processing. 3D variability analysis was performed on these particles with a mask covering

mtEF-TU, bL12m and MTG1 and a 10 Å low pass filter. Subsequent clustering (4 clusters)

revealed 2 clusters containing mtEF-TU/bL12m (17886 particles in total), one cluster

containing MTG1 (8233 particles) and one cluster containing all three proteins (13376

particles). The reconstructions derived from the MTG1- and the mtEF-Tu-containing particles

reached a resolution of 3.19 and 3.21 Å respectively.

**Model building and refinement**

Model building of the GTPBP5[KO] and GTPBP5[IP] mt-LSU assembly intermediate structures

was performed using *Coot* [36]. The structure of a previous mt-LSU assembly intermediate

(PDB 5OOL, [20]) was used as a starting model. MRM2 and MTERF4-NSUN4 were identified

by modelling secondary structure elements in *Coot*, and using the initial models for a

structural search using the DALI server [37]. MTG1, as well as mtEF-Tu and the bL12m C-

terminal domain in the GTPBP5-bound mt-LSU dataset, were identified using a density-based

fold-recognition pipeline [20]. Using SWISS-MODEL [38], we generated homology models for

the human GTPBP5 (template: PDB 4CSU chain 9 [25]), MTG1 (template: PDB 3CNL chain A

[39]), bL12m (template: PDB 1DD3 chain A [40]) and mtEF-TU (template: PDB 1D2E chain A

[41]). All the models, as well as the crystal structure of the human MTERF4-NSUN4 (template:

PDB 4FP9 chains A and B [18]), were fitted into the density map using *Coot* JiggleFit. The

MTG1 GTPase domain and the L17/12 stalk were excluded from atomic refinement and were

only subject to rigid body refinement. Metal ions and modifications were placed based on

map densities. Stereochemical refinement was performed using PHENIX [42]. Refinement

statistics are reported in Extended Data Table 2, while modeled proteins and rRNA are shown

in Extended Data Table 3. Validation of the final models was done via MolProbity [43]. Figures

were generated using ChimeraX [44].

**References**

1.    Hällberg, B. M. & Larsson, N.-G. Making Proteins in the Powerhouse. *Cell Metab.* **20**,

226–240 (2014).

2.    Greber, B. J. *et al.* The complete structure of the 55S mammalian mitochondrial

ribosome. *Science (80-. ).* **348**, 303–308 (2015).

3.    Jaskolowski, M. *et al.* Structural Insights into the Mechanism of Mitoribosomal Large

Subunit Biogenesis. *Mol. Cell* **79**, 629-644.e4 (2020).

4.    Saurer, M. *et al.* Mitoribosomal small subunit biogenesis in trypanosomes involves an

extensive assembly machinery. *Science (80-. ).* **365**, 1144–1149 (2019).

5.    Perez Boerema, A. *et al.* Structure of the chloroplast ribosome with chl-RRF and

hibernation-promoting factor. *Nat. Plants* **4**, 212–217 (2018).

6.    Amunts, A., Brown, A., Toots, J., Scheres, S. H. W. & Ramakrishnan, V. The structure

of the human mitochondrial ribosome. *Science (80-. ).* **348**, 95–98 (2015).

7.    Bogenhagen, D. F., Ostermeyer-Fay, A. G., Haley, J. D. & Garcia-Diaz, M. Kinetics

and Mechanism of Mammalian Mitochondrial Ribosome Assembly. *Cell Rep.* **22**,

1935–1944 (2018).

8.    Cipullo, M. *et al.* Human GTPBP5 is involved in the late stage of mitoribosome large

subunit assembly. *Nucleic Acids Res.* (2020) doi:10.1093/nar/gkaa1131.

9.    Lavdovskaia, E. *et al.* The human Obg protein GTPBP10 is involved in mitoribosomal

biogenesis. *Nucleic Acids Res.* **46**, 8471–8482 (2018).

10.   Maiti, P., Kim, H.-J., Tu, Y.-T. & Barrientos, A. Human GTPBP10 is required for

mitoribosome maturation. *Nucleic Acids Res.* (2018) doi:10.1093/nar/gky938.

11.   Maiti, P., Antonicka, H., Gingras, A.-C., Shoubridge, E. A. & Barrientos, A. Human

GTPBP5 (MTG2) fuels mitoribosome large subunit maturation by facilitating 16S

rRNA methylation. *Nucleic Acids Res.* **48**, 7924–7943 (2020).

12.   Kim, H.-J. & Barrientos, A. MTG1 couples mitoribosome large subunit assembly with

intersubunit bridge formation. *Nucleic Acids Res.* **46**, 8435–8453 (2018).

13.   Lavdovskaia, E. *et al.* Dual function of GTPBP6 in biogenesis and recycling of human

mitochondrial ribosomes. *Nucleic Acids Res.* **48**, 12929–12942 (2020).

14.   Brown, A. *et al.* Structure of the large ribosomal subunit from human mitochondria.

*Science (80-. ).* **346**, 718–722 (2014).

15.   Aibara, S., Singh, V., Modelska, A. & Amunts, A. Structural basis of mitochondrial
translation. *Elife* **9**, (2020).

16.   Cámara, Y. *et al.* MTERF4 Regulates Translation by Targeting the Methyltransferase
NSUN4 to the Mammalian Mitochondrial Ribosome. *Cell Metab.* **13**, 527–539 (2011).

17.   Metodiev, M. D. *et al.* NSUN4 Is a Dual Function Mitochondrial Protein Required for
Both Methylation of 12S rRNA and Coordination of Mitoribosomal Assembly. *PLoS*
*Genet.* **10**, e1004110 (2014).

18.   Spåhr, H., Habermann, B., Gustafsson, C. M., Larsson, N.-G. & Hallberg, B. M.
Structure of the human MTERF4–NSUN4 protein complex that regulates
mitochondrial ribosome biogenesis. *Proc. Natl. Acad. Sci.* **109**, 15253–15258 (2012).

19.   Yakubovskaya, E. *et al.* Structure of the Essential MTERF4:NSUN4 Protein Complex
Reveals How an MTERF Protein Collaborates to Facilitate rRNA Modification.
*Structure* **20**, 1940–1947 (2012).

20.   Brown, A. *et al.* Structures of the human mitochondrial ribosome in native states of
assembly. *Nat. Struct. Mol. Biol.* **24**, 866–869 (2017).

21.   Kim, H. J. & Barrientos, A. MTG1 couples mitoribosome large subunit assembly with
intersubunit bridge formation. *Nucleic Acids Res.* **46**, 8435–8453 (2018).

22.   Seffouh, A. *et al.* Structural consequences of the interaction of RbgA with a 50S
ribosomal subunit assembly intermediate. *Nucleic Acids Res.* **47**, 10414–10425 (2019).

23.   Lee, K. W. & Bogenhagen, D. F. Assignment of 2′-O-methyltransferases to
modification sites on the mammalian mitochondrial large subunit 16 S ribosomal RNA
(rRNA). *J. Biol. Chem.* **289**, 24936–24942 (2014).

24.   Rorbach, J. *et al.* MRM2 and MRM3 are involved in biogenesis of the large subunit of
the mitochondrial ribosome. *Mol. Biol. Cell* **25**, 2542–2555 (2014).

25.   Feng, B. *et al.* Structural and Functional Insights into the Mode of Action of a
Universally Conserved Obg GTPase. *PLoS Biol.* **12**, e1001866 (2014).

26.   Matsumoto, S. *et al.* Molecular Mechanism for Conformational Dynamics of Ras·GTP
Elucidated from In-Situ Structural Transition in Crystal. *Sci. Rep.* **6**, 25931 (2016).

27.   Desai, N. *et al.* Elongational stalling activates mitoribosome-associated quality control.
*Science (80-. ).* **370**, 1105–1110 (2020).

28.   Antonicka, H. *et al.* A High-Density Human Mitochondrial Proximity Interaction
Network. *Cell Metab.* **32**, 479-497.e9 (2020).

29.   Traut, R. R. *et al.* Location and domain structure of Escherichia coli ribosomal protein
L7/L12: site specific cysteine cross-linking and attachment of fluorescent probes.

*Biochem. Cell Biol.* **73**, 949–958 (1995).

30. Mustafi, M. & Weisshaar, J. C. Simultaneous Binding of Multiple EF-Tu Copies to Translating Ribosomes in Live Escherichia coli. *MBio* **9**, (2018).

31. Diaconu, M. *et al.* Structural Basis for the Function of the Ribosomal L7/12 Stalk in Factor Binding and GTPase Activation. *Cell* **121**, 991–1004 (2005).

32. Garone, C. *et al.* Defective mitochondrial rRNA methyltransferase MRM2 causes MELAS-like clinical syndrome. *Hum. Mol. Genet.* **26**, 4257–4266 (2017).

33. Solomon, B. D. *et al.* De novo deletion of chromosome 20q13.33 in a patient with tracheo-esophageal fistula, cardiac defects and genitourinary anomalies implicates GTPBP5 as a candidate gene. *Birth Defects Res. Part A Clin. Mol. Teratol.* **91**, 862–865 (2011).

34. Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data preprocessing with Warp. *Nat. Methods* **16**, 1146–1152 (2019).

35. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

36. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 486–501 (2010).

37. Holm, L. DALI and the persistence of protein shape. *Protein Sci.* **29**, 128–140 (2020).

38. Arnold, K., Bordoli, L., Kopp, J. & Schwede, T. The SWISS-MODEL workspace: a web-based environment for protein structure homology modelling. *Bioinformatics* **22**, 195–201 (2006).

39. Kim, D. J., Jang, J. Y., Yoon, H.-J. & Suh, S. W. Crystal structure of YlqF, a circularly permuted GTPase: Implications for its GTPase activation in 50 S ribosomal subunit assembly. *Proteins Struct. Funct. Bioinforma.* **72**, 1363–1370 (2008).

40. Wahl, M. C., Bourenkov, G. P., Bartunik, H. D. & Huber, R. Flexibility, conformational diversity and two dimerization modes in complexes of ribosomal protein L12. *EMBO J.* **19**, 174–186 (2000).

41. Andersen, G. R., Thirup, S., Spremulli, L. L. & Nyborg, J. High resolution crystal structure of bovine mitochondrial EF-tu in complex with GDP. *J. Mol. Biol.* **297**, 421–436 (2000).

42. Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix. *Acta Crystallogr. Sect. D Struct. Biol.* **75**, 861–877 (2019).

43.  Chen, V. B. *et al.* MolProbity□: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 12–21 (2010).
44.  Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization
and analysis. *Protein Sci.* **27**, 14–25 (2018).
45.  Rosenthal, P. B. & Henderson, R. Optimal Determination of Particle Orientation,
Absolute Hand, and Contrast Loss in Single-particle Electron Cryomicroscopy. *J. Mol.*
*Biol.* **333**, 721–745 (2003).
46.  Baker, N. A., Sept, D., Joseph, S., Holst, M. J. & McCammon, J. A. Electrostatics of
nanosystems: Application to microtubules and the ribosome. *Proc. Natl. Acad. Sci.* **98**,
10037–10041 (2001).
47.  Greber, B. J. *et al.* The complete structure of the 55S mammalian mitochondrial
ribosome. *Science (80-. ).* **348**, 303–308 (2015).

**Acknowledgments:** All cryo-EM data used in this work were collected at the Karolinska
Institutet's 3D-EM facility. The SciLifeLab cryo-EM facility (used for grid preparation and
initial screening) is funded by the Knut and Alice Wallenberg, Family Erling Persson, and
Kempe foundations. We acknowledge the support of: Knut and Alice Wallenberg Foundation
(KAW 2018.0080) to B.M.H and J.R.; the Swedish Research Council (2018-3808) to B.M.H.;
Karolinska Institutet and the Max Planck Institute to J.R. J.R. is a Fellow of the Knut and
Alice Wallenberg Foundation (WAF 2017).

**Author contributions:** M.C. with A.K and J.R. help performed cell biology, biochemistry
and sample preparations; G.V.G. and B.M.H. performed cryo-EM data collection; G.V.G. and
B.M.H. processed data and determined structures; G.V.G. built and refined models with help
from A.K. and B.M.H.; M.C. and G.V.G. wrote the manuscript with input from all authors;
J.R. and B.M.H. oversaw the project and edited the manuscript.

**Competing interests:** Authors declare no competing interests.

**Fig. 1: Overview of the GTPBP5[KO] and the GTPBP5[IP] mt-LSU assembly intermediates and comparison with the mature mt-LSU. a,** The GTPBP5[KO] is bound by MTERF4, NSUN4, MRM2, MTG1 and the MALSU1 module. Mitoribosomal proteins and 16S mt-rRNA are shown in grey. Helix-X bound to MTERF4 is highlighted as well as H89. **b,** The interface of the GTPBP5[IP] mt-LSU intermediate associated with MTERF4, NSUN4, MRM2, MTG1, MALSU1:L0R8F8:mt-ACP complex, GTPBP5 and mtEF-Tu. Helix-X bound to MTERF4 is shown in light blue. **c,** Secondary structure of the mature mt-LSU 16S mt-rRNA. Differences in the rRNA fold

519    of the GTPBP5[KO] mt-LSU intermediate are shown in the zoomed-in views. Dashed lines indicate regions that are

520    not modelled. MRM2 methylation site (H92) is indicated in red. The six 16S mt-rRNA domains are shown in

521    different colours. **d**, Positioning of helix-X (H68-71) and helices H89 and H90 in GTPBP5[KO] mt-LSU (left),

522    GTPBP5[IP] mt-LSU (middle) and the mature mt-LSU (right) (PDB:6ZSG [15]). In the GTPBP5[KO] and the

523    GTPBP5[IP] mt-LSU structures MRM2 is present.

**Fig. 2: MTERF4-NSUN4 and MRM2 interaction with the mt-LSU assembly intermediates. a**, Comparison
of the MTERF4-NSUN4 complex bound to the GTPBP5[KO/IP] mt-LSU (orange and green respectively) with the
MTERF4-NSUN4 crystal structure (PDB: 4FP9 [18]) (yellow and brown, respectively), and of uL2m from the
GTPBP5[KO/IP] mt-LSU (pink) with uL2m from the mature mt-LSU (blue) (PDB: 3J7Y [14]). The uL2m C-terminus
is indicated in both structures. Helix-X is not shown. **b**, MTERF4-NSUN4 complex bound to helix-X. Zoom-in
panels show the interactions of MTERF4 with helix-X. **c,** Schematic representation of MRM2 domains (NTD -
light pink, SAM MTase domain - red). **d**, MRM2 interaction with the domain IV rRNA (nucleotides 2644-2652,
green) and the A-loop (grey). The MRM2 methylation site (U3039) as well as the catalytic triad of MRM2 (K59,

D154, K194) are highlighted as sticks. **e,** Zoomed-in views showing MRM2 interactions with the A-loop in

different orientations.

**Fig. 3: GTPBP5 contributes to the maturation of the PTC region. a,** Schematic representation of GTPBP5

domains (Obg-domain dark blue, G-domain light blue). **b,** Overview of GTPBP5 interactions with the 16S

rRNA. The Obg-domain (dark blue) contacts helices that are in the PTC region: P-loop, A-loop, H89, H90, H93.

Helices a-f of GTPBP5 Obg-domain are indicated. The SRL and the NSUN4 N-terminus are shown. Boxes 1-3

show the remodelling of the PTC in GTPBP5[IP] mt-LSU (in color) compared with GTPBP5[KO] mt-LSU (in white,

higher panel) and with the mature mt-LSU (in white, lower panel) (PDB: 6ZSG[15]). **c**, Comparison of MRM2

(red) and the A-loop (grey) conformations between GTPBP5[KO] mt-LSU (left) and GTPBP5[IP] mt-LSU (right).

The N-terminal helices (pink) of MRM2 could not be modelled in the GTPBP5[IP] mt-LSU. The GTPBP5 Obg-

domain is shown in dark blue. **d**, Comparison of the P-loop and H89 conformations between GTPBP5[IP] mt-LSU

(lower panel) and GTPBP5[KO] mt-LSU structures (higher panel).

**Fig. 4: Interaction of mtEF-Tu with the mt-LSU assembly intermediate and model of the final steps of mt-LSU biogenesis. a**, Schematic representation of mtEF-Tu domains. **b**, mtEF-Tu interaction with GTPBP5, the MALSU1 module and the bL12m C-terminal domain. mtEF-Tu G-domain, domain II and domain III and the SRL are indicated. The six copies of bL12m N-terminal domain and uL10m are also highlighted. The yellow

dashed line indicates a hypothetical connection between bL12m CTD and one of the six copies of bL12m NTD,

not visible in the structure. **c**, Representation of the mtEF-Tu interaction with GTPBP5 and MALSU1. The upper

zoomed-in panel features interactions between the GTPBP5 G-domain and the mtEF-Tu domain III. The green

dashed line indicates interactions to the RNA phosphate backbone. The lower zoomed-in panel shows the mtEF-

Tu switch I interaction with MALSU1. **d**, Final steps of the mt-LSU assembly. The dashed arrow indicates that

biogenesis factors are released in an unknown order.