RESEARCH

Response inhibition in premotor cortex corresponds to a complex reshuffle of the mesoscopic information network

Giampiero Bardella[1],**Valentina Giuffrida**[2],**Franco Giarrocco**[2],**Emiliano Brunamonti**[1],**Pierpaolo Pani**[1] and **Stefano Ferraina**[1]

[1]Department of Physiology and Pharmacology, Sapienza University of Rome, 00185, Rome, Italy

[2]PhD Program in Behavioral Neuroscience, Sapienza University of Rome, 00185, Rome, Italy

Keywords: information theory, motor control, inhibition, brain networks, complexity,behavior

ABSTRACT

Recent studies have explored functional and effective neural networks in animal models; however, the dynamics of information propagation among functional modules under cognitive control remain largely unknown. Here, we addressed the issue using Transfer Entropy and graph theory methods on mesoscopic neural activities recorded in the dorsal premotor cortex of rhesus monkeys. We focused our study on the decision time of a Stop-signal task, looking for patterns in the network configuration that could influence motor plan maturation when the Stop signal is provided. When comparing trials with successful inhibition to those with generated movement, the nodes of the network resulted organized into four clusters, hierarchically arranged, and distinctly involved in information transfer. Interestingly, the hierarchies and the strength of information transmission between clusters varied throughout the task, distinguishing between generated movements and canceled ones and corresponding to measurable levels of network complexity. Our results suggest a putative mechanism for motor inhibition in premotor cortex: a topological reshuffle of the information exchanged among ensembles of neurons.

AUTHOR SUMMARY

In this study, we investigated the dynamics of information transfer among functionally identified neural modules during cognitive motor control. Our focus was on mesoscopic neural activities in the dorsal premotor cortex of rhesus monkeys engaged in a Stop-signal task. Leveraging multivariate Transfer Entropy and graph theory, we uncovered insights on how behavioral control shapes the topology of information transmission in a local brain network. Task phases modulated the strength and hierarchy of information exchange between modules, revealing the nuanced interplay between neural populations during generated and canceled movements. Notably, during successful inhibition, the network displayed a distinctive configuration, unveiling a novel mechanism for motor inhibition in the premotor cortex: a topological reshuffle of information among neuronal ensembles.

INTRODUCTION

The brain is a complex system with billions of neurons interacting across various spatial and temporal scales, forming interconnected modules of different sizes. At the small spatial scale, modules correspond to single neurons; at the large scale, modules correspond to specialized brain areas formed by many neurons. At the mesoscale, modules are neural populations of variable sizes, sometimes described as functional columns. Throughout the recent decades, neuroscience has attempted to characterize brain computations by relating neural activities to behavior. Understanding how network modules interact at the different scales and how information is shared and processed among parts is critical at all levels of investigation. Historically, most approaches have relied on single unit activity to describe the neural correlates of behavior in discrete populations of neurons recorded in a single area (Caminiti, Johnson, Galli, Ferraina, & Burnod, 1991; Fetz, 1992; Georgopoulos, Kalaska, Caminiti, & Massey, 1982; Munoz & Wurtz, 1993; Riehle, Grün, Diesmann, & Aertsen, 1997; Vaadia, Kurata, & Wise, 1988) being able to characterize larger networks only occasionally(Ferraina, Paré, & Wurtz, 2002; Wurtz, Sommer, Paré, & Ferraina, 2001). Recent developments in microelectrode fabrication (Hong & Lieber, 2019) have made it possible to record simultaneously from multiple channels and then investigate the dynamics of massive numbers of neurons and sites (Shenoy, Sahani, & Churchland, 2013). Nonetheless, the majority of studies focused on single-unit activity (**SUA**) , and on the lower bands of the local field potentials

(**LFPs**), which, however, indicate more the average synaptic input to the investigated area than the local

spiking activity (for a review see (Herreras, 2016)). Comparatively, few studies have looked at signals

that sample the average spiking activity of the population of neurons in close proximity to the recording

electrode (multiunit activity; **MUA** (Mattia et al., 2013; Stark & Abeles, 2007; Supèr & Roelfsema, 2005;

Trautmann et al., 2019)). Such mesoscopic signals, more stable than SUA, can provide crucial

information for investigating the local dynamics that enable brain function control.

In the arm motor domain, studies have indicated that neurons in the dorsal premotor (PMd) cortex of

monkeys are engaged in motor control, as evidenced by modulation patterns of both SUA and

mesoscopic signals (LFPs and MUA) obtained both at the single and population level (Cisek & Kalaska,

2010; Milekovic, Truccolo, Grün, Riehle, & Brochier, 2015; Shenoy et al., 2013; Wise, Boussaoud,

Johnson, & Caminiti, 1997). Some studies have employed the Stop signal task (Logan & Cowan, 1984),

that requires actively inhibiting an upcoming movement upon presentation of a Stop signal, to investigate

how neuronal activity in this area supports both movement generation and cancellation. According to

these studies, neuronal activity in PMd is strongly related to both the generation and the suppression of

an arm movement at different scales, from single neurons (Marcos et al., 2013; Mirabella, Pani, &

Ferraina, 2011) to population of neurons(Pani et al., 2022) and local mesoscopic signals (Bardella, Pani,

Brunamonti, Giarrocco, & Ferraina, 2020; Mattia et al., 2013; Pani et al., 2014). One of the proposed

mechanisms suggests that arm movements are generated when neurons, following a Go signal,

accumulate evidence up to a threshold, and inhibited when the accumulation is disrupted or postponed by

a competing process triggered if a Stop signal is presented early enough after the Go signal. This view, in

which neurons work as stochastic accumulators, stemmed from the analogy with the neural results in the

oculomotor domain (Hanes, Patterson, & Schall, 1998; Paré & Hanes, 2003) and the original behavioral

model (Logan & Cowan, 1984) based on a competition between accumulating processes. Recently (Pani

et al., 2022), it has been advanced the hypothesis that such evolution can be depicted as a dynamic in the

state-space, in which movement generation and inhibition correspond to distinct neural trajectories

hopping within specific subspaces. While these approaches have been fruitful in giving a mechanistic (or

quasi- mechanistic) account of movement control, the issue can be tackled also from another perspective.

For example, an alternative hypothesis is that movement generation necessitates the functional

interactions of clusters of neurons to build networks that efficiently handle information locally and send

information to other brain areas and/or the spinal cord (see Bassett & Sporns review (Bassett & Sporns,

2017)). Thus, one could also ask whether motor decisions are characterized by specific routings of

information across mesoscale neuronal "components". To answer, we studied the PMd's network

dynamics using the MUA as a mesoscopic measure of spiking activity and implementing a combined

information-theory and graph-theory approach to describe how the collective interaction of recorded

modules is linked to motor decision-making. We demonstrated that in the behavioral epoch when

decisions are made, mesoscopic modules may be classified into four distinct clusters that interact with

one another and are organized hierarchically. Interestingly, the hierarchical level of some of them, as well

as their strength of interaction and the overall communication rules, exhibited differences when

movements are normally executed and when the animal successfully canceled the planned movement.

Our results indicate that neurons in PMd have diverse opportunities to contribute to motor control at

different spatial scales. In general, these findings support the idea of complexity being a core feature of

the brain's ability in managing information at multiple levels of neural interaction.

MATERIALS AND METHODS

Subjects

The study used two male rhesus macaque monkeys (Macaca mulatta; Monkeys P and C), weighing 9 and

9.5 kg, respectively. The animals' care, housing, surgical operations, and experiments were all in

accordance with European (Directive 86/609/ECC and 2010/63/UE) and Italian (D.L. 116/92 and D.L.

26/2014) laws and were approved by the Italian Ministry of Health. The monkeys were housed in pairs

and given cage enrichment. During the recording days, they were fed regular primate chow supplemented

with nuts and fresh fruits as needed, and they were given a daily water supply.

Apparatus and task

The monkeys were placed in a soundproof, dimly lighted chamber in front of an isoluminant black

background with a brightness intensity of less than 0.1 cd/m2. They were shown stimuli on a 17-inch

touchscreen panel with an LCD display with an 800×600 pixel resolution. CORTEX, a non-commercial

software program available at http://www.nimh.gov.it, was used to control the presentation of stimuli and

recording of behavioral responses. Figure 1 depicts the overall task, which our group first described as a

stop-signal reaching task (Mirabella et al., 2011). The central target (CT), a red circle with a diameter of

1.9 cm, was displayed at the start of each trial. The CT had to be reached by the monkeys to initiate the

trial. A peripheral target (PT) randomly appeared on the left or right side of the screen after a variable

holding duration of 400–900 ms, in steps of 100 ms. At the end of the holding period the CT vanished,

signaling a Go signal. In the Go trials, the monkeys had to reach and hold on the PT for a range of

400–800 ms, in steps of 100 ms, to get a juice reward. The time between the presentation of the Go signal

and the beginning of the hand movement was used to compute the reaction time (RT). The sequence of

events in Stop trials was the same as in Go trials until the Go signal. The CT reappeared as the Stop

signal after a variable delay known as the Stop signal delay (SSD), and the monkeys had to hold the CT

until the end of the trial, which was between 800-1000 ms, to get the reward for a correct Stop trial. If the

monkey removed its hand before the trial ended, it was considered an incorrect response for a Stop trial.

For both correct Stop and correct Go trials, the same amount of juice was provided. The inter-trial

interval was set at 800 milliseconds. Stop trials accounted for approximately 25% of all Monkey P trials

and 32% of Monkey C trials. White rings around the central target provided input to the animals for the

touch. A staircase tracking approach was utilized to determine the duration of the SSDs, to obtain a

success rate of roughly 50% in Stop trials. The SSD rose by 100 ms in the succeeding Stop trial if the

monkey successfully withheld the movement, but it reduced by 100 ms if the monkey failed to cancel the

response.

Behavioral considerations

The stop-signal reaching task allows for the calculation of the Stop signal reaction time (SSRT), which is

recognised as a metric of efficiency in movement suppression. The race model(Logan & Cowan, 1984) is

used to describe the behavior in the Stop trials, which involves two stochastic processes racing towards a

threshold: the GO process, which is triggered by the Go signal and whose duration is represented by the

RT, and the STOP process, whose duration must be derived. If the GO process wins the race against the

STOP process, the movement is generated (resulting in wrong Stop trials). If instead the STOP process

prevails, the movement is restrained (resulting in correct Stop trials). We may estimate the SSRT using

the race model by considering the duration of the GO process, the probability of response, and the

stop-signal delays (SSDs). To estimate the SSRT, we used the integration approach, which has been

demonstrated to be the most trustworthy (Band, van der Molen, & Logan, 2003). This technique takes

into account the probability of response [p(response)] for a given SSD in Stop trials. The longer the SSD,

the higher the p(response), indicating a higher likelihood of a wrong Stop trial. By integrating

119 **Figure 1.** **Sequence of Behavioral events characterizing the task**. Go and Stop trials were randomly intermixed during each session. The epoch (**T**) of

120 analysis is shown as a gray bar for both trials. White circles are feedbacks for the touch provided to the animals. **RT**, reaction time; **SSD**, Stop signal delay;

121 **SSRT**, Stop signal reaction time; **Go**: time of Go signal appearance; **Stop**: Stop Signal appearance; **M_ON**: movement onset.

p(response) over the distribution of Go trials, we can derive the nth value Go RT. Subtraction of the SSD

from the nth Go RT yields the SSRT. To visualize the procedure, imagine an SSD where p (response) is

exactly 0.5. For such SSD, the Go and Stop process would have the same probability of winning the race,

meaning that the Stop and Go processes will both reach the threshold at the same time (from a

probabilistic point of view). The completion times of the Stop and Go processes are, respectively, the end

of the SSRT and the end of the RT (i.e., the beginning of movement). In this study, we compared the

evolution of brain activity during movement preparation in the Go trials to the inhibitory effect of the

Stop signal in correct Stop trials. To make this comparison effective, we considered an epoch before

movement generation in Go trials and an epoch starting before the Stop signal and ending at the

conclusion of the Stop process (SSRT duration) in correct Stop trials (see below). When adding wrong

Stop trials in the analysis, we used the same epoch duration and we aligned neural activities to the

beginning of movement. Because different SSDs were used in separate recording sessions, and because

RT distributions also varied, resulting in varying SSRTs across sessions, we chose an epoch that could

contain all estimated SSRTs in the various sessions (see Table S1). We decided for a duration of 400 ms

for the epoch of analysis (Fig.1, epoch T, gray bar). In Stop trials, the epoch T did not include the Go

signal when short SSDs were employed.

Extraction and processing of neuronal data

In the left PMd, a multielectrode array (Blackrock Microsystems, Salt Lake City) with 96 electrodes

spaced at 0.4 mm was surgically implanted. After opening the dura, the arcuate sulcus and precentral

dimple were employed as reference (for electrode location see Figure S1 in (Pani et al., 2022)). We

recorded the raw signal using a Tucker Davis Technologies (Alachua, FL) system, sampling at 24.4 kHz.

As a measure of neuronal activity at the population level, local spectral estimated MUA was extracted

offline from the raw signal. This was accomplished by computing the time-varying power spectra $P(\omega, t)$

from the signal's short-time Fourier transform in 5-ms sliding windows, as described in Mattia et al.

(Mattia et al., 2013). Relative spectra $R(\omega, t)$ were obtained normalizing $P(\omega, t)$ by their average $P_{ref}(\omega)$

across a fixed window (30 minutes) for the entire recording. Thus, the average $R(\omega, t)$ across the $\omega/2\pi$

band $[0.2, 1.5]$ kHz represents our mesoscopic measure of spiking activity. As explained in Mattia et al.

(Mattia et al., 2013), such estimation relies on two hypotheses. The first is that high ω components of the

raw signal result from the convolution of firing rates $\nu(t)$ of neurons that are close to the electrode tip

with a stereotypical single-unit waveform. The Fourier transform $K(\omega)$ of such an unknown waveform is

canceled out in $R(\omega, t)$, which is therefore a good approximation of the ratio of firing rate spectra

$|\nu(\omega,t)|^2 / |\nu_{ref}(\omega)|^2$. The second is that high ω power $|\nu(\omega,t)|^2$ is proportional to the firing rate $\nu(t)$ itself

(Mattia & Del Giudice, 2002). This entails that our MUA is proportional to $\nu(t)$, i.e. to the firing rate of

the sampled cortical volume. The aggregate spiking activity of neuronal populations in the vicinity of the

electrode, without spike detection, was estimated with similar methods in previous work (Choi, Koenig,

Jia, & Thakor, 2010; Mattia et al., 2013; Sharma et al., 2015; Stark & Abeles, 2007). As a last step,

logarithmically scaled MUAs were smoothed by a moving average (40 ms window, 5ms step which led to

a time series of continuous values with a dt=1 ms).

Quantifying information dynamics with Transfer Entropy

[174] Based on the MUA obtained from each electrode site, we used multivariate **Transfer Entropy** (TE)

[175] (Komárek, Hrnčíř, & Štěrbová, 2001; Schreiber, 2000) to infer patterns of local information dynamics in

[176] the PMd, the directed functional relationship between different neural populations (modules) and the

[177] collective state of the information flow through the network at the time of the motor decision. Recent and

[178] thriving literature (Bossomaier, Barnett, Harré, & Lizier, 2016; Goetze & Lai, 2019; Ito et al., 2011;

[179] Michael Wibral, 2014; Newman, Varley, Parakkattu, Sherrill, & Beggs, 2022; Ramos, Macau, Lopes, &

[180] Machado, 2017; Vicente, Wibral, Lindner, & Pipa, 2011; Wibral et al., 2011; Wollstadt,

[181] Martínez-Zarzuela, Vicente, Díaz-Pernas, & Wibral, 2014; Xiong, Faes, & Ivanov, 2017) has shown that

[182] TE, particularly in its multivariate formulation, is more accurate than other measures (Novelli & Lizier,

[183] 2021; Ursino, Ricci, & Magosso, 2020), also capturing the effect of nonlinear interactions in the system

[184] dynamics without assuming any generative model. Given an ensemble of M time series, the multivariate

[185] information transfer from a time series X to a series Y, conditioned to the remaining $Z_{k=1,..,M-2}$ series,

[186] can be quantified considering the present values of Y and the past values of both X and the Z through

[187] (Faes, Kugiumtzis, Nollo, Jurysta, & Marinazzo, 2015; Montalto, Faes, & Marinazzo, 2014; Xiong et al.,
2017):

$$TE_{X \to Y \mid Z} = H(Y_n \mid Y_n^-, Z_n^-) - H(Y_n \mid Y_n^-, X_n^-, Z_n^-) \,, \tag{1}$$

[188] where Y_n is the vector that represent the present state n of Y, $X_n^- = [X_{n-1}, X_{n-2}, ...]$,

[189] $Y_n^- = [Y_{n-1}, Y_{n-2}, ...]$ and $Z_n^- = [Z_{n-1}, Z_{n-2}, ...]$ are the vectors that represent the past values of X, Y

[190] and Z respectively. The vertical bar stands for conditional probability, e.g. $H(Y_n \mid Y_n^-, Z_n^-)$ is the

[191] entropy of the present state of Y conditioned to the knowledge of the past of Y and to the past of the

remaining Z. H is the Shannon entropy (Shannon, 1948), which in the case of Y is given by:

$$H(Y_n) = - \sum_n P(Y_n) \, log P(Y_n) \,, \tag{2}$$

[192] where P indicates the probability density associated with the vector Y_n. Using equation 2 expression 1

becomes:

$$TE_{X \to Y \mid Z} = \sum_n P(Y_n, Y_n^-, X_n^-, Z_n^-) \, log \frac{P(Y_n \mid Y_n^-, X_n^-, Z_n^-)}{P(Y_n \mid Y_n^-, Z_n^-)}. \tag{3}$$

[193] TE is ≥ 0 and increases if the history of X gives more information about the present of Y than the past

[194] of Y or any other series contained in Z. Because the past of X is used to predict the present of Y, TE is

[195] not symmetric (i.e. $TE_{X \to Y \mid Z} \neq TE_{Y \to X \mid Z}$) and defines an information transfer direction, which

[196] corresponds to a driver-to-target causal relationship. Approximating the vectors that represent the time

series' history, a process known as *embedding*, is critical when estimating TE. The optimal embedding should include the X_n^-, Y_n^- and Z_n^- components that are most informative in describing Y_n (Montalto et al., 2014). We used a non-uniform embedding scheme (Faes, Marinazzo, Montalto, & Nollo, 2014) and computed H with kernel estimators (for a detailed description of embedding methods and estimators for computing H, please refer to the works of Faes and colleagues and the references cited therein (Faes et al., 2015, 2014; Faes, Nollo, & Porta, 2011; Xiong et al., 2017)). Non-uniform embedding is a more reliable alternative to uniform embedding, which would choose the previous values X_n^-, Y_n^- and Z_n^- a *priori* and separately for each time series (Montalto et al., 2014; Xiong et al., 2017). Instead, non-uniform embedding selects past components based on maximum relevance and minimum redundancy requirements, gradually removing non-informative terms. Maximum relevance means the most significant predicted information components. If the non-uniform embedding procedure selects at least one component from the past, the resulting $TE_{X \to Y \mid Z}$ is > 0 and statistically significant. Otherwise, if no component from the past is selected, $TE_{X \to Y \mid Z}$ is deemed to be non-significant and equal to 0 (Montalto et al., 2014). In the selection of the past values, three parameters are important. The maximum number of prior values, or maximum lag l; the embedding lag τ, that gives the number of steps to take into the past and the offset u from where to start to count the steps (e.g. from the first past value, second etc..). In non-uniform embedding methods, the parameters are initialized at the start of the trimming procedure and then sequentially tested for significance. To avoid any bias due to an arbitrary choice of the past values, we initially fixed a "relaxed" uniform embedding vector containing the past up to *l*=50 ms with *tau* and *u* set to 1. Then, we let the non-uniform embedding procedure prune the non informative terms and find the optimal embedding. The statistical significance of past values was progressively assessed by cycling through the past components (up to *l*) and comparing them with a null distribution built from empirical data using a time shifting randomization (Faes, Nollo, & Chon, 2008; Faes et al., 2011; Montalto et al., 2014; Quian Quiroga, Kraskov, Kreuz, & Grassberger, 2002; Varley, Sporns, Schaffelhofer, Scherberger, & Dann, 2023). The components of the past of X, Y, and Z were retained if significant at $\alpha < 0.01$ after 300 shuffles. As expected, the procedure selected only a recent past (no older than 10ms for each n of equation 2), aligning with findings from similar studies (Antonello et al., 2022; Shimono & Beggs, 2015; N. M. Timme et al., 2016; Varley et al., 2023; Wollstadt et al., 2014). This is consistent with prior research, which indicates that neuronal populations in the cortex typically establish causal interactions within a time range of 1 to 20 ms (Barthó et al., 2004; Mason,

Nicoli, & Stratford, 1991; Swadlow, 1994). To estimate the probability densities necessary for the computation of H, we employed the Heaviside Kernel estimator. This estimator reconstructs the P at each data point by assigning weights based on the distance r from the reference point x_n to any other point in the time series. Such local exploration of the state space within the neighborhood points of the time series, makes it more robust to inaccurate estimations compared to other approaches (Xiong et al., 2017). r is usually chosen to be a fraction of the time series variance (Gómez et al., 2014; Richman & Moorman, 2000; Xiong et al., 2017) so that the dependencies of the estimated entropies from the amplitude of the process are removed. To measure the distance we used the Chebyshev distance (or maximum norm) which is the most commonly used metric (Xiong et al., 2017). The Heaviside Kernel with the Chebyshev distance is equivalent to the sample entropy (Richman & Moorman, 2000; Xiong et al., 2017) which is an established measure to infer complexity of dynamical systems from time series data (Grassberger & Procaccia, 1983; Pincus, 1991) that has been recognized to reduce the estimation bias of conditional entropies. We built single-trial information transfer networks for each behavioral condition of each recording session by customizing some functions of the Matlab MuTE toolbox (Montalto et al., 2014).

Since the application of information theory techniques in neuroscience is still in its early stages, it is worth briefly examining their use on invasive neurophysiological data. Gomez-Herrero (Gómez-Herrero et al., 2015), Bossomaier (Bossomaier et al., 2016), Wibral (Wibral et al., 2011), Newman (Newman et al., 2022) and colleagues have published technical discussions on this topic. Timme and Lapish landmark review (N. M. Timme & Lapish, 2018) is recommended for a full treatment of the critical issues in the field. As summarized in the latter, the main assumption made by experimenters concerns the stationarity of the neural processes under examination. Stationarity of a process means that the probability density function that generates the observable variables does not vary over time, and each observation contributes to the same probability distribution. In practice, this translates into viewing the experimental trials as multiple realizations of the same underlying mechanism. In other words, assuming stationarity implies that each observed realization helps to define the same underlying probability distribution, which finally becomes stable over time, i.e., the lifetime of the experiment (Wollstadt et al., 2014). In a behavioral neurophysiology experiment, the observations correspond to neural activity values, the realizations to repeated trials, and the probability distribution to the probability distribution characteristic of the investigated neural process that is supposed to generate the observed behavior. It is critical to remember that in any experimental paradigm, this distribution is not known a *a priori* and is almost never

accessible, in contrast to any theoretical study in which time series are generated based on many repetitions of processes whose analytical details are known. Furthermore, stationarity does not imply a flat variability across realizations, and some variance is expected, reflecting the natural fluctuation of the underlying (unknown) probability distribution (Wollstadt et al., 2014). For these reasons, when an experiment has a multi-trial structure, stationarity over realizations is assumed by default, provided that no slow drifts or trends affect the signal, being the latter one of the non-stationarity factors that most impinge on a correct estimation of TE (Xiong et al., 2017). These could arise during training or in an experiment that encourages learning, where the probability distribution of both the brain signal and the behavioral parameters changes over time by definition (Wollstadt et al., 2014).

As a first step in using TE on our data, we analyzed the single-trials MUA profiles of each module. Our estimate of MUA is baseline corrected so any potentially distorting slow trends are eliminated. In addition, to remove noisy outliers from our data, we excluded from the analysis the trials for which the MUAs showed peaks with an amplitude that exceeded the average of the activity by 2 standard deviations in any of the time bins of the epoch of interest and for over 80% of the channels (Bardella et al., 2020) (which accounted for less than 2% of the total number of trials). This ensures that artifacts caused by non-physiological peaks and/or segments of ulta-rapid oscillations of high variance are excluded from the analysis. These non-stationary artifacts are among those that have the most impact on trustworthy TE estimation since they could influence the probability distributions in equation 3 in such a way that outliers would dominate most of the signal variance, significantly biasing the estimation of the temporal link between the current and past states of the time series (Xiong et al., 2017).

Graph theoretical analysis

We obtained a single-trial TE matrix in the epoch T for each behavioral condition (Go, correct Stop trials and wrong Stop trials) for both monkeys. Assuming that the nodes are single modules (discrete neuronal populations producing MUAs) and the weighted edges are the TE values, we interpreted the TE matrix as the adjacency matrix of a directed weighted network. We disregarded the main diagonal of the matrix thus excluding self-loops from the networks. Then, we aggregated the trials of the same condition for each recording session to get the appropriate trial-averaged matrices on which we applied **percolation** (Bardella, Bifone, Gabrielli, Gozzi, & Squartini, 2016; Bardella et al., 2020; Bordier, Nicolini, & Bifone, 2017; Bordier, Nicolini, Forcellini, & Bifone, 2018; Callaway, Newman, Strogatz, & Watts, 2000;

Del Ferraro et al., 2018; Gallos, Makse, & Sigman, 2012; Mastrandrea et al., 2017, 2021; Nicolini & Bifone, 2016; Nicolini, Bordier, & Bifone, 2017; Vlasov & Bifone, 2017). During percolation, the iterative pruning of links based on their weights is used to monitor the stability of network topology. Percolation was used in this work for two reasons. First, to remove weaker links that were influenced by the statistical noise of trial averaging. Indeed, false positives TE values may emerge from the constraints of limited sample size, even when no direct link exists between the variables (Bossomaier et al., 2016; Novelli & Lizier, 2021). Our pipeline included percolation as a topology-based data-driven method for an additional fine-grained cleaning. This also remove the hindrance of optimal thresholding, a common problem when dealing with connectivity matrices (Bardella et al., 2016, 2020; Nicolini & Bifone, 2016), without the need for computationally intractable full error-correction inference of each link at the whole network level. Second, percolation was used to analyze how hierarchical information flow between nodes shapes the complexity of network topology. Percolation analysis produces a curve that plots the size of the largest **connected component** of the network as a function of link weights. Since the networks we studied are directed, the definition of connectedness for directed graphs must be considered. Without delving into much detail, two definitions for a giant component of a directed graph can be given: giant strongly connected component (GSCC) and giant weakly connected component (GWCC) (Graf, 2016; Squires et al., 2013). Very briefly, a GSCC is a subgraph in which every vertex is reachable from every other vertex following the direction of the links. Instead, in a GWCC this is not possible. In other words, in a GWCC a path between every two vertices exists only in the underlying undirected graph. The GSCC thus reveals the network configurations in which information is exchanged cyclically through paths connecting all nodes. We have previously shown that this configuration is important especially during Stop trials, when the PMd network is expected to express quantitative contributions to the inhibition process (Bardella et al., 2020). Hence, we used the GSCC. For simplicity, in the Results we will refer to the threshold at which the fragmentation of the GSCC begins (formally, the percolation threshold) as the "descent point" of the curves. Such a threshold achieves the best balance between the removal of spurious links and the consequent loss of information potentially encoded in those links (Bordier et al., 2017; Nicolini et al., 2017). The power of percolation lies in displaying the stepped pattern of the curve, which suggests a transitory stable network configuration composed of hierarchical connections between nodes. A steeper curve with no or few leaps, on the other hand, indicates a single incipient component that dominates the network and fragments swiftly in fewer or a single step. This process naturally

discloses simultaneously both network robustness and its hierarchical self-organization. Notably, the percolation threshold identifies the level of complexity of the network: a higher threshold means a stronger component; a stronger component means that more links need to be removed to fragment it; more links implies more paths between nodes and thus higher complexity. Thus, we could estimate the complexity of information flow-paths within the network (under different behavioral conditions) by evaluating fragmentation at higher thresholds (see (Bardella et al., 2016, 2020; Del Ferraro et al., 2018; Gallos et al., 2012) for additional details on the percolation analysis applied to brain networks). On each trial-averaged matrix, we measured the Vertex Degree (VD), which is the number of links to each node i:

$$VD_i = \sum_{j=1}^{N} a_{ij} \, , \tag{4}$$

where a_{ij} is the generic entry of the adjacency matrix and N is the number of nodes. In directed networks one can distinguish between in-degree (the number on inward links) and out-degree (the number of outward links). For each behavioral condition of each recording session, we estimated the in/out-degree probability for both animals (see supplementary Figure S1). If the variance of the degree distribution is significantly larger than its mean, tails in the distribution arise and network hubs are found. Hubs are thus nodes that significantly exceed the average degree of the network (here we used mean $\pm$ 2 standard deviations of the out-degree distributions (Albert & Barabási, 2002; Barabási, 2013; Barabási & Albert, 1999)). After hubs were identified, we sorted nodes into clusters to better understand their contribution to information transfer. We labeled the nodes based on the two possible outputs of the motor behaviors studied in the task: movement generation (correct Go trials) and movement inhibition (correct Stop trials). To summarize the total quantity of information exchanged between clusters we defined an interaction measure I:

$$I_{ij}^{(m,n)} = \rho \sum_{i}^{M} \sum_{j}^{M} \sum_{(m,n)} TE_{C_i^m \to C_j^n} \, , \tag{5}$$

where C is the module's cluster, M is the number of clusters and m and n run over the all possible combinations of nodes within each cluster. $\rho = \frac{1}{dim\,(C_i)}$ is a normalization factor that accounts for the heterogeneous number of nodes within each of the clusters. Thus, our I_{ij} is equivalent to a node strength computed on a renormalized graph formed by M clusters. The strength of a node in a weighted graph is the total of the weights of the links that connect to the node. The values and directions of I reflect the cluster's location in the network communication hierarchy. We enclosed all of the interactions defined by

342 the empirical TE matrix in a MxM matrix, with each node representing a functional cluster. I was

343 calculated for each recording session and behavioral condition, and the sessions-averaged element (mean

344 $\pm$ SEM) $\overline{I}_{ij}$ was derived (see Figure 7 and Tables S2 and S3).

A null model

We built a null network model to assess the statistical significance of the results beyond what was

attained by the randomization approach used in the TE computations. As noted in a recent work (Cimini

et al., 2019), selecting a suitable null model remains a thorny issue in network science. In this study, we

drew our conclusions exclusively in a data-driven fashion, with no other constraints. Thus, the simplest

null hypothesis is that the results are caused by non-trivial temporal dependencies among neural activities

and cannot be explained by the distribution of MUA values and the length of the time series alone. To test

this, given a recording session and a behavioral condition and trial, we generated a synthetic time series

for each recording site (node of our TE network) by randomly permuting the corresponding MUAs. In

this way each surrogate time series has the same basic statistical properties of the original one (mean,

variance and histogram distribution), while relationships across the series of different nodes are destroyed

(Faes, Pinna, Porta, Maestri, & Nollo, 2004). For each surrogate single-trial graph we then preserved the

link density of the corresponding empirical matrix and computed the trial-averaged null TE network.

Repeating the procedure 500 times for each condition and recording session, we obtained an

ensemble-averaged TE network with which we compared our empirical graph metrics (see Figure S2).

We chose this null model because it is the least restrictive and no assumptions are made about the

connection patterns and weight distributions of the synthetic networks. This was in line with our goal: to

test that the information patterns in the network could not be detected solely by constraining the activity

distribution at the node level.

RESULTS

Table S1 shows behavioral data from the 21 recording sessions used in the study, 12 from Monkey P and

9 from Monkey C. In order to compute the SSRT, we ensured that the race model's fundamental premise

was followed in all sessions. To be more specific, the Go process in the Stop trials must be the same as

the Go process in the Go trials to compute the SSRT. This assumption is usually satisfied by showing that

RTs in wrong Stop trials are shorter than RTs in Go trials, as shown in Table S1. This indicates that all

trials are drawn from the same distribution of RTs. Across all sessions, the average RT for Monkey P was

656 ms (SD = 129 ms) and 632 ms (SD = 119 ms) for Monkey C. The average SSRT was 217 ms (SD =

34 ms) for Monkey P and 213 ms (SD = 26 ms) for Monkey C.

Neural activity significantly contributes to motor control at the single-site level

Fig 2 illustrates the time course of the average MUA obtained from each cortical site separately during

Go trials (dark green traces), correct Stop trials (red traces), and wrong Stop trials (dark blue traces)

during the time (epoch T in Figure 1) preceding the movement onset (in Go trials and wrong Stop trials)

or the end of the SSRT (in correct Stop trials) for one example session of Monkey P (S6 of Table S1).

The activity in the wrong Stop trials appears to be indistinguishable from the activity in the Go trials for

all sites. In contrast, the presentation of the Stop signal (red vertical line) and the effective cancellation of

movement produce a distinct pattern in many of the locations, with the spiking activity modulated

(increasing or decreasing) compared to the activity during movement generation. In this specific session

we found a significant difference between the Go and correct Stop conditions in 78% of the recording

sites. Conversely, there was no significant difference in spiking activity between the Go and wrong Stop

trials at any of the sites. A similar trend was found in all recording sessions (Table S4), confirming

previous findings on the role of PMd in inhibitory motor control and the relevance of the epoch of

analysis in motor decisions during Stop trials (Bardella et al., 2020; Battaglia-Mayer et al., 2014;

Giarrocco et al., 2021; Mattia, Ferraina, & Del Giudice, 2010; Mirabella et al., 2011; Pani et al., 2014,

2022, 2018; Ramawat et al., 2022). This result, however, does not provide information on how the nodes

in the network interacted in the different behavioral conditions.

Fig.3 shows the TE matrix for each trial type for the data provided in Figure 2. The dense horizontal

stripes that change between Go, correct, and wrong Stop trials suggest that, depending on the behavioral

situation, a few of the sites might influence the activity of many others during the epoch where the motor

choice is taken. In this picture, any recording location is interpreted as a network node, and any

significant TE value as a link. To describe network topology we first calculated the vertex degree (VD)

for the average networks, thresholded via percolation analysis after removing the noisiest links (Bardella

et al., 2016, 2020; Bordier et al., 2017; Nicolini & Bifone, 2016; Nicolini et al., 2017) (see Section). A

high VD indicates that some modules are functionally related to many others and serve as hubs. Hubs

Figure 2. Neuronal activity modulation for different trial types. Neuronal MUAs (mean ± SEM) in the epoch T for the three behavioral conditions for all recording sites (modules) of S6 of Monkey P. Dark green traces show the average activity during Go trials aligned to the movement onset (gray vertical line). Dark blue traces show the average activity during wrong Stop trials aligned to the movement onset (gray vertical line). Red traces show the average activity during correct Stop trials aligned to the Stop signal presentation (light red vertical line) and plotted until the end of the SSRT. For each plot the Y axes are units of MUA while the X axes mark the time during the epoch T; ticks, every 100 ms, are indicated for panels in the top row only. Numbers above each panel indicate the recording site. The actual spatial mapping of the recording sites is not preserved and are here shown in cardinal order from 1 to 96.

only appeared in the out-degree (VD_{out}) distributions (see Figure S1 and S2). In each session, we labeled the nodes based on the behavioral condition in which they were hubs. Were classified as **C_1** or **C_2**, the nodes that were hubs only in trials where the movement was generated or successfully canceled, respectively. The **C_3**, included all the nodes that were hubs in both Go and correct Stop trials. The **C_4** included the nodes that were never hubs. Interestingly, no new hubs were found when movements were incorrectly inhibited after a Stop signal (wrong Stop trials). This implies that the same neuronal clusters

Figure 3. **Average** TE **matrices** of Go, wrong and correct Stop trials for data in Fig.2. Each entry of the matrix is color-coded according to the corresponding

TE value averaged over trials. Statistically significant driver $\rightarrow$ target relationships correspond to TE values > 0 for the pair of modules considered. Here, for

graphical purposes only, TE is normalized to the maximum across behavioral conditions so to obtain $TE \in [0, 1]$ (color bar). Axes labels indicate recording

sites.

(groups of nodes) recruited in Go and correct Stop trials are also involved when errors occur in Stop

trials. The percentage of nodes belonging to each cluster throughout all sessions is shown in Table S5.

One might wonder if there is any internal hierarchy between the clusters, specifically whether the

VD_{out} of the nodes in a cluster changes depending on the behavioral conditions. To answer, we kept the

cluster label fixed and studied how the VD_{out} of the nodes in each cluster differed across trial types. **C_3**

consistently exhibited the highest VD_{out} values across all three behavioral conditions across all sessions

(see Table S6), indicating that they are the primary hubs for the spreading of information

(Kruskal-Wallis, $p < 0.01$, Bonferroni corrected for multiple comparisons). Thus, **C_3** are at the highest

hierarchical level in the network because they regulate information transfer regardless of whether the

decision is to move or stop. On the other hand, **C_1** and **C_2** have distinct levels of engagement, with

lower VD_{out} values in Go and correct Stop trials, respectively, shifting their role based on the type of

motor choice being processed. In Go trials, the **C_2** had lower VD_{out} values than the **C_1**, whereas in

correct Stop trials, the **C_1** had lower VD_{out} values than the **C_2**. Interestingly, the **C_1** had larger VD_{out}

values during wrong Stop trials than during correct Stop trials, whereas the **C_2** had the opposite

(Kruskal-Wallis, $p < 0.01$ Bonferroni corrected for multiple comparisons). This reveals that during

wrong Stop trials, the neuronal processing of information is midway between Go and correct Stop trials,

but more comparable to Go trials. The presence of four distinct clusters with varying VD_{out} shows that

some nodes serve as driving hubs only in relation to a specific behavioral outcome. Such heterogeneity

implies that the patterns of information transmission are fine-grained and are adjusted according to

whether the planned movement is executed or withheld. The topology of the PMd network for data in

Fig.3 is depicted in Fig.4, highlighting the **C_1** and **C_2** hubs, that change between conditions (for the

networks including also the **C_3** see Figure S4; see Figure S5 for examples from other sessions). The

information network has a centralized organization in Go trials (and to a lesser extent in wrong Stop

trials), corroborating our previous study that showed how such functional conditions facilitate movement

generation Bardella et al. (2020). Fig.4 (and Supplementary Figures S3 and S4) show how the **C_1** act as

hubs during Go and wrong Stop trials, whereas the **C_2** are hubs during correct Stop trials only. The

optimal centralized transmission present during Go trials is disrupted during correct Stop trials, where the

links arrangement is more complex and compounded, with more intersections and less directness

compared to Go trials. This suggests that the network retains its core feature (the existence of hubs)

across trials but varies in the way they are overall deployed based on the required behavioral control.

449 ***The PMd network becomes more complex in Stop trials***

To better understand how the hierarchical relationships between the clusters relates to the overall network

organization, we analyzed the VD_{out} as a function of TE. This involved tracking the changes in VD_{out}

of each cluster as the percolation process progressed. In this way, the resilience of the information carried

by each cluster is compared to the information spread by the others. We expect that a cluster that is

functionally relevant to a specific behavioral condition should possess not only a large number of

connections (a high VD_{out} value) but also a substantial number of these connections should exhibit high

and robust information content (reflected in high TE values). Results are shown in Fig. 5 as the average

of all recorded sessions of each animal. The analysis confirmed the **C_3** as the highest hierarchical level

cluster in all three behavioral conditions. The highest values of VD_{out} as a function of the amount of

information processed, indicate that it had the most links with high and robust information content. The

C_1 and **C_2**, on the other hand, were placed between the **C_3** and the **C_4**, and their hierarchical level

changed depending on the trial type. Thus, the network's global structure is similar in both wrong and

correct Stop trials. Inspecting the percolation curves we could infer the level of network complexity

under the three behavioral settings (see Section). Figure 6 shows that the percolation curves of the Go

Figure 4. Information network of Go, wrong, and correct Stop trials. Data from the same recording session of Fig.2 and Fig.3. Each link is a significant TE

value (i.e., a non-zero entry in the TE matrix). The number of outgoing connections of each module reflects its VD_{out}: the number of modules on which

it acts as a driver. Each link is color-coded according to the cluster to which its source node belongs. Information-spreading hubs are observable. Here, for

illustrative purposes and to better highlight the differences between behavioral conditions, the links of the C_3 nodes are removed and for each network only

the 20% of the strongest links are shown (for the complete version including the C_3 see Figure S4). Recording sites 1 and 96 are marked in the Go trials graph.

The coordinates of each node on the circle are preserved across behavioral conditions.

trials evolve differently from those of the Stop trials (whether correct or wrong) and the null model for all

sessions. The curves for the Go trials begin their descent at lower TE values and have a more stepped

Figure 5. Clusters robustness analysis. Robustness is inferred by monitoring the evolution of VD_{out} (y axis) as a function of the information processed (TE, x axis). Panels show the robustness curves averaged (mean $\pm$ SEM) over recording sessions compared between clusters and across behavioral conditions. Curves are color coded accordingly to the corresponding cluster as in Fig.4.

.

pattern (normalized among conditions). Therefore, the Go trials have fewer paths belonging to the giant component itself, with most links with high information content heading outwards to the rest of the network. This is a symptom of an ideal network setup with more direct and organized communication than in the Stop trials. One important implication is that in the Go trials the number of redundant links is very low. In other words, there are just a few links that deviate from the main common structure. During Go trials most high-weight links are used for direct communication between hubs and the rest of the network, lowering the level of complexity compared to Stop trials. Conversely, in Stop trials, a complicated rearrangement happens, with denser communications and high-content information traveling more within the large component than outwards. The proximity of the Stop trial curves to the null model (NM) reveals that, although they differ from a purely random situation, they still adhere to some of its communication rules. Indeed, in the NM, which is inherently complex, one would anticipate a configuration where links are randomly arranged regardless of their weights. This results in a network that is dense with paths between nodes but lacks stable configurations, as indicated by the gray curve's abrupt descent without intermediate steps. Most notably, such a network would lack hubs (see also

Figure 6. **Percolation analysis. a:** percolation curves over recording sessions compared across behavioral conditions and vs the null model for both animals. Each curve is a recording session. As sketched, the level of complexity of the underlying networks increases traveling the curves from left to right, whereas the hierarchy can be explored climbing down the steps of the curves (the fraction of GSCC, z axis), measured in units of information (TE, x axis). **b:** comparison between the point of descent between behavioral conditions (mean $\pm$ SEM) and against the null model distribution. **c:** comparison between the measured slopes between behavioral conditions (mean $\pm$ SEM) and against the null model distribution. Stop trials resulted more complex and less hierarchical than Go trials while having comparable values between themselves (Kruskal-Wallis, $p < 0.01$ Bonferroni corrected for multiple comparisons). In all panels, for simplicity and homogeneity, TE is reported normalized to its maximum so to obtain TE $\in [0, 1]$. The null distribution is obtained from 500 runs of the null model (see Section). Since the null distributions for the different behavioral conditions were indistinguishable, we only reported one for brevity. (**GSCC**: Giant Strongly Connected Component; **NM**: Null Model).

Figure S1). The slope of the curves in panel c demonstrates that the quantity of information necessary to

obtain equivalent stable configurations (i.e.the same values of y in panel a) is smaller in Go trials than in

Stop trials. This suggests that the higher inherent complexity of the latter is related to increased resource

recruitment. These findings confirm how the local links of the clusters impact the overall state of the

network, changing how it manages information transmission.

A recap

In summary, we have demonstrated the existence of diverse neural modules/clusters (Figure 4) and have

described how these clusters interact and distribute information in various ways, thus contributing to the

state of the mesoscopic network and its level of complexity, each with its own distinct features (Figures 5

and 6). The wiring of information transmission within the network can be efficiently summarized in

Figure 7 using the interaction measure *I* (see Materials and Methods, Table S2 and S3 for further details).

The **C_3** cluster is the primary determinant of the global network state, while the **C_1** and **C_2** play

supporting roles. The strength and direction of the interactions between **C_1** and **C_2** hint at the

specialization of these clusters for behavioral conditions. Specifically, in Go trials, information travels

from **C_1** to **C_2** while in correct Stop trials from **C_2** to **C_1**. The flow of information from the **C_1** and

C_2 to the **C_3** followed the same pattern. The network topology during wrong Stop trials was, as

expected, a hybrid of the configurations observed in Go and correct Stop trials. We discovered that

correct and wrong Stop trials have the same overall topological organization but differ in the details and

hierarchies of information processing. In Wrong Stop trials, we observed a mixture of interactions

between clusters, similar to what was observed in Go trials, along with evidence of attempted inhibition.

Our findings revealed that motor control at the PMd level revolves around a well-defined hierarchical

scheme in which **C_3** are the pivotal nodes in the network and the **C_1** and **C_2** provide critical support in

movement generation and inhibition, respectively.

DISCUSSION

520 In this work, we investigated the patterns of local information transfer in a cortical network directly

521 involved in movement decision-making (the PMd). We used a combined TE and graph theory-based

522 approach to analyze the aggregate spiking activity of simultaneously recorded populations of neurons.

Figure 7. Information spreading among clusters before movement generation and cancellation. Arrows between groups of modules are scaled according to the corresponding average value of **I** (among sessions and animals). The level of hierarchy in which each cluster is ranked is graphically represented as the vertical positioning of the nodes; such level depends on the set of measures explored and could be efficiently summarized by the index I. Since the **C_3** is the dominant cluster in the network, its interactions are always shown. For the other clusters, only the interactions at least one order of magnitude greater than the others and statistically significant across behavioral conditions are shown. Wrong Stop trials are not shown; see Tables S5 and S6 for further details.

Our results contribute to moving forward the knowledge on the neural basis of motor control at different levels.

A topological approach to local information propagation in a motor decision task

Although TE is growing in popularity in modern neuroscience, its application with graph theory to study patterns of directed connectivity has been limited, and restricted mostly, to either single neurons (Gerhard, Pipa, Lima, Neuenschwander, & Gerstner, 2011; Shimono & Beggs, 2015; Varley et al., 2023) or in vitro (Buehlmann & Deco, 2010; Kajiwara et al., 2021; Newman et al., 2022; Nigam et al., 2016; Orlandi, Stetter, Soriano, Geisel, & Battaglia, 2014; Shimono & Beggs, 2015; Thivierge, 2014), and in silico studies (Ito et al., 2011). Its applications to behavioral neurophysiology have been instead extremely limited. However, in recent times, authors have started to explore more consistently the joint information theory-complex networks approach under various experimental settings (Kajiwara et al., 2021; Newman et al., 2022; Schroeder et al., 2016; Shimono & Beggs, 2015; N. Timme et al., 2014;

Varley et al., 2023). In an early contribution, Gerhard et al. (Gerhard et al., 2011) examined spike train recordings from a rhesus monkey's visual system during a fixation task, employing a different approach to measure directed connectivity. Besserve and colleagues (Besserve, Lowe, Logothetis, Schölkopf, & Panzeri, 2015)are credited with applying TE to invasive electrophysiological data from the visual system, although without incorporating graph theory techniques. Another significant contribution comes from Honey and colleagues (Honey, Kötter, Breakspear, & Sporns, 2007), who explored the relationship between the functional and structural properties of a large-scale interregional anatomical network in the macaque cortex using TE, albeit at different spatial and temporal scales. Wollstadt and collaborators (Wollstadt et al., 2014) considered larger spatial and temporal scales than those investigated here and introduced the "ensemble method" for estimating TE under nonstationary conditions, which utilizes the multi-trial structure in neuroscience data for more accurate time-resolved TE estimation. This represents a difference from our work, where, instead, the exchange of information is inferred in a fixed time window. The choice was motivated by our goal: characterize the state of the PMd network at a specific epoch of the task and compare it across behavioral conditions. To do so, we needed several network realizations (over trials) in order to obtain a statistically robust estimate of its topological properties. In contrast, the ensemble approach would have led to a single representation of the temporal evolution of the network. A temporally precise but unique one. It could be argued that this constitutes a limitation of our analysis. Instead, we believe it is the way to answer a different scientific question. We also believe that the ensemble method is extremely powerful and it could be of interest to use it on our data in future studies. An attempt to study voluntary action control through analysis of directed connectivity was made by Jahfari and colleagues (Jahfari et al., 2011) but on human fMRI data. Hence, to the best of our knowledge, this report is one of two studies that use graph theory to analyze the information transfer network of a specific cortical area at the mesoscale level *in vivo* and during cognitive motor control. The only other work on the topic known to us is the extremely recent paper from Varley et al.(Varley et al., 2023) who, using the same dataset as Dann et al.(Dann, Michaels, Schaffelhofer, & Scherberger, 2016), quantified information transmission between single neurons in the fronto-parietal areas AIP, F5, and M1 during various epochs of a delayed grasping task. The authors compared several components of information processing, including also a multi-variate estimate of Transfer Entropy very similar to ours, and inspected how they changed as the task unfolded (see next paragraph). The level of description obtained in our work is more detailed compared to previous ones on motor control. Indeed, we were able

to specify how the decision on whether to move or to stop is implemented in PMd at the meso-scale

population level, who are the (key) players that manage information transmission, and how information

propagation shape the hierarchies and the complexity of the network depending on the behavioral

conditions. Notably, in our framework, neither any a priori assumption nor a specific modeling technique

was needed. Our entirely data-driven approach, in addition to complementing the most recent models for

motor generation and suppression (Boucher, Palmeri, Logan, & Schall, 2007; Lo, Boucher, Paré, Schall,

& Wang, 2009) allows us to address their primary limitation, which involves the need to fine-tune

numerous biophysical parameters to achieve an accurate fit with experimental data. It is known that to

fully understand the neural mechanisms behind motor control future research should focus on

cortico-cortical and cortico-subcortical interactions through massive simultaneous recordings. In this

scenario, a topological information-based approach is unquestionably necessary and should be favored

over other common approaches that rely on the analysis of covariance between neurons or mesoscopic

signals(Chandrasekaran, Peixoto, Newsome, & Shenoy, 2017; Churchland et al., 2012; Clawson et al.,

2019; Kaufman et al., 2016; Mattia et al., 2013; Pani et al., 2022). On one hand, these methods enable a

precise temporal description of neural variability. In contrast, they lack the ability to describe information

dynamics between neurons or provide quantitative insights into network topology and hierarchy. Without

this spectrum of details the computational strategy underlying motor control (and neural circuitry

computation in general) would remain yet elusive.

Fine-grained information transfer underlies motor decisions in PMd

We found that neuronal activities could be organized in four clusters that actively participate and interact,

with distinct roles, both in movement execution and cancellation. This is a step forward in the

conceptualization of the neural processes underlying motor control. Indeed, other studies describe

movement generation as the result of accumulation processes toward a threshold, or as a transition of

neuronal trajectories across specific subspaces. Some studies describe the accumulation process as

instantiated by specific neuronal types (as for gaze-shifting neurons in the frontal eye field and in the

superior colliculus for saccades). Others consider the neuronal trajectory reflecting the dynamics of the

entire neuronal population without considering specific neuronal types. units (Boucher et al., 2007;

Kaufman, Churchland, Ryu, & Shenoy, 2014; Lo et al., 2009; Marcos et al., 2013; Pani et al., 2022;

Schall, Palmeri, & Logan, 2017; Verbruggen & Logan, 2008; Wei, Rubin, & Wang, 2015) Instead, we

showed that information is hierarchically transferred between actors with a well-identifiable role, e.g. the

cluster (C_3) acting as the most important hub for information spreading across behavioral

conditions.This reflects the presence of high-order complexity at the population level, even in small

cortical regions during behavioral control, regardless of the nature of neurons in each cluster, whether

excitatory or inhibitory. Our findings clearly show that hierarchical control is not only exerted across

brain regions, but it is also implemented locally by specific neuronal clusters (the C_3 cluster in our case)

over the others. This is in line with evidence provided in recent years that the brain is hierarchically

organized both globally and locally on a spatial (Bardella et al., 2016, 2020; Felleman & Van Essen,

1991; C. Hilgetag, O'Neill, & Young, 2000; C. C. Hilgetag & Kaiser, 2004; Kaiser, 2010; Kaiser, Görner,

& Hilgetag, 2007; Sporns, Honey, & Kötter, 2007; Sporns, Tononi, & Kötter, 2005; Tononi, Sporns, &

Edelman, 1994; Varley et al., 2023; Watanabe et al., 2014; Zamora-López, 2010; Zeki & Shipp, 1988)

(for a detailed review see Hilgetag et al., 2020(C. C. Hilgetag & Goulas, 2020)) and temporal scale

(Cirillo, Fascianelli, Ferrucci, & Genovesio, 2018; Fox et al., 2005; He, 2011; Morcos & Harvey, 2016;

Shine et al., 2016; Vidaurre, Smith, & Woolrich, 2017; Zalesky, Fornito, Cocchi, Gollo, & Breakspear,

2014). We revealed both condition-specific and nonspecific neuronal clusters, demonstrating that

information transfer and network complexity depend on behavioral outcomes. Our depiction aligns with

the current view by identifying specific clusters primarily involved in the generation (C_1 cluster) or

inhibition (C_2 cluster) of movements. Additionally, it introduces a high-order level of control (C_3

cluster) not previously proposed in other works.This fully portrays the interaction patterns underlying

inhibitory control at the PMd level (see Fig. 7). How does this relate to the current behavioral models?

According to the race model, the Stop process should be 'activated' only in the Stop trials following the

presentation of the Stop signal. However, increasing evidence (Drummond, Cressman, & Carlsen, 2017;

Dunovan, Lynch, Molesworth, & Verstynen, 2015) suggests that the Stop process is also active in Go

trials. Our results show that such co-activation corresponds to information revolving around four clusters

arranged in higher-order connectivity patterns. This is testified by the hierarchical levels in which the C_1

cluster and C_2 cluster are ranked in Go and Stop trials. Moreover, our results show that the C_2 cluster

exhibits a flow of information directed to the C_3 cluster that is significantly higher during correct Stop

trials compared to the other trial types. This suggests that some of the information that the C_2 cluster

handles in attempting to abort the movement is directed to influence the C_3 cluster, which are

extensively involved in facilitating the overall process. This evidence implies that the reason for the

unsuccessful attempts in the wrong Stop trials is associated with the strong flow directed from the C_1

cluster to the C_4 cluster, similar to what occurs in the Go trials This aligns with the limited involvement

of the C_2 cluster, which distributes significantly less information compared to properly inhibited

movements (see 7 and Tables S2, S3 and S6). The found subdivision in clusters and the presence of hubs

reveal that the cortical information dynamics behind motor control is extremely rich and must be

accounted for to generate more comprehensive behavioral models. The topological structure detected

across behavioral conditions exhibits stable rules for the information transmission: the one based on

hubs.This implies that the presence of high-degree nodes is a constituent feature of neural processing in a

cortical network directly involved in cognitive control, as is the PMd.. This is consistent with our

previous study (Bardella et al., 2020), in which we showed how the functional PMd network organization

differs between movement generation and inhibition in terms of hierarchy and centrality of nodes. It is

also in agreement with other works that found high degree nodes in silico (Gu, Qi, & Gong, 2019), in

cortical and hippocampal in vitro networks (Antonello et al., 2022; Gal et al., 2017; Nigam et al., 2016;

Perin, Berger, & Markram, 2011; Schroeter, Charlesworth, Kitzbichler, Paulsen, & Bullmore, 2015;

Shimono & Beggs, 2015; N. Timme et al., 2014), in vivo (Dann et al., 2016; Varley et al., 2023) and

anatomical structural networks (Honey et al., 2007). Within this topology, different information exchange

regimes are realized. On one hand, correct and wrong Stop trials resulted in similar levels of complexity.

On the other, during Go and wrong Stop trials clusters were recruited to exchange information in a

similar fashion. This indicates a transition of the information network from a centralized state during Go

trials to a distinct one during Stop trials. The latter is characterized by higher complexity in information

transfer and increased randomness. This change signals a strong association between the network state

and the behavioral condition. The structure found in the Go trials implies a high and very robust

communication strategy compared to that observed in trials in which a Stop is presented.Conversely, the

network configuration associated with a planned and the inhibited movement appears to be topologically

very similar to the previous one but with some degree of randomness 'injected' into the interactions. This

shows how a network engaged in a cognitive task can adapt its functionality by altering its complexity

and heterogeneity, utilizing different resources while maintaining the stable generating rules of its

topology. We interpret the reorganization occurring in correct Stop trials as the execution in the local

network, through the intervention of the C_2 cluster, of a command originating from other regions.

Indeed, as known, the PMd is part of a larger network subserving motor control based on frontal, parietal,

subcortical and spinal cord structures. It is reasonable to think that during Go trials the hubs serve to convey the command to move to other (and possibly anatomically lower) cortical, subcortical, and spinal circuits that will eventually promote muscle activation. What is observed during correct Stop trials could reflect the PMd collective reaction to the incoming inhibitory input that prevents the execution of the programmed movement. The volition to inhibit would be locally implemented as 'the attenuation of the movement state', achieved by adjusting the level of randomness in the network's wiring scheme, resulting in a more complex organization. This appears convenient and straightforward to implement at the network level. To transmit the message, it would only require the recruitment of dedicated resources (the C_2 cluster) and the redirection of information flow through different paths between nodes. The state of wrong Stop trials is topologically related to that of a correct Stop trial and, at the same time, exhibits the same patterns of information transfer as in Go trials. This may result from an erroneous transmission of inhibitory coding. It can be speculated that this is due to a persistent strong recruitment of the C_1 cluster and a contemporary weaker involvement of the C_2 cluster, as it happens during Go trials. This aligns with our recent paper where we analyzed the neural activity dynamics in the same task using a state-space approach (Pani et al., 2022). In that study, we demonstrated the existence of a subspace in which neural activities are confined before movement generation and during movement suppression.

One weakness of this study is that we cannot account for the information dynamics between PMd and other structures of the reaching network. Therefore, additional research, in a multi-area framework, will be needed to unambiguously clarify the interactions of the PMd network with other structures and to investigate to whom the clusters project or are connected to. The multi-area approach is instead a strength of the work of Varley's and colleagues (Varley et al., 2023) with which our results are only partially comparable. They did not record from the PMd and the task they analyzed did not include inhibition trials. For the shared part (the movement trials), the results are in agreement: the presence of hierarchical hubs also emerged as a core topological attribute in their case, further confirming the ubiquitous presence of this feature of neural organization in cognitive control. The 'processing units' they described, emerge from averaged interactions of macro ensembles of neurons, thus neglecting their decomposition into individual contributions (and their interactions) within the same area. Both works thus have their strengths and weaknesses, testifying how for a more complete understanding of behavioral control this type of approach needs to be systematized. Considering the nature of the mesoscopic signal we studied, it was not possible to resolve the contributions of individual neurons within each of the modules forming

the clusters. This relates to another question that remains unexplored in this article: whether and how the

excitatory/inhibitory nature of individual neurons could affect cluster composition and role. To this end,

the framework of information theory is essential, as extensions of the TE measure have recently been

proposed to track the excitatory or inhibitory signatures of information exchange (Goetze & Lai, 2019;

Kajiwara et al., 2021). Our conclusions suggest that the collective network organization we discovered

represents the neural implementation for the voluntary motor control at the PMd level.

ACKNOWLEDGMENTS

This work was in part supported by European Union Horizon 2020 Research and Innovation Programme

(Grant 945539; Human Brain Project Specific Grant Agreement 3, to SF) and received partial support

from the Italian National Recovery and Resilience Plan (NRRP), M4C2, funded by the European Union -

NextGenerationEU (Project IR0000011, CUP B51E22000150006, "EBRAINS-Italy", to SF), from EC

ERANET CHIST-ERA 2019 grant 'MUCCA' (to SF), from grant RM11916B89232364 (to PP) and from

grant 'Borse di Studio 2020' Fondazione Baroni (to GB).

REFERENCES

Albert, R., & Barabási, A.-L. (2002). Statistical mechanics of complex networks. *Reviews of Modern Physics*, *74*(1), 47–97.

Retrieved from https://link.aps.org/doi/10.1103/RevModPhys.74.47 doi:

10.1103/RevModPhys.74.47

Antonello, P. C., Varley, T. F., Beggs, J., Porcionatto, M., Sporns, O., & Faber, J. (2022). Self-organization of in vitro

neuronal assemblies drives to complex network topology. *eLife*, *11*. doi: 10.7554/ELIFE.74921

Band, G. P., van der Molen, M. W., & Logan, G. D. (2003). Horse-race model simulations of the stop-signal procedure.

Acta Psychologica, *112*(2), 105–142. doi: 10.1016/S0001-6918(02)00079-3

Barabási, A. L. (2013). *Network science* (Vol. 371) (No. 1987). Royal Society. Retrieved from

http://dx.doi.org/10.1098/rsta.2012.0375 doi: 10.1098/rsta.2012.0375

Barabási, A. L., & Albert, R. (1999). Emergence of scaling in random networks. *Science*, *286*(5439), 509–512. Retrieved

from https://www.science.org/doi/10.1126/science.286.5439.509 doi:

10.1126/science.286.5439.509

Bardella, G., Bifone, A., Gabrielli, A., Gozzi, A., & Squartini, T. (2016). Hierarchical organization of functional

connectivity in the mouse brain: A complex network approach. *Scientific Reports*, *6*(1), 1–11. Retrieved from

www.nature.com/scientificreports doi: 10.1038/srep32060

Bardella, G., Pani, P., Brunamonti, E., Giarrocco, F., & Ferraina, S. (2020). The small scale functional topology of

movement control: Hierarchical organization of local activity anticipates movement generation in the premotor cortex of

primates. *NeuroImage*, *207*. doi: 10.1016/j.neuroimage.2019.116354

Barthó, P., Hirase, H., Monconduit, L., Zugaro, M., Harris, K. D., & Buzsáki, G. (2004). Characterization of neocortical

principal cells and interneurons by network interactions and extracellular features. *Journal of neurophysiology*, *92*(1),

600–608. Retrieved from https://pubmed.ncbi.nlm.nih.gov/15056678/ doi: 10.1152/JN.01170.2003

Bassett, D. S., & Sporns, O. (2017). Network neuroscience. *Nature Neuroscience 2017 20:3*, *20*(3), 353–364. Retrieved

from https://www.nature.com/articles/nn.4502 doi: 10.1038/nn.4502

Battaglia-Mayer, A., Buiatti, T., Caminiti, R., Ferraina, S., Lacquaniti, F., & Shallice, T. (2014). *Correction and suppression*

of reaching movements in the cerebral cortex: Physiological and neuropsychological aspects (Vol. 42). Elsevier Ltd.

Retrieved from https://pubmed.ncbi.nlm.nih.gov/24631852/ doi: 10.1016/j.neubiorev.2014.03.002

Besserve, M., Lowe, S. C., Logothetis, N. K., Schölkopf, B., & Panzeri, S. (2015). Shifts of Gamma Phase across Primary

Visual Cortical Sites Reflect Dynamic Stimulus-Modulated Information Transfer. *PLoS Biology*, *13*(9), e1002257.

Retrieved from http://dx.doi.org/10.6084/

m9.figshare.1460872.Thetimeseriesofgamma-rangesignalsandofmulti-unitactivity doi:

10.1371/journal.pbio.1002257

Bordier, C., Nicolini, C., & Bifone, A. (2017). Graph analysis and modularity of brain functional connectivity networks:

Searching for the optimal threshold. *Frontiers in Neuroscience*, *11*(AUG), 441. doi: 10.3389/fnins.2017.00441

Bordier, C., Nicolini, C., Forcellini, G., & Bifone, A. (2018). Disrupted modular organization of primary sensory brain

areas in schizophrenia. *NeuroImage: Clinical*, *18*, 682–693. doi: 10.1016/j.nicl.2018.02.035

Bossomaier, T., Barnett, L., Harré, M., & Lizier, J. T. (2016). *An introduction to transfer entropy: Information flow in*

complex systems. Springer International Publishing. doi: 10.1007/978-3-319-43222-9/COVER

Boucher, L., Palmeri, T. J., Logan, G. D., & Schall, J. D. (2007). Inhibitory control in mind and brain: An interactive race

model of countermanding saccades. *Psychological Review*, *114*(2), 376–397. doi: 10.1037/0033-295X.114.2.376

Buehlmann, A., & Deco, G. (2010). Optimal information transfer in the cortex through synchronization. *PLoS*

Computational Biology, *6*(9), 1000934. Retrieved from www.ploscompbiol.org doi:

10.1371/journal.pcbi.1000934

Callaway, D. S., Newman, M. E., Strogatz, S. H., & Watts, D. J. (2000). Network Robustness and Fragility: Percolation on

Random Graphs. *Physical Review Letters*, *85*(25), 5468. Retrieved from

https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.85.5468 doi:

10.1103/PhysRevLett.85.5468

Caminiti, R., Johnson, P. B., Galli, C., Ferraina, S., & Burnod, Y. (1991). Making arm movements within different parts of

space: the premotor and motor cortical representation of a coordinate system for reaching to visual targets. *Journal of*

Neuroscience, *11*(5), 1182–1197. Retrieved from https://www.jneurosci.org/content/11/5/

1182https://www.jneurosci.org/content/11/5/1182.abstract doi:

10.1523/JNEUROSCI.11-05-01182.1991

Chandrasekaran, C., Peixoto, D., Newsome, W. T., & Shenoy, K. V. (2017). Laminar differences in decision-related neural

activity in dorsal premotor cortex. *Nature Communications 2017 8:1*, *8*(1), 1–16. Retrieved from

https://www.nature.com/articles/s41467-017-00715-0 doi: 10.1038/s41467-017-00715-0

Choi, Y. S., Koenig, M. A., Jia, X., & Thakor, N. V. (2010). Quantifying time-varying multiunit neural activity using

entropy-based measures. *IEEE Transactions on Biomedical Engineering*, *57*(11), 2771–2777. doi:

10.1109/TBME.2010.2049266

Churchland, M. M., Cunningham, J. P., Kaufman, M. T., Foster, J. D., Nuyujukian, P., Ryu, S. I., ... Shenoy, K. V. (2012).

Neural population dynamics during reaching. *Nature*, *487*(7405), 51–56. Retrieved from /pmc/articles/

PMC3393826/?report=abstracthttps://www.ncbi.nlm.nih.gov/pmc/articles/PMC3393826/

doi: 10.1038/nature11129

Cimini, G., Squartini, T., Saracco, F., Garlaschelli, D., Gabrielli, A., & Caldarelli, G. (2019). The statistical physics of

real-world networks. *Nature Reviews Physics*, *1*(1), 58–71. Retrieved from

https://www.nature.com/articles/s42254-018-0002-6 doi: 10.1038/s42254-018-0002-6

Cirillo, R., Fascianelli, V., Ferrucci, L., & Genovesio, A. (2018). Neural Intrinsic Timescales in the Macaque Dorsal

Premotor Cortex Predict the Strength of Spatial Response Coding. *iScience*, *10*, 203–210. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/30529952/ doi: 10.1016/J.ISCI.2018.11.033

Cisek, P., & Kalaska, J. F. (2010). Neural mechanisms for interacting with a world full of action choices. *Annual Review of*

Neuroscience, *33*, 269–298. Retrieved from https://pubmed.ncbi.nlm.nih.gov/20345247/ doi:

10.1146/annurev.neuro.051508.135409

Clawson, W., Vicente, A. F., Ferraris, M., Bernard, C., Battaglia, D., & Quilichini, P. P. (2019). Computing hubs in the

hippocampus and cortex. *Science Advances*, *5*(6), eaax4843. Retrieved from

http://advances.sciencemag.org/ doi: 10.1126/sciadv.aax4843

Dann, B., Michaels, J. A., Schaffelhofer, S., & Scherberger, H. (2016). Uniting functional network topology and oscillations

in the fronto-parietal single unit network of behaving primates. *eLife*, *5*(AUGUST). doi: 10.7554/eLife.15719

Del Ferraro, G., Moreno, A., Min, B., Morone, F., Pérez-Ramírez, U., Pérez-Cervera, L., ... Makse, H. A. (2018). Finding

influential nodes for integration in brain networks using optimal percolation theory. *Nature Communications 2018 9:1*,

9(1), 1–12. Retrieved from https://www.nature.com/articles/s41467-018-04718-3 doi:

10.1038/s41467-018-04718-3

Drummond, N. M., Cressman, E. K., & Carlsen, A. N. (2017). Go-activation endures following the presentation of a

stop-signal: evidence from startle. *J Neurophysiol*, *117*(2), 403–411. Retrieved from www.jn.org doi:

10.1152/jn.00567.2016.-It

Dunovan, K., Lynch, B., Molesworth, T., & Verstynen, T. (2015). Competing basal ganglia pathways determine the

difference between stopping and deciding not to go. *eLife*, *4*(September 2015). Retrieved from

/pmc/articles/PMC4686424//pmc/articles/PMC4686424/?report=abstracthttps://

www.ncbi.nlm.nih.gov/pmc/articles/PMC4686424/ doi: 10.7554/ELIFE.08723

Faes, L., Kugiumtzis, D., Nollo, G., Jurysta, F., & Marinazzo, D. (2015). Estimating the decomposition of predictive

information in multivariate systems. *PHYSICAL REVIEW E*, *91*, 32904. doi: 10.1103/PhysRevE.91.032904

Faes, L., Marinazzo, D., Montalto, A., & Nollo, G. (2014). Lag-specific transfer entropy as a tool to assess cardiovascular

and cardiorespiratory information transfer. *IEEE Transactions on Biomedical Engineering*, *61*(10), 2556–2568. Retrieved

from https://pubmed.ncbi.nlm.nih.gov/24835121/ doi: 10.1109/TBME.2014.2323131

Faes, L., Nollo, G., & Chon, K. H. (2008). Assessment of granger causality by nonlinear model identification: Application

to short-term cardiovascular variability. *Annals of Biomedical Engineering*, *36*(3), 381–395. Retrieved from

https://link.springer.com/article/10.1007/s10439-008-9441-z doi:

10.1007/S10439-008-9441-Z/FIGURES/8

Faes, L., Nollo, G., & Porta, A. (2011). Information-based detection of nonlinear Granger causality in multivariate

processes via a nonuniform embedding technique. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*,

83(5), 051112. Retrieved from https://link.aps.org/doi/10.1103/PhysRevE.83.051112 doi:

10.1103/PhysRevE.83.051112

Faes, L., Pinna, G. D., Porta, A., Maestri, R., & Nollo, G. (2004). Surrogate data analysis for assessing the significance of

the coherence function. *IEEE Transactions on Biomedical Engineering*, *51*(7), 1156–1166. doi:

10.1109/TBME.2004.827271

Felleman, D. J., & Van Essen, D. C. (1991). Distributed Hierarchical Processing in the Primate Cerebral Cortex. *Cerebral*

Cortex, *1*(1), 1–47. Retrieved from https://academic.oup.com/cercor/article/1/1/1/408896 doi:

10.1093/CERCOR/1.1.1-A

Ferraina, S., Paré, M., & Wurtz, R. H. (2002). Comparison of cortico-cortical and cortico-collicular signals for the

generation of saccadic eye movements. *Journal of Neurophysiology*, *87*(2), 845–858. doi: 10.1152/jn.00317.2001

Fetz, E. E. (1992). Are movement parameters recognizably coded in the activity of single neurons? *Behavioral and Brain*

Sciences, *15*(4), 679–690. Retrieved from

https://www.cambridge.org/core/journals/behavioral-and-brain-sciences/article/

abs/are-movement-parameters-recognizably-coded-in-the-activity-of-single-neurons/

875AA83CCD35507B762373869BC88D72 doi: 10.1017/S0140525X00072599

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005). The human brain is

intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of*

Sciences of the United States of America, *102*(27), 9673–9678. Retrieved from

www.pnas.orgcgidoi10.1073pnas.0504136102 doi: 10.1073/pnas.0504136102

Gal, E., London, M., Globerson, A., Ramaswamy, S., Reimann, M. W., Muller, E., . . . Segev, I. (2017). Rich

cell-type-specific network topology in neocortical microcircuitry. *Nature Neuroscience*, *20*(7), 1004–1013. Retrieved

from https://pubmed.ncbi.nlm.nih.gov/28581480/ doi: 10.1038/nn.4576

Gallos, L. K., Makse, H. A., & Sigman, M. (2012). A small world of weak ties provides optimal global integration of

self-similar modules in functional brain networks. *Proceedings of the National Academy of Sciences of the United States*

of America, *109*(8), 2825–2830. Retrieved from https://pubmed.ncbi.nlm.nih.gov/22308319/

https://pubmed.ncbi.nlm.nih.gov/22308319/?dopt=Abstract doi: 10.1073/pnas.1106612109

Georgopoulos, A. P., Kalaska, J. F., Caminiti, R., & Massey, J. T. (1982). On the relations between the direction of

two-dimensional arm movements and cell discharge in primate motor cortex. *The Journal of neuroscience : the official*

journal of the Society for Neuroscience, *2*(11), 1527–1537. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/7143039/ doi: 10.1523/JNEUROSCI.02-11-01527.1982

Gerhard, F., Pipa, G., Lima, B., Neuenschwander, S., & Gerstner, W. (2011). Extraction of Network Topology From

Multi-Electrode Recordings: Is there a Small-World Effect? *Frontiers in Computational Neuroscience*, *5*, 4. Retrieved

from http://journal.frontiersin.org/article/10.3389/fncom.2011.00004/abstract doi:

10.3389/fncom.2011.00004

Giarrocco, F., Bardella, G., Giamundo, M., Fabbrini, F., Brunamonti, E., Pani, P., & Ferraina, S. (2021). Neuronal dynamics

of signal selective motor plan cancellation in the macaque dorsal premotor cortex. *Cortex*, *135*, 326–340. doi:

10.1016/j.cortex.2020.09.032

Goetze, F., & Lai, P. Y. (2019). Reconstructing positive and negative couplings in Ising spin networks by sorted local

transfer entropy. *Physical Review E*, *100*(1), 012121. Retrieved from

https://journals.aps.org/pre/abstract/10.1103/PhysRevE.100.012121 doi:

10.1103/PHYSREVE.100.012121/FIGURES/6/MEDIUM

Gómez, C., Lizier, J. T., Schaum, M., Wollstadt, P., Grützner, C., Uhlhaas, P., . . . Wibral, M. (2014). Reduced predictable

information in brain signals in autism pectrum disorder. *Frontiers in Neuroinformatics*, *8*(FEB), 75547. doi:

10.3389/FNINF.2014.00009/BIBTEX

Gómez-Herrero, G., Wu, W., Rutanen, K., Soriano, M. C., Pipa, G., & Vicente, R. (2015). Assessing Coupling Dynamics

from an Ensemble of Time Series. *Entropy 2015, Vol. 17, Pages 1958-1970*, *17*(4), 1958–1970. Retrieved from

https://www.mdpi.com/1099-4300/17/4/1958/htmhttps://www.mdpi.com/1099-4300/17/4/

1958 doi: 10.3390/E17041958

Graf, A. (2016). *On the Strongly Connected Components of Random Directed Graphs with Given Degree Sequences* (Tech.

Rep.). University of Waterloo. Retrieved from http://hdl.handle.net/10012/10681

Grassberger, P., & Procaccia, I. (1983). Estimation of the Kolmogorov entropy from a chaotic signal. *Physical Review A*,

28(4), 2591. Retrieved from https://journals.aps.org/pra/abstract/10.1103/PhysRevA.28.2591

doi: 10.1103/PhysRevA.28.2591

Gu, Y., Qi, Y., & Gong, P. (2019). Rich-club connectivity, diverse population coupling, and dynamical activity patterns

emerging from local cortical circuits. *PLoS Computational Biology*, *15*(4), e1006902. Retrieved from

https://doi.org/10.1371/journal.pcbi.1006902 doi: 10.1371/journal.pcbi.1006902

Hanes, D. P., Patterson, W. F., & Schall, J. D. (1998). Role of frontal eye fields in countermanding saccades: Visual,

movement, and fixation activity. *Journal of Neurophysiology*, *79*(2), 817–834. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/9463444/ doi: 10.1152/jn.1998.79.2.817

He, B. J. (2011). Scale-free properties of the functional magnetic resonance imaging signal during rest and task. *Journal of*

Neuroscience, *31*(39), 13786–13795. Retrieved from

http://brainvis.wustl.edu/wiki/index.php/Caret: doi: 10.1523/JNEUROSCI.2111-11.2011

Herreras, O. (2016). Local Field Potentials: Myths and Misunderstandings. *Frontiers in Neural Circuits*, *10*(DEC), 101.

Retrieved from http://journal.frontiersin.org/article/10.3389/fncir.2016.00101/full

doi: 10.3389/fncir.2016.00101

Hilgetag, C., O'Neill, M. A., & Young, M. P. (2000). Hierarchical organization of macaque and cat cortical sensory systems

explored with a novel network processor. *Philosophical Transactions of the Royal Society of London. Series B: Biological*

Sciences, *355*(1393), 71–89. Retrieved from

https://royalsocietypublishing.org/doi/10.1098/rstb.2000.0550 doi: 10.1098/rstb.2000.0550

Hilgetag, C. C., & Goulas, A. (2020). 'Hierarchy' in the organization of brain networks. *Philosophical Transactions of the*

Royal Society B: Biological Sciences, *375*(1796), 20190319. Retrieved from

https://royalsocietypublishing.org/doi/10.1098/rstb.2019.0319 doi: 10.1098/rstb.2019.0319

Hilgetag, C. C., & Kaiser, M. (2004). *Clustered organization of cortical connectivity* (Vol. 2) (No. 3). Springer. Retrieved

from https://link.springer.com/article/10.1385/NI:2:3:353 doi: 10.1385/NI:2:3:353

Honey, C. J., Kötter, R., Breakspear, M., & Sporns, O. (2007). Network structure of cerebral cortex shapes functional

connectivity on multiple time scales. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(24), 10240–10245. Retrieved from https://pubmed.ncbi.nlm.nih.gov/17548818/ doi: 10.1073/pnas.0701519104

Hong, G., & Lieber, C. M. (2019). Novel electrode technologies for neural recordings. *Nature Reviews Neuroscience*, *20*(6), 330–345. Retrieved from www.nature.com/nrn doi: 10.1038/s41583-019-0140-6

Ito, S., Hansen, M. E., Heiland, R., Lumsdaine, A., Litke, A. M., & Beggs, J. M. (2011). Extending Transfer Entropy Improves Identification of Effective Connectivity in a Spiking Cortical Network Model. *PLoS ONE*, *6*(11), e27431. Retrieved from https://dx.plos.org/10.1371/journal.pone.0027431 doi: 10.1371/journal.pone.0027431

Jahfari, S., Waldorp, L., van den Wildenberg, W. P., Scholte, H. S., Ridderinkhof, K. R., & Forstmann, B. U. (2011). Effective connectivity reveals important roles for both the hyperdirect (fronto-subthalamic) and the indirect (fronto-striatal-pallidal) fronto-basal ganglia pathways during response inhibition. *Journal of Neuroscience*, *31*(18), 6891–6899. doi: 10.1523/JNEUROSCI.5253-10.2011

Kaiser, M. (2010). Hierarchy and dynamics of neural networks. *Frontiers in Neuroinformatics*, *4*(AUG), 112. Retrieved from http://journal.frontiersin.org/article/10.3389/fninf.2010.00112/abstract doi: 10.3389/fninf.2010.00112

Kaiser, M., Görner, M., & Hilgetag, C. C. (2007). Criticality of spreading dynamics in hierarchical cluster networks without inhibition. *New Journal of Physics*, *9*(5), 110. Retrieved from http://www.njp.org/ doi: 10.1088/1367-2630/9/5/110

Kajiwara, M., Nomura, R., Goetze, F., Kawabata, M., Isomura, Y., Akutsu, T., & Shimono, M. (2021). Inhibitory neurons exhibit high controlling ability in the cortical microconnectome. *PLOS Computational Biology*, *17*(4), e1008846. Retrieved from https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1008846 doi: 10.1371/JOURNAL.PCBI.1008846

Kaufman, M. T., Churchland, M. M., Ryu, S. I., & Shenoy, K. V. (2014). Cortical activity in the null space: permitting preparation without movement. *Nature neuroscience*, *17*(3), 440. Retrieved from /pmc/articles/PMC3955357//pmc/articles/PMC3955357/?report=abstracthttps://

www.ncbi.nlm.nih.gov/pmc/articles/PMC3955357/ doi: 10.1038/NN.3643

Kaufman, M. T., Seely, J. S., Sussillo, D., Ryu, S. I., Shenoy, K. V., & Churchland, M. M. (2016). The largest response

component in the motor cortex reflects movement timing but not movement type. *eNeuro*, *3*(4), 85–101. Retrieved from

http://dx.doi.org/10.1523/ENEURO.0085-16.2016 doi: 10.1523/ENEURO.0085-16.2016

Komárek, V., Hrnčíř, Z., & Štěrbová, K. (2001). Synchronization as adjustment of information rates: Detection from

bivariate time series. *Physical Review E*, *63*(4), 046211. Retrieved from

https://journals.aps.org/pre/abstract/10.1103/PhysRevE.63.046211 doi:

10.1103/PhysRevE.63.046211

Lo, C. C., Boucher, L., Paré, M., Schall, J. D., & Wang, X. J. (2009). Proactive inhibitory control and attractor dynamics in

countermanding action: A spiking neural circuit model. *Journal of Neuroscience*, *29*(28), 9059–9071. Retrieved from

https://www.jneurosci.org/content/29/28/9059https://www.jneurosci.org/content/29/

28/9059.abstract doi: 10.1523/JNEUROSCI.6164-08.2009

Logan, G. D., & Cowan, W. B. (1984). On the ability to inhibit thought and action: A theory of an act of control.

Psychological Review, *91*(3), 295–327. doi: 10.1037//0033-295x.91.3.295

Marcos, E., Pani, P., Brunamonti, E., Deco, G., Ferraina, S., & Verschure, P. (2013). Neural variability in premotor cortex is

modulated by trial history and predicts behavioral performance. *Neuron*, *78*(2), 249–255. doi:

10.1016/j.neuron.2013.02.006

Mason, A., Nicoli, A., & Stratford, K. (1991). Synaptic transmission between individual pyramidal neurons of the rat visual

cortex in vitro. *Journal of Neuroscience*, *11*(1), 72–84. Retrieved from https://www.jneurosci.org/

content/11/1/72https://www.jneurosci.org/content/11/1/72.abstract doi:

10.1523/JNEUROSCI.11-01-00072.1991

Mastrandrea, R., Gabrielli, A., Piras, F., Spalletta, G., Caldarelli, G., & Gili, T. (2017). Organization and hierarchy of the

human functional brain network lead to a chain-like core. *Scientific Reports 2017 7:1*, *7*(1), 1–13. Retrieved from

https://www.nature.com/articles/s41598-017-04716-3 doi: 10.1038/s41598-017-04716-3

Mastrandrea, R., Piras, F., Gabrielli, A., Banaj, N., Caldarelli, G., Spalletta, G., & Gili, T. (2021). The unbalanced

reorganization of weaker functional connections induces the altered brain network topology in schizophrenia. *Scientific*

Reports, *11*(1), 15400. Retrieved from /pmc/articles/PMC8319172//pmc/articles/PMC8319172/

?report=abstracthttps://www.ncbi.nlm.nih.gov/pmc/articles/PMC8319172/ doi:

10.1038/S41598-021-94825-X

Mattia, M., & Del Giudice, P. (2002). Population dynamics of interacting spiking neurons. *Physical Review E - Statistical*

Physics, Plasmas, Fluids, and Related Interdisciplinary Topics, *66*(5), 19. Retrieved from

https://journals.aps.org/pre/abstract/10.1103/PhysRevE.66.051917 doi:

10.1103/PhysRevE.66.051917

Mattia, M., Ferraina, S., & Del Giudice, P. (2010). Dissociated multi-unit activity and local field potentials: a theory

inspired analysis of a motor decision task. *NeuroImage*, *52*(3), 812–823. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/20100578/ doi: 10.1016/J.NEUROIMAGE.2010.01.063

Mattia, M., Pani, P., Mirabella, G., Costa, S., Del Giudice, P., & Ferraina, S. (2013). Heterogeneous attractor cell assemblies

for motor planning in premotor cortex. *Journal of Neuroscience*, *33*(27), 11155–11168. doi:

10.1523/JNEUROSCI.4664-12.2013

Michael Wibral, J. T. L., Raul Vicente (Ed.). (2014). *Directed Information Measures in Neuroscience — SpringerLink* (1st

ed.). Springer Berlin, Heidelberg. Retrieved from

https://link.springer.com/book/10.1007/978-3-642-54474-3 doi: 10.1007/978-3-642-54474-3

Milekovic, T., Truccolo, W., Grün, S., Riehle, A., & Brochier, T. (2015). Local field potentials in primate motor cortex

encode grasp kinetic parameters. *NeuroImage*, *114*, 338–355. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/25869861/ doi: 10.1016/J.NEUROIMAGE.2015.04.008

Mirabella, G., Pani, P., & Ferraina, S. (2011). Neural correlates of cognitive control of reaching movements in the dorsal

premotor cortex of rhesus monkeys. *Journal of Neurophysiology*, *106*(3), 1454–1466. Retrieved from

https://www.physiology.org/doi/10.1152/jn.00995.2010 doi: 10.1152/jn.00995.2010

Montalto, A., Faes, L., & Marinazzo, D. (2014). MuTE: A MATLAB toolbox to compare established and novel estimators

of the multivariate transfer entropy. *PLoS ONE*, *9*(10), e109462. Retrieved from http://dx.doi.org/10.6084/

m9.figshare.1005245;thecardiovasculardataareincludedinthetoolboxpackage;

theepilepsydataareavailableatthisaddress:http://math.bu.edu/people/kolaczyk/

datasets.html. doi: 10.1371/journal.pone.0109462

Morcos, A. S., & Harvey, C. D. (2016). History-dependent variability in population dynamics during evidence accumulation in cortex. *Nature Neuroscience*, *19*(12), 1672–1681. Retrieved from https://pubmed.ncbi.nlm.nih.gov/27694990/ doi: 10.1038/nn.4403

Munoz, D. P., & Wurtz, R. H. (1993). Fixation cells in monkey superior colliculus I. Characteristics of cell discharge. *Journal of Neurophysiology*, *70*(2), 559–575. Retrieved from https://pubmed.ncbi.nlm.nih.gov/8410157/ doi: 10.1152/jn.1993.70.2.559

Newman, E. L., Varley, T. F., Parakkattu, V. K., Sherrill, S. P., & Beggs, J. M. (2022). Revealing the Dynamics of Neural Information Processing with Multivariate Information Decomposition. *Entropy 2022, Vol. 24, Page 930*, *24*(7), 930. Retrieved from https://www.mdpi.com/1099-4300/24/7/930/htmhttps://www.mdpi.com/1099-4300/24/7/930 doi: 10.3390/E24070930

Nicolini, C., & Bifone, A. (2016). Modular structure of brain functional networks: breaking the resolution limit by Surprise. *Scientific Reports*, *6*. Retrieved from /pmc/articles/PMC4725862//pmc/articles/PMC4725862/?report=abstracthttps://www.ncbi.nlm.nih.gov/pmc/articles/PMC4725862/ doi: 10.1038/SREP19250

Nicolini, C., Bordier, C., & Bifone, A. (2017). Community detection in weighted brain connectivity networks beyond the resolution limit. *NeuroImage*, *146*, 28–39. doi: 10.1016/j.neuroimage.2016.11.026

Nigam, S., Shimono, M., Ito, S., Yeh, F. C., Timme, N., Myroshnychenko, M., . . . Beggs, J. M. (2016). Rich-club organization in effective connectivity among cortical neurons. *Journal of Neuroscience*, *36*(3), 655–669. Retrieved from https://pubmed.ncbi.nlm.nih.gov/26791200/ doi: 10.1523/JNEUROSCI.2177-15.2016

Novelli, L., & Lizier, J. T. (2021). Inferring network properties from time series using transfer entropy and mutual information: Validation of multivariate versus bivariate approaches. *Network Neuroscience*, *5*(2), 373–404. Retrieved from https://dx.doi.org/10.1162/netn_a_00178 doi: 10.1162/NETN_A_00178

Orlandi, J. G., Stetter, O., Soriano, J., Geisel, T., & Battaglia, D. (2014). Transfer entropy reconstruction and labeling of neuronal connections from simulated calcium imaging. *PLoS ONE*, *9*(6), 98842. Retrieved from www.plosone.org doi: 10.1371/journal.pone.0098842

Pani, P., Di Bello, F., Brunamonti, E., D'Andrea, V., Papazachariadis, O., & Ferraina, S. (2014). Alpha- and beta-band

oscillations subserve different processes in reactive control of limb movements. *Frontiers in behavioral neuroscience*,

8(November). Retrieved from https://pubmed.ncbi.nlm.nih.gov/25414649/ doi:

10.3389/FNBEH.2014.00383

Pani, P., Giamundo, M., Giarrocco, F., Mione, V., Fontana, R., Brunamonti, E., ... Ferraina, S. (2022). Neuronal population

dynamics during motor plan cancellation in nonhuman primates. *Proceedings of the National Academy of Sciences of the*

United States of America, *119*(28), e2122395119. Retrieved from

https://www.pnas.org/doi/abs/10.1073/pnas.2122395119 doi: 10.1073/pnas.2122395119

Pani, P., Giarrocco, F., Giamundo, M., Montanari, R., Brunamonti, E., & Ferraina, S. (2018). Visual salience of the stop

signal affects the neuronal dynamics of controlled inhibition. *Scientific Reports 2018 8:1*, *8*(1), 1–13. Retrieved from

https://www.nature.com/articles/s41598-018-32669-8 doi: 10.1038/s41598-018-32669-8

Paré, M., & Hanes, D. P. (2003). Controlled movement processing: Superior colliculus activity associated with

countermanded saccades. *Journal of Neuroscience*, *23*(16), 6480–6489. Retrieved from

https://www.jneurosci.org/content/23/16/6480https://www.jneurosci.org/content/23/

16/6480.abstract doi: 10.1523/jneurosci.23-16-06480.2003

Perin, R., Berger, T. K., & Markram, H. (2011). A synaptic organizing principle for cortical neuronal groups. *Proceedings*

of the National Academy of Sciences of the United States of America, *108*(13), 5419–5424. Retrieved from https://

www.pnas.org/content/108/13/5419https://www.pnas.org/content/108/13/5419.abstract

doi: 10.1073/pnas.1016051108

Pincus, S. M. (1991). Approximate entropy as a measure of system complexity. *Proceedings of the National Academy of*

Sciences of the United States of America, *88*(6), 2297–2301. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/11607165/ doi: 10.1073/PNAS.88.6.2297

Quian Quiroga, R., Kraskov, A., Kreuz, T., & Grassberger, P. (2002). Performance of different synchronization measures in

real data: A case study on electroencephalographic signals. *Physical Review E*, *65*(4), 041903. Retrieved from

https://journals.aps.org/pre/abstract/10.1103/PhysRevE.65.041903 doi:

10.1103/PhysRevE.65.041903

Ramawat, S., Mione, V., Di Bello, F., Bardella, G., Genovesio, A., Pani, P., ... Brunamonti, E. (2022). Different

Contribution of the Monkey Prefrontal and Premotor Dorsal Cortex in Decision Making During a Transitive Inference

Task. *Neuroscience*, *485*, 147–162. doi: 10.1016/J.NEUROSCIENCE.2022.01.013

Ramos, A. M. T., Macau, E. E. N., Lopes, A. M., & Machado, J. A. T. (2017). Minimum Sample Size for Reliable Causal

Inference Using Transfer Entropy. *Entropy 2017*, *19*(4), 150. Retrieved from https://www.mdpi.com/

1099-4300/19/4/150/htmhttps://www.mdpi.com/1099-4300/19/4/150 doi: 10.3390/E19040150

Richman, J. S., & Moorman, J. R. (2000). Physiological time-series analysis using approximate and sample entropy.

American Journal of Physiology - Heart and Circulatory Physiology, *278*(6 47-6), 2039–2049. Retrieved from

https://journals.physiology.org/doi/10.1152/ajpheart.2000.278.6.H2039 doi:

10.1152/AJPHEART.2000.278.6.H2039/ASSET/IMAGES/LARGE/AHEA40640005X.JPEG

Riehle, A., Grün, S., Diesmann, M., & Aertsen, A. (1997). Spike synchronization and rate modulation differentially

involved in motor cortical function. *Science*, *278*(5345), 1950–1953. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/9395398/ doi: 10.1126/science.278.5345.1950

Schall, J. D., Palmeri, T. J., & Logan, G. D. (2017). Models of inhibitory control. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *372*(1718), 20160193. Retrieved from

https://royalsocietypublishing.org/doi/10.1098/rstb.2016.0193 doi: 10.1098/rstb.2016.0193

1020

Schreiber, T. (2000). Measuring information transfer. *Physical Review Letters*, *85*(2), 461–464. Retrieved from

https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.85.461 doi:

10.1103/PhysRevLett.85.461

Schroeder, K. E., Irwin, Z. T., Gaidica, M., Bentley, J. N., Patil, P. G., Mashour, G. A., & Chestek, C. A. (2016). Disruption

of corticocortical information transfer during ketamine anesthesia in the primate brain. *NeuroImage*, *134*, 459–465.

Retrieved from /pmc/articles/PMC4912854/?report=abstracthttps://www.ncbi.nlm.nih.gov/

pmc/articles/PMC4912854/ doi: 10.1016/j.neuroimage.2016.04.039

Schroeter, M. S., Charlesworth, P., Kitzbichler, M. G., Paulsen, O., & Bullmore, E. T. (2015). Emergence of rich-club

topology and coordinated dynamics in development of hippocampal functional networks In vitro. *Journal of*

Neuroscience, *35*(14), 5459–5470. Retrieved from http://www.ru.nl/neuroimaging/fieldtrip/ doi:

10.1523/JNEUROSCI.4259-14.2015

Shannon, C. E. (1948). A Mathematical Theory of Communication. *The Bell System Technical Journal*, *27*, 623–656.

Sharma, G., Annetta, N., Friedenberg, D., Blanco, T., Vasconcelos, D., Shaikhouni, A., ... Bouton, C. (2015). Time

Stability and Coherence Analysis of Multiunit, Single-Unit and Local Field Potential Neuronal Signals in Chronically

Implanted Brain Electrodes. *Bioelectronic Medicine 2015 2:1*, *2*(1), 63–71. Retrieved from

https://bioelecmed.biomedcentral.com/articles/10.15424/bioelectronmed.2015.00010

doi: 10.15424/BIOELECTRONMED.2015.00010

Shenoy, K. V., Sahani, M., & Churchland, M. M. (2013). Cortical Control of Arm Movements: A Dynamical Systems

Perspective. *Annual Review of Neuroscience*, *36*(1), 337–359. Retrieved from

http://www.annualreviews.org/doi/10.1146/annurev-neuro-062111-150509 doi:

10.1146/annurev-neuro-062111-150509

Shimono, M., & Beggs, J. M. (2015). Functional clusters, hubs, and communities in the cortical microconnectome.

Cerebral Cortex, *25*(10), 3743–3757. Retrieved from https://pubmed.ncbi.nlm.nih.gov/25336598/ doi:

10.1093/cercor/bhu252

Shine, J. M., Bissett, P. G., Bell, P. T., Koyejo, O., Balsters, J. H., Gorgolewski, K. J., ... Poldrack, R. A. (2016). The

Dynamics of Functional Brain Networks: Integrated Network States during Cognitive Task Performance. *Neuron*, *92*(2),

544–554. Retrieved from https://pubmed.ncbi.nlm.nih.gov/27693256/ doi:

10.1016/j.neuron.2016.09.018

Sporns, O., Honey, C. J., & Kötter, R. (2007). Identification and Classification of Hubs in Brain Networks. *PLoS ONE*,

2(10), e1049. Retrieved from https://dx.plos.org/10.1371/journal.pone.0001049 doi:

10.1371/journal.pone.0001049

Sporns, O., Tononi, G., & Kötter, R. (2005). The Human Connectome: A Structural Description of the Human Brain. *PLoS*

Computational Biology, *1*(4), e42. Retrieved from

https://dx.plos.org/10.1371/journal.pcbi.0010042 doi: 10.1371/journal.pcbi.0010042

Squires, S., Sytwu, K., Alcala, D., Antonsen, T. M., Ott, E., & Girvan, M. (2013). Weakly explosive percolation in directed

networks. *PHYSICAL REVIEW E*, *87*, 52127. doi: 10.1103/PhysRevE.87.052127

Stark, E., & Abeles, M. (2007). Predicting movement from multiunit activity. *Journal of Neuroscience*, *27*(31), 8387–8394.

Retrieved from https://www.jneurosci.org/content/27/31/8387https://www.jneurosci.org/

content/27/31/8387.abstract doi: 10.1523/JNEUROSCI.1321-07.2007

Supèr, H., & Roelfsema, P. R. (2005). Chronic multiunit recordings in behaving animals: advantages and limitations.

Progress in Brain Research, *147*(SPEC. ISS.), 263–282. doi: 10.1016/S0079-6123(04)47020-4

Swadlow, H. A. (1994). Efferent neurons and suspected interneurons in motor cortex of the awake rabbit: axonal properties,

sensory receptive fields, and subthreshold synaptic inputs. *Journal of neurophysiology*, *71*(2), 437–453. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/8176419/ doi: 10.1152/JN.1994.71.2.437

Thivierge, J.-P. (2014). Scale-free and economical features of functional connectivity in neuronal networks. *PHYSICAL*

REVIEW E, *90*, 22721. doi: 10.1103/PhysRevE.90.022721

Timme, N., Ito, S., Myroshnychenko, M., Yeh, F.-C., Hiolski, E., Hottowy, P., & Beggs, J. M. (2014). Multiplex Networks

of Cortical and Hippocampal Neurons Revealed at Different Timescales. *PLoS ONE*, *9*(12), e115764. Retrieved from

https://dx.plos.org/10.1371/journal.pone.0115764 doi: 10.1371/journal.pone.0115764

Timme, N. M., Ito, S., Myroshnychenko, M., Nigam, S., Shimono, M., Yeh, F.-C., . . . Beggs, J. M. (2016). High-Degree

Neurons Feed Cortical Computations. *PLOS Computational Biology*, *12*(5), e1004858. Retrieved from

https://dx.plos.org/10.1371/journal.pcbi.1004858 doi: 10.1371/journal.pcbi.1004858

Timme, N. M., & Lapish, C. (2018). A Tutorial for Information Theory in Neuroscience. *eNeuro*, *5*(3). Retrieved from

https://www.eneuro.org/content/5/3/ENEURO.0052-18.2018https://www.eneuro.org/

content/5/3/ENEURO.0052-18.2018.abstract doi: 10.1523/ENEURO.0052-18.2018

Tononi, G., Sporns, O., & Edelman, G. M. (1994). A measure for brain complexity: Relating functional segregation and

integration in the nervous system. *Proceedings of the National Academy of Sciences of the United States of America*,

91(11), 5033–5037. Retrieved from https://www.pnas.org/content/91/11/5033https://

www.pnas.org/content/91/11/5033.abstract doi: 10.1073/pnas.91.11.5033

Trautmann, E. M., Stavisky, S. D., Lahiri, S., Ames, K. C., Kaufman, M. T., O'Shea, D. J., . . . Shenoy, K. V. (2019).

Accurate Estimation of Neural Population Dynamics without Spike Sorting. *Neuron*, *103*(2), 292–308. doi:

10.1016/j.neuron.2019.05.003

Ursino, M., Ricci, G., & Magosso, E. (2020). Transfer Entropy as a Measure of Brain Connectivity: A Critical Analysis

With the Help of Neural Mass Models. *Frontiers in Computational Neuroscience*, *14*, 534148. doi:

10.3389/FNCOM.2020.00045/BIBTEX

Vaadia, E., Kurata, K., & Wise, S. P. (1988). Neuronal activity preceding directional and nondirectional cues in the

premotor cortex of rhesus monkeys. *Somatosensory & Motor Research*, *6*(2), 207–230. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/3242346/ doi: 10.3109/08990228809144674

Varley, T. F., Sporns, O., Schaffelhofer, S., Scherberger, H., & Dann, B. (2023). Information-processing dynamics in neural

networks of macaque cerebral cortex reflect cognitive state and behavior. *Proceedings of the National Academy of*

Sciences of the United States of America, *120*(2), e2207677120. Retrieved from

https://www.pnas.org/doi/abs/10.1073/pnas.2207677120 doi:

10.1073/PNAS.2207677120/SUPPL_FILE/PNAS.2207677120.SD05.CSV

Verbruggen, F., & Logan, G. D. (2008). *Response inhibition in the stop-signal paradigm* (Vol. 12) (No. 11). Elsevier

Current Trends. doi: 10.1016/j.tics.2008.07.005

Vicente, R., Wibral, M., Lindner, M., & Pipa, G. (2011). Transfer entropy-a model-free measure of effective connectivity

for the neurosciences. *Journal of Computational Neuroscience*, *30*(1), 45–67. Retrieved from

https://link.springer.com/article/10.1007/s10827-010-0262-3 doi:

10.1007/s10827-010-0262-3

Vidaurre, D., Smith, S. M., & Woolrich, M. W. (2017). Brain network dynamics are hierarchically organized in time.

Proceedings of the National Academy of Sciences of the United States of America, *114*(48), 12827–12832. Retrieved from

https://www.pnas.org/content/114/48/12827https://www.pnas.org/content/114/48/

12827.abstract doi: 10.1073/pnas.1705120114

Vlasov, V., & Bifone, A. (2017). Hub-driven remote synchronization in brain networks. *Scientific Reports 2017 7:1*, *7*(1),

1–11. Retrieved from https://www.nature.com/articles/s41598-017-09887-7 doi:

10.1038/s41598-017-09887-7

Watanabe, T., Hirose, S., Wada, H., Imai, Y., Machida, T., Shirouzu, I., . . . Masuda, N. (2014). Energy landscapes of

resting-state brain networks. *Frontiers in Neuroinformatics*, *8*(FEB), 12. Retrieved from

http://journal.frontiersin.org/article/10.3389/fninf.2014.00012/abstract doi:

10.3389/fninf.2014.00012

Wei, W., Rubin, J. E., & Wang, X. J. (2015). Role of the indirect pathway of the basal ganglia in perceptual decision

making. *Journal of Neuroscience*, *35*(9), 4052–4064. Retrieved from https://www.jneurosci.org/content/

35/9/4052https://www.jneurosci.org/content/35/9/4052.abstract doi:

10.1523/JNEUROSCI.3611-14.2015

Wibral, M., Rahm, B., Rieder, M., Lindner, M., Vicente, R., & Kaiser, J. (2011). Transfer entropy in

magnetoencephalographic data: Quantifying information flow in cortical and cerebellar networks. *Progress in Biophysics*

and Molecular Biology, *105*(1-2), 80–97. doi: 10.1016/j.pbiomolbio.2010.11.006

Wise, S. P., Boussaoud, D., Johnson, P. B., & Caminiti, R. (1997). Premotor and parietal cortex: Corticocortical

connectivity and combinatorial computations. *Annual Review of Neuroscience*, *20*, 25–42. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/9056706/ doi: 10.1146/annurev.neuro.20.1.25

Wollstadt, P., Martínez-Zarzuela, M., Vicente, R., Díaz-Pernas, F. J., & Wibral, M. (2014). Efficient transfer entropy

analysis of non-stationary neural time series. *PLoS ONE*, *9*(7), 102833. Retrieved from www.plosone.org doi:

10.1371/journal.pone.0102833

Wurtz, R. H., Sommer, M. A., Paré, M., & Ferraina, S. (2001). Signal transformations from cerebral cortex to superior

colliculus for the generation of saccades. *Vision Research*, *41*(25-26), 3399–3412. doi: 10.1016/S0042-6989(01)00066-9

Xiong, W., Faes, L., & Ivanov, P. C. (2017). Entropy measures, entropy estimators, and their performance in quantifying

complex dynamics: Effects of artifacts, nonstationarity, and long-range correlations. *Physical Review E*, *95*(6), 1–37. doi:

10.1103/PhysRevE.95.062114

Zalesky, A., Fornito, A., Cocchi, L., Gollo, L. L., & Breakspear, M. (2014). Time-resolved resting-state brain networks.

Proceedings of the National Academy of Sciences of the United States of America, *111*(28), 10341–10346. Retrieved from

www.pnas.org/cgi/doi/10.1073/pnas.1400181111 doi: 10.1073/pnas.1400181111

Zamora-López. (2010). Cortical hubs form a module for multisensory integration on top of the hierarchy of cortical

networks. *Frontiers in Neuroinformatics*, *4*(MAR), 1. Retrieved from

http://journal.frontiersin.org/article/10.3389/neuro.11.001.2010/abstract doi:

10.3389/neuro.11.001.2010

—47—

1136   Zeki, S., & Shipp, S. (1988). The functional logic of cortical connections. *Nature*, *335*(6188), 311–317. Retrieved from

1137   https://pubmed.ncbi.nlm.nih.gov/3047584/ doi: 10.1038/335311a0