A two-hit epistasis model prevents core genome disharmony in recombining bacteria

Aidan J. Taylor[1], Koji Yahara[2], Ben Pascoe[3], Leonardos Mageiros[4,13], Evangelos Mourkas[3], Jessica K Calland[5], Santeri Puranen[6], Matthew D. Hitchings[4], Keith A. Jolley[3], Carolin M. Kobras[7], Sion Bayliss[8], Nicola J. Williams[9], Arnoud H. M. van Vliet[10], Julian Parkhill[11], Martin C. J. Maiden[3], Jukka Corander[6,7,12], Laurence D Hurst[13], Daniel Falush[14], Paul Keim[15], Xavier Didelot[16], David J. Kelly[17*], Samuel K. Sheppard[3*]

Affiliations

[1]University of Reading, Whiteknights, Reading, RG6 6AH, UK

[2]Antimicrobial Resistance Research Center, National Institute of Infectious Diseases, Tokyo, Japan.

[3]Department of Biology, University of Oxford, South Parks Road, Oxford, OX1 3RE. UK

[4]Swansea University Medical School, Institute of Life Science, Swansea SA2 8PP, UK

[5]Department of Biostatistics, University of Oslo, Oslo, Norway

[6]Department of Mathematics and Statistics, Helsinki Institute for Information Technology, University of Helsinki, Helsinki, Finland

[7]Sir William Dunn School of Pathology, University of Oxford, Oxford, OX1 3RE, UK

[8]Langford House, University of Bristol, Langford. BS405DU

[9]Department of Epidemiology and Population Health, Institute of Infection & Global Health, University of Liverpool, Leahurst Campus, Wirral, UK

[10]School of Veterinary Medicine, University of Surrey, Guildford GU2 7AL, UK

[11]Department of Veterinary Medicine, University of Cambridge, Cambridge CB3 0ES, UK

[12]Parasites and Microbes, Wellcome Sanger Institute, Cambridge, UK

[13]The Department of Biology and Biochemistry, University of Bath, UK

[14]The Centre for Microbes, Development and Health, Institut Pasteur of Shanghai, Shanghai,

China

[15]The Pathogen and Microbiome Institute, Northern Arizona University, Flagstaff, Arizona,

United States of America, Department of Biological Sciences, Northern Arizona University,

Flagstaff, Arizona, United States of America

[16]School of Life Sciences and Department of Statistics, University of Warwick, Coventry CV4

7AL, UK.

[17]School of Biosciences, University of Sheffield, Sheffield S10 2TN, United Kingdom;

*Corresponding author(s): David J. Kelly, d.kelly@sheffield.ac.uk; Samuel K. Sheppard,

s.k.sheppard@bath.ac.uk;

Significance Statement

Genetic exchange among bacteria shapes the microbial world. From the acquisition of antimicrobial resistance genes to fundamental questions about the nature of bacterial species, this powerful evolutionary force has preoccupied scientists for decades. However, the mixing of genes between species rests on a paradox. On one hand, promoting adaptation by conferring novel functionality, on the other potentially introducing disharmonious gene combinations (negative epistasis) that will be selected against. Taking an interdisciplinary approach to analyse natural populations of the enteric bacteria *Campylobacter*, an ideal example of long-range admixture, we demonstrate that genes can independently transfer across species boundaries and rejoin in epistasis in a recipient genome. This challenges conventional ideas and highlights the possibility of single-step evolution by saltation.

Abstract

Recombination of short DNA fragments via horizontal gene transfer (HGT) can both introduce beneficial alleles and create genomic disharmony through negative epistasis. For non-core (accessory) genes, the negative epistatic cost is likely to be minimal because the incoming genes have not co-evolved with the recipient genome. By contrast, for the core genome, interspecific recombination is expected to be rare because disruptive allelic replacement is likely to introduce negative epistasis. Why then is homologous recombination common in the core of bacterial genomes? To understand this enigma we take advantage of an exceptional model system, the common enteric pathogens *Campylobacter jejuni* and *Campylobacter coli*, that are known for very high magnitude interspecies gene flow in the core genome. As expected, HGT does indeed disrupt co-adapted allele pairings (negative epistasis). However, multiple HGT events enable recovery of

the genome's co-adaption between introgressing alleles, even in core metabolism genes (e.g.,

formate dehydrogenase). These findings demonstrate that, even for complex traits, genetic

coalitions can be decoupled, transferred and independently reinstated in a new genetic background

– facilitating transition between fitness peaks. In this example, the two-step recombinational

process is associated with *C. coli* that are adapted to the agricultural niche.

Introduction

Mutation is the engine of genetic novelty, but for most bacteria adaptation also involves the

acquisition of DNA from other strains and species through horizontal gene transfer (HGT) [1]. In

some well documented cases, a single nucleotide substitution or acquisition of a small number of

genes, can prompt new evolutionary trajectories with striking outcomes such as the evolution of

virulent or antibiotic resistant strains [2]. With such dynamic genomic architecture, it may be

tempting (and possibly useful [3]) to consider genes as independent units that 'plug and play'

innovation into recipient genomes. This is clearly an oversimplification. In fact, genes within

genomes are highly interactive wherein the effect of one allele depends on another (epistasis).

Therefore, it is likely that some horizontally introduced changes will disrupt gene networks and be

costly to the original coadapted genetic background, especially for complex phenotypes involving

multiple genes and more distant taxonomic relationships.

Understanding how epistasis influences the evolution of phenotype diversity has preoccupied

researchers since the origin of population genetics [4-10], with much emphasis placed upon the

relative amounts of recombination and epistatic effect sizes [11, 12]. In sexual populations, such

as outbreeding metazoans, genetic variation is shuffled at each generation. This the trend towards randomization between alleles (linkage equilibrium), means it is unlikely that multiple distinct epistatic allele combinations will be maintained in the same population [8]. In bacteria, however, rapid clonal reproduction allows multiple genomically distant beneficial allele combinations to rise to high frequency or even fixed in a single population. For example, in common enteric bacteria such as *Escherichia coli, Salmonella enterica* and *Campylobacter jejuni,* the doubling time in the wild has been estimated at around 24 hours or less [13, 14]. Therefore, though HGT occurs in these organisms [15], even in highly recombinogenic *C. jejuni* [13, 16], there will likely be many millions of bacterial generations between recombination events at a given locus. This allows mutations that are beneficial only in specific genetic backgrounds to establish in a single population and linkage disequilibrium to form between different epistatic pairs [17].

In a coadapted genomic landscape, recombination is expected to have two antagonistic effects. On one hand it could be beneficial, promoting adaptation by conferring novel functionality on the recipient genome [18-22] and though specialization reduce the competition between clones (clonal interference). On the other it could be detrimental, introducing disharmonious allele-combinations that will be discriminated against by selection [19]. The balance of these two effects is likely to be different for core and non-core HGT events. Non-core events, such as introduced accessory antibiotic resistance genes, can be immediately beneficial and, as they do not replace recipient sequence, need not break established epistatic interactions. By contrast, HGT replacing one allele for another (homologous recombination) in part of the core genome is likely to disrupt highly evolved co-adapted networks. For these reasons, negative epistatic interactions between core genes with different evolutionary histories have been proposed as a barrier to recombination [4-10, 23],

particularly between species. However, interspecies recombination is common in bacterial core

genomes [18, 24]. How then can a core genome, that is expected to build up extensive co-adapted

epistatic networks, be so accepting of HGT events?

The common animal gut bacterium *Campylobacter,* which is among the most prolific causes of

human bacterial gastroenteritis worldwide [25], provides an exceptional model system to study the

impact of HGT on the core genome for several reasons. First, introgression between the two most

important pathogenic species, *C. jejuni* and *C. coli,* has led to the evolution of a globally distributed

'hybrid' *C. coli* lineage [26] that is responsible for almost all livestock and human infections with

this species. The hybrids occupy a niche that was not available to the parental subtypes. Second,

up to 23% of the core genome of one common *C. coli* subtype has been recently introgressed from

C. jejuni [27], potentially disrupting epistatic interactions. Finally, because *C. jejuni* and *C. coli*

have undergone an extended period of independent evolution with only 85% average nucleotide

identity, recombined sequence is conspicuous in the genome.

Here, we investigate the disruptive effect of HGT on co-adapted bacterial core genomes by

examining covarying allele pairings and imported DNA sequence. Even though recombined

fragments enter the genome one-by-one, we find that most covariation in the core genome is

between sites where both alleles were imported. Having confirmed that allelic covariation is

indicative of epistasis, with laboratory mutagenesis and complementation assays, we conclude that

independent disruptive recombination events occur and persist until a second event restores the

functional link in a new genetic background. This process resembles the two-hit cancer model

where an initial mutation is actuated by a second [48]. Both bacterial and cancer cell lines are

asexual clones and a two-hit model for core genome HGT is consistent with a more general theory

in which negative effects from an initial event (MGT or mutation) need not preclude adaptive

evolution. As in cancer, the recombinant genotypes are under strong selection and occupy a unique

agricultural niche.

Results

***Campylobacter* populations are highly structured with intermediate sequence clusters**

Analysis of every automatically annotated gene from every genome, revealed a core genome of

631 gene orthologues in all *Campylobacter* isolates in this study. There were 1287 genes common

to all *C. jejuni* isolates and 895, 1021 and 1272 common to *C. coli* clades 1, 2 and 3, respectively.

Consistent with previous studies [27], neighbour-joining and ClonalFrameML trees based on

genes within concatenated core genome alignments revealed population structure in which *C.*

jejuni and *C. coli* clade 2 and 3 isolates each formed discrete clusters (Figure 1A, Figure S1).

Isolates designated as *C. coli* clade 1 were found in three clusters on the phylogeny: unintrogressed

ancestral strains, and the ST-828 and ST-1150 clonal complexes (CCs) which account for the great

majority of strains found in agriculture and human disease [27]. Pan-genome analysis quantified

the increase in unique gene discovery as the number of sampled genomes increased (Figure S2A).

For all phylogenetic clusters there was evidence of an open pan-genome with a trend towards

continued rapid gene discovery within clusters containing fewer isolates. There was considerable

accessory genome variation between species and clades, potentially associated with important

adaptive traits (Figure S3) and there was evidence that the average number of genes per genome

was greater in *C. coli* CC-828 isolates than in *C. jejuni* (Figure S2B).

There is substantial introgression within the *C. coli* core genome

The large genetic distances among *C. coli* clade 1 isolates (Figure 1A), have been shown to be a

consequence of the import of DNA from *C. jejuni* rather than accumulation of mutations during a

prolonged period of separate evolution [26]. Using chromosome painting to infer the co-ancestry

of core-genome haplotype data from CC-828 and CC-1150 isolates gave a detailed representation

of the recombination-derived segments from each *C. jejuni* donor group and clonal complex to

each recipient individual (Figure S4). The majority of introgressed SNPs were rare, occurring in

fewer than 50 recipient genomes (Figure 1B). However, a large proportion of the introgressed *C.*

coli clade 1 genomes contained DNA of *C. jejuni* ancestry in >98% of recipient isolates. Consistent

with previous estimates [27], these regions where introgressed DNA was largely fixed within the

C. coli population, occurred across the genome and comprised up to 15% and 28% of the CC-828

and CC-1150 isolate genomes, respectively (Figure 1B). After considering potential donor groups,

we find that the majority of introgressed DNA in *C. coli* involves genes that are present in multiple

C. jejuni lineages (core genes) (Figure S5A).

Having identified *C. jejuni* ancestry within *C. coli* genomes, we investigated the sequence of events

responsible for introgression. Most introgressed SNPs are found at low frequency in both CC-828

and CC-1150 clonal complexes. However, there was evidence of SNPs that are introgressed in

both complexes as well as high frequency lineage specific introgression in both CC-828 and CC-

1150 (Figure S5B). Specifically, 25% of the *C. jejuni* DNA found in >98% of CC-828 isolates was

also found in CC-1150 (Figure S5C), implying that this genetic material was imported by the

common ancestor(s) of both complexes. After the divergence of these two complexes,

introgression continued with nearly 75% of *C. jejuni* DNA present in one recipient clonal complex

and not the other. This stepwise model is consistent with an evolutionary history in which there

was a period of progressive species and clade divergence reaching approximately 12% at the

nucleotide level between *C. jejuni* and *C. coli* and around 4% between the three *C. coli* lineages.

More recently, changes to the patterns of gene flow led one *C. coli* clade 1 lineage to import

substantial quantities of *C. jejuni* DNA, and further lineage-specific introgression gave rise to two

clonal complexes (CC-828 and CC-1150) that continued to accumulate *C. jejuni* DNA,

independently creating the population structure observed today (Figure 1C).

The high magnitude introgression into *C. coli* clade 1 isolates has introduced thousands of

nucleotide changes to the core genome. However, divergence in bacteria may be uneven across

the genome for at least two reasons. First, recombination is more likely to occur in regions where

donor and recipient genomes have high nucleotide similarity [24, 28, 29]. Second, 'fragmented

speciation' [30], in which gene flow varies in different parts of the genome, such as regions

responsible for adaptive divergence, can result in phylogenetic incongruence among genes.

Consistent with previous estimates [27], we found that the three *C. coli* clades had similar high

divergence with *C. jejuni* across the genome, ranging from 68% to 98% nucleotide identity for

individual genes (Figure S5D), implying a period of divergence with low levels of gene flow. We

found no evidence that high genetic differentiation between the species prevented recombination.

While there was some evidence that more recombination occurred in regions of low nucleotide

divergence (between unintrogressed *C. coli* clade 1 and *C. jejuni*), introgression occurred across

the genome at sites with a wide range of nucleotide identity (Figure S5D). This level of

recombination has greatly increased overall genetic diversity across the genome in *C. coli* clade 1

and introduced changes that have potential functional significance.

Much of the putative epistasis occurs between SNPs in introgressed genes

ClonalFrameML analysis revealed the importance of homologous recombination in generating

sequence variation within the introgressed *C. coli*. Estimates of the relative frequency of

recombination versus mutation ($R/\theta=0.43$), mean recombination event length ($\delta=152$bp) and

average amount of polymorphism per site in recombined fragments ($\nu=0.07$), imply that

recombination has had an effect (*r/m*) 4.57 times higher than *de novo* mutation during the

diversification of CC-828. This is consistent with previous analysis and confirmed recombination

as the major driver of molecular evolution in *C. coli* [13, 27, 31]. The continuous time Markov

chain model for the joint evolution of pairs of biallelic sites on a phylogenetic tree (Figure S6) was

applied to investigate patterns of covariance for all pairs of sites >20kb apart (Figure S7). For most

biallelic sites there were few branches on the tree where substitutions occurred, so that their

evolution is compatible with separate evolution on the same clonal frame. However, 2874

covarying pairs evolved more frequently together than would be expected (*p*-value 10^{-8}) if they

had evolved independently based on the tree, and hence indicated patterns of putative epistasis.

Among them, the location of 2618 putative epistatic pairs of sites was compared to the inferred

ancestry (unintrogressed *C. coli* or *C. jejuni*) of sequence across the genome of CC-828 and CC-

1150 *C. coli* strains (Figure 2C, Data S1). Putative epistatic pairs in the recipient introgressed

genome represent a combination of the haplotypes in the ancestral populations (Figure 2A). For

each epistatic pair, the major and minor haplotype were defined, from these possible combinations,

if there was haplotype polymorphism between *C. jejuni* and *C. coli.* This allowed quantification

of the number of covarying sites that occurred between an ancestral *C. coli* (unintrogressed) and

an introgressed *C. jejuni* allele (*C. coli – C. jejuni*), two introgressed alleles (*C. jejuni – C. jejuni*),

and sites that do not segregate by species. Strikingly, the breakdown of the major and minor

haplotype combinations among the 2618 epistatic pairs (Figure 2B, Data S2) shows the major

haplotype for 83.5% of putative epistatic SNP pairs was *C. jejuni* indicating that both co-varying

sites had *C. jejuni* ancestry, consistent with epistasis between introgressed ancestral *C. jejuni*

sequence at divergent genomic positions. Investigation of the genes containing co-varying sites

revealed that 2187 SNP pairs were in 16 genes with just five genes accounting for 99.1% of them

(Figure 3A&B, Data S3, Figure S8).

Genomic context and physiological role of epistatically linked genes

The five genes accounting for the majority of epistatic interactions (*cj1167*, *cj1168c*, *cj1171c/ppi*,

cj1507c/modE and *cj1508c/fdhD*) were investigated for their physiological role in *C. jejuni*. FdhD

and ModE are proteins involved in the biogenesis of formate dehydrogenase (FDH). The FDH

complex (FdhABC) oxidises formate to bicarbonate to generate electrons that fuel cellular

respiration. Formate is an abundant electron donor produced by host microbiota and an important

energy source for *Campylobacter in vivo* [32, 33]. The remaining three genes, *cj1167* (annotated

incorrectly as *ldh*, lactate dehydrogenase), *cj1168c* and *cj1171c* (*ppi*) are also grouped together in

the genome, where *cj1167* and *cj1168c* are adjacent but with the open reading frames (ORFs)

convergent and overlapping, while *ppi* is upstream, separated by two non-epistatically linked genes

(*cj1169c* and *cj1170c*, Figure 3B). Considering the genomic arrangement, it is therefore clear that

the putative epistatic links uncovered in this study essentially occur across two loci in the genome

(*fdhD*/*modE* and *cj1167*/*cj1168c*/*ppi*), with each of the latter three genes linked with both *fdhD*

and *modE* (Figure 3B). Given the known function of *fdhD* and *modE* in biogenesis of the FDH

complex, we hypothesised that *cj1167*/*cj1168c*/*ppi* might also have some role in FDH biogenesis

or activity in order to form a functional epistatic connection. We therefore constructed deletion

mutants to investigate the possible role of these genes in FDH activity.

Initially, each of the mutants and their parental wildtype (*C. jejuni* NCTC11168) were grown in

rich media (Muller-Hinton broth) and their formate dependent oxygen consumption rates

determined (Figure 3C). *cj1167, cj1168c* and *ppi* mutants demonstrated wildtype levels of FDH

activity, while activity in both *fdhD* and *modE* mutants was abolished. To confirm that the

phenotype of the *fdhD* and *modE* mutants was not due to a polar effect on the surrounding *fdh*

locus, these mutants were genetically complemented by reintroduction of a second copy of the

wildtype gene into the rRNA locus, which restored near wildtype levels of FDH activity in both

cases (Figure 3C).

As neither *cj1167, cj1168c* or *ppi* mutants showed altered FDH activity in cells grown in rich

media, we considered that their function may be related to an FDH-specific nutrient requirement

as would likely be found *in vivo*. Since the formate oxidising subunit of FDH, FdhA, specifically

requires a molybdo- or tungsto-pterin (Mo/W) cofactor and a selenocysteine (SeC) residue for

catalysis [34], Mo, W or Se supply presented possible targets. *cj1168c* encodes a DedA family

integral membrane protein of unknown function. DedA proteins are solute transporters widespread

in bacteria but are mostly uncharacterized [35]. However, a homologue of *cj1168c* in the heavy

metal specialist beta-proteobacterium *Cupriavidus metallidurans* has been shown to be involved

in selenite (SeO_3^{2-}) uptake [36]. We therefore speculated that Cj1168 could be a selenium oxyanion

transporter that supplies Se for SeC biosynthesis. To test this, FDH activities were measured in

cj1168c mutant and parental wildtype strains grown in minimal media with limiting concentrations

of selenite or selenate (SeO_4^{2-}) (Figure S10). The data in Figure 3D shows that the *cj1168c* mutant

displayed significantly reduced FDH activity after growth with selenite in the low nM range, and

this phenotype was partially restored by genetic complementation. We therefore designated

cj1168c as *selF* (<u>se</u>lenium transporter for <u>f</u>ormate dehydrogenase). However, although this

phenotype does suggest that SelF is a selenium importer, another unrelated selenium transporter,

FdhT (Cj1500), has previously been documented in *C. jejuni* [37], which is not epistatic with *fdhD*

or *modE*. In contrast to this previous report, we found considerable residual FDH activity still

remained in an *fdhT* deletion mutant, which was fully restored to wildtype levels by

complementation (Figure S9).

Finally, we tested whether the residual FDH activity in our *fdhT* mutant was due to selenium uptake

by SelF. An *fdhT selF* double mutant was generated and assayed for FDH activity after growth in

minimal media containing limiting concentrations of selenite or selenate (Figure S9). The *fdhT*

selF double mutant demonstrated a significant additional reduction in FDH activity over the *fdhT*

single mutant, a phenotype that was partially restored by complementing the double mutant with

selF. Complementation of the double mutant with *fdhT* returned FDH activity to near wildtype

levels (Figure S9). Taken together, our data suggests that both FdhT and SelF facilitate selenium

acquisition in *C. jejuni*, possibly representing low and high affinity transporters, respectively

(Figure S11).

Discussion

To address the apparent contradiction of frequent core genome recombination in a co-adapted

genomic background we focused on *Campylobacter,* in which interspecies recombination is well

documented [26, 27, 38]. As in other studies, we found that a large proportion (15-28%) of the *C.*

coli core genome originated in *C. jejuni* despite the genetic distance (~85% nucleotide identity)

between the species. Investigating the likely disruptive impact this would have on coadapted

epistatic gene networks, we quantified the frequency of *C. jejuni – C. coli* (and *vice versa*) and *C.*

jejuni – C. jejuni covarying allele pairs in introgressed *C. coli*. Where recombination is minimally

disruptive there would be more *C. jejuni – C. coli* than *C. jejuni – C. jejuni*. However, consistent

with selection against disharmonious gene combinations we found that *C. jejuni – C. jejuni* allele

pairs constituted >83% of covarying introgressed haplotypes. It is possible that in some cases both

sites were introgressed in a single large recombination event [39-41] but in *Campylobacter* LD for

pairs of sites decreases with distance, and is approximately constant after 20kbp [42] consistent

with the independent acquisition of alleles that are >20kb apart. It follows, therefore, that the first

introgression event was not fatal to the recipient genome. Acquisition of the second member of the

pair then potentially restored the function of the integrated *C. jejuni – C. jejuni* coevolutionary

unit.

Statistical genetic analysis confirmed covariation, but to confirm coadaptation (*sensu stricto)* it

was necessary to confirm epistasis in the laboratory. The most common co-varying *C. jejuni – C.*

jejuni gene pairs were linked to FDH, a key enzyme allowing the utilisation of formate as an

electron donor *in vivo* [32, 33]. FdhD and ModE were shown to be essential for FDH activity.

While FdhD is a known sulfur-transferase required for cofactor insertion into FdhA [43] (Figure

S11), the abolished FDH activity in a *modE* mutant indicates functions for ModE in FDH

biogenesis, beyond the known role as a transcriptional repressor [44, 45]. In contrast, *cj1167, self*

and *ppi* mutants did not show reduced FDH activity. This apparently contradicted the epistasis

hypothesis until we considered the functional links between *fdhD*/*modE* and *cj1167*/*selF*/*ppi*. This

revealed that a *selF* mutant strain had significantly reduced FDH activity under conditions of

selenite limitation (Figure 3D). This phenotype is consistent with SelF being a Se oxyanion

transporter and functionally links SelF with FdhD/ModE. We suggest that *selF* rather than *fdhT* is

epistatic because SelF confers an additional benefit (SeC biosynthesis, essential for FDH activity)

under conditions of selenium limitation, for example as may be found in the host gut (Figure S11).

Direct functional association with FDH activity was more difficult to explain for *cj1167* and *ppi*.

cj1167 is currently incorrectly annotated as lactate dehydrogenase (Ldh) and it's actual function is

unknown [46]. While we found no evidence for a functional connection between Cj1167 and FDH

activity, the overlapping convergent gene arrangement of *cj1167* and *cj1168c* (*selF*) suggests a

transcriptional architecture that might form similar epistatic dependencies even if Cj1167 is not

required for FDH activity. Finally, the *cj1171c* (*ppi*) deletion mutant showed no growth defect or

reduction in FDH activity. *ppi* encodes a cytoplasmic peptidyl-prolyl *cis-trans* isomerase and

PPIases are general protein folding catalysts that often have pleiotropic and redundant functions

[47]. Therefore, it is possible that if Cj1171 does help promote the folding of FdhD or ModE but

analysis of a simple deletion mutant may not reveal this if another PPIase can substitute in that

genetic background.

Understanding the functional significance of core genome recombination has considerable

potential to explain the evolution of complex phenotypes. In *Campylobacter,* our results suggest

that an ancestral *C. coli* lineage colonized a new niche and surviving lineages (CC-828 and CC-

1150) gained access to *C. jejuni* DNA (Figure 4A&B). In this case, introgressed *C. coli* acquired

multiple genes allowing them to utilize the high levels of formate in the host gut as an energy

source. Specifically, the *fdh* and *ModE,* essential for FDH biogenesis, as well as *selF* that ensures

sufficient selenium for the enzyme to function. This example shows how as the adaptive landscape

of the genome changed, potentially decoupling epistatic interactions that were previously selected,

new gene combinations can be introduced by homologous recombination and tested in the *C. coli*

genetic background. These new recombinant genotypes are frequently isolated in the agricultural

setting and from clinical cases caused by agricultural products, arguing for their enhanced fitness

in this niche.

The notion that multiple events are necessary to achieve a phenotype with the first event potentially

being deleterious (causing negative epistatsis) is strongly reminiscent of the two-hit cancer model

in eukaryotes [48] in which the benefit to the tumour cell appears only after earlier non-

advantageous mutations. In bacteria, while the first hit (HGT event) in a core genome, is consistent

with fitness costs associated with negative epistasis (breaking co-adapted gene networks), this can

be rescued by a second HGT event. The second event, in effect restores a pre-existing co-adapted

allele pair from the donor species to the recipient. This type of genetic rewiring may indeed be

more common than previously thought [49-51] and, despite theoretical expectations, negative

epistasis is not an absolute barrier to genome-wide recombination in structured bacterial

populations. Multiple HGT events thus provide a solution to the classic evolutionary fitness peak

jumping problem.

Materials and Methods

Isolates, genome sequencing and assembly

A total of 973 isolates were used in this study, 827 from *C. coli* and a selection of 146 from a

diversity of *C. jejuni* clonal complexes (Data S4). Isolates were sampled mostly in the United

Kingdom to maximise environmental and riparian reservoirs and thus the representation of genetic

diversity in *C. coli*. Isolates were stored in a 20% (v/v) glycerol medium mix at -80°C and

subcultured onto *Campylobacter* selective blood-free agar (mCCDA, CM0739, Oxoid). Plates

were incubated at 42°C for 48 h under microaerobic conditions (5% CO_2, 5% O_2) generated using

a CampyGen (CN0025, Oxoid) sachet in a sealed container. Subsequent phenotype assays were

performed on Brucella agar (CM0271, Oxoid). Colonies were picked onto fresh plates and

genomic DNA extraction was carried out using the QIAamp® DNA Mini Kit (QIAGEN; cat.

number: 51306) according to the manufacturer's instructions. DNA was eluted in 100–200 µl of

the supplied buffer and stored at −20°C. DNA was quantified using a Nanodrop spectrophotometer

and high-throughput genome sequencing was performed on a MiSeq (Illumina, San Diego, CA,

USA), using the Nextera XT Library Preparation Kit with standard protocols involving

fragmentation of 2 µg genomic DNA by acoustic shearing to enrich for 600 bp fragments, A-

tailing, adapter ligation and an overlap extension PCR using the Illumina 3 primer set to introduce

specific tag sequences between the sequencing and flow cell binding sites of the Illumina adapter.

DNA cleanup was carried out after each step to remove DNA < 150 bp using a 1:1 ratio of

AMPure® paramagnetic beads (Beckman Coulter, Inc., USA). Short read paired-end data was

assembled using the *de novo* assembly algorithm, SPAdes (version 3.10.0) [52]. All novel genome

sequences (n=475) generated for use in this study are available on NCBI BioProjects

PRJNA689604 and PRJEB11972. These were augmented with 498 previously published genomes

and accession numbers for all genomes can be found in Data S4 [16, 27, 31, 42, 53-57].

Genome archiving, pan-genome content analyses and phylogenetic reconstruction

Contiguous genome sequence assemblies were individually archived on the web-based database

platform PubMLST [58] and sequence type (ST) and clonal complex (CC) designation were

assigned based upon the *C. jejuni* and *C. coli* multi-locus sequence typing scheme [59]. To examine

the full pan-genome content of the dataset, a reference pan-genome list was assembled as

previously described [60]. Briefly, genome assemblies from all 973 genomes in this study were

automatically annotated using the RAST/SEED platform[61], the BLAST algorithm was used to

determine whether coding sequences from this list were allelic variants of one another or 'unique'

genes, with two alleles of the same gene being defined as sharing >70% sequence identity on >10%

of the sequence length. The prevalence of each gene in the collection of 973 genomes was

determined using BLAST with a positive hit in a genome being defined as a local alignment of the

reference sequence with the genomic sequence of >70% identity on >50% of the length, as

previously described [62]. The resulting matrix was analysed for differentiating core and accessory

genome variation. Genes present in all genomes were concatenated to produce a core-genome

alignment, used for subsequent phylogenetic reconstructions. Phylogenetic trees were

reconstructed using an approximation of maximum-likelihood phylogenetics in FastTree2 [63].

This tree was used as an input for ClonalFrameML [64] to produce core genome phylogenies with

branch lengths corrected for recombination.

Inference of introgression

All 973 genomes were aligned to a full reference sequence of *C. coli* strain CVM29710. We

conducted imputation for polymorphic sites with missing frequency $\leq$10% using BEAGLE [65] as

previously reported [66]. A total of 286,393 gapless SNPs (~17% of the average *C. coli* genome

size) were used for recombination analyses. The coancestry of genome-wide haplotype data was

inferred based on alignments using chromosome painting and FineStructure [67] as previously

described[68]. Briefly, ChromoPainter was used to infer chunks of DNA donated from a list of 33

donor groups normalized for sample size to each of 677 ST-CC-828 and 12 CC-1150 recipient

haplotypes. Results were summarized into a coancestry matrix containing the number of

recombination-derived chunks from each donor to each recipient individual. FineStructure was

then used for 100,000 iterations of both the burn-in and Markov chain Monte Carlo chain to cluster

individuals based on the co-ancestry matrix. The results are visualized as a heat map with each cell

indicating the proportion of DNA "chunks" (a series of SNPs with the same expected donor) a

recipient receives from each donor.

Analysis of covariation in bacterial genomes

Non-random allele associations can result from selection and clonal population structure. To

control for the latter, our approach identified SNP combinations in independent genetic

backgrounds by accounting for the sequence variation associated with the inferred phylogeny.

Based on the alignment of 677 genomes of *C. coli* CC-828, a first phylogenetic tree was created

using PhyML [69]. ClonalFrameML [64] was then applied to correct the tree by accounting for

the effect of recombination, and also to infer the ancestral sequence of each node. Covariance was

assessed for pairs of biallelic sites across the genome using a Continuous Time Markov Chain

(CTMC) model as follows. Briefly, let A and a denote the two alleles of the first site and B and b

denote the two alleles of the second site, so that there are four states in total for the pair of sites

(ab, Ab, aB and AB). The four substitution rates from A to a, from a to A, from B to b and from b

to B are not assumed to be identical, to allow for differences in substitution rates in different parts

of the genome and also to allow for non-equal rates of forward and backward substitution (for

example as a result of recombination opportunities). Assuming no epistatic effect between the two

sites ($\varepsilon=1$), the model M_0 has four free parameters (α_1, α_2, β_1 and β_2) representing independent

substitutions at the two sites. We expand model M_0 with an additional fifth parameter $\varepsilon>1$ into

model M_1 which is such that the state AB where the first site is allele A and the second site is allele

B is favored relative to the other three sites $ab,$ aB and Ab. Specifically, the state AB has a

probability increased by a factor ε^2 in the stationary distribution of the CTMC of model M_1

compared to model M_0.

Both models M_0 (with 4 parameters) and M_1 (with 5 parameters) are fitted to the data using

maximum likelihood techniques, where the likelihood is equal to the product for every branch of

the tree of the state at the bottom of the branch given the state at the top. The two fitted models M_0

and M_1 are then compared using a likelihood-ratio test (LRT) as follows: since M_0 is nested with

M_1, two times their difference in log-likelihood is expected to be distributed according to a chi-

square distribution with number of degrees of freedom equal to the difference in their

dimensionality, which is one. This LRT returns a p-value for the significance of a covariation

effect, and a Bonferroni correction is applied to determine a conservative cutoff of significance

that accounts for multiple testing. Furthermore, the test is applied only to pairs of sites separated

by >20kb to reduce the chance that they were the result of a single recombination event, consistent

with estimates of the length of recombined DNA sequence in quantitative bacterial transformation

experiments [39, 70] and evidence from *Campylobacter* genome analyses that show that LD for

pairs of sites decreases with distance to approximately 20kbp and then remains at the same level

for very distant sites [42]. It is still possible of course that rare recombination events would stretch

20kbp [39-41], but for this to have an effect on the analysis of epistasis it would have to have

happened several times for the same pairs of sites against different genomic background which

becomes quite unlikely just by chance. This phylogenetically aware approach to testing for

covariance presents the advantage to naturally account for both population structure and the effect

of recombination [71]. The script implementing this coevolution test is available in R at:

https://github.com/xavierdidelot/campy.

Quantifying covariation between recombined and unrecombined genomic regions

The results of the introgression and covariation analyses were combined so that for each pair of

significantly covarying SNPs (*p*-value $<10^{-8}$), haplotype frequency was calculated among the 689

recipient introgressed *C. coli* clade-1 strains as well as among the donor *C. coli* (ancestral) and *C.*

jejuni strains, respectively. If the most frequent haplotype of the pair is the same between the donor

C. coli (ancestral) and *C. jejuni*, it was classified as 'no polymorphism'. Otherwise, if the most

frequent haplotype accounted for >90% among the recipients, it was classified as either '*C. jejuni*

(>90%)' or '*C. coli* (>90%)' if it was the same as that of donor *C. jejuni* or *C. coli* (ancestral) (inset

in Figure 2C). If the most frequent haplotype accounted for ≤90% among the recipients, the top

two most frequent haplotypes (written as major and minor haplotype in this manuscript) were

indicated as either "*C. jejuni* / *C. coli*", "*C. jejuni* / other", "*C. coli* / *C. jejuni*", "*C. coli* / other",

"other / *C. coli*", "other / *C. jejuni*", and "other / other", and the frequency of the major and minor

haplotypes were calculated. For example, where the haplotype frequencies were as follows,

AA=285, TA=192, TG=181, AG=27, A-=2, --=1, -A=1, AA is the major haplotype, frequency of

which is 41.3%

Mutagenesis and complementation cloning

Genes *cj1167*, *cj1168c* (here designated *selF* for <u>sel</u>enium transport for <u>f</u>ormate dehydrogenase),

cj1171c (*ppi*), *cj1507c* (*modE*), *cj1508c* (*fdhD*) and *cj1500* (*fdhT*) were deleted by allelic exchange

mutagenesis, with the majority of the open reading frame replaced by an antibiotic resistance

cassette. Mutagenesis plasmids were generated by the isothermal assembly method using the HiFi

system (NEB, UK). In brief, flanking regions of target genes were PCR amplified from genomic

DNA using primers with adaptors homologous to either the backbone vector pGEM3ZF or the

antibiotic resistance cassette (Data S5). pGEM3ZF was linearised by digestion with HincII. The

kanamycin and chloramphenicol resistance cassettes were PCR amplified from pJMK30 and

pAV35, respectively [72]. Four fragments consisting of linearised pGEM3ZF, antibiotic resistance

cassette and 2 flanking regions were combined in equimolar amounts and mixed with 2 x HiFi

reagent (NEB, UK) and incubated at 50°C for 1 hour. The fragments combine such that the gene

fragments flank the antibiotic resistance cassette, in the same transcriptional orientation, within

the vector. Mutagenesis plasmids were transformed into *C. jejuni* NCTC 11168 by electroporation.

Spontaneous double-crossover recombinants were selected for using the appropriate antibiotic and

correct insertion into the target gene confirmed by PCR screening. For genetic complementation

of mutants, genes *cj1168c* (*selF*), *cj1507c* (*modE*), *cj1508c* (*fdhD*) and *cj1500* (*fdhT*) were PCR

amplified from genomic DNA, restriction digested with MfeI and XbaI, then ligated into similarly

digested pRRA [73] (Data S5). The orientation of insertion allowed the target gene to be expressed

constitutively by a chloramphenicol resistance gene-derived promoter within the vector.

Complementation plasmids were transformed into *C. jejuni* by electroporation. Spontaneous

double-crossover recombinants were selected for using apramycin and correct insertion into the

ribosomal locus confirmed by PCR screening.

Growth of *C. jejuni*

Microaerobic growth cabinets (Don Whitley, UK) were maintained at 42°C with an atmosphere

of 10% O_2, 5% CO_2 and 85% N_2 (v/v). *C. jejuni* was grown on Columbia-base agar containing 5%

v/v defibrinated horse blood. Selective antibiotics were added to plates as appropriate at the

following concentrations: 50 µg ml^{-1} kanamycin, 20 µg ml^{-1} chloramphenicol, 60 µg ml^{-1}

apramycin. Muller-Hinton (MH) broth supplemented with 20 mM L-serine was used as a rich

medium. Minimal medium was prepared from a supplied MEM base (51200-38, Thermo

Scientific, UK) with the following additions: 20 mM L-serine, 0.5 mM sodium pyruvate, 50 µM

sodium metabisulfite, 4 mM L-cysteine. HCl, 2 mM L-methionine, 5 mM L-glutamine, 50 µM

ferrous sulfate, 100 µM ascorbic acid, 1 µM vitamin B12, 5 µM sodium molybdate, 1 µM sodium

tungstate. Selenium was then added as appropriate from stocks of sodium selenate or sodium

selenite prepared in dH_2O. For assays, cells were washed and suspended in sterile phosphate-

buffered saline (PBS, Sigma-Aldrich).

Respiration rates with formate

Cells were first grown in MH broth for 12 hours, then washed thoroughly in PBS before inoculating minimal media without an added selenium source. The appropriate concentrations were determined by serial dilution trials, and it was subsequently found that *C. jejuni* has a strong preference for selenite over selenate, as equivalent FDH activity requires some 1000-fold greater concentration of selenate than selenite (Figure S10). These cultures were grown for 8 hours before the cells were thoroughly washed again, then used to inoculate further minimal media, with a selenium source added as appropriate, and grown for 10 hours. This passaging was necessary to remove all traces of selenium from the inoculum, such that control cultures without selenium added had negligible (FDH) activity. Assay cultures were again thoroughly washed before the equivalent of 20 ml at an optical density of 0.8 at 600 nm was finally suspended in 1 ml of PBS. Formate-dependent oxygen consumption by whole cells was measured in a Clark-type electrode using 20 mM sodium formate as electron donor. The electrode was calibrated with air-saturated PBS assuming 220 nmol dissolved O_2 ml^{-1} at 37°C. In the electrode, 200 µl of the dense cell suspension was added to 800 µl air-saturated PBS for a final volume of 1 ml. The chamber was sealed and the suspension allowed to equilibrate for 2 minutes. The assay was initiated by the addition of 20 µl of 1 M sodium formate (prepared in PBS) and the rate of oxygen consumption recorded for 90 s. The total protein concentration of the cell suspensions was determined by Lowry assay and the specific rate of formate-dependent oxygen consumption expressed as nM oxygen consumed min^{-1} mg^{-1} total protein.

References

1.   Ochman H, Lawrence JG, Groisman EA. Lateral gene transfer and the nature of bacterial innovation. Nature 2000, 405:299-304.
2.   Koonin EV, Makarova KS, Aravind L. Horizontal gene transfer in prokaryotes: quantification and classification. Annu Rev Microbiol 2001, 55:709-742.
3.   Haldane JB. A defense of beanbag genetics. Perspect Biol Med 1964, 7:343-359.
4.   Bateson W. Heredity And Variation In Modern Lights. In *Darwin And Modern Science.* Edited by Seward AC. Cambridge: Cambridge University Press; 1909: 85–101
5.   Fisher RA. The correlation between relatives on the supposition of Mendelian inheritance. Trans R Soc Edinb 1918, 52:399–433.
6.   Wright S. Evolution in Mendelian populations. Genetics 1931, 16:97-159.
7.   Dobzhansky T. Studies on hybrid sterility. II. Localization of sterility factors in *Drosophila pseudoobscura* hybrids. Genetics 1936, 21:113-135.
8.   Dobzhansky T. Genetics And The Origin Of Species. New York: Columbia University Press; 1937.
9.   Muller HJ. Bearing of the *Drosophila* work on systematics. The new systematics 1940:185-268.
10. Muller H. Isolating mechanisms, evolution, and temperature. Biol Symp 1942, 6:71-125.
11. Kimura M. Attainment of quasi linkage equilibrium when gene frequencies are changing by natural selection. Genetics 1965, 52:875-890.
12. Neher RA, Shraiman BI. Competition between recombination and epistasis can cause a transition from allele to genotype selection. PNAS 2009, 106:6866-6871.
13. Wilson DJ, Gabriel E, Leatherbarrow AJH, Cheesbrough J, Gee S, Bolton E, Fox A, Hart CA, Diggle PJ, Fearnhead P. Rapid evolution and the importance of recombination to the gastroenteric pathogen *Campylobacter jejuni*. Mol Biol Evol 2009, 26:385-397.
14. Gibson B, Wilson DJ, Feil E, Eyre-Walker A. The distribution of bacterial doubling times in the wild. Proc R Soc B 2018, 285.
15. Vos M, Didelot X. A comparison of homologous recombination rates in bacteria and archaea. The ISME Journal 2009, 3:199-208.
16. Calland JK, Pascoe B, Bayliss SC, Mourkas E, Berthenet E, Thorpe HA, et al. Quantifying bacterial evolution in the wild: A birthday problem for Campylobacter lineages. PLoS Genet 2021, 17:e1009829.
17. Arnold BJ, Gutmann MU, Grad YH, Sheppard SK, Corander J, Lipsitch M, Hanage WP. Weak epistasis may drive adaptation in recombining bacteria. Genetics 2018, 208:1247-1260.
18. Sheppard SK, Guttman DS, Fitzgerald JR. Population genomics of bacterial host adaptation. Nat Rev Genet 2018, 19:549-565.
19. Andreani NA, Hesse E, Vos M. Prokaryote genome fluidity is dependent on effective population size. The ISME Journal 2017, 11:1719-1721.
20. Vos M, Eyre-Walker A. Are pangenomes adaptive or not? Nat Microbiol 2017, 2:1576-1576.
21. Shapiro BJ. The population genetics of pangenomes. Nat Microbiol 2017, 2:1574-1574.
22. McInerney JO, McNally A, O'Connell MJ. Why prokaryotes have pangenomes. Nat Microbiol 2017, 2:17040.

23. Orr HA. Dobzhansky, Bateson, and the genetics of speciation. Genetics 1996, 144:1331-1335.

24. Fraser C, Hanage WP, Spratt BG. Recombination and the nature of bacterial speciation. Science 2007, 315:476-480.

25. Sheppard SK, Dallas JF, Strachan NJ, MacRae M, McCarthy ND, Wilson DJ, Gormley FJ, Falush D, Ogden ID, Maiden MC, Forbes KJ. *Campylobacter* genotyping to determine the source of human infection. Clin Infect Dis 2009, 48:1072-1078.

26. Sheppard SK, McCarthy ND, Falush D, Maiden MC. Convergence of *Campylobacter* species: implications for bacterial evolution. Science 2008, 320:237-239.

27. Sheppard SK, Didelot X, Jolley KA, Darling AE, Pascoe B, Meric G, et al. Progressive genome-wide introgression in agricultural *Campylobacter coli*. Mol Ecol 2013, 22:1051-1064.

28. Cohan FM. Sexual Isolation And Speciation In Bacteria. In *Genetics of Mate Choice: From Sexual Selection to Sexual Isolation.* Springer; 2002: 359-370

29. Ansari MA, Didelot X. Inference of the properties of the recombination process from whole bacterial genomes. Genetics 2014, 196:253-265.

30. Retchless AC, Lawrence JG. Phylogenetic incongruence arising from fragmented speciation in enteric bacteria. PNAS 2010, 107:11453-11458.

31. Dearlove BL, Cody AJ, Pascoe B, Méric G, Wilson DJ, Sheppard SK. Rapid host switching in generalist *Campylobacter* strains erodes the signal for tracing human infections. The ISME Journal 2016, 10:721-729.

32. Weerakoon DR, Borden NJ, Goodson CM, Grimes J, Olson JW. The role of respiratory donor enzymes in *Campylobacter jejuni* host colonization and physiology. Microb Pathog 2009, 47:8-15.

33. Taylor AJ, Kelly DJ. The function, biogenesis and regulation of the electron transport chains in *Campylobacter jejuni*: New insights into the bioenergetics of a major food-borne pathogen. Adv Microb Physiol 2019, 74:239-329.

34. Smart JP, Cliff MJ, Kelly DJ. A role for tungsten in the biology of *Campylobacter jejuni*: tungstate stimulates formate dehydrogenase activity and is transported via an ultra-high affinity ABC system distinct from the molybdate transporter. Mol Microbiol 2009, 74:742-757.

35. Doerrler WT, Sikdar R, Kumar S, Boughner LA. New functions for the ancient DedA membrane protein family. J Bacteriol 2013, 195:3-11.

36. Ledgham F, Quest B, Vallaeys T, Mergeay M, Covès J. A probable link between the DedA protein and resistance to selenite. Res Microbiol 2005, 156:367-374.

37. Shaw FL, Mulholland F, Le Gall G, Porcelli I, Hart DJ, Pearson BM, van Vliet AHM. Selenium-dependent biogenesis of formate dehydrogenase in *Campylobacter jejuni* is controlled by the *fdhTU* accessory genes. J Bacteriol 2012, 194:3814-3823.

38. Sheppard SK, McCarthy ND, Jolley KA, Maiden MCJ. Introgression in the genus *Campylobacter*: generation and spread of mosaic alleles. Microbiology 2011, 157:1066-1074.

39. Croucher NJ, Harris SR, Barquist L, Parkhill J, Bentley SD. A high-resolution view of genome-wide pneumococcal transformation. PLoS Path 2012, 8:e1002745.

40. He M, Sebaihia M, Lawley TD, Stabler RA, Dawson LF, Martin MJ, et al. Evolutionary dynamics of *Clostridium difficile* over short and long time scales. PNAS 2010, 107:7527-7532.

41. Power PM, Bentley SD, Parkhill J, Moxon ER, Hood DW. Investigations into genome diversity of *Haemophilus influenzae* using whole genome sequencing of clinical isolates and laboratory transformants. BMC Microbiol 2012, 12:273.

42. Yahara K, Méric G, Taylor AJ, de Vries SP, Murray S, Pascoe B, et al. Genome-wide association of functional traits linked with *Campylobacter jejuni* survival from farm to fork. Environ Microbiol 2017, 19:361-380.

43. Arnoux P, Ruppelt C, Oudouhou F, Lavergne J, Siponen MI, Toci R, Mendel RR, Bittner F, Pignol D, Magalon A, Walburger A. Sulphur shuttling across a chaperone during molybdenum cofactor maturation. Nat Commun 2015, 6:6148.

44. Taveirne ME, Sikes ML, Olson JW. Molybdenum and tungsten in *Campylobacter jejuni*: their physiological role and identification of separate transporters regulated by a single ModE-like protein. Mol Microbiol 2009, 74:758-771.

45. Aguilar-Barajas E, Díaz-Pérez C, Ramírez-Díaz MI, Riveros-Rosas H, Cervantes C. Bacterial transport of sulfate, molybdate, and related oxyanions. BioMetals 2011, 24:687-707.

46. Thomas MT, Shepherd M, Poole RK, van Vliet AHM, Kelly DJ, Pearson BM. Two respiratory enzyme systems in *Campylobacter jejuni* NCTC 11168 contribute to growth on l-lactate. Environ Microbiol 2011, 13:48-61.

47. Ünal CM, Steinert M. Microbial peptidyl-prolyl cis/trans isomerases (PPIases): virulence factors and potential alternative drug targets. Microbiology and molecular biology reviews 2014, 78:544-571.

48. Knudson AG, Jr. Mutation and cancer: statistical study of retinoblastoma. Proc Natl Acad Sci U S A 1971, 68:820-823.

49. Oren Y, Smith MB, Johns NI, Kaplan Zeevi M, Biran D, Ron EZ, Corander J, Wang HH, Alm EJ, Pupko T. Transfer of noncoding DNA drives regulatory rewiring in bacteria. PNAS 2014, 111:16112-16117.

50. Arnold B, Sohail M, Wadsworth C, Corander J, Hanage WP, Sunyaev S, Grad YH. Fine-scale haplotype structure reveals strong signatures of positive selection in a recombining bacterial pathogen. Mol Biol Evol 2019, 37:417-428.

51. Wadsworth CB, Arnold BJ, Sater MRA, Grad YH. Azithromycin resistance through interspecific acquisition of an epistasis-dependent efflux pump component and transcriptional regulator in *Neisseria gonorrhoeae*. mBio 2018, 9:e01419-01418.

52. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. Journal of computational biology : a journal of computational molecular cell biology 2012, 19:455-477.

53. Sheppard SK, Didelot X, Meric G, Torralbo A, Jolley KA, Kelly DJ, Bentley SD, Maiden MCJ, Parkhill J, Falush D. Genome-wide association study identifies vitamin B5 biosynthesis as a host specificity factor in *Campylobacter*. PNAS 2013, 110:11923-11927.

54. Sheppard SK, Cheng L, Méric G, de Haan CP, Llarena AK, Marttinen P, et al. Cryptic ecology among host generalist *Campylobacter jejuni* in domestic animals. Mol Ecol 2014, 23:2442-2451.

55. Pascoe B, Méric G, Yahara K, Wimalarathna H, Murray S, Hitchings MD, et al. Local genes for local bacteria: evidence of allopatry in the genomes of transatlantic *Campylobacter* populations. Mol Ecol 2017, 26:4497-4508.

56. Pascoe B, Schiaffino F, Murray S, Méric G, Bayliss SC, Hitchings MD, et al. Genomic epidemiology of *Campylobacter jejuni* associated with asymptomatic pediatric infection in the Peruvian Amazon. PLoS Negl Trop Dis 2020, 14:e0008533.

57. Mourkas E, Taylor AJ, Méric G, Bayliss SC, Pascoe B, Mageiros L, et al. Agricultural intensification and the evolution of host specialism in the enteric pathogen *Campylobacter jejuni*. PNAS 2020, 117:11018-11028.

58. Jolley KA, Maiden MCJ. BIGSdb: scalable analysis of bacterial genome variation at the population level. BMC Bioinformatics 2010, 11:595.

59. Dingle KE, Colles FM, Wareing DR, Ure R, Fox AJ, Bolton FE, Bootsma HJ, Willems RJ, Urwin R, Maiden MC. Multilocus sequence typing system for *Campylobacter jejuni*. J Clin Microbiol 2001, 39:14-23.

60. Méric G, Yahara K, Mageiros L, Pascoe B, Maiden MCJ, Jolley KA, Sheppard SK. A reference pan-genome approach to comparative bacterial genomics: identification of novel epidemiological markers in pathogenic *Campylobacter*. PLOS One 2014, 9:e92798.

61. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST server: rapid annotations using subsystems technology. BMC Genomics 2008, 9:75.

62. Sheppard SK, Jolley KA, Maiden MCJ. A gene-by-gene approach to bacterial population genomics: whole genome MLST of *Campylobacter*. Genes 2012, 3:261-277.

63. Price MN, Dehal PS, Arkin AP. FastTree 2 – approximately maximum-likelihood trees for large alignments. PLOS One 2010, 5:e9490.

64. Didelot X, Wilson DJ. ClonalFrameML: efficient inference of recombination in whole bacterial genomes. PLoS Comp Biol 2015, 11:e1004041.

65. Browning BL, Browning SR. A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. Am J Hum Genet 2009, 84:210-223.

66. Yahara K, Furuta Y, Oshima K, Yoshida M, Azuma T, Hattori M, Uchiyama I, Kobayashi I. Chromosome painting *in silico* in a bacterial species reveals fine population structure. Mol Biol Evol 2013, 30:1454-1464.

67. Lawson DJ, Hellenthal G, Myers S, Falush D. Inference of population structure using dense haplotype data. PLoS Genet 2012, 8:e1002453.

68. Yahara K, Didelot X, Ansari MA, Sheppard SK, Falush D. Efficient inference of recombination hot regions in bacterial genomes. Mol Biol Evol 2014, 31:1593-1605.

69. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol 2010, 59:307-321.

70. Mell JC, Shumilina S, Hall IM, Redfield RJ. Transformation of natural genetic variation into *Haemophilus influenzae* genomes. PLoS Path 2011, 7:e1002151-e1002151.

71. Collins C, Didelot X. A phylogenetic method to perform genome-wide association studies in microbes that accounts for population structure and recombination. PLoS Comp Biol 2018, 14:e1005958.

72. van Vliet AH, Wooldridge KG, Ketley JM. Iron-responsive gene regulation in a *Campylobacter jejuni fur* mutant. J Bacteriol 1998, 180:5291-5298.

73. Cameron A, Gaynor EC. Hygromycin B and apramycin antibiotic resistance cassettes for use in *Campylobacter jejuni*. PLOS One 2014, 9:e95084.

Acknowledgments

Funding: Wellcome Trust grant 088786/C/09/Z (SS), Medical Research Council (MRC) grant

MR/M501608/1 (SS), Medical Research Council (MRC) grant MR/L015080/1 (SS), Food

Standards Agency project FS101087 (SS), Biotechnology & Biological Sciences Research

Council (BBSRC) grant BB/S014497/1 (DK).

Author contributions:

Conceptualization and study design: SS, XD, DF, KY, DK, AT

Sample collection: SS, AvV, NW

Laboratory work: AT, LM, MH, BP

Data archiving: BP, KJ

Data analysis: AT, XD, KY, LM, JKC, EM, SP, SB, SS, JC

Data interpretation: KY, XD, SS, MM, JP, DK, AT, DF

Writing: SS, CK, AT, DK

Competing Interest Statement: The authors declare that they have no competing interests.

Data and materials availability: Short-read sequence data for all isolates sequenced in this study

are deposited in the sequence read archive (SRA) and can be found associated with NCBI

BioProjects   PRJNA689604   (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA689604)   and

PRJEB11972 (https://www.ncbi.nlm.nih.gov/bioproject/PRJEB11972). These were augmented

with 498 previously published genomes and assembled genomes are available on Figureshare

(doi.org/10.6084/m9.Figureshare.13521683). Accession numbers for all genomes are included in

Data S4. Source data are provided for this paper (Data S2).

Figures and Tables

Figure 1. Genome-wide introgression from *C. jejuni* to *C. coli*. (A) Phylogenetic tree reconstructed using neighbour-joining on a whole-genome alignment of 973 *C. jejuni* and *C. coli* isolates. Introgressed *C. coli* clades are represented with red (CC-828, n=677) and purple (CC-1150, n=12) circles, unintrogressed clade 1* (n=35) is shown in pink, clade 2 (n=45) in yellow and clade 3 (n=58) in green. A set of 146 *C. jejuni* genomes belonging to 30 clonal complexes (4 to 5 isolates per ST) are shown in blue. Recipient and donor populations, used to infer introgression in chromosome painting analysis, are indicated. The scale bar represents the

number of substitutions per site. (B) Summary of introgressed *C. jejuni* SNPs in *C. coli* CC-828

(n=677, red) and CC-1150 (n=12, purple) genomes using ChromoPainter; the number of

introgressed core SNPs (coloured histograms; left y-axis) and core genes (white histograms; right

y-axis) for a range of recipient strains proportions (at least 1, more than 50% and more than 98%)

is shown. (C) Diagram of *Campylobacter* species and clade (C1*, C2, C3) divergence with arrows

indicating sequential introgression from *C. jejuni* into *C. coli* (I) clade 1, (II, III) CC-828 and CC-

1150, (IV, V) subsequent clonal expansion and ongoing introgression.

Figure 2. Covariation in introgressed *C. coli* genomes. (A) Possible SNP pairs in ancestral donor

C. jejuni and recipient *C. coli* genomes and potential recombinant (***R***) haplotype pairs in

introgressed *C. coli*. Possible genome haplotypes are designated with *C. jejuni* fixed SNPs (blue

X), other *C. jejuni* SNP (gray X), *C. coli* fixed SNP (red X), and other *C. coli* SNPs (gray X). (B)

The frequency of major and minor haplotype combinations among the 2578 pairs of covarying

SNPs in the 689 *C. coli* clade-1 recipient genomes, revealing that 83.5% of long-range covariation

was between introgressed *C. jejuni* sites. (C) Miami plot of each polymorphic site showing the

maximum *p*-value for covarying biallelic pairs (>20kb apart) and the frequency of introgression in

CC-828 and CC-1150.

Figure 3. Genomic context and physiological roles of introgressed epistatically linked genes involved in FDH biogenesis and activity. (A) The position of putative epistatic sites mapped on the *C. coli* CVM29710 reference for covarying *C. jejuni-C. jejuni* SNPs (blue) in 16 gene pairs (a to l), and other haplotype combinations (grey). (B) Genome organization of the most common co-varying introgressed *C. jejuni – C. jejuni* gene pairs linked to FDH. (C-D) FDH activity of whole cells determined by oxygen consumption rates in a Clark-type electrode (nmol oxygen consumed per minute per mg of total protein) for (C) cells grown in rich media (excess selenium), (D) cells grown in minimal media with 0.5 nM sodium selenite.

Figure 4. A two-hit model for core genome recombination in natural *C. coli* populations.
Fitness landscape upon which (i) unintrogressed *C. coli* (red) and *C. jejuni* (blue) co-exist. Their genomes (internal circles) harbour haplotype pairs (x-x) that segregate by species. (ii) Horizontal gene transfer, HGT, occurs (*R1*) disrupting covarying genetic elements and reducing the relative fitness of introgressed *C. coli* to varying degrees, few strains retain mixed *C. coli - C. jejuni* haplotypes. (iii) HGT continues (*R2*) and, where recombined mixed haplotypes survived, ancestral *C. jejuni* haplotype pairs are reinstated in introgressed *C. coli* – some of which outcompete unintrogressed strains.