Cross-species evolutionary rewiring in the enteric bacterium *Campylobacter*

Aidan J. Taylor[1]§, Guillaume Méric[2]†§, Koji Yahara[3], Ben Pascoe[2], Leonardos Mageiros[2,4], Evangelos Mourkas[2], Jessica K Calland[2], Santeri Puranen[5], Matthew D. Hitchings[6], Sion Bayliss[2], Keith A. Jolley[7], Carolin M. Kobras[1], Nicola J. Williams[8], Arnoud H. M. van Vliet[9], Julian Parkhill[10], Martin C. J. Maiden[7], Jukka Corander[5,11,12], Daniel Falush[13], Xavier Didelot[14], David J. Kelly[1*], Samuel K. Sheppard[2,7*]

[1]Department of Molecular Biology and Biotechnology, The University of Sheffield, Sheffield S10 2TN, United Kingdom;
[2]The Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK;
[3]Antimicrobial Resistance Research Centre, National Institute of Infectious Diseases, Tokyo, Japan;
[4]Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria 3004, Australia;
[5]Department of Mathematics and Statistics, Helsinki Institute for Information Technology, University of Helsinki, Helsinki, Finland;
[6]Swansea University Medical School, Institute of Life Science, Swansea SA2 8PP, UK;
[7]Department of Zoology, University of Oxford, South Parks Road, Oxford OX1 3PS, UK;
[8]Department of Epidemiology and Population Health, Institute of Infection & Global Health, University of Liverpool, Leahurst Campus, Wirral, UK;
[9]School of Veterinary Medicine, University of Surrey, Guildford GU2 7AL, UK;
[10]Department of Veterinary Medicine, University of Cambridge, Madingley Rd, Cambridge CB3 0ES, UK;
[11]Department of Biostatistics, University of Oslo, Oslo, Norway;
[12]Parasites and Microbes, Wellcome Sanger Institute, Cambridge, UK;
[13]The Centre for Microbes, Development and Health, Institut Pasteur of Shanghai, Shanghai, China;
[14]School of Life Sciences and Department of Statistics, University of Warwick, Coventry CV4 7AL, UK.

*Corresponding author(s): David J. Kelly, d.kelly@sheffield.ac.uk; Samuel K. Sheppard, s.k.sheppard@bath.ac.uk;
†Present address: Cambridge Baker Systems Genomics Initiative, Baker Heart and Diabetes Institute, 75 Commercial Rd, Melbourne 3004, Victoria, Australia.

§ Contributed equally.

Keywords: comparative genomics, introgression, admixture, epistasis, evolution, *Campylobacter,* microbiology

Running title: Epistasis and recombination in *Campylobacter*

Abstract

The lateral transfer of genes among bacterial strains and species has opposing effects, conferring potentially beneficial adaptations whilst introducing disharmony in coadapted genomes. The prevailing outcome will depend upon the fitness cost of disrupting established epistatic interactions between genes. It is challenging to understand this in nature because it requires population-scale analysis of recombination and genomic coadaptation, and laboratory confirmation of the functional significance of genotype variation. By assigning the ancestry of DNA in the genomes of two species of the world's most common enteric bacterial pathogen, we show that up to 28% of the *Campylobacter coli* genome has been recently introgressed from *Campylobacter jejuni*. Then, by quantifying covariation across the genome we show that >83% of putative epistatic links are between introgressed *C. jejuni* genes in divergent genomic positions (>20kb apart), consistent with independent acquisition. Much of this covariation is between 16 genes, with just 5 genes accounting for 99% of epistatic SNP pairs. Laboratory mutagenesis and complementation cloning assays demonstrated functional links between these genes, specifically related to formate dehydrogenase (FDH) activity. These findings suggest that the genetic confederations that define genomic species may be transient. Even for complex traits such as central metabolism in the bacterial cell, conditions can arise where epistatic genes can be decoupled, transferred and reinstated in a new genetic background.

Introduction

Complex biological life is made possible by the interaction of genes. In bacteria, a vast array of genes, found together in almost infinite combinations, has allowed colonization of virtually every conceivable habitat on earth. While mutation remains the engine of genetic novelty, for most bacteria adaptation also involves the acquisition of genes from other strains and species through horizontal gene transfer (HGT)[1] potentially conferring new phenotypes to future generations. In some well documented cases, a single nucleotide substitution or acquisition of a small number of genes, can prompt new evolutionary trajectories with striking outcomes such as the emergence of virulent or antibiotic resistant strains[2]. With such dynamic genomic architecture, it may be tempting (and possibly useful[3]) to consider genes as independent units that 'plug and play' innovation into recipient genomes. This is clearly an oversimplification. In fact, genomes are highly interactive wherein the effect of one gene depends on another (epistasis). Therefore, it is likely that some introduced changes will disrupt gene networks and be costly to the original coadapted genetic background, particularly for complex phenotype innovation involving multiple genes.

Understanding how epistasis influences the evolution of phenotype diversity has preoccupied researchers since the origin of population genetics[4-10], with much emphasis placed upon the relative amounts of recombination and epistatic effect sizes[11,12]. In sexual populations, such as outbreeding metazoans, genetic variation is shuffled at each generation so genes can only rise in frequency if they have high mean fitness across genetic backgrounds. This means that it is unlikely that multiple distinct epistatic allele combinations will be maintained in the same population and barriers to gene flow, such as geographic isolation, may be required for marked phenotypic diversification[8]. In bacteria, however, rapid clonal reproduction allows multiple independent beneficial allele pairs to rise to high frequency in a single population. For example, in common enteric bacteria such as *Escherichia coli, Salmonella enterica* and *Campylobacter jejuni* the doubling time in the wild has been estimated at around 24 hours or less[13,14]. Therefore, though HGT occurs in these organisms[15], even in highly recombinogenic *C. jejuni*[13,16], there will likely be many millions of bacterial generations between recombination events at a given locus. This allows mutations that are beneficial only in specific genetic backgrounds to establish in a single population and linkage disequilibrium to form between different epistatic pairs[17].

In this coadapted genomic landscape, recombination presents something of a paradox. On the one hand, it promotes adaptation by conferring novel functionality on the recipient genome[18] and reduces competition between clones that carry different beneficial mutations (clonal interference), on the other it potentially introduces disharmonious genes or gene-combinations that will be discriminated against by selection[19]. Indeed, negative epistatic interactions between genes with different evolutionary histories have been proposed as a barrier to recombination[4-10,20], particularly between species. However, interspecies recombination is common in bacteria[18,21]. One of the most conspicuous examples of this can be seen in the common animal gut bacterium *Campylobacter,* which is among the most prolific causes of human bacterial gastroenteritis worldwide[22]. Specifically, introgression between the two most important pathogenic species, *C. jejuni* and *C. coli,* has led to the emergence of a globally distributed 'hybrid' *C. coli* lineage[23] that is responsible for almost all livestock and human infections with this species.

This evolutionary scenario provides an ideal context for addressing the paradox of recombination in coadapted genomes. First, because up to 23% of the core genome of

common *C. coli* has been recently introgressed from *C. jejuni*[24], potentially disrupting
epistatic interactions. Second, because *C. jejuni* and *C. coli* have undergone an extended
period of independent evolution (85% average nucleotide identity), therefore recombined
sequence is conspicuous in the genome. Third, because the outcompetition of ancestral strains
by introgressed *C. coli* is inconsistent with traditional assumptions of selection against hybrid
lineages.
125
Here, we analyse the genomes of isolates representing known diversity within *C. jejuni* and
C. coli and perform chromosome painting analysis to assign ancestry across the genomes of
introgressed *C. coli*. We then perform a systematic screen for long-range covariation in the
core genome to identify SNPs that are in strong linkage disequilibrium in independent genetic
backgrounds and, therefore, are not explained by the clonal frame of the population. Finally,
we investigate the function of covarying genes that account for most of the interactions and
confirm epistasis among coadapted genes. By combining these analyses, we demonstrate how
multiple interacting genes from one species can transfer to another, modifying the co-adapted
genomic landscape of the recipient and promote a natural admixture-mediated genetic
revolution[19,25].
136
137
138 **Results**
139
***Campylobacter* populations are highly structured with intermediate sequence clusters**
Pan-genome analysis identified the presence and variation of every automatically annotated
gene from every genome. This revealed a core genome of 631 gene orthologues in all
Campylobacter isolates in this study. There were 1287 genes common to all *C. jejuni* isolates
and 895, 1021 and 1272 common to *C. coli* clades 1, 2 and 3, respectively. Consistent with
previous studies[24], neighbour-joining and ClonalFrameML trees based on genes within
concatenated core genome alignments revealed population structure in which *C. jejuni* and *C.*
coli clade 2 and 3 isolates each formed discrete clusters (Figure 1A, Supplementary Figure 1).
Isolates designated as *C. coli* clade 1 were found in three clusters on the phylogeny:
unintrogressed ancestral strains, and the ST-828 and ST-1150 clonal complexes which
account for the great majority of strains found in agriculture and human disease[24]. Pan-
genome analysis quantified the increase in unique gene discovery as the number of sampled
genomes increased (Figure 1B). For all sequence clusters there was evidence of an open pan-
genome with a trend towards continued rapid gene discovery within sequence clusters with
fewer isolates. There was considerable accessory genome variation between species and
clades, potentially associated with important adaptive traits (Supplementary Figure 2) and
there was evidence that the average number of genes per genome was greater in *C. coli* ST-
828 complex isolates than in *C. jejuni* (Figure 1C).
158
There is substantial introgression within the *C. coli* core genome
The large genetic distances among *C. coli* clade 1 isolates (Figure 1A), have been shown to
be a consequence of the import of DNA from *C. jejuni* rather than accumulation of mutations
during a prolonged period of separate evolution[23]. Using chromosome painting to infer the
coancestry of core-genome haplotype data from CC-828 and CC-1150 isolates gave a
detailed representation of the recombination-derived chunks from each *C. jejuni* donor group
to each recipient individual (Supplementary Figure 3). The majority of introgressed SNPs
were rare, occurring in fewer than 50 recipient genomes (Figure 2A). However, a large
proportion of the introgressed *C. coli* clade 1 genomes contained DNA of *C. jejuni* ancestry
in >98% of recipient isolates. Consistent with previous estimates[24], these regions where

introgressed DNA was largely fixed within the *C. coli* population, occurred across the
genome and comprised up to 15% and 28% of the CC-828 and CC-1150 isolate genomes
respectively (Figure 2A). When considering donor groups, the majority of introgressed DNA
in *C. coli* involves genes that are present in multiple *C. jejuni* lineages (core genes) (Figure
2B).
174
Having identified *C. jejuni* ancestry within *C. coli* genomes, we investigated the sequence of
events responsible for introgression. Most introgressed SNPs are found at low frequency in
both clonal complexes. However, there was evidence of SNPs that are introgressed in both
complexes as well as high frequency lineage specific introgression in both CC-828 and CC-
1150 (Figure 2C). Specifically, 25% of the *C. jejuni* DNA found in >98% of CC-828 isolates
was also found in CC-1150 (Figure 2D), implying that this genetic material was imported by
the common ancestor(s) of both complexes. Subsequent to the divergence of these two
complexes, introgression continued with nearly 75% of *C. jejuni* DNA present in one
recipient clonal complex and not the other. This is consistent with an evolutionary history in
which there was a period of progressive species and clade divergence reaching approximately
12% at the nucleotide level between *C. jejuni* and *C. coli* and around 4% between the
three *C. coli* lineages. More recently, changes to the patterns of gene flow led one
C. coli clade 1 lineage to import substantial quantities of *C. jejuni* DNA, and further lineage-
specific introgression gave rise to two clonal complexes (CC-828 and CC-1150) that
continued to accumulate *C. jejuni* DNA, independently creating the population structure
observed today (Figure 2E).
191
The high magnitude introgression into *C. coli* clade 1 isolates has introduced thousands of
nucleotide changes to the core genome. However, divergence in bacteria may be uneven
across the genome. First, because recombination is more likely to occur in regions where
donor and recipient genomes have high nucleotide similarity[21,26,27]. Second, because of
'fragmented speciation'[28], in which gene flow varies in different parts of the genome, such as
regions responsible for adaptive divergence, leading to phylogenetic incongruence among
genes. Consistent with previous estimates[24], we found that the three *C. coli* clades had similar
high divergence with *C. jejuni* across the genome, ranging from 68% to 98% nucleotide
identity for individual genes (Figure 2F), implying a period of divergence with low levels of
gene flow. We found no evidence that high genetic differentiation between the species
prevented recombination. While there was some evidence that more recombination occurred
in regions of low nucleotide divergence (between unintrogressed *C. coli* clade 1 and *C.
jejuni*), introgression occurred across the genome at sites with varying levels of nucleotide
identity (Figure 2F). This level of recombination has greatly increased overall genetic
diversity across the genome in *C. coli* clade 1 and introduced changes that have potential
functional significance.
208
Much of the putative epistasis occurs between SNPs in introgressed genes
ClonalFrameML analysis revealed the importance of homologous recombination in
generating sequence variation within the introgressed *C. coli*. Estimates of the relative
frequency of recombination versus mutation (R/θ=0.43), mean recombination event length
(δ=152bp) and average amount of polymorphism per site in recombined fragments (v=0.07),
imply that recombination has had an effect (*r/m*) 4.57 times higher than *de novo* mutation
during the diversification of CC-828. This is consistent with previous analysis and confirmed
recombination as the major driver of molecular evolution in *C. coli*[13,24,29]. The continuous
time Markov chain model for the joint evolution of pairs of biallelic sites on a phylogenetic
tree (Supplementary Figure 4) was applied to investigate patterns of covariance for all pairs

of sites >20kb apart (Figure 3A). For most biallelic sites there were few branches on the tree
where substitutions occurred, so that their evolution is compatible with separate evolution on
the same clonal frame. However, 2874 covarying pairs evolved more frequently together than
would be expected (p-value 10^{-8}) if they had evolved independently based on the tree, and
hence indicated patterns of putative epistasis.
224
Among them, the location of 2618 putative epistatic pairs of sites was compared to the
inferred ancestry (unintrogressed *C. coli* or *C. jejuni*) of sequence across the genome of CC-
828 and CC-1150 *C. coli* strains (Figure 3B, Supplementary Data 1). For each epistatic pair,
the major and minor haplotype were defined if there was haplotype polymorphism between
C. jejuni and CC-828 and CC-1150 *C. coli.* This allowed quantification of the number of
covarying sites that occurred between an ancestral *C. coli* (unintrogressed) and an
introgressed *C. jejuni* allele, two introgressed alleles, and sites that do not segregate by
species. Strikingly, the breakdown of the major and minor haplotype combinations among the
2618 epistatic pairs (Figure 3C, Source Data) shows the major haplotype for 83.5% of
putative epistatic SNP pairs was *C. jejuni* indicating that both co-varying sites had *C. jejuni*
ancestry, consistent with epistasis between introgressed ancestral *C. jejuni* sequence at
divergent genomic positions. Investigation of the genes containing co-varying sites revealed
that 2187 SNP pairs were in 16 genes with just five genes accounting for 99.1% of them
(Figure 3D, Supplementary Data 2, Supplementary Figure 5).
239
Genomic context and physiological role of epistatically linked genes
The five genes accounting for the majority of epistatic interactions (*cj1167*, *cj1168c*,
cj1171c/ppi, *cj1507c/modE* and *cj1508c/fdhD*) were investigated for their physiological role
in *C. jejuni*. FdhD and ModE are proteins involved in the biogenesis of formate
dehydrogenase (FDH). The FDH complex (FdhABC) oxidises formate to bicarbonate to
generate electrons that fuel cellular respiration. Formate is an abundant electron donor
produced by host microbiota and an important energy source for *Campylobacter in vivo*[30,31].
The remaining three genes, *cj1167* (annotated incorrectly as *ldh*, lactate dehydrogenase),
cj1168c and *cj1171c* (*ppi*) are also grouped together on the genome, where *cj1167* and
cj1168c are adjacent but with the open reading frames (ORFs) convergent and overlapping,
while *ppi* is upstream, separated by two non-epistatically linked genes (*cj1169c* and *cj1170c*,
Figure 4A). Considering the genomic arrangement, it is therefore clear that the putative
epistatic links uncovered in this study essentially occur across two loci in the genome
(*fdhD/modE* and *cj1167/cj1168c/ppi*), with each of the latter three genes linked with both
fdhD and *modE* (Figure 4A). Given the known function of *fdhD* and *modE* in biogenesis of
the FDH complex, we hypothesised that *cj1167/cj1168c/ppi* might also have some role in
FDH biogenesis or activity in order to form a functional epistatic connection. We therefore
constructed deletion mutants to investigate the possible role of these genes in FDH activity.
258
Initially, each of the mutants and their parental wildtype (*C. jejuni* NCTC11168) were grown
in rich media (Muller-Hinton broth) and their formate dependent oxygen consumption rates
determined (Figure 4B). *cj1167, cj1168c* and *ppi* mutants demonstrated wildtype levels of
FDH activity, while activity in both *fdhD* and *modE* mutants was abolished. In order to
confirm that the phenotype of the *fdhD* and *modE* mutants was not due to a polar effect on the
surrounding *fdh* locus, these mutants were genetically complemented by reintroduction of a
second copy of the wildtype gene into the rRNA locus, which restored near wildtype levels of
FDH activity in both cases (Figure 4B).
267

As neither *cj1167, cj1168c* or *ppi* mutants showed altered FDH activity in cells grown in rich
media, we considered that their function may be related to an FDH-specific nutrient
requirement as would likely be found *in vivo*. Since the formate oxidising subunit of FDH,
FdhA, specifically requires a molybdo- or tungsto-pterin (Mo/W) cofactor and a
selenocysteine (SeC) residue for catalysis[32], Mo, W or Se supply presented possible targets.
cj1168c encodes a DedA family integral membrane protein of unknown function. DedA
proteins are solute transporters widespread in bacteria but are mostly uncharacterised[33].
However, a homologue of *cj1168c* in the heavy metal specialist beta-proteobacterium
Cupriavidus metallidurans has been shown to be involved in selenite (SeO_3^{2-}) uptake[34]. We
therefore speculated that Cj1168 could be a selenium oxyanion transporter that supplies Se
for SeC biosynthesis. To test this, FDH activities were measured in *cj1168c* mutant and
parental wildtype strains grown in minimal media with limiting concentrations of selenite or
selenate (SeO_4^{2-}). The data in Figure 4C shows that the *cj1168c* mutant displayed
significantly reduced FDH activity after growth with selenite in the low nM range, and this
phenotype was partially restored by genetic complementation. We therefore designated
cj1168c as *selF* (selenium transporter for formate dehydrogenase). However, although this
phenotype does suggest that SelF is a selenium importer, another unrelated selenium
transporter, FdhT (Cj1500), has previously been documented in *C. jejuni*[35], which is not
epistatic with *fdhD* or *modE*. In contrast to this previous report we found considerable
residual FDH activity still remained in an *fdhT* deletion mutant, which was fully restored to
wildtype levels by complementation (Figure 4D).
289
Finally, we tested whether the residual FDH activity in our *fdhT* mutant was due to selenium
uptake by SelF. An *fdhT selF* double mutant was generated and assayed for FDH activity
after growth in minimal media containing limiting concentrations of selenite or selenate
(Figure 4D). The *fdhT selF* double mutant demonstrated a significant additional reduction in
FDH activity over the *fdhT* single mutant, a phenotype that was partially restored by
complementing the double mutant with *selF*. Complementation of the double mutant with
fdhT returned FDH activity to near wildtype levels (Figure 4D). Taken together, our data
suggests that both FdhT and SelF facilitate selenium acquisition in *C. jejuni*, possibly
representing low and high affinity transporters, respectively (Figure 4E).
299
300 **Discussion**
301
Hybridization between distantly related organisms can bring together new gene combinations
and traits that potentially allow adaptation in a single evolutionary leap[36]. However,
introducing disharmony among genes that have coevolved in epistasis can lead to reduced
hybrid fitness, limiting the chance of this type of evolution by saltation[8,10]. Evidence for these
contrasting paradigms comes largely from eukaryotes, but among prokaryotes, widespread
HGT may seem to contradict assumptions about selection against hybrid lineages. For
example, the transfer of mobile resistance genes between bacterial strains and species can
confer a clear adaptive benefit, in a single step, allowing the recipient lineages to proliferate
in the presence of antibiotics. Clearly, therefore, epistasis is not an absolute barrier to long-
range recombination in bacteria.
312
There is ongoing debate about the extent to which the galaxy of accessory genes that are
variously present or absent in many bacterial genomes constitute a cache of mobile genes
from which innovation can be drawn and transferred between strains or species[37-40].
However, considering HGT of accessory genes does not fully address the extent to which
recombination is constrained by epistatic fitness interactions between genes. First, because

genes or gene clusters introduced as autonomous elements, encoding specific novel traits,
may cause minimal disruption to other essential cellular functions. Second, even when
introduced genes result in a concomitant change elsewhere in the genome[41], the fitness cost
may be outweighed by the benefit in a given niche. Third, and most importantly, because
most recombination in bacteria is between homologous sequences[21], where a given gene is
replaced by another version of the same gene.

Campylobacter is an ideal model organism for considering barriers to gene flow as
interspecies recombination is well documented within the core genome. Consistent with
previous studies[23,24,42], we found that a large proportion (15-28%) of the core genome of the
two common *C. coli* lineages found in livestock and human disease (CC-828 and CC-1150)
originated in *C. jejuni*. It is challenging to explain this level of genome-wide gene flow
between species after a prolonged period of diversification (~15% nucleotide divergence) as
it would likely disrupt coadapted gene networks. The evident success of hybrid *C. coli* in the
agricultural niche suggests that the accumulation of *C. jejuni* DNA was not detrimental, but
to specifically address if genome coadaptation was a barrier to recombination requires
quantification of covariation among alleles.

Most bacteria, including *Campylobacter,* have highly genetically structured populations
reflecting both neutral and adaptive evolutionary processes[18]. Therefore, a phylogenetically
naive statistical model of coadaptation would afford equal significance to linkage
disequilibrium resulting from selection and common ancestry. Even in recombinogenic
bacteria such as *C. coli*, HGT is not sufficiently common to scramble the genome and abolish
non-random SNP associations resulting from clonal population structure. Consistent with
other models[43], the statistical test developed here accounts for the amount of covariation that
would be expected based upon the clonal frame and identifies the same combinations of
alleles in independent genetic backgrounds, thus providing evidence for coadaptation.

By combining the results from the chromosome painting and covariation models we were
able to quantify the frequency of *C. coli* and *C. jejuni* SNPs in covarying allele pairs across
the genome of introgressed *C. coli*. Recent interspecies admixture results in allele pairs, that
correspond to *C. jejuni – C. coli* (and *vice versa*) and *C. jejuni – C. jejuni.* Comparing
haplotype frequency provides the opportunity to contrast the disruptive and beneficial effects
of homologous recombination in the bacterial genome. Under a neutral model, where
recombination does not disrupt beneficial epistatic interactions, there would be more *C. jejuni*
– C. coli than *C. jejuni – C. jejuni* covarying allele pairs in the recipient genome. Clearly,
relatively low levels of covariation between ancestral *C. coli* and introgressed *C. jejuni* SNPs
is expected, consistent with selection against disharmonious gene combinations in the
coadapted recipient genome. However, the finding that *C. jejuni – C. jejuni* allele pairs
constituted >83% of covarying introgressed haplotypes was striking. It is possible that in
some cases both sites were introgressed in a single recombination event as bacteria can
import very large pieces of DNA (>100kb)[44-46] but in *Campylobacter* LD for pairs of sites
decreases with distance to approximately 20kbp and then remains at the same level for very
distant sites[47]. Therefore, the divergent genome position of allele pairs (>20kb) implies that
they were acquired independently. It follows, therefore, that the acquisition of the first
introgression event was not fatal to the recipient genome, and was either mildly detrimental,
neutral or beneficial. Acquisition of the second member of the pair then potentially enhanced
the fitness restoring the integrated *C. jejuni – C. jejuni* coevolutionary unit.

Investigating covariation provides clues about genome plasticity and the potential benefit of
introgression for niche adaptation. However, to directly address if genes are in epistasis it is
necessary to test their function and confirm coadaptation. We demonstrated a functional link
amongst the top scoring co-varying gene pairs with FDH, a key enzyme allowing the
utilisation of formate as an electron donor *in vivo*[30,31]. In this study, FdhD and ModE were
shown to be essential for FDH activity. While FdhD is a sulfur-transferase known to be
required for the insertion of the pterin cofactor into FdhA[48] (Figure 4E), ModE is a
transcriptional repressor that has been shown previously only to regulate the Mo/W uptake
genes *mod* and *tup*[49,50]. However, the unexpected abolished FDH activity in a *modE* mutant
indicates further functions for ModE in FDH biogenesis which warrant future investigation.
Searching for functional links between *fdhD/modE* and *cj1167/selF/ppi*, revealed that a *selF*
mutant strain had significantly reduced FDH activity under conditions of selenite limitation, a
phenotype consistent with SelF being a Se oxyanion transporter and that functionally links
SelF with FdhD/ModE. We suggest that *selF* rather than *fdhT* is epistatic because SelF
confers an additional benefit for SeC biosynthesis (essential for FDH activity) under
conditions of selenium limitation, for example as may be found in the host (Figure 4E).
cj1167 encodes a cytoplasmic NADPH dependent 2-oxoacid dehydrogenase but the current
genome database annotation as lactate dehydrogenase (Ldh) is incorrect and its function is
unknown[51]. There is no precedent for the involvement of such an enzyme in bacterial FDH or
SeC biogenesis and we obtained no evidence for a functional connection between Cj1167 and
FDH activity. However, the overlapping convergent gene arrangement of *cj1167* and *cj1168c*
(*selF*) suggests a transcriptional architecture that might dictate these genes both form similar
epistatic dependencies even if Cj1167 is not required for FDH activity. Finally, *cj1171c* (*ppi*)
encodes a cytoplasmic peptidyl-prolyl *cis-trans* isomerase of the cyclophilin family. PPIases
are general protein folding catalysts that often have pleiotropic and redundant functions[52] and
we note that our *C. jejuni ppi* deletion mutant showed no growth defect as well as no
reduction in FDH activity. It is possible that if Cj1171 does help promote the folding of e.g.
FdhD or ModE, analysis of a simple deletion mutant may not reveal this if another PPIase
can substitute in that genetic background.
396
Explaining the aberrant genome architecture among introgressed *C. coli* is challenging.
Understanding the selective value of genes that promote proliferation depends on the overall
genetic environment[19]. Our findings are consistent with an evolutionary scenario where an
ancestral *C. coli* lineage underwent a niche transition and the surviving lineages (CC-828 and
CC-1150) gained access to *C. jejuni* DNA (Figure 3E&F). As the adaptive landscape of the
genome changed, potentially decoupling epistatic interactions that were previously selected,
new gene combinations could be introduced by recombination and tested in the *C. coli*
genetic background. Recent studies have shown that this type of genetic rewiring may be
more common than previously thought[53-55] and when it leads to the proliferation of a hybrid
organism it can be associated with the colonization of a new niche. Intensive livestock
systems represent a possible explanation for the genetic revolution in *C. coli*. Host ecology
can dramatically affect the evolution of gut-dwelling organisms, including *Campylobacter*[56]
and colonization of this niche could promote the conditions necessary to promote
introgressed *C. coli*. Whether host ecology is a factor or not, it is clear that conditions can
arise where coadapted genes in a highly interactive bacterial genome can be transferred
between species and reinstated as a single evolutionary unit in a new genome. This suggests
that epistasis is not an absolute barrier to genome-wide recombination in structured bacterial
populations.
415
416   **Methods**

417
Isolates, genome sequencing and assembly
A total of 973 isolates were used in this study, 827 from *C. coli* and a selection of 146 from a
diversity of *C. jejuni* clonal complexes (Supplementary Data 3). Isolates were sampled
mostly in the United Kingdom to maximise environmental and riparian reservoirs and thus
the representation of genetic diversity in *C. coli*. Isolates were stored in a 20% (v/v) glycerol
medium mix at -80°C and subcultured onto *Campylobacter* selective blood-free agar
(mCCDA, CM0739, Oxoid). Plates were incubated at 42°C for 48 h under microaerobic
conditions (5% CO_2, 5% O_2) generated using a CampyGen (CN0025, Oxoid) sachet in a
sealed container. Subsequent phenotype assays were performed on Brucella agar (CM0271,
Oxoid). Colonies were picked onto fresh plates and genomic DNA extraction was carried out
using the QIAamp® DNA Mini Kit (QIAGEN; cat. number: 51306) according to the
manufacturer's instructions. DNA was eluted in 100–200 µl of the supplied buffer and stored
at −20°C. DNA was quantified using a Nanodrop spectrophotometer and high-throughput
genome sequencing was performed on a MiSeq (Illumina, San Diego, CA, USA), using the
Nextera XT Library Preparation Kit with standard protocols involving fragmentation of 2 µg
genomic DNA by acoustic shearing to enrich for 600 bp fragments, A-tailing, adapter ligation
and an overlap extension PCR using the Illumina 3 primer set to introduce specific tag
sequences between the sequencing and flow cell binding sites of the Illumina adapter. DNA
cleanup was carried out after each step to remove DNA < 150 bp using a 1:1 ratio of
AMPure® paramagnetic beads (Beckman Coulter, Inc., USA). Short read paired-end data was
assembled using the *de novo* assembly algorithm, SPAdes (version 3.10.0)[57]. All novel
genome sequences (n=475) generated for use in this study are available on NCBI BioProjects
PRJNA689604 and PRJEB11972. These were augmented with 498 previously published
genomes and accession numbers for all genomes can be found in Supplementary Data
3[16,24,29,47,56,58-61].
443
444
Genome archiving, pan-genome content analyses and phylogenetic reconstruction
Contiguous genome sequence assemblies were individually archived on the web-based
database platform BIGSDB[62] and sequence type (ST) and clonal complex (CC) designation
were assigned based upon the *C. jejuni* and *C. coli* multi-locus sequence typing scheme[63]. To
examine the full pan-genome content of the dataset, a reference pan-genome list was
assembled as previously described[64]. Briefly, genome assemblies from all 973 genomes in
this study were automatically annotated using the RAST/SEED platform[65], the BLAST
algorithm was used to determine whether coding sequences from this list were allelic variants
of one another or 'unique' genes, with two alleles of the same gene being defined as sharing
>70% sequence identity on >10% of the sequence length. The prevalence of each gene in the
collection of 973 genomes was determined using BLAST with a positive hit in a genome
being defined as a local alignment of the reference sequence with the genomic sequence of
>70% identity on >50% of the length, as previously described[66]. The resulting matrix was
analysed for differentiating core and accessory genome variation. Genes present in all
genomes were concatenated to produce a core-genome alignment, used for subsequent
phylogenetic reconstructions. Phylogenetic trees were reconstructed using an approximation
of maximum-likelihood phylogenetics in FastTree2[67]. This tree was used as an input for
ClonalFrameML[68] to produce core genome phylogenies with branch lengths corrected for
recombination.
464
Inference of introgression

All 973 genomes were aligned to a full reference sequence of *C. coli* strain CVM29710. We
conducted imputation for polymorphic sites with missing frequency ≤10% using BEAGLE[69]
as previously reported[70]. A total of 286,393 gapless SNPs (~17% of the average *C. coli*
genome size) were used for recombination analyses. The coancestry of genome-wide
haplotype data was inferred based on alignments using chromosome painting
and FineStructure[71] as previously described[72]. Briefly, ChromoPainter was used to infer
chunks of DNA donated from a list of 33 donor groups normalised for sample size to each of
677 ST-CC-828 and 12 CC-1150 recipient haplotypes. Results were summarised into a
coancestry matrix containing the number of recombination-derived chunks from each donor
to each recipient individual. FineStructure was then used for 100,000 iterations of both the
burn-in and Markov chain Monte Carlo chain to cluster individuals based on the co-ancestry
matrix. The results are visualized as a heat map with each cell indicating the proportion of
DNA "chunks" (a series of SNPs with the same expected donor) a recipient receives from
each donor.
480
Analysis of covariation in bacterial genomes
Non-random allele associations can result from selection and clonal population structure. To
control for the latter, our approach identified SNP combinations in independent genetic
backgrounds by accounting for the sequence variation associated with the inferred phylogeny.
Based on the alignment of 677 genomes of *C. coli* CC-828, a first phylogenetic tree was
created using PhyML[73]. ClonalFrameML[68] was then applied to correct the tree by accounting
for the effect of recombination, and also to infer the ancestral sequence of each node.
Covariance was assessed for pairs of biallelic sites across the genome using a Continuous
Time Markov Chain (CTMC) model as follows. Briefly, let *A* and *a* denote the two alleles of
the first site and *B* and *b* denote the two alleles of the second site, so that there are four states
in total for the pair of sites (*ab*, *Ab*, *aB* and *AB*). The four substitution rates from *A* to *a*, from
a to *A*, from *B* to *b* and from *b* to *B* are not assumed to be identical, to allow for differences in
substitution rates in different parts of the genome and also to allow for non-equal rates of
forward and backward substitution (for example as a result of recombination opportunities).
Assuming no epistatic effect between the two sites ($\varepsilon=1$), the model M_0 has four free
parameters (α_1, α_2, β_1 and β_2) representing independent substitutions at the two sites. We
expand model M_0 with an additional fifth parameter $\varepsilon>1$ into model M_1 which is such that the
state *AB* where the first site is allele *A* and the second site is allele *B* is favored relative to the
other three sites *ab, aB* and *Ab*. Specifically, the state *AB* has a probability increased by a
factor ε^2 in the stationary distribution of the CTMC of model M_1 compared to model M_0.
501
Both models M_0 (with 4 parameters) and M_1 (with 5 parameters) are fitted to the data using
maximum likelihood techniques, where the likelihood is equal to the product for every branch
of the tree of the state at the bottom of the branch given the state at the top. The two fitted
models M_0 and M_1 are then compared using a likelihood-ratio test (LRT) as follows: since M_0
is nested with M_1, two times their difference in log-likelihood is expected to be distributed
according to a chi-square distribution with number of degrees of freedom equal to the
difference in their dimensionality, which is one. This LRT returns a *p*-value for the
significance of a covariation effect, and a Bonferroni correction is applied to determine a
conservative cutoff of significance that accounts for multiple testing. Furthermore, the test is
applied only to pairs of sites separated by >20kb to reduce the chance that they were the
result of a single recombination event, consistent with estimates of the length of recombined
DNA sequence in quantitative bacterial transformation experiments[44,74] and evidence from
Campylobacter genome analyses that show that LD for pairs of sites decreases with distance
to approximately 20kbp and then remains at the same level for very distant sites[47]. It is still

possible of course that rare recombination events would stretch 20kbp[44-46], but for this to
have an effect on the analysis of epistasis it would have to have happened several times for
the same pairs of sites against different genomic background which becomes quite unlikely
just by chance. This phylogenetically aware approach to testing for covariance presents the
advantage to naturally account for both population structure and the effect of
recombination[75]. The script implementing this coevolution test is available in R at:
https://github.com/xavierdidelot/campy.
523
Quantifying covariation between recombined and unrecombined genomic regions
The results of the introgression and covariation analyses were combined so that for each pair
of significantly covarying SNPs (p-value $<10^{-8}$), haplotype frequency was calculated among
the 689 recipient introgressed *C. coli* clade-1 strains as well as among the donor *C. coli*
(ancestral) and *C. jejuni* strains, respectively. If the most frequent haplotype of the pair is the
same between the donor *C. coli* (ancestral) and *C. jejuni*, it was classified as 'no
polymorphism'. Otherwise, if the most frequent haplotype accounted for >90% among the
recipients, it was classified as either '*C. jejuni* (>90%)' or '*C. coli* (>90%)' if it was the same
as that of donor *C. jejuni* or *C. coli* (ancestral) (inset in Figure 3C). If the most frequent
haplotype accounted for ≤90% among the recipients, the top two most frequent haplotypes
(written as major and minor haplotype in this manuscript) were indicated as either "*C. jejuni* /
C. coli", "*C. jejuni* / other", "*C. coli* / *C. jejuni*", "*C. coli* / other", "other / *C. coli*", "other / *C.
jejuni*", and "other / other", and the frequency of the major and minor haplotypes were
calculated. For example, where the haplotype frequencies were as follows, AA=285,
TA=192, TG=181, AG=27, A-=2, --=1, -A=1, AA is the major haplotype, frequency of
which is 41.3%
540
Mutagenesis and complementation cloning
Genes *cj1167*, *cj1168c* (here designated *selF* for <u>sel</u>enium transport for <u>f</u>ormate
dehydrogenase), *cj1171c* (*ppi*), *cj1507c* (*modE*), *cj1508c* (*fdhD*) and *cj1500* (*fdhT*) were
deleted by allelic exchange mutagenesis, with the majority of the open reading frame
replaced by an antibiotic resistance cassette. Mutagenesis plasmids were generated by the
isothermal assembly method using the HiFi system (NEB, UK). In brief, flanking regions of
target genes were PCR amplified from genomic DNA using primers with adaptors
homologous to either the backbone vector pGEM3ZF or the antibiotic resistance cassette
(Supplementary Data 4). pGEM3ZF was linearised by digestion with HincII. The kanamycin
and chloramphenicol resistance cassettes were PCR amplified from pJMK30 and pAV35,
respectively[76]. Four fragments consisting of linearised pGEM3ZF, antibiotic resistance
cassette and 2 flanking regions were combined in equimolar amounts and mixed with 2 x
HiFi reagent (NEB, UK) and incubated at 50°C for 1 hour. The fragments combine such that
the gene fragments flank the antibiotic resistance cassette, in the same transcriptional
orientation, within the vector. Mutagenesis plasmids were transformed into *C. jejuni* NCTC
11168 by electroporation. Spontaneous double-crossover recombinants were selected for
using the appropriate antibiotic and correct insertion into the target gene confirmed by PCR
screening. For genetic complementation of mutants, genes *cj1168c* (*selF*), *cj1507c* (*modE*),
cj1508c (*fdhD*) and *cj1500* (*fdhT*) were PCR amplified from genomic DNA, restriction
digested with MfeI and XbaI, then ligated into similarly digested pRRA[77] (Supplementary
Data 4). The orientation of insertion allowed the target gene to be expressed constitutively by
a chloramphenicol resistance gene-derived promoter within the vector. Complementation
plasmids were transformed into *C. jejuni* by electroporation. Spontaneous double-crossover
recombinants were selected for using apramycin and correct insertion into the ribosomal
locus confirmed by PCR screening.

566
Growth of *C. jejuni*
Microaerobic growth cabinets (Don Whitley, UK) were maintained at 42^{o}C with an atmosphere of 10% O_2, 5% CO_2 and 85% N_2 (v/v). *C. jejuni* was grown on Columbia-base agar containing 5% v/v defibrinated horse blood. Selective antibiotics were added to plates as appropriate at the following concentrations: 50 µg ml^{-1} kanamycin, 20 µg ml^{-1} chloramphenicol, 60 µg ml^{-1} apramycin. Muller-Hinton (MH) broth supplemented with 20 mM L-serine was used as a rich medium. Minimal medium was prepared from a supplied MEM base (51200-38, Thermo Scientific, UK) with the following additions: 20 mM L-serine, 0.5 mM sodium pyruvate, 50 µM sodium metabisulfite, 4 mM L-cysteine. HCl, 2 mM L-methionine, 5 mM L-glutamine, 50 µM ferrous sulfate, 100 µM ascorbic acid, 1 µM vitamin B12, 5 µM sodium molybdate, 1 µM sodium tungstate. Selenium was then added as appropriate from stocks of sodium selenate or sodium selenite prepared in dH$_2$O. For assays, cells were washed and suspended in sterile phosphate-buffered saline (PBS, Sigma-Aldrich).

Respiration rates with formate
Cells were first grown in MH broth for 12 hours, then washed thoroughly in PBS before inoculating minimal media without an added selenium source. The appropriate concentrations were determined by serial dilution trials and it was subsequently found that *C. jejuni* has a strong preference for selenite over selenate, as equivalent FDH activity requires some 1000-fold greater concentration of selenate than selenite (Supplementary Figure 6). These cultures were grown for 8 hours before the cells were thoroughly washed again, then used to inoculate further minimal media, with a selenium source added as appropriate, and grown for 10 hours. This passaging was necessary to remove all traces of selenium from the inoculum, such that control cultures without selenium added had negligible (FDH) activity. Assay cultures were again thoroughly washed before the equivalent of 20 ml at an optical density of 0.8 at 600 nm was finally suspended in 1 ml of PBS. Formate-dependent oxygen consumption by whole cells was measured in a Clark-type electrode using 20 mM sodium formate as electron donor. The electrode was calibrated with air-saturated PBS assuming 220 nmol dissolved O_2 ml^{-1} at 37 °C. In the electrode, 200 µl of the dense cell suspension was added to 800 µl air-saturated PBS for a final volume of 1 ml. The chamber was sealed and the suspension allowed to equilibrate for 2 minutes. The assay was initiated by the addition of 20 µl of 1 M sodium formate (prepared in PBS) and the rate of oxygen consumption recorded for 90 s. The total protein concentration of the cell suspensions was determined by Lowry assay and the specific rate of formate-dependent oxygen consumption expressed as nmol oxygen consumed min^{-1} mg^{-1} total protein.

Data availability
Short-read sequence data for all isolates sequenced in this study are deposited in the sequence read archive (SRA) and can be found associated with NCBI BioProjects PRJNA689604 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA689604) and PRJEB11972 (https://www.ncbi.nlm.nih.gov/bioproject/PRJEB11972). These were augmented with 498 previously published genomes and assembled genomes are available on Figshare (doi.org/10.6084/m9.figshare.13521683). Accession numbers for all genomes are included in Supplementary Data 3. Source data are provided for this paper.

Acknowledgements

This work was supported by Wellcome Trust grants 088786/C/09/Z and Medical Research
Council (MRC) grants MR/M501608/1 and MR/L015080/1 awarded to SKS. Bacterial
sampling was funded by Food Standards Agency project FS101087. AJT was supported by a
Biotechnology and Biological Sciences Research Council grant (BB/S014497/1) awarded to
DJK. GM was supported by a NISCHR Health Research Fellowship (HF-14-13). JC was
supported by the ERC grant no. 742158. Computational calculations were performed using
computer servers from the MRC CLIMB project (UK), at National Institute of Genetics
(Japan) and the Human Genome Center of the Institute of Medical Science (University of
Tokyo, Japan) and at HPC Wales (UK).

Author Contributions
S.S., X.D. D.F. and K.Y. conceived and designed the study. S.S., N.W., A.V. and M.M.
collected samples. G.M., A.T., L.M., M.H. and B.P. carried out Laboratory work. B.P., M.H.,
K.J., M.M., J.P. and S.S. supported data archiving. G.M., A.T., X.D., K.Y., L.M., S.P., S.S.
and J.C. analysed the data. M.M., J.P., J.C., D.K. and D.F. contributed to data interpretation.
S.S., B.P., G.M., C. K., A.T. and D.K. wrote the paper.

Competing Interests
The authors declare no competing interests.

632
633
634

 Figure legends

 Figure 1. Population genomics of *C. jejuni* and *C. coli.* (A) Phylogenetic tree reconstructed
 using neighbour-joining on a whole-genome alignment of 973 *C. jejuni* and *C. coli* isolates.
 Introgressed *C. coli* clades are represented with red (CC-828, n=677) and purple (CC-1150,
 n=12) circles, unintrogressed clade 1 (n=35) is shown in pink, clade 2 (n=45) in yellow and
 clade 3 (n=58) in green. A set of 146 *C. jejuni* genomes belonging to 30 clonal complexes (4
 to 5 isolates per ST) are shown in blue. Recipient and donor populations, used to infer
 introgression in chromosome painting analysis, are indicated. The scale bar represents the
 number of substitutions per site. (B) Accumulation curves of the clade-specific pan-genome
 content of *Campylobacter* lineages, using the same colour code as panel A. Randomized
 genome sampling was used to obtain the average number of genes for each comparison (plain
 lines) and standard deviations (dotted lines). (C) The average number of genes/genome,
 identified by BLAST, is shown for *C. jejuni* and the different *C. coli* clades. Asterisks denote
 significant difference between distributions as inferred from a Dunn's multiple comparisons
 test after a Kruskal-Wallis test, as follows: ***: $p<0.001$; ****: $p<0.0001$.

 Figure 2. Genome-wide introgression from *C. jejuni* to *C. coli*. (A) Summary of
 introgressed *C. jejuni* SNPs in *C. coli* CC-828 (n=677, red) and CC-1150 (n=12, purple)
 genomes using ChromoPainter; the number of introgressed core SNPs (coloured histograms;
 left y-axis) and core genes (white histograms; right y-axis) for a range of recipient strains
 proportions (at least 1, more than 50% and more than 98%) is shown. (B) The number of
 genes with different frequencies of maximum SNP introgression/gene in *C. coli* as a function
 of gene frequency in *C. jejuni*. Highly introgressed genes in CC-828 and CC-1150 tend to be
 core in *C. jejuni*. (C) Density plot (n=1000 bins) of specific and shared introgression events
 in CC-828 (x-axis) and CC-1150 (y-axis). The frequency of SNP introgression/gene is shown
 for both lineages. Close blue lines denote a high density of points. (D) Shared introgression
 between *C. coli* CC-828 and CC-1150. The number of SNPs being shared between the two
 lineages at various frequencies is shown in y-axis. (E) Diagram of *Campylobacter* species
 and clade (C1*,C2, C3) divergence with arrows indicating introgression from *C. jejuni* into
 C. coli (i) clade 1, (ii, iii) CC-828 and CC-1150, (iv, v) subsequent clonal expansion and
 ongoing introgression. (F) Pairwise nucleotide identity between *C. jejuni* and ancestral
 (unintrogressed) clade 1 *C. coli* core genes (black circles). Genes found to be introgressed in
 clade 1 CC-828 are highlighted in blue.

 Figure 3. Covariation in introgressed *C. coli* genomes. (A) CC-828 and CC-1150 *C. coli*
 genomes were analysed using a continuous time Markov chain (CTMC) model and
 covariation was assessed for pairs of biallelic sites separated by at least 20kb along the
 genome while accounting for the effect of population structure and recombination. There are
 many biallelic sites that do not change often on the tree and few that do. Putative epistatic
 sites change more frequently than average with biallelic pairs found together on multiple
 branches. (B) Miami plot of each polymorphic site showing the maximum *p*-value for
 covarying biallelic pairs (>20kb apart) and the frequency of introgression in CC-828 and CC-
 1150. (C) The frequency of major and minor haplotype combinations (inset) among the 2578
 pairs of covarying SNPs in the 689 *C. coli* clade-1 recipient genomes*,* revealing that the
 majority of long range covariation was between introgressed *C. jejuni* sites. (D) The position
 of putative epistatic sites mapped on the *C. coli* CVM29710 reference for covarying *C.*
 jejuni-C. jejuni SNPs (red) in 16 gene pairs (a to l), and other haplotype combinations (grey).
 (E and F) An evolutionary scenario for the observed patterns of covariation and introgression
 in natural *C. coli* populations: (i) *C. jejuni* (blue) and unintrogressed *C. coli* (red) co-exist

with genomes (internal circles) harbouring haplotype pairs (x-x) that segregate by species; (ii)
Horizontal gene transfer, HGT, occurs (R1) disrupting covarying genetic elements and
reducing the relative fitness of introgressed *C. coli* to varying degrees (grey arrow), few
strains retain mixed *C. coli - C. jejuni* haplotypes; (iii) HGT continues (R2) and, where
recombined mixed haplotypes survived, ancestral *C. jejuni* haplotype pairs are reinstated in
introgressed *C. coli;* (iv,v) introgressed *C. coli* outcompete unintrogressed strains.

Figure 4. Genomic context and physiological roles of introgressed epistatically linked
genes. (A) Genome organisation and percentage of co-varying SNP pairs (internal legend).
(B-D) FDH activity of whole cells determined by oxygen consumption rates in a Clark-type
electrode (nmol oxygen consumed per minute per mg of total protein) for (B) cells grown in
rich media (excess selenium), (C) cells grown in minimal media with 0.5 nM sodium selenite,
and (D) cells grown in minimal media with either 5 nM sodium selenite (left, open bars) or 5
µM sodium selenate (right, hashed bars). All data are means of at least 4 independent
determinations and error bars are SD. *** denotes *p* value of <0.001 by students *t*-test. (E)
Model for epistatically linked genes involving FDH biogenesis and activity. Host derived
formate is converted to bicarbonate in the periplasm by the FDH complex to release electrons
which are transferred from the iron sulfur (FeS) cluster of FdhA to the *b*-type hemes of FdhC
and into the menaquinone (MK) pool where they can ultimately be used to reduce molecular
oxygen via terminal oxidases. FdhA contains a selenocysteine residue and Mo/W-pterin
cofactor (W/MoCo) at its active site, both of which are essential for catalysis. ModE is a
DNA binding regulator which represses the Mo and W transporters Mod and Tup to regulate
the cellular pool of Mo/W. W/MoCo is generated by the Moa pathway and inserted into apo-
FdhA by the sulphur-transferase FdhD. Environmental selenite (the most abundant oxyanion)
or selenate diffuses into the periplasm where it must be actively imported to the cytoplasm by
FdhT and SelF. Putatively, FdhT is low affinity and functions efficiently when ample
selenium is available. When selenium is limited, SelF can import sufficient selenium to
maintain FDH production and activity. Cytoplasmic selenium is converted to
selenophosphate by SelD, which is used by SelA to generate tRNA-SeC from tRNA-Ser.
During translation of FdhA, tRNA-SeC is incorporated by the specific elongation factor SelB.
Apo-FdhA with W/MoCo and SeC inserted is then transported to the periplasm by the TAT
translocation system and incorporated into the FDH complex (See [31] for a review of FDH in
C. jejuni).

718
719
720
721

References

722
1   Ochman, H., Lawrence, J. G. & Groisman, E. A. Lateral gene transfer and the nature
of bacterial innovation. *Nature* **405**, 299-304 (2000).
2   Koonin, E. V., Makarova, K. S. & Aravind, L. Horizontal gene transfer in
prokaryotes: quantification and classification. *Annu. Rev. Microbiol.* **55**, 709-742
(2001).
3   Haldane, J. B. A defense of beanbag genetics. *Perspect Biol Med* **7**, 343-359 (1964).
4   Bateson, W. in *Darwin And Modern Science*   85–101 (Cambridge University Press,
1909).
5   Fisher, R. A. The correlation between relatives on the supposition of Mendelian
inheritance. *Trans. R. Soc. Edinb* **52**, 399–433 (1918).
6   Wright, S. Evolution in Mendelian populations. *Genetics* **16**, 97-159 (1931).
7   Dobzhansky, T. Studies on hybrid sterility. II. Localization of sterility factors in
Drosophila Pseudoobscura hybrids. *Genetics* **21**, 113-135 (1936).
8   Dobzhansky, T. *Genetics And The Origin Of Species*. (Columbia University Press,
1937).
9   Muller, H. J. Bearing of the *Drosophila* work on systematics. *The new systematics*,
185-268 (1940).
10   Muller, H. Isolating mechanisms, evolution, and temperature. *Biol. Symp.* **6**, 71-125
(1942).
11   Kimura, M. Attainment of quasi linkage equilibrium when gene frequencies are
changing by natural selection. *Genetics* **52**, 875-890 (1965).
12   Neher, R. A. & Shraiman, B. I. Competition between recombination and epistasis can
cause a transition from allele to genotype selection. *Proc. Natl. Acad. Sci.* **106**, 6866-
6871 (2009).
13   Wilson, D. J. *et al.* Rapid evolution and the importance of recombination to the
gastroenteric pathogen *Campylobacter jejuni*. *Mol. Biol. Evol.* **26**, 385-397 (2009).
14   Gibson, B., Wilson, D. J., Feil, E. & Eyre-Walker, A. The distribution of bacterial
doubling times in the wild. *Proc. R. Soc. B.* **285** (2018).
15   Vos, M. & Didelot, X. A comparison of homologous recombination rates in bacteria
and archaea. *Isme J* **3**, 199-208 (2009).
16   Calland, J. K. *et al.* Quantifying bacterial evolution in the wild: a birthday problem for
Campylobacter lineages. *Preprint at*
https://www.biorxiv.org/content/10.1101/2020.12.02.407999v1 (2020).
17   Arnold, B. J. *et al.* Weak epistasis may drive adaptation in recombining bacteria.
Genetics **208**, 1247-1260 (2018).
18   Sheppard, S. K., Guttman, D. S. & Fitzgerald, J. R. Population genomics of bacterial
host adaptation. *Nat. Rev. Genet.* **19**, 549-565 (2018).
19   Mayr, E. in *Evolution as a Process*   157–180 (Allen & Unwin, 1954).
20   Orr, H. A. Dobzhansky, Bateson, and the genetics of speciation. *Genetics* **144**, 1331-
1335 (1996).
21   Fraser, C., Hanage, W. P. & Spratt, B. G. Recombination and the nature of bacterial
speciation. *Science* **315**, 476-480 (2007).
22   Sheppard, S. K. *et al. Campylobacter* genotyping to determine the source of human
infection. *Clin. Infect. Dis.* **48**, 1072-1078 (2009).
23   Sheppard, S. K., McCarthy, N. D., Falush, D. & Maiden, M. C. Convergence of
Campylobacter species: implications for bacterial evolution. *Science* **320**, 237-239
(2008).
24   Sheppard, S. K. *et al.* Progressive genome-wide introgression in agricultural
Campylobacter coli. *Mol. Ecol.* **22**, 1051-1064 (2013).

25 Provine, W. B. in *Genetics, Speciation And The Founder Principle*   43-76 (Oxford
University Press, 1989).
26 Cohan, F. M. in *Genetics of Mate Choice: From Sexual Selection to Sexual Isolation*
359-370 (Springer, 2002).
27 Ansari, M. A. & Didelot, X. Inference of the properties of the recombination process
from whole bacterial genomes. *Genetics* **196**, 253-265 (2014).
28 Retchless, A. C. & Lawrence, J. G. Phylogenetic incongruence arising from
fragmented speciation in enteric bacteria. *Proc. Natl. Acad. Sci.* **107**, 11453-11458
(2010).
29 Dearlove, B. L. *et al.* Rapid host switching in generalist *Campylobacter* strains erodes
the signal for tracing human infections. *Isme J* **10**, 721-729 (2016).
30 Weerakoon, D. R., Borden, N. J., Goodson, C. M., Grimes, J. & Olson, J. W. The role
of respiratory donor enzymes in *Campylobacter jejuni* host colonization and
physiology. *Microb. Pathog.* **47**, 8-15 (2009).
31 Taylor, A. J. & Kelly, D. J. The function, biogenesis and regulation of the electron
transport chains in *Campylobacter jejuni*: New insights into the bioenergetics of a
major food-borne pathogen. *Adv. Microb. Physiol.* **74**, 239-329 (2019).
32 Smart, J. P., Cliff, M. J. & Kelly, D. J. A role for tungsten in the biology of
Campylobacter jejuni: tungstate stimulates formate dehydrogenase activity and is
transported via an ultra-high affinity ABC system distinct from the molybdate
transporter. *Mol. Microbiol.* **74**, 742-757 (2009).
33 Doerrler, W. T., Sikdar, R., Kumar, S. & Boughner, L. A. New functions for the
ancient DedA membrane protein family. *J. Bacteriol.* **195**, 3-11 (2013).
34 Ledgham, F., Quest, B., Vallaeys, T., Mergeay, M. & Covès, J. A probable link
between the DedA protein and resistance to selenite. *Res. Microbiol.* **156**, 367-374
(2005).
35 Shaw, F. L. *et al.* Selenium-dependent biogenesis of formate dehydrogenase in
Campylobacter jejuni is controlled by the *fdhTU* accessory genes. *J. Bacteriol.* **194**,
3814-3823 (2012).
36 Rieseberg, L. H. Hybrid origins of plant species. *Annu. Rev. Ecol. Syst.* **28**, 359-389
(1997).
37 Andreani, N. A., Hesse, E. & Vos, M. Prokaryote genome fluidity is dependent on
effective population size. *Isme J* **11**, 1719-1721 (2017).
38 Vos, M. & Eyre-Walker, A. Are pangenomes adaptive or not? *Nature Microbiol.* **2**,
1576-1576 (2017).
39 Shapiro, B. J. The population genetics of pangenomes. *Nature Microbiol.* **2**, 1574-
1574 (2017).
40 McInerney, J. O., McNally, A. & O'Connell, M. J. Why prokaryotes have
pangenomes. *Nature Microbiol.* **2**, 17040 (2017).
41 Levin, B. R., Perrot, V. & Walker, N. Compensatory mutations, antibiotic resistance
and the population genetics of adaptive evolution in bacteria. *Genetics* **154**, 985-997
(2000).
42 Sheppard, S. K., McCarthy, N. D., Jolley, K. A. & Maiden, M. C. J. Introgression in
the genus *Campylobacter*: generation and spread of mosaic alleles. *Microbiology* **157**,
1066-1074 (2011).
43 Pensar, J. *et al.* Genome-wide epistasis and co-selection study using mutual
information. *Nucleic Acids Res.* **47**, e112-e112 (2019).
44 Croucher, N. J., Harris, S. R., Barquist, L., Parkhill, J. & Bentley, S. D. A high-
resolution view of genome-wide pneumococcal transformation. *PLoS Path.* **8**,
e1002745 (2012).

45   He, M. *et al.* Evolutionary dynamics of *Clostridium difficile* over short and long time
scales. *Proc. Natl. Acad. Sci.* **107**, 7527-7532 (2010).
46   Power, P. M., Bentley, S. D., Parkhill, J., Moxon, E. R. & Hood, D. W. Investigations
into genome diversity of *Haemophilus influenzae* using whole genome sequencing of
clinical isolates and laboratory transformants. *BMC Microbiol.* **12**, 273 (2012).
47   Yahara, K. *et al.* Genome-wide association of functional traits linked with
Campylobacter jejuni survival from farm to fork. *Environ. Microbiol.* **19**, 361-380
(2017).
48   Arnoux, P. *et al.* Sulphur shuttling across a chaperone during molybdenum cofactor
maturation. *Nat. Commun.* **6**, 6148 (2015).
49   Taveirne, M. E., Sikes, M. L. & Olson, J. W. Molybdenum and tungsten in
Campylobacter jejuni: their physiological role and identification of separate
transporters regulated by a single ModE-like protein. *Mol. Microbiol.* **74**, 758-771
(2009).
50   Aguilar-Barajas, E., Díaz-Pérez, C., Ramírez-Díaz, M. I., Riveros-Rosas, H. &
Cervantes, C. Bacterial transport of sulfate, molybdate, and related oxyanions.
BioMetals **24**, 687-707 (2011).
51   Thomas, M. T. *et al.* Two respiratory enzyme systems in *Campylobacter jejuni* NCTC
11168 contribute to growth on l-lactate. *Environ. Microbiol.* **13**, 48-61 (2011).
52   Ünal, C. M. & Steinert, M. Microbial peptidyl-prolyl cis/trans isomerases (PPIases):
virulence factors and potential alternative drug targets. *Microbiol. Mol. Biol. Rev.* **78**,
544-571 (2014).
53   Oren, Y. *et al.* Transfer of noncoding DNA drives regulatory rewiring in bacteria.
Proc. Natl. Acad. Sci. **111**, 16112-16117 (2014).
54   Arnold, B. *et al.* Fine-scale haplotype structure reveals strong signatures of positive
selection in a recombining bacterial pathogen. *Mol. Biol. Evol.* **37**, 417-428 (2019).
55   Wadsworth, C. B., Arnold, B. J., Sater, M. R. A. & Grad, Y. H. Azithromycin
resistance through interspecific acquisition of an epistasis-dependent efflux pump
component and transcriptional regulator in *Neisseria gonorrhoeae*. *mBio* **9**, e01419-
01418 (2018).
56   Mourkas, E. *et al.* Agricultural intensification and the evolution of host specialism in
the enteric pathogen *Campylobacter jejuni*. *Proc. Natl. Acad. Sci.* **117**, 11018-11028
(2020).
57   Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications
to single-cell sequencing. *Journal of computational biology : a journal of*
computational molecular cell biology **19**, 455-477 (2012).
58   Sheppard, S. K. *et al.* Genome-wide association study identifies vitamin B5
biosynthesis as a host specificity factor in *Campylobacter*. *Proc. Natl. Acad. Sci.* **110**,
11923-11927 (2013).
59   Sheppard, S. K. *et al.* Cryptic ecology among host generalist *Campylobacter jejuni* in
domestic animals. *Mol. Ecol.* **23**, 2442-2451 (2014).
60   Pascoe, B. *et al.* Local genes for local bacteria: evidence of allopatry in the genomes
of transatlantic *Campylobacter* populations. *Mol. Ecol.* **26**, 4497-4508 (2017).
61   Pascoe, B. *et al.* Genomic epidemiology of *Campylobacter jejuni* associated with
asymptomatic pediatric infection in the Peruvian Amazon. *PLoS Negl. Trop. Dis.* **14**,
e0008533 (2020).
62   Jolley, K. A. & Maiden, M. C. J. BIGSdb: scalable analysis of bacterial genome
variation at the population level. *BMC Bioinformatics* **11**, 595 (2010).
63   Dingle, K. E. *et al.* Multilocus sequence typing system for *Campylobacter jejuni*. *J.*
Clin. Microbiol. **39**, 14-23 (2001).

64   Méric, G. *et al.* A reference pan-genome approach to comparative bacterial genomics:
identification of novel epidemiological markers in pathogenic *Campylobacter*. *PLOS*
One **9**, e92798 (2014).
65   Aziz, R. K. *et al.* The RAST server: rapid annotations using subsystems technology.
BMC Genomics **9**, 75 (2008).
66   Sheppard, S. K., Jolley, K. A. & Maiden, M. C. J. A gene-by-gene approach to
bacterial population genomics: whole genome MLST of *Campylobacter*. *Genes* **3**,
261-277 (2012).
67   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – approximately maximum-
likelihood trees for large alignments. *PLOS One* **5**, e9490 (2010).
68   Didelot, X. & Wilson, D. J. ClonalFrameML: efficient inference of recombination in
whole bacterial genomes. *PLoS Comp. Biol.* **11**, e1004041 (2015).
69   Browning, B. L. & Browning, S. R. A unified approach to genotype imputation and
haplotype-phase inference for large data sets of trios and unrelated individuals. *Am. J.*
Hum. Genet. **84**, 210-223 (2009).
70   Yahara, K. *et al.* Chromosome painting *in silico* in a bacterial species reveals fine
population structure. *Mol. Biol. Evol.* **30**, 1454-1464 (2013).
71   Lawson, D. J., Hellenthal, G., Myers, S. & Falush, D. Inference of population
structure using dense haplotype data. *PLoS Genet.* **8**, e1002453 (2012).
72   Yahara, K., Didelot, X., Ansari, M. A., Sheppard, S. K. & Falush, D. Efficient
inference of recombination hot regions in bacterial genomes. *Mol. Biol. Evol.* **31**,
1593-1605 (2014).
73   Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood
phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* **59**, 307-321 (2010).
74   Mell, J. C., Shumilina, S., Hall, I. M. & Redfield, R. J. Transformation of natural
genetic variation into *Haemophilus influenzae* genomes. *PLoS Path.* **7**, e1002151-
e1002151 (2011).
75   Collins, C. & Didelot, X. A phylogenetic method to perform genome-wide association
studies in microbes that accounts for population structure and recombination. *PLoS*
Comp. Biol. **14**, e1005958 (2018).
76   van Vliet, A. H., Wooldridge, K. G. & Ketley, J. M. Iron-responsive gene regulation
in a *Campylobacter jejuni fur* mutant. *J. Bacteriol.* **180**, 5291-5298 (1998).
77   Cameron, A. & Gaynor, E. C. Hygromycin B and apramycin antibiotic resistance
cassettes for use in *Campylobacter jejuni*. *PLOS One* **9**, e95084 (2014).