A two-hit epistasis model prevents core genome disharmony in recombining bacteria

Aidan J. Taylor[1]†, Guillaume Méric[2]†‡, Koji Yahara[3], Ben Pascoe[2], Leonardos Mageiros[2], Evangelos Mourkas[2], Jessica K Calland[2], Santeri Puranen[4], Matthew D. Hitchings[5], Keith A. Jolley[6], Carolin M. Kobras[1], Nicola J. Williams[7], Arnoud H. M. van Vliet[8], Julian Parkhill[9], Martin C. J. Maiden[6], Jukka Corander[4,10,11], Laurence D Hurst[2], Daniel Falush[12], Xavier Didelot[13], David J. Kelly[1*], Samuel K. Sheppard[2,6*]

Affiliations

[1]School of Biosciences, University of Sheffield, Sheffield S10 2TN, United Kingdom;

[2]The Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK;

[3]Antimicrobial Resistance Research Centre, National Institute of Infectious Diseases, Tokyo, Japan;

[4]Department of Mathematics and Statistics, Helsinki Institute for Information Technology, University of Helsinki, Helsinki, Finland;

[5]Swansea University Medical School, Institute of Life Science, Swansea SA2 8PP, UK;

[6]Department of Zoology, University of Oxford, 11a Mansfield Road, Oxford OX1 3SZ, UK;

[7]Department of Epidemiology and Population Health, Institute of Infection & Global Health, University of Liverpool, Leahurst Campus, Wirral, UK;

[8]School of Veterinary Medicine, University of Surrey, Guildford GU2 7AL, UK;

[9]Department of Veterinary Medicine, University of Cambridge, Madingley Rd, Cambridge CB3 0ES, UK;

[10]Department of Biostatistics, University of Oslo, Oslo, Norway;

[11]Parasites and Microbes, Wellcome Sanger Institute, Cambridge, UK;

[12]The Centre for Microbes, Development and Health, Institut Pasteur of Shanghai, Shanghai, China;

[13]School of Life Sciences and Department of Statistics, University of Warwick, Coventry CV4 7AL, UK.

*Corresponding author(s): David J. Kelly, d.kelly@sheffield.ac.uk; Samuel K. Sheppard, s.k.sheppard@bath.ac.uk;

‡Present address: Cambridge Baker Systems Genomics Initiative, Baker Heart and Diabetes Institute, 75 Commercial Rd, Melbourne 3004, Victoria, Australia.

† Contributed equally.

Abstract

Recombination of short DNA fragments via horizontal gene transfer (HGT) can both introduce beneficial alleles and create disharmony in coadapted genomes (negative epistasis). Owing to a lack of protracted intragenomic co-evolution, negative epistatic costs of HGT into non-core (accessory) bacterial genomes are likely to be minimal. By contrast, for the core genome, recombination is expected to be rare because disruptive allelic replacement is likely to introduce negative epistasis. Why then is homologous recombination common in the core of bacterial genomes? To understand this enigma, we take advantage of an exceptional model system, the common enteric pathogens *Campylobacter jejuni* and *Campylobacter coli*, that are known for very high magnitude interspecies gene flow in the core genome. As expected, HGT does indeed disrupt co-adapted allele pairings (negative epistasis). However, multiple HGT events enable recovery of the recipient genome's co-adaption between the alleles, even in core metabolism genes (e.g. formate dehydrogenase). These findings demonstrate that, even for complex traits, genetic coalitions can be decoupled, transferred and independently reinstated in a new genetic background. There is a strong resemblance to the two-hit cancer model. Whether it be by multiple mutations or multiple HGT events, immediate harmful effects from an initial event need not preclude adaptive evolution. The fitness peak jumping problem in asexual lineages is thus less of problem than envisaged.

 Introduction

Mutation is the engine of genetic novelty, but for most bacteria adaptation also involves the

acquisition of DNA from other strains and species through horizontal gene transfer (HGT)[1]. In

some well documented cases, a single nucleotide substitution or acquisition of a small number of

genes, can prompt new evolutionary trajectories with striking outcomes such as the evolution of

virulent or antibiotic resistant strains[2]. With such dynamic genomic architecture, it may be

tempting (and possibly useful[3]) to consider genes as independent units that 'plug and play'

innovation into recipient genomes. This is clearly an oversimplification. In fact, genomes are

highly interactive wherein the effect of one allele depends on another (epistasis). Therefore, it is

likely that some horizontally introduced changes will disrupt gene networks and be costly to the

original coadapted genetic background, especially for complex phenotypes involving multiple

genes.

Understanding how epistasis influences the evolution of phenotype diversity has preoccupied

researchers since the origin of population genetics[4-10], with much emphasis placed upon the

relative amounts of recombination and epistatic effect sizes[11, 12]. In sexual populations, such as

outbreeding metazoans, genetic variation is shuffled at each generation. As a consequence of the

trend towards randomization between alleles (linkage equilibrium) it is unlikely that multiple

distinct epistatic allele combinations will be maintained in the same population and barriers to

gene flow, such as geographic isolation, may be required for marked phenotypic diversification[8].

In bacteria, rapid clonal reproduction allows multiple genomically distant beneficial allele

combinations to rise to high frequency in a single population. For example, in common enteric

bacteria such as *Escherichia coli, Salmonella enterica* and *Campylobacter jejuni* the doubling

time in the wild has been estimated at around 24 hours or less[13, 14]. Therefore, though HGT

occurs in these organisms[15], even in highly recombinogenic *C. jejuni*[13, 16], there will likely be

many millions of bacterial generations between recombination events at a given locus. This

allows mutations that are beneficial only in specific genetic backgrounds to establish in a single

population and linkage disequilibrium to form between different epistatic pairs[17].

In a coadapted genomic landscape, recombination is expected to have two antagonistic effects.

On the one hand, it could promote adaptation by conferring novel functionality on the recipient

genome[18-22] and reducing competition between clones that carry different beneficial mutations

(clonal interference). On the other, owing to the small average size of HGT transfer events and

their relatively low rate, it is also likely to introduce disharmonious allele-combinations that will

be discriminated against by selection[19].

The balance of these two effects is likely to be different for core and non-core HGT events.

Non-core events, such as the introduction of accessory antibiotic resistance genes, can be

immediately beneficial and, as they do not replace extant sequence, need not break epistatic

interactions. By contrast, HGT replacing one allele for another in part of the core genome is

unlikely to introduce great novelty but is likely to disrupt highly evolved co-adapted networks.

For these reasons, negative epistatic interactions between core genes with different evolutionary

histories have been proposed as a barrier to recombination[4-10, 23], particularly between species.

As bacteria typically have very high effective population sizes and hence efficient selection, the

expectation would be that HGT affecting core genes would then be selected against[19]. However,

interspecies recombination is common in bacterial core genomes[18, 24]. How, then, can a core

genome, that is expected to build up extensive co-adapted epistatic networks, be so accepting of

HGT events?

The common animal gut bacterium *Campylobacter,* which is among the most prolific causes of

human bacterial gastroenteritis worldwide[25], provides an exceptional model system to study the

impact of HGT on the core genome for several reasons. Introgression between the two most

important pathogenic species, *C. jejuni* and *C. coli,* has led to the evolution of a globally

distributed 'hybrid' *C. coli* lineage[26] that is responsible for almost all livestock and human

infections with this species. Up to 23% of the core genome of common *C. coli* has been recently

introgressed from *C. jejuni*[27], potentially disrupting epistatic interactions. As *C. jejuni* and *C. coli*

have undergone an extended period of independent evolution (85% average nucleotide identity),

recombined sequence is conspicuous in the genome.

Here, we investigate the disruptive effect of HGT on co-adapted bacterial core genomes by

examining covarying allele pairings and imported DNA fragments. Even though recombined

fragments enter the genome one-by-one, we find that most covariation in the core genome is

between sites where both alleles were imported. Having confirmed that allelic covariation is

indicative of epistasis, with laboratory mutagenesis and complementation assays, we conclude

that independent disruptive recombination events occur and persist until a second event restores

the functional link in a new genetic background. This process resembles the two-hit cancer

model. Both bacterial and cancer cell lines are asexual clones and a two-hit model for core

genome HGT is consistent with a more general theory in which, whether by multiple mutations

or multiple HGT events, immediate harmful effects from an initial event need not preclude

adaptive evolution.

Results

***Campylobacter* populations are highly structured with intermediate sequence clusters**

Analysis every automatically annotated gene from every genome, revealed a core genome of 631

gene orthologues in all *Campylobacter* isolates in this study. There were 1287 genes common to

all *C. jejuni* isolates and 895, 1021 and 1272 common to *C. coli* clades 1, 2 and 3, respectively.

Consistent with previous studies [27], neighbour-joining and ClonalFrameML trees based on genes

within concatenated core genome alignments revealed population structure in which *C. jejuni*

and *C. coli* clade 2 and 3 isolates each formed discrete clusters (Figure 1A, Figure S1). Isolates

designated as *C. coli* clade 1 were found in three clusters on the phylogeny: unintrogressed

ancestral strains, and the ST-828 and ST-1150 clonal complexes (CCs) which account for the

great majority of strains found in agriculture and human disease[27]. Pan-genome analysis

quantified the increase in unique gene discovery as the number of sampled genomes increased

(Figure S2A). For all sequence clusters there was evidence of an open pan-genome with a trend

towards continued rapid gene discovery within sequence clusters with fewer isolates. There was

considerable accessory genome variation between species and clades, potentially associated with

important adaptive traits (Figure S3) and there was evidence that the average number of genes

per genome was greater in *C. coli* CC-828 isolates than in *C. jejuni* (Figure S2B).

There is substantial introgression within the *C. coli* core genome

The large genetic distances among *C. coli* clade 1 isolates (Figure 1A), have been shown to be a

consequence of the import of DNA from *C. jejuni* rather than accumulation of mutations during a

prolonged period of separate evolution[26]. Using chromosome painting to infer the co-ancestry of

core-genome haplotype data from CC-828 and CC-1150 isolates gave a detailed representation

of the recombination-derived chunks from each *C. jejuni* donor group and clonal complex to

each recipient individual (Figure S4). The majority of introgressed SNPs were rare, occurring in

fewer than 50 recipient genomes (Figure 1B). However, a large proportion of the introgressed *C.*

coli clade 1 genomes contained DNA of *C. jejuni* ancestry in >98% of recipient isolates.

Consistent with previous estimates[27], these regions where introgressed DNA was largely fixed

within the *C. coli* population, occurred across the genome and comprised up to 15% and 28% of

the CC-828 and CC-1150 isolate genomes respectively (Figure 1B). When considering donor

groups, the majority of introgressed DNA in *C. coli* involves genes that are present in multiple *C.*

jejuni lineages (core genes) (Figure S5A).

Having identified *C. jejuni* ancestry within *C. coli* genomes, we investigated the sequence of

events responsible for introgression. Most introgressed SNPs are found at low frequency in both

clonal complexes. However, there was evidence of SNPs that are introgressed in both complexes

as well as high frequency lineage specific introgression in both CC-828 and CC-1150 (Figure

S5B). Specifically, 25% of the *C. jejuni* DNA found in >98% of CC-828 isolates was also found

in CC-1150 (Figure S5C), implying that this genetic material was imported by the common

ancestor(s) of both complexes. After the divergence of these two complexes, introgression

continued with nearly 75% of *C. jejuni* DNA present in one recipient clonal complex and not the

other. This is consistent with an evolutionary history in which there was a period of progressive

species and clade divergence reaching approximately 12% at the nucleotide level

between *C. jejuni* and *C. coli* and around 4% between the three *C. coli* lineages. More recently,

changes to the patterns of gene flow led one *C. coli* clade 1 lineage to import substantial

quantities of *C. jejuni* DNA, and further lineage-specific introgression gave rise to two clonal

complexes (CC-828 and CC-1150) that continued to accumulate *C. jejuni* DNA, independently

creating the population structure observed today (Figure 1C).

The high magnitude introgression into *C. coli* clade 1 isolates has introduced thousands of

nucleotide changes to the core genome. However, divergence in bacteria may be uneven across

the genome for at least two reasons. First, recombination is more likely to occur in regions where

donor and recipient genomes have high nucleotide similarity[24, 28, 29]. Second, 'fragmented

speciation'[30], in which gene flow varies in different parts of the genome, such as regions

responsible for adaptive divergence, can result in phylogenetic incongruence among genes.

Consistent with previous estimates[27], we found that the three *C. coli* clades had similar high

divergence with *C. jejuni* across the genome, ranging from 68% to 98% nucleotide identity for

individual genes (Figure S5D), implying a period of divergence with low levels of gene flow.

We found no evidence that high genetic differentiation between the species prevented

recombination. While there was some evidence that more recombination occurred in regions of

low nucleotide divergence (between unintrogressed *C. coli* clade 1 and *C. jejuni*), introgression

occurred across the genome at sites with varying levels of nucleotide identity (Figure S5D). This

level of recombination has greatly increased overall genetic diversity across the genome

in *C. coli* clade 1 and introduced changes that have potential functional significance.

Much of the putative epistasis occurs between SNPs in introgressed genes

ClonalFrameML analysis revealed the importance of homologous recombination in generating sequence variation within the introgressed *C. coli*. Estimates of the relative frequency of recombination versus mutation (R/θ=0.43), mean recombination event length (δ=152bp) and average amount of polymorphism per site in recombined fragments (ν=0.07), imply that recombination has had an effect (*r/m*) 4.57 times higher than *de novo* mutation during the diversification of CC-828. This is consistent with previous analysis and confirmed recombination as the major driver of molecular evolution in *C. coli*[13, 27, 31]. The continuous time Markov chain model for the joint evolution of pairs of biallelic sites on a phylogenetic tree (Figure S6) was applied to investigate patterns of covariance for all pairs of sites >20kb apart (Figure 2A). For most biallelic sites there were few branches on the tree where substitutions occurred, so that their evolution is compatible with separate evolution on the same clonal frame. However, 2874 covarying pairs evolved more frequently together than would be expected (*p*-value 10^{-8}) if they had evolved independently based on the tree, and hence indicated patterns of putative epistasis.

Among them, the location of 2618 putative epistatic pairs of sites was compared to the inferred ancestry (unintrogressed *C. coli* or *C. jejuni*) of sequence across the genome of CC-828 and CC-1150 *C. coli* strains (Figure 2B, Data S1). For each epistatic pair, the major and minor haplotype were defined if there was haplotype polymorphism between *C. jejuni* and CC-828 and CC-1150 *C. coli*. This allowed quantification of the number of covarying sites that occurred between an ancestral *C. coli* (unintrogressed) and an introgressed *C. jejuni* allele, two introgressed alleles, and sites that do not segregate by species. Strikingly, the breakdown of the major and minor haplotype combinations among the 2618 epistatic pairs (Figure 2C, Data S2) shows the major

231 haplotype for 83.5% of putative epistatic SNP pairs was *C. jejuni* indicating that both co-varying

232 sites had *C. jejuni* ancestry, consistent with epistasis between introgressed ancestral *C. jejuni*

233 sequence at divergent genomic positions. Investigation of the genes containing co-varying sites

234 revealed that 2187 SNP pairs were in 16 genes with just five genes accounting for 99.1% of them

235 (Figure 2D, Data S3, Figure S7).

Genomic context and physiological role of epistatically linked genes

The five genes accounting for the majority of epistatic interactions (*cj1167*, *cj1168c*,

cj1171c/ppi, *cj1507c/modE* and *cj1508c/fdhD*) were investigated for their physiological role in

C. jejuni. FdhD and ModE are proteins involved in the biogenesis of formate dehydrogenase

(FDH). The FDH complex (FdhABC) oxidises formate to bicarbonate to generate electrons that

fuel cellular respiration. Formate is an abundant electron donor produced by host microbiota and

an important energy source for *Campylobacter in vivo*[32, 33]. The remaining three genes, *cj1167*

(annotated incorrectly as *ldh*, lactate dehydrogenase), *cj1168c* and *cj1171c* (*ppi*) are also grouped

together in the genome, where *cj1167* and *cj1168c* are adjacent but with the open reading frames

(ORFs) convergent and overlapping, while *ppi* is upstream, separated by two non-epistatically

linked genes (*cj1169c* and *cj1170c*, Figure 3A). Considering the genomic arrangement, it is

therefore clear that the putative epistatic links uncovered in this study essentially occur across

two loci in the genome (*fdhD/modE* and *cj1167/cj1168c/ppi*), with each of the latter three genes

linked with both *fdhD* and *modE* (Figure 3A). Given the known function of *fdhD* and *modE* in

biogenesis of the FDH complex, we hypothesised that *cj1167/cj1168c/ppi* might also have some

role in FDH biogenesis or activity in order to form a functional epistatic connection. We

therefore constructed deletion mutants to investigate the possible role of these genes in FDH

activity.

Initially, each of the mutants and their parental wildtype (*C. jejuni* NCTC11168) were grown in

rich media (Muller-Hinton broth) and their formate dependent oxygen consumption rates

determined (Figure 3B). *cj1167, cj1168c* and *ppi* mutants demonstrated wildtype levels of FDH

activity, while activity in both *fdhD* and *modE* mutants was abolished. To confirm that the

phenotype of the *fdhD* and *modE* mutants was not due to a polar effect on the surrounding *fdh*

locus, these mutants were genetically complemented by reintroduction of a second copy of the

wildtype gene into the rRNA locus, which restored near wildtype levels of FDH activity in both

cases (Figure 3B).

As neither *cj1167, cj1168c* or *ppi* mutants showed altered FDH activity in cells grown in rich

media, we considered that their function may be related to an FDH-specific nutrient requirement

as would likely be found *in vivo*. Since the formate oxidising subunit of FDH, FdhA, specifically

requires a molybdo- or tungsto-pterin (Mo/W) cofactor and a selenocysteine (SeC) residue for

catalysis[34], Mo, W or Se supply presented possible targets. *cj1168c* encodes a DedA family

integral membrane protein of unknown function. DedA proteins are solute transporters

widespread in bacteria but are mostly uncharacterised[35]. However, a homologue of *cj1168c* in

the heavy metal specialist beta-proteobacterium *Cupriavidus metallidurans* has been shown to be

involved in selenite (SeO_3^{2-}) uptake[36]. We therefore speculated that Cj1168 could be a selenium

oxyanion transporter that supplies Se for SeC biosynthesis. To test this, FDH activities were

measured in *cj1168c* mutant and parental wildtype strains grown in minimal media with limiting

concentrations of selenite or selenate (SeO_4^{2-}) (Figure S8). The data in Figure 3C shows that the

cj1168c mutant displayed significantly reduced FDH activity after growth with selenite in the

low nM range, and this phenotype was partially restored by genetic complementation. We

therefore designated *cj1168c* as *selF* (<u>se</u>lenium transporter for <u>f</u>ormate dehydrogenase).

However, although this phenotype does suggest that SelF is a selenium importer, another

unrelated selenium transporter, FdhT (Cj1500), has previously been documented in *C. jejuni* [37],

which is not epistatic with *fdhD* or *modE*. In contrast to this previous report we found

considerable residual FDH activity still remained in an *fdhT* deletion mutant, which was fully

restored to wildtype levels by complementation (Figure 3D).

Finally, we tested whether the residual FDH activity in our *fdhT* mutant was due to selenium

uptake by SelF. An *fdhT selF* double mutant was generated and assayed for FDH activity after

growth in minimal media containing limiting concentrations of selenite or selenate (Figure 3D).

The *fdhT selF* double mutant demonstrated a significant additional reduction in FDH activity

over the *fdhT* single mutant, a phenotype that was partially restored by complementing the

double mutant with *selF*. Complementation of the double mutant with *fdhT* returned FDH

activity to near wildtype levels (Figure 3D). Taken together, our data suggests that both FdhT

and SelF facilitate selenium acquisition in *C. jejuni*, possibly representing low and high affinity

transporters, respectively (Figure S9).

Discussion

To address the apparent contradiction of frequent core genome recombination in a co-adapted

background we focused on *Campylobacter,* in which interspecies recombination is well

documented[26, 27, 38]. As in other studies, we found that a large proportion (15-28%) of the *C. coli*

core genome originated in *C. jejuni* despite the genetic distance (~85% nucleotide identity)

between the species. Investigating the likely disruptive impact on coadapted epistatic gene

networks, we quantified the frequency of *C. jejuni – C. coli* (and *vice versa*) and *C. jejuni – C.*

jejuni covarying allele pairs in introgressed *C. coli*. Where recombination is minimally disruptive

there would be more *C. jejuni – C. coli* than *C. jejuni – C. jejuni*. However, consistent with

selection against disharmonious gene combinations we found that *C. jejuni – C. jejuni* allele

pairs constituted >83% of covarying introgressed haplotypes. It is possible that in some cases

both sites were introgressed in a single large recombination event[39-41] but in *Campylobacter* LD

for pairs of sites decreases with distance and is approximately constant after 20kbp[42] consistent

with the independent acquisition of allele pairs (>20kb). It follows, therefore, that the first

introgression event was not fatal to the recipient genome. Acquisition of the second member of

the pair then potentially restored the function of the integrated *C. jejuni – C. jejuni*

coevolutionary unit.

Coadaptation and epistasis were confirmed for the most common co-varying *C. jejuni – C. jejuni*

gene pairs linked to FDH, a key enzyme allowing the utilisation of formate as an electron donor

in vivo[32, 33]. FdhD and ModE were shown to be essential for FDH activity. While FdhD is a

sulfur-transferase known to be required for the insertion of the pterin cofactor into FdhA[43]

(Figure S9), ModE is a transcriptional repressor that has been shown previously only to regulate

the Mo/W uptake genes *mod* and *tup*[44, 45]. However, the unexpected abolished FDH activity in a

modE mutant indicates further functions for ModE in FDH biogenesis which warrant future

investigation. Searching for functional links between *fdhD/modE* and *cj1167/selF/ppi*, revealed

that a *selF* mutant strain had significantly reduced FDH activity under conditions of selenite

limitation, a phenotype consistent with SelF being a Se oxyanion transporter and that

functionally links SelF with FdhD/ModE. We suggest that *selF* rather than *fdhT* is epistatic

because SelF confers an additional benefit for SeC biosynthesis (essential for FDH activity)

under conditions of selenium limitation, for example as may be found in the host (Figure S9).

cj1167 encodes a cytoplasmic NADPH dependent 2-oxoacid dehydrogenase but the current

genome database annotation as lactate dehydrogenase (Ldh) is incorrect and its function is

unknown[46]. There is no precedent for the involvement of such an enzyme in bacterial FDH or

SeC biogenesis and we obtained no evidence for a functional connection between Cj1167 and

FDH activity. However, the overlapping convergent gene arrangement of *cj1167* and *cj1168c*

(*selF*) suggests a transcriptional architecture that might dictate these genes both form similar

epistatic dependencies even if Cj1167 is not required for FDH activity. Finally, *cj1171c* (*ppi*)

encodes a cytoplasmic peptidyl-prolyl *cis-trans* isomerase of the cyclophilin family. PPIases are

general protein folding catalysts that often have pleiotropic and redundant functions[47] and we

note that our *C. jejuni ppi* deletion mutant showed no growth defect as well as no reduction in

FDH activity. It is possible that if Cj1171 does help promote the folding of e.g. FdhD or ModE,

analysis of a simple deletion mutant may not reveal this if another PPIase can substitute in that

genetic background.

Understanding the functional significance of core genome recombination has potential to explain

the evolution of complex phenotypes. In *Campylobacter,* our results suggest that an ancestral *C.*

coli lineage colonized a new niche and surviving lineages (CC-828 and CC-1150) gained access

to *C. jejuni* DNA (Figure 4A&B). As the adaptive landscape of the genome changed, potentially

decoupling epistatic interactions that were previously selected, new gene combinations could be

introduced by homologous recombination and tested in the *C. coli* genetic background.

The notion that multiple events are necessary to achieve a phenotype with the first event

potentially being deleterious (causing negative epistatsis) is strongly reminiscent of the two-hit

cancer model in eukaryotes[48] in which the benefit to the tumour cell appears only after earlier

non-advantageous mutations. In bacteria, while the first hit (HGT event) in a core genome, is

consistent with fitness costs associated with negative epistasis (breaking co-adapted gene

networks), this can be rescued by a second HGT event. The second event, in effect restores a

pre-existing co-adapted allele pair from the donor species to the recipient. This type of genetic

rewiring may indeed be more common than previously thought [49-51] and, despite theoretical

expectations, negative epistasis is not an absolute barrier to genome-wide recombination in

structured bacterial populations. Multiple HGT events thus provide a solution to the peak

jumping problem.

Materials and Methods

Isolates, genome sequencing and assembly

A total of 973 isolates were used in this study, 827 from *C. coli* and a selection of 146 from a

diversity of *C. jejuni* clonal complexes (Data S4). Isolates were sampled mostly in the United

Kingdom to maximise environmental and riparian reservoirs and thus the representation of

genetic diversity in *C. coli*. Isolates were stored in a 20% (v/v) glycerol medium mix at -80°C

and subcultured onto *Campylobacter* selective blood-free agar (mCCDA, CM0739, Oxoid).

Plates were incubated at 42°C for 48 h under microaerobic conditions (5% CO_2, 5% O_2)

generated using a CampyGen (CN0025, Oxoid) sachet in a sealed container. Subsequent

phenotype assays were performed on Brucella agar (CM0271, Oxoid). Colonies were picked

onto fresh plates and genomic DNA extraction was carried out using the QIAamp® DNA Mini

Kit (QIAGEN; cat. number: 51306) according to the manufacturer's instructions. DNA was

eluted in 100–200 μl of the supplied buffer and stored at −20°C. DNA was quantified using a

Nanodrop spectrophotometer and high-throughput genome sequencing was performed on a

MiSeq (Illumina, San Diego, CA, USA), using the Nextera XT Library Preparation Kit with

standard protocols involving fragmentation of 2 μg genomic DNA by acoustic shearing to enrich

for 600 bp fragments, A-tailing, adapter ligation and an overlap extension PCR using the

Illumina 3 primer set to introduce specific tag sequences between the sequencing and flow cell

binding sites of the Illumina adapter. DNA cleanup was carried out after each step to remove

DNA < 150 bp using a 1:1 ratio of AMPure® paramagnetic beads (Beckman Coulter, Inc.,

USA). Short read paired-end data was assembled using the *de novo* assembly algorithm, SPAdes

(version 3.10.0)[52]. All novel genome sequences (n=475) generated for use in this study are

available on NCBI BioProjects PRJNA689604 and PRJEB11972. These were augmented with

498 previously published genomes and accession numbers for all genomes can be found in Data

S4[16, 27, 31, 42, 53-57].

Genome archiving, pan-genome content analyses and phylogenetic reconstruction

Contiguous genome sequence assemblies were individually archived on the web-based database

platform BIGSDB[58] and sequence type (ST) and clonal complex (CC) designation were assigned

based upon the *C. jejuni* and *C. coli* multi-locus sequence typing scheme[59]. To examine the full

pan-genome content of the dataset, a reference pan-genome list was assembled as previously

described[60]. Briefly, genome assemblies from all 973 genomes in this study were automatically

annotated using the RAST/SEED platform[61], the BLAST algorithm was used to determine

whether coding sequences from this list were allelic variants of one another or 'unique' genes,

with two alleles of the same gene being defined as sharing >70% sequence identity on >10% of

the sequence length. The prevalence of each gene in the collection of 973 genomes was

determined using BLAST with a positive hit in a genome being defined as a local alignment of

the reference sequence with the genomic sequence of >70% identity on >50% of the length, as

previously described[62]. The resulting matrix was analysed for differentiating core and accessory

genome variation. Genes present in all genomes were concatenated to produce a core-genome

alignment, used for subsequent phylogenetic reconstructions. Phylogenetic trees were

reconstructed using an approximation of maximum-likelihood phylogenetics in FastTree2[63]. This

tree was used as an input for ClonalFrameML[64] to produce core genome phylogenies with branch

lengths corrected for recombination.

Inference of introgression

All 973 genomes were aligned to a full reference sequence of *C. coli* strain CVM29710. We

conducted imputation for polymorphic sites with missing frequency $\leq$10% using BEAGLE[65] as

previously reported[66]. A total of 286,393 gapless SNPs (~17% of the average *C. coli* genome

size) were used for recombination analyses. The coancestry of genome-wide haplotype data was

inferred based on alignments using chromosome painting and FineStructure[67] as previously

described[68]. Briefly, ChromoPainter was used to infer chunks of DNA donated from a list of 33

donor groups normalised for sample size to each of 677 ST-CC-828 and 12 CC-1150 recipient

haplotypes. Results were summarised into a coancestry matrix containing the number of

recombination-derived chunks from each donor to each recipient individual. FineStructure was

then used for 100,000 iterations of both the burn-in and Markov chain Monte Carlo chain to

cluster individuals based on the co-ancestry matrix. The results are visualized as a heat map with

each cell indicating the proportion of DNA "chunks" (a series of SNPs with the same expected

donor) a recipient receives from each donor.

Analysis of covariation in bacterial genomes

Non-random allele associations can result from selection and clonal population structure. To

control for the latter, our approach identified SNP combinations in independent genetic

backgrounds by accounting for the sequence variation associated with the inferred phylogeny.

Based on the alignment of 677 genomes of *C. coli* CC-828, a first phylogenetic tree was created

using PhyML[69]. ClonalFrameML[64] was then applied to correct the tree by accounting for the

effect of recombination, and also to infer the ancestral sequence of each node. Covariance was

assessed for pairs of biallelic sites across the genome using a Continuous Time Markov Chain

(CTMC) model as follows. Briefly, let *A* and *a* denote the two alleles of the first site and *B* and *b*

denote the two alleles of the second site, so that there are four states in total for the pair of sites

(*ab*, *Ab*, *aB* and *AB*). The four substitution rates from *A* to *a*, from *a* to *A*, from *B* to *b* and from *b*

to *B* are not assumed to be identical, to allow for differences in substitution rates in different

parts of the genome and also to allow for non-equal rates of forward and backward substitution

(for example as a result of recombination opportunities). Assuming no epistatic effect between

the two sites ($\varepsilon=1$), the model M_0 has four free parameters (α_1, α_2, β_1 and β_2) representing

independent substitutions at the two sites. We expand model M_0 with an additional fifth

parameter $\varepsilon>1$ into model M_1 which is such that the state *AB* where the first site is allele *A* and

the second site is allele *B* is favored relative to the other three sites *ab, aB* and *Ab*. Specifically,

the state *AB* has a probability increased by a factor ε^2 in the stationary distribution of the CTMC

of model M_1 compared to model M_0.

Both models M_0 (with 4 parameters) and M_1 (with 5 parameters) are fitted to the data using

maximum likelihood techniques, where the likelihood is equal to the product for every branch of

the tree of the state at the bottom of the branch given the state at the top. The two fitted models

M_0 and M_1 are then compared using a likelihood-ratio test (LRT) as follows: since M_0 is nested

with M_1, two times their difference in log-likelihood is expected to be distributed according to a

chi-square distribution with number of degrees of freedom equal to the difference in their

dimensionality, which is one. This LRT returns a *p*-value for the significance of a covariation

effect, and a Bonferroni correction is applied to determine a conservative cutoff of significance

that accounts for multiple testing. Furthermore, the test is applied only to pairs of sites separated

by >20kb to reduce the chance that they were the result of a single recombination event,

consistent with estimates of the length of recombined DNA sequence in quantitative bacterial

transformation experiments[39, 70] and evidence from *Campylobacter* genome analyses that show

that LD for pairs of sites decreases with distance to approximately 20kbp and then remains at the

same level for very distant sites[42]. It is still possible of course that rare recombination events

would stretch 20kbp[39-41], but for this to have an effect on the analysis of epistasis it would have

to have happened several times for the same pairs of sites against different genomic background

which becomes quite unlikely just by chance. This phylogenetically aware approach to testing

for covariance presents the advantage to naturally account for both population structure and the

effect of recombination[71]. The script implementing this coevolution test is available in R at:

https://github.com/xavierdidelot/campy.

Quantifying covariation between recombined and unrecombined genomic regions

The results of the introgression and covariation analyses were combined so that for each pair of

significantly covarying SNPs (p-value $<10^{-8}$), haplotype frequency was calculated among the 689

recipient introgressed *C. coli* clade-1 strains as well as among the donor *C. coli* (ancestral) and

C. jejuni strains, respectively. If the most frequent haplotype of the pair is the same between the

donor *C. coli* (ancestral) and *C. jejuni*, it was classified as 'no polymorphism'. Otherwise, if the

most frequent haplotype accounted for >90% among the recipients, it was classified as either '*C.*

jejuni (>90%)' or '*C. coli* (>90%)' if it was the same as that of donor *C. jejuni* or *C. coli*

(ancestral) (inset in Figure 2C). If the most frequent haplotype accounted for ≤90% among the

recipients, the top two most frequent haplotypes (written as major and minor haplotype in this

manuscript) were indicated as either "*C. jejuni* / *C. coli*", "*C. jejuni* / other", "*C. coli* / *C. jejuni*",

"*C. coli* / other", "other / *C. coli*", "other / *C. jejuni*", and "other / other", and the frequency of

the major and minor haplotypes were calculated. For example, where the haplotype frequencies

were as follows, AA=285, TA=192, TG=181, AG=27, A-=2, --=1, -A=1, AA is the major

haplotype, frequency of which is 41.3%

Mutagenesis and complementation cloning

Genes *cj1167*, *cj1168c* (here designated *selF* for <u>se</u>lenium transport for <u>f</u>ormate dehydrogenase),
cj1171c (*ppi*), *cj1507c* (*modE*), *cj1508c* (*fdhD*) and *cj1500* (*fdhT*) were deleted by allelic
exchange mutagenesis, with the majority of the open reading frame replaced by an antibiotic
resistance cassette. Mutagenesis plasmids were generated by the isothermal assembly method
using the HiFi system (NEB, UK). In brief, flanking regions of target genes were PCR amplified
from genomic DNA using primers with adaptors homologous to either the backbone vector
pGEM3ZF or the antibiotic resistance cassette (Data S5). pGEM3ZF was linearised by digestion
with HincII. The kanamycin and chloramphenicol resistance cassettes were PCR amplified from
pJMK30 and pAV35, respectively[72]. Four fragments consisting of linearised pGEM3ZF,
antibiotic resistance cassette and 2 flanking regions were combined in equimolar amounts and
mixed with 2 x HiFi reagent (NEB, UK) and incubated at 50°C for 1 hour. The fragments
combine such that the gene fragments flank the antibiotic resistance cassette, in the same
transcriptional orientation, within the vector. Mutagenesis plasmids were transformed into *C.
jejuni* NCTC 11168 by electroporation. Spontaneous double-crossover recombinants were
selected for using the appropriate antibiotic and correct insertion into the target gene confirmed
by PCR screening. For genetic complementation of mutants, genes *cj1168c* (*selF*), *cj1507c*
(*modE*), *cj1508c* (*fdhD*) and *cj1500* (*fdhT*) were PCR amplified from genomic DNA, restriction
digested with MfeI and XbaI, then ligated into similarly digested pRRA[73] (Data S5). The
orientation of insertion allowed the target gene to be expressed constitutively by a
chloramphenicol resistance gene-derived promoter within the vector. Complementation plasmids
were transformed into *C. jejuni* by electroporation. Spontaneous double-crossover recombinants
were selected for using apramycin and correct insertion into the ribosomal locus confirmed by
PCR screening.

Growth of *C. jejuni*

Microaerobic growth cabinets (Don Whitley, UK) were maintained at 42°C with an atmosphere of 10% O_2, 5% CO_2 and 85% N_2 (v/v). *C. jejuni* was grown on Columbia-base agar containing 5% v/v defibrinated horse blood. Selective antibiotics were added to plates as appropriate at the following concentrations: 50 µg ml^{-1} kanamycin, 20 µg ml^{-1} chloramphenicol, 60 µg ml^{-1} apramycin. Muller-Hinton (MH) broth supplemented with 20 mM L-serine was used as a rich medium. Minimal medium was prepared from a supplied MEM base (51200-38, Thermo Scientific, UK) with the following additions: 20 mM L-serine, 0.5 mM sodium pyruvate, 50 µM sodium metabisulfite, 4 mM L-cysteine. HCl, 2 mM L-methionine, 5 mM L-glutamine, 50 µM ferrous sulfate, 100 µM ascorbic acid, 1 µM vitamin B12, 5 µM sodium molybdate, 1 µM sodium tungstate. Selenium was then added as appropriate from stocks of sodium selenate or sodium selenite prepared in dH_2O. For assays, cells were washed and suspended in sterile phosphate-buffered saline (PBS, Sigma-Aldrich).

Respiration rates with formate

Cells were first grown in MH broth for 12 hours, then washed thoroughly in PBS before inoculating minimal media without an added selenium source. The appropriate concentrations were determined by serial dilution trials and it was subsequently found that *C. jejuni* has a strong preference for selenite over selenate, as equivalent FDH activity requires some 1000-fold greater concentration of selenate than selenite (Figure S8). These cultures were grown for 8 hours before the cells were thoroughly washed again, then used to inoculate further minimal media, with a selenium source added as appropriate, and grown for 10 hours. This passaging was necessary to

remove all traces of selenium from the inoculum, such that control cultures without selenium

added had negligible (FDH) activity. Assay cultures were again thoroughly washed before the

equivalent of 20 ml at an optical density of 0.8 at 600 nm was finally suspended in 1 ml of PBS.

Formate-dependent oxygen consumption by whole cells was measured in a Clark-type electrode

using 20 mM sodium formate as electron donor. The electrode was calibrated with air-saturated

PBS assuming 220 nmol dissolved O_2 ml^{-1} at 37°C. In the electrode, 200 µl of the dense cell

suspension was added to 800 µl air-saturated PBS for a final volume of 1 ml. The chamber was

sealed and the suspension allowed to equilibrate for 2 minutes. The assay was initiated by the

addition of 20 µl of 1 M sodium formate (prepared in PBS) and the rate of oxygen consumption

recorded for 90 s. The total protein concentration of the cell suspensions was determined by

Lowry assay and the specific rate of formate-dependent oxygen consumption expressed as nM

oxygen consumed min^{-1} mg^{-1} total protein.

References

1.   Ochman, H., Lawrence, J.G., and Groisman, E.A. (2000). Lateral gene transfer and the nature of bacterial innovation. Nature *405*, 299-304.

2.   Koonin, E.V., Makarova, K.S., and Aravind, L. (2001). Horizontal gene transfer in prokaryotes: quantification and classification. Annu. Rev. Microbiol. *55*, 709-742.

3.   Haldane, J.B. (1964). A defense of beanbag genetics. Perspect Biol Med *7*, 343-359.

4.   Bateson, W. (1909). Heredity And Variation In Modern Lights. In Darwin And Modern Science, A.C. Seward, ed. (Cambridge: Cambridge University Press), pp. 85–101.

5.   Fisher, R.A. (1918). The correlation between relatives on the supposition of Mendelian inheritance. Trans. R. Soc. Edinb *52*, 399–433.

6.   Wright, S. (1931). Evolution in Mendelian populations. Genetics *16*, 97-159.

7.   Dobzhansky, T. (1936). Studies on hybrid sterility. II. Localization of sterility factors in *Drosophila pseudoobscura* hybrids. Genetics *21*, 113-135.

8.   Dobzhansky, T. (1937). Genetics And The Origin Of Species, (New York: Columbia University Press).

9.   Muller, H.J. (1940). Bearing of the *Drosophila* work on systematics. The new systematics, 185-268.

10.   Muller, H. (1942). Isolating mechanisms, evolution, and temperature. Biol. Symp. *6*, 71-125.

11.   Kimura, M. (1965). Attainment of quasi linkage equilibrium when gene frequencies are changing by natural selection. Genetics *52*, 875-890.

12.   Neher, R.A., and Shraiman, B.I. (2009). Competition between recombination and epistasis can cause a transition from allele to genotype selection. Proc. Natl. Acad. Sci. U.S.A. *106*, 6866-6871.

13.   Wilson, D.J., Gabriel, E., Leatherbarrow, A.J.H., Cheesbrough, J., Gee, S., Bolton, E., Fox, A., Hart, C.A., Diggle, P.J., and Fearnhead, P. (2009). Rapid evolution and the importance of recombination to the gastroenteric pathogen *Campylobacter jejuni*. Mol. Biol. Evol. *26*, 385-397.

14.   Gibson, B., Wilson, D.J., Feil, E., and Eyre-Walker, A. (2018). The distribution of bacterial doubling times in the wild. Proc. R. Soc. B. *285*.

15.   Vos, M., and Didelot, X. (2009). A comparison of homologous recombination rates in bacteria and archaea. Isme J *3*, 199-208.

16.   Calland, J.K., Pascoe, B., Bayliss, S.C., Mourkas, E., Berthenet, E., Thorpe, H.A., Hitchings, M.D., Feil, E.J., Corander, J., Blaser, M.J., et al. (2021). Quantifying bacterial evolution in the wild: A birthday problem for Campylobacter lineages. PLoS Genet. *17*, e1009829.

17.   Arnold, B.J., Gutmann, M.U., Grad, Y.H., Sheppard, S.K., Corander, J., Lipsitch, M., and Hanage, W.P. (2018). Weak epistasis may drive adaptation in recombining bacteria. Genetics *208*, 1247-1260.

18.   Sheppard, S.K., Guttman, D.S., and Fitzgerald, J.R. (2018). Population genomics of bacterial host adaptation. Nat. Rev. Genet. *19*, 549-565.

19.   Andreani, N.A., Hesse, E., and Vos, M. (2017). Prokaryote genome fluidity is dependent on effective population size. Isme J *11*, 1719-1721.

20.   Vos, M., and Eyre-Walker, A. (2017). Are pangenomes adaptive or not? Nature Microbiol. *2*, 1576-1576.

21.   Shapiro, B.J. (2017). The population genetics of pangenomes. Nature Microbiol. *2*, 1574-1574.

22. McInerney, J.O., McNally, A., and O'Connell, M.J. (2017). Why prokaryotes have pangenomes. Nature
Microbiol. *2*, 17040.
23. Orr, H.A. (1996). Dobzhansky, Bateson, and the genetics of speciation. Genetics *144*, 1331-1335.
24. Fraser, C., Hanage, W.P., and Spratt, B.G. (2007). Recombination and the nature of bacterial speciation.
Science *315*, 476-480.
25. Sheppard, S.K., Dallas, J.F., Strachan, N.J., MacRae, M., McCarthy, N.D., Wilson, D.J., Gormley, F.J.,
Falush, D., Ogden, I.D., Maiden, M.C., et al. (2009). *Campylobacter* genotyping to determine the source of
human infection. Clin. Infect. Dis. *48*, 1072-1078.
26. Sheppard, S.K., McCarthy, N.D., Falush, D., and Maiden, M.C. (2008). Convergence of *Campylobacter*
species: implications for bacterial evolution. Science *320*, 237-239.
27. Sheppard, S.K., Didelot, X., Jolley, K.A., Darling, A.E., Pascoe, B., Meric, G., Kelly, D.J., Cody, A.,
Colles, F.M., Strachan, N.J., et al. (2013). Progressive genome-wide introgression in agricultural
Campylobacter coli. Mol. Ecol. *22*, 1051-1064.
28. Cohan, F.M. (2002). Sexual Isolation And Speciation In Bacteria. In Genetics of Mate Choice: From
Sexual Selection to Sexual Isolation. (Springer), pp. 359-370.
29. Ansari, M.A., and Didelot, X. (2014). Inference of the properties of the recombination process from whole
bacterial genomes. Genetics *196*, 253-265.
30. Retchless, A.C., and Lawrence, J.G. (2010). Phylogenetic incongruence arising from fragmented speciation
in enteric bacteria. Proc. Natl. Acad. Sci. U.S.A. *107*, 11453-11458.
31. Dearlove, B.L., Cody, A.J., Pascoe, B., Méric, G., Wilson, D.J., and Sheppard, S.K. (2016). Rapid host
switching in generalist *Campylobacter* strains erodes the signal for tracing human infections. Isme J *10*,
721-729.
32. Weerakoon, D.R., Borden, N.J., Goodson, C.M., Grimes, J., and Olson, J.W. (2009). The role of respiratory
donor enzymes in *Campylobacter jejuni* host colonization and physiology. Microb. Pathog. *47*, 8-15.
33. Taylor, A.J., and Kelly, D.J. (2019). The function, biogenesis and regulation of the electron transport
chains in *Campylobacter jejuni*: New insights into the bioenergetics of a major food-borne pathogen. Adv.
Microb. Physiol. *74*, 239-329.
34. Smart, J.P., Cliff, M.J., and Kelly, D.J. (2009). A role for tungsten in the biology of *Campylobacter jejuni*:
tungstate stimulates formate dehydrogenase activity and is transported via an ultra-high affinity ABC
system distinct from the molybdate transporter. Mol. Microbiol. *74*, 742-757.
35. Doerrler, W.T., Sikdar, R., Kumar, S., and Boughner, L.A. (2013). New functions for the ancient DedA
membrane protein family. J. Bacteriol. *195*, 3-11.
36. Ledgham, F., Quest, B., Vallaeys, T., Mergeay, M., and Covès, J. (2005). A probable link between the
DedA protein and resistance to selenite. Res. Microbiol. *156*, 367-374.
37. Shaw, F.L., Mulholland, F., Le Gall, G., Porcelli, I., Hart, D.J., Pearson, B.M., and van Vliet, A.H.M.
(2012). Selenium-dependent biogenesis of formate dehydrogenase in *Campylobacter jejuni* is controlled by
the *fdhTU* accessory genes. J. Bacteriol. *194*, 3814-3823.
38. Sheppard, S.K., McCarthy, N.D., Jolley, K.A., and Maiden, M.C.J. (2011). Introgression in the genus
Campylobacter: generation and spread of mosaic alleles. Microbiology *157*, 1066-1074.
39. Croucher, N.J., Harris, S.R., Barquist, L., Parkhill, J., and Bentley, S.D. (2012). A high-resolution view of
genome-wide pneumococcal transformation. PLoS Path. *8*, e1002745.
40. He, M., Sebaihia, M., Lawley, T.D., Stabler, R.A., Dawson, L.F., Martin, M.J., Holt, K.E., Seth-Smith,
H.M., Quail, M.A., Rance, R., et al. (2010). Evolutionary dynamics of *Clostridium difficile* over short and
long time scales. Proc. Natl. Acad. Sci. U.S.A. *107*, 7527-7532.
41. Power, P.M., Bentley, S.D., Parkhill, J., Moxon, E.R., and Hood, D.W. (2012). Investigations into genome
diversity of *Haemophilus influenzae* using whole genome sequencing of clinical isolates and laboratory
transformants. BMC Microbiol. *12*, 273.
42. Yahara, K., Méric, G., Taylor, A.J., de Vries, S.P., Murray, S., Pascoe, B., Mageiros, L., Torralbo, A.,
Vidal, A., Ridley, A., et al. (2017). Genome-wide association of functional traits linked with
Campylobacter jejuni survival from farm to fork. Environ. Microbiol. *19*, 361-380.
43. Arnoux, P., Ruppelt, C., Oudouhou, F., Lavergne, J., Siponen, M.I., Toci, R., Mendel, R.R., Bittner, F.,
Pignol, D., Magalon, A., et al. (2015). Sulphur shuttling across a chaperone during molybdenum cofactor
maturation. Nat. Commun. *6*, 6148.
44. Taveirne, M.E., Sikes, M.L., and Olson, J.W. (2009). Molybdenum and tungsten in *Campylobacter jejuni*:
their physiological role and identification of separate transporters regulated by a single ModE-like protein.
Mol. Microbiol. *74*, 758-771.

45. Aguilar-Barajas, E., Díaz-Pérez, C., Ramírez-Díaz, M.I., Riveros-Rosas, H., and Cervantes, C. (2011).
Bacterial transport of sulfate, molybdate, and related oxyanions. BioMetals *24*, 687-707.

46. Thomas, M.T., Shepherd, M., Poole, R.K., van Vliet, A.H.M., Kelly, D.J., and Pearson, B.M. (2011). Two
respiratory enzyme systems in *Campylobacter jejuni* NCTC 11168 contribute to growth on l-lactate.
Environ. Microbiol. *13*, 48-61.

47. Ünal, C.M., and Steinert, M. (2014). Microbial peptidyl-prolyl cis/trans isomerases (PPIases): virulence
factors and potential alternative drug targets. Microbiology and molecular biology reviews *78*, 544-571.

48. Knudson, A.G., Jr. (1971). Mutation and cancer: statistical study of retinoblastoma. Proc Natl Acad Sci U S
A *68*, 820-823.

49. Oren, Y., Smith, M.B., Johns, N.I., Kaplan Zeevi, M., Biran, D., Ron, E.Z., Corander, J., Wang, H.H., Alm,
E.J., and Pupko, T. (2014). Transfer of noncoding DNA drives regulatory rewiring in bacteria. Proc. Natl.
Acad. Sci. U.S.A. *111*, 16112-16117.

50. Arnold, B., Sohail, M., Wadsworth, C., Corander, J., Hanage, W.P., Sunyaev, S., and Grad, Y.H. (2019).
Fine-scale haplotype structure reveals strong signatures of positive selection in a recombining bacterial
pathogen. Mol. Biol. Evol. *37*, 417-428.

51. Wadsworth, C.B., Arnold, B.J., Sater, M.R.A., and Grad, Y.H. (2018). Azithromycin resistance through
interspecific acquisition of an epistasis-dependent efflux pump component and transcriptional regulator in
Neisseria gonorrhoeae. mBio *9*, e01419-01418.

52. Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,
Nikolenko, S.I., Pham, S., Prjibelski, A.D., et al. (2012). SPAdes: a new genome assembly algorithm and
its applications to single-cell sequencing. Journal of computational biology : a journal of computational
molecular cell biology *19*, 455-477.

53. Sheppard, S.K., Didelot, X., Meric, G., Torralbo, A., Jolley, K.A., Kelly, D.J., Bentley, S.D., Maiden,
M.C.J., Parkhill, J., and Falush, D. (2013). Genome-wide association study identifies vitamin B5
biosynthesis as a host specificity factor in *Campylobacter*. Proc. Natl. Acad. Sci. U.S.A. *110*, 11923-11927.

54. Sheppard, S.K., Cheng, L., Méric, G., de Haan, C.P., Llarena, A.K., Marttinen, P., Vidal, A., Ridley, A.,
Clifton-Hadley, F., Connor, T.R., et al. (2014). Cryptic ecology among host generalist *Campylobacter
jejuni* in domestic animals. Mol. Ecol. *23*, 2442-2451.

55. Pascoe, B., Méric, G., Yahara, K., Wimalarathna, H., Murray, S., Hitchings, M.D., Sproston, E.L., Carrillo,
C.D., Taboada, E.N., Cooper, K.K., et al. (2017). Local genes for local bacteria: evidence of allopatry in
the genomes of transatlantic *Campylobacter* populations. Mol. Ecol. *26*, 4497-4508.

56. Pascoe, B., Schiaffino, F., Murray, S., Méric, G., Bayliss, S.C., Hitchings, M.D., Mourkas, E., Calland,
J.K., Burga, R., Yori, P.P., et al. (2020). Genomic epidemiology of *Campylobacter jejuni* associated with
asymptomatic pediatric infection in the Peruvian Amazon. PLoS Negl. Trop. Dis. *14*, e0008533.

57. Mourkas, E., Taylor, A.J., Méric, G., Bayliss, S.C., Pascoe, B., Mageiros, L., Calland, J.K., Hitchings,
M.D., Ridley, A., Vidal, A., et al. (2020). Agricultural intensification and the evolution of host specialism
in the enteric pathogen *Campylobacter jejuni*. Proc. Natl. Acad. Sci. U.S.A. *117*, 11018-11028.

58. Jolley, K.A., and Maiden, M.C.J. (2010). BIGSdb: scalable analysis of bacterial genome variation at the
population level. BMC Bioinformatics *11*, 595.

59. Dingle, K.E., Colles, F.M., Wareing, D.R., Ure, R., Fox, A.J., Bolton, F.E., Bootsma, H.J., Willems, R.J.,
Urwin, R., and Maiden, M.C. (2001). Multilocus sequence typing system for *Campylobacter jejuni*. J. Clin.
Microbiol. *39*, 14-23.

60. Méric, G., Yahara, K., Mageiros, L., Pascoe, B., Maiden, M.C.J., Jolley, K.A., and Sheppard, S.K. (2014).
A reference pan-genome approach to comparative bacterial genomics: identification of novel
epidemiological markers in pathogenic *Campylobacter*. PLOS One *9*, e92798.

61. Aziz, R.K., Bartels, D., Best, A.A., DeJongh, M., Disz, T., Edwards, R.A., Formsma, K., Gerdes, S., Glass,
E.M., Kubal, M., et al. (2008). The RAST server: rapid annotations using subsystems technology. BMC
Genomics *9*, 75.

62. Sheppard, S.K., Jolley, K.A., and Maiden, M.C.J. (2012). A gene-by-gene approach to bacterial population
genomics: whole genome MLST of *Campylobacter*. Genes *3*, 261-277.

63. Price, M.N., Dehal, P.S., and Arkin, A.P. (2010). FastTree 2 – approximately maximum-likelihood trees for
large alignments. PLOS One *5*, e9490.

64. Didelot, X., and Wilson, D.J. (2015). ClonalFrameML: efficient inference of recombination in whole
bacterial genomes. PLoS Comp. Biol. *11*, e1004041.

65. Browning, B.L., and Browning, S.R. (2009). A unified approach to genotype imputation and haplotype-
phase inference for large data sets of trios and unrelated individuals. Am. J. Hum. Genet. *84*, 210-223.

66. Yahara, K., Furuta, Y., Oshima, K., Yoshida, M., Azuma, T., Hattori, M., Uchiyama, I., and Kobayashi, I.
(2013). Chromosome painting *in silico* in a bacterial species reveals fine population structure. Mol. Biol.
Evol. *30*, 1454-1464.
67. Lawson, D.J., Hellenthal, G., Myers, S., and Falush, D. (2012). Inference of population structure using
dense haplotype data. PLoS Genet. *8*, e1002453.
68. Yahara, K., Didelot, X., Ansari, M.A., Sheppard, S.K., and Falush, D. (2014). Efficient inference of
recombination hot regions in bacterial genomes. Mol. Biol. Evol. *31*, 1593-1605.
69. Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of
PhyML 3.0. Syst. Biol. *59*, 307-321.
70. Mell, J.C., Shumilina, S., Hall, I.M., and Redfield, R.J. (2011). Transformation of natural genetic variation
into *Haemophilus influenzae* genomes. PLoS Path. *7*, e1002151-e1002151.
71. Collins, C., and Didelot, X. (2018). A phylogenetic method to perform genome-wide association studies in
microbes that accounts for population structure and recombination. PLoS Comp. Biol. *14*, e1005958.
72. van Vliet, A.H., Wooldridge, K.G., and Ketley, J.M. (1998). Iron-responsive gene regulation in a
Campylobacter jejuni fur mutant. J. Bacteriol. *180*, 5291-5298.
73. Cameron, A., and Gaynor, E.C. (2014). Hygromycin B and apramycin antibiotic resistance cassettes for use
in *Campylobacter jejuni*. PLOS One *9*, e95084.

Acknowledgments

Funding: Wellcome Trust grant 088786/C/09/Z (SS), Medical Research Council (MRC) grant

MR/M501608/1 (SS), Medical Research Council (MRC) grant MR/L015080/1 (SS), Food

Standards Agency project FS101087 (SS), Biotechnology & Biological Sciences Research

Council (BBSRC) grant BB/S014497/1 (DK).

Author contributions:

Conceptualization and study design: SS, XD, DF, KY, DK, AT

Sample collection: SS, AvV, NW

Laboratory work: GM, AT, LM, MH, BP

Data archiving: BP, KJ

Data analysis: AT, XD, KY, LM, JKC, EM, SP, SB, SS, JC

Data interpretation: KY, GM, XD, SS, MM, JP, DK, AT, DF

Writing: SS, GM, BP, CK, AT, DK

Competing Interest Statement: The authors declare that they have no competing interests.

Data and materials availability: Short-read sequence data for all isolates sequenced in this

study are deposited in the sequence read archive (SRA) and can be found associated with NCBI

BioProjects   PRJNA689604   (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA689604)   and

PRJEB11972 (https://www.ncbi.nlm.nih.gov/bioproject/PRJEB11972). These were augmented

with 498 previously published genomes and assembled genomes are available on Figureshare

(doi.org/10.6084/m9.Figureshare.13521683). Accession numbers for all genomes are included in

Data S4. Source data are provided for this paper (Data S2).

Figureures and Tables

Figure 1. Genome-wide introgression from *C. jejuni* to *C. coli*. (A) Phylogenetic tree

reconstructed using neighbour-joining on a whole-genome alignment of 973 *C. jejuni* and *C. coli*

isolates. Introgressed *C. coli* clades are represented with red (CC-828, n=677) and purple (CC-

1150, n=12) circles, unintrogressed clade 1 (n=35) is shown in pink, clade 2 (n=45) in yellow

and clade 3 (n=58) in green. A set of 146 *C. jejuni* genomes belonging to 30 clonal complexes (4

to 5 isolates per ST) are shown in blue. Recipient and donor populations, used to infer

introgression in chromosome painting analysis, are indicated. The scale bar represents the

number of substitutions per site. (B) Summary of introgressed *C. jejuni* SNPs in *C. coli* CC-828

(n=677, red) and CC-1150 (n=12, purple) genomes using ChromoPainter; the number of

introgressed core SNPs (coloured histograms; left y-axis) and core genes (white histograms; right

y-axis) for a range of recipient strains proportions (at least 1, more than 50% and more than

98%) is shown. (C) Diagram of *Campylobacter* species and clade (C1*, C2, C3) divergence with

arrows indicating introgression from *C. jejuni* into *C. coli* (i) clade 1, (ii, iii) CC-828 and CC-

1150, (iv, v) subsequent clonal expansion and ongoing introgression.

Figure 2. Covariation in introgressed *C. coli* genomes. (A) CC-828 and CC-1150 *C. coli*

genomes were analysed using a continuous time Markov chain (CTMC) model and covariation

was assessed for pairs of biallelic sites separated by at least 20kb along the genome while

accounting for the effect of population structure and recombination. There are many biallelic

sites that do not change often on the tree and few that do. Putative epistatic sites change more

frequently than average with biallelic pairs found together on multiple branches. (B) Miami plot

of each polymorphic site showing the maximum *p*-value for covarying biallelic pairs (>20kb

apart) and the frequency of introgression in CC-828 and CC-1150. (C) The frequency of major

and minor haplotype combinations (inset) among the 2578 pairs of covarying SNPs in the 689 *C.*

coli clade-1 recipient genomes, revealing that the majority of long range covariation was

801   between introgressed *C. jejuni* sites. (D) The position of putative epistatic sites mapped on the *C.*

802   *coli* CVM29710 reference for covarying *C. jejuni-C. jejuni* SNPs (red) in 16 gene pairs (a to l),

803   and other haplotype combinations (grey).

Figure 3. Genomic context and physiological roles of introgressed epistatically linked genes involved in FDH biogenesis and activity. (A) Genome organisation and percentage of co-varying SNP pairs (internal legend). (B-D) FDH activity of whole cells determined by oxygen consumption rates in a Clark-type electrode (nmol oxygen consumed per minute per mg of total protein) for (B) cells grown in rich media (excess selenium), (C) cells grown in minimal media with 0.5 nM sodium selenite, and (D) cells grown in minimal media with either 5 nM sodium selenite (left, open bars) or 5 µM sodium selenate (right, hashed bars). All data are means of at

824   least 4 independent determinations and error bars are SD. *** denotes p value of <0.001 by

825   students t-test. Assays suggest that both FdhT and SelF facilitate selenium acquisition, possibly

826   representing low and high affinity transporters, respectively.

Figure 4. A two-hit model for core genome recombination in natural *C. coli* populations.

A&B: (i) *C. jejuni* (blue) and unintrogressed *C. coli* (red) co-exist with genomes (internal circles) harbouring haplotype pairs (x-x) that segregate by species; (ii) Horizontal gene transfer, HGT, occurs (R1) disrupting covarying genetic elements and reducing the relative fitness of introgressed *C. coli* to varying degrees (grey arrow), few strains retain mixed *C. coli - C. jejuni* haplotypes; (iii) HGT continues (R2) and, where recombined mixed haplotypes survived, ancestral *C. jejuni* haplotype pairs are reinstated in introgressed *C. coli;* (iv,v) introgressed *C. coli* outcompete unintrogressed strains.