Title: Rationally designed immunogens enable immune focusing to the SARS-CoV-2 receptor binding motif

Authors: Blake M. Hauser[1], Maya Sangesland[1], Kerri St. Denis[1], Jared Feldman[1], Evan C. Lam[1], Ty Kannegieter[1], Alejandro B. Balazs[1], Daniel Lingwood[1], Aaron G. Schmidt[1,2*]

[1]Ragon Institute of MGH, MIT and Harvard, Cambridge, MA, 02139, USA

[2]Department of Microbiology, Harvard Medical School, Boston, MA 02115, USA

Key words: immunogen design, glycan, immune focusing, SARS-CoV-2, coronavirus

Correspondence:
Aaron G. Schmidt
Tel: 857-268-7118; E-mail: aschmidt@crystal.harvard.edu

ABSTRACT

Eliciting antibodies to surface-exposed viral glycoproteins can lead to protective responses that

ultimately control and prevent future infections. Targeting functionally conserved epitopes may

help reduce the likelihood of viral escape and aid in preventing the spread of related viruses with

pandemic potential. One such functionally conserved viral epitope is the site to which a receptor

must bind to facilitate viral entry. Here, we leveraged rational immunogen design strategies to

focus humoral responses to the receptor binding motif (RBM) on the SARS-CoV-2 spike. Using

glycan engineering and epitope scaffolding, we find an improved targeting of the serum response

to the RBM in context of SARS-CoV-2 spike imprinting. Furthermore, we observed a robust

SARS-CoV-2-neutralizing serum response with increased potency against related sarbecoviruses,

SARS-CoV and WIV1-CoV. Thus, RBM focusing is a promising strategy to elicit breadth across

emerging sarbecoviruses and represents an adaptable design approach for targeting conserved

epitopes on other viral glycoproteins.

One Sentence Summary: SARS-CoV-2 immune focusing with engineered immunogens

MAIN TEXT

Humoral responses elicited by vaccination or infection predominantly target surface-exposed viral glycoproteins. These responses can often provide protection against future infections to the same or closely related viral variants. However, in some instances, such as influenza and HIV, the elicited responses are often poorly protective as they target variable epitopes (*1, 2*). Furthermore, waning of responses (i.e., durability), as is the case for common cold-causing coronaviruses, results in susceptibility to reinfections (*3-6*). For SARS-CoV-2 (SARS-2) it remains unclear whether current vaccines will confer long-term protection. Furthermore, it is increasingly apparent that humoral immunity elicited by vaccination or natural infection may provide reduced protection against emerging SARS- 2 variants (*7-9*). Thus, implementing rational design strategies aimed at directing the immune response to conserved viral epitopes may help reduce the likelihood of viral escape and lead to more broadly protective responses (*10, 11*).

Two immunogen design strategies used to direct humoral responses include "masking" epitopes via engineering putative N-linked glycosylation sites (PNGs) and the design of protein scaffolds to present broadly protective epitopes (*12, 13*); these strategies have been used previously for viral glycoproteins RSV F, influenza hemagglutinin and HIV envelope (*14-16*). Applying these approaches to the SARS-2 spike provides an opportunity to potentially improve serum neutralization potency, efficacy against variants, and cross-reactivity of antibody responses. A potential target of these efforts is the angiotensin converting enzyme 2 (ACE2) receptor binding motif (RBM) of the receptor binding domain (RBD) (*17, 18*). Indeed, several potently neutralizing RBM-directed antibodies that interfere with ACE2 binding are protective and some can also neutralize related sarbecoviruses (*10, 11, 18-20*). Here, we show that hyperglycosylation of the RBD and a "resurfacing" approach that grafts the RBM from SARS-2 onto heterologous coronavirus RBDs focuses serum responses to the RBM. This immune-focused response is potently neutralizing with breadth across SARS-2 variants and other coronaviruses.

The RBM of SARS-2 and related sarbecoviruses, SARS-CoV (SARS-1) and WIV1-CoV (WIV1), is a contiguous sequence spanning residues 437-507 (SARS-2 numbering) of the spike protein. In an effort to elicit RBM-specific responses only, we first asked whether the RBM itself could be recombinantly expressed in absence of the rest of the RBD (**Fig. 1A**). While the SARS-2 RBM

could indeed be overexpressed, it failed to both engage the conformation-specific RBM-directed antibody B38 and bind to cell-surface expressed ACE2 (**Fig. S1**). These results likely suggest that the RBM is conformationally flexible, and that the RBD serves as a structural "scaffold" to stabilize the RBM in its binding-compatible conformation. To circumvent the considerable hurdle of *de novo* scaffold design for RBM presentation, we asked whether heterologous sarbecovirus RBDs from SARS-1 and WIV1 and the more distantly related merbecovirus MERS-CoV (MERS) could serve as scaffolds (**Fig. 1A**) —variations of this approach were used previously to modulate ACE2 binding properties (*21, 22*). In context of immunizations, we hypothesized that these heterologous RBDs would present the SARS-2 RBM while removing any other SARS-2-specific epitopes. The SARS-1, WIV-1 and MERS RBDs share a pairwise amino acid identity with SARS-2 of 73.0%, 75.4% and 19.5%, respectively. The RBM is less conserved despite have a shared ACE2 receptor for SARS-1 and WIV1 with only 49.3% and 52.1% identity, respectively; as MERS uses DPP4 as a receptor, its RBM shares no notable identity (*23*). While we were unable to "resurface" MERS RBD with the SARS-2 RBM, the related SARS-1 and WIV-1 RBDs successfully accepted the RBM transfer. These resurfaced constructs, rsSARS-1 and rsWIV-1 retained binding to the SARS-2 RBM-specific B38 antibody as well as effectively engaged ACE2 (**Fig. S2**) (*19*). These data suggest that there are sequence and structural constraints within the RBD required for successful RBM grafting; such an approach may be facilitated by using CoV RBDs that use the same receptor for viral entry.

We next used these resurfaced RBDs as templates for further modification using glycan engineering. This approach aimed to mask conserved, cross-reactive epitopes shared between the SARS-1, SARS-2, and WIV1 RBDs. There are two evolutionarily conserved PNGs at positions 331 and 343; SARS-1 and WIV1 have an additional conserved PNG at position 370 (SARS-2 numbering). To further increase overall surface glycan density, we introduced novel PNGs onto wildtype SARS-2 as well as rsSARS-1 and rsWIV1 RBDs. Based on structural modeling, we identified 5 potential sites on rsSARS-1 and rsWIV1 as well as 6 on SARS-2. Including the native PNGs, all constructs had a total of 8 glycans (**Fig. 1B-D, S3**)— we denote these hyperglycosylated (hg) constructs as SARS-2^{hg}, rsSARS-1^{hg}, and rsWIV1^{hg}. We expressed these constructs in mammalian cells to ensure complex glycosylation in order to maximize any glycan "shielding" effect. We subsequently characterized these constructs using the RBM-directed antibody B38, as

well as ACE2 binding, to ensure that the engineered PNGs did not adversely affect the RBM

conformation. Overall, the hyperglycosylated constructs were largely comparable in affinity for

B38, with only ~2-fold decrease, and still effectively engaged ACE2 (**Fig. S4**). These results

confirm a conformational and functionally intact RBM.

Next, we assessed whether the engineered PNGs abrogated binding to sarbecovirus cross-reactive

antibodies S309 and CR3022—both antibodies were isolated from SARS-1 convalescent

individuals (*24, 25*). The CR3022 contact residues on SARS-1 and WIV1 differ only at a single

residue while SARS-2 differs at 5 residues across both CR3022 and S309 epitopes (*26*).

Importantly, these epitopic regions were shown to be a significant portion of the SARS-2 RBD-

directed response in murine immunizations and thus any RBM focusing would require masking of

these regions (**Fig. S4**) (*24, 25, 27*). While SARS-2^{hg} effectively abrogated S309 and CR3022

binding, the engineered PNGs at the antibody:antigen interface on rsSARS-1^{hg} and rsWIV1^{hg} did

not completely abrogate S309 and CR3022 binding. We therefore incorporated unique mutations

on rsSARS-1^{hg} and rsWIV1^{hg} so that any elicited antibodies would be less likely to cross-react

between these two constructs. To that end, we found K378A and the engineered glycan at residue

383 (SARS-2 numbering) completely abrogated CR3022 binding in both rsSARS-1^{hg} and

rsWIV1^{hg} (**Fig. S4**). For S309, mutations P337D in rsSARS-1^{hg} and G339W in rsWIV1^{hg} in

addition to glycans at residues 441 and 354 (SARS-2 numbering) were sufficient to disrupt binding

(**Fig. S4**). We made two additional mutations, G381R, M430K on rsSARS-1^{hg} and K386A, T430R

on rsWIV1^{hg}, to further increase the antigenic distance between these scaffolds (**Fig. 1C, D**).

We then tested the immunogenicity and antigenicity of our optimized constructs and assessed their

RBM immune-focusing properties, in the murine model. In order to increase avidity and to

minimize any off-target tag-specific responses, we generated trimeric versions of each immunogen

using our previously characterized hyperglycosylated, cysteine-stabilized GCN4 tag (hgGCN4^{cys})

(*27, 28*). We first primed all cohorts with SARS-2 spike to reflect pre-existing SARS-2 immunity

and to imprint an initial RBM response that may be recalled and selectively expanded by our

immunogens. To test potential RBM immune-focusing, one cohort was sequentially immunized

with SARS-2^{hg} trimers ("Trimer^{hg} cohort") and a second cohort was immunized with SARS-2^{hg}

trimers followed by a cocktail of rsSARS-1^{hg} and rsWIV1^{hg} ("Cocktail^{hg} cohort") (**Fig. 2A**). In

order to facilitate comparisons and establish the efficacy of RBM-focusing, we included a "ΔRBM

cohort". This cohort was immunized with a modified SARS-2 RBD (ΔRBM) with four novel

glycans engineered at positions 448, 475, 496, and 501 within the RBM. These PNGs effectively

abrogate RBM-directed B38 antibody binding and engagement of ACE2 (27) and should restrict

elicited humoral response to this epitope. Finally, as a control cohort, we included a SARS-2 spike

prime followed with sequential immunizations with wildtype (i.e., unmodified) SARS-2 RBD

trimer ("Trimer cohort").

Overall, we find that all cohorts elicit robust serum responses to wildtype SARS-2 RBD (**Fig. 2B-**

C, S5). In order to specifically evaluate the RBM-directed responses, we compared serum ELISA

titers to wildtype SARS-2 RBD and the SARS-2 ΔRBM RBD construct. We find that the Trimer^{hg}

and Cocktail^{hg} cohorts had a significant increase in serum titers to wildtype SARS-2 RBD relative

to SARS-2 ΔRBM RBD; this was in contrast to the ΔRBM and Trimer cohorts (**Fig. 2B,C, S5A,B**).

Across the Trimer^{hg} and Cocktail^{hg} cohorts, the median binding loss to the SARS-2 ΔRBM RBD

relative to wildtype SARS-2 RBD was 82%, indicating that ~82% of serum antibodies are RBM-

directed by this metric. The Cocktail^{hg} cohort had a slight increase in RBM focusing relative to the

Trimer^{hg} cohort which received two SARS-2^{hg} boosts. This observed increase may be due to

increasing the overall antigenic distance (i.e., sequence difference) between the WIV1 and SARS-

1 RBDs relative to SARS-2 while maintaining the identical SARS-2 RBM epitope. Additionally,

we find that the Trimer^{hg} and Cocktail^{hg} cohorts had significantly lower titers to SARS-1 and WIV1

RBDs as compared to SARS-2 RBD (**Fig. 2B**). This difference was most pronounced in the

Cocktail^{hg} cohort, suggesting that the hyperglycosylation and engineered mutations within the

RBD effectively dampened responses to these conserved, cross-reactive epitopes that are present

outside the RBM and shared between SARS-1, WIV1, and SARS-2. Furthermore, serum titers

against the rsSARS-1 and rsWIV1 RBDs were comparable to SARS-2 RBD, indicating that there

is minimal antibody response directed towards wildtype SARS-1 and WIV1 RBD epitopes in

comparison to the SARS-2 RBM (**Fig. 2D, S5C**). We observed no significant glycan-dependent

serum response in either cohort that used hyperglycosylation (**Fig. S6**). Collectively, these data

confirm an enhanced SARS-2 RBM-focused serum response elicited by our engineered

immunogens.

We next compared the neutralization potency of all cohorts using SARS-1, SARS-2, and WIV1

pseudoviruses (*29, 30*). While all cohorts elicited a potent SARS-2 neutralizing response, notably,

the Trimer[hg] and Cocktail[hg] cohorts also exhibited potent SARS-1 and WIV1 pseudovirus

neutralization relative to the control cohorts (**Fig. 3A, S7, S5B**). This is particularly noteworthy

for the Trimer[hg] cohort as it did not include SARS-1 or WIV1 RBDs in the immunization regimen.

WIV1 in this instance is broadly representative of possible future emerging sarbecoviruses with

pandemic potential (*31*). To further epitope map the RBM-focused responses, we performed

ELISA-based antibody competition using cross-reactive antibodies CR3022, S309, ADI-55689,

and ADI-56046 and WIV1 RBD (**Fig. 3B-C**). The latter two antibodies bind a conserved

sarbecovirus RBM epitope also targeted by the antibody ADG-2, which is currently in clinical

development (*10, 11*). Competition ELISAs suggest that the cross reactive WIV1-directed

responses in the Trimer[hg] and Cocktail[hg] cohorts focus to the ADG-2-like epitope, as well as to the

CR3022 and S309 epitopes in the Cocktail[hg] cohort (**Fig. 3B-C**). Thus, SARS-2[hg], rsSARS-1[hg],

and rsWIV1[hg] RBDs can induce not only potent SARS-2 neutralizing antibodies, but also cross-

reactive antibodies that bind to a conserved RBM epitope (**Fig. S8**). Notably, these results are in

contrast to our previous work showing that a cocktail of sarbecovirus that included SARS-1 and

WIV1 RBDs could predominantly focus the antibody response towards the conserved CR3022

and S309 epitopic regions (*27*).

Many SARS-2 variants of concern include mutations within the RBM including B.1.1.7, B.1.351

and P.1 first detected in the United Kingdom, South Africa, and Brazil, respectively (**Fig. 4A**). We

therefore asked what the consequence was of enhanced focusing to the RBM and whether the

elicited responses elicited were sensitive to these mutations. Interestingly, serum from the Trimer[hg]

and Cocktail[hg] cohorts showed no significant loss of binding to the B.1.351 RBD compared to the

wildtype SARS-2 RBD (**Fig. 4A, B**). This is in contrast to the control Trimer cohort, which showed

a significant loss of binding and parallels the observation of reduced serum binding from human

subjects immunized with current SARS-2 vaccines (*9, 32, 33*). Second, we tested all sera for

neutralization against SARS-2 variant pseudoviruses: B.1.1.7, B.1.351 and P.1. While the control

Trimer cohort could still neutralize P.1 and B.1.1.7 pseudoviruses, there was a significant loss of

neutralization to the B.1.351 variant, consistent with our ELISA data. In contrast, we find no

significant loss of neutralization against these variants in the Trimer[hg], Cocktail[hg], and ΔRBM

cohorts (**Fig. 4C**). For ΔRBM cohort, the elicited responses were likely focused to neutralizing

epitopes within the RBD (e.g., CR3022, S309) and therefore were not sensitive to these RBM

mutations. However, the neutralizing response observed in Trimer^{hg} and Cocktail^{hg} cohorts

potentially indicate that immune-focusing to the RBM may allow for greater recognition (i.e.,

accommodation) of mutations compared to the RBM-directed antibody response elicited via

infection or vaccination (*32, 34*).

Collectively, our results demonstrate immunogen design approaches that can be leveraged to

enhance an RBM-focused humoral response. It is a strategy that maintains protective SARS-2

neutralization while also eliciting humoral responses that recognize emerging variants and

coronaviruses with pandemic potential. Importantly, these design strategies are not limited to

coronaviruses and are adaptable to other viruses as a general approach to elicit protective responses

to conserved epitopes.

Acknowledgments: We thank members of the Schmidt Laboratory for helpful discussions. We
thank Timothy Caradonna and Catherine Jacob-Dolan for critical reading of the manuscript. We
thank Dr. Jason McLellan from University of Texas, Austin for the spike plasmid. We thank Nir
Hacohen and Michael Farzan for the kind gift of the ACE2 expressing 293T cells.
214
Funding: We acknowledge funding from NIH R01s AI146779 (AGS), AI124378, AI137057 and
AI153098 (DL), and a Massachusetts Consortium on Pathogenesis Readiness (MassCPR) grant
(AGS); training grants: NIGMS T32 GM007753 (BMH and TMC); T32 AI007245 (JF); F31
Al138368 (MS). A.B.B. is supported by the National Institutes for Drug Abuse (NIDA) Avenir
New Innovator Award DP2DA040254, the MGH Transformative Scholars Program as well as
funding from the Charles H. Hood Foundation (ABB). This independent research was supported
by the Gilead Sciences Research Scholars Program in HIV (ABB).
222
223
224   **Author contributions:** Conceptualization, BMH, AGS; Methodology, BMH, ECL, ABB, DL,
225   AGS; Investigation, BMH, MS, KS, ECL, JF, TK; Writing – Original Draft, BMH and AGS;
226   Writing – Review and Editing, all authors; Funding Acquisition, ABB, DL, AGS; Supervision,
227   ABB, DL, AGS.; **Competing interests:** Authors declare no competing interests.; and **Data and**
228   **materials availability:** All data is available in the main text or in the supplementary materials.
229

Fig. 1. Resurfacing and hyperglycosylation approaches for immune-focusing. (A) Design
schematic for resurfacing SARS-1 (rsSARS-1) and WIV1 (rsWIV1
) with the SARS-2 receptor binding motif (RBM). Design schematic for hyperglycosylating
SARS-2 **(B)**, rsSARS-1 **(C)** and rsWIV1 **(D)** receptor binding domains (RBDs). Non-native
engineered glycans and native glycans are modeled; native SARS-2 RBM glycan at position 331
is omitted in the schematic. Mutations in the WIV1 and SARS-1 RBDs are shown in red and
italicized in the linear diagram. All images were created using PDB 6M0J.

Fig. 2. Serum analysis from cohorts. (A) Schematic of immunization cohorts; N= number of mice in each cohort **(B, C)** Serum following immunizations was assayed in ELISA at day 56 with different coronavirus antigens. Statistical significance was determined using Kruskal-Wallis test with post-hoc analysis using Dunn's test corrected for multiple comparisons or Mann-Whitney U test (* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$). **(D)** Day 56 serum samples assayed against rsSARS-1 and rsWIV1 RBDs no longer show statistically significant differences in binding compared to SARS-2 RBD as determined using Kruskal-Wallis test with post-hoc analysis using Dunn's test corrected for multiple comparisons.

Fig. 3. Potency and characterization of SARS-like coronavirus neutralization response. (A)

Day 56 serum from all mice was assayed for neutralization against SARS-2, SARS-1, and WIV1

pseudoviruses. Statistical significance was determined using the Kruskal-Wallis test with post-hoc

analysis using Dunn's test corrected for multiple comparisons (* = $p < 0.05$, ** = $p < 0.01$, ns =

not significant). **(B)** Approximate locations of representative antibody epitopes from each of the

four SARS-2 RBD-directed antibody classes (*18*) and ADG-2-like antibodies on the SARS-2

RBD. (PDB: 6M0J) **(C)** Antibody competition ELISAs with WIV1 RBD as the coating antigen.

The Trimer^{hg} and Cocktail^{hg} were independently analyzed (first two panels) and statistically

combined (last panel) to highlight observed RBM-focusing. Statistical significance was

determined the Friedman test with post-hoc analysis using Dunn's test corrected for multiple

comparisons (* = $p < 0.05$, ** $p < 0.01$).

Fig. 4. Binding and neutralization of SARS-2 variants. (A) Structural depiction of SARS-2

variant RBD mutations for 501Y.V2 (red), as well as ACE2 contact residues (cyan). (PDB: 6M0J)

Sequences depict all spike mutations across select variants. **(B)** Day 56 serum was assayed in

ELISA against SARS-2 RBD (WT) and SARS-2 RBD with K417N, E484K, and N501Y mutations

(B.1.351). Statistical significance was determined using the ratio paired t-test (* = $p < 0.05$, ** =

$p < 0.01$; ns = not significant). **(C)** Day 56 serum was assayed against SARS-2 variant

pseudoviruses for neutralization. Statistical significance was determined using the Kruskal-Wallis

test with post-hoc analysis using Dunn's test corrected for multiple comparisons (* = $p < 0.05$).

Lower limit of detection for each cohort is shown with a dotted line, and y-axis values were

selected accordingly. Pseudovirus neutralization for WT SARS-2 was also performed with the

$\sim 10^2$ limit of detection for the ΔRBM, Trimer^{hg}, and Cocktail^{hg} cohorts (**Fig. S7**), but NT50 values

at the limit of detection were adjusted to match the lower limit of detection ($\sim 10^1$) for the variant

pseudoviruses to facilitate fair comparison.

METHODS

<u>Immunogen and Coating Protein Expression and Purification</u>

The SARS-CoV-2 (Genbank MN975262.1), SARS-CoV (Genbank ABD72970.1), WIV1-CoV (Genbank AGZ48828.1) RBDs were used as the basis for constructing these immunogens. To graft the SARS-2 RBM onto SARS-1 and WIV1 scaffolds to create the rsSARS-1 and rsWIV1 monomers, boundaries of SARS-2 residues 437 – 507 were used. All constructs were codon optimized by Integrated DNA Technologies and purchased as gblocks. Gblocks were then cloned into pVRC and sequence confirmed via Genewiz. Monomeric constructs for serum ELISA coating contained C-terminal HRV 3C-cleavable 8xHis and SBP tags. Trimeric constructs also included C-terminal HRV 3C-cleavable 8xHis tags, in addition to a previously published hyperglycosylated GCN4 tag with two engineered C-terminal cystines (*27, 28*). Dr. Jason McLellan at the University of Texas, Austin provided the spike plasmid, which contained a non-cleavable foldon trimerization domain in addition to C-terminal HRV 3C cleavable 6xHis and 2xStrep II tags. The SARS-2 ΔRBM RBD construct was generated as previously described with four additional engineered putative N-linked glycosylation sites at positions 448, 475, 496, and 501 (*27*).

Expi 293F cells (ThermoFisher) were used to express proteins. Transfections were performed with Expifectamine reagents per the manufacturer's protocol. After 5-7 days, transfections were harvested and centrifuged for clarification. Cobalt-TALON resin (Takara) was used to perform immobilized metal affinity chromatography via the 8xHis tag. Proteins were eluted using imidazole, concentrated, and passed over a Superdex 200 Increase 10/300 GL (GE Healthcare) size exclusion column. Size exclusion chromatography was performed in PBS (Corning). For immunogens, HRV 3C protease (ThermoScientific) cleavage of affinity tags was performed prior to immunization. Cobalt-TALON resin was used for a repurification to remove the His-tagged HRV 3C protease, cleaved tag, and remaining uncleaved protein.

<u>Fab and IgG Expression and Purification</u>

The variable heavy and light chain genes for each antibody were codon optimized by Integrated DNA Technologies, purchased as gblocks, and cloned into pVRC constructs which already contained the appropriate constant domains as previously described (*35, 36*). The Fab heavy chain

vector contained a HRV 3C-cleavable 8xHis tag, and the IgG heavy chain vector contained HRV

3C-cleavable 8xHis and SBP tags. The same transfection and purification protocol as used for the

immunogens and coating proteins was used for the Fabs and IgGs.

<u>Biolayer Interferometry</u>

Biolayer interferometry (BLI) experiments were performed using a BLItz instrument (Fortebio)

with FAB2G biosensors (Fortebio). All proteins were diluted in PBS. Fabs were immobilized to

the biosensors, and coronavirus proteins were used as the analytes. To determine binding affinities,

single-hit measurements were performed starting at 10 µM to calculate an approximate K_D in order

to evaluate which concentrations should be used for subsequent titrations. Measurements at a

minimum of three additional concentrations were performed. Vendor-supplied software was used

to generate a final K_D estimate via a global fit model with a 1:1 binding isotherm.

<u>Immunizations</u>

All immunizations were performed using female C57BL/6 mice (Jackson Laboratory) aged 6-10

weeks. Mice received 20 µg of protein adjuvanted with 50% w/v Sigma adjuvant in 100 µL of

inoculum via the intraperitoneal route. Following an initial prime (day 0), boosts occurred at days

21 and 42. Serum samples were collected for characterization on day 56 from all cohorts, in

addition to day 35 for the HG Trimer and HG Cocktail cohorts. All experiments were conducted

with institutional IACUC approval (MGH protocol 2014N000252).

<u>Serum ELISAs</u>

Serum ELISAs were executed using 96-well, clear, flat-bottom, high bind microplates (Corning).

These plates were coated with 100 µL of protein, which were adjusted to a concentration of 5

µg/mL (in PBS). Plates were incubated overnight at 4°C. After incubation, plates had their coating

solution removed and were blocked using 1% BSA in PBS with 1% Tween. This was done for 60

minutes at room temperature. This blocking solution was removed, and sera was diluted 40-fold

in PBS. A 5-fold serial dilution was then performed. CR3022 IgG, similarly serially diluted (5-

fold) from a 5 µg/mL starting concentration, was used as a positive control. 40 µL of primary

antibody solution was used per well. Following this, samples were incubated for 90 minutes at

room temperature. Plates were washed three times using PBS-Tween. 150 µL of HRP-conjugated

rabbit anti-mouse IgG antibody, sourced commercially from Abcam (at a 1:20,000 dilution in

PBS), was used for the secondary incubation. Secondary incubation was performed for one hour,

similarly at room temperature. Plates were subsequently washed three times using PBS-Tween.

1xABTS development solution (ThermoFisher) was used according to the manufacturer's

protocol. Development was abrogated after 30 minutes using a 1% SDS solution, and plates were

read using a SectraMaxiD3 plate reader (Molecular Devices) for absorbance at 405 nm.

Competition ELISAs

A similar protocol to the serum ELISAs was used for the competition ELISAs. For the primary

incubation, 40 µL of the relevant IgG at 1 µM was used at room temperature for 60 minutes. Mouse

sera were then spiked in such that the final concentration of sera fell within the linear range for the

serum ELISA titration curve for the respective coating antigen, and an additional 60 minutes of

room temperature incubation occurred. After removing the primary solution, plates were washed

three times with PBS-Tween. Secondary incubation consisted of HRP-conjugated goat anti-mouse

IgG, human/bovine/horse SP ads antibody (Southern Biotech) at a concentration of 1:4000. The

remaining ELISA procedure (secondary incubation, washing, developing) occurred as described

for the serum ELISAs.

ACE2 Cell Binding Assay

ACE2 expressing 293T cells (37) (a kind gift from Nir Hacohen and Michael Farzan) were

harvested. A wash was performed using PBS supplemented with 2% FBS. 200,000 cells were

allocated to each labelling condition. Primary incubation occurred using 100 µL of 1 µM antigen

in PBS on ice for 60 minutes. Two washes were performed with PBS supplemented with 2% FBS.

Secondary incubation was performed using 50 µL of 1:200 streptavidin-PE (Invitrogen) on ice for

30 mins. Two washes were performed with PBS supplemented with 2% FBS, and then cells were

resuspended in 100 µL of PBS supplemented with 2% FBS. A Stratedigm S1000Exi Flow

Cytometer was used to perform flow cytometry. FlowJo (version 10) was used to analyze FCS

files.

Pseudovirus Neutralization Assay

Serum neutralization against SARS-CoV-2, SARS-CoV, and WIV1-CoV was assayed using pseudotyped lentiviral particles expressing spike proteins described previously (*29*). Transient transfection of 293T cells was used to generate lentiviral particles. Viral supernatant titers were measured using flow cytometry of 293T-ACE2 cells (*37*) and utilizing the HIV-1 p24^{CA} antigen capture assay (Leidos Biomedical Research, Inc.). 384-well plates (Grenier) were used to perform assays on a Tecan Fluent Automated Workstation. For mouse sera, samples underwent primary dilutions of 1:3 or 1:9 followed by serial 3-fold dilutions. 20 µL each of sera and pseudovirus (125 infectious units) were loaded into each well. Plates were then incubated for 1 hour at room temperature. Following incubation, 10,000 293T-ACE2 cells (*37*) in 20 µL of media containing 15 µg/mL polybrene was introduced to each well. The plates were then further incubated at 37°C for 60-72 hours.

Cells were lysed using assay buffers described previously (*38*). Luciferase expression was quantified using a Spectramax L luminometer (Molecular Devices). Neutralization percentage for each concentration of serum was calculated by deducting background luminescence from cells-only sample wells and subsequently dividing by the luminescence of wells containing both virus and cells. Nonlinear regressions were fitted to the data using GraphPad Prism (version 9), allowing IC50 values to be calculated via the interpolated 50% inhibitory concentration. IC50 values were calculated with a neutralization values greater than or equal to 80% at maximum serum concentration for each sample. NT50 values were then calculated using the reciprocal of IC50 values. Serum neutralization potency values were calculated by dividing the NT50 against a particular pseudovirus by the endpoint titer against the respective RBD. For samples with NT50 values below the limit of detection, the lowest limit of detection across all neutralization assays was used as the NT50 value to calculate neutralization potency. This prevents a higher limit of detection from skewing neutralization potency results. Endpoint titers were normalized relative to a CR3022 IgG control, which was run in every serum ELISA.

In comparing NT50 values for the various cohorts across the wildtype and variant pseudoviruses, the lowest limit of detection across all neutralization assays performed for a given cohort was used for any NT50 values that fell below the limit of detection. This prevents a higher limit of detection in some assays from skewing the comparison results.

Statistical Analysis

Curve fitting and statistical analyses were performed with GraphPad Prism (version 9). Non-parametric statistics were used throughout. To compare multiple populations, the Kruskal-Wallis non-parametric ANOVA was used with post hoc analysis using Dunn's test for multiple comparisons. The Mann-Whitney U test was used to compare two populations without consideration for paired samples. The ratio-paired t-test was used to compare two populations with consideration for paired samples and evidence of normality. P values in ANOVA analyses were corrected for multiple comparisons. A p value < 0.05 was considered significant.

REFERENCES

1. J. Overbaugh, L. Morris, The Antibody Response against HIV-1. *Cold Spring Harb Perspect Med* **2**, a007039 (2012).
2. F. Krammer, The human antibody response to influenza A virus infection and vaccination. *Nat Rev Immunol* **19**, 383-397 (2019).
3. K. A. Callow, H. F. Parry, M. Sergeant, D. A. Tyrrell, The time course of the immune response to experimental coronavirus infection of man. *Epidemiol Infect* **105**, 435-446 (1990).
4. J. O. Hendley, H. B. Fishburne, J. M. Gwaltney, Jr., Coronavirus infections in working adults. Eight-year study with 229 E and OC 43. *Am Rev Respir Dis* **105**, 805-811 (1972).
5. A. S. Monto, S. K. Lim, The Tecumseh study of respiratory illness. VI. Frequency of and relationship between outbreaks of coronavirus infection. *J Infect Dis* **129**, 271-276 (1974).
6. R. Eguia *et al.*, A human coronavirus evolves antigenically to escape antibody immunity. *bioRxiv*, (2020).
7. C. K. Wibmer *et al.*, SARS-CoV-2 501Y.V2 escapes neutralization by South African COVID-19 donor plasma. *bioRxiv*, (2021).
8. P. Wang *et al.*, Increased Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7 to Antibody Neutralization. *bioRxiv*, (2021).
9. Z. Wang *et al.*, mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. *bioRxiv*, (2021).
10. A. Z. Wec *et al.*, Broad neutralization of SARS-related viruses by human monoclonal antibodies. *Science* **369**, 731-736 (2020).
11. C. G. Rappazzo *et al.*, Broad and potent activity against SARS-like viruses by an engineered human monoclonal antibody. *Science* **371**, 823-829 (2021).
12. G. Bajic *et al.*, Influenza Antigen Engineering Focuses Immune Responses to a Subdominant but Broadly Protective Viral Epitope. *Cell Host Microbe* **25**, 827-835 e826 (2019).
13. G. Bajic *et al.*, Structure-Guided Molecular Grafting of a Complex Broadly Neutralizing Viral Epitope. *ACS Infect Dis* **6**, 1182-1191 (2020).
14. M. Crispin, A. B. Ward, I. A. Wilson, Structure and Immune Recognition of the HIV Glycan Shield. *Annu Rev Biophys* **47**, 499-523 (2018).
15. G. Ofek *et al.*, Elicitation of structure-specific antibodies by epitope scaffolds. *Proc Natl Acad Sci U S A* **107**, 17880-17887 (2010).
16. B. E. Correia *et al.*, Proof of principle for epitope-focused vaccine design. *Nature* **507**, 201-206 (2014).
17. L. Piccoli *et al.*, Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike Receptor-Binding Domain by Structure-Guided High-Resolution Serology. *Cell* **183**, 1024-1042 e1021 (2020).
18. C. O. Barnes *et al.*, SARS-CoV-2 neutralizing antibody structures inform therapeutic strategies. *Nature* **588**, 682-687 (2020).
19. Y. Wu *et al.*, A noncompeting pair of human neutralizing antibodies block COVID-19 virus binding to its receptor ACE2. *Science* **368**, 1274-1278 (2020).
20. J. Hansen *et al.*, Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail. *Science* **369**, 1010-1014 (2020).

21. M. Letko, A. Marzi, V. Munster, Functional assessment of cell entry and receptor usage
for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol* **5**, 562-569
(2020).
22. J. Shang *et al.*, Structural basis of receptor recognition by SARS-CoV-2. *Nature* **581**,
221-224 (2020).
23. V. S. Raj *et al.*, Dipeptidyl peptidase 4 is a functional receptor for the emerging human
coronavirus-EMC. *Nature* **495**, 251-254 (2013).
24. D. Pinto *et al.*, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290-295 (2020).
25. M. Yuan *et al.*, A highly conserved cryptic epitope in the receptor binding domains of
SARS-CoV-2 and SARS-CoV. *Science* **368**, 630-633 (2020).
26. N. C. Wu *et al.*, A natural mutation between SARS-CoV-2 and SARS-CoV determines
neutralization by a cross-reactive antibody. *PLoS Pathog* **16**, e1009089 (2020).
27. B. M. Hauser *et al.*, Engineered receptor binding domain immunogens elicit pan-
coronavirus neutralizing antibodies. *bioRxiv*, (2020).
28. K. Sliepen, T. van Montfort, M. Melchers, G. Isik, R. W. Sanders, Immunosilencing a
highly immunogenic protein trimerization domain. *J Biol Chem* **290**, 7436-7442 (2015).
29. W. F. Garcia-Beltran *et al.*, COVID-19-neutralizing antibodies predict disease severity
and survival. *Cell* **184**, 476-488 e411 (2021).
30. K. H. D. Crawford *et al.*, Protocol and Reagents for Pseudotyping Lentiviral Particles
with SARS-CoV-2 Spike Protein for Neutralization Assays. *Viruses* **12**, (2020).
31. V. D. Menachery *et al.*, SARS-like WIV1-CoV poised for human emergence. *Proc Natl
Acad Sci U S A* **113**, 3048-3053 (2016).
32. D. Zhou *et al.*, Evidence of escape of SARS-CoV-2 variant B.1.351 from natural and
vaccine induced sera. *Cell*, (2021).
33. W. F. Garcia-Beltran *et al.*, Circulating SARS-CoV-2 variants escape neutralization by
vaccine-induced humoral immunity. *bioRxiv*, (2021).
34. M. Yuan *et al.*, Structural and functional ramifications of antigenic drift in recent SARS-
CoV-2 variants. *bioRxiv*, (2021).
35. A. G. Schmidt *et al.*, Immunogenic Stimulus for Germline Precursors of Antibodies that
Engage the Influenza Hemagglutinin Receptor-Binding Site. *Cell Rep* **13**, 2842-2850
(2015).
36. A. G. Schmidt *et al.*, Viral receptor-binding site antibodies with diverse germline origins.
Cell **161**, 1026-1034 (2015).
37. M. J. Moore *et al.*, Retroviruses pseudotyped with the severe acute respiratory syndrome
coronavirus spike protein efficiently infect cells expressing angiotensin-converting
enzyme 2. *J Virol* **78**, 10628-10635 (2004).
38. E. Siebring-van Olst *et al.*, Affordable luciferase reporter assay for cell-based high-
throughput screening. *J Biomol Screen* **18**, 453-461 (2013).
499