Early Spring Snowmelt and Summer Droughts Strongly Impair the Resilience of Key

Microbial Communities in Subalpine Grassland Ecosystems

Farhan Hafeez[1,2], Lionel Bernard[3], Jean-Christophe Clément[4], Franck Poly[1], Thomas

Pommier[1*]

[1]*Laboratoire d'Ecologie Microbienne LEM, INRA UMR 1418, CNRS UMR 5557, Université*

de Lyon, F-69622 Villeurbanne Cedex, France

[2]*Department of Environmental Sciences, COMSATS University Islamabad, Abbottabad*

Campus, Tobe Camp University Road Abbottabad 22060 Pakistan.

 [3]*Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LECA, 38000 Grenoble, France*

[4]*Univ. Savoie Mont Blanc, INRAE, CARRTEL, 74200 Thonon-les-Bains, France*

Corresponding author :

Thomas Pommier

Laboratoire d'Ecologie Microbienne LEM, INRA UMR 1418, CNRS UMR 5557, Université

de Lyon, F-69622 Villeurbanne Cedex, France

Email : thomas.pommier@univ-lyon1.fr

ABSTRACT

Subalpine grassland ecosystems are important from biodiversity, agriculture, and touristic perspectives but their resilience to seasonally occurring climatic extremes is increasingly challenged with climate change, accelerating their vulnerability to tipping points. Microbial communities, which are central in ecosystem functioning, are usually considered as more resistant and highly resilient to such extreme events due to their functional redundancy and strong selection in residing habitats. To investigate this, we explored the soil microbial responses upon recurrent summer droughts associated with early snowmelt in grasslands mesocosms set-up at the Lautaret Pass (French Alps). Potential respiration, nitrification and denitrification were monitored over a period of two growing seasons along with quantification of community gene abundances of total bacteria as well as (de)nitrifiers. Results revealed that droughts had a low and short-term impact on bacterial total respiration supporting their hypothesized high resistance and ability to recover. Nitrification and abundances of the corresponding functional guilds showed relatively strong resistance to summer droughts but declined in response to early snowmelt. This triggered a cascading effect on denitrification but also on the abundances of denitrifying communities which could recover from all climatic extremes except from the summer droughts where nitrifiers were collapsed. Denitrification and the respective functional groups faced high impact of applied stresses with strong reduction in the abundance and activity of this specialized community. Although, the consequently lower microbial competition for nitrate may be positive for plant biomass production, warnings exist when considering the potential nitrogen leaching from these ecosystems as well as risks of greenhouses gases emission such as N_2O.

Key words: climate change, grasslands, (de)nitrification, weather extremes, snowmelt, N_2O

1. Introduction

Subalpine grassland ecosystems are important for biodiversity, agriculture, and a wide range of key services (Pecher et al., 2017; Zoderer et al., 2016; Tappeiner et al., 2008; Grêt-Regamey et al., 2012). These grasslands are under extreme pressure of climatic changes drastically affecting their functioning and disrupting the linked ecosystem services, *e.g.,* plants productivity, nutrient cycling etc. (Berauer et al., 2019; Bernard et al., 2019). Considering the continuous human perturbations and global changes (Berauer et al., 2019), it is direly required to investigate whether these grasslands are highly vulnerable, have developed resilience (resistance: ability to maintain the existing state, and recovery characteristics: to recover from disturbances) (Ingrisch and Bahn, 2018; Hodgson et al., 2015) or have attained an altered recovery state (Seabloom et al., 2020) over the time which is crucial in conserving the ecosystems functions (Donohue et al., 2016).

The multiple effects of the global climatic changes not only impact the biodiversity and ecosystem services (Berauer et al., 2019; Schirpke et al., 2013, Engler et al., 2011) but also their associated resistance (Bernard et al., 2019; Wipf and Rixen, 2010) and recovery in Alpine grasslands (Ingrisch et al., 2020; Karlowsky et al., 2018). The effect of the recurrent of climatic extremes on microbial functioning in these grasslands needs to be assessed as they may shift the fate of particular biogeochemical functions especially when resilience limits are crossed and tipping points (where a potentially irreversible change is about to occur or is already occurring *sensu* (Schnellhuber et al. (2006)) are reached. Such estimations are crucial to understand and possibly protect the stability of ecosystems and inhabiting species (de Vries and Shade, 2013; Griffiths and Philippot, 2013), but also to forecast the unforeseen consequences of adverse climatic extremes. Where weather extremes including droughts are predicted to be severe (Fuchslueger et al., 2019; Karlowsky et al., 2018), alpine grassland ecosystems are expected to face less snow falls that ultimately lead to subsequent droughts

(Klein et al., 2016; Gobiet et al., 2014). However, their vulnerability to collapse or support a particular function towards the combined (cumulative) impact of different weather extremes is poorly documented.

Microbial communities are crucial actors in ecosystem functioning (Chen et al., 2020; Bardgett and van der Putten, 2014) and respond strongly to environmental variability (Piton et al., 2020), thus are critical in estimating the functional stability perspectives (de Vries and Shade 2013; Griffiths et al., 2000). They are highly diverse, functionally redundant, and can be more resilient to climatic fluctuations as compared to many other environmental actors albeit the strong selection resulting from the local harsh climatic conditions (Capdeville et al., 2019; Griffiths and Philippot, 2013, Benot et al., 2014). Despite these yet known inherent capabilities, whether these microbial communities can withstand to correlated acceleration of weather extremes is largely unknown. Especially, the assessment of the functional boundaries of the subalpine grasslands and the linked ecosystems services is critical since they are susceptible to extreme events (Bernard et al., 2019).

The characteristics of existing plant species and nutrient availability as well as their interaction with microorganisms to ultimately respond to climatic changes are important in assessing the soil functioning resilience (Bernard et al., 2020). For example, any shift in the native plant community can influence the soil nutrient usage leading to the selection in the pool of residing microbes hence the corresponding functions (Legay et al., 2016; de Vries et al., 2012). Similarly, the weather extremes directly influencing the plant community may also impact the microbes (Bardgett et al., 2013; de Vries and Shade, 2013, Saccone et al., 2013). Earlier, summer droughts in subalpine grasslands are reported to increase leaf senescence while reducing plant productivity (Benot et al., 2014), as well as affecting N cycling-related microbial community and functions (Dai et al., 2020; Cantarel et al., 2012).

Here we determined the short-term effect of drought, snow removal, and their additive (recurrent) effect on microbial activities and size of the involved functional guilds and the following responses, to measure the resistance and recovery, respectively. Since the microbes, driving broad (*e.g.,* respiration, decomposition) and specialized soil processes (*e.g.* (de)nitrification), may respond differentially to various perturbations (Dai et al., 2020; Jusselme et al., 2016; Chaer et al., 2009; Wertz et al., 2007), we conducted our work both at broad and functional scales. Investigating the response of these grasslands is crucial since the variable impact of natural and manmade perturbations might have considerably influenced the nutrient cycling including that of N (Galloway et al., 2004), but also the involved microbial functional guilds. First, we tested how total bacterial community (16S *rRNA*) and related substrate-induced respiration (SIR) were influenced under droughts, snow removal and their cumulative impact and how they recovered from these stresses. Then we investigated the nitrogen (N) cycling (that is critical in ecosystem sustainability (Chen et al., 2020; Kuypers et al., 2018)) processes by determining the abundances (nitrifiers (*nxrA*, *NS*), denitrifiers (*nirK*, *nirS*, *nosZ*)) and activities (nitrification and denitrification enzyme activities -NEA and DEA, respectively) of the involved microbial communities, since they carry implications for plant productivity (loss of N - an important plant nutrient) and environment (N_2O emissions-global warming) (Dai et al., 2020; Galloway et al., 2004).

The current work was carried out on functionally contrasted and assembled grasslands set up in a common garden at Lautaret Pass with various sampling campaigns during two growing seasons. Based on an assumed functional redundancy (Jia and Whalen 2020; Griffiths and Philippot 2013), we hypothesized that the broad SIR and the relevant microbial community should resist and/or recover once the stresses/extreme events are over. SIR being representative of the total microbial communities is expected to be influenced by the soil moisture and abundance of *16S rRNA* (Hallin et al., 2009). However, the more specific

functions of N-cycling should be more sensitive due to comparatively less functional

redundancy and, despite being able to resist to single stress, the combined effects (SR+D)

may trigger a tipping point, disturbing the various steps of these processes. Moreover, any

shift in soil moisture, substrate availability and abundance of denitrifiers can strongly impact

DEA through affecting the micro-environment conditions (Philippot et al., 2007) both under

drought and no drought while NEA might be sensitive to ammonium availability (Jia and

Conrad 2009) and snow removal. To address these questions, we quantified the immediate

responses of the microbial communities' abundances and processes to droughts and early

snowmelt extreme events to explore the resistance and the following recovery across

subalpine grassland ecosystems.

2. Materials and methods

2.1. Experimental site, Mesocosms and Treatments' implementation

The study was set up at Lautaret pass proximity in southeast France (2,100 m a.s.l,

45°02'13''N, 6°24'01''E) with a multifactorial experimental design where 128 mesocosms

(n=128), 50-cm width and 40-cm depth, were installed in a common garden beside subalpine

grasslands. Mesocosms were filled with previously sieved (5-mm) *in-situ* soil and planted

with varying relative abundances of four species ranging along a plant economics spectrum.

Species richness and functional composition remained stable throughout the whole

experimentation due to low plant mortality. No snow removal (NSR) and snow removal (SR)

treatments (n=64+64) were performed and for each of the NSR and SR there was an

additional (sub) climate change treatment, i.e., maintained average soil moisture (ND) or

drought (D) (n=32+32). Treatments receiving no snow removal with maintained average soil

moisture were named as control (CTRL). All the 128 Mesocosms were installed inside 8

blocks (Fig. 1). Within each of the 8 blocks with or without snow removal, there were two

sub-blocks either maintained at long-term average soil moisture or subjected to summer

drought. Within each sub-block 8 mesocosms were installed; and 4 were submitted to plant

cuttings and fertilization (CF) while the other half remained untreated (NCF) (n=16+16).

During spring 2014, slow-release fertilizer was applied as pellets in the CF mesocosms (150

kg N/ha). To simulate early snowmelt, snow cover was removed one month earlier than the

average snowmelt period which ultimately simulated a shorter snow-covered period followed

by an early spring drought. The drought treatment was implemented using rainout shelters

during spring, summer and fall. These shelters were thus in use during the whole vegetation

season and all mesocosms were watered manually to maintain long-term average soil

moisture. This manual watering was discontinued to achieve drought that reduced soil

moisture for more than one month in August - September 2013 and June - July 2014.

Fig. 1. Experimental timeline showing the implementation of various treatments as well as the

sampling dates.

2.2. Soil sampling campaigns

Five sampling campaigns were carried out in June 2013 (T0), September 2013 (T1),

May 2014 (T2), July 2014 (T3) and September 2014 (T4). 1st and 2nd D were applied in

August - September 2013 and June - July 2014 respectively. The SR treatment was applied before T2 sampling campaign. Resistance to 1^{st} D of the microbial communities was measured at T1, resistance to 1^{st} SR and recovery from 1^{st} D at T2, resistance to 2^{nd} D and recovery from 1^{st} SR at T3; and recovery from overall D and SR treatments was evaluated at T4 sampling campaign (Fig. 1). During all sampling campaigns composite soil samples (5 cores mixed together, 0-10 cm depth) were collected from all 128 mesocosms (128x5: n=640). Fresh soil samples were sieved at 2-mm, homogenized and brought to the laboratory to measure the enzyme activities in fresh soil while a sub-sample was stored at -20°C for DNA extraction. Soil properties including soil moisture content, SOM% (soil organic matter), soil pH, soil %C, soil %N, total dissolved nitrogen-TDN, and the dissolved organic nitrogen-DON, soil NH_4^+, and soil NO_3^- were also measured in an associated work with focus on the effects of local environmental changes on subalpine grasslands functioning (Bernard 2017). Specifically, subsamples of 5 g fresh soil were dried at 70°C for 1 week to determine soil moisture content (in g.g-1 dw calculated as the 70°C dry soil weight relative to the fresh mass), followed by 4 h at 550° C to determine SOM (calculated as the 550°C soil weight relative to the 70°C dry mass). 10 mg soil subsample were air dried, ground to powder and analyzed for soil total C and N concentrations using a C/N elemental analyzer (FlashEA 1112, ThermoFisher Scientific, MA, USA). Soil pH was measured in a 1:4 (air dry soil/distilled water) solution. Soil $N-NH_4^+$ and $N-NO_3^-$, N-TDN and N-DON were determined in separated 10 mg fresh soil subsamples extracts (K_2SO_4, 0.5 M) using an FS-IV colorimetric chain (OI-Analytical, College Station, TX, USA) (Jones and Willett, 2006).

2.3. Soil DNA extractions and quantifications

DNA was extracted from soil stored at -20°C using the PowerSoil® DNA Isolation Kit (Mo Bio Laboratories Inc. USA) and DNA concentration was determined by using the Quant-iTTM PicoGreen dsDNA Assay Kit (Invitrogen, Carlsbad, CA, USA).

 2.4. Microbial community abundances

The abundance of the total bacterial communities was estimated in all DNA extracts

using a *16S rRNA* primer-based quantitative PCR (qPCR) assays as described by Lane (1991).

Final reaction volume was 20-µl, with 300-nM of forward and reverse primers; 1 X

LightCycler DNA Master Mix SYBR® Green I (Roche Diagnostics); 1-ng of sample DNA or

10^8 –10^2 copies of the standard DNA pQuantAlb16S plasmid (Zouacheet al., 2012). The

abundances of the *NS, nxrA, nirS, nirK and nosZ* genes were estimated using the primers

developed previously (Attard et al., 2011; Henry et al., 2004; Henry et al., 2006; Kandeler et

al., 2006). The total volume of the reaction was made 20 µl for *nxrA* and *nirK*, 500 nM for

nxrA while 1µM for *nirK* for both the forward as well as reverse primers; 10 µL of the

SYBR® Green I kit was used for the *nxrA* but also for the *nirK*, 0.4 µg of T4 protein

(Qbiogene, Carlsbad, CA USA) for *nirK*. The total volume of the reaction was made 25 µl for

NS, *nirS* but also for the *nosZ* while 400 nM with *NS* while 1µM for *nirS* and *nosZ* for both

the forward as well as reverse primers; 12.5 µL of the SYBR® Green I kit was used for the

NS, *nirS* but also with *nosZ*; for *nirS* the T4 protein was 0.5 µg while it was 0.4 µg for *nosZ*.

The curves for the standards for *NS*, *nxrA*, *nirS, nosZ and also for the nirK* were acquired

after the quantitative PCR assays with regular dilutions with already known concentrations of

the plasmid DNA with corresponding gene (10^7 - 10^1 number of gene copies). Quantification

of all gene community abundances was conducted out on a lightcycler 480 (Roche

Diagnostics). The amplification conditions used for qPCR assays were: 10 min at 95°C for

each 40 cycles for *16S rRNA*, 45 for *nxrA*, *NS*, *nirK* and *nirS* and 40 for *nosZ* (after 6 cycles of

touchdown); 95°C for 15 s for *16S rRNA*, *nirK*, *nirS*, *nosZ* and for 30 s for *nxrA* and NS; 63°C

for 1 min for *16S rRNA*, 55°C for 45 s for *nxrA* and 66°C for 30 s for *NS*, 30 s for *nirK* and

nirS, 60°C for 15 s for *nosZ*. 72°C for 30 s for 16S *rRNA*, *nirK*, *nirS* and *nosZ*, 45 s for *nxrA*,

1 min for NS; and additional step of 80°C for 15 s *nirS* and *nosZ*. Then 95°C for 5 s for 16S

rRNA and *nxrA*, 1 s for *NS*, *nirK* and *nirS* and 15 s for *nosZ*; 65°C for 1 m for *16S rRNA* and

nxrA, 68°C for 20 s for *NS* and *nirK*, 60°C for 20 s for *nirS* and 15 s for *nosZ*. And a

continuous increase to 98°C for *16S rRNA*, *nxrA*, *NS, nirK* and *nirS* and 95°C for *nosZ* to

determine the melting point; and finally, 10 s at 40°C for *16S rRNA*, *nxrA*, *NS*, *nirK* and *nirS*

while 30 s for *nosZ* for cooling. At least two independent quantitative PCRs were done for

every sample while the average was used. Co-amplification of the standards and the dilution

of samples were used to control any inhibition in the quantification.

2.5. Microbial activities

Susbtrate Induced Respiration MicroResp. (SIR) was measured as CO_2 production

using MicroResp plates (Chapman et al., 2007). Soil was added to the wells of 96-well plates.

0.5-ml of a nutritive solution including glucose (1.2-mg of C-glucose g of dried soil) then

incubated at 28°C. CO_2 concentrations were measured using a TECAN spectrophotometer

after 6-hours of incubation and the slope of the linear regression was used to estimate aerobic

respiration as the CO_2 produced (ppmV CO_2 h^{-1} g^{-1}).

Nitrifying enzyme activity (NEA) was measured on fresh soil samples as described by

Patra et al. (2005). In brief, the fresh soil sample (equal to 3-g dry weight) was incubated with

3 ml solution of $N-NH_4$ (200 µg $N-(NH_4)_2SO_4$ g^{-1}) in the flask for 72-hours at 180-rpm and

28°C. The flask was sealed with parafilm® (Pechiney Plastic Packaging, Menasha, WI, USA)

to produce the aerobic conditions. With distilled water, the total volume of the suspension was

made up to 15-ml. During the incubation, at regular intervals of 5, 24, 48 and 72-hours soil

slurry was filtered with a pore size of 0.2-µm. The rates of production of NO_2/NO_3 were

measured by ionic chromatography (DX120. Dionex, Salt Lake City, UT, USA).

Denitrification enzyme activities (DEA) was measured by incubating the fresh soil

sample in a 150-ml airtight plasma-flask sealed by a rubber stopper for 4-hours at 28°C

followed by the N_2O measured each 30 minutes for 3-hours a gas chromatograph coupled to a

micro-catharometer detector (lGC-R3000; SRA instruments) using the method adapted from

Dassonville et al. (2011). Briefly, a substrate solution (2-ml) containing glucose (0.5-mg of C-

glucose g^{-1} of dried soil), glutamic acid (0.5-mg of C-glutamic acid g^{-1} of dried soil) and

potassium nitrate (50-µg of N-KNO$_3$ g^{-1} of dried soil) were added to the soil. A 90:10 He-

C_2H_2 atmosphere provided anaerobic conditions and inhibited N_2O-reductase activity using

the acetylene inhibition technique that blocks the last step in the denitrification pathway

resulting in the accumulation of N_2O (Yoshinari et al., 1977). The soil was incubated for 4-

hours at 28°C and after incubation the amount of N_2O was measured each 30 minutes for 3-

hours. The amount of N_2O produced was measured using a gas chromatograph coupled to a

micro-catharometer detector (lGC-R3000; SRA instruments).

2.6. Statistics and Structural Equation Model

The data obtained were statically analyzed using JMP8® (SAS Institute Inc., SAS

Campus Drive, NC, USA). Normally distributed data was analyzed using the Student's test

and Tukey pairwise comparison. Where the data was not normally distributed after a log

transformation further analysis was carried using the Kruskal-Wallis and Kruskal-Wallis

pairwise comparison. Structural Equation Model (SEM) was constructed using the AMOS

(Amos Development Corporation, Crawfordville, Florida, USA) to link the environmental and

microbiological variables to investigate the relationship between these parameters under

different climatic extremes. The fitness of the model was assessed by using the χ2 test and the

model was considered as correct if the χ2 test produced a non-significant p-value - p>0.05.

Since the factors controlling the microbial functioning under the drought and/or snow removal

treatments were changing upon these treatments, the hypothesized models were changed to

increase the level of significance and to fit the model. SEM were constructed to understand

the relationship and path coefficients among the various causal environmental and microbial

factors under D, SR, SR+D treatments. We also carried out linear regression modeling for the

studied microbial parameters and associated soil properties and the SEM were fitted

considering the parameters showing strong correlation.

3. Results

3.1. Changes in Broad Scale Function and Community

The trajectory followed for the soil moisture at all sampling campaigns compared to

CTRL showed the significant effect of the D, SR and SR+D treatments (Fig. 2a; Fig. S1). The

resistance and recovery of the N-related microbial activities both for nitrification and

denitrification and the abundances of corresponding genes were calculated as the relative

change in D, SR, and SR+D treatments in comparision to CTRL. It was observed that the SIR

was found to be significantly (p=0.001) lowered upon both droughts and was recorded up to

89% of CTRL (43.83 ppmV CO_2 h^{-1} g^{-1} ± 0.58) but recovered ultimately (p=0.639). Although

the interaction of SR+D treatments marginally (p=0.0123) affected the SIR, it was not

affected by SR alone (p=0.625). It was noteworthy that the abundance of the total bacterial

community (as expressed by 16S *rRNA* gene copies) followed the similar trend as SIR and

was significantly (p=0.0006) reduced up to 56.3% of CTRL (4.87 x 10^8 copies per g of dry

soil) with a complete recovery observed at T4 (Fig. 2; Fig. S1).

Fig. 2. Relative change compared to CTRL for different treatments at various sampling dates
a) in soil moisture; **b)** in SIR-MicroResp™; **c)** in abundance of total bacterial community
(number of 16S *rRNA* gene copies g⁻¹ dry soil); Light grey dotted lines show variations in
time of the variables in the snow removal (SR) treatment relative to CTRL; dark grey dotted
lines show variations of the Drought (D) treatment; and the black dotted lines show the
variations of cumulative SR+D treatment.

3.2. Response of N Cycling Processes to Droughts and Snowmelt

Nitrification (NEA) showed resistance to both droughts at T1 and T3 and thus remained independent of the change in moisture content. Whereas there was a significant reduction in NEA (up to 76% of CTRL) as observed upon SR treatment at T2 (p=0.01, Fig. 3; Fig. S2). To assess the linkages, the multiple regression analyses were carried out for the parameters in addition to moisture contents and the findings showed that the NEA was positively correlated to NH_4, %N, TDN and *nxrA* (r = 0.56, 0.20, 0.31, 0.20; p < 0.005). On the contrary, DEA was significantly affected by both droughts and was found to be lowered up to 85% and 68% of CTRL upon first and second drought, respectively (Fig. 4; Fig. S3). Recovery from the first drought was observed for DEA at T2 except for the interaction of SR+D treatment. However, the second drought (at T3, Fig. 4) strongly (p=0.0001) impacted DEA which did not recover from SR. Hence, at T4 there was an incomplete recovery in denitrification acitivity as the DEA for the SR+D mesocosms could not return to the initial levels (0.362 µg $N-N_2O$/g dry soil/h ± 0.022).

3.3. Impact on N Cycling Functional Guilds

The results revealed that the abundances of the N cycling functional guilds including the nitrifiers and denitrifiers were strongly impacted by both D and SR climatic extremes (p=0.001). We observed that the *Nitrospira* and *Nitrobacter* (*nxrA* and *NS)* abundances were significantly (p=0.001) reduced upon applied stresses which could recover only for *nxrA* (Fig. 3). The denitrifiers' abundances were particularly impacted by the second drought, that strongly lowered the gene abundances of this functional group. At T4, the community gene abundances of the *nirS* and *nosZ* denitrifiers were completely recovered, but the *nirK* denitrifiers could not recover which was in agreement to the observations as recorded for DEA (Fig. 4; Fig. S3). The multiple correlation analyses as discussed in the next section

showed the strong positive correlations of DEA with the %C, %N, DON, SOM, soil moisture, NEA as well as with the abundances of denitrifying genes. However, these correlations were found more prominent in the climate change treatments than the CTRL especially for soil moisture. For example, the Spearman correlation coefficient showed a non-significant correlation between the soil moisture and *nirK* denitrifiers (r = 0.04; p > 0.652) only for the CTRL whereas it was positive in the SR+D treatment (r = 0.33; p < 0.005).

3.4. Microbial Variables and Climatic Treatments, a Structural Equation Modeling

Fig. 5 represents how the various environmental (soil moisture, NH_4, NO_3 etc.) and microbial parameters (gene abundances) influenced the global and functional activities of microbial communities at different levels of drought and snow removal as well as by their interaction. This initial SEM was constructed for each sampling interval to observe how the various environmental and microbial factors influenced the global and functional microbial activities over each period of time. Afterwards, we separately analyzed the most correlated parameters from the results of the linear regression modeling and these parameters were linked to fit the SEM for the drought, snow removal treatments but also for their interaction. The SEM revealed that the SIR was linked to the soil moisture and *16S rRNA* for various treatments at different sampling times except at T1 and T2 (χ^2 = 14.35, p = 0.35; Fig. 5). In contrast, NEA was never associated to the soil moisture and was found to be mainly linked with soil NH_4 concentration and community gene abundances of the *NS* nitrifiers, except for the SR treatment. Interestingly, under droughts, major part of the DEA variance (up to 55%) was explained by the soil moisture with *nirS* as dominant denitrifiers except at T3 when it was *nosZ*. It was also observed that the overall impact of snow removal and drought was variable for the different denitrifiers with severe impact on *nirK* which could not recover even by the end of experiment. No significant effects of plant functional composition and fertilization on microbial parameters were recorded.

Fig. 3. a) Relative change compared to CTRL in **a)** nitrification enzyme activities (µgN-(NO₂+NO₃)/g dry soil/h); **b)** in abundance of *nxrA* nitrite reductases (number of *nxrA* gene copies g⁻¹ dry soil); **c)** in abundance of *NS* nitrite reductases (number of *NS* gene copies g⁻¹ dry soil); Light grey dotted lines show variations in time of the variables in the snow removal (SR) treatment relative to CTRL; dark grey dotted lines show variations of the Drought (D) treatment; and the black dotted lines show the variations of cumulative SR+D treatment.

Fig. 4. a) Relative change compared to CTRL in **a)** denitrification enzyme activities (µg N-N_2O/g dry soil/h); **b)** in abundance of *nirK* denitrifiers (number of *nirK* gene copies g^{-1} dry soil); **c)** in abundance of *nosZ* denitrifiers (number of *nosZ* gene copies g^{-1} dry soil); Light grey dotted lines show variations in time of the variables in the snow removal (SR) treatment relative to CTRL; dark grey dotted lines show variations of the Drought (D) treatment; and the black dotted lines show the variations of cumulative SR+D treatment.

4. Discussion

Determining the resilience including the resistance and recovery capabilities at subalpine grassland ecosystems is crucial to understand since these ecosystems are under drastic influence of global climatic changes which ultimately may disrupt their functioning and the linked ecosystem services (Berauer et al., 2019; Bernard et al., 2019; Schirpke et al., 2013; Wipf and Rixen, 2010; Vittoz et al., 2009). This work demonstrates the impact of successive droughts and early snowmelt climatic extremes on subalpine grassland ecosystem functioning focusing on nitrification and denitrification as model processes since these give valuable insights into overall N cycling (Chen et al., 2020; Dai et al., 2020).

The quantification of resistance and recovery by comparing the relative change in D, SR, and SR+D treatments inferred how the microbial activities and the corresponding functional groups responded upon different climate change treatments. Broad scale function – SIR and the size of the total bacterial community - showed low resistance but high recovery for all treatments which strengthened the fact that the wide range of microbial communities are carrying this function generally exhibit a fast recovery (Ingrisch et al., 2020). SIR represents the broad scale microbial activity thus, in our study, disturbing a part of the global community could not affect the overall function greatly. Yet, the short term reduction in SIR upon droughts can be associated to the low microbial activity induced by the water deficit conditions (Bernard et al., 2019; Bloor and Bardgett, 2012). Earlier, Chaer et al. (2009) elaborated that the broad scale functions are less sensitive to disturbance. This is similar to the observations of weak impact of climatic stresses for decompostion processes in a joint experiment (Bernard et al., 2019), and high recovery of microbial biomass in other mountainous grasslands (Piton et al., 2020). These are in accordance to the functional redundancy concepts (Jia and Whalen, 2020; Griffiths and Philippot, 2013).

Fig. 5. Structural equation models (SEM) bridging the effects of different microbial and environmental factors. SEMs including the Substrate induced respiration MicroResp (SIR), nitrification (NEA), denitrification (DEA), gene community abundances (16S *rRNA*, *nxrA*, *NS*, *nirK*, *nirS*, *nosZ*) and environmental factors (soil moisture, NH₄, NO₃), in **a, b, c** are built using the samples from treatments; SR, D, SR+D, respectively. The values shown beside the

arrows are the path coefficients and correspond to the standardized coefficients calculated

based on the analysis of correlation matrices. Values only for the significant relations (that fit

the model, $P > 0.05$) are indicated in the SEMs.

    The impact of extreme events including droughts on microbial N cycling are yet less

clear (Fuchslueger et al., 2019), however the climate change treatments here showed

contrasting impacts on N cycling processes with denitrification (community gene abundance

and activity) showing less resistance to D as compared to nitrification (community gene

abundance and activity). Moreover, at T3 when snow removal and droughts occurred in the

same mesocosms (SR+D), they strongly reduced the DEA followed by an incomplete

recovery, which was in contrast to the NEA that recovered compeletly at T4. This confirmed a

strong influence of droughts and associated soil moisture deficits on DEA (Di et al., 2020;

Philippot et al., 2009), which was especially pronounced for SR+D indicating the cascading

effect of SR and declined nitrification upon snowmelt. The collapse in microbial pool size

upon snowmelt has also been observed while studying the niche differentiation perspectives

in soils (Sorensen et al., 2020). A strong turnover by these nitrifying and denitrifying

microbial communities was previously shown for similar subalpine grasslands upon natural

variations in snowpack depth (Jusselme et al., 2016). Such multiple cascading effects on N-

related microbial activities are important since these can adversely impact the greenhouse gas

emissions (i.e., N_2O in particular), soil fertility and plant productivity hence ask for adaptive

management strategies (Griffis et al., 2014; Di et al., 2014; Bouwman et al., 2009).

    The reduction in gene abundances of both nitrifiers and denitrifiers upon D and SR

delineated the strong impact of studied climate change treatments on grasslands functioning.

Although denitrifiers are reported as more tolerant to droughts (Hammerl et al., 2019) than

nitrifiers (Austin and Strauss, 2010), here in agreement to DEA steadiness, the denitrifiers'

abundances were also negatively impacted by droughts, especially the second one. In line

with DEA, the *nirK* denitrifiers' abundances were severely lowered and could not show a

complete recovery at T4. Such correlations were previously recorded between DEA and the

abundance of *nirK* denitrifiers upon variable snow depths under similar subalpine grasslands

(Jusselme et al., 2016). Conversely, Wertz et al. (2007) found that denitrifiers were less

affected by the heat stress in an agricultural soil and it was associated to the high functional

redundancy. Besides, resilience of microbial communities is also influenced by the inherent

soil edaphic characteristics (Berard et al., 2015). Although we found strong positive

correlations between the DEA and %C, %N, DON, SOM, soil moisture, NEA, denitrifiers'

abundances, interestingly these controlling factors shifted during the studied time. For

example, in SR+D treatment, soil moisture along with *nirK*, *nirS* and *nosZ* were correlated to

DEA (r = 0.33, 0.15, 0.32, 0.38; p < 0.005), while soil moisture and *nirK* had not shown any

significant correlation with DEA (r = 0.04; p > 0.64) in the CTRL. This also shows the

variable dominance of different denitrifiers upon D, SR and CTRL treatments. These different

variables, across a range of environments, are shown to be differently correlated to DEA

(Chen et al., 2020; Chen et al., 2019; Petersen et al., 2012; Chroňáková et al., 2009, Dong et

al., 2009). At T4, DEA could not recover reflecting a severe impact of the subsequent

warming caused both by D and the additive effect through SR+D. Possibly the system, (as

described by Schnellnhuber et al. (2006)) upon disturbance was 'pulsed' (respond on

alteration but return) and 'pressed' (respond on sustained alteration so shift) where 'press'

brings the ecosystems to a tipping point (as observed for DEA in our case) and may shift to an

irreversible new state (Wall, 2007), or at least may lead to altered recovery patterns

(Seabloom et al., 2020). Furthermore, such changes in the relative abundances of the

denitrifiers may also result in the selection of certain stable/unstable groups, which may

consequently impact (increase) the N losses (NO_3 leaching or N_2O emissions) (de Vries and

Shade 2013). In addition, studies in other ecosystems have also suggested that these

functional guilds may have contrasting resilience abilities against various disturbances

(Capdeville et al., 2019).

Though SIR was related to soil moisture and *16S rRNA* when the SEM was fitted,

NEA was highly linked to the soil NH_4^+ and the abundance of the *NS* nitrifiers hence showing

a weak effect of soil moisture when droughts occurred. Higher abundance of *NS* nitrifiers than

nxrA as well as the dominant role of *NS* nitrifiers under drought was earlier shown in low

activity soils (Attard et al., 2010). However, DEA was related both to soil moisture and

abundances of *nirS* denitrifiers throughout the experiment except at T3 when it was *nosZ*.

After the second drought at T3, a cascading effect of SR might have shifted the community

dynamics from *nirS* to *nosZ* with better adaptation of the *nosZ* harboring denitrifiers. The

denitrifiers responded differentially, *e.g.,* in contrast to *nirK*, the *nirS* and *nosZ* were

recovered, showing a community specific impact (Karlowsky et al., 2018) of drought and

snow removal. The *nirK* were previously reported to be less abundant (Peterson et al., 2012),

and relatively more sensitive to environmental change than *nirS* in agricultural soils (Szukics

et al., 2019; Hallin et al., 2009). Though the findings implied by Dai et al. (2020) contrasted

with these observations, seasonally driven drought effects have also been reported for *nirK*,

nirS and *nosZ* dynamics in grassland ecosystems (Hammerl et al., 2019). Moreover, the

denitrifiers harboring *nirK* genes here might not contain *nosZ* genes, in agreement to the

discovery that not all denitrifiers contain *nosZ* (Hallin et al., 2018).

We did not find any significant effect of plant functional composition and fertilization

on microbial parameters; hence these were not the main drivers of the studied processes but

there might be a trait-centered influence of different variables (Griffin-Nolan et al., 2019). For

example, leaf or root specific traits, rather than plant composition, might have affected the

nutrient availability and ultimately associated microbial processes. Interestingly under similar

grasslands, no direct effect of fertilization was observed but only specific plant traits were affecting the same microbial activities (Legay et al., 2016). This may also suggest a competition for nutrients between plants and microbes resulting in a non-significant fertilization effect on the microbial parameters during the study period. However, such management strategies including fertilization practices may also favor certain microbial communities consequently causing their fast resilience to stress conditions (Piton et al., 2020). It is noteworthy that due to the climatic extremes, the responses showed by the various nitrifiers' and denitrifiers' groups were different to CTRL conditions indicating an unanticipated internal competition for initially available nutrients. Importance of such competitions will be crucial to further investigate especially in ecosystems facing climatic extremes since these are suggested to be critical in controlling the ecosystem N cycling (Legay et al., 2013; Grigulis et al., 2013, Le Roux et al., 2013).

5. Conclusions

This work clearly demonstrated a strong and unprecedented impact of early snowmelt and recurrent droughts, alone or in combination, on soil microbial communities both at broad (i.e., soil respiration) and functional (nitrifiers' and denitrifiers' activities and abundances) scales. The droughts showed a low and short-term impact on bacterial total respiration supporting their assumed high resistance and ability to recover. Though the nitrification and community gene abundances of the corresponding functional guilds exhibited relatively strong resistance to summer droughts, it declined in response to early snowmelt leaving the strong impact on denitrification. The findings inferred that the predicted climatic extremes may carry severe consequences for the soil microbial community functioning in subalpine grasslands ultimately resulting in shifts in N availability, plant community diversity, and associated ecosystem services.

Conflict of Interest

The authors of this article declare that they have no financial conflict of interest with the

content of this article.

Acknowledgements

This work was part of the ERA-Net BiodivERsA project REGARDS (ANR-12-EBID-004-

01). Quantitative PCR were carried out at the platform DTAMB (IFR 41, Université Lyon 1)

and N cycling activities at the platform AME (UMR5557-UMR1418). We want to thank all

the people who helped us in the field or in the lab: C. Arnoldi, A. Foulquier, N. Guillaumaud.

This research was conducted at 'Jardin du Lautaret'(UMS 3370 Univ. Grenoble Alpes –

CNRS), a member of the Fédération FREE-Alpes and of the AnaEE-France (ANR-11-INBS-

0001AnaEE-Services), and on the LTSER platform 'Zone Atelier Alpes', a member of the

ILTER-Europe network.

References

Attard, E., Recous, S., Chabbi, A., Berranger, C.D., Guillaumaud, N., Labreuche, J.,

Philippot, L., Schimid, B., Le Roux, X., 2011. Soil environmental conditions rather

than denitrifier abundance and diversity drive potential denitrification after changes in

land uses. Glob. Chang. Biol. 17, 1975-1989.

Attard, E., Poly F., Commeaux, C., Laurent, F., Terada, A., Smets, B.F., Recous, S., Le Roux,

X., 2010. Shifts between Nitrospira- and Nitrobacter-like nitrite oxidizers underlie the

response of soil potential nitrite oxidation to changes in tillage practices. Environ.

Microbiol. 12(2), 315-26.

Austin, B.J., Strauss,, E.A., 2011. Nitrification and denitrification response to varying periods

of desiccation and inundation in a western Kansas stream. Hydrobiol. 658, 183-195.

Bardgett, R.D., Manning, P., Morriën, E., De Vries, F.T., 2013. Hierarchical responses of

plant–soil interactions to climate change: consequences for the global carbon cycle. J.

Ecol. 101, 334-343.

Bardgett, R.D., van der Putten, W.H., 2014. Belowground biodiversity and ecosystem

functioning. Nat. 515, 505–511.

Benot M.L., Saccone, P., Pautrat, E., Vicente, R., Colace, M.P., Karl, G., Clément, J.C.,

Lavorel, S., 2014. Stronger Short-Term Effects of Mowing Than Extreme Summer

Weather on a Subalpine Grassland. Ecosys. 17, 458-472.

Berauer, B.J., Wilfahrt, P.A., Arfin-Khan, M.A.S., Eibes, P., Heßberg, A.V., Ingrisch, J.,

Schloter, M., Schuchardt, M.A., Jentsch, A., 2019. Low resistance of montane and

alpine grasslands to abrupt changes in temperature and precipitation regimes. Arct.

Antarc. Alp. Res. 51:1, 215-231.

Bernard, L., Foulquier, A., Gallet, C., Lavorel, S., Clément, J.C., 2019. Effects of snow pack

reduction and drought on litter decomposition in subalpine grassland communities.

Plant Soil. 435, 225–238.

Bernard, L., Decau, M.L., Morvan-Bertrand, A., Lavorel, S., Clément, J.C., 2020.

Water-soluble carbohydrates in Patzkea paniculata (L.): a plant strategy to tolerate

snowpack reduction and spring drought in subalpine grasslands. Plant Biol. 22, 441-

449.

Bernard L., 2017. Mécanismes fonctionnels de résilience des prairies subalpines au

changement globale. PhD thesis, Université Grenoble-Alpes.

Berard, A, Ben Sassi, M, Kaisermann, A, Renault, P., 2015. Soil microbial community

responses to heat wave components: drought and high temperature. Clim. Res.

66:243–264

Bloor, J.M.G., Bardgett, R.D., 2012. Stability of above-ground and below-ground processes
to extreme drought in model grassland ecosystems: interactions with plant species
diversity and soil nitrogen availability. Perspect. Plant Ecol. Evol. Syst. 14: 193–204

Bouwman, A.F., Beusen, A.H.W., Billen, G., 2009. Human alteration of the global nitrogen
and phosphorus soil balances for the period 1970–2050. Glob. Biogeochem. Cyc. 23,
GB0A04.

Cantarel, A., Bloor, J.M.G., Pommier, T., Guillaumaud, N., Moirot, C., Soussana, J.F., Poly,
F., 2012. Four years of experimental climate change modifies the microbial drivers of
N_2O fluxes in an upland grassland ecosystem. Glob. Chang. Biol. 18, 2520-2531.

Capdeville, C., Pommier, T., Gervaix, J., Fromard, F., Rols, J.L., Leflaive, J., 2019.
Mangrove Facies Drives Resistance and Resilience of Sediment Microbes Exposed to
Anthropic Disturbance. Front. Microbiol. 9:3337

Chaer, G., Fernandes, M., Myrold, D., Bottomley, P., 2009. Comparative resistance and
resilience of soil microbial communities and enzyme activities in adjacent native
forest and agricultural soils. Microb. Ecol. 58, 414-424.

Chapman, S.J., Campbell, C.D., Artz, R.R.E., 2007. Assessing CLPPs using MicroResp™. J.
Soils Sedim. 7, 406-410.

Chen, Q.L., Ding, J., Li, C.Y., Yan, Z.Z., He, J.Z., Hu, H.W., 2020. Microbial functional
attributes, rather than taxonomic attributes, drive top soil respiration, nitrification and
denitrification processes, Sci. Tot. Environ. 734,139479.

Chen, J., Li, J., Shen, W., Xu, H., Li, Y., Luo, T., 2019. The Structure and Species Co-
Occurrence Networks of Soil Denitrifying Bacterial Communities Differ Between A
Coniferous and A Broadleaved Forests. Microorg. 7(9), 361.

Chroňáková, A., Radl, V., Čuhel, J., Šimek, M., Elhottová, D., Engel, M., Schloter, M., 2009.
Overwintering management on upland pasture causes shifts in an abundance of

denitrifying microbial communities, their activity and N2O-reducing ability. Soil Biol.

Biochem. 41, 1132-1138.

Dai, Z., Yu, M., Chen, H., Zhao, H., Huang, Y., Su, W., Xia, F., Chang, S.X., Brookes, P.C.,

Dahlgren, R.A., 2020. Elevated temperature shifts soil N cycling from microbial

immobilization to enhanced mineralization, nitrification and denitrification across

global terrestrial ecosystems. Glob. Chang. Biol. 26, 5267-5276.

Dassonville, N., Guillaumaud, N., Piola, F., Meerts, P., Poly, F., 2011. Niche construction by

the invasive Asian knotweeds (species complex Fallopia): impact on activity,

abundance and community structure of denitrifiers and nitrifiers. Biolog. Invas. 13,

1115-1133.

de Vries, F.T., Liiri, M.E., Bjørnlund, L., Setälä H.M., Christensen S., Bardgett R.D., 2012.

Legacy effects of drought on plant growth and the soil food web. Oecolog. 170, 821–

833.

De Vries, F.T., Shade, A., 2013. Controls on soil microbial community stability under climate

change. Front. Microbiol. 4, 265.

Di, H.J., Cameron, K.C., Podolyan, A., Robinson, A., 2014. Effect of soil moisture status and

a nitrification inhibitor, dicyandiamide, on ammonia oxidizer and denitrifier growth

and nitrous oxide emissions in a grassland soil. Soil Biol. Biochem. 73, 59–68

Dong, L.F., Smith, C.J., Papaspyrou, S., Stott A., Osborn A.M., Nedwell D.B., 2009. Changes

in Benthic Denitrification, Nitrate Ammonification, and Anammox Process Rates and

Nitrate and Nitrite Reductase Gene Abundances along an Estuarine Nutrient Gradient

(the Colne Estuary, United Kingdom). App. Environ. Microbiol. 75, 3171-3179.

Donohue, I., Hillebrand, H., Montoya, J.M., Petchey, O.L., Pimm, S.L., Fowler, M.S., Yang,

Q., Jackson, A.L., Lurgi, M., McClean, D., O'Connor, N.E., O'Gorman, E.J., Yang,

Q., 2016. Navigating the complexity of ecological stability. Ecol. Lett. 19:1172–85

Engler, R., Randin, C.F., Thuiller, W., et al., 2011. 21st century climate change threatens mountain flora unequally across Europe. Glob. Chang. Biol. 17:2330–2341.

Fuchslueger, L., Wild, B., Mooshammer, M., Takriti, M., Kienzl, S., Knoltsch, A., Hofhansl, F., Bahn, M., Richter, A., 2019. Microbial carbon and nitrogen cycling responses to drought and temperature in differently managed mountain grasslands. Soil Biol. Biochem. 135, 144–153.

Galloway, J.N., Dentener, F.J., Capone, D.G., Boyer,E.W., Howarth, R.W., Seitzinger,S.P., Asner, G.P., Cleveland,C.C., Green, P.A., Holland, E.A., Karl, D.M., Michaels, A.F., Porter, J.H., Townsend, A.R., Vöosmarty, C.J., 2004. Nitrogen cycles: past, present, and

future. Biochem. 70, 153–226.

Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., Stoffel, M., 2014. 21st century climate change in the European Alps—a review. Sci. Tot. Environ. 493:1138– 1151.

Grêt-Regamey, A., Brunner, S.H. Kienast, F., 2012. Mountain Ecosystem Services: Who Cares? Mount. Res. Develop. 32, S23–S34

Griffin-Nolan, R.J., Blumenthal, D.M., Collins, S.L., Farkas, T.E., Hofman, A.M., Mueller, K.E., Ocheltree, T.W., Smith, M.D., Whitney, K.D., Knapp, A.K., 2019. Shifts in plant functional composition following longterm drought in grasslands. J. Ecol. 107:2133–2148.

Griffis, T.J., Chen, Z., Baker, J.M., Wood, J.D., Millet, D.B., Lee, X., Venterea, R.T., Turner, P.A., 2014. Nitrous oxide emissions enhanced in a warmer world. PINAS. 114, 12081-12085

Griffiths, B.S., Philippot, L., 2013. Insights into the resistance and resilience of the soil microbial community. FEMS Microbiol. Rev. 37, 112-129.

Griffiths, B.S., Ritz, K., Bardgett, R.D., Cook, R., Christensen, S., Ekelund, F., Sørensen,

S.J., Bååth. E., Bloem, J., De Ruiter, P.C., Dolfing, J., Nicolardot, B., 2000.

Ecosystem response of pasture soil communities to fumigation-induced microbial

diversity reductions: an examination of the biodiversity–ecosystem function

relationship. Oikos. 90, 279-294.

Grigulis, K., Lavorel, S., Krainer, U., Legay, N., Baxendale, C., Dumont, M., Kastl, E.,

Arnoldi, C., Bardgett, R.D., Poly, F., Pommier, T., Schloter, M., Tappeiner, U., Bahn,

M., Clément, J.C., 2013. Relative contributions of plant traits and soil microbial

properties to mountain grassland ecosystem services. J. Ecol. 101, 47-57.

Hallin, S., Philippot, L., Löffler, F.E., Sanford, R.A., Jones, C.M., 2018. Genomics and

ecology of novel N_2O-reducing microorganisms. Trend Microbiol. 26, 43–55

Hallin, S., Jones, C.M., Schloter, M., Philippot, L., 2009. Relationship between N-cycling

communities and ecosystem functioning in a 50-year-old fertilization experiment. Int.

Soc. Microb. Ecol. 3, 597-605.

Hammerl, V., Kastl, E., Schloter, M., Kublik, S., Schmidt, H., Welzl, G., Jentsch, A.,

Beierkuhnlein, C., Gschwendtner, S., 2019. Influence of rewetting on microbial

communities involved in nitrification and denitrification in a grassland soil after a

prolonged drought period. Sci. Rep. 9, 2280

Henry, S., Baudoin, E., Lopez-Gutierrez, J.C., Martin-Laurent, F., Brauman, A., Philippot, L.,

2004. Quantification of denitrifying bacteria in soils by nirK gene targeted real-time

PCR. J. Microbiol. Method. 59, 327-335.

Henry, S., Bru, D., Stres, B., Hallet, S., Philippot, L., 2006. Quantitative detection of the nosZ

gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S

rRNA, narG, nirK, and nosZ genes in soils. App. Environ. Microbiol. 72, 5181-5189.

Hodgson, D., McDonald, J.L., Hosken, D.J., 2015. What do you mean, "resilient"? Trend.
638   Ecol. Evol. 30: 9, 503-506.

639   Ingrisch, J., Karlowsky, S., Anadon-Rosell, A., Hasibeder, R., Konig, A., Augusti, A.,
640   Gleixner, G., Bahn, M., 2018. Land Use Alters the Drought Responses of Productivity
641   and CO2 Fluxes in Mountain Grassland. Ecosys. 21, 689–703.

642   Ingrisch, J., Karlowsky, S., Hasibeder, R., Gleixner, G., Bahn, M., 2020. Drought
643   andrecovery effects on belowground respiration dynamics and the partitioning
644   ofrecent carbon in managed and abandoned grassland. Glob. Chang. Biol. 26, 4366-
645   4378.

646   Jia, Z., Conrad, R., 2009. Bacteria rather than Archaea dominate microbial ammonia
647   oxidation in an agricultural soil. Environ. Microb. 11, 1658-1671.

648   Jia, Y., Whalen, J.K., 2020. A new perspective on functional redundancy and
649   phylogeneticniche conservatism in soil microbial communities. Pedosph. 30, 18–24

650   Jones, D.L., Willett, V.B., 2006. Experimental evaluation of methods to quantify dissolved
651   organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. Soil Biol.
652   Biochem. 38, 991-999.

653   Jusselme, M.D., Saccone, P., Zinger, L., Faure, M., Le Roux, X., Guillaumaud, N., Bernard,
654   L., Clement, J.C., Poly, F., 2016. Variations in snow depth modify N-related soil
655   microbial abundances and functioning during winter in subalpine grassland. Soil Biol.
656   Biochem. 92, 27–37.

657   Kandeler, E., Deiglmayr, K., Tscherko, D., Bru, D., Philippot, L., 2006. bundance of narG,
658   nirS, nirK, and nosZ Genes of Denitrifying Bacteria during Primary Successions of a
659   Glacier Foreland. App. Env. Microbiol. 5957-5962.

660   Karlowsky, S., Augusti, A., Ingrisch, J., Akanda, M.K.U., Bahn, M., Gleixner, G., 2018.

661   Drought-Induced Accumulation of Root Exudates Supports Post-drought Recovery of

662   Microbes in Mountain Grassland. Front. Plant Sci. 2018;9:1593.

663   Klein, G., Vitasse, Y., Rixen, C., Marty, C., Rebetez, M., 2016. Shorter snow cover duration

664   since 1970 in the Swiss Alps due to earlier snowmelt more than to later snow onset.

665   Climat. Chang. 139:637–49.

666   Kuypers, M.M., Marchant, H.K., Kartal, B., 2018. The microbial nitrogen-cycling network

667   Marcel. Nat. Rev. Microbiol. 16, 263–276

668   Lane, D.J., 1991. 16S/23S rRNA Sequencing. In: Nucleic acid Techniques in Bacterial

669   Systematics (eds Stackebrandt E, Goodfellow M), pp. 115–175. John Wiley & Sons,

670   New York City, New York.

671   Le Roux, X., Schmid, B., Poly, F., Barnard, R.L., Niklaus, P.A., Guillaumaud, N., Habekost,

672   M., Oelmann, Y., Philippot, L., Salles, J.F., Schloter, M., Steinbeiss, S., Weigelt, A.,

673   2013. Soil environmental conditions and microbial build-up mediate the effect of plant

674   diversity on soil nitrifying and denitrifying enzyme activities in temperate grasslands.

675   PLoS One. 8, e61069.

676   Legay, N., Grassein, F., Binet, M.N., Arnoldi, C., Personeni E., Perigon, S., Poly, F.,

677   Pommier, T., Puissant, J., Clément, J.C., Lavorel, S., Mouhamadou, B., 2016. Plant

678   species identities and fertilization influence on arbuscular mycorrhizal fungal

679   colonisation and soil bacterial activities. App. Soil Ecol. 98, 132-139.

680   Legay, N., Grassein, F., Robson, T.M., Personeni, E., Bataillé, M.P., Lavorel, S., Clément,

681   J.C., 2013. Comparison of inorganic nitrogen uptake dynamics following snowmelt

682   and at peak biomass in subalpine grasslands. Biogeosci. 10, 7631-7645.

683   Patra, A.K., Abbadie, L., Clays-Josserand, A., Degrange, V., Grayston, S.J., Loiseau, P.,

684   Louault, F., Mahmood, S., Nazaret, S., Philippot, L., Poly, F., Prosser, J.I., Richaume,

A., Le Roux, X., 2005. Effects of grazing on microbial functional groups involved in soil n dynamics. Ecologic. Monograph. 75, 65-80.

Pecher, C., Bacher, M., Tasser, E., Tappeiner, U., 2017. Agricultural landscapes between intensification and abandonment: the expectations of the public in a Central-Alpine cross-border region. Landscap. Res. 1-15.

Petersen, D.G., Blazewicz, S.J., Firestone, M., Herman, D.J., Turetsky, M., Waldrop, M., 2012. Abundance of microbial genes associated with nitrogen cycling as indices of biogeochemical process rates across a vegetation gradient in Alaska. Environ. Microb. 14, 993-1008.

Philippot, L., Cuhel, J., Saby, N.P., Chèneby, D., Chronáková, A., Bru, D., Arrouays, D., Martin-Laurent, F., Simek, M., 2009. Mapping field-scale spatial patterns of size and activity of the denitrifier community. Environ. Microb. 11, 1518-1526.

Philippot, L., Hallin, S., Schloter, M., 2007. Ecology of Denitrifying Prokaryotes in Agricultural Soil. In: Advanc. Agron. pp Page., Academic Press.

Piton, G, Legay, N, Arnoldi, C, Lavorel, S, Clément, J-C, Foulquier, A., 2020. Using proxies of microbial community-weighted means traits to explain the cascading effect of management intensity, soil and plant traits on ecosystem resilience in mountain grasslands. J. Ecol. 108: 876– 893.

Saccone, P., Morin, S., Baptist, F., Bonneville, J.M., Colace, M.P., Domine, F., Faure, M., Geremia, R., Lochet, J., Poly, F., Lavorel, S. Clément, J.C., 2013. The effects of snowpack properties on litter decomposition during winter in subalpine meadows. Plant Soil. 363:215-229.

Seabloom, E.W., Borer, E.T. and Tilman, D., 2020. Grassland ecosystem recovery after soil disturbance depends on nutrient supply rate. Ecol. Lett. 23(12) 1756-1765.

Schirpke, U., Leitinger, G., Tasser, E., Schermer, M., Steinbacher, M., Tappeiner, U., 2013. Multiple ecosystem services of a changing Alpine landscape: past, present and future. Int. J. Biodiv. Sci. Ecosys. Serv. Manag. 9, 123-135.

Schnellnhuber, H.J., Cramer, W., Nakicenovic, N., Wigley, T. Yohe, G., 2006. Avoiding dangerous climate change. Cambridge, UK, Cambridge University Press.

Sorensen, P.O., Beller, H.R., Bill, M., Bouskill, N.J., Hubbard, S.S., Karaoz, U., Polussa, A., Steltzer, H., Wang, S., Williams, K.H., Wu, Y., Brodie, E.L., 2020. The Snowmelt Niche Differentiates Three Microbial Life Strategies That Influence Soil Nitrogen Availability During and After Winter. Front. Microbiol. 11:871.

Szukics, U., Grigulis, K., Legay, N., Kastl. E.M., Baxendale, C., Bardgett, R.D., Clément, J.C., Lavorel, S., Schloter, M., Bahn, M., 2019. Management versus site effects on the abundance of nitrifiers and denitrifiers in European mountain grasslands. Sci. Tot. Environ. 648, 745-753.

Tappeiner, U., Tasser, E., Leitinger, G., Cernusca, A., Tappeiner, G., 2008. Effects of Historical and Likely Future Scenarios of Land Use on Above- and Belowground Vegetation Carbon Stocks of an Alpine Valley. Ecosys. 11, 1383–1400

Vittoz, P., Randin, C., Dutoit, A., Bonnet, F., Hegg, O., 2009. Low impact of climate change on subalpine grasslands in the Swiss Northern Alps. Glob. Chang. Biol. 15, 209-220.

Wall, D.H., 2007. Global change tipping points: above- and below-ground biotic interactions in a low diversity ecosystem. Philosoph. Transact. Roy. Soc. B: Biologic. Sci. 362, 2291-2306.

Wertz, S., Degrange, V., Prosser, J.I., Poly, F., Commeaux, C., Guillaumaud, N., Le Roux, X., 2007. Decline of soil microbial diversity does not influence the resistance and resilience of key soil microbial functional groups following a model disturbance. Environ. Microb. 9, 2211-2219.

Wipf, S., Rixen, C., 2010. A review of snow manipulation experiments in Arctic and alpine

tundra ecosystems. Polar Res. 29, 95-109.

Yoshinari, T., Hynes, R., Knowles, R., 1977. Acetylene inhibition of nitrous oxide reduction

and measurement of denitrification and nitrogen fixation in soil. Soil Biol. Biochem.

9, 177-183.

Zouache, K.,   Rory, J.M.,   Failloux, A.B.,   Genevieve, L.G., Mavingui, P., 2012.

Chikungunya virus impacts the diversity of symbiotic bacteria in mosquito vector.

Mol. Ecol. 21: 9, 2297-2309.

Zoderer, B.M., Lupo Stanghellini, P.S., Tasser, E., Walde, J., Wieser, H., Tappeiner, U.,

2016. Exploring socio-cultural values of ecosystem service categories in the Central

Alps: the influence of sociodemographic factors and landscape type. Reg. Environ.

Chang. 16:2033–2044.

  Supplementary Figures

Fig. S1. a) Soil moisture in percentage of dry soil; **b)** SIR-MicroResp™ (ppmV CO_2 h^{-1} g^{-1});

c) Abundance of total bacterial community (number of 16S *rRNA* gene copies g^{-1} dry soil).

764 **Fig. S2. a)** Nitrification enzyme activities quantified in µgN-(NO₂+NO₃)/g dry soil/h for

765 different treatments at various sampling dates; **b)** Abundance of *nxrA* nitrite reductase

766 (number of *nxrA* gene copies g⁻¹ dry soil) for different treatments at various sampling dates; **c)**

767 Abundance of *NS* nitrite reductase (number of *NS* gene copies g⁻¹ dry soil) for different

768 treatments at various sampling dates.

Fig. S3. a) Denitrification enzyme activities determined in µg N-N$_2$O/g dry soil/h for different

treatments at various sampling dates; **b)** Abundance of *nirK* denitrifiers (number of *nirK* gene

copies g^{-1} dry soil); **c)** Abundance of *nirS* denitrifiers (number of *nirS* gene copies g^{-1} dry

soil); **d)** Abundance of *nosZ* denitrifiers (number of *nosZ* gene copies g^{-1} dry soil) for different

treatments at various sampling dates.