Early Spring Snowmelt and Summer Droughts Strongly Impair the Resilience of Key

Microbial Communities in Subalpine Grassland Ecosystems

Farhan Hafeez[1,2], Lionel Bernard[3], Franck Poly[1], Jean-Christophe Clément[4], Thomas

Pommier[1*]

[1]*Laboratoire d'Ecologie Microbienne LEM, INRA UMR 1418, CNRS UMR 5557, Université*

de Lyon, F69622 Villeurbanne Cedex, France

[2]*Department of Environmental Sciences, COMSATS University Islamabad, Abbottabad*

Campus, Tobe Camp University Road Abbottabad 22060 Pakistan.

[3]*Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LECA, 38000 Grenoble, France*

[4]*Univ. Savoie Mont Blanc, INRAE, CARRTEL, 74200 Thonon-les-Bains, France*

Corresponding author :

Thomas Pommier

Laboratoire d'Ecologie Microbienne LEM, INRA UMR 1418, CNRS UMR 5557, Université

de Lyon, F69622 Villeurbanne Cedex, France

Email: thomas.pommier@univ-lyon1.fr

ABSTRACT

Subalpine grassland ecosystems are important from biodiversity, agriculture, and touristic perspectives but their resilience to seasonally occurring climatic extremes is increasingly challenged with climate change, accelerating their vulnerability to tipping points. Microbial communities, which are central in ecosystem functioning, are usually considered as more resistant and highly resilient to such extreme events due to their functional redundancy and strong selection in residing habitats. To investigate this, we explored soil microbial responses upon recurrent summer droughts associated with early snowmelt in subalpine grasslands mesocosms set-up at the Lautaret Pass (French Alps). Potential respiration, nitrification and denitrification were monitored over a period of two growing seasons along with quantification of community gene abundances of total bacteria as well as (de)nitrifiers. Results revealed that droughts had a low and short-term adverse impact on bacterial total respiration supporting their hypothesized high resilience, i.e., resistance and ability to recover. Nitrification and abundances of the corresponding functional guilds showed relatively strong resistance to summer droughts but declined in response to early snowmelt. This triggered a cascading effect on denitrification but also on abundances of denitrifying communities which recovered from all climatic extremes except from the summer droughts where nitrifiers were collapsed. Denitrification and respective functional groups faced high impact of applied stresses with strong reduction in abundance and activity of this specialized community. Although, consequently lower microbial competition for nitrate may be positive for plant biomass production, warnings exist when considering the potential nitrogen leaching from these ecosystems as well as risks of greenhouses gases emission such as N_2O.

Key words: climate change, grasslands, (de)nitrification, weather extremes, snowmelt, N_2O

1. Introduction

Subalpine grassland ecosystems are important for biodiversity, agriculture, and a wide range of key services (Pecher et al., 2017; Zoderer et al., 2016; Tappeiner et al., 2008; Grêt-Regamey et al., 2012). These grasslands are under extreme pressure of climatic changes drastically affecting their functioning and disrupting the linked ecosystem services, *e.g.,* plants productivity, nutrient cycling etc. (Berauer et al., 2019; Bernard et al., 2019). Considering the continuous human perturbations and global changes (Berauer et al., 2019), it is direly required to investigate whether these grasslands are highly vulnerable, have developed resilience (comprising of resistance: ability to maintain the existing state, and recovery characteristics: to recover from disturbances) (Ingrisch and Bahn, 2018; Hodgson et al., 2015) or have attained an altered recovery state (Seabloom et al., 2020) over time which is crucial in conserving ecosystems functions (Donohue et al., 2016).

The multiple effects of global climatic changes not only impact biodiversity and ecosystem services (Berauer et al., 2019; Schirpke et al., 2013, Engler et al., 2011) but also their associated resilience including both resistance (Bernard et al., 2019; Wipf and Rixen, 2010) and recovery in Alpine grasslands (Ingrisch et al., 2020; Karlowsky et al., 2018). The effect of recurring climatic extremes on microbial functioning in these grasslands needs to be assessed as they may shift the fate of particular biogeochemical functions especially when resilience limits are crossed and tipping points (where a potentially irreversible change is about to occur or is already occurring *sensu*, Schellnhuber et al. 2006) are reached. Such estimations are crucial to understand and possibly protect the stability of ecosystems and inhabiting species (de Vries and Shade, 2013; Griffiths and Philippot, 2013), but also to forecast unforeseen consequences of adverse climatic extremes. Where weather extremes including droughts are predicted to be severe (Fuchslueger et al., 2019; Karlowsky et al., 2018), Alpine grassland ecosystems are expected to face less snow falls that ultimately lead to

subsequent droughts (Klein et al., 2016; Gobiet et al., 2014). However, their vulnerability to collapse or support a particular function towards the combined (cumulative) impact of different weather extremes is poorly documented.

Microbial communities are crucial actors in ecosystem functioning (Chen et al., 2020; Bardgett and van der Putten, 2014) and respond strongly to environmental variability (Piton et al., 2020). These are critical actors in estimating the functional stability (de Vries and Shade 2013; Griffiths et al., 2000), and they are highly diverse and functionally redundant across a range of environments. Consequently, microbial communities are usually expected to be highly resilient to climatic fluctuations as compared to many other environmental actors albeit the strong selection under local harsh climatic conditions (Capdeville et al., 2019; Griffiths and Philippot, 2013, Benot et al., 2014). Despite these yet known inherent capabilities, whether these microbial communities can withstand to correlated acceleration of weather extremes is largely unknown. Especially, assessment of the functional boundaries of the subalpine grasslands and linked ecosystems services is critical since they are susceptible to extreme events (Bernard et al., 2019).

The characteristics of existing plant species and nutrient availability as well as their interaction with microorganisms to ultimately respond to climatic changes are important in assessing soil functioning resilience (Bernard et al., 2020). For example, any shift in native plant community can influence the soil nutrient usage leading to selection in the pool of residing microbes hence the corresponding functions (Legay et al., 2016; de Vries et al., 2012). Similarly, weather extremes directly influencing plant community may also impact microbes (Bardgett et al., 2013; de Vries and Shade, 2013, Saccone et al., 2013). Earlier, summer droughts in subalpine grasslands were reported to increase leaf senescence while reducing plant productivity (Benot et al., 2014), as well as affecting N cycling-related microbial community and functions (Dai et al., 2020; Cantarel et al., 2012).

Here we determined the short-term effect of drought, snow removal, and their additive

(recurrent) effect on microbial activities and size of involved functional guilds and following

responses, to measure the resistance and recovery, respectively. Since microbes, driving broad

(*e.g.,* respiration, decomposition) and specialized soil processes (*e.g.* (de)nitrification), may

respond differentially to various perturbations (Dai et al., 2020; Jusselme et al., 2016; Chaer

et al., 2009; Wertz et al., 2007), we conducted our work both at broad and functional scales.

Investigating the response of these grasslands is crucial since the variable impact of natural

and manmade perturbations might have considerably influenced nutrient cycling including

that of N (Galloway et al., 2004), but also the involved microbial functional guilds. First, we

tested how total bacterial community (16S *rRNA*) and related substrate-induced respiration

(SIR) were influenced under droughts, snow removal and their cumulative impact and how

much they recovered from these stresses. Then we investigated nitrogen (N) cycling (that is

critical in ecosystem sustainability (Chen et al., 2020; Kuypers et al., 2018)) by determining

the abundances (nitrifiers (*nxrA*, *NS*), denitrifiers (*nirK*, *nirS*, *nosZ*)) and activities

(nitrification and denitrification enzyme activities -NEA and DEA, respectively) of involved

microbial communities, since they carry implications for plant productivity (loss of N - an

important plant nutrient) and environment (N_2O emissions-global warming) (Dai et al., 2020;

Galloway et al., 2004).

The current work was carried out on functionally contrasted and assembled grasslands

set up in a common garden at Lautaret Pass with various sampling campaigns during two

growing seasons. Based on an assumed functional redundancy (Jia and Whalen 2020;

Griffiths and Philippot 2013), we hypothesized that broad scale function i.e., SIR and relevant

microbial community should resist and/or recover once the stresses/extreme events are over.

SIR being representative of total microbial communities is expected to be influenced by soil

moisture and abundance of *16S rRNA* (Hallin et al., 2009). However, more specific functions

of N-cycling should be more sensitive because we expect comparatively less functional

redundancy and, despite being able to resist to single stress, the combined effects (SR+D)

may trigger a tipping point, disturbing various steps of these processes. Moreover, any shift in

soil moisture, substrate availability and abundance of denitrifiers can strongly impact DEA

through affecting micro-environment conditions (Philippot et al., 2007) both under drought

and no drought, while NEA might be sensitive to ammonium availability (Jia and Conrad

2009) and snow removal. To address these questions, we quantified the immediate responses

of microbial communities' abundances and processes to droughts and early snowmelt extreme

events to explore the resistance and the following recovery across subalpine grassland

ecosystems.

2. Materials and methods

2.1. Experimental site, Mesocosms and Treatments' implementation

The study was set up at Lautaret pass proximity in southeast France (2,100 m a.s.l,

45°02'13''N, 6°24'01''E) with a multifactorial experimental design where 128 mesocosms

(n=128), 50-cm width and 40-cm depth, were installed in a common garden beside subalpine

grasslands. Mesocosms were filled with previously sieved (5-mm) *in-situ* soil and planted

with varying relative abundances of four species ranging along a plant economics spectrum.

Species richness and functional composition remained stable throughout the whole

experimentation due to low plant mortality. No snow removal (NSR) and snow removal (SR)

treatments (n=64+64) were performed and for each of the NSR and SR there was an

additional (sub) climate change treatment, i.e., maintained average soil moisture (ND) or

drought (D) (n=32+32). Treatments receiving no snow removal with maintained average soil

moisture were named as control (CTRL). All the 128 Mesocosms were installed inside 8

blocks. Within each of the 8 blocks with or without snow removal, there were two sub-blocks

either maintained at long-term average soil moisture or subjected to summer drought. Within

each sub-block 8 mesocosms were installed and 4 were submitted to plant cuttings and fertilization (CF), while the other half remained untreated (NCF) (n=16+16). During spring 2014, slow-release fertilizer was applied as pellets in the CF mesocosms (150 kg N/ha). To simulate early snowmelt, snow cover was removed one month earlier than the average snowmelt period which ultimately simulated a shorter snow-covered period followed by an early spring drought. The drought treatment was implemented using rainout shelters during spring, summer and fall. These shelters were thus in use during the whole vegetation season and all mesocosms were watered manually to maintain long-term average soil moisture. This manual watering was discontinued to achieve drought that reduced soil moisture for more than one month in August - September 2013 and June - July 2014.

Fig. 1. Experimental timeline showing the implementation of various treatments as well as the sampling dates. R and Rc represent resistance and recovery respectively.

2.2. Soil sampling campaigns

Five sampling campaigns in total were carried out at different time intervals including June 2013 (T0), September 2013 (T1), May 2014 (T2), July 2014 (T3) and September 2014 (T4). 1st and 2nd D were applied in August - September 2013 and June - July 2014

respectively (Fig. 1). The SR treatment was applied before T2 sampling campaign. Resistance

to 1^{st} D of the microbial communities was measured at T1, resistance to 1^{st} SR and recovery

from 1^{st} D at T2, resistance to 2^{nd} D and recovery from 1^{st} SR at T3; and recovery from

overall D and SR treatments was evaluated at T4 sampling campaign (Fig. 1). No difference

at T0 observed since no treatment was applied, and control was already placed at each date -

T1 to T4, so we choose to compare only these on graph. During all sampling campaigns

composite soil samples (5 cores mixed together, 0-10 cm depth) were collected from all 128

mesocosms (128x5: n=640). Fresh soil samples were sieved at 2-mm, homogenized, and

brought to the laboratory to measure the enzyme activities in fresh soil while a sub-sample

was stored at -20°C for DNA extraction. Soil properties including soil moisture content,

SOM% (soil organic matter), soil pH, soil %C, soil %N, total dissolved nitrogen-TDN, and

the dissolved organic nitrogen-DON, soil NH_4^+, and soil NO_3^- were also measured in an

associated work with focus on the effects of local environmental changes on subalpine

grasslands functioning (Bernard 2017). Specifically, subsamples of 5 g fresh soil were dried at

70°C for 1 week to determine soil moisture content (in g.g-1 dw calculated as the 70°C dry

soil weight relative to the fresh mass), followed by 4 h at 550° C to determine SOM

(calculated as the 550°C soil weight relative to the 70°C dry mass). 10 mg soil subsample

were air dried, ground to powder and analyzed for soil total C and N concentrations using a

C/N elemental analyzer (FlashEA 1112, ThermoFisher Scientific, MA, USA). Soil pH was

measured in a 1:4 (air dry soil/distilled water) solution. Soil $N-NH_4^+$ and $N-NO_3^-$, N-TDN and

N-DON were determined in separated 10 mg fresh soil subsamples extracts (K_2SO_4, 0.5 M)

using an FS-IV colorimetric chain (OI-Analytical, College Station, TX, USA) (Jones and

Willett, 2006).

2.3. Soil DNA extractions and quantifications

DNA was extracted from soil stored at -20°C using the PowerSoil® DNA Isolation Kit

(Mo Bio Laboratories Inc. USA) and DNA concentration was determined by using the Quant-

iTTM PicoGreen dsDNA Assay Kit (Invitrogen, Carlsbad, CA, USA).

2.4. Microbial community abundances

The abundance of the total bacterial communities was estimated in all DNA extracts

using a *16S rRNA* primer-based quantitative PCR (qPCR) assays as described by Lane (1991).

Final reaction volume was 20-µl, with 300-nM of forward and reverse primers; 1 X

LightCycler DNA Master Mix SYBR® Green I (Roche Diagnostics); 1-ng of sample DNA or

10^8 –10^2 copies of the standard DNA pQuantAlb16S plasmid (Zouacheet al., 2012). The

abundances of the *NS, nxrA, nirS, nirK and nosZ* genes were estimated using the primers

developed previously (Attard et al., 2011; Henry et al., 2004; Henry et al., 2006; Kandeler et

al., 2006). The total volume of the reaction was made 20 µl for *nxrA* and *nirK* each, with

concentration of 500 nM for *nxrA* while 1µM for *nirK* for both forward as well as reverse

primers; 10 µL of the SYBR® Green I kit was used for the *nxrA* and *nirK*, and 0.4 µg of T4

protein (Qbiogene, Carlsbad, CA USA) was used for *nirK*. The total volume of the reaction

was made 25 µl for *NS, nirS* and *nosZ*, with concentration of 400 nM for *NS* while 1µM for

nirS and *nosZ* both for forward as well as reverse primers; 12.5 µL of the SYBR® Green I kit

was used for the *NS, nirS* and *nosZ*; for *nirS* the T4 protein was 0.5 µg while it was 0.4 µg for

nosZ. The curves for the standards for *NS, nxrA, nirS, nosZ* and *nirK* were acquired after the

quantitative PCR assays with regular dilutions of already known concentrations of the

plasmid DNA with corresponding genes (10^7 - 10^1 number of gene copies). Quantification of

all gene community abundances was conducted out on a lightcycler 480 (Roche Diagnostics).

The amplification conditions used for qPCR assays were: 10 min at 95°C for each 40 cycles

for *16S rRNA*, 45 for *nxrA, NS, nirK* and *nirS* and 40 for *nosZ* (after 6 cycles of touchdown);

95°C for 15 s for *16S rRNA*, *nirK*, *nirS*, *nosZ* and for 30 s for *nxrA* and NS; 63°C for 1 min

for *16S rRNA*, 55°C for 45 s for *nxrA* and 66°C for 30 s for *NS*, 30 s for *nirK* and *nirS*, 60°C

for 15 s for *nosZ*. 72°C for 30 s for 16S *rRNA*, *nirK*, *nirS* and *nosZ*, 45 s for *nxrA*, 1 min for

NS; and additional step of 80°C for 15 s *nirS* and *nosZ*. Then 95°C for 5 s for 16S *rRNA* and

nxrA, 1 s for *NS*, *nirK* and *nirS* and 15 s for *nosZ*; 65°C for 1 m for *16S rRNA* and *nxrA*, 68°C

for 20 s for *NS* and *nirK*, 60°C for 20 s for *nirS* and 15 s for *nosZ*. And a continuous increase

to 98°C for *16S rRNA*, *nxrA*, *NS*, *nirK* and *nirS* and 95°C for *nosZ* to determine the melting

point; and finally, 10 s at 40°C for *16S rRNA*, *nxrA*, *NS*, *nirK* and *nirS* while 30 s for *nosZ* for

cooling. At least two independent quantitative PCRs were done for every sample while the

average was used. Co-amplification of the standards and the dilution of samples were used to

control any inhibition in the quantification.

2.5. Microbial activities

Susbtrate Induced Respiration MicroResp. (SIR) was measured as CO_2 production

using MicroResp plates (Chapman et al., 2007). Soil was added to the wells of 96-well plates.

0.5-ml of a nutritive solution including glucose (1.2-mg of C-glucose g of dried soil) was

added then incubated at 28°C. CO_2 concentrations were measured using a TECAN

spectrophotometer after 6-hours of incubation and the slope of the linear regression was used

to estimate aerobic respiration as the CO_2 produced (ppmV CO_2 h^{-1} g^{-1}).

Nitrifying enzyme activity (NEA) was measured on fresh soil samples as described by

Patra et al. (2005). In brief, the fresh soil sample (equal to 3-g dry weight) was incubated with

3 ml solution of N-NH$_4$ (200 µg N-(NH$_4$)$_2$SO$_4$ g^{-1}) in the flask for 72-hours at 180-rpm and

28°C. The flask was sealed with parafilm® (Pechiney Plastic Packaging, Menasha, WI, USA)

to produce the aerobic conditions. With distilled water, the total volume of the suspension was

made up to 15-ml. During the incubation, at regular intervals of 5, 24, 48 and 72-hours soil

slurry was filtered with a pore size of 0.2-µm. The rates of production of NO_2/NO_3 were

measured by ionic chromatography (DX120. Dionex, Salt Lake City, UT, USA).

Denitrification enzyme activities (DEA) was measured by incubating the fresh soil

sample in a 150-ml airtight plasma-flask sealed by a rubber stopper for 4-hours at 28°C

followed by the N_2O measured each 30 minutes for 3-hours a gas chromatograph coupled to a

micro-catharometer detector (lGC-R3000; SRA instruments) using the method adapted from

Dassonville et al. (2011). Briefly, a substrate solution (2-ml) containing glucose (0.5-mg of C-

glucose g^{-1} of dried soil), glutamic acid (0.5-mg of C-glutamic acid g^{-1} of dried soil) and

potassium nitrate (50-µg of N-KNO_3 g^{-1} of dried soil) were added to the soil. A 90:10 He-

C_2H_2 atmosphere provided anaerobic conditions and inhibited N_2O-reductase activity using

the acetylene inhibition technique that blocks the last step in the denitrification pathway

resulting in the accumulation of N_2O (Yoshinari et al., 1977). The soil was incubated for 4-

hours at 28°C and after incubation the amount of N_2O was measured each 30 minutes for 3-

hours. The amount of N_2O produced was measured using a gas chromatograph coupled to a

micro-catharometer detector (lGC-R3000; SRA instruments).

2.6. Statistics and Structural Equation Model

The data obtained were analyzed using JMP8® (SAS Institute Inc., SAS Campus

Drive, NC, USA). First, we did t-test to know the distribution of data (which was mostly not

normally distributed except for SIR) and then ANOVA (for normally distributed data) and

Kruskal-Wallis pairwise test (suited for non-normally distributed data) were applied to know

which treatment is different from the other for studied parameters at various sampling times.

Similarly, Kruskal-Wallis pairwise comparison was carried out also for the data which

remained non-normally distributed even after a log transformation. In addition to linear

modeling, for block effect, and repeated measures considerations (to consider time component

or sampling from same place at different times), we used Cluster analyses approach using R

(the package of R for hierarchical cluster analysis is 'cluster' and the function is hclust and to

make dendogram we used function dendogram) to determine the effect of treatments within

each block (to investigate the block effect) at different sampling times (to investigate the

repeated measures) on each other. Structural Equation Model (SEM) was constructed using

AMOS (Amos Development Corporation, Crawfordville, Florida, USA) to link the

environmental and microbiological variables to investigate the relationship between these

parameters under different climatic extremes. The fitness of the model was assessed by using

the $\chi 2$ test and the model was considered as correct if the $\chi 2$ test produced a non-significant p-

value - $p > 0.05$. Since the factors controlling the microbial functioning under the drought

and/or snow removal treatments were changing upon these treatments, the hypothesized

models were changed to increase the level of significance and to fit the model. SEM were

constructed to understand the relationship and path coefficients among the various causal

environmental and microbial factors under D, SR, SR+D treatments. The choices for SEMs

were made considering the parameters showing strong correlations as obtained through

analyses of linear regression modeling. Initial SEMs were constructed for each sampling

interval at each period of time. Then the strongly correlated parameters were linked through

SEMs to understand the variable impact of treatments for different parameters hence

variability was also considered.

3. Results

3.1. Changes in Broad Scale Function and Community

Soil moisture in CTRL showed the significant effect of the D, SR and SR+D

treatments at all sampling times (Fig. 2a; Fig. S1). The resistance and recovery of the N-

related microbial activities both for nitrification and denitrification and the abundances of

corresponding genes were calculated as the relative change in D, SR, and SR+D treatments in

comparision to CTRL. SIR was found to be significantly (p=0.001) lowered upon both

droughts and was recorded up to 89% of CTRL (43.83 ppmV CO_2 h^{-1} g^{-1} ± 0.58) but

recovered ultimately (p=0.639). Although the interaction of SR+D treatments marginally

(p=0.0123) affected the SIR, it was not affected by SR alone (p=0.625). It was noteworthy

that the abundance of the total bacterial community (as expressed by 16S *rRNA* gene copies)

followed a similar trend as SIR and was significantly (p=0.0006) reduced up to 56.3% of

CTRL (4.87 x 10^8 copies per g of dry soil) with a complete recovery observed at T4 (Fig. 2;

Fig. S1).

Fig. 2. Relative change compared to CTRL for different treatments at various sampling dates
a) in soil moisture; **b)** in SIR-MicroResp™; **c)** in abundance of total bacterial community
(number of 16S *rRNA* gene copies g^{-1} dry soil); Light grey dotted lines show variations in
time of the variables in the snow removal (SR) treatment relative to CTRL; dark grey dotted
lines show variations of the Drought (D) treatment relative to CTRL; and the black dotted
lines show the variations of cumulative SR+D treatment relative to CTRL.

3.2. Response of N Cycling Processes to Droughts and Snowmelt

Nitrification (NEA) showed resistance to both droughts at T1 and T3 and thus remained independent of the change in moisture content. Whereas, there was a significant reduction in NEA (up to 76% of CTRL) as observed upon SR treatment at T2 (Fig. 3; Fig. S2). To assess linkages, multiple regression analyses were carried out for the parameters in addition to moisture contents and the findings showed that the NEA was positively correlated to NH_4, %N, TDN and *nxrA* (r = 0.56, 0.20, 0.31, 0.20; p < 0.005). On the contrary, DEA was significantly affected by both droughts and was found to be lowered up to 85% and 68% of CTRL upon first and second drought, respectively (Fig. 4; Fig. S3). Recovery from the first drought was observed for DEA at T2 except for the interaction of SR+D treatment. However, the second drought (at T3, Fig. 4) strongly (p=0.0001) impacted DEA which did not recover from SR. Hence, at T4 there was an incomplete recovery in denitrification acitivity as the DEA for SR+D mesocosms could not return to the initial levels (0.362 µg $N-N_2O$/g dry soil/h ± 0.022).

3.3. Impact on N Cycling Functional Guilds

Our results revealed that the abundances of N cycling functional guilds including nitrifiers and denitrifiers were strongly impacted by both D and SR climatic extremes (p=0.001). We observed that the *Nitrospira* and *Nitrobacter* (*nxrA* and *NS)* abundances were significantly (p=0.001) reduced upon applied stresses and could recover only for *nxrA* (Fig. 3). The denitrifiers were particularly impacted by the second drought, that strongly lowered their gene abundances. At T4, the community gene abundances of *nirS* and *nosZ* denitrifiers were completely recovered. However, *nirK* denitrifiers could not recover in agreement to the observations as recorded for DEA (Fig. 4; Fig. S3). The multiple correlation analyses as discussed in the next section showed the strong positive correlations of DEA with the %C,

%N, DON, SOM, soil moisture, NEA as well as with the abundances of denitrifying genes.

However, these correlations were found more prominent in the climate change treatments than

the CTRL especially for soil moisture. For example, the Spearman correlation coefficient

showed a non-significant correlation between soil moisture and *nirK* denitrifiers ($r = 0.04$; $p >$

0.652) only for the CTRL whereas it was positive in SR+D treatment ($r = 0.33$; $p < 0.005$).

3.4. Microbial Variables and Climatic Treatments, a Structural Equation Modeling

Fig. 5 represents how the various environmental (soil moisture, NH_4, NO_3 etc.) and

microbial parameters (gene abundances) influenced the broad scale and N-related functional

activities of microbial communities at different levels of drought and snow removal as well as

by their interaction. The initial SEM was constructed for each sampling interval to observe

how the various environmental and microbial factors influenced the broad scale (SIR) and N-

related (NEA, DEA) functional microbial activities over each period of time. Afterwards, we

separately analyzed the most correlated parameters from the results of the linear regression

modeling and these parameters were linked to fit the SEM for drought and snow removal

treatments but also for their interaction. The SEM revealed that the SIR was linked to soil

moisture and *16S rRNA* for various treatments at different sampling times except at T1 and T2

($\chi^2 = 14.35$, $p = 0.35$; Fig. 5). In contrast, NEA was never associated to soil moisture and was

found to be mainly linked with soil NH_4 concentration and community gene abundances of

the *NS* nitrifiers, except for the SR treatment. Interestingly, under droughts, major part of the

DEA variance (up to 55%) was explained by soil moisture with *nirS* as dominant denitrifiers

except at T3 when it was *nosZ*. It was also observed that the overall impact of snow removal

and drought was variable for different denitrifiers with severe impact on *nirK* which could not

recover even by the end of the experiment. No significant effects of plant functional

composition and fertilization on microbial parameters were recorded.

Fig. 3. a) Relative change compared to CTRL in **a)** nitrification enzyme activities (µgN-

(NO$_2$+NO$_3$)/g dry soil/h); **b)** in abundance of *nxrA* nitrite reductases (number of *nxrA* gene

copies g^{-1} dry soil); **c)** in abundance of *NS* nitrite reductases (number of *NS* gene copies g^{-1}

dry soil); Light grey dotted lines show variations in time of the variables in the snow removal

(SR) treatment relative to CTRL; dark grey dotted lines show variations of the Drought (D)

treatment relative to CTRL; and the black dotted lines show the variations of cumulative

SR+D treatment relative to CTRL.

Fig. 4. a) Relative change compared to CTRL in **a)** denitrification enzyme activities (μg N-N$_2$O/g dry soil/h); **b)** in abundance of *nirK* denitrifiers (number of *nirK* gene copies g^{-1} dry soil); **c)** in abundance of *nosZ* denitrifiers (number of *nosZ* gene copies g^{-1} dry soil); Light grey dotted lines show variations in time of the variables in the snow removal (SR) treatment relative to CTRL; dark grey dotted lines show variations of the Drought (D) treatment relative

to CTRL; and the black dotted lines show the variations of cumulative SR+D treatment

relative to CTRL.

4. Discussion

This work reported the resilience including the resistance and recovery capabilities at

subalpine grassland ecosystems. Investigating the impact of applied climate change treatments

was crucial since these ecosystems are under drastic influence of global climatic changes

which may ultimately disrupt their functioning and the linked ecosystem services (Berauer et

al., 2019; Bernard et al., 2019; Schirpke et al., 2013; Wipf and Rixen, 2010; Vittoz et al.,

2009). Our results demonstrated the impact of successive droughts and early snowmelt

climatic extremes on subalpine grassland ecosystem functioning including nitrification and

denitrification as model processes and such findings give valuable insights into overall N

cycling resilience (Chen et al., 2020; Dai et al., 2020).

The quantification of resistance and recovery by comparing the relative change in D,

SR, and SR+D treatments over time compared to control showed the strong impact of

different climate change treatments on microbial activities and the corresponding functional

groups. Cluster analyses approach used to determine the effect of treatments within each

block at different sampling times revealed five different clusters. In each cluster, specific

treatments grouped separately to show that effect of one specific treatment does not influence

any block. The further effect of outliers was analyzed to know the difference in descriptive

values after non-consideration of outliers. Overall, broad scale function – SIR and the size of

total bacterial community - showed low resistance but high recovery for all treatments. The

finding strengthened the fact that this function is carried out by a wide range of microbial

communities which generally exhibit a fast recovery (Ingrisch et al., 2020). SIR represents the

broad scale microbial activity hence disturbing a part of the community could not affect the

overall function on the long term. Yet, short term reduction in SIR upon droughts can be

associated to lower microbial activities induced by water deficit conditions (Bernard et al.,

2019; Bloor and Bardgett, 2012). Moreover, broad scale functions are reported to be less

sensitive to applied disturbance (Chaer et al., 2009). Like respiration, decomposition

processes are generally driven by broad scale communities and weak impact of climatic

stresses for these processes have been reported in a joint experiment (Bernard et al., 2019),

while high recovery of microbial biomass to climatic stress have been observed in other

mountainous grasslands (Piton et al., 2020). These observations are also in accordance with

our assumptions on functional redundancy concepts (Jia and Whalen, 2020; Griffiths and

Philippot, 2013).

Fig. 5. Structural equation models (SEM) bridging the effects of microbial and environmental

factors.   SEMs including the Substrate induced respiration MicroResp (SIR), nitrification

(NEA), denitrification (DEA), gene community abundances (16S *rRNA*, *nxrA*, *NS*, *nirK*, *nirS*,

nosZ) and environmental factors (soil moisture, NH₄, NO₃) in **a, b, c** are built using the data

from treatments; SR, D, SR+D, respectively. The values shown beside the arrows are the path

coefficients and correspond to the standardized coefficients calculated based on the analysis

of correlation matrices. Values only for the significant relations (that fit the model, $P > 0.05$)

are indicated in the SEMs.

415   The impacts of extreme climatic events including droughts on microbial N cycling are

less clear (Fuchslueger et al., 2019). Climate change treatments showed contrasting impacts

on N cycling processes with denitrification (community gene abundance and activity)

showing less resistance to drought (D) as compared to nitrification (both for community gene

abundances and activities). For example, at T3 where snow removal and droughts occurred in

the same mesocosms (SR+D), DEA was first strongly reduced followed by an incomplete

recovery. Such observations were in contrast to NEA which was affected by snow removal

(SR) but recovered completely at T4. Our findings confirmed a strong influence of droughts

and associated soil moisture deficits specifically for DEA (Di et al., 2020; Philippot et al.,

2009). This adverse influence of applied stresses was pronounced when both climatic stresses

(SR+D) occurred indicating the additive effect of SR with declined nitrification upon earlier

snowmelt. Interestingly, the observations were both stress and community specific, with

variable response for microbial community size and functions. The collapse in microbial pool

size upon snowmelt was observed while studying the niche differentiation perspectives in

soils (Sorensen et al., 2020). In nearby subalpine grasslands, a strong turnover by nitrifying

and denitrifying microbial communities was previously shown upon natural variations in

snowpack depth (Jusselme et al., 2016). In wake of climatic extremes and considering the

functional stability scenarios, such multiple cascading effects are important to consider not

only from microbial community perspectives but also for related ecosystem services. For

example, these cascading effects can be crucial for N-related microbial activities which can

adversely impact greenhouse gas emissions (i.e., N_2O in particular), soil fertility and plant

productivity, hence ask for adaptive management strategies to counter the unforeseen

consequences (Griffis et al., 2014; Di et al., 2014; Bouwman et al., 2009).

The reduction in gene abundances of nitrifiers and denitrifiers upon drought (D) or

snow removal (SR) delineated the strong impact of climate change treatments in studied

grasslands. Denitrifiers are usually reported as more tolerant to droughts (Hammerl et al.,

2019) than nitrifiers (Austin and Strauss, 2010). Here, in line with DEA, denitrifiers'

abundances were also negatively impacted by droughts, especially after the second one. Like

DEA, *nirK* denitrifiers' abundances were severely lowered and could not show a complete

recovery at T4. DEA seems to be related to various denitrifying communities including *nirK*.

In agreement, DEA and abundance of *nirK* denitrifiers have been found to be correlated upon

variable snow depths under similar subalpine grasslands (Jusselme et al., 2016). Conversely,

Wertz et al. (2007) found that the abundances of denitrifiers were less affected by heat stress

in an agricultural soil, and this was associated to high functional redundancy.

Soil abiotic properties are crucial in estimating soil N cycling processes, and resilience

of microbial communities upon applied stresses may be influenced by inherent soil edaphic

characteristics (Berard et al., 2015). In our study, although we found strong positive

correlations of DEA with %C, %N, DON, SOM, soil moisture, NEA, and denitrifiers'

abundances, interestingly these factors shifted during the studied time. For example, when

grasslands were exposed to combined snow removal and drought (SR+D), soil moisture along

with *nirK*, *nirS* and *nosZ* were correlated to DEA ($r = 0.33, 0.15, 0.32, 0.38; p < 0.005$). While

soil moisture and *nirK* were not correlated to DEA ($r = 0.04; p > 0.64$) in the CTRL. These

observations showed the variable dominance of different denitrifiers in D, SR and CTRL

treatments. Similarly, DEA has been associated differently to these variables across a range of

environments (Chen et al., 2020; Chen et al., 2019; Petersen et al., 2012; Chroňáková et al.,

2009, Dong et al., 2009). At T4, DEA could not recover demonstrating a severe impact

caused both by drought (D) and the additive effect with snow removal (SR+D). Moreover, the

impact on nitrification during SR significantly passed on to DEA which collapsed upon the

second drought showing the cascading effect for SR+D.

These pronounced impacts on DEA and corresponding denitrifiers' abundances were

also explained through SEM (Fig.5). Possibly the system, (as described by Schnellnhuber et

al. (2006)) upon disturbance was 'pulsed' (respond on alteration but return) and 'pressed'

(respond on sustained alteration so shift) where 'press' brings the ecosystems to a tipping

point (as observed for DEA in our study). This may shift the system to an irreversible new

state (Wall, 2007), or at least may lead to altered recovery patterns (Seabloom et al., 2020).

Furthermore, such changes in denitrifiers' relative abundances may also result in selection of

certain stable/unstable groups. Such extreme climatic events may consequently impact

(increase) N losses through NO_3 leaching or N_2O emissions (de Vries and Shade 2013). In

addition, studies in other ecosystems have suggested that these functional guilds may have

contrasting resilience abilities against various disturbances (Capdeville et al., 2019).

Though SIR was related to soil moisture and *16S rRNA* when SEM was fitted, NEA

was highly linked to both soil NH_4^+ and *NS* nitrifiers' abundance hence showed a weak effect

of soil moisture on NEA. Therefore, *NS* communities might have played the role as dominant

nitrifiers in these grasslands. Higher abundance of *NS* nitrifiers than *nxrA* under drought as

well as the dominant role of *NS* nitrifiers in these soils was earlier shown in other soils with

low activity (Attard et al., 2010). However, here DEA was related both to soil moisture and

abundances of *nirS* denitrifiers throughout the experiment except at T3 when it was *nosZ*,

suggesting community selection or adaptation is an important phenomenon under changing

climate or stress conditions. After the second drought at T3, a cascading effect of snow

removal (SR) might have shifted the community dynamics from *nirS* to *nosZ* with better

adaptation of the *nosZ* denitrifiers. The denitrifiers responded differentially, *e.g.,* in contrast to

nirK; *nirS* and *nosZ* abundances recovered, showing a community specific impact (Karlowsky

et al., 2018) of drought and snow removal. The *nirK* denitrifiers were reported as less

abundant (Peterson et al., 2012), and relatively more sensitive to environmental change than

nirS denitrifiers in subalpine and agricultural soils (Szukics et al., 2019; Hallin et al., 2009).

Though the findings implied by Dai et al. (2020) contrasted with these observations,

seasonally driven drought effects have been reported for *nirK*, *nirS* and *nosZ* abundances in

grassland ecosystems (Hammerl et al., 2019). Moreover, the denitrifiers harboring *nirK* genes

might not contain *nosZ* genes, in agreement to the discovery that not all denitrifiers contain

nosZ (Hallin et al., 2018), which may ultimately disturb the overall nitrous oxide budget from

studied soils.

We did not find any significant effect of plant functional composition and fertilization on the

investigated soil microbial parameters, suggesting they were not the main drivers of the

studied processes in our experimental set-up. Consequently, the trait-centered influence of

abiotic variables on broad and N-related microbial abundances and activities, in response to

climatic stresses, was not observed here but, instead, their direct effect (Griffin-Nolan et al.,

2019). Yet, leaf or root specific traits, rather than plant community composition, might have

affected the nutrient availability and ultimately associated microbial processes. This could

explain why we found that soil nutrient contents along with community abundances were

driving the N cycling processes in our study. Interestingly in similar grasslands, no direct

effect of fertilization was observed but only specific plant traits were affecting the same

microbial activities (Legay et al., 2016). This suggests a competition for nutrients between

plants and microbes, as observed following snowmelt in nearby subalpine grasslands (Legay

et al. 2013), resulting here in a non-significant fertilization effect on microbial parameters

during the study period. However, such management strategies including fertilization

practices may favor certain microbial communities ultimately causing fast resilience to stress

conditions (Piton et al., 2020). It is noteworthy that due to the climatic extremes, the

responses showed by the various nitrifiers' and denitrifiers' groups were different to CTRL

indicating an unanticipated internal competition for initially available nutrients. Importance of

such competitions will be crucial to further investigate, especially in ecosystems facing

climatic extremes, since these are suggested to be critical in controlling ecosystem N cycling

(Legay et al., 2013; Grigulis et al., 2013, Le Roux et al., 2013).

5. Conclusions

This work clearly demonstrated a strong and unprecedented impact of climate

extremes such as early snowmelt and recurrent droughts, alone or in combination, on soil

microbial communities both at broad (i.e., soil respiration) and N-related (nitrifiers' and

denitrifiers' activities and abundances) scales. Droughts showed a low and short-term impact

on bacterial total respiration supporting their hypothesized high resilience (resistance and

ability to recover). Though nitrification activity and community gene abundances of

corresponding functional guilds exhibited relatively strong resistance to summer droughts,

they declined in response to early snowmelt leaving strong impact on denitrification. Our

findings inferred that predicted climatic extremes may carry severe consequences for soil

microbial community functioning in subalpine grasslands ultimately resulting in shifts in N

availability, plant community diversity, and associated ecosystem services.

Conflict of Interest

The authors of this article declare that they have no financial conflict of interest with the

content of this article.

Acknowledgements

This work was part of the ERA-Net BiodivERsA project REGARDS (ANR-12-EBID-004-01). Quantitative PCRs were carried out at the platform DTAMB (IFR 41, Université Lyon 1) and N cycling activities at the platform AME (UMR5557-UMR1418). We want to thank all the people who helped us in the field or in the lab: C. Arnoldi, A. Foulquier, N. Guillaumaud. This research was conducted at 'Jardin du Lautaret' (UMS 3370 Univ. Grenoble Alpes – CNRS), a member of the DIPEE Grenoble-Chambery, the FREE-Alpes Federation (FR n°2001-CNRS), and of the AnaEE-France (ANR-11-INBS- 0001AnaEE-Services). This research was conducted within the Long-Term Socio-Ecological Research (LTSER) platform Zone Atelier Alpes, a member of the eLTER network.

References

Attard, E., Recous, S., Chabbi, A., Berranger, C.D., Guillaumaud, N., Labreuche, J., Philippot, L., Schimid, B., Le Roux, X., 2011. Soil environmental conditions rather than denitrifier abundance and diversity drive potential denitrification after changes in land uses. Glob. Chang. Biol. 17, 1975-1989.

Attard, E., Poly F., Commeaux, C., Laurent, F., Terada, A., Smets, B.F., Recous, S., Le Roux, X., 2010. Shifts between Nitrospira□ and Nitrobacter□like nitrite oxidizers underlie the response of soil potential nitrite oxidation to changes in tillage practices. Environ. Microbiol. 12(2), 315-26.

Austin, B.J., Strauss,, E.A., 2011. Nitrification and denitrification response to varying periods of desiccation and inundation in a western Kansas stream. Hydrobiol. 658, 183-195.

Bardgett, R.D., Manning, P., Morriën, E., De Vries, F.T., 2013. Hierarchical responses of plant–soil interactions to climate change: consequences for the global carbon cycle. J. Ecol. 101, 334-343.

Bardgett, R.D., van der Putten, W.H., 2014. Belowground biodiversity and ecosystem

functioning. Nat. 515, 505–511.

Benot M.L., Saccone, P., Pautrat, E., Vicente, R., Colace, M.P., Karl, G., Clément, J.C.,

Lavorel, S., 2014. Stronger Short-Term Effects of Mowing Than Extreme Summer

Weather on a Subalpine Grassland. Ecosys. 17, 458-472.

Berauer, B.J., Wilfahrt, P.A., Arfin-Khan, M.A.S., Eibes, P., Heßberg, A.V., Ingrisch, J.,

Schloter, M., Schuchardt, M.A., Jentsch, A., 2019. Low resistance of montane and

alpine grasslands to abrupt changes in temperature and precipitation regimes. Arct.

Antarc. Alp. Res. 51:1, 215-231.

Bernard, L., Foulquier, A., Gallet, C., Lavorel, S., Clément, J.C., 2019. Effects of snow pack

reduction and drought on litter decomposition in subalpine grassland communities.

Plant Soil. 435, 225–238.

Bernard, L., Decau, M.L., Morvan☐Bertrand, A., Lavorel, S., Clément, J.C., 2020.

Water☐soluble carbohydrates in Patzkea paniculata (L.): a plant strategy to tolerate

snowpack reduction and spring drought in subalpine grasslands. Plant Biol. 22, 441-

449.

Bernard L., 2017. Mécanismes fonctionnels de résilience des prairies subalpines au

changement globale. PhD thesis, Université Grenoble-Alpes.

Berard, A, Ben Sassi, M, Kaisermann, A, Renault, P., 2015. Soil microbial community

responses to heat wave components: drought and high temperature. Clim. Res.

66:243–264

Bloor, J.M.G., Bardgett, R.D., 2012. Stability of above-ground and below-ground processes

to extreme drought in model grassland ecosystems: interactions with plant species

diversity and soil nitrogen availability. Perspect. Plant Ecol. Evol. Syst. 14: 193–204

Bouwman, A.F., Beusen, A.H.W., Billen, G., 2009. Human alteration of the global nitrogen

and phosphorus soil balances for the period 1970–2050. Glob. Biogeochem. Cyc. 23,

GB0A04.

Cantarel, A., Bloor, J.M.G., Pommier, T., Guillaumaud, N., Moirot, C., Soussana, J.F., Poly,

F., 2012. Four years of experimental climate change modifies the microbial drivers of

N_2O fluxes in an upland grassland ecosystem. Glob. Chang. Biol. 18, 2520-2531.

Capdeville, C., Pommier, T., Gervaix, J., Fromard, F., Rols, J.L., Leflaive, J., 2019.

Mangrove Facies Drives Resistance and Resilience of Sediment Microbes Exposed to

Anthropic Disturbance. Front. Microbiol. 9:3337

Chaer, G., Fernandes, M., Myrold, D., Bottomley, P., 2009. Comparative resistance and

resilience of soil microbial communities and enzyme activities in adjacent native

forest and agricultural soils. Microb. Ecol. 58, 414-424.

Chapman, S.J., Campbell, C.D., Artz, R.R.E., 2007. Assessing CLPPs using MicroResp™. J.

Soils Sedim. 7, 406-410.

Chen, Q.L., Ding, J., Li, C.Y., Yan, Z.Z., He, J.Z., Hu, H.W., 2020. Microbial functional

attributes, rather than taxonomic attributes, drive top soil respiration, nitrification and

denitrification processes, Sci. Tot. Environ. 734,139479.

Chen, J., Li, J., Shen, W., Xu, H., Li, Y., Luo, T., 2019. The Structure and Species Co-

Occurrence Networks of Soil Denitrifying Bacterial Communities Differ Between A

Coniferous and A Broadleaved Forests. Microorg. 7(9), 361.

Chroňáková, A., Radl, V., Čuhel, J., Šimek, M., Elhottová, D., Engel, M., Schloter, M., 2009.

Overwintering management on upland pasture causes shifts in an abundance of

denitrifying microbial communities, their activity and N2O-reducing ability. Soil Biol.

Biochem. 41, 1132-1138.

Dai, Z., Yu, M., Chen, H., Zhao, H., Huang, Y., Su, W., Xia, F., Chang, S.X., Brookes, P.C.,

Dahlgren, R.A., 2020. Elevated temperature shifts soil N cycling from microbial

immobilization to enhanced mineralization, nitrification and denitrification across

global terrestrial ecosystems. Glob. Chang. Biol. 26, 5267-5276.

Dassonville, N., Guillaumaud, N., Piola, F., Meerts, P., Poly, F., 2011. Niche construction by

the invasive Asian knotweeds (species complex Fallopia): impact on activity,

abundance and community structure of denitrifiers and nitrifiers. Biolog. Invas. 13,

1115-1133.

de Vries, F.T., Liiri, M.E., Bjørnlund, L., Setälä H.M., Christensen S., Bardgett R.D., 2012.

Legacy effects of drought on plant growth and the soil food web. Oecolog. 170, 821–

833.

De Vries, F.T., Shade, A., 2013. Controls on soil microbial community stability under climate

change. Front. Microbiol. 4, 265.

Di, H.J., Cameron, K.C., Podolyan, A., Robinson, A., 2014. Effect of soil moisture status and

a nitrification inhibitor, dicyandiamide, on ammonia oxidizer and denitrifier growth

and nitrous oxide emissions in a grassland soil. Soil Biol. Biochem. 73, 59–68

Dong, L.F., Smith, C.J., Papaspyrou, S., Stott A., Osborn A.M., Nedwell D.B., 2009. Changes

in Benthic Denitrification, Nitrate Ammonification, and Anammox Process Rates and

Nitrate and Nitrite Reductase Gene Abundances along an Estuarine Nutrient Gradient

(the Colne Estuary, United Kingdom). App. Environ. Microbiol. 75, 3171-3179.

Donohue, I., Hillebrand, H., Montoya, J.M., Petchey, O.L., Pimm, S.L., Fowler, M.S., Yang,

Q., Jackson, A.L., Lurgi, M., McClean, D., O'Connor, N.E., O'Gorman, E.J., Yang,

Q., 2016. Navigating the complexity of ecological stability. Ecol. Lett. 19:1172–85

Engler, R., Randin, C.F., Thuiller, W., et al., 2011. 21st century climate change threatens

mountain flora unequally across Europe. Glob. Chang. Biol. 17:2330–2341.

Fuchslueger, L., Wild, B., Mooshammer, M., Takriti, M., Kienzl, S., Knoltsch, A., Hofhansl,

F., Bahn, M., Richter, A., 2019. Microbial carbon and nitrogen cycling responses to

drought and temperature in differently managed mountain grasslands. Soil Biol.

Biochem. 135, 144–153.

Galloway, J.N., Dentener, F.J., Capone, D.G., Boyer,E.W., Howarth, R.W., Seitzinger,S.P.,

Asner, G.P., Cleveland,C.C., Green, P.A., Holland, E.A., Karl, D.M., Michaels, A.F.,

Porter, J.H., Townsend, A.R., Vöosmarty, C.J., 2004. Nitrogen cycles: past, present,

and

future. Biochem. 70, 153–226.

Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., Stoffel, M., 2014. 21st

century climate change in the European Alps—a review. Sci. Tot. Environ. 493:1138–

1151.

Grêt-Regamey, A., Brunner, S.H. Kienast, F., 2012. Mountain Ecosystem Services: Who

Cares? Mount. Res. Develop. 32, S23–S34

Griffin-Nolan, R.J., Blumenthal, D.M., Collins, S.L., Farkas, T.E., Hofman, A.M., Mueller,

K.E., Ocheltree, T.W., Smith, M.D., Whitney, K.D., Knapp, A.K., 2019. Shifts in

plant functional composition following longterm drought in grasslands. J. Ecol.

107:2133–2148.

Griffis, T.J., Chen, Z., Baker, J.M., Wood, J.D., Millet, D.B., Lee, X., Venterea, R.T., Turner,

P.A., 2014. Nitrous oxide emissions enhanced in a warmer world. PINAS. 114,

12081-12085

Griffiths, B.S., Philippot, L., 2013. Insights into the resistance and resilience of the soil

microbial community. FEMS Microbiol. Rev. 37, 112-129.

Griffiths, B.S., Ritz, K., Bardgett, R.D., Cook, R., Christensen, S., Ekelund, F., Sørensen,

S.J., Bååth. E., Bloem, J., De Ruiter, P.C., Dolfing, J., Nicolardot, B., 2000.

Ecosystem response of pasture soil communities to fumigation-induced microbial

diversity reductions: an examination of the biodiversity–ecosystem function

relationship. Oikos. 90, 279-294.

Grigulis, K., Lavorel, S., Krainer, U., Legay, N., Baxendale, C., Dumont, M., Kastl, E.,

Arnoldi, C., Bardgett, R.D., Poly, F., Pommier, T., Schloter, M., Tappeiner, U., Bahn,

M., Clément, J.C., 2013. Relative contributions of plant traits and soil microbial

properties to mountain grassland ecosystem services. J. Ecol. 101, 47-57.

Hallin, S., Philippot, L., Löffler, F.E., Sanford, R.A., Jones, C.M., 2018. Genomics and

ecology of novel N_2O-reducing microorganisms. Trend Microbiol. 26, 43–55

Hallin, S., Jones, C.M., Schloter, M., Philippot, L., 2009. Relationship between N-cycling

communities and ecosystem functioning in a 50-year-old fertilization experiment. Int.

Soc. Microb. Ecol. 3, 597-605.

Hammerl, V., Kastl, E., Schloter, M., Kublik, S., Schmidt, H., Welzl, G., Jentsch, A.,

Beierkuhnlein, C., Gschwendtner, S., 2019. Influence of rewetting on microbial

communities involved in nitrification and denitrification in a grassland soil after a

prolonged drought period. Sci. Rep. 9, 2280

Henry, S., Baudoin, E., Lopez-Gutierrez, J.C., Martin-Laurent, F., Brauman, A., Philippot, L.,

2004. Quantification of denitrifying bacteria in soils by nirK gene targeted real-time

PCR. J. Microbiol. Method. 59, 327-335.

Henry, S., Bru, D., Stres, B., Hallet, S., Philippot, L., 2006. Quantitative detection of the nosZ

gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S

rRNA, narG, nirK, and nosZ genes in soils. App. Environ. Microbiol. 72, 5181-5189.

Hodgson, D., McDonald, J.L., Hosken, D.J., 2015. What do you mean, "resilient"? Trend.

Ecol. Evol. 30: 9, 503-506.

Ingrisch, J., Karlowsky, S., Anadon-Rosell, A., Hasibeder, R., Konig, A., Augusti, A.,

Gleixner, G., Bahn, M., 2018. Land Use Alters the Drought Responses of Productivity

and CO2 Fluxes in Mountain Grassland. Ecosys. 21, 689–703.

Ingrisch, J., Karlowsky, S., Hasibeder, R., Gleixner, G., Bahn, M., 2020. Drought

andrecovery effects on belowground respiration dynamics and the partitioning

ofrecent carbon in managed and abandoned grassland. Glob. Chang. Biol. 26, 4366-

4378.

Jia, Z., Conrad, R., 2009. Bacteria rather than Archaea dominate microbial ammonia

oxidation in an agricultural soil. Environ. Microb. 11, 1658-1671.

Jia, Y., Whalen, J.K., 2020. A new perspective on functional redundancy and

phylogeneticniche conservatism in soil microbial communities. Pedosph. 30, 18–24

Jones, D.L., Willett, V.B., 2006. Experimental evaluation of methods to quantify dissolved

organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. Soil Biol.

Biochem. 38, 991-999.

Jusselme, M.D., Saccone, P., Zinger, L., Faure, M., Le Roux, X., Guillaumaud, N., Bernard,

L., Clement, J.C., Poly, F., 2016. Variations in snow depth modify N-related soil

microbial abundances and functioning during winter in subalpine grassland. Soil Biol.

Biochem. 92, 27–37.

Kandeler, E., Deiglmayr, K., Tscherko, D., Bru, D., Philippot, L., 2006. bundance of narG,

nirS, nirK, and nosZ Genes of Denitrifying Bacteria during Primary Successions of a

Glacier Foreland. App. Env. Microbiol. 5957-5962.

Karlowsky, S., Augusti, A., Ingrisch, J., Akanda, M.K.U., Bahn, M., Gleixner, G., 2018.

Drought-Induced Accumulation of Root Exudates Supports Post-drought Recovery of

Microbes in Mountain Grassland. Front. Plant Sci. 2018;9:1593.

Klein, G., Vitasse, Y., Rixen, C., Marty, C., Rebetez, M., 2016. Shorter snow cover duration

since 1970 in the Swiss Alps due to earlier snowmelt more than to later snow onset.

Climat. Chang. 139:637–49.

Kuypers, M.M., Marchant, H.K., Kartal, B., 2018. The microbial nitrogen-cycling network

Marcel. Nat. Rev. Microbiol. 16, 263–276

Lane, D.J., 1991. 16S/23S rRNA Sequencing. In: Nucleic acid Techniques in Bacterial

Systematics (eds Stackebrandt E, Goodfellow M), pp. 115–175. John Wiley & Sons,

New York City, New York.

Le Roux, X., Schmid, B., Poly, F., Barnard, R.L., Niklaus, P.A., Guillaumaud, N., Habekost,

M., Oelmann, Y., Philippot, L., Salles, J.F., Schloter, M., Steinbeiss, S., Weigelt, A.,

2013. Soil environmental conditions and microbial build-up mediate the effect of plant

diversity on soil nitrifying and denitrifying enzyme activities in temperate grasslands.

PLoS One. 8, e61069.

Legay, N., Grassein, F., Binet, M.N., Arnoldi, C., Personeni E., Perigon, S., Poly, F.,

Pommier, T., Puissant, J., Clément, J.C., Lavorel, S., Mouhamadou, B., 2016. Plant

species identities and fertilization influence on arbuscular mycorrhizal fungal

colonisation and soil bacterial activities. App. Soil Ecol. 98, 132-139.

Legay, N., Grassein, F., Robson, T.M., Personeni, E., Bataillé, M.P., Lavorel, S., Clément,

J.C., 2013. Comparison of inorganic nitrogen uptake dynamics following snowmelt

and at peak biomass in subalpine grasslands. Biogeosci. 10, 7631-7645.

Patra, A.K., Abbadie, L., Clays-Josserand, A., Degrange, V., Grayston, S.J., Loiseau, P.,

Louault, F., Mahmood, S., Nazaret, S., Philippot, L., Poly, F., Prosser, J.I., Richaume,

A., Le Roux, X., 2005. Effects of grazing on microbial functional groups involved in

soil n dynamics. Ecologic. Monograph. 75, 65-80.

Pecher, C., Bacher, M., Tasser, E., Tappeiner, U., 2017. Agricultural landscapes between

intensification and abandonment: the expectations of the public in a Central-Alpine

cross-border region. Landscap. Res. 1-15.

Petersen, D.G., Blazewicz, S.J., Firestone, M., Herman, D.J., Turetsky, M., Waldrop, M.,

2012. Abundance of microbial genes associated with nitrogen cycling as indices of

biogeochemical process rates across a vegetation gradient in Alaska. Environ. Microb.

14, 993-1008.

Philippot, L., Cuhel, J., Saby, N.P., Chèneby, D., Chronáková, A., Bru, D., Arrouays, D.,

Martin-Laurent, F., Simek, M., 2009. Mapping field-scale spatial patterns of size and

activity of the denitrifier community. Environ. Microb. 11, 1518-1526.

Philippot, L., Hallin, S., Schloter, M., 2007. Ecology of Denitrifying Prokaryotes in

Agricultural Soil. In: Advanc. Agron. pp Page., Academic Press.

Piton, G, Legay, N, Arnoldi, C, Lavorel, S, Clément, J-C, Foulquier, A., 2020. Using proxies

of microbial community☐weighted means traits to explain the cascading effect of

management intensity, soil and plant traits on ecosystem resilience in mountain

grasslands. J. Ecol. 108: 876– 893.

Saccone, P., Morin, S., Baptist, F., Bonneville, J.M., Colace, M.P., Domine, F., Faure, M.,

Geremia, R., Lochet, J., Poly, F., Lavorel, S. Clément, J.C., 2013. The effects of

snowpack properties on litter decomposition during winter in subalpine meadows.

Plant Soil. 363:215-229.

Seabloom, E.W., Borer, E.T. and Tilman, D., 2020. Grassland ecosystem recovery after soil

disturbance depends on nutrient supply rate. Ecol. Lett. 23(12) 1756-1765.

Schirpke, U., Leitinger, G., Tasser, E., Schermer, M., Steinbacher, M., Tappeiner, U., 2013.

Multiple ecosystem services of a changing Alpine landscape: past, present and future.

Int. J. Biodiv. Sci. Ecosys. Serv. Manag. 9, 123-135.

Schnellnhuber, H.J., Cramer, W., Nakicenovic, N., Wigley, T. Yohe, G., 2006. Avoiding

dangerous climate change. Cambridge, UK, Cambridge University Press.

Sorensen, P.O., Beller, H.R., Bill, M., Bouskill, N.J., Hubbard, S.S., Karaoz, U., Polussa, A.,

Steltzer, H., Wang, S., Williams, K.H., Wu, Y., Brodie, E.L., 2020. The Snowmelt

Niche Differentiates Three Microbial Life Strategies That Influence Soil Nitrogen

Availability During and After Winter. Front. Microbiol. 11:871.

Szukics, U., Grigulis, K., Legay, N., Kastl. E.M., Baxendale, C., Bardgett, R.D., Clément,

J.C., Lavorel, S., Schloter, M., Bahn, M., 2019. Management versus site effects on the

abundance of nitrifiers and denitrifiers in European mountain grasslands. Sci. Tot.

Environ. 648, 745-753.

Tappeiner, U., Tasser, E., Leitinger, G., Cernusca, A., Tappeiner, G., 2008. Effects of

Historical and Likely Future Scenarios of Land Use on Above- and Belowground

Vegetation Carbon Stocks of an Alpine Valley. Ecosys. 11, 1383–1400

Vittoz, P., Randin, C., Dutoit, A., Bonnet, F., Hegg, O., 2009. Low impact of climate change

on subalpine grasslands in the Swiss Northern Alps. Glob. Chang. Biol. 15, 209-220.

Wall, D.H., 2007. Global change tipping points: above- and below-ground biotic interactions

in a low diversity ecosystem. Philosoph. Transact. Roy. Soc. B: Biologic. Sci. 362,

2291-2306.

Wertz, S., Degrange, V., Prosser, J.I., Poly, F., Commeaux, C., Guillaumaud, N., Le Roux,

X., 2007. Decline of soil microbial diversity does not influence the resistance and

resilience of key soil microbial functional groups following a model disturbance.

Environ. Microb. 9, 2211-2219.

Wipf, S., Rixen, C., 2010. A review of snow manipulation experiments in Arctic and alpine

tundra ecosystems. Polar Res. 29, 95-109.

Yoshinari, T., Hynes, R., Knowles, R., 1977. Acetylene inhibition of nitrous oxide reduction

and measurement of denitrification and nitrogen fixation in soil. Soil Biol. Biochem.

9, 177-183.

Zouache, K., Rory, J.M., Failloux, A.B., Genevieve, L.G., Mavingui, P., 2012.

Chikungunya virus impacts the diversity of symbiotic bacteria in mosquito vector.

Mol. Ecol. 21: 9, 2297-2309.

Zoderer, B.M., Lupo Stanghellini, P.S., Tasser, E., Walde, J., Wieser, H., Tappeiner, U.,

2016. Exploring socio-cultural values of ecosystem service categories in the Central

Alps: the influence of sociodemographic factors and landscape type. Reg. Environ.

Chang. 16:2033–2044.

800   **Supplementary Figures**

801
802

804   **Fig. S1. a)** Soil moisture in percentage of dry soil; **b)** SIR-MicroResp™ (ppmV CO_2 h^{-1} g^{-1});

805   **c)** Abundance of total bacterial community (number of 16S *rRNA* gene copies g^{-1} dry soil).

806
807

809 **Fig. S2. a)** Nitrification enzyme activities quantified in μgN-(NO_2+NO_3)/g dry soil/h for

810 different treatments at various sampling dates; **b)** Abundance of *nxrA* nitrite reductase

811 (number of *nxrA* gene copies g^{-1} dry soil) for different treatments at various sampling dates; **c)**

812 Abundance of *NS* nitrite reductase (number of *NS* gene copies g^{-1} dry soil) for different

813 treatments at various sampling dates.

819 **Fig. S3. a)** Denitrification enzyme activities determined in µg N-N$_2$O/g dry soil/h for different

820 treatments at various sampling dates; **b)** Abundance of *nirK* denitrifiers (number of *nirK* gene

821 copies g^{-1} dry soil); **c)** Abundance of *nirS* denitrifiers (number of *nirS* gene copies g^{-1} dry

822 soil); **d)** Abundance of *nosZ* denitrifiers (number of *nosZ* gene copies g^{-1} dry soil) for different

823 treatments at various sampling dates.