Title

Neither wood ash application nor phosphorus-fertilisation mitigated the negative effects of whole-tree

harvesting on a temperate oligotrophic forest.

Authors

Laurent Augusto [a,*]

Florent Beaumont [a]

Christophe Nguyen [a]

Jean-Yves Fraysse [b]

Pierre Trichet [a]

Céline Meredieu [c,d]

David Vidal [a]

Valérie Sappin-Didier [a]

a INRAE, Bordeaux Sciences Agro, ISPA, F-33140 Villenave d'Ornon, France

b FCBA, F-33610 Cestas, France

c INRAE, UEFP, F-33610 Cestas, France

d INRAE, Univ. Bordeaux, BIOGECO, F-33610 Cestas, France

Abstract

<u>Background and Aims:</u> Concerns about climate change and carbon economy have prompted the

promotion of alternative energy sources, including forest-based bioenergy. An evaluation of the

environmental consequences of intensive harvests (stumps and roots, and also branches and foliage) for

energy wood supply, and use of wood-ash recycling as a compensatory practice, helps in the analysis of

the use of forest biomass for energy production.

Methods: After 11 years, we made use of records from a split-plot experimental site crossing four

different intensities of Biomass Harvesting (Stem-Only Harvest [SOH], Aboveground Additional

Harvest [AAH], Belowground Additional Harvest [BAH], and Whole-Tree Harvest [WTH]) and three

Compensation Methods (control [C], wood ash application [A] and phosphorus fertilisation [P]) to

evaluate their effects on tree growth, soil fertility, chemical properties and soil carbon. This site is

located in a maritime pine forest on a poor soil, under a warm temperate climate (SW France).

Key results: Despite their low additional biomass exports (+10% for AAH to +34% for WTH), the non-

conventional harvest practices exported much higher quantities of nutrients than the conventional SOH

technique (for example +145% for N and K in WTH). Consequently, these treatments had several effects

on the soil nutritive status. Additional biomass harvests impacted the soil organic matter content, with

negative effects on $P_{organic}$ and soil cation exchange capacity. However, data suggested that tree growth

and foliage nutrient content had not yet been significantly impacted by harvest treatments, whereas tree

nutritional status was improved by P-fertiliser or wood ash. As expected, we observed a positive effect

of wood ash application on soil acidity and nutrient content but, like additional harvests, wood ash

application decreased the pool of soil organic carbon (~10% of the initial stock with ~7% of N_{total}

losses).

Conclusions: Overall, this factorial experiment showed that exporting more forest biomass due to the

additional harvesting of tree canopies, stumps and roots had negative consequences on the ecosystem.

Additional harvests have aggravated the poverty of the already oligotrophic soil, which in turn may

decrease tree growth and the soil organic carbon content in the future (but without any impact on either

soil acidity or on trace metal contents). Importantly, applying nutrients as fertiliser or wood ash did not

compensate for the negative impact of biomass exports and the method of wood ash recycling in forests

could even decrease the soil organic carbon.

Introduction

Most recent reports of the Intergovernmental Panel on Climate Change have concluded that climate change is accelerating and requires rapid mitigation policies (IPCC 2014). These policies rely on two main and complementary approaches: the emission of less greenhouse gases (GHG), and the sequestration of more carbon dioxide (CO_2) from the atmosphere. Forest ecosystems can do both, depending on whether they are disturbed or not (Thom and Seidl 2016). Due to their large carbon contents (Pan et al. 2011), and because they represent a large proportion of land (Bastin et al. 2017), forests are pivotal ecosystems with respect to the Earth's carbon cycle. However, although there is a consensus that forests constitute an important leverage for climate change mitigation (IPCC 2014), there is a heated debate on how forests should be managed for mitigation purposes (Lindner and Karjalainen 2007): the *old-growth* strategy consisting of sequestrating carbon in the ecosystem, conflicts with the *intensive biomass harvest* strategy, which proposes to harvest more woody biomass to replace fossil C in the production of manufactured objects and energy (Bright et al. 2012; Schulze et al. 2012; Haberl et al. 2013; Luyssaert et al. 2018). In particular, additional harvesting of forest biomass, such as foliage, branches, stumps, and roots (Nicholls et al. 2009), for energy production can impoverish forest ecosystems because these tree compartments dedicated to energy wood supply –the so-called *harvest residues* or *logging residues*– are rich in nutrients (Andre et al. 2010; Achat et al. 2015a; Augusto et al. 2015a). Review studies indicate that intensive biomass harvesting –as compared with conventional harvesting– can have negative consequences on ecosystem functioning, such as soil nutrients and organic matter pools, and consequently on the level of soil productivity for following forest establishment (Thiffault et al. 2011; Wall 2012; Achat et al. 2015a). Extracting more biomass from forests without any compensatory fertilisation seems to be an unsustainable leverage for climate change mitigation (Garcia et al. 2018). Therefore, it has been proposed that intensive harvests should be associated with compensatory practices (Nohrstedt 2001; Ranius et al. 2018; Ventura et al. 2019).

Applying wood ash is often promoted as a good method to compensate for the negative effects of intensive harvests on the ecosystem nutrient budget (Hannam et al. 2018; Ranius et al. 2018; Ventura et al. 2019). Indeed, wood ash has a high nutrient content (Aronsson and Ekelund 2004), can reduce soil acidity (Reid and Watmough 2014) and contributes to the concept of *circular bioeconomy* as wood ash

returns some of the nutrients that were exported by the removal of logging residues (Pitman 2006). On

the other hand, applying wood ash might contaminate ecosystems because of its relatively high content

in several micronutrients or non-essential metals (Vance 1996; Nnadi et al. 2019). However, other

studies conclude that, based on the metal content of wood ash, a single application per forest rotation

may not result in a significant increase in soil metal content and might even improve the micronutrient

status of trees (Pitman 2006). Unfortunately, while several meta-analyses have been carried out on

intensive biomass harvests (Thiffault et al. 2011; Achat et al. 2015a) or wood ash application (Augusto

et al. 2008a; Reid and Watmough 2014), a combination of these two practices at the time of clear-cutting

and prior to reforestation have rarely been studied together to assess their interactive consequences

(Hagerberg and Wallander 2002; Wang et al. 2010). Therefore, we conducted a field experiment whose

main objective was to evaluate their interactions.

Located in south-western France, the *Landes de Gascogne* forest, where our study took place, has several

advantages for evaluating the environmental consequences of intensive harvests and their compensatory

practices. Firstly, it is a large man-made pine forest almost entirely dedicated to intensive forestry. As

such, the area is already subjected to collection of harvest residues (mainly stumps and roots, and to a

lesser extent also branches and needles (Mora et al. 2014; Banos and Dehez 2017)) for energy wood

consumption (Augusto et al. 2010, 2015a), which in turn produces large amounts of wood ash (Alvarez-

Alvarez et al. 2018). Secondly, because this forest is strongly oligotrophic (Augusto et al. 2010), it has

been diagnosed to be particularly sensitive to the potential negative consequences of intensive harvest

(Durante et al. 2019). Finally, it is a warm–temperate forest. Indeed, knowledge about wood ash

application and removal of logging residues is based on studies carried out in boreal forests (Augusto et

al. 2008a; Walmsley and Godbold 2010), whereas the consequences of these management practices are

probably climate-dependent. For instance, Achat *et al.* (2015b) found that exporting tree canopies

generally decreases the pool of soil organic carbon (SOC) under temperate climates, but not under cold

climates. Similarly, several Nordic studies indicated that stump harvests had no effect on SOC while

studies conducted in warmer latitudes concluded that the impact was highly negative (Zabowski et al.

2008; Stromgren et al. 2013; Jurevics et al. 2016).

Our initial expectations were based on the current knowledge about intensive biomass harvest or wood

ash application, in comparison with the local conditions of climate, soils, and forestry. More explicitly,

we expected the following effects (Table 1).

Hypothesis H1: the soil phosphorus, the most limiting nutrient in the sandy acidic soils of the study

region (see Materials and Methods), would be decreased by intensive harvests (Achat et al. 2015a) and

would be only partly compensated for wood ash by which has a low phosphorus content and low

bioavailability (Steenari and Lindqvist 1997; Fransson et al. 1999).

H2: the sequestration rate of C into forest biomass (estimated as the growth rate of the subsequent forest

stand) would be decreased by intensive harvests because the local soils are nutrient poor (Achat et al.

2009, 2015a), and would not be compensated for wood ash due to its low value as a P-fertiliser (see H1;

Table 1).

H3: the size of the soil organic matter pool (and related organic carbon and nitrogen (N)) would be

decreased by intensive harvests as for most temperate forests (Achat et al. 2015a, b), but might remain

unaffected by wood ash application as observed in boreal forests (Augusto et al. 2008a).

H4: the soil would be acidified by intensive harvest, but neutralised by wood ash (Reid and Watmough

2014; Achat et al. 2015a). The local soils are already quite acidic (Augusto et al. 2010), and we expected

the liming effect to prevail over the acidifying effect. The durability of neutralisation after a single ash

input in this type of forest remains to be determined.

H5: the trace metal content in the ecosystem, and its bioavailability, would not be affected either by

biomass harvests or by wood ash application. These expectations were based on two rationales. Firstly,

the local pine trees contain moderate amounts of trace metals in their biomass (Saur et al. 1992; Trichet

et al. 2018), which implies moderate exports through biomass harvest, and small returns through the

dose of wood ash applied in our field trial. This assumption is in line with current knowledge, mostly

based on Nordic podzols. Secondly, as the application of wood ash is expected to induce an increase in

pH, the bioavailability of metals might decrease (Cappuyns and Swennen 2008).

Materials and Methods

Study region and experimental site

The *Landes de Gascogne* area is a flat region ~1.3 Mha in south-western France (between 43.5 and 45.5°N, and 1.5°W and 0.3°E). The climate is oceanic, usually with a dry season in summer and a wet season in winter and/or in spring (annual rainfall is around 900 mm per year, with a North-South gradient). Soils are sandy podzols (9–14% fine sands; 80–89% coarse sands), which are classified as more or less hydromorphic and humic (an "entic podzol" in the WRB soil classification), or with a hardpan (an "ortsteinic podzol"; (Augusto et al. 2010)). They developed in Aeolian deposits of quaternary sands (the *Sables des Landes*) and are characterized by their coarse particle size and their high quartz content (Bertran et al. 2011). Soils are acidic and poor in nutrients, particularly in phosphorus (P) and they have been identified as being among the poorest soils in P in the world (in this region = 60 µg-P_{total} g^{-1}, and a global mean value ≈ 500 µg-P_{total} g^{-1}; (Achat et al. 2009; Bredoire et al. 2016)). The vegetation was formally a mosaic of heathlands and wetlands, probably derived from human-degraded forests (Jolivet et al. 2007). During the 18[th] and 19[th] centuries, the French central government promoted the installation of big farms, most of which failed to survive due to the harsh local conditions (poor acidic soils, and extremely low soil water contents in summer alternating with periods of waterlogging). From 1857, a forest of ~1.0 Mha forest was progressively created thanks to a law inciting the sowing of a local species, the Maritime pine (*Pinus pinaster* Ait.), combined with the use of ditches for drainage. Initially grown for its resin, Maritime pine is now used for wood production and is often P-fertilised at plantation to alleviate the main nutritional limitation of the region (Trichet et al. 2009). As a result of two storms (in 1999 and 2009), with the development of stump removal practices and an incentive policy for biomass power plant installation, the practice of collecting *harvest residues* as biomass-based energy sources is currently spreading (Banos and Dehez 2017).

The site used for this study is called the *Forêt du Nezer* (44.57°N, 1.05°W; elevation = 21 m asl; slope < 1%; soil type predominantly an entic podzol). Its name came from Daniel Nezer, a Swiss banker who initiated the first attempt to develop a large agricultural complex in 1766 (the *Colonie de Nezer*), but which eventually ruined him (Sargos 1997). After this short farming period, the site returned to heathlands before becoming a pine forest during the second half of the 19[th] century (see above).

Experimental design and field procedures

The experimental site is a split-plot factorial design, crossing four different intensities of *Biomass Harvesting* (BH factor in main plots) and four *Compensation Methods* by applying wood ash or phosphorus fertilisation (CM factor in subplots nested in BH plots), resulting in sixteen treatments replicated in three blocks (Figure S1). Contrary to the CM treatments, the BH treatments had to be applied to large plots (main plots) because they required the use of large machines.

After the clear-cut of a 45-year-old stand in April 2007, twelve large square plots, 100×100 m (1 ha large), were established in order to apply the three blocks $\times$ four *Biomass Harvesting* treatments, described below:

1- Stem-Only Harvest ('SOH'; stem harvest down to 7 cm of diameter above bark) without any residue harvest (exports = stems: this has been the local conventional practice for decades),

2- Aboveground Additional Harvest ('AAH'), which included stem harvest with additional harvesting of branches with part of their foliage (see below; exports = stems + canopies; canopy = branches + twigs + foliage),

3- Belowground Additional Harvest ('BAH'), which included stem (without branches, twigs, and foliage) harvest with additional harvesting of stumps and roots (exports = stems + stumps + roots),

4- Whole-Tree Harvest ('WTH') with additional harvest of all other tree compartments (exports = stems + branches + twigs + foliage + stumps + roots).

The conventional harvesting (clear-cut) was carried out in April 2007, with particular attention to the slash (branches and needles) which was gathered in each of the four 1 ha plots (Figure S2a). In June 2007, the branches were bundled, forwarded from the six main plots (AAH-WTH plots $\times$ 3 blocks; Figure S2b-d), and finally transported to the power plant. The time-lapse between this harvest and the clear-cut has enabled a partial fall of the foliage (needles, in unquantified proportions). Then, in July-August 2007, the stumps were excavated with their main roots, fragmented into a few pieces, and left on the ground in the six main plots (BAH-WTH plots $\times$ 3 blocks; Figure S2e-g), enabling the natural removal of the soil particles by the rain. These fragments were harvested in February 2008. Two years after clear-cutting, in January 2009, the area was strip ploughed to a depth of 0.3 m,

and was replanted with 1,250 per hectare (4×2 m) of one-year-old maritime pine seedlings of local

origin.

Forty-eight small sub-plots, 50×50 m (0.25 ha) wide, were established within the two-year-old

plantation. Four sub-plots of compensation methods (CM factor) were nested in every main plot

(dedicated to a BH treatment; Figure S1). The *Compensation Methods* were applied in July 2011 for the

wood ash and in December 2011 for the other fertilisers, as follows:

1- Control: no nutrient application ('C'),

2- Phosphorus fertilisation ('P'), considered as standard in this forest area (application of 60 kg-P_2O_5

per hectare in the form of superphosphate (30% P_2O_5)),

3- Phosphate-potassium fertilisation (the same P treatment as above with the addition of 60 kg-K_2O per

hectare in the form of potassium sulphate (50% K_2O)). It should be noted that this CM treatment was

not included in this study.

4- Wood ash application ('A') consisted of the application of 5 Mg of loose ash per hectare (provided

by a local power biomass plant). Taking into account the total phosphorus content of the wood ash, it

corresponded to a dose of 62 kg-P_2O_5 per hectare (regardless of P forms). The ash composition is shown

in the Table S1.

The WTH-C combined treatment was included to simulate the most severe harvest condition where all

forest residues were removed from the site and with no nutrient application. Although unusual, this

harvest management without any compensation system could be implemented if the objective is to

maximize biomass production in the case of high prices for energy wood. Conversely, the SOH-P

treatment is considered as the current usual stand management of local forest owners.

Plant measurements and field sampling

Estimates of biomass and nutrient exports (*Biomass Harvesting* factor):

The quantity of biomass, and associated nutrients, exported from the plots due the Biomass Harvest

treatments were estimated, based on a combination of measurements and models. In practice, before

clear-cutting the standing pine stand, all trees were identified and each diameter was measured at breast

height ($\approx$ 1.3 m).

Biomass estimates

Then, we selected 14 trees that represented the range of stem size of the forest. All these trees were cut

down and studied intensively:

(i) the stem diameter was measured, and wood disks were sampled, at several heights (0 m, 0.5 m, 1.3 m,

and then every 2 m); the wet mass and the dry mass of the stem disks (wood and bark, separately) were

measured,

(ii) the diameter of all branches was measured at 10 cm from their insertion level,

(iii) the total dry mass of three branch classes (living branches, treetop [*i.e.* the upper part of the stem,

at diameter < 7 cm], dead branches) were weighed (all branches of each class were pooled),

(iv) the dry mass of 5 living branches (wood + bark + foliage) per tree was weighed, and

(v) the dry mass values of the stem wood, and the stem bark, were measured separately.

The total dry mass by compartment of each sampled tree was then calculated. The tree size

measurements (diameter at breast height) were used to estimate the biomass of the different tree

compartments using allometric relationships (unpublished FCBA models, calibrated with 45 trees

distributed in 5 other pine stands of a similar age). The estimated biomass values of the tree

compartments were consistent with the measured values, so the models were used to calculate the

biomass at the plot scale, based on the inventory of all trees (measurements of the diameter at breast

height).

The effective export rate of the different tree compartments was estimated in different ways, depending

on the compartment studied:

(i) stem wood was assumed to be harvested in totality (export rate = 100%),

(ii) stem bark harvests were quantified by collecting all the bark that remained on the floor of a given

area, after the stems exported from five other stands of the region (FCBA unpublished data; export rate

$\approx$ 80%),

(iii) the effective exports of the canopies (*i.e.* branches + twigs + foliage) and of the tree belowground

(stump + roots) were estimated by direct measurements of the harvests realised. In practice, the

harvested canopies were gathered into ~2.5 m long bundles (n = 465 in total), labelled per plot, and

taken to the local power biomass plant. Bundles were weighed individually (fresh weight = 225-253 kg;

mean weight = 240 kg), and 20 bundles were taken at random to estimate their dry matter content (mean

dry content = 67% of the initial mass). All bundles were crushed to provide pellets for the power plant.

During the crushing stage, 36 samples of biomass were taken at regular time intervals (Figure S2) to

estimate the dry matter content. This procedure enabled us to estimate the effective harvest rate of the

canopy. Conversely, it was not logistically feasible to measure the effective rates for foliage, twigs, and

large branches. Based on visual observations and on field studies (Stupak et al. 2008), we assumed that

(i) the twig harvest rate was 30% higher than that of the foliage, and (ii) the branch harvest rate was

50% higher than that of the twigs. Finally, based on these ratios and on the measured harvest rate of the

whole canopy, the harvest rate values were adjusted. The values of the nutrient content in the different

canopy compartments (*i.e.* needles, twigs, branches, stembark, stemwood) were the mean values

measured on samples from the 45 trees used to build allometric relationships (see above). These mean

values were within the ranges commonly observed in the study region (Augusto et al. 2008b; Trichet et

al. 2018).

A similar approach to that used for branches was used for the belowground biomass. In practice, the

potentially available belowground biomass (*i.e.* stump + roots) was estimated for each tree based on

allometric relationships that use the stem diameter at breast height as predictive variable (Augusto et al.

2015a). The realised belowground biomass harvest was estimated from the weight of all the fresh

biomass per plot (by weighing the trucks, with and without the transported biomass). The water content

of the biomass was measured on 36 samples in order to calculate the dry biomass (Figure S2g; sampling

at regular intervals during the crushing flow; mean value per plot of the dry content = 65-72%). These

samples were used also for nutrient analyses. The measured values of nutrient content in stumps and

roots were within the ranges commonly observed (Augusto et al. 2015a), except for the potassium

content, which was higher in the present study (1.50 versus 0.99 mg g^{-1}).

All these measurements and analyses enabled us to quantify the standing biomass before the Biomass

Harvest treatments, and the effective exports due to harvests in terms of biomass and nutrients.

 Field measurements in the subsequent forest plantation

Tree growth and needle sampling:

The growth rate of the new plantation –established after biomass harvest– was assessed simply by

measuring the standing biomass of the young trees. Indeed, because the survival rate of the planted

seedlings was nearly 100%, and not affected by experimental treatments, the stand density was the

almost the same in all subplots. The individual size of trees was estimated by measuring at least 50

pines, chosen at random, per subplot. Tree height and stem circumference at breast height were measured

in December 2015 and 2019 (*i.e.* on 7- and 11-year-old trees). The aboveground biomass was then

estimated based on allometric relationships (Vidal et al. 2019).

Needles were sampled to assess the nutritive status of trees, and their degree of contamination by trace

elements. For this, 8 pines were first selected at random in each subplot. Current-year needles were

collected in December 2019 using a telescopic pruner. Needles were collected in the upper third of the

canopy, where sunlight is always directly available, and in two opposite directions.

The needles were cleaned in a bath of demineralised water before being dried with absorbent paper and

then placed in an oven for ten days at a temperature of 40°C. In order to have the most homogeneous

needle composite samples, and to avoid the operator effect, about 10% of the weight of the overall

sample was taken at random. For this, ten needles were taken from each of the 8 batches of needles.

Understorey survey:

The understorey composition was assessed in May 2019, using the "*phytovolume*" approach (Gonzalez

et al. 2013; Vidal et al. 2019). In practice, the same pair of operators surveyed all the surface area of the

studied subplot, and then estimated the soil cover percentage of the main plant functional types (*i.e.* the

perennial herb *Molinia caerulea*, the bracken fern *Pteridium aquilinum*, ericaceous shrubs (*Calluna*

vulgaris and *Erica scoparia*), and the common gorse *Ulex europaeus*). To help the operators' work, we

printed patterns representing surface areas with defined cover values (assessed in increasing classes:

1%, 2%, 3%, 5%, 7%, 10%, and then +5% up to 95%). The mean height of each plant functional type

was estimated based on three height measurements representative of the plant height range in the

surveyed area (Figure S3). The phytovolume value was computed as the product of the ground cover

(estimated in squares of 16 m^2) and the plant height. Finally, the phytovolume value was converted into

estimates of plant aboveground biomass using dedicated allometric relationships that were previously

calibrated with destructive measurements of the standing biomass (Gonzalez et al. 2013; Vidal et al.

2019).

Soil sampling:

The topsoil was sampled in all of the 36 subplots studied. To do this, a systematic grid of 8 sampling

points was used per subplot: 4 points on the tree ridges, and 4 points in the furrows, to capture the spatial

distribution of the topsoil induced by the regular design of the strip ploughing and tree plantation and

by subsequent operations of understorey control (*i.e.* a bladed roller passing in the furrows). At each

sampling point, the forest floor layer was gently removed. Then the topsoil was sampled using a corer

(length = 15 cm; diameter = 8 cm). All samples were bulked in the field, and continuously homogenised

by hand until the composite sample showed no heterogeneity. The soil bulk density (kg L^{-1}) was

estimated based on a pedo-transfer function specifically calibrated for the local soils (Augusto et al.

2010).

Laboratory analyses

Plant analyses:

The needle samples were dried at 40°C for 7 days to constant weight and ground (< 1μm) using a IKA

M20 (mill in titanium to avoid contamination of the sample).

Aliquots of powdered samples (0.25g) were digested using a mixture of 1 ml HNO$_3$ (69% vol Aristar®

for trace element analysis, VWR Chemicals) and 4 ml H$_2$O$_2$ (30% vol, Ultratrace®, ppttrace analysis

grade,Scharlau) in DigiPREP System in dry bath blocks. Then, the solution was filtered to 0.45 μm and

the volume adjusted at 50 ml with ultra-pure water. The concentrations of K, Ca, Mg, Na, Fe, Mn, Al,

Cr, Cu, Ni, Zn, and Co in the extracts were analysed by ICP-AES (inductively coupled plasma atomic

emission spectrometry) and Cd, Pb, Tl, and Mo by ICP-MS (inductively coupled plasma mass

spectrometry) (Servicio Central de Análisis de Bizkaia, Leioa-Bizkaia). The validity and accuracy of

the procedures were checked using standard reference materials ("Standard Reference Material®

1573a" certified by the National Institute of Standards and Technology).

332   Soil characteristics:

Soils were sieved to < 2 mm and air-dried. All analyses (except pH) were carried by INRAE soil testing

laboratory (INRAE-LAS, Arras, France), according to French standardised procedures or international

procedures. Soil pH was determined in a distilled water (1:5 v/v soil to solution ratio) and 0.01 M CaCl2

extract (1:10 v/v soil to solution ratio; standard NF ISO 10390:2005). The soil organic carbon content

(SOC) and total nitrogen (N_{-total}) were determined by dry combustion after correction for carbonate

(SOC: NF ISO 10694:1995, N_{-total}: NF ISO 13878). The available phosphorus (P_{-Olsen}) was determined

using the Olsen method (NF ISO 11263; sodium bicarbonate extraction). The cation exchange capacity

(CEC) and exchangeable cations (K^+, Ca^{2+}, Mg^{2+}, Na^+, Fe^{2+}, Al^{3+}, and Mn^{2+}) were measured at soil pH

using the cobaltihexamine chloride method (NF X 31–130:1999). The soil texture (5 fractions) was

determined using the Robinson pipette method (NF X 31–107:1983), but only on three soil samples

(composite samples corresponding to the three blocks of the site).

Total soil mineral element contents (major and trace) were quantified after solubilisation by fluorhydric

and perchloric acids (NF X 31–147:1996). After complete dissolution, the concentrations of K, Ca, Mg,

Na, Fe, Mn, Al, Cr, Cu, Ni, Zn, and Co in the extracts were analysed by ICP-AES (inductively coupled

plasma atomic emission spectrometry) and Cd, Pb, Tl, and Mo by ICP-MS (inductively coupled plasma

mass spectrometry).

350   Soil solution extraction and analyses:

Two chemical extractions were realised for the soils to estimate the availability of trace metals: ultra-

pure water and ammonium nitrate 1M (NH_4-NO_3). Extraction allows the quantification of the content of

elements adsorbed onto the soil with variable energy depending on the chemical nature of the extractant.

Water extractions make it possible to estimate the content of elements that would be found in the soil

solution. Ammonium nitrate 1 M extractions quantify the elements weakly adsorbed onto the solid phase

by ion exchange at pH=6.

Ultra-pure water extraction at soil pH was performed at 1:5 (M/M) soil to solution ratio, stirring on a

roller-shaking table for one hour. At the end of the extraction, the solution was extracted using a

Terumo® 10 ml syringe and filtered using a Sartorius Minisart® 0.22 µm filter and then part of the

solution was acidified at 2 % with HNO_3 for trace metal analysis. The second part was reserved at 4°C

to measure pH (using a microelectrode (pHC4000-8, Radiometer Analytical)), the concentration of

dissolved organic (DOC) and inorganic (IC) carbon (by oxidative combustion (TOC-VCSH, Shimadzu))

and the anions (PO_4^{3-} (phosphate anion), NO_2^- (nitrogen dioxide) and NO_3^- (nitrate anion)) and the cation

NH_4^+ (ammonium) by colorimetric method (Technicon auto analyser II).

The NH_4-NO_3 extractions were performed at 1:2.5 (M/M) soil to solution ratio (Symeonides and McRae

1977). The suspensions were shaken on a roller-shaking table for one hour and the solution was extracted

using a Terumo® 10 ml syringe and filtered using a Sartorius Minisart® 0.22 µm filter, and then

acidified with 2 % HNO_3.

The concentrations of P, K, Ca, Mg, Na, Fe, Mn, and Zn in extracts from two extractions were assayed

by ICP-AES and Cd and that of Pb and Cd by GF-AAS.

Data analyses

Data processing was performed with R, version 3.6.1. (R Core Team 2019). The split-plot design was

analysed by a mixed effect model (*lme* function, package nlme 3.1-143). The biomass harvest factor and

the compensation method factor were fixed effects (main and interaction effects). The block and biomass

harvest factor nested in blocks were random effects associated with the intercept. When detected,

heteroscedasticity was corrected by modelling the variance with the *varIdent*() function (nlme package)

which allows different variances by level of the biomass harvest and compensation methods. Least-

squares means were calculated (emmeans package 1.4.2) and linear contrasts were used to test for

significant differences between treatments and their corresponding controls. When necessary,

the *P*=0.05 significance probability was adjusted for multiple pairwise comparisons (Tukey's test).

Mean values are shown ± one standard error. Data and scripts are available at

https://doi.org/10.15454/LCU6OZ.

Preliminary analyses of the results showed that the interactions between the BH factor and the CM factor

were almost never statistically significant, and that is why we analysed these factors separately. The lack

of interactions between biomass residues removal and wood ash application has already been reported

(Hagerberg and Wallander 2002).

It should be noted that, due to the split-plot design, the statistical power is greater for the CM factor

nested in BH (12 replicates = 3 blocks × 4 BH treatments) compared to the BH factor (3 replicates = 3

blocks). The main consequence of this split-plot design was greater difficulty in detecting significant

effects of the BH factor.

Results

Initial stand biomass and nutrient content, and effective harvests

The standing biomass of the stand prior to harvests was 179.1 ± 3.5 Mg ha^{-1} of which 40.1 ± 0.7 Mg ha^{-1}

were belowground (Table S2). The standing biomass was distributed homogeneously among the

experimental plots, with a coefficient of variation of 1% among the WTH plots to 8% among the SOH

plots (CV=6% at the site scale). This biomass contained nutrient contents of 24.7 ± 0.5 kg-P ha^{-1}

(phosphorus) and 311 ± 6 kg-N ha^{-1} (nitrogen; Table S2). The harvesting treatments did not export all

this biomass and its associated nutrients: the effective rate of harvest was measured as 71% for

belowground biomass (in BAH and WTH treatments) and 91% for aboveground biomass (in AAH and

WTH treatments). Within the aboveground biomass pool, the measured harvest rate of canopies

(foliage + twigs + branches) was 53%, with estimated rates of 34%, 44%, and 66% for foliage, twigs

and branches, respectively. The realised export of biomass was 115.32 Mg ha^{-1} for the conventional

harvest (SOH), with moderately increased values for more intensive harvests (from +10% for AAH to

+34% for WTH; Figure 1a). Despite their low additional biomass exports, due to the differences in

nutrient contents between tree parts, the non-conventional harvest practices exported a lot more nutrients

than the conventional SOH technique. This was particularly the case for the canopy export of nitrogen

(+74% in AAH; Figure 1b) and the belowground export of potassium (+123% in BAH). In the most

impacting treatment (WTH), the increase in nutrient exports was between +64% for P (Figure 1c) and

+145% for N and K (Figure 1b and 1d).

Soil properties and fertility (hypothesis H1)

The main soil characteristics were representative of the sandy podzols of the region. For instance, the

mean soil particle size distribution showed the dominance of coarse sand ($85.7\pm0.2\%$, $7.5\pm0.5\%$,

$0.7\pm0.3\%$, $2.4\pm0.2\%$ and $3.6\pm0.3\%$, for the coarse sand, fine sand, coarse silt, fine silt and clay,

fractions, respectively), and a low initial value of the total soil P content of 44 ± 17 µg-P g^{-1} (see Augusto

et al. (2010) for comparisons).

The biomass harvesting treatments had many effects on the soil nutritive status, but the effects depended

on the form of the nutrients and micronutrients. While the AAH treatment showed increases in values

of K, Na, Fe and Mn total contents (Table 2), the pattern was quite different for exchangeable

cations (Table 3) and extractable nutrients (Table 4). Indeed, the nutrients that constitute the so-called

basic cations of the CEC (*i.e.* K^+, Ca^{2+}, Mg^{2+}, and Na^+) were depleted to varying extents (except for K^+)

by the non-conventional harvest practices (AAH, BAH, and WTH) as compared with the SOH

treatment (Table 3). The same pattern applied to Ca, Mg, and to a lesser extent K, that were NH_4-NO_3

extracted (Table 4). The losses in these nutrients (Ca, Mg, Mn, and Zn) were generally partly

compensated for by the application of wood ash (Table S1), as shown by the significant increases in

their soil content as compared with the control treatment (Tables 3 and 4).

The effects of the treatments on P also depended on the chemical form considered. The soil P content

as measured using the sodium bicarbonate extractant (P$_{-Olsen}$) showed no significant differences, even if

the non-conventional harvest treatments resulted in a decrease in this soil content (Table 2). This not-

significant trend became significant when considering the P content as measured using the NH_4-NO_3

extractant (Table 4). As expected, the application of wood ash did not modify the soil P status.

Surprisingly the P-fertiliser treatment did not improve this P status either.

The soil N$_{-total}$ content followed the same, but insignificant, trend as for P with low values in the

treatments that involved canopy harvesting (Table 2). The C/N ratio was remarkably irresponsive to all

experimental treatments, but the soil N content was reduced by wood ash application.

Vegetation response (H2)

The *Biomass Harvesting* treatments and the *Compensatory Methods* treatments showed radically

different effects on the tree foliage composition. While the intensity of biomass harvesting had no

consequences on foliage nutrient content, applying phosphorus or wood ash influenced most nutrients.

As expected, P-fertiliser improved the foliage P content, and the wood ash increased the content of many

nutrients and micronutrients (N, P, K, Mg, B, Cu, and Fe; Table 5). Conversely, we did not expect the

absence of a significant effect of wood ash on Ca content and, above all, the significant positive effect

of the P fertilisation on most elements.

The total aboveground biomass of the spontaneous vegetation remained unaffected by the *Biomass*

Harvesting treatments, but changed due to the *Compensatory Methods* treatments with an increasing

standing biomass following the ranking: Control ≤ P fertilisation ≤ Ash (Figure S4). For both factors,

the proportion of the prominent plant species was influenced by treatments. For the *Biomass Harvesting*

factor (BH), the proportion of species adapted to oligotrophic-acidic moorlands (*i.e. Molinia caerulea*

and ericaceous species) was increased by the intensive harvests as compared with the conventional SOH

treatment (Figure S4). The opposite pattern was observed in the wood ash treatment (*Compensation*

Methods factor, CM), which strongly decreased the abundance of those moorland-adapted species. The

wood ash application and, to a lesser extent, the P fertilisation treatments enhanced the growth of the N-

fixers (almost entirely composed of the spiny shrub *Ulex europaeus*).

As expected, the most severe treatments of the *Biomass Harvesting* factor (*i.e.* AAH and WTH) tended

to depress the tree growth (field observations and Figure 2), but these trends remained statistically not-

significant. Conversely we measured an unexpected significant difference between two treatments of

the *Compensatory Methods* factor, with a negative effect of P fertilisation (Figure 2). In addition, the

wood ash application tended to improve tree growth.

Soil organic matter and CEC (H3)

The soil Cation Exchange Capacity (CEC) was strongly and linearly correlated with the quantity of

elements that are commonly found as organic matter in acidic soils, which are organic carbon (SOC)

and total nitrogen (N_{-total}). In the case of organic carbon, its soil content explained 76% of the variance

of the exchange capacity (CEC = $0.088 \times$ SOC; $r^2 = 0.76$).

Due to the tight proportional relationship between CEC and SOC, the observed effects of the

experimental treatments on these variables were quite similar. The harvest of the belowground biomass

and the application of wood ash both tended to decrease the soil organic matter content and the soil CEC

(Tables 2 and 3). The dissolved organic carbon (DOC) followed a similar pattern with a significant

decrease in concentration after whole-tree harvesting (WTH) and wood ash application (Table 2).

Soil acidity status (H4)

Harvesting more biomass than conventionally practiced did not modify any of the soil variables used to

assess the acidic status of a soil (pH-H_2O, pH-$CaCl_2$, Base Saturation of the CEC; Table 3). The absence

of effect on the Base Saturation was the consequence of the concomitant decrease in *basic cations* (K^+,

Ca^{2+}, Mg^{2+}, and Na^+) and CEC.

Conversely to biomass harvesting, applying wood ash modified the soil acidity status. The soil pH-H_2O

value increased by +0.24 unit, and the Base Saturation increased from 39% to 46% of the CEC (Table 3).

The P fertilisation had no effect on soil acidity.

Ecosystem content in trace metals (H5)

Overall, the *Biomass Harvesting* treatments generally had no significant effect on the trace elements

(As, Cd, Mo, Ni, Pb, Tl, and Zn) in the ecosystem (Tables 2, 4, 5, 6, S4). The exceptions were a decrease

in soil Pb and Zn contents due to WTH.

The *Compensatory Methods* also had only a few significant effects on the trace element distribution in

the ecosystem (Tables 2, 4, 5, 6, S4). The most noticeable exceptions were increases of soil Zn content

due to wood ash application (Tables 4 and S4), an effect of P fertilisation on Cd distribution between

the soil and trees, and an improvement of Cu content of tree foliage after P or wood ash

application (Table 5).

Discussion

Importance of the study design and the study region for data interpretation

Implementing a careful harvest of adult tree compartments over small areas was hardly feasible because

the machines that collect stumps or canopies need space to work (Figure S2). Consequently, the *Biomass*

Harvesting factor could be studied only by using large plots (*i.e.* one ha per plot). With 16 experimental

treatments and three blocks, building a complete factorial experimental design (*i.e.* with 48 large plots)

was not possible because the surface area of 48 ha required was incompatible with the premise of an

area that was initially homogeneous in its properties and past-management. Therefore, we built our

experiment based on a split-plot design. Such a design had one major consequence on data analysis.

Indeed, the number of the subplots –dedicated to the *Compensation Methods* treatments– were four-fold

more numerous than the experimental areas dedicated to *Biomass Harvesting* treatments (Figure S1).

The power of the statistical tests was consequently much lower for the *Biomass Harvesting* factor, so

isolating significant effects was sometimes difficult.

The forest region where our experiment took place is atypical from a biogeochemical perspective. As a

matter of fact, the soils of the *Landes de Gascogne* region are among the poorest soils in phosphorus in

the world. While in non-agricultural systems the soil P_{-total} content ranges between ~10 and 2,000 µg g^{-1}

^{1} (Achat et al. 2009; Yang and Post 2011; Bredoire et al. 2016; Augusto et al. 2017), the study region

has a mean value of ~50 µg g^{-1} and is within a range of ~10-100 µg g^{-1} (Achat et al. 2009; Augusto et

al. 2010). As such, the study region is a hotspot of P limitation and is more similar to ancient landscapes

than to most European forests (Lambers et al. 2010; Vitousek et al. 2010). Being also acidic and poor in

other nutrients and micronutrients (Augusto et al. 2010; Trichet et al. 2018), the soils of the study region

have been identified as having the highest level of vulnerability to biomass exports according to the

national system of forest evaluation (Durante et al. 2019). Poorest of the poor, the study region is also

expected to respond strongly and negatively to additional harvests of biomass, which implies that results

should be extrapolated to other regions with caution.

Consequences of harvesting more biomass on the functioning of an oligotrophic forest

Exporting tree canopies generally enhances the amount of harvested biomass (~ +20% to +50%), but

this is at the expense of losing huge quantities of nutrients (~ +100% to +250%; (Achat et al. 2015a)).

Our results (10-34% extra biomass, and 37-145% extra nutrients) fit well with this general pattern. On

the other hand, the quantities of exported nutrients of this field experiment are about half of the values

used in previous modelling studies (Augusto et al. 2015b; Achat et al. 2018). This was the direct

consequences of the harvesting techniques that were purposely used to reduce the losses of nutrients:

letting branches shed their needles (Stupak et al. 2008), and harvesting only stumps and coarse roots

(Augusto et al. 2015a). Despite these precautions, harvesting more biomass, and above all harvesting

the whole-tree, reduced the most available forms of nutrients (K, Ca, Mg, and P) in the soil nutrient

pools (-15% to -58%; Table 4). Additional biomass harvests likely also impacted the soil content in

organic matter, with negative ripple effects on organic P and CEC, the latter being entirely dependent

on organic matter in these sandy soils (Augusto et al. 2010). In the study region the organic forms of P

are important for seedling and tree nutrition (Jonard et al. 2009; Achat et al. 2013), which would explain

why trees of the subsequent stand have groww more slowly after intensive biomass harvests (even if a

longer monitoring is needed to confirm the observed trend; see Table S3). The absence of change in the

tree foliage composition might be considered as surprising as foliage nutrient content is often used to

assess plant nutrition (CSIRO 1997; Mellert and Göttlein 2012). Nevertheless, maintaining the nutrient

content of foliage at the expense of growth is a response commonly observed after the harvest of

additional biomass (Achat et al. 2015a), and our results are consistent with this general pattern.

Fertilisation and wood ash application as possible mitigation practices

Because phosphorus is the first-order factor that limits tree growth in the study region (Trichet et al.

2009), we initially expected that the application of P-fertiliser to mitigate the negative effects of

additional harvests, but this was not the case in our experiment. Because the soil was also impoverished

in other major nutrients, and because they are considered as second-order nutritional limitations in this

context (Trichet et al. 2008), tree growth may have been constrained in our experiment by a multi-

element scarcity. Multi-element fertilisation is often put forward as a major advantage of applying wood

ash in forests (Ranius et al. 2018). Both wood ash application and P fertilisation were tested in the

present study.

The major effect of applying wood ash was expected to be a change in the acidity status of soils (Reid

and Watmough 2014). A simple approximation states that two tons of ash have the same neutralising

effect as one ton of calcium carbonate (Vance 1996). Our experiment was not an exception and the soil

pH, base saturation and concomitant available calcium and magnesium, were improved by wood ash

application (for results in a similar context to ours, see Gomez-Rey et al. (2013)). The available soil

micronutrient contents of Mn and Zn also logically increased because wood ash contains them in

substantial quantities (Table S1). Whereas the effects of wood ash were expected and straight forward

to interpret, the P fertilisation had surprising outcomes: the soil P status was not improved, and the soil

contents of available Ca and Mg actually decreased by P fertilisation. These results may be explained

by the tree foliage composition that was modified by both P fertilisation and wood ash application, with

increases in concentration values for most nutrients and micronutrients (and even some non-essential

trace metals). Hence, it seemed that the two compensatory methods improved tree nutrition but, for P

fertilisation, it was at the expense of the soil reserves in Ca and Mg.

In turn, these changes in the soil properties modified the dynamics of the spontaneous vegetation and of

the tree plantation. The vegetation experienced a shift of species prominence after the wood ash

application, with a decrease in the species typical of acidic moorlands and heathlands (*i.e. Molinia*

caerulea and ericaceous species) to the benefit of the local main N-fixer (the shrub *Ulex europaeus*).

The decrease of the abundance of the acidophilus species is logical since wood ash decreased the soil

acidity. The increase of the N-fixer shrub was expected as we are used to observing a positive response

of this leguminous species to P fertilisation (*e.g.* Delerue et al. (2015)), but although the N-fixers

abundance was increased by P fertilisation, it was relatively low compared with previous experiments

(Augusto et al. 2005; Vidal et al. 2019), and lower than results after the wood ash application. This effect

has already been reported in the literature (Campillo et al. 2005) and may be the positive consequence

of reducing the ambient acidity on the symbionts of the leguminous species (Slattery and Coventry

1995).

Trees that were planted responded differently to the *Compensatory Methods*. Firstly, trees that received

wood ash tended to grow the fastest. This result is consistent with our interpretation that plant growth

in this site was limited by several elements at the same time. Since wood ash application improved the

availability of many nutrients and micronutrients, it was logical that plant growth was enhanced. On the

other hand, the depressive effect of P fertilisation was surprising considering that this practice has proven

its efficiency in the study region for more than half a century (Trichet et al. 2009). As explained above,

it seems that P fertilisation improved the tree foliage composition at the expense of soil fertility, and we

speculate that this soil impoverishment might be at the origin of subsequent degraded tree growth.

However, tis atypical phenomenon requires further investigation.

Considering the positive effects of wood ash application on soil acidity and nutrient content, and its

positive repercussion on tree growth, one might conclude that this compensatory method should be

promoted in oligotrophic forests submitted to intensive biomass harvestings. Nevertheless, this

conclusion does not take into account one major drawback of wood ash application in our experiment,

which was a strong decrease of the soil organic matter content, with negative consequences on organic

carbon and total nitrogen contents and soil exchange capacity. Even after correcting for the change in

soil bulk density, the losses of organic carbon were ~10-11% of the initial stock of the topsoil (N_{total}

losses were ~7%). We interpret this decrease as the result of the acidity alleviation by wood ash, which

probably enhanced the soil microbial activity and respiration (Bååth and Arnebrant 1994; Jokinen et al.

2006; Omil et al. 2013).

In a meta-analysis, we previously concluded that wood ash had no influence on soil organic carbon

(Augusto et al. 2008a), but this study was almost entirely based on Nordic, or cold temperate

experiments (mean annual temperature < 8.5°C; *e.g.* (Feldkirchner et al. 2003)). Under a warm

temperate climate, wood ash application seems to increase SOC stocks when applied in recent

afforestation of former croplands (Sartori et al. 2007; Ventura et al. 2019), which can be explained by

the initial soil conditions (*i.e.* low SOC values and high pH values) and improved necromass production

by plants (*i.e.* litterfall and fine root turnover). In areas with a long history of forest occupation (*i.e.* high

SOC values and often low pH values), to our knowledge, our experiment is only the second study under

a warm temperate climate that has quantified the effect of wood ash on SOC: both studies observed a

possible decrease in organic carbon soil contents (Solla-Gullon et al. 2006).

Initial expectations and future anticipation

Most, but not all, of our initial expectations were confirmed by the field experiment. Firstly, as

anticipated, harvesting more biomass aggravated the P scarcity of the local soils, and the application of

wood ash did not modify the soil P content (Hypothesis H1; Table S5). Still in line with our expectations,

the impoverishment of the soil caused by additional harvests of biomass could result in reduced growth

of the subsequent forest stand; wood ash application had positive effects on early tree growth (H2). The

fact that, unexpectedly, P application did not enable trees to overcome P limitation suggested that other

nutritional limitations –such as Ca and Mg– prevented the experimental fertilisation from having an

effect. If our two first hypotheses were fairly well supported by results, the third and fourth hypotheses

received mixed support. Indeed, both high rates of biomass harvesting and wood ash application

decreased the pool of soil organic carbon, but only the former treatment was supposed to do so (H3).

The SOC destocking caused by wood ash application opened perspectives on a large scale (see below).

Similarly, while wood ash application logically improved the acidity status of the soil to a moderate

extent, intensive biomass exports did not deteriorate it, as in many other studies (H4; see above for

possible explanations). Finally, none of the experimental factors had the anticipated major effects on the

trace metal distribution within the ecosystem (H5).

Overall, this factorial experiment showed that exporting more forest biomass though the additional

harvests of tree canopies, stumps, and roots had negative consequences on the ecosystem. Additional

harvests have aggravated the poverty of the already oligotrophic soil, which in turn may decrease tree

growth and the soil content of organic carbon in the future (but without any impact either on soil acidity

or on trace metals content). Importantly, applying nutrients as fertiliser or wood ash did not fully

compensate for the negative impact of biomass exports.

Beyond the management consequences for the *Les Landes de Gascogne* region, this study may have an

interesting repercussion on research about climate change mitigation. Indeed, the *intensive biomass*

harvest strategy (which proposes to harvest more woody biomass to replace fossil C in the production

of energy), in conjunction with the application of wood ash to return part of the exported nutrients, can

be seen as a winning approach in boreal forests because neither biomass harvests (Achat et al. 2015b)

nor wood ash (Augusto et al. 2008a) destock soil carbon in this biome. In temperate forests, intensive

biomass harvests are already known to decrease soil organic carbon (Achat et al. 2015b), but wood ash

was not suspected to do so, probably because current knowledge is based on research carried out in

Nordic countries (Augusto et al. 2008a). This study is among the first field experiments to have tested

the effect of wood ash application on soil organic carbon under a warm temperate climate. Although we

acknowledge that our study is based on only one field experiment and should be followed by others, we

anticipate that, if the negative impact of wood ash on soil carbon sequestration was confirmed in several

other temperate forests, the *intensive biomass harvest* strategy would be inefficient in such ecosystems

and should be replaced by the *old-growth* strategy –consisting of sequestrating carbon in the ecosystem–

to mitigate climate change.

Acknowledgements

We are grateful to the ADEME, GIS G-PMF and the Council of the Nouvelle-Aquitaine region for their

financial support (D.V. PhD grant; SYLVOGENE and PINASTER projects). Similarly, we thank the

ISPA lab for providing the F.B. Master grant. Finally, we are grateful to the Forest Experimental Facility

(UEFP-https://doi.org/10.15454/1.5483264699193726E12) for their assistance during the successive

field campaigns, and to Sylvie Bussière, Cécile Coriou, and Catherine Lambrot for their help during the

sample analyses.

References

Achat DL, Bakker MR, Augusto L, et al (2009) Evaluation of the phosphorus status of P-deficient

podzols in temperate pine stands: combining isotopic dilution and extraction methods.

Biogeochemistry 92:183–200. doi: 10.1007/s10533-008-9283-7

Achat DL, Bakker MR, Augusto L, Morel C (2013) Contributions of microbial and physical-chemical

processes to phosphorus availability in Podzols and Arenosols under a temperate forest. Geoderma

211:18–27. doi: DOI 10.1016/j.geoderma.2013.07.003

Achat DL, Deleuze C, Landmann G, et al (2015a) Quantifying consequences of removing harvesting

residues on forest soils and tree growth–A meta-analysis. For Ecol Manage 348:124–141. doi: doi:

10.1016/j.foreco.2015.03.042

Achat DL, Fortin M, Landmann G, et al (2015b) Forest soil carbon is threatened by intensive biomass

harvesting. Sci Rep 5:15991

Achat DL, Martel S, Picart D, et al (2018) Modelling the nutrient cost of biomass harvesting under

different silvicultural and climate scenarios in production forests. For Ecol Manage 429:642–653.

doi: 10.1016/j.foreco.2018.06.047

Alvarez-Alvarez P, Pizarro C, Barrio-Anta M, et al (2018) Evaluation of tree species for biomass energy

production in Northwest Spain. FORESTS 9:. doi: 10.3390/f9040160

Andre F, Jonard M, Ponette Q (2010) Biomass and nutrient content of sessile oak (Quercus petraea

(Matt.) Liebl.) and beech (Fagus sylvatica L.) stem and branches in a mixed stand in southern

Belgium. Sci Total Environ 408:2285–2294. doi: DOI 10.1016/j.scitotenv.2010.02.040

Aronsson KA, Ekelund NGA (2004) Biological effects of wood ash application to forest and aquatic

ecosystems. J Environ Qual 33:1595–1605. doi: 10.2134/jeq2004.1595

Augusto L, Achat DL, Bakker MR, et al (2015a) Biomass and nutrients in tree root systems–sustainable

harvesting of an intensively managed Pinus pinaster (Ait.) planted forest. Glob Chang Biol

Bioenergy 7:231–243. doi: doi: 10.1111/gcbb.12127

Augusto L, Achat DL, Jonard M, et al (2017) Soil parent material - A major driver of plant nutrient

limitations in terrestrial ecosystems. Glob Chang Biol 23:3808–3824

Augusto L, Bakker MR, Meredieu C (2008a) Wood ash applications to temperate forest ecosystems -

potential benefits and drawbacks. Plant Soil 306:181–198. doi: 10.1007/s11104-008-9570-z

Augusto L, Bakker MR, Morel C, et al (2010) Is "grey literature" a reliable source of data to characterize

soils at the scale of a region? A case study in a maritime pine forest in southwestern France. Eur J Soil Sci 61:807–822

Augusto L, Crampon N, Saur E, et al (2005) High rates of nitrogen fixation of Ulex species in the understory of maritime pine stands and the potential effect of phosphorus fertilization. Can J For Res Can Rech For 35:1183–1192. doi: 10.1139/x05-054

Augusto L, De Schrijver A, Vesterdal L, et al (2015b) Influences of evergreen gymnosperm and deciduous angiosperm tree species on the functioning of temperate and boreal forests. Biol Rev 90:444–466

Augusto L, Meredieu C, Bert D, et al (2008b) Improving models of forest nutrient export with equations that predict the nutrient concentration of tree compartments. Ann For Sci 65:. doi: 10.1051/forest:2008059

Bååth E, Arnebrant K (1994) Growth rate and response of bacterial communities to pH in limed and ash treated forest soils. Soil Biol Biochem 26:995–1001

Banos V, Dehez J (2017) Le bois-énergie dans la tempête, entre innovation et captation ? Les nouvelles ressources de la forêt landaise. Natures Sci Sociétés 25:122–133

Bastin JF, Berrahmouni N, Grainger A, et al (2017) The extent of forest in dryland biomes. Science (80-) 356:635–638

Bertran P, Bateman MD, Hernandez M, et al (2011) Inland aeolian deposits of south-west France: facies, stratigraphy and chronology. J Quat Sci 26:374–388

Bredoire F, Bakker MR, Augusto L, et al (2016) What is the P value of Siberian soils? Soil phosphorus status in south-western Siberia and comparison with a global data set. Biogeosciences 13:2493–2509. doi: 10.5194/bg-13-2493-2016

Bright RM, Cherubini F, Astrup R, et al (2012) A comment to "Large-scale bioenergy from additional harvest of forest biomass is neither sustainable nor greenhouse gas neutral": Important insights beyond greenhouse gas accounting. Glob Chang Biol Bioenergy 4:617–619

Campillo R, Urquiaga S, Undurraga P, et al (2005) Strategies to optimise biological nitrogen fixation in legume/grass pastures in the southern region of Chile. Plant Soil 273:57–67

Cappuyns V, Swennen R (2008) The application of pHstat leaching tests to assess the pH-dependent

release of trace metals from soils, sediments and waste materials. J Hazard Mater 158:185–195

CSIRO (1997) Plant analysis: an interpretation manual. CSIRO Publishing, Reuter, D. J. and Robinson,

J. B. (Eds), Collingwood, Australia

Delerue F, Gonzalez M, Michalet R, et al (2015) Weak Evidence of Regeneration Habitat but Strong

Evidence of Regeneration Niche for a Leguminous Shrub. PLoS One 10:. doi:

10.1371/journal.pone.0130886

Durante S, Augusto L, Achat DL, et al (2019) Diagnosis of forest soil sensitivity to harvesting residues

removal - A transfer study of soil science knowledge to forestry practitioners. Ecol Indic 104:512–

523. doi: 10.1016/j.ecolind.2019.05.035

Feldkirchner DC, Wang C, Gower ST, et al (2003) Effects of nutrient and paper mill biosolids

amendments on the growth and nutrient status of hardwood forests. For Ecol Manage 177:95–116.

doi: 10.1016/S0378-1127(02)00318-3

Fransson AM, Bergkvist B, Tyler G (1999) Phosphorus solubility in an acid forest soil as influenced by

form of applied phosphorus and liming. Scand J For Res 14:538–544. doi:

10.1080/02827589950154014

Garcia W de O, Amann T, Hartmann J (2018) Increasing biomass demand enlarges negative forest

nutrient budget areas in wood export regions. Sci Rep 8:. doi: 10.1038/s41598-018-22728-5

Gomez-Rey MX, Madeira M, Coutinho J (2013) Soil C and N dynamics, nutrient leaching and fertility

in a pine plantation amended with wood ash under Mediterranean climate. Eur J For Res 132:281–

295. doi: 10.1007/s10342-012-0674-x

Gonzalez M, Augusto L, Gallet-Budynek A, et al (2013) Contribution of understory species to total

ecosystem aboveground and belowground biomass in temperate Pinus pinaster Ait. forests. For

Ecol Manage 289:38–47. doi: DOI 10.1016/j.foreco.2012.10.026

Haberl H, Schulze ED, Korner C, et al (2013) Response: complexities of sustainable forest use. Glob

Chang Biol Bioenergy 5:1–2

Hagerberg D, Wallander H (2002) The impact of forest residue removal and wood ash amendment on

the growth of the ectomycorrhizal external mycelium. FEMS Microbiol Ecol 39:139–146. doi:

10.1016/S0168-6496(01)00207-0

Hannam KD, Venier L, Allen D, et al (2018) Wood ash as a soil amendment in Canadian forests: what

are the barriers to utilization? Can J For Res 48:442–450. doi: 10.1139/cjfr-2017-0351

IPCC (2014) Climate Change 2014: Synthesis Report. Fifth Assessment Report. Geneva, Switzerland

Jokinen H, Kiikkilä O, Fritze H (2006) Exploring the mechanisms behind elevated microbial activity

after wood ash application. Soil Biol Biochem 38:2285–2291

Jolivet C, Augusto L, Trichet P, Arrouays D (2007) Les sols du massif forestier des Landes de

Gascogne : formation, histoire, propriétés et variabilité spatiale. Rev For Fr 59:7–30

Jonard M, Augusto L, Morel C, et al (2009) Forest floor contribution to phosphorus nutrition:

experimental data. Ann For Sci 66:510. doi: 10.1051/forest/2009039

Jurevics A, Peichl M, Olsson BA, et al (2016) Slash and stump harvest have no general impact on soil

and tree biomass C pools after 32-39 years. For Ecol Manage 371:33–41. doi:

10.1016/j.foreco.2016.01.008

Lambers H, Brundrett MC, Raven JA, Hopper SD (2010) Plant mineral nutrition in ancient landscapes:

high plant species diversity on infertile soils is linked to functional diversity for nutritional

strategies. Plant Soil 334:11–31

Lindner M, Karjalainen T (2007) Carbon inventory methods and carbon mitigation potentials of forests

in Europe: a short review of recent progress. Eur J For Res 126:149–156

Luyssaert S, Marie G, Valade A, et al (2018) Trade-offs in using European forests to meet climate

objectives. Nature 562:259–262

Mellert KH, Göttlein A (2012) Comparison of new foliar nutrient thresholds derived from van den

Burg's literature compilation with established central European references. Eur J For Res

131:1461–1472

Mora O, Banos V, Regolini M, Carnus JM (2014) Using scenarios for forest adaptation to climate

change: a foresight study of the Landes de Gascogne Forest 2050. Ann For Sci 71:313–324

Nicholls D, Monserud RA, Dykstra DP (2009) International bioenergy synthesis-Lessons learned and

opportunities for the western United States. For Ecol Manage 257:1647–1655

Nnadi EO, Mbah CN, Nweke AI, Njoku C (2019) Physicochemical properties of an acid ultisol

subjected to different tillage practices and wood-ash amendment: Impact on heavy metal

concentrations in soil and Castor plant. Soil Tillage Res 194:104288

Nohrstedt HO (2001) Response of coniferous forest ecosystems on mineral soils to nutrient additions:

A review of Swedish experiences. Scand J For Res 16:555–573. doi: 10.1080/02827580152699385

Omil B, Pineiro V, Merino A (2013) Soil and tree responses to the application of wood ash containing

charcoal in two soils with contrasting properties. For Ecol Manage 295:199–212. doi:

10.1016/j.foreco.2013.01.024

Pan YD, Birdsey RA, Fang JY, et al (2011) A large and persistent carbon sink in the world's forests.

Science (80-) 333:988–993

Pitman RM (2006) Wood ash use in forestry - a review of the environmental impacts. FORESTRY

79:563–588. doi: 10.1093/forestry/cpl041

R Core Team (2019) R: A language and environment for statistical computing. R Foundation for

Statistical Computing; Vienna; Austria. URL https://www.R-project.org

Ranius T, Hamalainen A, Egnell G, et al (2018) The effects of logging residue extraction for energy on

ecosystem services and biodiversity: A synthesis. J Environ Manage 209:409–425. doi:

10.1016/j.jenvman.2017.12.048

Reid C, Watmough SA (2014) Evaluating the effects of liming and wood-ash treatment on forest

ecosystems through systematic meta-analysis. Can J For Res 44:867–885. doi: 10.1139/cjfr-2013-

0488

Sargos J (1997) Histoire de la forêt landaise. L'horizon Chimérique (Ed.), Bordeaux, France.

Sartori F, Lal R, Ebinger MH, Miller RO (2007) Tree species and wood ash affect soil in Michigan's

Upper Peninsula. Plant Soil 298:125–144. doi: 10.1007/s11104-007-9345-y

Saur E, Ranger J, Lemoine B, Gelpe J (1992) Micronutrient Distribution in 16-Year-Old Maritime Pine.

Tree Physiol 10:307–316

Schulze ED, Korner CI, Law BE, et al (2012) Large-scale bioenergy from additional harvest of forest

biomass is neither sustainable nor greenhouse gas neutral. Glob Chang Biol Bioenergy 4:611–616

Slattery JF, Coventry DR (1995) Acid-tolerance and symbiotic effectiveness of Rhizobium-

Leguminosarum by Trifoloo isolated from subterranean clover growing in permanent pastures.

Soil Biol Biochem 27:111–115. doi: 10.1016/0038-0717(94)00143-O

Solla-Gullon F, Santalla M, Rodriguez-Soalleiro RJ, Merino A (2006) Nutritional status and growth of

a young Pseudotsuga menziesii plantation in a temperate region after application of wood-bark

ash. For Ecol Manage 237:312–321. doi: 10.1016/j.foreco.2006.09.054

Steenari BM, Lindqvist O (1997) Stabilisation of biofuel ashes for recycling to forest soil. Biomass

Bioenergy 13:39–50. doi: 10.1016/S0961-9534(97)00024-X

Stromgren M, Egnell G, Olsson BA (2013) Carbon stocks in four forest stands in Sweden 25 years after

harvesting of slash and stumps. For Ecol Manage 290:59–66. doi: 10.1016/j.foreco.2012.06.052

Stupak I, Nordfjell T, Gundersen P (2008) Comparing biomass and nutrient removals of stems and fresh

and predried whole trees in thinnings in two Norway spruce experiments. Can J For Res Can Rech

For 38:2660–2673

Symeonides C, McRae SG (1977) The assessment of plant-available cadmium in soils. J Environ Qual

6:120–123

Thiffault E, Hannam KD, Pare D, et al (2011) Effects of forest biomass harvesting on soil productivity

in boreal and temperate forests - A review. Environ Rev 19:278–309

Thom D, Seidl R (2016) Natural disturbance impacts on ecosystem services and biodiversity in

temperate and boreal forests. Biol Rev 91:760–781

Trichet P, Bakker MR, Augusto L, et al (2009) Fifty years of fertilization experiments on Pinus pinaster

in southwest France: the importance of phosphorus as a fertilizer. For Sci 55:390–402

Trichet P, Cheval N, Lambrot C, et al (2018) Using a dune forest as a filtering ecosystem for water

produced by a treatment plant - One decade of environmental assessment. Sci Total Environ

640:849–861. doi: 10.1016/j.scitotenv.2018.05.263

Trichet P, Loustau D, Lambrot C, Linder S (2008) Manipulating nutrient and water availability in a

maritime pine plantation: effects on growth, production, and biomass allocation at canopy closure.

Ann For Sci 65:

Vance ED (1996) Land application of wood-fired and combination boiler ashes: An overview. J Environ

Qual 25:937–944. doi: 10.2134/jeq1996.00472425002500050002x

Ventura M, Panzacchi P, Muzzi E, et al (2019) Carbon balance and soil carbon input in a poplar short

rotation coppice plantation as affected by nitrogen and wood ash application. NEW For 50:969–

990. doi: 10.1007/s11056-019-09709-w

Vidal DF, Trichet P, Puzos L, et al (2019) Intercropping N-fxing shrubs in pine plantation forestry as

an ecologically sustainable management option. For Ecol Manage 437:175–187

Vitousek PM, Porder S, Houlton BZ, Chadwick OA (2010) Terrestrial phosphorus limitation:

mechanisms, implications, and nitrogen-phosphorus interactions. Ecol Appl 20:5–15

Wall A (2012) Risk analysis of effects of whole-tree harvesting on site productivity. For Ecol Manage

282:175–184

Walmsley JD, Godbold DL (2010) Stump harvesting for bioenergy - A review of the environmental

impacts. Forestry 83:17–38

Wang P, Olsson BA, Arvidsson H, Lundkvist H (2010) Short-term effects of nutrient compensation

following whole-tree harvesting on soil and soil water chemistry in a young Norway spruce stand.

Plant Soil 336:323–336. doi: 10.1007/s11104-010-0484-1

Yang X, Post WM (2011) Phosphorus transformations as a function of pedogenesis: A synthesis of soil

phosphorus data using Hedley fractionation method. Biogeosciences 8:2907–2916

Zabowski D, Chambreau D, Rotramel N, Thies WG (2008) Long-term effects of stump removal to

control root rot on forest soil bulk density, soil carbon and nitrogen content. For Ecol Manage

255:720–727

847   **Figure 1** – Realised exports of biomass, nitrogen, and phosphorus

848
849

850
SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest;
WTH = whole-tree harvest.
All data (including data about calcium, and magnesium) are presented in Table S2.
The methods used to quantify the biomass and nutrient exports are detailed in the manuscript in the section
"*Plant measurements and field sampling*", in particular in the subsection "*Estimates of biomass and nutrient*
exports".
857
858

861 **Figure 2** – Tree size as affected by biomass harvest and compensation methods
862

SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground
additional harvest; WTH = whole-tree harvest. Control = no nutrient application.
P values for differences between treatments of reference (*i.e.* SOH and Control) on the one
hand, and other treatments on the other hand, are indicated in the graph.
Trees were measured at 11 years old.
Other tree metrics are shown in Table S3.
869
870

873 **Table 1** – Initial expected consequences of intensive biomass harvesting and wood ash
874 application on a temperate oligotrophic forest
875 (➔: means "has no effect on…"; ↘: means "decreases..."; ↗: means "increases..."; **?**: means "unknown
876 effect on…")

Ecosystem trait		Intensive harvests	Wood ash
Soil fertility (phosphorus) {hypothesis H1}	*General knowledge*	↘ soil P	➔ soil P
	Local conditions	P-limited soils	
	Expected results	↘ (aggravated P-limitation)	➔ (or ↗) soil P
Tree growth (carbon sequestration) {H2}	*General knowledge*	↘ tree growth	➔ (or ↗) tree growth
	Local conditions	oligotrophic P-limited ecosystem	
	Expected results	↘ tree growth	➔ tree growth
Soil Organic Matter SOM (carbon and nitrogen) {H3}	*General knowledge*	➔ boreal forests ↘ temperate forests	➔ boreal forests **?** temperate forests
	Local conditions	warm temperate	
	Expected results	↘ SOM	➔ SOM
Soil acidity (pH, base saturation) {H4}	*General knowledge*	↘ acido-basic status	↗ acido-basic status
	Local conditions	acidic soils	
	Expected results	↘ acido-basic status	↗ acido-basic status
Trace metals in ecosystem (micronutrients: [B, Cu, Fe, Mn, Mo, Zn] and toxic trace metals [As, Pb, Cd]) {H5}	*General knowledge*	➔ or ↘ trace metals (due to exports)	➔ for podzol soils (boreal soils in Nordic areas)
	Local conditions	low content in tree biomass growing on podzol soils	
	Expected results	➔ trace metals	➔ or ↘ bioavailability at moderate ash dose + increase pH

877
878
879

881 **Table 2** – Soil properties related to site fertility as affected by biomass harvest and compensation methods

	SOC (mg g^{-1})	DOC (mg L^{-1})	N (mg g^{-1})	C/N unit less	P$_{Olsen}$ (µg g^{-1})	K (mg g^{-1})	Ca (mg g^{-1})	Mg (mg g^{-1})	Na (mg g^{-1})	Fe (mg g^{-1})	Mn (µg g^{-1})	Mo (µg g^{-1})	Zn (µg g^{-1})
Biomass Harvesting													
SOH	38.6±3.9	44±2	1.41±0.15	27.7±0.8	29.3±2.2	4.07±0.2	0.82±0.06	0.17±0.09	1.17±0.09	1.8±0.4	41.5±3.5	0.16±0.01	6.60±0.29
AAH	35.4±3.9	42±4	1.46±0.15	24.5±0.8	25.3±2.5	5.40±0.2 *	0.98±0.06	0.25±0.09	1.65±0.08 *	3.1±0.3 *	62.0±3.5 *	0.21±0.01	7.37±0.48
BAH	30.2±3.9	39±2	1.14±0.15	26.3±0.8	25.3±2.9	4.31±0.2	0.71±0.04	0.15±0.08	1.22±0.08	2.2±0.3	43.6±3.5	0.15±0.01	5.77±0.36
WTH	31.1±3.9	31±2 *	1.10±0.15	28.2±0.8	21.0±3.5	4.09±0.2	0.72±0.03	0.11±0.04	1.11±0.05	1.5±0.3	35.4±3.5	0.13±0.01	3.85±0.54 *
Compensation Methods													
Control	37.1±2.9	39±1	1.36±0.11	27.2±0.7	27.1±2.4	4.47±0.13	0.85±0.03	0.18±0.10	1.31±0.05	2.1±0.3	46.6±3.2	0.16±0.01	5.69±0.31
Phosphorus	33.6±2.9	42±2	1.28±0.11	26.3±0.7	24.5±2.4	4.45±0.26	0.74±0.03 *	0.16±0.08	1.26±0.09	2.0±0.3	45.0±4.9	0.17±0.01	5.57±0.32
Ash	30.8±2.9 *	36±1 *	1.18±0.11 *	26.5±0.7	24.0±2.4	4.50±0.15	0.85±0.03	0.17±0.09	1.31±0.06	2.3±0.3	45.2±2.0	0.16±0.01	6.42±0.36

883
884 All values are for the total content of the elements in the soil (except for P). SOC = soil organic carbon; DOC = dissolved organic carbon. For phosphorus (P), values are extractible P using the
885 Olsen method.
886 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.
887 Treatments with an asterisk differ at *P*<0.05 from their reference treatment (SOH or Control, respectively; see Methods).
888
889

892 **Table 3** – Soil acidity status as affected by biomass harvest and compensation methods
893

	pH-H$_2$O unit less	pH-CaCl$_2$ unit less	CEC$_m$ (cmol$_c$ kg^{-1})	BS (%)	K (cmol$_c$ kg^{-1})	Ca (cmol$_c$ kg^{-1})	Mg (cmol$_c$ kg^{-1})	Na (cmol$_c$ kg^{-1})
Biomass Harvesting								
SOH	4.43±0.07	3.43±0.08	3.51±0.23	43±2	0.06±0.00	0.92±0.04	0.36±0.02	0.06±0.00
AAH	4.68±0.07	3.69±0.08	2.90±0.24	38±2	0.06±0.00	0.67±0.03 *	0.21±0.01 *	0.05±0.00
BAH	4.49±0.07	3.57±0.08	2.51±0.23 *	42±2	0.05±0.00	0.65±0.04 *	0.23±0.02 *	0.04±0.00 *
WTH	4.51±0.07	3.46±0.08	2.90±0.24	39±2	0.06±0.01	0.69±0.06	0.28±0.02 *	0.04±0.00 *
Compensation Methods								
Control	4.41±0.06	3.47±0.07	3.45±0.21	39±2	0.06±0.00	0.74±0.05	0.29±0.01	0.05±0.00
Phosphorus	4.53±0.05	3.51±0.07	2.58±0.19 *	38±2	0.06±0.00	0.57±0.04 *	0.22±0.02 *	0.04±0.00 *
Ash	4.65±0.05 *	3.64±0.07	2.84±0.19 *	46±2 *	0.06±0.00	0.89±0.04 *	0.30±0.02	0.05±0.00

894
SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.
CEC$_m$ = cationic exchange capacity (measured value); BS = base saturation of the CEC, which is the proportion of charges occupied by non-acidic exchangeable cations (K,
Ca, Mg, Na).
For consistency between the different terms of its equation, BS was calculated using the effective CEC (CEC$_e$: calculated as the sum all the charges of all the analysed cations),
whose values differ slightly from CEC$_m$ values.
Treatments with an asterisk differ at *P*<0.05 from their reference treatment (SOH or Control, respectively; see Methods).
901
902
903
904

907 **Table 4** – Soil composition in NH_4-NO_3 extractable elements as affected by biomass harvest and compensation methods

	P (µg g^{-1})	K (µg g^{-1})	Ca (µg g^{-1})	Mg (µg g^{-1})	Fe (µg g^{-1})	Mn (µg g^{-1})	Zn (µg g^{-1})
Biomass Harvesting							
SOH	1.2±0.3	27±2	178±8	47±2	7.0±1.2	1.5±0.1	0.60±0.05
AAH	0.5±0.2 *	21±2 (*)	117±10 *	25±2 *	9.7±1.2	1.3±0.1	0.48±0.04
BAH	0.8±0.2	21±2 (*)	128±8 *	30±2 *	8.2±1.2	1.2±0.1	0.47±0.04
WTH	0.9±0.2	21±2 (*)	151±12	38±2	6.1±1.2	1.2±0.1	0.45±0.04
Compensation Methods							
Control	0.9±0.2	23±1	131±9	35±2	7.2±1.3	1.1±0.1	0.44±0.02
Phosphorus	0.7±0.2	22±1	117±9	30±2 *	8.5±0.9	1.1±0.1	0.48±0.03
Ash	0.9±0.1	22±1	183±9 *	40±1 *	7.5±0.5	1.8±0.1 *	0.58±0.06 *

910 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.

911 Treatments with an asterisk differ at $P<0.05$ (except for K: P value < 0.10; symbol in brackets) from their reference treatment (SOH or Control, respectively; see Methods).

917 **Table 5** – Tree nutritional status (needle composition) as affected by biomass harvest and compensation methods

	N (mg g^{-1})	P (mg g^{-1})	K (mg g^{-1})	Ca (mg g^{-1})	Mg (mg g^{-1})	B (µg g^{-1})	Cu (µg g^{-1})	Fe (µg g^{-1})	Mn (µg g^{-1})	Mo (µg g^{-1})	Zn (µg g^{-1})
Biomass Harvesting											
SOH	12.8±0.7	0.64±0.04	4.6±0.1	2.3±0.1	1.4±0.1	16.3±1.0	1.8±0.2	27.8±2.0	96.3±11.0	0.011±0.003	29.8±2.5
AAH	13.9±0.7	0.69±0.04	4.4±0.1	2.2±0.1	1.5±0.1	16.7±1.0	1.8±0.2	32.1±2.1 *	104.2±10.9	0.014±0.002	32.1±2.5
BAH	12.1±0.7	0.65±0.04	4.5±0.1	2.2±0.1	1.4±0.1	17.2±1.0	2.0±0.2	27.1±2.0	92.9±11.7	0.009±0.003	30.5±2.5
WTH	12.6±0.7	0.65±0.05	4.6±0.1	2.3±0.1	1.4±0.1	17.2±1.0	1.8±0.3	28.9±2.9	81.0±11.7	0.012±0.002	32.1±2.5
Compensation Methods											
Control	12.3±0.7	0.64±0.04	4.2±0.1	2.2±0.1	1.4±0.1	16.0±0.9	1.7±0.1	27.0±2.1	92.3±9.8	0.011±0.002	29.1±2.2
Phosphorus	13.6±0.7 *	0.66±0.04 *	4.5±0.1 *	2.3±0.1	1.5±0.1 *	18.0±0.9 *	1.9±0.1	30.4±2.1 *	100.1±9.4 *	0.012±0.002	32.7±2.2 *
Ash	12.7±0.7 *	0.67±0.04 *	4.8±0.1 *	2.3±0.1	1.4±0.1	16.6±0.9 *	2.0±0.1	29.6±2.0 *	88.4±9.8	0.012±0.002	31.7±2.2 *

919
920 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.
921 Treatments with an asterisk differ at *P*<0.05 from their reference treatment (SOH or Control, respectively; see Methods).
922
923

926 **Table 6** – Ecosystem content in non-essential trace metals

	Cadmium			Lead			Arsenic
	Soil (μg-Cd kg^{-1})	*Extractable* (μg-Cd kg^{-1})	*Foliage* (μg-Cd g^{-1})	*Soil* (μg-Pb g^{-1})	*Extractable* (μg-Pb g^{-1})	*Foliage* (μg-Pb g^{-1})	*Foliage* (μg-As kg^{-1})
Biomass Harvesting							
SOH	33±2	9.4±0.6	0.17±0.01	9.45±0.79	0.24±0.01	0.17±0.01	19±3
AAH	33±2	8.9±0.6	0.18±0.01	9.53±0.58	0.17±0.01	0.18±0.01	21±3
BAH	30±2	8.4±0.6	0.16±0.01	7.59±0.45	0.17±0.01	0.16±0.01	15±3
WTH	26±2	7.6±0.6	0.17±0.01	6.76±0.24 *	0.16±0.01	0.07±0.01	21±3
Compensation Methods							
Control	31±1	8.2±0.5	0.16±0.01	8.20±0.39	0.20±0.01	0.16±0.01	16±2
Phosphorus	28±1 *	8.2±0.4	0.18±0.01 (*)	8.24±0.39	0.18±0.01	0.18±0.01 (*)	20±4
Ash	32±1	9.4±0.7	0.17±0.01	8.57±0.39	0.17±0.01	0.17±0.01	20±4

928
929 "*Soil*" = soil substrate (total content of Cd and Pb); "*Extractable*" = soil extractable with 1M NH_4-NO_3 extractant; "*Foliage*" = tree foliage (*i.e.* pine needles) content. It should
930 be noted that the units used differ among Cd variables.
931 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.
932 Treatments with an asterisk differ at $P<0.05$ (for P value < 0.10, the symbol is in brackets) from their reference treatment (SOH or Control, respectively; see Methods).
933 Total soil content of nickel (Ni) and thallium (Tl) are not presented. No significant differences were observed for these variables (Ni=1.49 μg g^{-1}; Tl=0.11 μg g^{-1}).
934
935
936