Single cell RNA sequencing of the *Strongylocentrotus purpuratus* larva reveals the blueprint of major cell types and nervous system of a non-chordate deuterostome

Periklis Paganos[1], Danila Voronov[1], Jacob Musser[2], Detlev Arendt[2] and Maria I. Arnone[1*]

[1]Stazione Zoologica Anton Dohrn, Department of Biology and Evolution of Marine Organisms, Villa Comunale, 80121, Naples, Italy

[2]European Molecular Biology Laboratory, Developmental Biology Unit, Meyerhofstrasse 1, 69117 Heidelberg, Germany

*corresponding author: <u>miarnone@szn.it</u>

Abstract

Identifying the molecular fingerprint of organismal cell types is key for understanding their function and evolution. Here, we use single cell RNA sequencing (scRNA-seq) to survey the cell types of the sea urchin early pluteus larva, representing an important developmental transition from non-feeding to feeding larva. We identify 21 distinct cell clusters, representing cells of the digestive, skeletal, immune, and nervous systems. Further subclustering of these reveal a highly detailed portrait of cell diversity across the larva, including the identification of neuronal cell types. We then validate important gene regulatory networks driving sea urchin development and reveal new domains of activity within the larval body. Focusing on neurons that co-express *Pdx-1* and *Brn1/2/4*, we identify an unprecedented number of genes shared by this population of neurons in sea urchin and vertebrate endocrine pancreatic cells. Using differential expression results from Pdx-1 knockdown experiments, we show that Pdx1 is necessary for the acquisition of the neuronal identity of these cells. We hypothesize that a network similar to the one orchestrated by Pdx1 in the sea urchin neurons was active in an ancestral cell type and then inherited by neuronal and pancreatic developmental lineages in sea urchins and vertebrates.

Introduction

Multicellular organisms consist of numerous cell types, specialized in performing different tasks that guide growth and survival. During embryonic development, cells go through rounds of proliferation, specification and differentiation into cell types with distinct functions. The information for this developmental diversification lies in the genome and the spatio-temporal expression of regulatory genes that specifies the molecular fingerprint of a given cell type (Fu et al., 2017).

The identity of each cell type is established, controlled and maintained by distinct Gene Regulatory Networks (GRNs). GRNs are logical maps of the regulatory inputs and outputs active in a cell at a given place and time, and are enacted by transcription factors, signaling molecules and terminal differentiation genes (Davidson et al., 2003, Davidson

and Erwin, 2006). GRNs have been studied in a variety of organisms ranging from plants to animals in order to analyze the gene interactions at a specific time and place during the life of an organism (Krouk et al., 2013), and have been used for understanding the relationship between genome, development and evolution (Davidson and Erwin, 2006). Therefore, understanding the genetic mechanisms that provide cell types with a specific identity, and the conservation of this identity across animal taxa, is essential for understanding cell type function and evolutionary history (Arendt, 2008, Arnone et al., 2016).

Until recently, most approaches for comparing cell types relied on the description of distinct morphological features, that is linked to the functionality of a given cell type, the identification of molecular markers, perturbation of gene expression and fate mapping. However, technological advances in microfluidics and nucleic acid barcoding now allow the high-through-put recognition of an organism's cell types at a single cell level. In particular, single cell RNA sequencing (scRNA-seq) technology, developed during the last decade, is a powerful method used to unravel the transcriptional content of individual cells, resulting in the identification of distinct cell types in an unbiased manner (Tang et al., 2009, Klein et al., 2015). ScRNA-seq involves dissociation of an organism, organ or tissue into single cells, isolation and capture of the single cells into droplets, specific barcoding of individual mRNAs, and sequencing of transcriptomic content of each cell. Computational analysis can then identify putative cell type families by clustering cells with similar transcriptional profiles. Further analysis of such cell clusters can lead to the identification of distinct cell types.

Within deuterostomes, echinoderms are a member of the phylogenetic sister group to chordates, making them an ideal model for understanding the origin and diversification of deuterostome and chordate cell types (Arnone et al., 2015). Sea urchin embryos and larvae have also been extensively used to unravel the general mechanisms of cell type specification and differentiation during development (Cameron and Davidson, 1991, Davidson et al., 1998, McClay, 2011, Lyons et al., 2012, McClay et al., 2020). The main reason for this lies in the ease with which different cell types and biological processes can be observed in the optically transparent embryos and larvae. Among the most well-studied sea urchin cell types are those comprising the nervous (Bisgrove and Burke, 1987, Burke et al., 2006a, McClay et al., 2018), immune (Rast et al., 2006, Ho et al., 2017) and digestive systems (Annunziata et al., 2014, Annunziata and Arnone, 2014, Perillo and Arnone, 2014, Perillo et al., 2016), and of both musculature (Andrikou et al., 2013, Andrikou et al., 2015) and skeleton (Okazaki, 1965, Duloquin et al., 2007, Rafiq et al., 2012, Sun and Ettensohn, 2017). For these, the developmental origins and gene regulatory wiring has been described in great detail, making the sea urchin an ideal model for GRN comparative analyses in development and evolution (Cary et al., 2020).

Here we take advantage of the detailed characterization of the sea urchin cell types performed over the years, the available cell type specific molecular markers, and the ease with which the sea urchin larvae are dissociated into single cells, to perform scRNA-seq and generate a comprehensive atlas of sea urchin larval cell type families. Our findings suggest that the larva consists of 21 genetically-distinct cell clusters, representing distinct cell type families (Shekhar and Menon, 2019), which we validate using fluorescent *in situ* hybridization (FISH) and immunohistochemistry (IHC). Based on previous studies tracing

developmental lineage, we assign cell type families to the specific embryonic germ layers
they derived from. Comparing the transcription factor content of the grouped cell clusters
reveals that most transcription factors are expressed pleiotropically independently of their
germ layer origins, yet tend to be cell type family-specific within a germ layer. In addition,
we illustrate how single-cell data complement and validate previously studied GRNs, and
also reveal novel cellular domains where these GRNs are likely also activated. Lastly, we
investigate neuronal diversity in the sea urchin larva, identifying 12 distinct neuronal cell
types. Among these, we recover a unique neurosecretory type controlled by *Sp-Pdx1* and
Sp-Brn1/2/4 exhibiting a pancreatic-like gene expression signature (Perillo et al., 2018).
Our results confirm and extend this pancreatic-like signature, suggesting that an ancestral
neuron in early deuterostomes may have given rise to the endocrine cells in the vertebrate
pancreas. Supporting this, knockdown of *Sp-Pdx1* shows it is necessary for differentiation
of this pancreatic-like neuronal endocrine population, indicating it has an evolutionary
conserved role as a mediator of endocrine fate.

Results

Building a cell type atlas of the sea urchin larva with single-cell transcriptomics

Sea urchin early pluteus larvae were cultured and collected at 3 dpf. We performed single-
cell RNA sequencing on six samples from four independent biological replicates.
Individual samples were dissociated into single cells using a gentle enzyme-free
dissociation protocol and using the 10x Chromium scRNA-seq system (Figure 1A). In
total, transcriptomes from 19,699 cells were included in the final analysis. To identify sea
urchin larval cell types, we used Louvain graph clustering as implemented in the Seurat
pipeline (see methods). This revealed 21 genetically-distinct cell clusters (Figure 1B,
supplement 1A and B), each representing an individual cell type or set of closely related
cell types in the early pluteus larva.

Next, we set out to explore the identity of our initial 21 cell clusters. We first assigned
preliminary identities to each cluster based on the expression of previously described cell
type markers, benefiting from the unique and rich knowledge on sea urchin
developmental lineages: ciliary band (*Btub2*) (Harlow and Nemer, 1987), apical plate
(*Hbn*) (Burke et al., 2006a), aboral ectoderm (*Spec2a*) (Yuh et al., 2001), lower oral
ectoderm (*Bra*) (Wei et al., 2012), upper oral ectoderm (*Gsc*) (Wei et al., 2012), neurons
(*SynB*) (Burke et al., 2006a), esophageal muscles (*Mhc*) (Andrikou et al., 2013), coelomic
pouches (*Nan2*) (Juliano et al., 2010), blastocoelar cells (185/333) (Ho et al., 2017),
immune cells (*Gcm*) (Materna et al., 2013), skeleton (*Msp130*) (Harkey et al., 1992), anus
(*Hox11/13b*), intestine (*Cdx*), pyloric sphincter (*Pdx-1*), different stomach domains (*Chp,*
ManrC1a, Endo16) (Annunziata and Arnone, 2014), exocrine pancreas-like domain
(*Ptf1a*) (Perillo et al., 2016), cardiac sphincter (*Trop1*) (Yaguchi et al., 2017) and
esophagus (*Brn1/2/4*) (Cole and Arnone, 2009). Further, we grouped the cell clusters
according to their embryonic germ layer origin (Figure 1C) using knowledge from previous
lineage tracing experiments (Angerer and Davidson, 1984, Cameron et al., 1987).

Figure 1. Cell type family atlas of the 3 dpf *S. purpuratus* larva. A) Single-cell RNA
sequencing pipeline from gamete fertilization to computational analysis. **B)** UMAP showing 3 dpf
larval cells colored by their assignment to the initial set of 21 distinct cell clusters. **C)** UMAP with
cells colored by germ layer of origin: endoderm (yellow), mesoderm (red), and ectoderm (blue).

D) Dotplot of gene markers specific to cell clusters. **E)** Illustration depicting location of cell type families on different larval domains. Color-code is the same as in Figure 1B.

Figure 2. Validation of scRNA-seq predictions and novel expression domains. A) FISH of *S. purpuratus* 3 dpf larvae with antisense probes for *Sp-FbsL_2* (A1), *Sp-Chrna9_4* (A2), *Sp-Frizz5/8* (A3), *Sp-FoxABL* (A4), *Sp-Bra* (A5), *SPU_006199* (A6), *Sp-FcolI/II/III* (A7), *Sp-*

Hypp_1249 (A8), *Sp-Hypp_2386* (A9), *Sp-Mlckb* (A10), *Sp-MsxI* (A11), *SPU_008104* (A12), *Sp-*
Ahrl (A13), *Sp-Serp2; Sp-Serp3* (A14), *Sp-Hnf4* (A15), *Sp-Cyp2L42* (A16), *Sp-Rfxc1l* (A17), *Sp-*
Pdx1 (A17) and *Sp-Cdx* (A18). Color-code indicates germ layer embryonic origin: endoderm
(yellow), mesoderm (red), ectoderm (blue). Immunofluorescent detection of acetylated tubulin in
ciliary band (green). **B)** Dotplot of *Sp-Sip1*, *Sp-SoxC*, *Sp-Hbn* and *Sp-Fgf9/16/20* expression
showing previously described and novel expression domains. **C)** FISH of *S. purpuratus* 3 dpf
larvae with antisense probes for *Sp-SoxC* (C1-C2), *Sp-Hbn* (C4-C5) and *Sp-Fgf9/16/20* (C7-C8).
Illustrations depicting all the expression domains of *Sp-SoxC* (C3), *Sp-Hbn* (C6) and *Sp-*
Fgf9/16/20 (C9). Illustrations depicting all the expression domains of *Sp-SoxC* (C3), *Sp-Hbn* (C6)
and *Sp-Fgf9/16/20* (C9). Nuclei are labelled with DAPI (in blue). All images are stacks of merged
confocal Z sections. **D)** FISH of *S. purpuratus* 3 dpf larvae with antisense probes for *Sp-SoxC*
and Sp-Fgf9/16/20 (D1-D4), for *Sp-Hbn* combined with immunohistochemical detection for the
skeletal cells marker Msp130 (D5-D9) and *Sp-Fgf9/16/20* with immunohistochemical detection for
the midgut and hindgut protein Endo1 (D10-D13). A, Anus; Cs, Cardiac sphincter; I, Intestine; M,
Mouth; PMCs, Primary mesenchyme cells; St, Stomach.

To validate these identities, we identified all genes expressed in each cell type, totaling
in 15,578 WHL genes (transcriptome models) and 12,924 genes (SPU gene models)
(Supplementary file 1), and performed *in situ* hybridization on a selected set of these
together in combination with previously described markers. Based on this, we mapped 20
of the 21 clusters to distinct larval domains and confirmed their identity (Figure 1E).
Notably, the resulting expression patterns validated the initial predictions (Figure 2- Figure
supplement 2), verifying the high quality of the single-cell dataset. Importantly, this
approach identified various new markers for each cell type family, including *Sp-FbsL_2*
(ciliary band; Figure 2A1*), Sp-hypp_2386* (skeletal cells; Figure 2A9), and *Sp-Serp2/3*
(exocrine pancreas-like cells; Figure 2A14). The 21st cluster, which had a poorly-defined
molecular signature as judged by the low number and expression level of total and marker
genes combined with the lack of specific localization, likely represents not fully
differentiated cells (Figure 1- Figure supplement D & F) and we refer to this cluster as
undefined.

Our scRNA-seq analysis and *in situ* hybridization protocol unraveled novel expression
domains for several previously described cell type markers. For instance, the transcription
factors *Sp-SoxC* and *Sp-Hbn*, previously described in early neuronal specification
(Garner et al., 2016, Wei et al., 2016, Yaguchi et al., 2016), were predicted by our scRNA-
seq analysis to also be expressed in skeletal cells (Figure 2B &,). Likewise, the FGF
signaling ligand, *Sp-Fgf9/16/20*, is known to be involved in skeletal formation and is
expressed in specific populations of PMCs (Adomako-Ankomah and Ettensohn, 2014).
ScRNA-seq indicates it is also expressed in oral ectoderm, cardiac sphincter, intestine
and anus (Figure 2B). Using *in situ* hybridization and immunohistochemical detection
methods we confirmed the predictions and we found *Sp-SoxC* expressed in *Sp-*
Fgf9/16/20 positive PMCs and *Sp-Hbn* expressed in skeletal cells located in the vertex
(Figures 2C1-3, 2C4-6 2D1-4 & 2D5-9). Moreover, we found *Sp-Fgf9/16/20* transcripts
localized in distinct gut domains corresponding to the cardiac sphincter, intestinal and
anal regions (Figure 2C7-9, 2D10-13).

We compared the limits of detection by *in situ* hybridization versus single cell RNA
sequencing, using the coelomic pouch cell cluster as a case study. The coelomic pouches
are derived from the mesoderm and the left coelomic pouch contributes to the formation

of the rudiment and juvenile sea urchin after metamorphosis (Strathmann, 1987, Smith et
al., 2008). The formation of the coelomic pouches is complex, and includes contributions
from the small micromeres, and a mesodermal cell population that is set aside during
early development (Pehrson and Cohen, 1986, Strathmann, 1987). Previous attempts to
characterize this population had involved screening of genes active in germline
determination and maintenance in other species and revealed that, while some germ line-
specific transcripts and proteins were found exclusively expressed in the small
micromeres and the coelomic pouch of the sea urchin embryo (Juliano et al., 2006), the
majority of the genes tested by *in situ* hybridization were not enriched in this cell type.
Interestingly, plotting the Juliano and coauthors' gene list, alongside previously described
coelomic pouch specific genes (Luo and Su, 2012, Martik and McClay, 2015), they were
all found in our analysis to be expressed in the same cell cluster (Figure 2- Figure
supplement 3). This suggests a higher sensitivity of single cell RNA sequencing
compared to the *in situ* hybridization, adding crucial missing information on the molecular
fingerprint of a complex cell type.

Lastly, to determine which cells in the larva were undergoing active proliferation, we
plotted expression of cell division markers in sea urchin, including *pcna*, DNA
polymerases, DNA ligases, condensins, and centromere proteins (Perillo et al., 2020).
The majority of cell proliferation genes were found to be enriched in the ciliary band, apical
plate, coelomic pouch, immune and skeletal cell clusters (Figure 2- Figure supplement
4A). We also observed Cdk genes enriched in several endodermally derived cell
populations (Figure 2-supplement 4A). Validating this, we observed S-phase cells in
ciliary band, apical plate, oral ectoderm, endodermal and skeletal cells using Edu pulse
labelling (Figure 2- Figure supplement 4B). In contrast, we did not observe Edu
fluorescence in cell clusters that lacked expression of proliferation markers, such as the
aboral ectoderm, suggesting that indeed they may not be proliferating.

Shared lineage information of the larval cell type families

We next compared transcription factor profiles of early larval cell type families. In general,
cell populations derived from the same germ layer shared more factors with each other
than with cell populations from other germ layers (Figure 3, Figure 3-Figure supplement
1), consistent with the finding that cell type expression programs often retain information
about their developmental lineage (Sladitschek et al., 2020). Further analysis however
revealed very few transcription factors specific to cells from a single germ layer,
contrasting with many regulators expressed in derivatives of more than one germ layer
(Figure 3A), and nearly one third (n = 187) shared by cell type families from all three
layers. Unexpectedly, mesodermal cell populations share expression of more
transcription factors with ectodermal than with endodermal cell populations, even though
they are more closely linked to endodermal lineage in development. We also noted that
cell type families derived from the same germ layer share up to one third of transcription
factors, while the majority are cell type-specific (Figure 3B, 3D & 3E). In general,
neighboring cell populations and those with common developmental origins share a larger
number of TFs (Figure 3D), compared to cell populations with different developmental
histories (Figure 3C).

Figure 3. Regulatory states of the 3 dpf *S. purpuratus* larva. A) Comparison of the
transcription factor content per germ layer. Venn diagram showing the shared and unique
transcription factors per germ layer. Ectodermally derived cell type families are shown in blue,
mesodermally derived in red, and endodermally derived in yellow. **B)** Comparison of the
transcription factor content across mesodermal lineages and cell type families. Venn diagram
showing the shared and unique transcription factors per comparison. **C)** Transcription factor
content comparison of pyloric sphincter (endodermally derived) and skeletal cells (mesodermally
derived), used as a negative control of our comparison. **D)** Comparison of the transcription factor
content per endodermal lineage and endodermally derived cell type families. Venn diagram
showing the shared and unique transcription factors per comparison. **E)** TF signature comparison
of ectodermally derived cell type families. Venn diagram showing the shared and unique
transcription factors per comparison. Cartoons indicated the relative position of each cell type
family/lineage. Mesodermal cell type families/lineages are shown in shades of red, endodermal
ones in shades of yellow and endodermal ones in shades of blue.

To further characterize the regulatory profile of larval cell type families we set out to
identify the expression profiles of members of major transcription factor families (Figure
4 and Figure 4 supplement 1). In *S. purpuratus*, most homeobox transcription factors were
previously found expressed at the gastrula stage (2 dpf), with several members
expressed in domains derived from all three germ layers (Howard-Ashby et al., 2006).
Our single cell analysis, although at a later developmental time point, supports these
findings, and further refines our understanding of their expression to specific cell type
families. In the early pluteus larva, most homeobox class transcription factors are
enriched in ectodermally derived cell populations, such as the apical plate and neurons.
In contrast, ANTP Class and HNF class transcription factors are enriched in endodermal
derivatives (Figure 4). Other major transcription factor families, such as the Forkhead,
Ets, and Zinc-finger families, members of which are expressed throughout sea urchin
embryogenesis (Tu et al., 2006, Rizzo et al., 2006, Materna et al., 2006), are also
expressed across a spectrum of cell populations. Forkhead and zinc-finger transcription
factors are highly expressed in specific cell type families of all three germ layer
derivatives, whereas Ets family TFs are enriched in ectodermal and mesodermal
derivatives (Figure 4).

The active regulatory state of a given cell type family is an immediate consequence of the
gene regulatory network active at this time point. Previous research in sea urchin has
described in detail many regulatory networks active during embryonic and larval
development. Our scRNA-seq data is broadly consistent with previous studies, yet also
identifies new domains and cell populations in which these regulatory networks may be
active. For instance, we plotted all transcription factors known to be active in specifying
coelomic pouch cells (Martik and McClay, 2015). Our data confirmed their co-expression
in coelomic pouch, but also revealed their co-expression in the apical plate (Figure 5A).
Similarly, plotting genes involved in the aboral ectoderm gene regulatory network (Ben-
Tabou de-Leon et al., 2013), we found all genes in both the aboral ectoderm cluster as
well as in the apical plate cells (Figure 5B). On the other hand, plotting members of the
pre-gastrula skeletogenic mesoderm regulatory network (Figure 5C) revealed most were
still active in the pluteus larva and specific to skeletal cells (Oliveri et al., 2008, Shashikant
and Ettensohn, 2019, Khor et al., 2019, Ettensohn, 2020).

Figure 4. Localization of major transcription factor family members. Dotplot showing the average scaled expression of members of the Homeobox, Forkhead and Ets transcription factor families. The developmental origins of each cell type family are shown in blue for ectodermally derived, red for mesodermally derived and yellow for endodermally derived ones.

Figure 5. Validation of preexisting GRNs and putative novel function of specific gene
regulatory modules. A) Dotplot showing the mRNA localization of genes involved in the homing
of small micromeres to the coelomic pouch and novel apical plate domain. **B)** Dotplot of aboral
ectoderm regulatory module genes showing novel apical plate expression. **C)** Pre-gastrula gene
regulatory network enriched in skeletal cells of the sea urchin pluteus larva. Asterisks indicate
larval genes involved in biomineralization, putative members of this GRN.

These findings illustrate the immediate benefit of our dataset to drastically expand our
knowledge of larval regulatory networks. Finally, our scRNA-seq recreates a nearly
identical 3 dpf endoderm expression pattern atlas as that published previously by our
group using more traditional methods (Annunziata et al., 2014), providing additional
information on each gene's average expression level and the percentage of cells
expressing each marker (Figure 5- Figure supplement 1).
293
Unravelling the neuronal diversity and molecular signature of the nervous system

The classification of neuronal cell type diversity is an important step for unravelling the
evolution and function of the nervous system. The sea urchin free swimming larva is
equipped with a nervous system consisting of interconnected ganglia (Burke et al., 2006a)
that allows the animal to respond to environmental stimuli and coordinate its swimming
behavior (Soliman, 1983, Katow et al., 2010). Several neuronal types, including apical
and ciliary band neurons, as well as neurons along the digestive tube, have been
previously identified and their specification described in detail (Burke et al., 2006a, Burke
et al., 2006b, Wei et al., 2009, Wei et al., 2011, Garner et al., 2016, Wei et al., 2016,
McClay et al., 2018, Perillo et al., 2018, Wood et al., 2018).

Our initial clustering analysis resolved single clusters for neuronal cells, as well as for
PMCs and immune cells. However, expression of known markers suggested the
presence of distinct subclusters in each of these cell type family groups. In order to
investigate this, we independently performed subclustering and re-analysis of the
neuronal, immune, and PMC cells. Subclustering of each of these initial major clusters
revealed 12 neuronal, 8 immune, and 5 PMC subclusters, each likely representing distinct
cell types (Figure 6A, Figure 6- Figure Supplement 1 & 2). Two of the immune subclusters
expressed polyketide synthase 1 (*Sp-Pks1*), suggesting these represent sea urchin
pigment cell populations (Calestani and Rogers, 2010). We also found a subcluster of
immune cells that expresses the membrane attack complex/perforin family gene (*Sp-*
MacpfA2), suggesting this corresponds to immune system globular cells (Figure 6-
supplement 2). Notably, our finding of 5 PMC subclusters corroborates previous reports
showing five distinct groups of PMC cells along the syncytium (Sun and Ettensohn, 2014)
(Figure 6- Figure Supplement 1).
To annotate the 12 neuronal cell types revealed via subclustering we took advantage of
the extensive previous work investigating neurogenesis and neuronal differentiation in
sea urchin. Plotting neuronal markers, we resolved unique molecular signatures for each
subcluster and assigned each a putative identity and location in the larva (Figures 6B-D).
To validate this, we then conducted in-situ hybridization experiments for gene markers
labeling these specific neuronal populations (Figure 6D), including genes encoding
transcription factors (*SoxC*, *Delta*, *Ngn*, *Prox1*, *Isl*, *Hbn*, *SoxB2*, *Otx*, *NeuroD1*, *Six3*), and
members of neurotransmitter (*Ddc*, *Nacha6*, *Chrna9_4*, *Tph*), and neuropeptidergic
signaling pathways (*An*, *Salmfap*, *Trh*).

Figure 6. Neuronal complexity of the 3 dpf *S. purpuratus* larva. A) (From left to right and top
to bottom) UMAP highlighting the neurons cluster (green), immunohistochemical detection for the
paneuronal sea urchin marker 1E11 (green), UMAP showing the 12 distinct neuronal subclusters.
B) Schematic representation of the 3 dpf pluteus larva showing the localization of neuronal
subclusters (colors as in A). **C)** Dotplot of signaling molecules, transcription factors, and
neurotransmitters involved in sea urchin neuronal function and neurogenesis (colors as in A). **D)**
FISH of *S. purpuratus* 3 dpf larvae with antisense probes for the neuronal genes *Sp-Delta* (D1),
Sp-SoxC (D2), *Sp-Brn1/2/4* (D3), *Sp-Ddc* (D4), *Sp-Ngn* (D5), *Sp-Prox1* (D6), *Sp-Nacha6* (D7),
Sp-Isl (D8), *Sp-An* (D8 and D14), *Sp-Chrna9_4* (D9), *Sp-Hbn* (D10), *Sp-SoxB2* (D11), *Sp-Otx*
(D12), *Sp-Tph* (A13), *Sp-Salmfap* (D14), *Sp-NeuroD1*(D15), *Sp-Six3* (D16), *Sp-Trh* (D17), *Sp-Kp*
(D17) and *Sp-Rhox3* (D18). FISH shown in figures D1-3 are paired with immunohistochemical
detection of the neuropeptide Sp-An. Nuclei are labelled with DAPI (in blue). All images are stacks
of merged confocal Z sections. A, anus; M, mouth.

The sea urchin larva neuronal differentiation proceeds via stepwise differentiation,
including transient expression of the Notch ligand Delta, followed by expression of the
transcription factors *SoxC* and *Brn1/2/4* (Garner et al., 2016). During the final stages of
neurogenesis, the transcription factors *Sip1*, *Z167*, *Ngn* and *Otp* regulate differentiation
of diverse neuronal populations, including apical and ciliary band neurons (Wei et al.,
2016, McClay et al., 2018). In our data, we observed *Sp-Delta*, *Sp-SoxC* and *Sp-Brn1/2/4,*
as well different combinations of the transcription factors mentioned above, co-localize in
three neuronal populations (subclusters 1, 2 and 4), indicating neuronal differentiation is
taking place in those three subclusters (Figure 6C). Interestingly, in one of these
populations (subcluster 2) we found expression of the transcription factors *Sp-Rx*, *Sp-*
Hbn (Figure 6D10) and *Sp-Six3* (Figure 6D16), which are known to be expressed in the
periphery of the larva's apical domain (Burke et al., 2006a, Wei et al., 2009). This
suggests that this population is located in the periphery of the apical plate and not within
the apical organ. In the apical domain, we also detected a subcluster (number 6), which
coexpress *Sp-Trh* and *Sp-Salmfap* neuropeptides (Wood et al., 2018), as well as *Sp-Kp*
(Kissepeptin) (Figure 6D17).

In total, we identified three neuronal subclusters located in the apical domain (subclusters
2, 3 and 4) of the larva that express Tryptophan hydroxylase (*Tph*), which encodes a key
enzyme in the serotonin biosynthesis pathway, suggesting these represent serotonergic
neurons in the larva. Within the ciliary band, which comprises the larva's peripheral
nervous system (Slota et al., 2020), we identified two distinct cholinergic subclusters (7
and 8) expressing the enzyme involved in acetylcholine biosynthesis (*Sp-Chat*) (Figure
6C), one of which (subcluster 8) expresses also two nicotinic acetylcholine receptors
(*Nacha6*, *Chrna9*). Moreover, we identified a neuronal subcluster in close proximity with
the ciliary band, which corresponds to the lateral and post-oral neurons (subcluster 9).
This population has been previously characterized by our group and was found to co-
express *Sp-Pdx1*, *Sp-Brn1/2/4*, and the neuropeptide Sp-An (Perillo et al., 2018). Using
gene markers that mark differentiated neurons expressed in the rim of the larva's mouth,
including *Sp-Nkx3.2* (Wei et al., 2011), *Sp-Isl* (Perillo et al., 2018), the neuropeptide *Sp-*
Salmfap (Wood et al., 2018), and the enzyme *tyrosine hydroxylase* (*Sp-Th*) involved in
the dopaminergic pathway, we identified two distinct mouth neurons subtypes
(subclusters 10 and 11; Figure 6C). Lastly, we found one neuronal population that, based
on its molecular signature, and in particular the co-presence of *Sp-Fgfr1* and *Sp-*
Fgf9/16/20 (Figure 6-Figure supplement 3)*,* could correspond to neurons associated to

the cardiac sphincter within the esophageal endoderm (subcluster 12). Overall, our
subclustering analysis increases the resolution of the different neuronal subtypes present
at this developmental stage, describing new neuronal subtypes and providing novel
markers and gene candidates for future studies of these cell types and their gene
regulatory networks.

Characterizing a neuroendocrine neuronal population controlled by Pdx-1

Previous studies from our group suggested that the nervous system of the sea urchin
larva displays a strong pre-pancreatic signature, with neurons expressing genes that are
involved in endocrine cell differentiation in the vertebrate pancreas (Perillo et al., 2018).
To investigate this, we focused on the post-oral and lateral neuron subcluster, which co-
expresses *Sp-Pdx1*, *Sp-Brn1/2/4* and *Sp-An* (Figure 6). Double immunohistochemical
staining of the neuronal marker 1E11 and Sp-An shows that these neurons lie in close
proximity to the ciliary band, and project axons towards both the apical plate and ciliary
band (Figure 7 A1-A3). Double FISH of *Sp-An* and *Sp-FbsL_2,* a ciliary band marker
revealed by this study*,* as well as double IHC of Sp-An and acetylated tubulin (labeling
cilia)*,* further indicate their distribution relative to the ciliary band (Figure 7 A4-A6).
Moreover, we found that the post-oral Sp-An neurons are found in close proximity to cells
of both ciliary band subclusters (Figure 7 A7-A9), and project axons to the cell bodies of
the *Sp-Prox1* positive neurons (Figure 7 A9). We also observed close proximity between
An positive neurons in the post oral arms with the cells of the ventral-lateral cluster of
PMCs (Figure 7 A10), and with immune globular cells (Figure 7 A11). Next, we set out to
validate whether novel genes predicted by our single cell analysis to be expressed in this
neuronal population can be validated *in vivo*. Among the genes predicted to be expressed
in this population are the transcription factors *Sp-Nk1* and *Sp-Otp,* as well as the
catecholaminergic and cholinergic neuronal markers Sp-Th and Sp-Chat (Slota and
McClay, 2018), respectively. Double fluorescent *in situ* hybridization of Sp-Nk1 and the
neuropeptide Sp-An, as well as fluorescent *in situ* hybridization of *Sp-Otp* combined with
the immunohistochemical detection of Sp-An, reveal co-localization of these three genes
in the post-oral and lateral neuronal population (Figure 7 A13-A14), verifying the single-
cell data. Additionally, double immunostainings of the anti-Sp-An with anti-Th (Figure
7A9) and anti-Chat antibodies suggest that these two key enzymes, involved in different
neurotransmitter biosynthesis pathways, are co-produced in the Sp-*Pdx1/Sp-Brn1/2/4*
neurons (Figure 7 A12 and A15).

The vertebrate orthologue of *Sp-Pdx1* is essential for pancreas development, β-cell
differentiation, and maintaining mature β-cell function (Kaneto et al., 2007). As previously
described by our group, knockdown of the pancreatic transcription factor Sp-Pdx1 results
in severe downregulation of the Sp-An neuropeptide, compromising the neuroendocrine
fate of this neuronal type (Perillo et al., 2018). To further characterize the *Sp-Pdx1/Sp-*
Brn1/2/4 neuronal population, we performed a comprehensive analysis of genes involved
in pancreatic development and β-cell differentiation, as well as gene markers related to
neuroendocrine fate. We identified a total of 20 genes, all involved in the formation and
proper function of vertebrate endocrine pancreas, which are differentially enriched in the
Sp-Pdx1/Sp-Brn1/2/4 neurons (Figure 7B).

Figure 7. ScRNA-seq reveals a Pdx-1 dependent neuroendocrine cell type. **A)** Molecular

characterization of a *Sp-Pdx1/Sp-Brn1/2/4* double positive neuronal population. A1) Double FISH of *S. purpuratus* 3 dpf larvae with specific antisense probes for *Sp-Pdx1* and *Sp-An*. A2) Double immunohistochemical detection of the Sp-An and Synaptotagmin (1E11) proteins. A3) Close up caption of the Sp-An PON neurons shown in A2. A4) Double FISH of *S. purpuratus* 3 dpf larvae with specific antisense probes for *Sp-FbsL_2* and Sp-An. Double immunohistochemical detection of the Sp-An and acetylated tubulin proteins. A6) Close up caption of the Sp-An PON neurons shown in A5. A7) FISH of *S. purpuratus* 3 dpf larvae with a specific antisense probe for *Sp-Chrna9_4* paired with immunodetection of Sp-An. A8) FISH of *S. purpuratus* 3 dpf larvae with a specific antisense probe for *Sp-Prox1* paired with immunohistochemical detection of Sp-An. A9) Close up caption of the Sp-An PON neurons shown in A8. A10) Double immunohistochemical staining for the neuropeptide Sp-An and the skeletal cells marker Msp130. A11) FISH of *S. purpuratus* 3 dpf larvae with a specific antisense probe for *Sp-MacpfA2* paired with immunohistochemical detection of Sp-An. A12) Double immunohistochemical staining for the neuropeptide Sp-An and the enzyme Sp-TH. A13) Double FISH of *S. purpuratus* 3 dpf larvae with specific antisense probes for *Sp-Nk1* and Sp-An. A14) FISH of *S. purpuratus* 3 dpf larvae with a specific antisense probe for *Sp-Otp* paired with immunohistochemical detection of Sp-An. Double immunohistochemical staining for the neuropeptide Sp-An and the enzyme Sp-Chat. Nuclei are labelled with DAPI (in blue). All images are stacks of merged confocal Z sections. LGN, lateral ganglion neurons; M, Mouth; PON, Post-oral neurons. **B)** Dotplot of genes important in endocrine pancreas differentiation and function in vertebrates. **C)** Bar plot of selected Sp-Pdx1 target genes in the *Sp-Pdx1/Sp-Brn1/2/4* positive population as revealed by differential RNA sequencing analysis of *Sp-Pdx1* knockdown larvae. **D)** Schematic representation of the Sp-Pdx1/Sp-Brn1/2/4 neurons genetic wiring as revealed by the combination of scRNA-seq and differential RNA-seq analysis after *Sp-Pdx1* knockdown, highlighting an important role of *Sp-Pdx1* in the differentiation of this cell type.

Next, we intersected our scRNA-seq with bulk RNA sequencing data derived from Pdx1 morphants assayed at the same developmental stage in a previous study (Annunziata and Arnone, 2014). By coupling knowledge of cell type-specific expression programs with genes differentially expressed in the Pdx1 knockdown mutants, we were able to identify and refine likely gene targets specific to the *Sp-Pdx1/Sp-Brn1/2/4* neuronal cell type. In total, we found 249 genes belonging to the *Sp-Pdx1/Sp-Brn1/2/4* neuron subcluster (9) that were differentially expressed in the *Pdx1* knockdown dataset, with 65% of the targets being downregulated (Figure 7C). Among the downregulated genes, we found key transcription factors involved in neuronal differentiation, including *Sp-Brn1/2/4* and *Sp-Otp*, as well as terminal differentiation genes important in neuronal signaling, such as *Sp-An*, *Sp-Ngfffap*, *Sp-Th* and *Sp-Chat* (Figure 7C).

Discussion

Cell type identity is determined by the differential use of genomic information among cells. Unraveling the distinct transcriptomic signatures of cell types yields valuable insight into their function, as well as their evolutionary and developmental origins. In recent years, single cell transcriptomics has emerged as a powerful and unbiased approach for characterizing cell type diversity across a wide variety of animal taxa, with studies spanning from insects (Davie et al., 2018, Severo et al., 2018, Cho et al., 2020), to cnidarians (Sebe-Pedros et al., 2018, Siabert et al. 2019), planarians (Swapna et al. 2018, Zeng et al. 2018) and sea squirts (Sharma et al., 2019, Cao et al., 2019), as well as

vertebrates such as zebrafish (Wagner et al., 2018, Chestnut et al., 2020), mice
(Nestorowa et al., 2016, Jung et al., 2019, Ximerakis et al., 2019, Yu et al., 2019, Qi et
al., 2020) and humans (Yu et al., 2019, Qi et al., 2020, Esaulova et al., 2020, Zhao et al.,
2020).

The sea urchin embryo has served as a valuable model for understanding cell type
molecular specification and differentiation via gene regulatory networks. Despite this,
knowledge of later stages of development, including larval cell types, is limited. Here we
used single cell RNA sequencing to generate a detailed atlas of cell types of the early
pluteus larva, and to unravel the neuronal diversity at this critical stage that marks the
end of embryogenesis and the beginning of the larval life cycle.

Cellular diversity of the *S. purpuratus* pluteus larva

Conducting scRNA-seq on isolated *S. purpuratus* early pluteus cells we initially identified
21 genetically distinct cell clusters from 19.699 cells (Figure 1 A), that consist of
derivatives from all three germ layers. Similar ectodermal, mesodermal and endodermal
representatives have also been reported at a single cell resolution in previous studies,
carried out at either earlier developmental stages or different sea urchin species,
highlighting the power of this technique to capture fine and dynamic developmental
processes during sea urchin development (Foster et al. 2020, Massri et al. 2021). The
cells comprising the 21 cell type families are expressing a total of 15,578 WHL genes
(transcriptome models) and 12,924 genes (SPU gene models). Notably, the
computationally identified number of cells per cluster did not correlate well with their
actual distribution in the larva. This could be a bias linked to the dissociation process, as
for instance skeletal structures due to their calcite and syncytial nature are the last to
dissociate and hence cells can get trapped within the debris and thus be underestimated
in our datasets. Nonetheless, the cell type families and cell types identified in our study
include all known larval cell types, suggesting we obtained a sufficient number of cells to
comprehensively survey cell diversity at this developmental stage. Further, the total
number of genes expressed in our data was 15.579, in relatively close agreement to the
16.500 genes expressed at the end of *S. purpuratus* embryogenesis (Tu et al., 2014).
Our results reveal a rich tapestry of cell type families within the early pluteus larva. In
particular, our study reveals unprecedented transcriptional diversity among cells of the
pluteus larva that relates to feeding and digestion, as seen by the large representation of
such clusters in our dataset. This includes two distinct oral ectoderm cell clusters, as well
as distinct cell expression programs for the esophagus, cardiac and pyloric sphincters,
exocrine pancreas-like cells, three distinct stomach cell clusters, intestine and anus. We
also identified the mesodermally derived esophageal muscle cell type that ensures the
proper function of the digestive apparatus by regulating the flow of the water containing
food within the different compartments of the gut.

Cell type family tree reconstruction reveals two well supported groups (Figure 3 – Figure
supplement 1). One group represents the endoderm. It comprises a core of closely related
cell type families (stomach cells 1,2,3, pyloric and cardiac sphincter, and exocrine
pancreas-like cells) and the more distant subgroup of intestinal and anal cells. The
coherence of this group is also apparent from the transcription factor heatmap (Figure 4–
Figure supplement 2), which reveals numerous factors shared between subsets or the

entire endodermal group such as *Klf15* that is known to control lipid and glucose
metabolism in the liver (Oishi and Manabe 2018). The other group represents ectoderm.
It contains apical plate, ciliary band, the aboral and upper oral ectoderm cells, and
neurons. Again, this group manifests in shared transcription factor expression including
Klf2/4, Tbx2/3, and *SoxB1* and *SoxB2*. Unexpectedly, this group firmly comprises some
cell type families that are usually not considered ectodermal, such as the immune cells
and the esophageal cells (Figure 3 – Figure supplement 1). Outside these two groups,
the remaining cell type families do not assemble into a well-supported group. Instead,
they exhibit molecular affinities to either the endodermal or ectodermal clades.

Beyond our initial cell clusters, we used subclustering to uncover diversity among
neuronal, PMC, and immune cells in the pluteus larva. Among immune cells, the presence
of two pigment cell subclusters is in line with findings by Perillo and colleagues that
revealed two such populations (Perillo et al., 2020). However, in our study we found that
only one cluster is *Sp-Gcm* positive, in contrast to their findings which found both of their
clusters expressed *Sp-Gcm*. This difference could be a result of the different approaches
used to identify the different cell types or due to transient expression of *Sp-Gcm* in the
additional pigment cell type.

The larval nervous system has been among the first echinoderm cell types to be
characterized at a molecular level and yet the exact number of neuronal subtypes is still
not clear. Extensive work has been done on identifying the molecular pathways guiding
neuronal specification and the genes active during neuronal differentiation, however this
is limited to describing general neuronal categories. Most of the current information on
the different neuronal types relies on detection of specific neuropeptides,
neurotransmitters and enzymes involved in their biosynthesis. The most recent estimate
of neuronal diversity used neuropeptidergic content to identify seven distinct neuronal
types (Wood et al., 2018). Our study supports and refines this earlier work by providing a
comprehensive and unbiased survey of neuronal diversity in the early pluteus larva of *S.*
purpuratus.

Neurons in *S. purpuratus* arise from two ectodermal neurogenic regions (ciliary band and
apical domain) and from the anteriormost part of the foregut (Garner et al., 2016, McClay
et al., 2018, Wei et al., 2011), which is derived from endoderm (mouth neurons and
esophageal neurons, see Figure 6A). Different neuronal types from these domains arise
at different developmental time points, although by 3 dpf most larval neurons are thought
to be present and patterning diverse larval domains. Notably, our initial clustering analysis
recovered neurons as a single cluster, suggesting they share a common molecular
signature regardless of their developmental origin. Subclustering of these revealed twelve
distinct neuronal populations, which we were able to trace back on the larva. By doing so,
we identified i) one cluster consisting of neurons that apparently are undergoing
differentiation as judged by the combinatorial expression of Delta, Soxc, Brn1/2/4 and
terminal differentiation gene markers, ii) four associated with the apical domain, one of
which is expressing the same combination of TFs mentioned above, suggesting it might
also be undergoing differentiation, iii) one matching the distal apical plate neurons, iv) two
corresponding to ciliary band neurons, v) two located in the rim of the mouth, vi) one
associated with esophageal structures, and vii) one corresponding to the post-oral and
lateral neurons. Thus, our study greatly enhances our knowledge of neuronal diversity in

the *S. purpuratus* larva, nearly doubling the number of known neuronal cell types at this
developmental stage.

Lastly, we also found one cell cluster (undefined) that could not be traced back to a
specific domain. Based on the numbers and distributions of the features and transcript
numbers that are comparable to the ones of real cell type families, such as the
esophageal cell type family, we speculate that this represents a real cell type family and
not empty droplets. (Figure1- Figure supplement 1E). Overall, this cluster exhibited
greatest transcriptional similarity to cell types derived from the ectoderm (Figure 1- Figure
supplement 1F), suggesting it may be of ectodermal origin. Consistent with this, a recent
study revealed a similar uncharacterized ectodermal cell type (Perillo et al., 2020), further
suggesting this cell population exists and is not solely an artifact of our analysis. Taking
into account the great plasticity and regeneration capability of the sea urchin larva, it is
also possible that this non-differentiated ectodermal cell type is a progenitor population in
stasis, waiting to be activated. Future studies are necessary to validate its identity,
function, and origin during larval development.

Sp-Pdx1 as a regulator of neuroendocrine fate

Morphogenesis and organogenesis rely on the hierarchical control of gene expression as
encompassed in the GRNs. Although sea urchins diverged from chordates prior to the
origin of the pancreas, we have previously demonstrated that the neurogenic and
neuronal territories of the sea urchin embryo and larva have a strong pancreatic-like
molecular signature (Perillo et al., 2018). Interestingly, one neuronal population was found
to express *Sp-Pdx1* and *Sp-Brn1/2/4*, as well as the echinoderm-specific neuropeptide
An. *Pdx1* in mammals is essential for proper pancreatic formation, β-cell differentiation,
and regulation of the mature β-cells physiology and function (Hui and Perfetti, 2002). On
the other hand, the vertebrate orthologue of *Sp-Brn1/2/4*, *Brn4* is expressed in the α
pancreatic cells, where it acts as a key differentiation factor of this lineage (Hussain et al.,
2002).

The gene regulatory cascade leading to the endocrine pancreas formation in vertebrates
has been described in great detail and several transcription factors (Hes, Ascl1, Pax6,
Hb9, Mafs, Gbx2), and signaling molecules (Notch) have been characterized as essential
for this process (Ishibashi et al., 1995, de la Pompa et al., 1997, Li et al., 1999, Iso et al.,
2003, Mizusawa et al., 2004, Zaret and Grompe, 2008, Arkhipova et al., 2012, Hart et al.,
2013, Tritschler et al., 2017, Mitchell et al., 2017, Buckle et al., 2018) . (, de la Pompa et
al., 1997, Iso et al., 2003).

Here, we dissected the molecular fingerprint of the *Sp-Pdx1*/*Sp-Brn1/2/4* neuronal type
and identified the presence of genes involved in pancreatic development as well as of
genes known to be expressed in both endocrine pancreatic cells and neurons. From our
scRNA-seq analysis it is evident that *Sp-Notch*, *Sp-Pax6*, *Sp-Hb9*, *Sp-Hes*, *Sp-Ac/Sc* (the
orthologue of ASCL1), *Sp-Mafs* and the recently re-annotated *Sp-Gbx2* (previously
annotated as *Sp-Nk7*- https://new.echinobase.org) are all expressed in the *Sp-Pdx1*/*Sp-*
Brn1/2/4 neurons. This suggests that these neurons of a non-chordate deuterostome
have a gene regulatory machinery similar to the endocrine pancreas cells.

It has also been demonstrated that both endocrine pancreas and neuronal cells share
similar features and are able to produce and respond to several neuronal genes and
neurotransmitters. Interestingly, we were able to identify these shared components in our
Sp-Pdx1/*Sp-Brn1/2/4* neurons. For instance, the neural cell adhesion molecule *Ncam*
known to be produced in the nervous system and endocrine cells of the rat is also
expressed in this cluster (Langley et al., 1989). Moreover, these neurons also express
genes encoding members of the glutamate signaling pathway (*Sp-Glt*: glutamate
synthase; *Sp-Vglut1*: glutamate transporter; *Sp-Glur1*: glutamate receptor), which in
mammals are involved in glucose-responsive insulin secretion (Gonoi et al., 1994,
Maechler and Wollheim, 1999), and tyrosine hydroxylase (*Th*), the rate-limiting enzyme
of catecholamine biosynthesis, which is present in the endocrine pancreas of multiple
species (Teitelman et al., 1993, Iturriza and Thibault, 1993). Furthermore, *Sp-Pdx1*/*Sp-*
Brn1/2/4 neurons express choline acetyltransferase (*Chat*), which has been found to be
highly expressed in human pancreatic islets and may be essential for the stimulation of
insulin secretion by the neighboring β-cells (Rodriguez-Diaz et al., 2011).

Additionally, serotonergic signaling is believed to be involved in the regulation of insulin
secretion as several serotonin receptors have been found to be expressed in human
pancreatic islets (Amisten et al., 2015). Transcripts of the sea urchin serotonin receptor
Sp-5ht7 are present in *Sp-Pdx1*/*Sp-Brn1/2/4* neuronal population suggesting the
serotonin might regulate the neurotransmitter/neuropeptide secretion in a similar way as
for insulin. Similarly, transcripts of the Thyroid hormone receptor B are also found in this
population, suggesting that thyroid hormone signaling may play a similar role to its
differentiation similar to the one in murine endocrine pancreas differentiation (Aiello et al.,
2014). The hormone secretion mediated by pancreatic endocrine cells depends on their
ability to detect changes in extracellular glucose levels. To this end, they are equipped
with Glucose transporters and co-transporters (Navale and Paranjape, 2016, Berger and
Zdzieblo, 2020). Our analysis revealed that the *Sp-Pdx1*/*Sp-Brn1/2/4* neurons produce
transcripts of three glucose co-transporter genes (*Sp-Slc2a1*, *Sp-Slc5a9* and *Sp-*
Slc5a9L) proposing that they are able to detect such changes in glucose levels similarly
to the endocrine pancreas cells.

Evolutionary considerations

Taken together, our data show that the sea urchin *Sp-Pdx1/Sp-Brn1/2/4* neurons uniquely
express a combination of key genes that are necessary for the endocrine pancreas
differentiation and function in the vertebrates (Figure7 & Figure 7- figure supplement 1).
These shared molecular features of *Sp-Pdx1/Sp-Brn1/2/4* neurons and pancreatic cells
suggest these represent features of a cell type present in the deuterostome ancestor.
What was the nature of these cells and what happened in the evolution of the descendant
lineages so that the molecular characteristics of these cells are found in extant cell types
as divergent as ectodermal neurons and pancreatic cells?

One possibility is that the ancestral deuterostomes already possessed a *Sp-Pdx1/Sp-*
Brn1/2/4 neuronal cell type similar to the one found in today's sea urchin, and that the
regulatory program of this cell type was then co-opted into the endodermal lineage in the
vertebrates. The co-option and incorporation of gene regulatory elements to different
developmental or morphogenetic context has been postulated before and has the

potential to give rise to diverse cell types (Monteiro, 2012, Preger-Ben Noon and Frankel,
2015, Martik and McClay, 2015, Hu et al., 2018, Morgulis et al., 2019, McQueen and
Rebeiz, 2020, Cary et al., 2020). In particular, it has been hypothesized that β pancreatic
cells arose during evolution by co-option of a preexisting neuronal cell type program into
the pancreas developmental lineage (Arntfield and van der Kooy, 2011) based on the
many physiological, morphological and molecular features endocrine pancreatic cells
share with neurons (Alpert et al., 1988, Eberhard, 2013). However, given the long list of
genes shared between neurons and pancreatic cells, this would have necessarily
represented a massive co-option event profoundly altering the identity and function of the
receiving cells. Notably, a flexibility towards the co-option of a neuroendocrine phenotype
has also been reported in the vertebrate lineage in the case of the adrenal medulla
derived PC12 cell line, that upon stimulation with NGF, adopts a neurosecretory fate
(Fujita et., 1989).

A second possibility is that both ectodermal neurons and endodermal pancreatic cells are
direct evolutionary descendants of the same precursor cell, i.e. represent sister cell types
(Arendt, 2008). This explanation would avoid co-option; instead, it postulates a direct
evolutionary link between cell types derived from different germ layers - ectoderm versus
endoderm. Surprising at first, a similar link between digestive and neuronal-type
expression profiles has been reported repeatedly. For example, it appears to hold true for
choanocytes in the early-branching sponges (Musser et al., 2019), indicating that the
family of sponge choanocytes might be related to the hypothesized digestive-neuronal
precursors. In the sea anemone *Nematostella vectensis*, pharyngeal cells give rise to
both digestive cells and neurons (Steinmetz et al., 2017), and neuronal cell types and
secretory gland cells exhibit related molecular profiles (Sebe-Pedros et al., 2018). In fact,
based on these findings a close evolutionary link between endoderm and ectoderm has
been postulated (Steinmetz et al., 2017). Finally, motor neurons in the vertebrate ventral
neural tube and pancreatic islet cells share a specific combination of transcription factors
including *pax6, nkx6*, and *hb9* (reviewed in Arendt, 2021). Overall, this second possibility
is also in line with the hypothesis that at least subsets of neuron types in animal nervous
systems may have originated from a mucociliary sole that initially harbored a
neurosecretory network of cells with digestive and communicative functions (Arendt et
al., 2015; Arendt, 2021). Via division of labor, these precursor cell types may have then
given rise to both digestive as well as neuronal cell types in different animal phyla.

Materials and Methods

1. Animal husbandry and culture of embryos
Adult *Strongylocentrotus purpuratus* individuals were obtained from Patrick Leahy
(Kerckhoff Marine Laboratory, California Institute of Technology, Pasadena, CA, USA)
and maintained in circulating seawater aquaria at Stazione Zoologica Anton Dohrn in
Naples. Gametes were obtained by vigorous shaking of the animals. Embryos and larvae
were cultured at 15°C in filtered Mediterranean Sea water diluted 9:1 with deionized
water.

2. Larvae dissociation
Dissociation of the 3 dpf *Strongylocentrotus purpuratus* plutei into single cells was
performed according to adaptation of several protocols (McClay, 1986, McClay, 2004,
Juliano et al., 2014). Larvae were collected, concentrated using a 40 μm Nitex mesh filter
and spun down at 500 g for 5 min. Sea water was removed and larvae were resuspended
in Ca^{2+} Mg^{2+}free artificial sea water. Larvae were spun down at 500 g for 5 min and
resuspended in dissociation buffer containing 1M glycine and 0.02 M EDTA in Ca^{2+}
Mg^{2+}Free artificial sea water. Larvae were incubated for 10 min on ice and mixed gently
via pipette aspiration every 2 min. From that point and onwards the progress of
dissociation was monitored. Dissociated cells were spun down at 700 g for 5 min and
washed several times with Ca^{2+} Mg^{2+}Free artificial sea water. Cell viability was assessed
via using Propidium Iodide and Fluorescein diacetate and only specimens with cell
viability ≥ 90% were further processed. Single cells were counted using a hemocytometer
and diluted according to the manufacturer's protocol (10x Genomics). Throughout this
procedure samples were kept at 4°C.

3. Single cell RNA sequencing and data analysis
Single cell RNA sequencing was performed using the 10x Genomics single cell capturing
system. Specimens from four independent biological replicates, ranging from 6000-
20.000 cells, were loaded on the 10X Genomics Chromium Controller. Single cell cDNA
libraries were prepared using the Chromium Single Cell 3' Reagent Kit (Chemistries v2
and v3). Libraries were sequenced by GeneCore (EMBL, Heidelberg, Germany) for 75
bp paired-end reads (Illumina NextSeq 500), resulting in a mean of 88M reads. Cell
Ranger Software Suite 3.0.2 (10x Genomics) was used for the alignment of the single-
cell RNA-seq output reads and generation of feature, barcode and matrices. The genomic
index was made in Cell Ranger using the S. purpuratus genome version 3.1 (Sea Urchin
Genome Sequencing et al., 2006, Kudtarkar and Cameron, 2017). Cell Ranger output
matrices for four biological and two technical replicates were used for further analysis in
Seurat v3.0.2 R package (Stuart et al., 2019). The analysis was performed according to
the Seurat scRNA-seq R package documentation (Butler et al., 2018, Stuart et al., 2019).
Genes that are transcribed in less than three cells and cells that have less than a minimum
of 200 transcribed genes were excluded from the analysis. The cutoff number of
transcribed genes was determined based on feature scatter plots and varies depending
on the replicate. 19.699 cells out of the 29,130 cells estimated by Cell Ranger passed the
quality checks and were further analyzed. Datasets were normalized and variable genes
were found using the vst method with a maximum of 2000 variable features. Data
integration was performed via identification of anchors between the six different objects.
Next the datasets were scaled and principal component (PCA) analysis was performed.
Nearest Neighbor (SNN) graph was computed with 20 dimensions (resolution 1.0) to
identify the clusters. Uniform Manifold Approximate and Projection (UMAP) was used to
perform clustering dimensionality reduction. Cluster markers were found using the genes
that are detected in at least 0.01 fraction of min.pct cells in the two clusters. Transcripts
of all genes per cell type were identified by converting a Seurat DotPlot with all these

transcripts as features into a table (ggplot2 3.2.0 R package). Subclustering analysis was
performed by selecting a cell type of interest and performing similar analysis as described
above. All resulting tables containing the genes transcribed within different cell type
families were further annotated adding PFAM terms (Trapnell et al., 2010, Finn et al.,
2014) for associated proteins, gene ontology terms and descriptions from Echinobase
(Kudtarkar and Cameron, 2017).

Cell type tree reconstruction was performed in R (version 4.0.3) using the neighbor-joining
tree function in package 'ape' (version 5.4-1). Two final trees were constructed, one that
derived pairwise expression distances between clusters using all expressed genes, and
a second in the analysis was restricted to only expressed transcription factors. A gene
was defined as "expressed" when it had an average expression of greater than 3
transcripts per million in at least one of the 21 major clusters. For each tree, pairwise
distances were calculated as 1 – spearman correlation between each of the 20 major
clusters, excluding the undefined cells, using the spearman.dist function in the bioDist
package (version 1.60.0). To assess the support of each bifurcation in the tree, we
performed 10,000 bootstrap replicates using the boot.phylo function in 'ape'. Although not
reported in this manuscript, we also tested the effect of using different distance metrics
and normalization methods on tree topology. Specifically, we reconstructed trees using
distances derived from Pearson correlations of either log or square root normalized
average expression values, and Euclidean distances of scaled log-normalized expression
values. In general, similar topologies were found in all analyses, recovering clades of
endodermal and ectodermal derivatives. Mesodermal derivatives often, but not always,
grouped together, and the position of the neuron cluster moved in and out of the clade of
ectodermal derivatives. This latter result may possibly arise because of conflicting signals
resulting from cell type heterogeneity (i.e. different neuron types) within the cluster.

In order to assess the significance of identifying the homologs of the vertebrate genes
involved in the development of the pancreas in the "post oral and lateral neurons" sub-
population of neuronal cells, the normalized counts from all the transcripts present in the
neuronal subcluster were extracted in the form of a matrix using GetAssayData command
from Seurat v3.0.2. The counts within this matrix were then randomized with 100000
iterations using the randomizeMatrix function from the picante v1.8.2 package (Kembel
et al., 2010) with the independent swap null model. A new Seurat object was created,
which is the same as the neuronal subclustering object, however the actual transcript
normalized counts were replaced with the randomized ones. Then the average
expression of the 20 pancreatic genes was extracted from both from actual neuronal
subclustering Seurat object and the one with randomized counts using
AverageExpression function to visualize the expression of these pancreatic genes in the
"post oral and lateral neurons" subcluster in actual data and randomized data. This
process was looped 500 times for randomized data to then calculate the mean number
of pancreatic genes expressed in the "post oral and lateral neurons" subcluster to obtain
an estimate of the expected number of genes expressed, rounded to the closest integer,
if their expression was found in this subcluster by random. The number of genes

expressed in the actual data and the randomized data was then used to generate
proportions of the number of expressed and not expressed pancreatic genes in this
neuronal subcluster. Then, we have performed a chi-squared test within R (Hope, 1968)
to show that the proportion from the actual data proportion is significantly different to the
randomized data proportion with p-value less than 0.002.

4. Whole mount RNA Fluorescent *in situ* hybridization
Fluorescent *in situ* hybridization was performed as outlined in (Perillo et al., 2021).
Fluorescent signal was developed via using fluorophore conjugated tyramide technology
(Perkin Elmer, Cat. #NEL752001KT). Antisense probes were transcribed from linearized
DNA and labeled either during transcription via using digoxigenin-11-UTP nucleotides, or
post-transcriptionally by using Fluorescein (Mirus Bio, Cat. #MIR3200) or DNP (Mirus Bio,
Cat. #MIR3825) following the manufacturer's instructions. Probes for *Sp-Pdx1, Sp-Cdx,*
Sp-ManrC1A, Sp-Six1/2, Sp-Fgf9/16/20, Sp-Brn1/2/4, Sp-Ngn, Sp-Isl, Sp-NeuroD1, Sp-
Pks1, Sp-Soxb2, Sp-An, Sp-Trh and Sp-Salmfap were produced as previously published
[*Sp-Pdx1, Sp-Cdx* (Cole et al., 2009), *Sp-ManrC1A* (Annunziata et al., 2014), *Sp-Six1/2,*
Sp-Fgf9/16/20 (Andrikou et al., 2015), *Sp-Brn1/2/4* (Cole and Arnone, 2009), *Sp-Ngn, Sp-*
Isl, Sp-NeuroD1 (Perillo et al., 2018), *Sp-Pks1* (Perillo et al, 2020), *Sp-SoxB2*
(Anishchenko et al., 2018), *Sp-An, Sp-Trh, Sp-Salmfap* (Wood et al., 2018)]. Primer
sequences used for cDNA isolation and probes synthesis are in Supplementary file 2.
Specimens were imaged using a Zeiss LSM 700 confocal microscope.
789

5. Immunohistochemistry (IHC)
Immunohistochemical staining or IHC paired with FISH was performed as described in
(Perillo et al., 2021). Briefly 3 dpf plutei were fixed in 4 % paraformaldehyde (PFA) in
filtered sea water (FSW) for 15 min at room temperature (RT). FSW was removed and
samples were incubated in 100% methanol for 1 min at RT, washed multiple times with
phosphate buffer saline with 0.1% Tween 20 (PBST) and incubated blocking solution
containing 1 mg/ml Bovine Serum Albumin (BSA) and 4% sheep in PBST for 1h. Primary
antibodies were added in the appropriate dilution and incubated for 1h and 30 min at
37°C. Anti-acetylated alpha tubulin (Sigma-Aldrich T67930) was used to label cilia and
microtubules (1:200), Anti-Msp130 (gift from Dr. David R. McClay) to label skeletogenic
cells (undiluted), 1E11 (gift from Dr. Robert Burke) to mark the nervous system (1:20),
5c7 (gift from Dr. David R. McClay) to label the endoderm (undiluted), Sp-An to label the
post-oral and lateral neurons (1:250),   Sp-Th (Sigma-Aldrich AB152) to label
catecholaminergic neurons (1:100) and Sp-Chat (GeneTex GXGTX113164S) to label
cholinergic neurons (1:100). Specimens were washed multiple times with PBST and
incubated for 1h with the appropriate secondary antibody (AlexaFluor) diluted 1:1000 in
PBST. Larvae were washed several times with PBST and imaged using a Zeiss LSM 700
confocal microscope.
808

6. EdU labelling paired with immunohistochemistry
In order to understand the spatial distribution of proliferating cells across the putative
broad cell types, cell proliferation assays were carried out using Click-It EdU Cell

Proliferation Kit for Imaging Alexa Flour^{TM} 647 (Thermo Fisher Scientific). Larvae were
treated with EdU at a final concentration of 10 µM in FSW and let to grow for 2 hours.
Samples were fixed in 4% PFA in FSW for 15 min (RT) and washed several times with
PBST. PBST was removed, replaced by 100% Methanol for 1 min (RT) and followed by
several washes with PBST. After this step one can continue with either developing the
EdU signal or performing immunohistochemistry as described above. In order to develop
the EdU signal the Click-iT ^{TM} reaction mix was prepared according to the manufacturer's
guidelines. PBST was removed and the reaction mix was added to the samples for 30
min (RT). Larvae were washed several times with PBST, mounted and imaged using a
Zeiss LSM 700 confocal microscope
822

823 **7. Gene regulatory network draft**
824 Gene regulatory modules and networks were drafted using the interactive tool for
825 building and visualizing GRNs BioTapestry (Longabaugh, 2012).

Competing interests

828 The authors declare no competing interests, or appearance of competing interests, in the
829 production and dissemination of this work.
830

Acknowledgements

The authors would like to thank Prof. Paola Oliveri (UCL) for providing critical input on the
analysis and structure of the manuscript. We are grateful to Drs. David McClay and Robert
Burke for kindly providing antibodies and Dr. Francesco Lamanna (ZMBH) for helping
with the computational analysis. We also would like to thank Davide Caramiello for taking
care of the adult sea urchins, Dr. Giovanna Benvenuto (SZN) for microscopy assistance
and the Arnone lab members Maria Cocurullo and Inés Fournon Berodia for their help in
gene cloning and preparation of several RNA probes. We also thank Drs. Vladimir Benes
and Bianka Baying (GeneCore, Heidelberg, Germany) for sequencing of our single cell
datasets. Also we are grateful to the animal technician Emily Savage (EMBL, Heidelberg,
Germany) for her precious assistance.
842

Funding

844 This project has been supported by the European Union's Horizon 2020 research and
845 innovation programme under the Marie Skłodowska-Curie grant agreement no. 766053
846 (EvoCELL: grant to MIA and DA, fellowship to PP) and the Advanced grant
847 'NeuralCellTypeEvo' 788921 by the European Commission (grant to DA and JM).

References

ADOMAKO-ANKOMAH, A. & ETTENSOHN, C. A. 2014. Growth factors and early mesoderm morphogenesis: insights from the sea urchin embryo. *Genesis,* 52**,** 158-72.

AIELLO, V., MORENO-ASSO, A., SERVITJA, J. M. & MARTIN, M. 2014. Thyroid hormones promote endocrine differentiation at expenses of exocrine tissue. *Exp Cell Res,* 322**,** 236-48.

AMISTEN, S., NEVILLE, M., HAWKES, R., PERSAUD, S. J., KARPE, F. & SALEHI, A. 2015. An atlas of G-protein coupled receptor expression and function in human subcutaneous adipose tissue. *Pharmacol Ther,* 146**,** 61-93.

ANDRIKOU, C., IOVENE, E., RIZZO, F., OLIVERI, P. & ARNONE, M. I. 2013. Myogenesis in the sea urchin embryo: the molecular fingerprint of the myoblast precursors. *Evodevo,* 4**,** 33.

ANDRIKOU, C., PAI, C. Y., SU, Y. H. & ARNONE, M. I. 2015. Logics and properties of a genetic regulatory program that drives embryonic muscle development in an echinoderm. *Elife,* 4.

ANGERER, R. C. & DAVIDSON, E. H. 1984. Molecular indices of cell lineage specification in sea urchin embryos. *Science,* 226**,** 1153-60.

ANISHCHENKO, E., ARNONE, M. I. & D'ANIELLO, S. 2018. SoxB2 in sea urchin development: implications in neurogenesis, ciliogenesis and skeletal patterning. *Evodevo,* 9**,** 5.

ANNUNZIATA, R., ANDRIKOU, C., PERILLO, M., CUOMO, C. & ARNONE, M. I. 2019. Development and evolution of gut structures: from molecules to function. *Cell Tissue Res,* 377**,** 445-458.

ANNUNZIATA, R. & ARNONE, M. I. 2014. A dynamic regulatory network explains ParaHox gene control of gut patterning in the sea urchin. *Development,* 141**,** 2462-72.

ANNUNZIATA, R., PERILLO, M., ANDRIKOU, C., COLE, A. G., MARTINEZ, P. & ARNONE, M. I. 2014. Pattern and process during sea urchin gut morphogenesis: the regulatory landscape. *Genesis,* 52**,** 251-68.

ARENDT, D. 2008. The evolution of cell types in animals: emerging principles from molecular studies. *Nat Rev Genet,* 9**,** 868-82.

ARENDT, D, BENITO-GUTIERREZ, E., BRUNET, T., MARLOW H. 2015. Gastric pouches and the mucociliary sole: setting the stage for nervous system evolution. Phil. Trans. R. Soc. B 370: 20150286
ARENDT, D. 2021.Elementary nervous systems. Phil. Trans. R. Soc. B 376: 20200347.

ARKHIPOVA, V., WENDIK, B., DEVOS, N., EK, O., PEERS, B. & MEYER, D. 2012. Characterization and regulation of the hb9/mnx1 beta-cell progenitor specific enhancer in zebrafish. *Dev Biol,* 365**,** 290-302.

ARNONE, M.I., BYRNE, M., MARTINEZ, P. 2015. Echinodermata. In: Wanninger A. (eds) Evolutionary Developmental Biology of Invertebrates 6. Springer, Vienna.

ARNONE, M. I., ANDRIKOU, C. & ANNUNZIATA, R. 2016. Echinoderm systems for gene regulatory studies in evolution and development. *Curr Opin Genet Dev,* 39**,** 129-137.

ARNTFIELD, M. E. & VAN DER KOOY, D. 2011. beta-Cell evolution: How the pancreas borrowed from the brain: The shared toolbox of genes expressed by neural and pancreatic endocrine cells may reflect their evolutionary relationship. *Bioessays,* 33**,** 582-7.

BEN-TABOU DE-LEON, S., SU, Y. H., LIN, K. T., LI, E. & DAVIDSON, E. H. 2013. Gene regulatory control in the sea urchin aboral ectoderm: spatial initiation, signaling inputs, and cell fate lockdown. *Dev Biol,* 374**,** 245-54.

BERGER, C. & ZDZIEBLO, D. 2020. Glucose transporters in pancreatic islets. *Pflugers Arch,* 472**,** 1249-1272.

BISGROVE, B. W. & BURKE, R. D. 1987. Development of the Nervous-System of the Pluteus Larva of Strongylocentrotus-Droebachiensis. *Cell and Tissue Research,* 248**,** 335-343.

BUCKLE, A., NOZAWA, R. S., KLEINJAN, D. A. & GILBERT, N. 2018. Functional characteristics of novel pancreatic Pax6 regulatory elements. *Hum Mol Genet,* 27**,** 3434-3448.

BURKE, R. D., ANGERER, L. M., ELPHICK, M. R., HUMPHREY, G. W., YAGUCHI, S., KIYAMA, T., LIANG, S., MU, X., AGCA, C., KLEIN, W. H., BRANDHORST, B. P., ROWE, M., WILSON, K., CHURCHER, A. M., TAYLOR, J. S., CHEN, N., MURRAY, G., WANG, D., MELLOTT, D., OLINSKI, R., HALLBOOK, F. & THORNDYKE, M. C. 2006a. A genomic view of the sea urchin nervous system. *Dev Biol,* 300**,** 434-60.

BURKE, R. D., MOLLER, D. J., KRUPKE, O. A. & TAYLOR, V. J. 2014. Sea urchin neural development and the metazoan paradigm of neurogenesis. *Genesis,* 52**,** 208-21.

BURKE, R. D., OSBORNE, L., WANG, D., MURABE, N., YAGUCHI, S. & NAKAJIMA, Y. 2006b. Neuron-specific expression of a synaptotagmin gene in the sea urchin Strongylocentrotus purpuratus. *J Comp Neurol,* 496**,** 244-51.

BUTLER, A., HOFFMAN, P., SMIBERT, P., PAPALEXI, E. & SATIJA, R. 2018. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol,* 36**,** 411-420.

CALESTANI, C. & ROGERS, D. J. 2010. Cis-regulatory analysis of the sea urchin pigment cell gene polyketide synthase. *Dev Biol,* 340**,** 249-55.

CAMERON, R. A. & DAVIDSON, E. H. 1991. Cell type specification during sea urchin development. *Trends Genet,* 7**,** 212-8.

CAMERON, R. A., HOUGH-EVANS, B. R., BRITTEN, R. J. & DAVIDSON, E. H. 1987. Lineage and fate of each blastomere of the eight-cell sea urchin embryo. *Genes Dev,* 1**,** 75-85.

CAO, C., LEMAIRE, L. A., WANG, W., YOON, P. H., CHOI, Y. A., PARSONS, L. R., MATESE, J. C., WANG, W., LEVINE, M. & CHEN, K. 2019. Comprehensive single-cell transcriptome lineages of a proto-vertebrate. *Nature,* 571**,** 349-354.

CARY, G. A., MCCAULEY, B. S., ZUEVA, O., PATTINATO, J., LONGABAUGH, W. & HINMAN, V. F. 2020. Systematic comparison of sea urchin and sea star developmental gene regulatory networks explains how novelty is incorporated in early development. *Nat Commun,* 11**,** 6235.

CHESTNUT, B., CASIE CHETTY, S., KOENIG, A. L. & SUMANAS, S. 2020. Single-cell
transcriptomic analysis identifies the conversion of zebrafish Etv2-deficient
vascular progenitors into skeletal muscle. *Nat Commun,* 11**,** 2796.
CHO, B., YOON, S. H., LEE, D., KORANTENG, F., TATTIKOTA, S. G., CHA, N., SHIN,
M., DO, H., HU, Y., OH, S. Y., LEE, D., VIPIN MENON, A., MOON, S. J.,
PERRIMON, N., NAM, J. W. & SHIM, J. 2020. Single-cell transcriptome maps of
myeloid blood cell lineages in Drosophila. *Nat Commun,* 11**,** 4483.
COLE, A. G. & ARNONE, M. I. 2009. Fluorescent in situ hybridization reveals multiple
expression domains for SpBrn1/2/4 and identifies a unique ectodermal cell type
that co-expresses the ParaHox gene SpLox. *Gene Expr Patterns,* 9**,** 324-8.
COLE, A. G., RIZZO, F., MARTINEZ, P., FERNANDEZ-SERRA, M. & ARNONE, M. I.
2009. Two ParaHox genes, SpLox and SpCdx, interact to partition the posterior
endoderm in the formation of a functional gut. *Development,* 136**,** 541-9.
DAVIDSON, E. H., CAMERON, R. A. & RANSICK, A. 1998. Specification of cell fate in
the sea urchin embryo: summary and some proposed mechanisms.
Development, 125**,** 3269-90.
DAVIDSON, E. H. & ERWIN, D. H. 2006. Gene regulatory networks and the evolution of
animal body plans. *Science,* 311**,** 796-800.
DAVIDSON, E. H., MCCLAY, D. R. & HOOD, L. 2003. Regulatory gene networks and
the properties of the developmental process. *Proc Natl Acad Sci U S A,* 100**,**
1475-80.
DAVIE, K., JANSSENS, J., KOLDERE, D., DE WAEGENEER, M., PECH, U., KREFT,
L., AIBAR, S., MAKHZAMI, S., CHRISTIAENS, V., BRAVO GONZALEZ-BLAS,
C., POOVATHINGAL, S., HULSELMANS, G., SPANIER, K. I., MOERMAN, T.,
VANSPAUWEN, B., GEURS, S., VOET, T., LAMMERTYN, J., THIENPONT, B.,
LIU, S., KONSTANTINIDES, N., FIERS, M., VERSTREKEN, P. & AERTS, S.
2018. A Single-Cell Transcriptome Atlas of the Aging Drosophila Brain. *Cell,* 174**,**
982-998 e20.
DE LA POMPA, J. L., WAKEHAM, A., CORREIA, K. M., SAMPER, E., BROWN, S.,
AGUILERA, R. J., NAKANO, T., HONJO, T., MAK, T. W., ROSSANT, J. &
CONLON, R. A. 1997. Conservation of the Notch signalling pathway in
mammalian neurogenesis. *Development,* 124**,** 1139-48.
DULOQUIN, L., LHOMOND, G. & GACHE, C. 2007. Localized VEGF signaling from
ectoderm to mesenchyme cells controls morphogenesis of the sea urchin embryo
skeleton. *Development,* 134**,** 2293-302.
ESAULOVA, E., CANTONI, C., SHCHUKINA, I., ZAITSEV, K., BUCELLI, R. C., WU, G.
F., ARTYOMOV, M. N., CROSS, A. H. & EDELSON, B. T. 2020. Single-cell
RNA-seq analysis of human CSF microglia and myeloid cells in
neuroinflammation. *Neurol Neuroimmunol Neuroinflamm,* 7.
ETTENSOHN, C. A. 2020. The gene regulatory control of sea urchin gastrulation. *Mech*
Dev, 103599.
FINN, R. D., BATEMAN, A., CLEMENTS, J., COGGILL, P., EBERHARDT, R. Y.,
EDDY, S. R., HEGER, A., HETHERINGTON, K., HOLM, L., MISTRY, J.,
SONNHAMMER, E. L., TATE, J. & PUNTA, M. 2014. Pfam: the protein families
database. *Nucleic Acids Res,* 42**,** D222-30.

FOSTER, S., OULHEN, N., & WESSEL, G. 2020. A single cell RNA sequencing
resource for early sea urchin development. Development (Cambridge, England),
147(17), dev191528.
FU, X., HE, F., LI, Y., SHAHVERANOV, A. & HUTCHINS, A. P. 2017. Genomic and
molecular control of cell type and cell type conversions. *Cell Regen,* 6**,** 1-7.
FUJITA, K., LAZAROVICI, P. & GUROFF, G. 1989. Regulation of the differentiation of
PC12 pheochromocytoma cells. Environ Health Perspect. 80:127-42.
GARNER, S., ZYSK, I., BYRNE, G., KRAMER, M., MOLLER, D., TAYLOR, V. &
BURKE, R. D. 2016. Neurogenesis in sea urchin embryos and the diversity of
deuterostome neurogenic mechanisms. *Development,* 143**,** 286-97.
GONOI, T., MIZUNO, N., INAGAKI, N., KUROMI, H., SEINO, Y., MIYAZAKI, J. &
SEINO, S. 1994. Functional neuronal ionotropic glutamate receptors are
expressed in the non-neuronal cell line MIN6. *J Biol Chem,* 269**,** 16989-92.
HARKEY, M. A., WHITELEY, H. R. & WHITELEY, A. H. 1992. Differential expression of
the msp130 gene among skeletal lineage cells in the sea urchin embryo: a three
dimensional in situ hybridization analysis. *Mech Dev,* 37**,** 173-84.
HARLOW, P. & NEMER, M. 1987. Coordinate and selective beta-tubulin gene
expression associated with cilium formation in sea urchin embryos. *Genes Dev,*
1**,** 1293-304.
HART, A. W., MELLA, S., MENDRYCHOWSKI, J., VAN HEYNINGEN, V. & KLEINJAN,
D. A. 2013. The developmental regulator Pax6 is essential for maintenance of
islet cell function in the adult mouse pancreas. *PLoS One,* 8**,** e54173.
HO, E. C., BUCKLEY, K. M., SCHRANKEL, C. S., SCHUH, N. W., HIBINO, T., SOLEK,
C. M., BAE, K., WANG, G. & RAST, J. P. 2017. Perturbation of gut bacteria
induces a coordinated cellular immune response in the purple sea urchin larva.
Immunol Cell Biol, 95**,** 647.
HOPE, A. C. A. 1968. 'A Simplified Monte Carlo Significance Test Procedure', *Journal*
of the Royal Statistical Society: Series B (Methodological), 30(3), pp. 582–598
HOWARD-ASHBY, M., MATERNA, S. C., BROWN, C. T., CHEN, L., CAMERON, R. A.
& DAVIDSON, E. H. 2006. Identification and characterization of homeobox
transcription factor genes in Strongylocentrotus purpuratus, and their expression
in embryonic development. *Dev Biol,* 300**,** 74-89.
HU, Y., SCHMITT-ENGEL, C., SCHWIRZ, J., STROEHLEIN, N., RICHTER, T.,
MAJUMDAR, U. & BUCHER, G. 2018. A morphological novelty evolved by co-
option of a reduced gene regulatory network and gene recruitment in a beetle.
Proc Biol Sci, 285.
HUI, H. & PERFETTI, R. 2002. Pancreas duodenum homeobox-1 regulates pancreas
development during embryogenesis and islet cell function in adulthood. *Eur J*
Endocrinol, 146**,** 129-41.
HUSSAIN, M. A., MILLER, C. P. & HABENER, J. F. 2002. Brn-4 transcription factor
expression targeted to the early developing mouse pancreas induces ectopic
glucagon gene expression in insulin-producing beta cells. *J Biol Chem,* 277**,**
16028-32.
ISHIBASHI, M., ANG, S. L., SHIOTA, K., NAKANISHI, S., KAGEYAMA, R. &
GUILLEMOT, F. 1995. Targeted disruption of mammalian hairy and Enhancer of
split homolog-1 (HES-1) leads to up-regulation of neural helix-loop-helix factors,

premature neurogenesis, and severe neural tube defects. *Genes Dev,* 9**,** 3136-48.

ISO, T., KEDES, L. & HAMAMORI, Y. 2003. HES and HERP families: multiple effectors of the Notch signaling pathway. *J Cell Physiol,* 194**,** 237-55.

ITURRIZA, F. C. & THIBAULT, J. 1993. Immunohistochemical investigation of tyrosine-hydroxylase in the islets of Langerhans of adult mice, rats and guinea pigs. *Neuroendocrinology,* 57**,** 476-80.

JULIANO, C., SWARTZ, S. Z. & WESSEL, G. 2014. Isolating specific embryonic cells of the sea urchin by FACS. *Methods Mol Biol,* 1128**,** 187-96.

JULIANO, C. E., VORONINA, E., STACK, C., ALDRICH, M., CAMERON, A. R. & WESSEL, G. M. 2006. Germ line determinants are not localized early in sea urchin development, but do accumulate in the small micromere lineage. *Dev Biol,* 300**,** 406-15.

JULIANO, C. E., YAJIMA, M. & WESSEL, G. M. 2010. Nanos functions to maintain the fate of the small micromere lineage in the sea urchin embryo. *Dev Biol,* 337**,** 220-32.

JUNG, M., WELLS, D., RUSCH, J., AHMAD, S., MARCHINI, J., MYERS, S. R. & CONRAD, D. F. 2019. Unified single-cell analysis of testis gene regulation and pathology in five mouse strains. *Elife,* 8.

KANETO, H., MIYATSUKA, T., KAWAMORI, D., YAMAMOTO, K., KATO, K., SHIRAIWA, T., KATAKAMI, N., YAMASAKI, Y., MATSUHISA, M. & MATSUOKA, T. A. 2008. PDX-1 and MafA play a crucial role in pancreatic beta-cell differentiation and maintenance of mature beta-cell function. *Endocr J,* 55**,** 235-52.

KATOW, H., SUYEMITSU, T., OOKA, S., YAGUCHI, J., JIN-NAI, T., KUWAHARA, I., KATOW, T., YAGUCHI, S. & ABE, H. 2010. Development of a dopaminergic system in sea urchin embryos and larvae. *J Exp Biol,* 213**,** 2808-19.

KEMBEL, S. W., COWAN, P. D., HELMUS, M. R., CORNWELL, W. K., MORTON, H., ACKERLY, D. D., BLOMBERG, S. P. & WEBB, C. O. .2010. 'Picante: R tools for integrating phylogenies and ecology', *Bioinformatics*, 26(11), pp. 1463–1464.

KHOR, J.M., GUERRERO-SANTORO, J. & ETTENSOHN C.A. 2019. Genome-wide identification of binding sites and gene targets of Alx1, a pivotal regulator of echinoderm skeletogenesis. Development. 146(16):dev180653.KLEIN, A. M., MAZUTIS, L., AKARTUNA, I., TALLAPRAGADA, N., VERES, A., LI, V., PESHKIN, L., WEITZ, D. A. & KIRSCHNER, M. W. 2015. Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells. *Cell,* 161**,** 1187-1201.

KROUK, G., LINGEMAN, J., COLON, A. M., CORUZZI, G. & SHASHA, D. 2013. Gene regulatory networks in plants: learning causality from time and perturbation. *Genome Biology,* 14.

KUDTARKAR, P. & CAMERON, R. A. 2017. Echinobase: an expanding resource for echinoderm genomic information. *Database (Oxford),* 2017.

LI, H., ARBER, S., JESSELL, T. M. & EDLUND, H. 1999. Selective agenesis of the dorsal pancreas in mice lacking homeobox gene Hlxb9. *Nat Genet,* 23**,** 67-70.

LONGABAUGH, W. J. 2012. BioTapestry: a tool to visualize the dynamic properties of gene regulatory networks. *Methods Mol Biol,* 786**,** 359-94.

LUO, Y. J. & SU, Y. H. 2012. Opposing nodal and BMP signals regulate left-right asymmetry in the sea urchin larva. *PLoS Biol,* 10**,** e1001402.

LYONS, D. C., KALTENBACH, S. L. & MCCLAY, D. R. 2012. Morphogenesis in sea urchin embryos: linking cellular events to gene regulatory network states. *Wiley Interdiscip Rev Dev Biol,* 1**,** 231-52.

MAECHLER, P. & WOLLHEIM, C. B. 1999. Mitochondrial glutamate acts as a messenger in glucose-induced insulin exocytosis. *Nature,* 402**,** 685-9.

MARTIK, M. L. & MCCLAY, D. R. 2015. Deployment of a retinal determination gene network drives directed cell migration in the sea urchin embryo. *Elife,* 4.

MASSRI, A. J., GREENSTREET, L., AFANASSIEV, A., ESCOBAR, A. B., WRAY, G. M., SCHIEBINGER, G. & MCCLAY, D. R. 2021. Developmental Single-cell transcriptomics in the Lytechinus variegatus Sea Urchin Embryo. Development. 148(19):dev198614.

MATERNA, S. C., HOWARD-ASHBY, M., GRAY, R. F. & DAVIDSON, E. H. 2006. The C2H2 zinc finger genes of Strongylocentrotus purpuratus and their expression in embryonic development. *Dev Biol,* 300**,** 108-20.

MATERNA, S. C., RANSICK, A., LI, E. & DAVIDSON, E. H. 2013. Diversification of oral and aboral mesodermal regulatory states in pregastrular sea urchin embryos. *Dev Biol,* 375**,** 92-104.

MCCLAY, D. R. 1986. Embryo dissociation, cell isolation, and cell reassociation. *Methods Cell Biol,* 27**,** 309-23.

MCCLAY, D. R. 2004. Methods for embryo dissociation and analysis of cell adhesion. *Methods Cell Biol,* 74**,** 311-29.

MCCLAY, D. R. 2011. Evolutionary crossroads in developmental biology: sea urchins. *Development,* 138**,** 2639-48.

MCCLAY, D. R., MIRANDA, E. & FEINBERG, S. L. 2018. Neurogenesis in the sea urchin embryo is initiated uniquely in three domains. *Development,* 145.

MCCLAY, D. R., WARNER, J., MARTIK, M., MIRANDA, E. & SLOTA, L. 2020. Gastrulation in the sea urchin. *Curr Top Dev Biol,* 136**,** 195-218.

MCQUEEN, E. & REBEIZ, M. 2020. On the specificity of gene regulatory networks: How does network co-option affect subsequent evolution? *Curr Top Dev Biol,* 139**,** 375-405.

MITCHELL, R. K., NGUYEN-TU, M. S., CHABOSSEAU, P., CALLINGHAM, R. M., PULLEN, T. J., CHEUNG, R., LECLERC, I., HODSON, D. J. & RUTTER, G. A. 2017. The transcription factor Pax6 is required for pancreatic beta cell identity, glucose-regulated ATP synthesis, and Ca(2+) dynamics in adult mice. *J Biol Chem,* 292**,** 8892-8906.

MIZUSAWA, N., HASEGAWA, T., OHIGASHI, I., TANAKA-KOSUGI, C., HARADA, N., ITAKURA, M. & YOSHIMOTO, K. 2004. Differentiation phenotypes of pancreatic islet beta- and alpha-cells are closely related with homeotic genes and a group of differentially expressed genes. *Gene,* 331**,** 53-63.

MONTEIRO, A. 2012. Gene regulatory networks reused to build novel traits: co-option of an eye-related gene regulatory network in eye-like organs and red wing patches on insect wings is suggested by optix expression. *Bioessays,* 34**,** 181-6.

MORGULIS, M., GILDOR, T., ROOPIN, M., SHER, N., MALIK, A., LALZAR, M., DINES, M., BEN-TABOU DE-LEON, S., KHALAILY, L. & BEN-TABOU DE-LEON, S.

2019. Possible cooption of a VEGF-driven tubulogenesis program for biomineralization in echinoderms. *Proc Natl Acad Sci U S A,* 116**,** 12353-12362.

NAVALE, A. M. & PARANJAPE, A. N. 2016. Glucose transporters: physiological and pathological roles. *Biophys Rev,* 8**,** 5-9.

MUSSER J., SCHIPPERS K.J., NICKELI M., MIZZON G., KOHN A.B., PAPE C, HAMMELI J.U., WOLF F., LIANG C., HERNANDEZ-PLAZA A., ACHIM K., SCHIEBER N.L., FRANCIS W.R., VARGAS S. R., KLING S., RENKERT M., FEUDA R., GASPAR I., BURKHARDT P., BORK P., BECK M., KRESHUK A., WORHEIDE G., HUERTA-CEPAS J., Schwab Y., Moroz L.L., Arendt D. bioRxiv 758276

NESTOROWA, S., HAMEY, F. K., PIJUAN SALA, B., DIAMANTI, E., SHEPHERD, M., LAURENTI, E., WILSON, N. K., KENT, D. G. & GOTTGENS, B. 2016. A single-cell resolution map of mouse hematopoietic stem and progenitor cell differentiation. *Blood,* 128**,** e20-31.

OISHI, Y & MANABE, I. Krüppel-Like Factors in Metabolic Homeostasis and Cardiometabolic Disease. Front Cardiovasc Med.5:69.

OKAZAKI, K. 1965. Skeleton formation of sea urchin larvae. V. Continuous observation of the process of matrix formation. *Exp Cell Res,* 40**,** 585-96.

OLIVERI, P., TU, Q. & DAVIDSON E.H. 2008. Global regulatory logic for specification of an embryonic cell lineage. Proceedings of the National Academy of Sciences, 105 (16) 5955-5962

PEHRSON, J. R. & COHEN, L. H. 1986. The fate of the small micromeres in sea urchin development. *Dev Biol,* 113**,** 522-6.

PERILLO, M. & ARNONE, M. I. 2014. Characterization of insulin-like peptides (ILPs) in the sea urchin Strongylocentrotus purpuratus: insights on the evolution of the insulin family. *Gen Comp Endocrinol,* 205**,** 68-79.

PERILLO, M., OULHEN, N., FOSTER, S., SPURRELL, M., CALESTANI, C. & WESSEL, G. 2020. Regulation of dynamic pigment cell states at single-cell resolution. *Elife,* 9.

PERILLO, M., PAGANOS, P., MATTIELLO, T., COCURULLO, M., OLIVERI, P. & ARNONE, M. I. 2018. New Neuronal Subtypes With a "Pre-Pancreatic" Signature in the Sea Urchin Stongylocentrotus purpuratus. *Front Endocrinol (Lausanne),* 9**,** 650.

PERILLO, M., PAGANOS, P., SPURRELL, M., ARNONE, M. I. & WESSEL, G. M. 2021. Methodology for Whole Mount and Fluorescent RNA In Situ Hybridization in Echinoderms: Single, Double, and Beyond. *Methods Mol Biol,* 2219**,** 195-216.

PERILLO, M., WANG, Y. J., LEACH, S. D. & ARNONE, M. I. 2016. A pancreatic exocrine-like cell regulatory circuit operating in the upper stomach of the sea urchin Strongylocentrotus purpuratus larva. *BMC Evol Biol,* 16**,** 117.

PREGER-BEN NOON, E. & FRANKEL, N. 2015. Evolving Genital Structures: A Deep Look at Network Co-option. *Dev Cell,* 34**,** 485-6.

QI, F., QIAN, S., ZHANG, S. & ZHANG, Z. 2020. Single cell RNA sequencing of 13 human tissues identify cell types and receptors of human coronaviruses. *Biochem Biophys Res Commun,* 526**,** 135-140.

RAFIQ, K., CHEERS, M. S. & ETTENSOHN, C. A. 2012. The genomic regulatory control of skeletal morphogenesis in the sea urchin. *Development,* 139**,** 579-90.

RAST, J. P., SMITH, L. C., LOZA-COLL, M., HIBINO, T. & LITMAN, G. W. 2006.
Genomic insights into the immune system of the sea urchin. *Science,* 314**,** 952-6.
RIZZO, F., FERNANDEZ-SERRA, M., SQUARZONI, P., ARCHIMANDRITIS, A. &
ARNONE, M. I. 2006. Identification and developmental expression of the ets
gene family in the sea urchin (Strongylocentrotus purpuratus). *Dev Biol,* 300**,** 35-
48.
RODRIGUEZ-DIAZ, R., DANDO, R., JACQUES-SILVA, M. C., FACHADO, A., MOLINA,
J., ABDULREDA, M. H., RICORDI, C., ROPER, S. D., BERGGREN, P. O. &
CAICEDO, A. 2011. Alpha cells secrete acetylcholine as a non-neuronal
paracrine signal priming beta cell function in humans. *Nat Med,* 17**,** 888-92.
SEA URCHIN GENOME SEQUENCING, C., SODERGREN, E., WEINSTOCK, G. M.,
DAVIDSON, E. H., CAMERON, R. A., GIBBS, R. A., ANGERER, R. C.,
ANGERER, L. M., ARNONE, M. I., BURGESS, D. R., BURKE, R. D., COFFMAN,
J. A., DEAN, M., ELPHICK, M. R., ETTENSOHN, C. A., FOLTZ, K. R.,
HAMDOUN, A., HYNES, R. O., KLEIN, W. H., MARZLUFF, W., MCCLAY, D. R.,
MORRIS, R. L., MUSHEGIAN, A., RAST, J. P., SMITH, L. C., THORNDYKE, M.
C., VACQUIER, V. D., WESSEL, G. M., WRAY, G., ZHANG, L., ELSIK, C. G.,
ERMOLAEVA, O., HLAVINA, W., HOFMANN, G., KITTS, P., LANDRUM, M. J.,
MACKEY, A. J., MAGLOTT, D., PANOPOULOU, G., POUSTKA, A. J., PRUITT,
K., SAPOJNIKOV, V., SONG, X., SOUVOROV, A., SOLOVYEV, V., WEI, Z.,
WHITTAKER, C. A., WORLEY, K., DURBIN, K. J., SHEN, Y., FEDRIGO, O.,
GARFIELD, D., HAYGOOD, R., PRIMUS, A., SATIJA, R., SEVERSON, T.,
GONZALEZ-GARAY, M. L., JACKSON, A. R., MILOSAVLJEVIC, A., TONG, M.,
KILLIAN, C. E., LIVINGSTON, B. T., WILT, F. H., ADAMS, N., BELLE, R.,
CARBONNEAU, S., CHEUNG, R., CORMIER, P., COSSON, B., CROCE, J.,
FERNANDEZ-GUERRA, A., GENEVIERE, A. M., GOEL, M., KELKAR, H.,
MORALES, J., MULNER-LORILLON, O., ROBERTSON, A. J., GOLDSTONE, J.
V., COLE, B., EPEL, D., GOLD, B., HAHN, M. E., HOWARD-ASHBY, M.,
SCALLY, M., STEGEMAN, J. J., ALLGOOD, E. L., COOL, J., JUDKINS, K. M.,
MCCAFFERTY, S. S., MUSANTE, A. M., OBAR, R. A., RAWSON, A. P.,
ROSSETTI, B. J., GIBBONS, I. R., HOFFMAN, M. P., LEONE, A., ISTRAIL, S.,
MATERNA, S. C., SAMANTA, M. P., STOLC, V., et al. 2006. The genome of the
sea urchin Strongylocentrotus purpuratus. *Science,* 314**,** 941-52.
SEBE-PEDROS, A., SAUDEMONT, B., CHOMSKY, E., PLESSIER, F., MAILHE, M. P.,
RENNO, J., LOE-MIE, Y., LIFSHITZ, A., MUKAMEL, Z., SCHMUTZ, S.,
NOVAULT, S., STEINMETZ, P. R. H., SPITZ, F., TANAY, A. & MARLOW, H.
2018. Cnidarian Cell Type Diversity and Regulation Revealed by Whole-
Organism Single-Cell RNA-Seq. *Cell,* 173**,** 1520-1534 e20.
SEVERO, M. S., LANDRY, J. J. M., LINDQUIST, R. L., GOOSMANN, C., BRINKMANN,
V., COLLIER, P., HAUSER, A. E., BENES, V., HENRIKSSON, J., TEICHMANN,
S. A. & LEVASHINA, E. A. 2018. Unbiased classification of mosquito blood cells
by single-cell genomics and high-content imaging. *Proc Natl Acad Sci U S A,*
115**,** E7568-E7577.
SHARMA, S., WANG, W. & STOLFI, A. 2019. Single-cell transcriptome profiling of the
Ciona larval brain. *Dev Biol,* 448**,** 226-236.

SHEKHAR, K. & MENON, V. 2019. Identification of Cell Types from Single-Cell Transcriptomic Data. *Methods Mol Biol,* 1935**,** 45-77.

SHASHIKANT, T., & ETTENSOHN, C. A. 2019. Genome-wide analysis of chromatin accessibility using ATAC-seq. Methods in cell biology, 151, 219–235.

SIABERT, S., FARRELL, J.A., CAZET, J.F., ABEYKOON, Y., PRIMACK, A.S., SCHNITZLER, C.E., JULIANO, CE. 2019. Stem cell differentiation trajectories in Hydra resolved at single-cell resolution. Science. 365(6451)SLADITSCHEK, H. L., FIUZA, U. M., PAVLINIC, D., BENES, V., HUFNAGEL, L. & NEVEU, P. A. 2020. MorphoSeq: Full Single-Cell Transcriptome Dynamics Up to Gastrulation in a Chordate. *Cell,* 181**,** 922-935 e21.

SLOTA, L. A. & MCCLAY, D. R. 2018. Identification of neural transcription factors required for the differentiation of three neuronal subtypes in the sea urchin embryo. *Dev Biol,* 435**,** 138-149.

SLOTA, L. A., MIRANDA, E., PESKIN, B. & MCCLAY, D. R. 2020. Developmental origin of peripheral ciliary band neurons in the sea urchin embryo. *Dev Biol,* 459**,** 72-78.

SMITH, M. M., CRUZ SMITH, L., CAMERON, R. A. & URRY, L. A. 2008. The larval stages of the sea urchin, Strongylocentrotus purpuratus. *J Morphol,* 269**,** 713-33.

SOLIMAN, S. 1983. Pharmacological control of ciliary activity in the young sea urchin larva. Effects of monoaminergic agents. *Comp Biochem Physiol C Comp Pharmacol Toxicol,* 76**,** 181-91.

STEINMETZ PRH, AMAN A, KRAUS JEM, TECHNAU U. 2017. Gut-like ectodermal tissue in a sea anemone challenges germ layer homology. Nat Ecol Evol. Oct;1(10):1535-1542

STRATHMANN, M. F. 1987. *Reproduction and development of marine invertebrates of the northern Pacific coast: data and methods for the study of eggs, embryos, and larvae,* Seattle, University of Washington Press.

STUART, T., BUTLER, A., HOFFMAN, P., HAFEMEISTER, C., PAPALEXI, E., MAUCK, W. M., 3RD, HAO, Y., STOECKIUS, M., SMIBERT, P. & SATIJA, R. 2019. Comprehensive Integration of Single-Cell Data. *Cell,* 177**,** 1888-1902 e21.

SUN, Z. & ETTENSOHN, C. A. 2014. Signal-dependent regulation of the sea urchin skeletogenic gene regulatory network. *Gene Expr Patterns,* 16**,** 93-103.

SUN, Z. & ETTENSOHN, C. A. 2017. TGF-beta sensu stricto signaling regulates skeletal morphogenesis in the sea urchin embryo. *Dev Biol,* 421**,** 149-160.

SWAPNA, L.S., MOLINARO, A.M., LINDSAY-MOSHER, N., PEARSON, B.J., PARKINSON, J. 2018. Comparative transcriptomic analyses and single-cell RNA sequencing of the freshwater planarian Schmidtea mediterranea identify major cell types and pathway conservation. Genome Biol 19, 124

TANG, F., BARBACIORU, C., WANG, Y., NORDMAN, E., LEE, C., XU, N., WANG, X., BODEAU, J., TUCH, B. B., SIDDIQUI, A., LAO, K. & SURANI, M. A. 2009. mRNA-Seq whole-transcriptome analysis of a single cell. *Nat Methods,* 6**,** 377-82.

TEITELMAN, G., ALPERT, S., POLAK, J. M., MARTINEZ, A. & HANAHAN, D. 1993. Precursor cells of mouse endocrine pancreas coexpress insulin, glucagon and the neuronal proteins tyrosine hydroxylase and neuropeptide Y, but not pancreatic polypeptide. *Development,* 118**,** 1031-9.

TRAPNELL, C., WILLIAMS, B. A., PERTEA, G., MORTAZAVI, A., KWAN, G., VAN
BAREN, M. J., SALZBERG, S. L., WOLD, B. J. & PACHTER, L. 2010. Transcript
assembly and quantification by RNA-Seq reveals unannotated transcripts and
isoform switching during cell differentiation. *Nat Biotechnol,* 28**,** 511-5.
TRITSCHLER, S., THEIS, F. J., LICKERT, H. & BOTTCHER, A. 2017. Systematic
single-cell analysis provides new insights into heterogeneity and plasticity of the
pancreas. *Mol Metab,* 6**,** 974-990.
TU, Q., BROWN, C. T., DAVIDSON, E. H. & OLIVERI, P. 2006. Sea urchin Forkhead
gene family: phylogeny and embryonic expression. *Dev Biol,* 300**,** 49-62.
TU, Q., CAMERON, R. A. & DAVIDSON, E. H. 2014. Quantitative developmental
transcriptomes of the sea urchin Strongylocentrotus purpuratus. *Dev Biol,* 385**,**
160-7.
WAGNER, D. E., WEINREB, C., COLLINS, Z. M., BRIGGS, J. A., MEGASON, S. G. &
KLEIN, A. M. 2018. Single-cell mapping of gene expression landscapes and
lineage in the zebrafish embryo. *Science,* 360**,** 981-987.
WEI, Z., ANGERER, L. M. & ANGERER, R. C. 2016. Neurogenic gene regulatory
pathways in the sea urchin embryo. *Development,* 143**,** 298-305.
WEI, Z., ANGERER, R. C. & ANGERER, L. M. 2011. Direct development of neurons
within foregut endoderm of sea urchin embryos. *Proc Natl Acad Sci U S A,* 108**,**
9143-7.
WEI, Z., RANGE, R., ANGERER, R. & ANGERER, L. 2012. Axial patterning interactions
in the sea urchin embryo: suppression of nodal by Wnt1 signaling. *Development,*
139**,** 1662-9.
WEI, Z., YAGUCHI, J., YAGUCHI, S., ANGERER, R. C. & ANGERER, L. M. 2009. The
sea urchin animal pole domain is a Six3-dependent neurogenic patterning center.
Development, 136**,** 1179-89.
WESSEL, G. M. & MCCLAY, D. R. 1987. Gastrulation in the sea urchin embryo requires
the deposition of crosslinked collagen within the extracellular matrix. *Dev Biol,*
121**,** 149-65.
WOOD, N.J., T. M., WARD, E.T.M., ROWE, M.L., PERILLO, M., ARNONE, M.I. ,
ELPHICK, M.R. & OLIVERI, P. 2018. Neuropeptidergic systems in pluteus larvae
of the sea urchin Strongylocentrotus purpuratus: neurochemical complexity in a
"simple" nervous system. *Fronties in Endocrinology*.
XIMERAKIS, M., LIPNICK, S. L., INNES, B. T., SIMMONS, S. K., ADICONIS, X.,
DIONNE, D., MAYWEATHER, B. A., NGUYEN, L., NIZIOLEK, Z., OZEK, C.,
BUTTY, V. L., ISSERLIN, R., BUCHANAN, S. M., LEVINE, S. S., REGEV, A.,
BADER, G. D., LEVIN, J. Z. & RUBIN, L. L. 2019. Single-cell transcriptomic
profiling of the aging mouse brain. *Nat Neurosci,* 22**,** 1696-1708.
YAGUCHI, J., TAKEDA, N., INABA, K. & YAGUCHI, S. 2016. Cooperative Wnt-Nodal
Signals Regulate the Patterning of Anterior Neuroectoderm. *PLoS Genet,* 12**,**
e1006001.
YAGUCHI, S., YAGUCHI, J. & TANAKA, H. 2017. Troponin-I is present as an essential
component of muscles in echinoderm larvae. *Sci Rep,* 7**,** 43563.
YU, Z., LIAO, J., CHEN, Y., ZOU, C., ZHANG, H., CHENG, J., LIU, D., LI, T., ZHANG,
Q., LI, J., YANG, X., YE, Y., HUANG, Z., LONG, X., YANG, R. & MO, Z. 2019.

Single-Cell Transcriptomic Map of the Human and Mouse Bladders. *J Am Soc*
Nephrol, 30**,** 2159-2176.
YUH, C. H., LI, X., DAVIDSON, E. H. & KLEIN, W. H. 2001. Correct Expression of
spec2a in the sea urchin embryo requires both Otx and other cis-regulatory
elements. *Dev Biol,* 232**,** 424-38.
ZARET, K. S. & GROMPE, M. 2008. Generation and regeneration of cells of the liver
and pancreas. *Science,* 322**,** 1490-4.
ZENG, A., LI, H., GUO, L., GAO, X., MCKINNEY, S., WANG, Y., YU, Z., PARK, J.,
SEMERAD, C., ROSS, E., CHENG, LC., DAVIES, E., LEI, K., WANG, W.,
PERERA, A., HALL, K., PEAK, A., BOX, A.A, SANCHEZ ALVARADO, A. 2018.
Prospectively Isolated Tetraspanin+ Neoblasts Are Adult Pluripotent Stem Cells
Underlying Planaria Regeneration. Cell. 2018 Jun 14;173(7):1593-1608.e20.
ZHAO, J., ZHANG, S., LIU, Y., HE, X., QU, M., XU, G., WANG, H., HUANG, M., PAN,
J., LIU, Z., LI, Z., LIU, L. & ZHANG, Z. 2020. Single-cell RNA sequencing reveals
the heterogeneity of liver-resident immune cells in human. *Cell Discov,* 6**,** 22.

1317

Figure supplements

Figure 1 – Figure supplement 1. Overlap of the different replicates and characterization of
the undefined cluster. A) UMAPs split by original identity showing the overlap of the different
biological and technical replicates. **B)** Overlay UMAP showing the overlap of the different
biological and technical replicates. **C)** Distribution of cells per putative broad cell type. Color-code
is the same as indicated on the UMAP in Figure 1B. **C)** Table showing the percentage of WHL
gene models expressed per cell type family compared to the total number of genes we identified
(15,578). **E)** Violin plot showing the number of genes (nFeature_RNA) and number of molecules
(nCount_RNA) across the cell type families. **F)** Dotplot of the total marker genes (69) expressed
in the undefined cluster, that shows high similarity with ectodermally derived cell type families.

1328

Figure 2 – Figure supplement 1. Expression pattern of genes used to annotate the clusters.
FISH of S. purpuratus larvae with specific probes detecting the mRNAs for *Sp-Alpi* **(A)**, *Sp-IrxA*
(B), *SPU_002797* **(C)**, *SPU_021898* **(D)**, *Sp-ManrC1A* **(E)**, *Sp-Nkx6.1* **(F)**, *Sp-FoxP* **(G)**, *Sp-Emx*
(H), *Sp-Pks1* **(I)**, *SPU_016308* **(J)**, *Sp-Six1/2* **(K)**, *Sp-MacpfA2* **(L)**, *SPU_010640* **(M)**, *Sp-FoxJ2*
(N), *Sp-Chordin* **(O)** and *Sp-Tob* **(P)**. Nuclei are labelled with DAPI (in blue). All images are stacks
of merged confocal Z sections.

Figure 2 – Figure supplement 2. ScRNA-seq predicted expression patterns of genes used
to annotate the clusters. A) Dotplot showing the average expression of genes used as markers
to identify the cell type families. Cell type families are grouped based on their developmental
origins: ectodermally derived, in blue, mesodermally derived, in red, and endodermally derived,
in yellow. **B)** FISH using the antisense probe of *Sp-Hbn* (B1) paired with immunofluorescent
detection of Msp130 (B2). Merged channels of both signals show localization of Sp-Hbn
transcripts in Msp130 positive PMCs (B3). Nuclei are labelled with DAPI (in blue). All images are
stacks of merged confocal Z sections. PMCs: Primary mesenchyme cells.

Figure 2 – Figure supplement 3. ScRNA-seq is able to detect expression patterns,
previously undetectable by *in situ* hybridization. Dotplot of genes known for their involvement
in germline determination and maintenance showing overall enrichment of their average
expression in the coelomic pouches cell type family.

Figure 2 – Figure supplement 4. Proliferation status and dynamics of the larval cell type
families. A) Dotplot showing the average expression of genes encoding *cdk1*, *pcna*, DNA
polymerases, DNA ligases, condensins, and centromere proteins. The cell type families are
grouped based on their developmental origins: ectodermally derived, in blue, mesodermally
derived, in red and endodermally derived cell types, in yellow. **B)** Proliferating cells as indicated
by EdU labelling. B1) Larva labelled with EdU and the skeletogenic marker Msp130. B2) Selected
confocal sections of larva in oral view labelled with EdU. B3) Selected confocal sections showing
the internal part of the same larva as in B2 in oral view labelled with EdU. B4) Larva labelled with
EdU and the endodermal marker 5c7. Abo, Aboral ectoderm; Ap, Apical plate; Cb, Ciliary band;
Es, Esophagus; Oe, Oral ectoderm; St, Stomach.

Figure 3 – Figure supplement 1. Cell type family trees of the 3 dpf pluteus larva. Cell type
family trees reconstructed using all expressed genes **(A)** and only transcription factors **(B).**

Figure 4– Figure supplement 1. Distribution of the Zinc finger TF family members across
the cell type families. Dotplot showing the mRNA average expression of the zinc finger
transcription family members. The cell type families are grouped based on their developmental
origins: ectodermally derived, in blue, mesodermally derived, in red and endodermally derived, in
yellow.

Figure 4– Figure supplement 2. Differentially expressed TFs across cell type families.
Dotplot showing the average scaled expression of differentially expressed TFs with p-value less
than 0.5 among the 21 clusters. The cluster order is according to the transcription factor-based
tree as depicted in Figure 3-Figure supplement 1B.

Figure 5– Figure supplement 1. Reconstruction of the molecular signature of the gut at a
single cell resolution. Dotplot showing the regionalized average expression of marker genes
labeling specific endodermal domains. The scRNA-seq prediction recapitulates the previously
described compartmentalization of gene expression along the digestive tract of the 3 dpf sea
urchin larva.

Figure 6- Figure supplement 1. Subclustering of the skeletal cell type family. Dotplot
showing the mRNA average expression of 22 skeletal gene markers and their distribution among
5 skeletal subtypes.

Figure 6- Figure supplement 2. Subclustering of the immune cells cluster. A) UMAP showing
8 distinct immune cells populations as revealed by our subclustering analysis. **B)** Dotplot showing
gene markers expressed in the different immune cell population. Pigment cells gene markers are
highlighted in purple (box), while non pigmented globular cells are evident in yellow (box).

Figure 6- Figure supplement 3. Co-localization of *Sp-Fgf9/16/20* and *Sp-FgfR1* in the
cardiac sphincter region. FISH using antisense probes for *Sp-Fgf9/16/20* (magenta) and *Sp-*
FgfR1 (green). Nuclei are labelled with DAPI (in blue). All images are stacks of merged confocal
Z sections. Cs, cardiac sphincter; Es, esophagus; M, mouth.

Figure 7- Figure supplement 1. Pancreatic genes expression significance testing in the
"post oral and lateral neurons" subcluster. Barplots showing the number of genes expressed
in the actual data and the randomized data.