

Metagenomic analysis reveals the signature of gut microbiota associated with human chronotypes

Shaqed Carasso^a, Bettina Fishman^b, Liel Stelmach Lask^c, Tamar Shochat^c, Naama Geva and Eran Tauber^b ^a Department of Cell Biology and Cancer Science, Rappaport Faculty of Medicine,

Rappaport Technion Integrated Cancer Center (TICC), Haifa 3525422, Israel; ^b Department of Evolutionary and Environmental Biology and Institute of Evolution, University of Haifa, Haifa 3498838, Israel; ^c Cheryl Spencer Department of Nursing, Faculty of Social Welfare and Health Sciences, University of Haifa, Haifa 3498838, Israel. ^d Canadian Institute for Advanced Research (CIFAR), MaRS Centre, West Tower 661 University Ave., Suite 505 Toronto, ON M5G 1M1, Canada

Corresponding authors: Eran Tauber*; Naama Geva-Zatorsky; Tamar Shochat

Email: etauber@univ.haifa.ac.il; naama_gz@technion.ac.il; tshochat@univ.haifa.ac.il

* lead contact

Summary

Patterns of diurnal activity differ substantially between individuals, with early risers and late sleepers being examples of extreme chronotypes. Growing evidence suggests that the late chronotype significantly impacts the risk of developing mood disorders, obesity, diabetes, and other chronic diseases. Despite the vast potential of utilizing chronotype information for precision medicine, the factors that shape chronotypes remain poorly understood. Here, we assessed whether chronotypes are associated with different gut microbiome composition. Using metagenomic sequencing, we established a distinct signature associated with chronotype that involves two bacterial genera *Alistipes* (elevated in “larks”) and *Lachnospira* (elevated in “owls”). We have identified four metabolic pathways (e.g. gluconeogenesis) that were associated with early chronotype.

Results and Discussion

Numerous biological, mental, and behavioral functions exhibit circadian oscillations that are orchestrated by a central pacemaker in the brain. While the periodicity of these rhythms under natural conditions is uniform (24 hr), the phase of these oscillations shows considerable inter-individual variability. Phase variation is often manifested as the tendency of individuals to be active during specific times of the day and is referred to as their chronotype. There is a genetic predisposition for chronotype¹, which changes over the course of development and determines individual variation in diurnal timing of various functions, such as sleep, cognitive and physical activities².

The gut microbiome has been shown to regulate metabolic processes and to modulate brain functions and behavior via immune, endocrine and neural pathways along the gut-brain axis³. Diurnal oscillations in the function and composition of the gut microbiome have been reported^{4,5}, and in turn, metabolites of gut microbes have been shown to mediate gene expression of the circadian clock⁵. Furthermore, circadian misalignment with external light conditions induced by simulating jet lag in mice was associated with the disruption of feeding patterns, loss of diurnal microbiota rhythmicity and dysbiosis, and exacerbated metabolic imbalance including weight gain and glucose intolerance⁴. Travel-induced jetlag in humans was also associated with microbial dysbiosis, and fecal transfer from jet lagged mice and humans to germ free mice created similar metabolic impairments⁴. Overall, these findings allude to the bidirectional associations between gut microbiota and the circadian clock and begs the question as to whether variation in chronotype is also linked to distinct microbiota composition. Here, we sought to explore this link using metagenomic sequencing.

To investigate changes in the gut microbiome compositions of different chronotypes, we collected fecal samples from 133 individuals from across Israel (65 females, mean \pm SD age, 32.1 \pm 10.2 years, body mass index (BMI): 23.6 \pm 3.5). The distribution of chronotypes is shown in Figure 1A. The mid sleep time on free-days (MSF) distribution centered on 4:33 (\pm SD :34 hr) and did not differ significantly between males and females (Watson-Williams, F test, $p=0.2$). For the analysis of microbiome composition, participants were divided into three groups based on their MSF: "Early" (n = 18), "Intermediate" (n = 29) and "Late" (n = 42) chronotypes.

DNA from the fecal samples was extracted in order to conduct shotgun metagenomics sequencing. The mean library size was 3,091,458 reads (range, 174,934 – 19,829,176). Samples were filtered to include the taxa with a minimum relative abundance of >0.01% and were identified in 20% of the samples. The identified taxa in the filtered dataset were distributed into 7 phyla, 14 classes, 17 orders, 28 families, 46 genera, and 87 species. The most ubiquitous genera were the Bacteroides, Faecalibacterium, Parabacteroides, and Eubacterium (all detected in more than 96% of the samples). The ten most abundant genera are presented in Figure 1B.

We observed a higher α -diversity (Shannon index) in participants from the early compared to the late chronotype ($p < 0.05$, Kruskal-Wallis rank sum test) indicating a less complex microbiota community in the latter. There were no significant differences between early and intermediate or between intermediate and late chronotypes (Figure 1C). Chao1 indices, measuring the richness of the samples, were not significantly different between the chronotypes (Figure 1D).

Differences in individuals' gut bacteria relative abundances were tested using a feature-wise association approach, controlling for age, BMI, and gender. Two genera were significantly different between the early and late chronotypes, *Alistipes* (pFDR = 0.2, Figure 2A) and *Lachnospira* (pFDR = 0.2, Figure 2B). Within those genera, two species were significantly different between groups. *Alistipes fingoldii* were more abundant in early compared to both

intermediate and late chronotypes (pFDR = 0.2, Figure 2C). *Lachnospira pectinoschiza* were more abundant in the late compared to early chronotypes (pFDR = 0.2, Figure 2D).

Reads were also assigned to microbial metabolic pathways and differences in relative abundances were tested. In total, 15 pathways were significantly different between the groups (pFDR < 0.2), 11 of which showed a difference between intermediate to either early or late chronotypes. The four pathways that were different between early and late chronotypes (Figure 3) included the super-pathway of histidine, purine, and pyrimidine biosynthesis (pathway ID PRPP-PWY), anaerobic energy metabolism (PWY-7383), super-pathway of pyrimidine deoxyribonucleotides de novo biosynthesis (PWY-7211), and gluconeogenesis (PWY66-399). All four pathways were more abundant in early compared to late chronotypes (Figure 3).

Since a substantial factor shaping the gut microbiome composition is the diet⁶, we sought to analyze food consumption by the participants. The participants were asked to complete a questionnaire that assessed their food eating frequency. Participants reported the weekly consumption frequency of 13 types of food (e.g., beef, chicken, pastry, fruits, and vegetables) and drinks (water, sugary or diet drinks). The counts of the food items were analyzed using principal component analysis (PCA) (Figure 4A). The first principal component (PC1, 17.7%) revealed a major dietary difference between participants, where a healthy diet was found in some participants (fruits, vegetables, water), while a non-healthy diet (high-sugar, high fat, sugary drinks) was found in others. A small but highly significant correlation was found between chronotype and PC1 scores ($p < 0.01$, Figure 4B). As was reported in previous studies⁷, early chronotype individuals exhibited a higher adherence towards a healthy diet.

Overall, our analysis reveals modest but consistent differences between chronotypes. The finding that lower α -diversity, a marker of dysbiosis, was observed in late chronotypes, is consistent with growing evidence of increased cardio-metabolic morbidity and mortality risk in this group⁸. Indeed, previous studies showed that jet-lag in both humans and mice is associated with dysbiosis⁴, and since late chronotype individuals are predisposed to social jet-lag (i.e., misalignment between social and endogenous circadian clocks), the decreased microbial diversity that we observed is expected.

We identified two bacterial genera that differed between early and late chronotypes. Interestingly, *Alistipes*, which was enriched in early chronotypes, has previously been reported as being overrepresented in older mice⁹, and humans¹⁰. This may explain the well-known observation that chronotype becomes earlier with age, with MSF advancing by nearly two hours between ages 20 to 70 years¹¹. Furthermore, *Lachnospira* was more abundant in late chronotypes. In a recent human study on microbiota and eating behavior during the day¹², a higher abundance of *Lachnospira* was found when a greater percentage of energy was consumed after 2pm. This finding aligns with our data portraying *Lachnospira* as a biomarker for the late chronotype, as energy consumption is expected to be delayed in these individuals¹³.

Finally, our study underscores the role of several metabolic pathways that differ between chronotypes, particularly gluconeogenesis (Figure 3). In fact, Gluconeogenesis gluconeogenesis is known to be under tight circadian regulation of the host, where expression of gluconeogenesis associated genes is stimulated upon anticipation of nightly fasting¹⁴. This suggests an interaction between the host and the microbiome metabolic pathways that is likely to be chronotype-dependent.

Materials and Methods

This study was approved by the local ethics committees at the University of Haifa (#283/18). Participants provided written informed consent prior to participation.

The study population consisted of 133 individuals (51% males), evaluated for body mass index (BMI) and medication intake. Chronotype was assessed by computing mid sleep time on free-days (MSF) (2). Stool samples were collected by the participants using the DNA/RNA Shield Fecal Collection tubes (Zymo research). DNA extraction of the samples was performed using the PureLink™ Microbiome DNA Purification Kit (Invitrogen) according to the manufacturer's instructions. The genomic DNA was sheared to an average size of 300 bp with an M220 ultrasonicator (Covaris, Woburn, MA, USA). Sheared DNA samples were used for paired-end indexed library construction using Ovation Ultralow library systems V2 (NuGEN, San Carlos, CA, USA), according to the manufacturer instructions.

Next generation sequencing (Paired end, 2X150bp) was performed by the University of Illinois at Chicago, Core for Research, using the Illumina NextSeq500. Metagenomic sequencing data was processed with bioBakery workflows utilizing the bioBakery 3 tools¹⁵. Briefly, sequence data quality control, including removal of human reads, was conducted using KneadData. Taxonomic profiles were generated using MetaPhlAn v3.0 and functional profiles were generated with HUMAnN v3.0 using MetaCyc pathway definitions¹⁶. Relative proportions data was transformed by centered-log ratio. Microbial communities were compared between different chronotypes. The primary analysis was performed using MicrobiomeAnalyst¹⁷ followed by comprehensive analysis using the R packages Phyloseq¹⁸, Vegan, and MaAsLin2 v1.06. Alpha diversity indices (Chao1 and Shannon) were compared using the Wilcoxon rank sum test. Beta diversity distance matrices (Aitchison) were compared using the Vegan package's function ADONIS, a multivariate analysis of variance based on dissimilarity tests. Differences in microbial taxa and functional modules were assessed using differential abundance analyses with MaAsLin2. Results were visualized using the ggplot2 R package. The matrices were visualized using principal coordinate analysis (PCA).

Raw metagenomic data are deposited in the National Center for Biotechnology Information sequence read archive: BioProject accession is PRJNA714678.

Acknowledgments

We thank Bengisu Sezen Subaşı for her technical assistance, and the N.G-Z lab for discussions and help. This study was supported by a research grant from the Technion- University of Haifa joint projects funded the Milgrom family to E.T, T.S and N. G-Z. The Israel Science Foundation grant (1737/17) to Etfor research supplies. N.G-Z is supported by CIFAR, Azrieli-Global Scholar; the Technion, "Hanasi", "Cathedral", "Alon" and "Horev"; and Human Frontiers (HFSP CDA00025/2019-C).

Author Contributions: ET, TS and NGZ conceived the project, NGZ contributed to the design and implementation of the research. BF and TS recruited study participants, BF carried out data collection (stool samples and questionnaires) and ran the experiments. SC carried out the bioinformatics analysis. All authors discussed the results and contributed to the final manuscript.

Declaration of Interests: The authors declare no competing interest.

References

1. Jones, S.E., Lane, J.M., Wood, A.R., van Hees, V.T., Tyrrell, J., Beaumont, R.N., Jeffries, A.R., Dashti, H.S., Hillsdon, M., Ruth, K.S., et al. (2019). Genome-wide association analyses of chronotype in 697,828 individuals provides insights into circadian rhythms. *Nat. Commun.* 10, 343.
2. Roenneberg, T., Wirz-Justice, A., and Mellow, M. (2003). Life between clocks: Daily temporal patterns of human chronotypes. *J. Biol. Rhythms* 18, 80–90.
3. Mu, C., Yang, Y., and Zhu, W. (2016). Gut microbiota: The brain peacekeeper. *Front. Microbiol.* 7.
4. Thaïss, C.A., Zeevi, D., Levy, M., Zilberman-Schapira, G., Suez, J., Tengeler, A.C., Abramson, L., Katz, M.N., Korem, T., Zmora, N., et al. (2014). Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. *Cell* 159, 514–529.
5. Leone, V., Gibbons, S.M., Martinez, K., Hutchison, A.L., Huang, E.Y., Cham, C.M., Pierre, J.F., Heneghan, A.F., Nadimpalli, A., Hubert, N., et al. (2015). Effects of diurnal variation of gut microbes and high-fat feeding on host circadian clock function and metabolism. *Cell Host Microbe* 17, 681–689.
6. Rothschild, D., Weissbrod, O., Barkan, E., Kurilshikov, A., Korem, T., Zeevi, D., Costea, P.I., Godneva, A., Kalka, I.N., Bar, N., et al. (2018). Environment dominates over host genetics in shaping human gut microbiota. *Nature* 555, 210–215.
7. Maukonen, M., Kanerva, N., Partonen, T., Kronholm, E., Kontinen, H., Wennman, H., and Männistö, S. (2016). The associations between chronotype, a healthy diet and obesity. *Chronobiol. Int.* 33, 972–981.
8. Knutson, K.L., and von Schantz, M. (2018). Associations between chronotype, morbidity and mortality in the UK Biobank cohort. *Chronobiol. Int.* 35, 1045–1053.
9. Langille, M.G.I., Meehan, C.J., Koenig, J.E., Dhanani, A.S., Rose, R.A., Howlett, S.E., and Beiko, R.G. (2014). Microbial shifts in the aging mouse gut. *Morgan. Microbiome* 2, 50.
10. Claesson, M.J., Jeffery, I.B., Conde, S., Power, S.E., O'connor, E.M., Cusack, S., Harris, H.M.B., Coakley, M., Lakshminarayanan, B., O'sullivan, O., et al. (2012). Gut microbiota composition correlates with diet and health in the elderly. *Nature* 488, 178–184.
11. Fischer, D., Lombardi, D.A., Marucci-Wellman, H., and Roenneberg, T. (2017). Chronotypes in the US – Influence of age and sex. *PLoS One* 12.
12. Kaczmarek, J.L., Musaad, S.M., and Holscher, H.D. (2017). Time of day and eating behaviors are associated with the composition and function of the human gastrointestinal microbiota. *Am. J. Clin. Nutr.* 106, ajcn156380.
13. Xiao, Q., Garaulet, M., and Scheer, F.A.J.L. (2019). Meal timing and obesity: interactions with macronutrient intake and chronotype. *Int. J. Obes.* 43, 1701–1711.
14. Panda, S., Antoch, M.P., Miller, B.H., Su, A.I., Schook, A.B., Straume, M., Schultz, P.G., Kay, S.A., Takahashi, J.S., and Hogenesch, J.B. (2002). Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell* 109, 307–320.
15. McIver, L.J., Abu-Ali, G., Franzosa, E.A., Schwager, R., Morgan, X.C., Waldron, L., Segata, N., and Huttenhower, C. (2018). BioBakery: A meta'omic analysis environment.

Bioinformatics 34, 1235–1237.

16. Caspi, R., Billington, R., Fulcher, C.A., Keseler, I.M., Kothari, A., Krummenacker, M., Latendresse, M., Midford, P.E., Ong, Q., Ong, W.K., et al. (2018). The MetaCyc database of metabolic pathways and enzymes. *Nucleic Acids Res.* 46, D633–D639.
17. Dhariwal, A., Chong, J., Habib, S., King, I.L., Agellon, L.B., and Xia, J. (2017). MicrobiomeAnalyst: A web-based tool for comprehensive statistical, visual and meta-analysis of microbiome data. *Nucleic Acids Res.* 45, W180–W188.
18. McMurdie, P.J., and Holmes, S. (2015). Shiny-phyloseq: Web application for interactive microbiome analysis with provenance tracking. *Bioinformatics* 31, 282–283.

Figures




Figure 1. (A) Chronotype distribution. Chronotype were divided into 7 categories, Extremely early (EE): <01:29, Moderately early (ME) 1:30-2:30, SE 2:30-3:30, Intermediate (IM) 3:30-4:30, Slightly late (SL) 4:30-5:30, Moderately late (ML) 5:30-6:30, Extremely late (EL) > 6:30. **(B)** Microbiome composition (genus level) in the different chronotype. The relative abundance of main taxa in early (E) intermediate (IM) and late (L) chronotype is shown. **(C)+(D)** Alpha diversity was measured via Chao1 index (C) and Shannon index (D) in early (E) intermediate (IM) and late (L) chronotype. Data represent the median (line in box), IQR (box), and minimum/maximum (whiskers). Statistical comparisons shown for separate matched-pairs tests with the Kruskal-Wallis rank sum test corrected using the Benjamini–Hochberg FDR method.




Figure 2. Relative abundances of significantly different taxa in early (E) intermediate (IM) and late (L) chronotype. Data represent the median (line in box), IQR (box), and minimum/maximum (whiskers).




Figure 3. Relative abundances of significantly different metabolic pathways in in early (E) intermediate (IM) and late (L) chronotype. Data represent the median (line in box), IQR (box), and minimum/maximum (whiskers).




Figure 4. Principal component analysis (PCA) of food and drink items consumed by the participants. B. Correlation between chronotype (MSF) with scores of PC1 from the PCA analysis showing a significant trend ($p < 0.01$). Fruits, vegetables (and water as a drink) are prominent in early chronotypes, while meat, pastry and sugary drinks are associated with late chronotypes.