1 **Rat ultrasonic vocalizations and novelty-induced social and non-social investigation**

2 **behavior in a seminatural environment**

4 Indrek Heinla[1], Xi Chu[2], Anders Ågmo[1], Eelke Snoeren[1]

5 [1] Department of Psychology, UiT The Arctic University of Norway, Norway

6 [2] Department of Psychology, Norwegian University of Science and Technology, Norway

20 Corresponding author: Eelke M.S. Snoeren

21 Current address: Department of Psychology, UiT the Arctic University of Norway, 9037
22 Tromsø, Norway

23 E-mail: eelke.snoeren@uit.no

Abstract

Although rats are known to emit ultrasonic vocalizations (USVs), it remains unclear whether these calls serve an auditory communication purpose. For USVs to be part of communication, the vocal signals will need to be a transfer of information between two or more conspecifics, and with the possibility to induce changes in the behavior of the recipient. Therefore, the aim of our study was to investigate the role of USVs in adult rats' social and non-social investigation strategies when introduced into a large novel environment with unfamiliar conspecifics. We quantified a wide range of social and non-social behaviors in the seminatural environment, which could be affected by subtle signals, including USVs. We found that during the first hour in the seminatural environment the ability to vocalize did not affect how quickly adult rats met each other, their overall social investigation behavior, their passive social behavior nor their aggressive behavior. Furthermore, the non-social exploratory behaviors and behaviors reflecting anxiety/stress-like states were also unaffected. These results demonstrated that a disability to vocalize did not result in significant disadvantages (or changes) compared to intact conspecifics regarding social and non-social behaviors. This suggests that other (multi)sensory cues are more relevant in social interactions than USVs.

Keywords: Rat, Ultrasonic vocalization, Social, Behavior, Seminatural environment

Introduction

Many animals communicate through vocalization, and the understanding of how and why animals communicate has long been fascinating to scientists [1]. Information encoded by vocal cues has diverse behavioral significance depending on the species. They can, for instance, serve a role in mating rituals, act as warning calls, convey location of food sources, or play a role in influencing the behavior of an interacting partner (reviewed in [2]). The fact that rats can produce vocal signals as audible squeals in the range of 2-4 kHz and ultrasonic vocalizations (USVs, up to ~80 kHz) has been known for a long time [3]. However, researchers are still attempting to understand the structure and function of these calls.

Adult rats emit two main types of ultrasonic vocalizations: the low 22 kHz and the high 50 kHz calls. The 22 kHz calls are assumed to function as alarm calls, since they have been observed mostly in aversive situations/contexts (reviewed in [4]). The 50 kHz calls (ranging between 30-80 kHz), on the other hand, reflecting appetitive calls, are emitted in the presence of for instance a sexual partner and during copulation [5-7], during social [8] and play behavior [9, 10], or after administration of hedonic drugs [11, 12].

Although USVs are reported to be emitted before, during, and/or after certain events, the exact role of these vocalizations to the relevant event is not self-explanatory. Many researchers have proposed that the USVs serve a communicative role, but in order for the vocalizations to be part of *communication*, the vocal signals will need to be a transfer of information between two or more conspecifics, and with the possibility to induce changes in the behavior of the recipient. So far, the empirical evidence remains unsatisfying on whether USVs play a communicative role. Some evidence pointing in the direction of a communicative function is shown by the fact that playback of pre-recorded 22 kHz resulted in defensive behavior of the recipient [13]. With regards to 50 kHz calls, though, different studies have found opposite results. Some have shown that the playback of 50 kHz calls

induces transient approach behavior in rats, especially juveniles [14-17], while others have

demonstrated that the playback of vocalizations from a conspecific of the opposite sex does

not induce approach behavior in male nor female adult rats [18, 19]. In addition, it was found

that when the emission or receiving of the USVs is disrupted (e.g. by devocalization or

deafening), rats hardly elicit different patterns of behavior in their partners [20-23]. Only in

juvenile rats, different patterns of play behavior have been found in dyads of silent versus

vocalizing rats [24].

In addition, rodents constantly interact with their conspecifics, using different means

of communication. Their social behavior consists of more different elements of behaviors

besides the approach or play behaviors mentioned above. In a broad sense, social behaviors

can be defined as any modality of communication and/or interaction between conspecifics of

a given species (see review [25]). Social behavior displayed at the inappropriate time or place

or of unfitting intensity can have negative consequences to the individuals even to a social

group as a whole. These interactions involve active detection and response to cues from

multiple sensory modalities, and a continuous exchange of social information perceived from

sensory cues produces an important feedback loop that could change the behavioral responses

again. Since the complexity of interactions depends on the potential communication space

between individuals, social behaviors are among the most complex behaviors. Unlike some

other communication modalities, USV communication has strong directivity, low energy

consumption, thus they can be effective over a wide range of distances [26], which makes

USVs an interesting candidate for a communicative function in social behavior in rats.

Surprisingly, studies on the role of USVs in social interaction in rats are rare, and the

studies that are performed (mainly studying play behavior in juveniles) make use of

traditional test settings in which rats are placed in a small arena without the opportunity to

express their full repertoire of behavior or interact with multiple conspecifics [24, 27, 28]. As

it has been suggested that USVs are used as social-locational cues (providing information

about the other conspecifics nearby and their whereabouts) [29], one could argue that for

USVs to play a communicative role in social behavior, more space might need to be required

than is available in traditional set-ups, for some of these cues to have any significance.

Though, previously we have reported that silencing rats with devocalization

procedures did not significantly affect sexual behavior or social interactions, via sniffing

behavior, in rat tested in a seminatural environment [21]. As sexual behavior is probably one

of the most relevant behavior in which social-locational cues should play a major role, this

suggests that USVs do not play an essential role in social interaction. However, the rats in this

study were already present the environment for 7 days and were therefore already familiar

with each other at the moment of testing. It is hypothetically possible that the rats had already

adapted to the communication limitations and modified their interaction behaviors. In

addition, individuals with disabled social and communication abilities could perform

normally in some situations, whereas, when posed with novel situations, they might

experience higher levels of stress and need longer time to adjust to the circumstances. In

combination with the idea that appropriate communication and social interaction is probably

most important upon first encounter, it would be interesting to look at the role of USVs when

rats are introduced to a novel seminatural environment with unfamiliar conspecifics.

Therefore, the aim of our study was to investigate the role of USVs in rats' social and

non-social investigation strategies when introduced into a novel large environment with

unfamiliar conspecifics. We quantified a wide range of social and non-social behaviors in the

seminatural environment, which could be affected by subtle signals, including USVs. As

tracking of the individual's USVs within a group of rats comes with its own challenges,

especially in a large arena, our current study used devocalized and sham-operated vocalizing

male and female rats. Another advantage of this approach is that we were able to investigate a

batch in which some rats were completely silent. If the emission of USVs plays a role in

social investigation behavior, our test conditions should be ideal to detect differences in

behavior. Based on our previous findings, we expected that devocalized rats would overall

show similar social investigation patterns as sham-operated vocalizing controls in our

naturalistic set-up. However, at the same time, we expected that *if* USVs are indeed used as

means of communication, it would be most visible during the first encounters with unfamiliar

rats. Devocalized rats should then for instance be approached less by others than vocalizing

rats.

Methods

The data was collected from video recordings obtained in a previously performed experiment, resulting in the same materials and methods described previously [21]. The differences between the current and previous study are the behavioral scoring scheme that were used and timing of the observations. In the previous study, the role of USVs in sexual behavior were investigated, while the current study focuses on the role of USVs in other social and non-social behaviors. In addition, in the current study we analyzed the behavior during the first hour after introduction into the seminatural environment when the environment and conspecifics are still novel, whereas the previous study investigated the behaviors on day 7, after they had been familiarized to the new environment.

Animals

A total of 16 female and 12 male Wistar rats (250–300g upon arrival at an age of ca. 3 months) were obtained from Charles River (Sulzfeld, Germany). Before testing, the animals were housed in same-sex pairs in Macrolon IV open cages (so all the animals were used to hearing vocalizations in the animal room) with tap water and commercial rat pellets available *ad libitum*. All rats had obtained one sexual experience in a copulation test prior to the experiment [21, 30]. The experiment was conducted in accordance with European Union directive 2010/63/EU and was approved by the National Animal Research Authority (ID 5441). The rats were around an age of 3 months at the start of the experiment.

Surgeries

The procedures were described previously in [21]. Briefly, all females were ovariectomized upon arrival. Operations were done under isoflurane anesthesia and afterwards rats were checked twice daily for 3 days and treated with 0.05 mg/kg buprenorphine every 12 hours (subcutaneously). After obtaining one session of sexual

experience two weeks after ovariectomy, seven females and five males were devocalized 3

weeks before they entered the seminatural environment (DEV). Two-centimeter incision was

made on the ventral surface of the neck, sternohyoideus muscles were separated and trachea

exposed. Next, recurrent laryngeal nerves were cleared from fascia and bilaterally 3mm

section of the nerve was removed. The control rats (CTR) received sham surgery (similar

procedure, but the nerve was left intact). All animals recovered well from the surgeries.

Seminatural environment

The seminatural environment (2.4 x 2.1 x 0.75 meters) setup is previously described

and illustrated in [31-34]. It consists of a burrow system and an open field area, which are

connected by four 8 x 8 cm openings. The burrow system consists of an interconnected tunnel

maze (7.6 cm wide and 8 cm high) with 4 nest boxes (20 x 20 x 20 cm) attached, and is

covered with Plexiglas. The open area has 75cm high walls, and contains two partitions (40 x

75 cm) to simulate obstacles in nature. A light blocking wall (made of light blocking cloth)

between the burrow and the open field allows the light intensity for both arenas to be

controlled separately. The burrow system remained in total darkness for the duration of the

experiment, while a day-night cycle was simulated in the open area with a lamp 2.5 m above

the center that provided 180 lux from 22.45h to 10.30h and approximately 1 lux from 10.30h

to 11.00h (the equivalent of moonlight). The light gradually increased/decreased during 30

minutes between 1 and 180 lux.

The floors of both the open area and on the burrow system were covered with a 2 cm

layer of aspen wood chip bedding (Tapvei, Harjumaa, Estonia). In addition, the nest boxes

were provided with 6 squares of nesting material each (nonwoven hemp fibres, 5 x 5 cm, 0.5

cm thick, Datesend, Manchester, UK), and the open area was equipped with 3 red

polycarbonate shelters (15 x 16.5 x 8.5 cm, Datesend, Manchester, UK) and 12 aspen wooden

sticks (2 x 2 x 10 cm, Tapvei, Harjumaa, Estonia). Food was provided in one large pile of

approximately 2 kg in the open area close to the water supply. Water was available ad libitum

in four water bottles.

Two video cameras (VCC-6592; Sanyo, Tokyo, Japan) equipped with a zoom lens

(T6Z5710-CS 5.7–34.2 mm; Computar, San Jose, CA, USA) were mounted on the ceiling 2

meters above the seminatural environment to capture the whole environment: one above the

open field and another above the burrow system. Infrared lamps provided light for the video

camera centered above the burrow.

Procedure and design

Shortly before (circa 72 hours) being introduced into the seminatural environment, the

sham and devocalized males and females were tested for the presence or absence of

vocalizations, respectively. As previously described, the male and female rats (who were

sexually receptive at this point) were placed in two adjacent chambers covered with sound-

absorbing isolation material of extruded polyethylene foam and separated by a wire mesh. A

high-frequency sensible microphone (Metris, Hoofddorp, Netherlands) was placed above each

chamber and adjusted so that all sounds from within the chamber were recorded, while sounds

from the adjacent chamber were not captured by the microphone. The microphone was

connected to a computer with the Sonotrack sound analysis system. All devocalized rats used

in this experiment did not emit any USV, while the sham animals did. In addition, on another

experimental day we have also confirmed the presence of USVs in the seminatural

environment.

The day before introduction to the seminatural environment, the subjects' backs were

shaved and tails marked for individual recognition. A rectangle, about 2 x 3 cm, was carefully

shaved on the back of the rats the day before introduction into the environment. One female

had the rectangle close to the tail, another in the middle of the back, and a third had it close to

the neck. The fourth female was not marked on the back. In addition, the tail was marked with

one, two, or three transversal, thick black lines. The fourth female was not marked. Males

were marked exactly as the females, except that the tail marks were made larger to distinguish

between males and females.

Four cohorts of four females and three males were used (resulting in a total number of

9 control females, 7 devocalized females, 7 control males and 5 devocalized males; see

Supplementary Table 1). Animals in each cohort came from different cages to ensure that

they were previously unfamiliar to each other.

Each cohort lived in the seminatural environment for a total of 8 days with full-time

recording of all behaviors. After the experiment, the rats were removed from the seminatural

environment, the environment was thoroughly cleaned and bedding/nesting materials and

food were changed, before a new cohort was introduced.

Behavioral observation

An experienced observer, blinded for the treatment of rats, scored the behavioral

activity of each rat with Noldus Observer XT (Netherlands) during the first 60 minutes after

introduction to the seminatural environment. One of 18 different behaviors (see Table 1) was

assigned to each rat at any time. Where possible, up to four clarifying modifiers were added:

(1) the location where the behavior took place, (2) the partner/recipient of the behavior, (3) if

there was a tactile contact with another rat or not and (4) if the given animal initiated the

behavior or responded to another rat.

Table 1 Description of recorded behaviors

Behavior	Description
Walking/running	Walking or running through the environment
Chasing	Running forward in the direction of a conspecific
Non-social exploration	Exploring the environment by sniffing, usually when slowly walking or sitting still
Interacting with environment	Digging, pushing or carrying bedding/nesting/food material
Passive alone	Sitting or sleeping with minimal movement of the head without other rats in close vicinity
Passive socially	Sitting or sleeping with minimal movement of the head with at least 1 other rat on maximum 1 rat body length away
Hiding alone	Being in the shelter alone
Hiding socially	Being in the shelter with at least one other rat
Allogrooming	Grooming any part of a conspecific's body, usually on the head or in the neck region
Sniffing anogenitally	Sniffing the anogenital region of the conspecific
Sniffing nose-to-nose	Sniffing the facial region of the conspecific
Sniffing body/head	Sniffing any part of the conspecifics body or head, except for the anogenital and nose region
Fighting	Kicking, pouncing, pushing, grabbing, boxing or wrestling another rat
Nose-off	Facing another rat and aggressively posturing towards it
Self-grooming	Grooming itself
Rearing supported	Raising itself upright on its hind paws, facing a wall or an object
Rearing unsupported	Raising itself upright on its hind paws, not facing a wall or an object
Any other behavior	Behaviors that do not fit any of the other categories (e.g. mounting, drinking, etc)

235
236 **Table 2 Description of behavioral clusters**

Cluster	Behaviors within clusters
Social investigation	Sniffing anogenitally, sniffing nose-to-nose, sniffing body/head, and allogrooming
Non-social investigation	Walking/running, non-social exploration
Conflict behaviors	Nose-off, fighting
Passive behaviors	Passive alone, passive socially
Social passive behaviors	Passive socially, hiding socially,
Non-social passive behaviors	Passive alone, hiding alone
All passive behaviors	Social and non-social passive behaviors
Hiding	Hiding alone, hiding socially
All sniffing	Sniffing anogenitally, sniffing nose-to-nose, sniffing body/head
All rearing	Rearing supported, rearing unsupported

Data preparation and analysis

From the behavioral output file, the following behavioral clusters were generated (see Table 2): *social investigation* (consisting of sniffing anogenitally, sniffing nose-to-nose, sniffing body/head and allogrooming), *non-social investigation* (consisting of walking/running and non-social exploration), *conflict behaviors* (consisting of fighting and nose-off), *passive behaviors* (consisting of passive alone and passive socially), *social passive behaviors* (consisting of passive socially and hiding socially), *non-social passive behaviors* (consisting of passive alone and hiding alone), *all passive behaviors* (consisting of passive alone, passive socially, hiding alone and hiding socially), *hiding* (consisting of hiding alone and hiding socially), *all sniffing* (consisting of sniffing anogenitally, sniffing nose-to-nose and sniffing body/head) and *all rearing* (consisting of rearing supported and rearing unsupported). For these behavioral clusters, but also for the individual behaviors, the latency, frequency, duration and mean duration of episode of each behavior or behavioral cluster for the whole hour in the whole arena were calculated, along with the same parameters separated by location (burrow area versus open area) and into 10-minute time-bins.

To further investigate social behavior, it was also calculated how fast each rat met each of their conspecifics, the mean duration of their social interactions, how much time overall did they spend in tactile contact with conspecifics, ratio of social activity (time in social behaviors/overall time), ratio of active non-social behavior (non social investigation/immobility), ratios of different types of sniffing, percentage of unsupported rearing (unsupported rearing/all rearing), and how much time they spent on open arena doing non-social behaviors.

For analysis of the data of the whole hour, a linear mixed model with rat as subject and *treatment* and *sex* as factors was used (IBM SPSS Statistics 26). A modified Benjamini-Hochberg procedure was used (instead of using all possible comparisons, which would yield

too strict criteria for behavioral data, we only used p-values of four predetermined clusters: all

sniffing, non-social investigation, self-grooming and conflict behavior, in addition to all

behaviors with $p<0.05$) to correct for multiple comparison analysis. The data separated into

10-minute time-bins was analyzed with a repeated measures ANOVA with *time* as a within-

subject factor and *treatment* and *sex* as between-subject factors. If Mauchly's test of sphericity

yielded $p<0.05$, Greenhouse-Geisser test of within-subjects effects is reported, otherwise if

Mauchly's test of sphericity yielded n.s., sphericity assumed test of within-subjects effects is

reported.

One devocalized female rat was excluded from the analysis because she spent an

overwhelming majority of the time passively (87% of the overall time, in comparison to

others with on average $2.8\pm .3\%$). The reason remains unclear, but therefore the data

throughout the manuscript is presented without this rat.

Statement Open Science Framework (OSF)

The design of our study was preregistered on OSF on the 17th of December 2019

(https://osf.io/gzkjw). We refrained from the analysis of entry and re-entry latencies of

different parts of the environment, because first rats were entered into the environment before

starting the videos and therefore we were not able to collect complete data; otherwise there

were no changes in analysis.

Results

Since our data analysis generated a lot of data, only the most relevant findings from the total environment in this section are reported. For more details on different aspects of the data, or the data from the open area and burrow alone, please turn to the supplementary Tables 2-5.

Social investigation

As mentioned in the introduction, social behavior is a complex behavior that involves multiple aspects. Besides the different categories of social behavior, it also involves the interaction between two or more animals and thus the differentiation in whether a rat is the initiator or responder to a social interaction. To investigate the role of USVs in social behavior, parameters linked to social behavior were explored. The time it took to meet all new conspecifics, the frequency and duration of social behaviors in total as well as initiator or responder, the length of social interaction bouts and the frequency, duration and average time they were being socially investigated were calculated. In addition, it was analyzed how much of these episodes contained actual tactile contact. Interestingly, no differences were found in any of these parameters between silent (DEV) and vocalizing (CTR) female and/or male rats.

First of all, it was found that cohort members met very quickly, as most animals had actively sniffed more than half of their new cohort members within the first minute and had mostly approached all six of their new cohabitants within the first 5 minutes. No differences were found between CTR and DEV animals in terms of latency to approach new conspecifics or being approached by conspecifics (effect of treatment $F_{(1,23)}=.196$; n.s. Fig 1A). In addition, no differences were found between time spent on social investigation behavior (effect of treatment on social investigation $F_{(1,23)=.}039$; n.s. Fig 1B) or its separate subcomponents of social behaviors between CTR and DEV rats (Fig S1A, B, C, Supplementary Table 2). Male

rats only spent in general more time sniffing the anogenital or body regions than female rats,

but no significant treatment*sex interaction effect was found (see Supplementary Tables

2&3). Similar results were found in the time receiving social investigation behaviors (or its

subcomponents) from conspecifics (without necessarily responding to it: effect of treatment

on social investigation behavior $F_{(1,23)}=.007$; n.s. Fig 1C) or with regard to the length of the

social interaction bouts (effect of treatment: dyads with DEV rat $F_{(1,23)}<0.01$, n.s.; dyads with

CTR rat $F_{(1,23)}=0.81$, n.s., DEV-DEV vs CTR-CTR dyads $F_{(1,23)}=1.923$; n.s. Fig S1L, M). Not

even when only the first 10 encounters were analyzed separately effect of treatment

$F_{(1,23)}=2.298$; n.s. Fig S1H, Supplementary Table 2).

Also when the social behaviors were divided in the episodes in which a rat was the

initiator versus the responder or with/without tactile contact, DEV rats initiated (effect of

treatment $F_{(1,23)}=.496$; n.s. Fig 1D) and spent a similar amount of time on initiated social

behaviors (effect of treatment $F_{(1,23)}=.218$; n.s. Fig S1D) as CTR rats. Similarly, there were no

differences in episodes of responding to others (effect of treatment $F_{(1,23)}=.011$; n.s. Fig 1E) or

duration of responding to others ($F_{(1,23)}==.001$; n.s.) in social investigation behavior. It should

be mentioned, though, that it is sometimes unclear in a seminatural environment which animal

initiates the interaction. This limitation was solved by scoring both participants of the social

interaction as initiators. Moreover, it was found that the overall time spent *with* tactile contact

(effect of treatment $F_{(1,23)}=,016$; n.s. Fig 1F) and the average length of these interactions were

not different in CTR and DEV rats (Fig S1I, Supplementary Table 2).

Furthermore, the data revealed no differences in any other behavior involving a

conspecific that could have been affected by devocalization, such as following, passive

socially and conflict behavior. There was no difference between vocalizing and silent animals

in how much time they were being followed (effect of treatment $F_{(1,23)}=.024$; n.s. Fig 1G) or

how much they followed others (effect of treatment $F=.005$; n.s. Fig 1H). Also, when the data

at whom they follow (behaviors following DEV and following CTR rats are corrected

according to the number of available partners in a given cohort; Fig S1J, K) was analyzed, no

significant differences were found. The data analysis of following behavior only revealed a

significant sex effects in that female rats were more often being followed (effect of sex

$F_{(1,23)}=4.96$; p=.04) and males doing most of the following (effect of sex $F_{(1,23)}=17.32$;

p<.001). However, there was no significant interaction effect between treatment and sex

(Supplementary Tables 2&3). Additionally, it was found that silent DEV rats spent a

comparable amount of time on passive social behavior (and its subcomponents) to vocalizing

CTR rats (effect of treatment $F_{(1,23)}=.085$; n.s. Fig 1I), neither were there differences on the

time spent on conflict behavior of DEV and CTR rats, neither as an active partner nor as

receiving the conflict (effect of treatment as aggressive party $F_{(1,23)}=.413$; n.s. Fig 1J and

effect of treatment as recipient $F_{(1,23)}=.024$; n.s. Fig 1K, refer to the Supplementary Table 2

for mean values).

Figure 1. *Social behavior of devocalized (DEV, n=11) and sham-operated control (CTR, n=16) rats. (A) The number of different conspecifics that were met within 1, 3 ,5, 10 and 20 minutes. (B) Time spent on social investigation (sniffing anogenitally, sniffing nose-to-nose, sniffing body/head, and allogrooming). (C) Time being socially investigated by conspecifics. (D) Number of initiated social interactions. (E) Number of responses to a social interaction initiated by a conspecific. (F) Time spent in tactile contact with other rats. (G) Time being followed by a conspecific. (H) Time spent on following other rats. (I) Time spent on being passive socially (hiding and passive socially). (J) Time spent on conflict behavior (fighting or nose-off) with conspecifics. (K) Time receiving conflict behaviors from conspecifics. Data are shown with individual data points (females in grey, males in blue) with the lines representing the group means. Error bars are representing standard error of the mean SEM. s = seconds, no = number of episodes.*

Non-social investigation and other behaviors

Besides social behaviors, USVs could also affect emotional state of the vocalizing

animal itself, which could then influence their non-social investigation patterns in a novel

environment or their stress-coping behavior. For example, if USVs had a comforting effect on

the rat itself, one could hypothesize that CTR rats might feel safer to explore the novel

environment than a DEV rats. Therefore, our study also investigated the non-social

investigation strategies of the rats, in addition to parameters like self-grooming, rearing, and

time spent in the open area.

However, analysis of the overall time spent investigating the environment (effect of

treatment $F_{(1,23)}$=.612; n.s. Fig 2A), in addition to the separate subcomponents

walking/running (effect of treatment $F_{(1,23)}$=.30; n.s.) and non-social exploration (effect of

treatment $F_{(1,23)}$=.33; n.s.), did not reveal any differences between CTR and DEV rats. There

was, though, a sex effect showing that females spent more time on non-social investigation

than males (effect of sex $F_{(1,23)}$=14.27; p=.001), but no interaction effect between sex and

treatment was found.

Rearing to hind legs provides superior vantage point to investigate the surrounding

social and physical environment. A distinction between supported and unsupported rearing

was made with the idea that unsupported rearing has been shown to be modulated by anxiety-

like states [35]. If emitting USVs would induce a comforting effect, one could assume that

CTR rats show more unsupported rearing. In our experiment, however, no effect was found

on supported rearing nor unsupported rearing (supported rearing: effect of treatment F=.087;

n.s. Fig 2B; unsupported rearing: effect of treatment $F_{(1,23)}$=.703; n.s. Fig 2C). Also when

unsupported and supported rearing were combined, no differences between CTR and DEV

were found (effect of treatment $F_{(1,23)}$=.267; n.s. Fig S1N), except that females rear more often

than males (effect of sex $F_{(1,23)}$=5.786; p=.025).

Other behaviors that could be linked to anxiety-like states and could thus theoretically

be affected by USVs if these play a role on emotional state, are behaviors like digging,

transporting the bedding material, nesting material and food (combined in the cluster

"interaction with the environment"), self-grooming and the time spent in open arena. Data

analysis revealed, though, that there were no effects of the absence of USVs on interacting

with the environment (effect of treatment $F_{(1,23)}=.173$; n.s. Fig 2D), nor on the amount of time

spent in the open arena including (effect of treatmnet $F_{(1,23)}=.839$; n.s.) or excluding the

episodes in which they participated in social interactions (effect of treatment $F_{(1,23)}=.735$;

n.s.). It was found, though, that male rats spent in general more time in open area compared to

females (including social interactions: effect of sex $F_{(1,23)}=17.008$; $p<.001$; excluding social

interaction: effect of sex $F_{(1,23)}=14.403$; $p=.001$; Fig 2E). Female rats, on the other hand, spent

more time in the burrow (tunnels and nestboxes; effect of sex $F_{(1,23)}=16.432$; $p<.001$), but no

effects of treatment were found between CTR and DEV rats (effect of treatment $F_{(1,23)}=.879$;

n.s. Fig 2G).

With regard to self-grooming, a potential measure for stress-coping behavior [36-38],

male rats self-groomed more ($F_{(1,23)}=13.68$; $p=.001$) and longer ($F_{(1,23)}=13.41$; $p<.001$ Fig 2F)

than female rats, but no effect of treatment (number of episodes $F_{(1,23)}=.164$; n.s.; time spent

$F_{(1,23)}=.92$; n.s.) or interaction effects of sex*treatment were found.

It should be mentioned, though, that anxiety-like states can be accompanied by

behavioral inhibition, which can manifest in delayed onset of natural maintenance and

exploratory behaviors. But when we compared the latencies to start self-grooming (effect of

treatment $F_{(1,23)}=.337$; $p=.57$, Fig S1O), unsupported rearing (effect of treatment

$F_{(1,23)}=.09$; $p=.77$) or other behaviors (Supplementary Table 4), no differences between CTR

and DEV rats were found.

412

Figure 2. *Non-social behavior of devocalized (DEV, n=11) and sham-operated control (CTR,*

414 *n=16) rats. (A) Time spent on non-social investigation behavior. (B-C) Time spent on rearing*

415 *suppoted and unsupported. (D)Time spent on interacting with the environment. (E) Time*

416 *spent in the open area (OA) (excluding social interactions)(F) Time spent on self-grooming.*

417 *(G) Relative time spent in the different areas of the environment. The height of the colored*

418 *box represents the proportion of time the rats of the given group on average spent in*

419 *respective area. Data in A-F are shown with individual data points (females in grey, males in*

420 *blue) with the lines representing the group means. Error bars are representing standard error*

421 *of the mean SEM. s = seconds.*

422

423 *Behavioral patterns during the course of an hour*

424   At last, it was investigated how the behavioral patterns of the rats changed over the

425 course of the hour to detect if there are any deviations in how devocalized animals habituate

426 to the novel social and non-social environment. Therefore, the data was divided into six 10-

427 minute time-bins and analyzed the behavioral patterns cumulatively.

428   As expected, some behaviors were performed more or less in the beginning than in the

429 end. The amount of time spent on social investigation (effect of time $F_{(5,115)}=6.74$; $p<0.001$;

430 Fig 3A), being socially investigated (time effect $F_{(5,115)}=5.899$; $p<.001$; Fig S2A, Fig

S3C&D), and non-social investigation (effect of time $F_{(3.322,76.399)}=12.03$; p<0.001; Fig 3B)

slightly decreased over the course of an hour, whereas the time spent on rearing (effect of

time $F_{(5,115)}=2.013$; n.s. Fig S2D, especially unsupported rearing: effect of time $F_{(5,115)}=2.726$;

p=0.023), self-grooming (effect of time $F_{(3.039,69.896)}=13.26$; p<0.001; Fig S2B), and passive

behavior ($F_{(2.908,66.874)}=4.20$; p=0.009; Fig S2C) increased over the course of an hour.

Some sex differences were found in these behavioral patterns: male rats showed a

steeper decrease in time being socially investigated over the hour (time*sex interaction effect

$F_{(1.223,28.132)}=6.274$; p=.014) and a faster increase in self-grooming behavior (effect of

time*sex interaction $F_{(1.419,32.647)}=14.82$; p<001) compared to females, while females declined

faster in the time spent on non-social investigation (effect of time*sex interaction

$F_{(1.562,35.917)}=13.57$; p<001). With regard to rearing, females reared more at the beginning of

the experiment and less near the end (effect of time*sex interaction $F_{(1.621,37.294)}=3.636$;

p=.045; post-hoc: males vs females for first ten minutes and second ten minutes p=.037; for

40-50 minutes p=.021), while males were initially rearing less with support compared to

females (effect of time*sex interaction $F_{(1.851,42.572)}=5.213$; p=.011). But no remarkable

interaction effects with treatment (CTR versus DEV) were found.

Only in terms of the amount of time rats spent on social investigation behavior, it was

found that DEV rats spent slightly less time on these behaviors within the first 10 minutes

compared to CTR rats (p=.012), but this effect disappeared immediately and resulted in an

overall lack of interaction effect over the course of an hour (time*treatment interaction

$F_{(1.172,26.949)}=.11$; n.s.). Besides, none of the subcomponents of time spent on social

investigation showed differences between CTR and DEV rats when analyzed separately.

When the data was further divided into 1-minute time-bins, it became clear that the tendency

towards a difference in social investigation behavior between CTR and DEV rats occurs in the

minutes between 3 and 12 (Fig S3A&B), after which the DEV rats catch up again with the

CTR rats.

With regard to rearing, there was no overall time*treatment effect (effect of time*

treatment interaction $F_{(1.621,37.294)}=.03$; n.s.). However, silent rats did rear significantly more

within the first ten minutes compared to vocalizing rats (p=.007). This effect was probably

caused by supported rearing (p=.006). Further analysis into 1-minute time-bins revealed that

the difference in supported rearing between CTR and DEV rats was present around the 1st to

10th minute, after which they show comparable amount of rearing again (Fig S3E&F).

A summary of the main findings described below can be found in Table 3.

Figure 3. Behavioral patterns during the course of an hour in devocalized (DEV, n=11) and

sham-operated control (CTR, n=16) rats. (A) The cumulative time spent on social

investigation behavior. (B) The cumulative time spent on non-social investigation behavior.

Data are shown in mean±standard error of the mean per 10-miute time-bins. s = seconds, *

p<0.05 CTR versus DEV.

Table 3. Summary of main findings

No effects were found between CTR and DEV rats on the following parameters of social behaviors:
- latency to approach new conspecifics
- time spent on social investigation behavior
- time receiving social investigation behavior from conspecifics
- length of social investigation bouts
- time spent on social behavior as initiator or responder
- overall time spent with tactile contact
- average length of social interactions
- time spent on social passive behavior
- time spent on conflict behavior
- time spent on following behavior

No effects were found between CTR and DEV rats on the following parameters of non-social behaviors:
- overall time spent investigating the environment
- time spent rearing
- time spent interacting with the environment
- time spent self-grooming
- time spent in open area during non-social behaviors

Discussion

In our study, we investigated the role of USVs in social interactions and non-social

investigation of a novel environment with unfamiliar conspecifics in adult rats. Our findings

show that silent and vocalizing rats behave very similarly in the first hour of exposure to the

new environment. We found no differences in social interaction and non-social investigation

behaviors between sham and devocalized rats. Silent rats spent comparable amount of the

time on social interactions as vocalizing rats, independent of whether they were the initiator

or the receiver. In addition, silent and vocalizing rats also familiarized in the same way with

new neighbor conspecifics and novel environment, respectively.

This is in line with our hypothesis that was based on the findings of previous studies in

which devocalization did not have an effect on sociosexual behavior with familiar rats [20-

23]. Interestingly, though, our study is also in line with another recent study by Redecker et

al. who have studied the social behavior and USV production of heterozygous (Cacna1c+/−)

and wildtype (Cacna1c+/+) rats [27], a genetic modification of calcium voltage-gated channel

subunit that have been linked to deficits in social behavior in mice [39]. Upon their

expectations, they found that Cacna1c+/− rats emitted less USVs during social interactions

than the controls. However, although their auditory cues were reduced, the rats, both mutated

and wild type, did not show any differences in social behavior, measured as sniffing,

following, social grooming and crawling under/over [27]. Our studies therefore both show

that a reduction (or depletion) of USVs does not affect social interaction behavior.

Our experiment revealed that the emission of USVs did not affect rats' approach

behavior in the seminatural environment. This is somewhat contradicting with previous

studies that have shown a temporary approach behavior to the playback of 50 kHz calls [14,

15, 40, 41]. However, we have not been able to replicate these findings on approach behavior

even in a smaller arena [18, 19], and when rats were able to choose between an intact or

devocalized conspecific, the silent rats were just as much approached and preferred as play or

sexual partner as the vocalizing rats in traditional test settings [6, 19, 40, 42]. Therefore, it

remains unclear what the function and significance of this sort of short-lasting approach

behavior is.

If USVs indeed modulate rats' social interactions and induce approach behavior, it

would mean that rats that are incapable of vocalizing should be approached less than intact

conspecifics. Additionally, *if* USVs could act as a reinforcer of the behavior, the bouts of

social interaction between two vocalizing rats should last longer than bouts between dyads

from which one or both are devocalized. Another possibility could be that differences would

have been found in the approach behavior towards which part of the body (anogenital region,

body, nose) is targeted in devocalized and vocalizing rats. Consequently, if USVs played a

role in modulating rats' social interactions, vocalizing rats should perform and/or receive

more interactions compared to devocalized rats. However, in our experiment, there were no

differences in how quickly devocalized and vocalizing control rats met their cohort members

nor in any parameters regarding approach. Even though the vocalizing animals showed a

tendency towards increased social investigation early in the experiment, devocalized animals

displayed comparable amount of social investigation at the beginning and throughout the

hour. It seems that the transient approach behavior, which has been reported in several

playback studies, is not visible in a more naturalistic settings with adult rats. Interestingly,

these previously reported approach behaviors have only been seen once per each individual

and lasted only short, which would then be in line with our findings. Since the effect on

approach has previously been found strongly in juvenile rats, and we have indeed replicated

this approach (not published), it is also possible that the role of USVs could differ during

lifetime. In general, social behavior of the adult rats in our experiment was not affected by

their own nor their partner's ability to vocalize. This supports the idea that USVs do not play

a significant role in modulating communication in adult rats.

This conclusion makes us wonder what the function of these calls then could be.

Could, for instance, the ability to vocalize modulate rat's own or partner's emotional state?

One study has shown how rats, that have been trained to react to different sounds to either

earn a positive reward (sucrose) or to avoid unpleasant loud white noise, treat an ambiguous

cue as positive (predicting reward) if it is preceded by the playback of 50 kHz calls and treat

similar cue as negative (predicting unpleasant white noise) if preceded by 22 kHz calls [43].

This implies that 22 kHz and 50 kHz calls are indeed involved in inducing negative and

positive responses. In our study, however, we investigated whether potential feedback from

vocalizing rat would change dynamics of the social interaction such as length of the

interaction, preference for tactile contact or escalation to aggression. Interestingly, in the

study by Redecker et al. it was found that the Cacna1c+/− rats, who have reduced USVs

emission, did spend more time in physical contact than the Cacna1c+/+ rats [27]. However,

our findings did not show any signs of changes in physical contact, type of contact or

escalation to aggression upon devocalization. The differences in results could then be

explained by the use of a large seminatural environment in which rats are able to choose the

type of interaction they prefer in the moment, instead of being forced into a certain behavior.

Another possibility is that vocalizing itself can have a comforting effect on a rat and

that devocalization could thus influence their stress-coping and/or non-social investigation

behavior in the novel environment. For example, if USVs modulate anxiety/stress-like states

of the emitter, one could hypothesize that vocalizing rats should be more comfortable to

initially explore the environment, rear more frequently, and spend more time in anxiogenic

parts of the enclosure, and/or self-groom less than devocalized rats with less experiencing the

comforting effect of emitting USVs. Such self-comforting effect was indeed reported in the

study of Cacna1c+/− rats, as the mutated animals self-groom more, show less digging

behavior and rearings when interacting in pairs than the control rats [27]. In the current

context, however, we again found no differences between vocalizing and silent rats in terms

of self-grooming or manipulating the environment (including digging). This does not

necessarily contradict the previous findings, since the knockout strain Cacna1c+/− rats with a

dysfunctional calcium voltage-gated channel subunit alpha 1 C could have different

underlying reasons for this change in behavior. However, our findings at least suggest that the

emission of USVs does not modulate stress-related behaviors. At the same time, it is still

possible that USV emission is initiated by the same internal state that also facilitates the given

behaviors, something we would not be able to see in our devocalized rats. The emitted USVs

could then also in theory be a by-product of the given behaviors, which would then indicate

that a change in these behaviors result in a reduced number of USVs, but not the other way

around.

In our data, we did find an initial increase in supported rearing in devocalized rats. Our

exploratory methods are not suitable to explain whether this increase in exploratory rearing is

related to the reduced social investigation in the same time window (they can only perform

one behavior at the same time), and could then just as well be explained as an unfortunate

artifact. Unsupported rearing, which is linked to susceptibility to acute stress [35], was not

affected by devocalization. Along with the lack of effects in the other behavioral parameters

that could reflect anxiety/stress-like states such as self-grooming (time spent and latency to

start) and the time spent in the anxiogenic parts of the environment, our data suggests that the

ability to vocalize does not modulate rats' anxiety/stress-like states.

Previously, we and others have suggested that this could mean that USVs may be

purely a by-product of the arousal linked to the behaviors [30, 44, 45], and that 50-kHz USVs

are just a by-product of locomotion and breathing. It should be mentioned, though, that the

advances in research techniques have now made it possible to study this possibility in more

detail and resulted in the conclusion that USVs are not just simply a byproduct. Evidence

showed that the emitted USVs are indeed tightly linked to locomotion [29], breathing [46]

and cardiovascular function [45], and they are even interlocked with active sniffing [47].

However, the fact that they also actively sniff without the emission of USVs [47], and can

both vocalize without movement and move without vocalizing [29] weakens the argument for

a by-product effect. Besides, the vocal production apparently increases before locomotion

begins [29], and a new call type can be started at any point during the exhalation phase [46,

48, 49]. It should be taken into account, though, that *if* USVs are more than just a by-product

of arousal, there should be more information in nuances of the vocal communication, as they

have an extensive USV 'vocabulary' [50, 51]). So far, many studies have neglected the

existence of this vocabulary, and the possible role different type of calls, and the sequence of

calls, must have if USVs serve a communicative role after all. Moreover, some studies in

juvenile rats that did investigate different types of calls did not find an one-to-one relationship

between any movements and specific vocalizations, nor did they show a direct change in the

conspecific's behavior [10, 52]. Thus, altogether we conclude that USVs are unlikely

functioning for communication, neither are they involved in regulating non-social exploring

behaviors.

It is important to mention, though we failed to find USVs' effect, that our study does

not *exclude* the possibility that vocalizations play a communicative role in social and non-

social behavior. It could simply be the case that other (multi)sensory cues are more relevant in

these interactions, and compensate for the lack of vocalizations, something that we have

shown before with approach behavior in a sexual context [53]. It could still be possible that if

rats are never exposed to auditory stimuli, they would fail to socially interact normally. It was

shown by Kisko and colleagues that not only devocalized juvenile rats played less, but also

that intact rats housed with devocalized rats showed reduced levels of play behavior [24].

They suggested that rats could have a critical period in which the lack of exposure to

vocalizations could determine their behavior later in life. This is an interesting theory that

should be explored in the future, but our findings at least support the notion that vocalizations

are not the most essential way of communication later in life.

In conclusion, our data show that devocalized adult rats do not show altered social

interaction behaviors due to their inability to vocalize. Silent and vocalizing rats show similar

patterns and types of social interactions, and do not use other social and non-social

investigation strategies when introduced to a novel environment with unfamiliar conspecifics.

Our data, therefore, does not provide any evidence that USVs play a communicative role in

social behavior, nor do they serve a role in regulating non-social investigation behaviors.

Although it cannot be excluded that USVs play some unrevealed role in social behavior, it is

clear that other non-USV sensory cues are more relevant in these interactions and could have

compensated for the lack of vocalizations. New interesting research techniques using complex

algorithms to link behaviors to distinct pattens of USVs, as those used nowadays for mice

[54], are needed in the future to explore the potential role of USVs in social behavior in

naturalistic environments.

Acknowledgements

We also would like to thank Ragnhild Osnes, Carina Sørensen, Nina Løvhaug, Katrine

Harjo, and Remi Osnes for their excellent care of the animals.

References:

[1] Darwin, C. The descent of man, and selection in relation to sex. New York,: D. Appleton and company; 1871.

[2] Witzany, G. Biocommunication of animals. 1st ed. Dordrecht: Springer Netherlands : Imprint: Springer,; 2014. p. 1 online resource (XIII, 420 pages 68 illustrations, 44 illustrations in color.

[3] Anderson, J. W. The production of ultrasonic sounds by laboratory rats and other mammals. Science. 1954,119:808.

[4] Wöhr, M., Schwarting, R. K. W. Affective communication in rodents: Ultrasonic vocalizations as a tool for research on emotion and motivation. Cell and Tissue Research. 2013,354:81-97.

[5] Bialy, M., Rydz, M., Kaczmarek, L. Precontact 50-khz vocalizations in male rats during acquisition of sexual experience. Behavioral Neuroscience. 2000,114:983-90.

[6] Snoeren, E. M., Ågmo, A. The role of odors and ultrasonic vocalizations in female rat (rattus norvegicus) partner choice. J Comp Psychol. 2014,128:367-77.

[7] Snoeren, E. M., Helander, L. R., Iversen, E. E., Ågmo, A. On the role of individual differences in female odor and ultrasonic vocalizations for male's choice of partner. Physiol Behav. 2014,132:17-23.

[8] Brudzynski, S. M., Pniak, A. Social contacts and production of 50-khz short ultrasonic calls in adult rats. J Comp Psychol. 2002,116:73-82.

[9] Knutson, B., Burgdorf, J., Panksepp, J. Anticipation of play elicits high-frequency ultrasonic vocalizations in young rats. J Comp Psychol. 1998,112:65-73.

[10] Burke, C. J., Kisko, T. M., Swiftwolfe, H., Pellis, S. M., Euston, D. R. Specific 50-khz vocalizations are tightly linked to particular types of behavior in juvenile rats anticipating play. PLoS One. 2017,12:e0175841.

[11] Mu, P., Fuchs, T., Saal, D. B., Sorg, B. A., Dong, Y., Panksepp, J. Repeated cocaine exposure induces sensitization of ultrasonic vocalization in rats. Neurosci Lett. 2009,453:31-5.

[12] Williams, S. N., Undieh, A. S. Brain-derived neurotrophic factor signaling modulates cocaine induction of reward-associated ultrasonic vocalization in rats. J Pharmacol Exp Ther. 2010,332:463-8.

[13] Fendt, M., Brosch, M., Wernecke, K. E. A., Willadsen, M., Wöhr, M. Predator odour but not tmt induces 22-khz ultrasonic vocalizations in rats that lead to defensive behaviours in conspecifics upon replay. Scientific reports. 2018,8:11041.

[14] Wöhr, M., Schwarting, R. K. W. Ultrasonic communication in rats: Can playback of 50-khz calls induce approach behavior? PLOS ONE. 2007,2:e1365.

[15] Schönfeld, L.-M., Zech, M.-P., Schäble, S., Wöhr, M., Kalenscher, T. Lesions of the rat basolateral amygdala reduce the behavioral response to ultrasonic vocalizations. Behav Brain Res. 2020,378:112274.

[16] Willadsen, M., Seffer, D., Schwarting, R. K., Wöhr, M. Rodent ultrasonic communication: Male prosocial 50-khz ultrasonic vocalizations elicit social approach behavior in female rats (rattus norvegicus). J Comp Psychol. 2014,128:56-64.

[17] Melo-Thomas, L., Tonelli, L. C., Müller, C. P., Wöhr, M., Schwarting, R. K. W. Playback of 50-khz ultrasonic vocalizations overcomes psychomotor deficits induced by sub-chronic haloperidol treatment in rats. Psychopharmacology (Berl). 2020,237:2043-53.

[18] Snoeren, E. M. S., Ågmo, A. The incentive value of males' 50-khz ultrasonic vocalizations for female rats (rattus norvegicus). Journal of Comparative Psychology. 2014,128:40-55.

[19] Snoeren, E. M., Ågmo, A. Female ultrasonic vocalizations have no incentive value for male rats. Behav Neurosci. 2013,127:439-50.

[20] Snoeren, E. M. S., Ågmo, A. The role of odors and ultrasonic vocalizations in female rat (rattus norvegicus) partner choice. Journal of Comparative Psychology. 2014,128:367-77.

[21] Chu, X., Snoeren, E., Agmo, A. Functions of vocalization in sociosexual behaviors in rats (rattus norvegicus) in a seminatural environment. J Comp Psychol. 2017,131:10-8.

[22] Thomas, D. A., Takahashi, L. K., Barfield, R. J. Analysis of ultrasonic vocalizations emitted by intruders during aggressive encounters among rats (rattus norvegicus). J Comp Psychol. 1983,97:201-6.

[23] White, N. R., Barfield, R. J. Role of the ultrasonic vocalization of the female rat (rattus norvegicus) in sexual behavior. US: American Psychological Association; 1987. p. 73-81.

[24] Kisko, T. M., Wöhr, M., Pellis, V. C., Pellis, S. M. From play to aggression: High-frequency 50-khz ultrasonic vocalizations as play and appeasement signals in rats. In: Wöhr M, Krach S, eds. Social behavior from rodents to humans: Neural foundations and clinical implications. Cham: Springer International Publishing; 2017. p. 91-108.

[25] Chen, P., Hong, W. Neural circuit mechanisms of social behavior. Neuron. 2018,98:16-30.

[26] Titze, I. R., Palaparthi, A. Radiation efficiency for long-range vocal communication in mammals and birds. J Acoust Soc Am. 2018,143:2813.

[27] Redecker, T. M., Kisko, T. M., Schwarting, R. K. W., Wöhr, M. Effects of cacna1c haploinsufficiency on social interaction behavior and 50-khz ultrasonic vocalizations in adult female rats. Behav Brain Res. 2019,367:35-52.

[28] Ågmo, A., Snoeren, E. M. S. Silent or vocalizing rats copulate in a similar manner. PLOS ONE. 2015,10:e0144164.

[29] Laplagne, D. A., Elías Costa, M. Rats synchronize locomotion with ultrasonic vocalizations at the subsecond time scale. Frontiers in behavioral neuroscience. 2016,10:184-.

[30] Ågmo, A., Snoeren, E. M. Silent or vocalizing rats copulate in a similar manner. PLoS One. 2015,10:e0144164.

[31] Chu, X., Agmo, A. Sociosexual behaviours in cycling, intact female rats (rattus norvegicus) housed in a seminatural environment. Behaviour. 2014,151:1143-84.

[32] Heinla, I., Heijkoop, R., Houwing, D. J., Olivier, J. D. A., Snoeren, E. M. S. Third-party prosocial behavior in adult female rats is impaired after perinatal fluoxetine exposure. Physiol Behav. 2020,222:112899.

[33] Houwing, D. J., Heijkoop, R., Olivier, J. D. A., Snoeren, E. M. S. Perinatal fluoxetine exposure changes social and stress-coping behavior in adult rats housed in a seminatural environment. Neuropharmacology. 2019,151:84-97.

[34] Snoeren, E. M., Antonio-Cabrera, E., Spiteri, T., Musatov, S., Ogawa, S., Pfaff, D. W., et al. Role of oestrogen alpha receptors in sociosexual behaviour in female rats housed in a seminatural environment. J Neuroendocrinol. 2015,27:803-18.

[35] Sturman, O., Germain, P. L., Bohacek, J. Exploratory rearing: A context- and stress-sensitive behavior recorded in the open-field test. Stress. 2018,21:443-52.

[36] Spruijt, B. M., van Hooff, J. A., Gispen, W. H. Ethology and neurobiology of grooming behavior. Physiological reviews. 1992,72:825-52.

[37] Dunn, A. J., Berridge, C. W., Lai, Y. I., Yachabach, T. L. Crf-induced excessive grooming behavior in rats and mice. Peptides. 1987,8:841-4.

[38] Choleris, E., Thomas, A. W., Kavaliers, M., Prato, F. S. A detailed ethological analysis of the mouse open field test: Effects of diazepam, chlordiazepoxide and an extremely low frequency pulsed magnetic field. Neurosci Biobehav Rev. 2001,25:235-60.

[39] Kabir, Z. D., Che, A., Fischer, D. K., Rice, R. C., Rizzo, B. K., Byrne, M., et al. Rescue of impaired sociability and anxiety-like behavior in adult cacna1c-deficient mice by pharmacologically targeting eif2α. Molecular psychiatry. 2017,22:1096-109.

[40] Seffer, D., Rippberger, H., Schwarting, R. K. W., Wöhr, M. Pro-social 50-khz ultrasonic communication in rats: Post-weaning but not post-adolescent social isolation leads to social impairments-phenotypic rescue by re-socialization. Frontiers in behavioral neuroscience. 2015,9:102-.

[41] Willuhn, I., Tose, A., Wanat, M. J., Hart, A. S., Hollon, N. G., Phillips, P. E. M., et al. Phasic dopamine release in the nucleus accumbens in response to pro-social 50 khz ultrasonic vocalizations in rats. The Journal of Neuroscience. 2014,34:10616.

[42] Snoeren, E. M., Ågmo, A. The incentive value of males' 50-khz ultrasonic vocalizations for female rats (*rattus norvegicus*). J Comp Psychol. 2014,128:40-55.

[43] Saito, Y., Yuki, S., Seki, Y., Kagawa, H., Okanoya, K. Cognitive bias in rats evoked by ultrasonic vocalizations suggests emotional contagion. Behavioural processes. 2016,132:5-11.

[44] Blumberg, M. S., Alberts, J. R. Ultrasonic vocalizations by rat pups in the cold: An acoustic by-product of laryngeal braking? Behavioral neuroscience. 1990,104:808-17.

[45] Blumberg, M. S., Sokoloff, G. Do infant rats cry? Psychological Review. 2001,108:83-95.

[46] Riede, T., Schaefer, C., Stein, A. Role of deep breaths in ultrasonic vocal production of sprague-dawley rats. J Neurophysiol. 2020,123:966-79.

[47] Sirotin, Y. B., Costa, M. E., Laplagne, D. A. Rodent ultrasonic vocalizations are bound to active sniffing behavior. Front Behav Neurosci. 2014,8:399-.

[48] Riede, T. Rat ultrasonic vocalization shows features of a modular behavior. J Neurosci. 2014,34:6874-8.

[49] Hernandez, C., Sabin, M., Riede, T. Rats concatenate 22 khz and 50 khz calls into a single utterance. The Journal of experimental biology. 2017,220:814-21.

[50] Wright, J. M., Gourdon, J. C., Clarke, P. B. S. Identification of multiple call categories within the rich repertoire of adult rat 50-khz ultrasonic vocalizations: Effects of amphetamine and social context. Psychopharmacology (Berl). 2010,211:1-13.

[51] Takahashi, N., Kashino, M., Hironaka, N. Structure of rat ultrasonic vocalizations and its relevance to behavior. PLOS ONE. 2010,5:e14115.

[52] Burke, C. J., Kisko, T. M., Pellis, S. M., Euston, D. R. Avoiding escalation from play to aggression in adult male rats: The role of ultrasonic calls. Behavioural processes. 2017,144:72-81.

[53] Agmo, A., Snoeren, E. M. A cooperative function for multisensory stimuli in the induction of approach behavior of a potential mate. PLoS One. 2017,12:e0174339.

[54] Sangiamo, D. T., Warren, M. R., Neunuebel, J. P. Ultrasonic signals associated with different types of social behavior of mice. Nature neuroscience. 2020,23:411-22.