TITLE; Manganese toxicity disrupts indole acetic acid homeostasis and suppresses

CO₂ assimilation reaction in rice plants

RUNNING TITLE; *Mechanism of symplastic Mn toxicity in rice*

AUTHORS; Daisuke Takagi[1,2]*, Keiki Ishiyama[2], Mao Suganami[2 ‡], Tomokazu Ushijima[1],

Takeshi Fujii[1], Youshi Tazoe[2, †], Michio Kawasaki[1], Ko Noguchi[3], Amane Makino[2]

AUTHORS AFFILIATIONS;

[1] Faculty of Agriculture, Setsunan University, Hirakata, Osaka 573-0101, Japan

[2] Graduate School of Agricultural Science, Tohoku University, Sendai, Miyagi 980-8572,

Japan

[3] Department of Applied Life Science, School of Life Sciences, Tokyo University of

Pharmacy and Life Sciences, Hachioji, Tokyo, 192-0392 Japan

[†] Present address; Faculty of Agro-Food Science, Niigata Agro-Food University, Tainai,

Niigata, 959-2702 Japan

[‡] Present address; Faculty of Food and Agricultural Sciences, Fukushima University,

Kanayagawa, Fukushima, 960-1296, Japan

*Corresponding author

E-MAIL ADDRESSES

Daisuke   Takagi   (daisuke.takagi@setsunan.ac.jp);   Keiki   Ishiyama

(keiki.ishiyama.e5@tohoku.ac.jp); Mao Suganami (mao.suganami@agri.fukushima-u.ac.jp);

Tomokazu   Ushijima   (tomokazu.ushijima@setsunan.ac.jp);   Takeshi   Fujii

(takeshi.fujii@setsunan.ac.jp); Youshi Tazoe (yushi-tazoe@nafu.ac.jp); Michio Kawasaki

(michio.kawasaki@setsunan.ac.jp); Ko Noguchi (knoguchi@toyaku.ac.jp); Amane Makino

(amanemakino@tohoku.ac.jp);

○ **Date of submission:** 17 March 2021

○ **Number of tables and figures:** no tables and 9 figures

○ **Word counts**: 5739 words (excluding Materials and Methods)

○ **Number of supplemental tables and figures:** 2 tables and 4 figures

HIGHLIGHT (18 words)

Increased Mn concentration lowers auxin concentrations in rice leaves, which suppresses photosynthesis by changing stomatal function and development.

ABSTRACT (176 words)

Despite the essentiality of Mn in terrestrial plants, its excessive accumulation in plant tissues causes growth defects, known as Mn toxicity. Mn toxicity can be divided into apoplastic and symplastic types depending on its onset. For growth defects, symplastic rather than apoplastic Mn toxicity is hypothesised to be more critical. However, details of the relationship between growth defects and symplastic Mn toxicity remains elusive. In this study, we aimed to elucidate the molecular mechanisms of symplastic Mn toxicity in rice plants. We found that under excess Mn conditions, CO_2 assimilation was inhibited by stomatal closure, and both carbon anabolic and catabolic activities were decreased. In addition to stomatal dysfunction, stomatal and leaf anatomical development were also altered by excess Mn accumulation. Furthermore, the indole acetic acid (IAA) concentration was decreased, and auxin-responsive gene expression analyses showed IAA-deficient symptoms in leaves due to excess Mn accumulation. These results suggest that excessive Mn accumulation causes IAA deficiency, and low IAA concentrations suppress plant growth by suppressing stomatal opening and leaf anatomical development for efficient CO_2 assimilation in leaves.

KEYWORDS

amino acid metabolism, auxin, bundle-sheath cell, leaf anatomy, manganese, photorespiration, photosynthesis, rice, respiration, stomata,

ABBREVIATIONS

AOX, alternative oxidase; ARF, auxin response factor; AUX/IAA, auxin/indoleacetic acid; Cc, chloroplastic CO_2 partial pressure; COX, cytochrome *c* oxidase; Ci, internal CO_2 concentration in leaves; DAO, dioxygenase for auxin oxidation; D.W., dry weight; F.W., fresh weight; Fv/Fm, maximum quantum yield of photosystem II; GC-MS, gas chromatography-mass spectrometry; g_s, stomatal conductance; IAA, indole acetic acid; ICP-OES, inductively coupled plasma optical emission spectroscopy; HPLC, high-performance liquid chromatography; PODs, class III peroxidases; PS, photosystem; ROS, reactive oxygen species; SEM, scanning electron microscope; SOD, superoxide dismutase; TCA, tricarboxylic acid; Y(II), quantum yield of photosystem II; Y(I), quantum yield of photosystem I; Y(NPQ), quantum yield of non-photochemical quenching; Y(NO), quantum yield of non-radiative decay; Y(ND), quantum yield of non-photochemical quenching at the donor-side of photosystem I; Y(NA), quantum yield of non-photochemical quenching at the acceptor-side of photosystem I

INTRODUCTION

Mn is an essential nutrient for both terrestrial plants and animals (Shao *et al.*, 2017). With regard to terrestrial plants, Mn was first discovered in their ash, and McHargue (1922) proved that Mn is an essential nutrient. Manganese possesses a wide variety of physiological functions in plant cells. For example, Mn activates more than 35 different enzymes such as chloroplast RNA-polymerase, and several enzymes involved in the tricarboxylic acid (TCA) cycle and shikimic acid pathway (Broadley *et al.*, 2012). Mn is also directly involved in the physiological function of Mn-superoxide dismutase (SOD) as a cofactor to detoxify reactive oxygen species (ROS) and oxalate oxidase (Broadley *et al.*, 2012). Among these physiological functions, the Mn cluster within the oxygen-evolving complex of photosystem II (PSII) is crucial for driving photosynthesis in terrestrial plants (Suga *et al.*, 2015).

Although Mn is indispensable for terrestrial plants, excess Mn accumulation in leaves reduces their growth and crop yield, through what is known as Mn toxicity (Foy *et al.*, 1978; Li *et al.*, 2019). The toxic effects of Mn are observed in various terrestrial plants, but the critical concentration for expressing toxicity varies depending on plant species and genotype (Foy *et al.*, 1978; Horst *et al.*, 1999; Horiguchi*, 1987; Millaleo *et al.*, 2010). For example, *Zea mays* L. shows signs of Mn toxicity at an accumulation of 200 μg Mn g^{-1} (dry weight; D.W.) in the leaves; in contrast, *Lupinus albus* L. or woody Mn hyperaccumulator species, such as *Gossia bidwillii*, can accumulate more than 10,000 μg Mn g^{-1} (D.W.) without Mn toxicity symptoms (Blamey *et al.*, 1986, 2015; Fernando *et al.*, 2006). These differences in the critical concentration for Mn toxicity are derived from the different capacities of Mn compartments within cells (Blamey *et al.*, 2015). Excessive Mn is sequestrated to vacuoles in a chelated form with organic acids (malate/citrate) to maintain the Mn concentration within cells (Blamey *et al.*, 2015). However, when the vacuole capacity for storing Mn reaches the upper limit, the Mn concentration increases in the symplastic and apoplastic regions, where Mn toxicity occurs (Horiguchi, 1987; Führs *et al.*, 2009). In fact, Mn-sensitive plants show a smaller vacuole capacity for retaining Mn than do Mn-tolerant

plants (Horst *et al*., 1999; Blamey *et al*., 2015). This interpretation has been validated by

loss-of-function and overexpression mutants of tonoplast-localised Mn transporter proteins

such as metal tolerant protein (MTP) 8 and calcium exchanger (CAX) 2, which show a

susceptible and tolerant phenotype to Mn toxicity, respectively (Hirschi *et al*., 2000; Delhaize

et al., 2003; Chen *et al*., 2013; Eroglu *et al*., 2016). In addition to these mechanisms,

Mn-tolerant plants, such as *Helianthus annuus* L., use trichomes as Mn storage tissue to

avoid an excessive increase in Mn concentration within cells (Blamey *et al*., 1986, 2015).

However, this mechanism depends on the plant species and is not a general strategy to

prevent Mn toxicity (Blamey *et al.,* 2015).

Mn toxicity can be divided into two categories depending on the cell part of its

onset: apoplastic and symplastic (Fernando *et al*., 2016). The primary symptoms of

apoplastic Mn toxicity in terrestrial plants are brown spots containing oxidised phenolics,

oxidised Mn [Mn^{3+} and Mn^{4+}], and callose in the leaves (Kenten and Mann, 1957; Horiguchi,

1987; Wissemeir and Horst, 1987, 1992; Horst *et al*., 1999; Fecht-Christoffers *et al*., 2003*a,*

b; Blamey *et al*., 2015). Mn accumulation in the apoplast stimulates the expression of class

III peroxidases (PODs), which undertake both H_2O_2 production through NADH-peroxidase

activity and H_2O_2 consumption through guaiacol-peroxidase activity (Fecht-Christoffers *et al.,*

2003a, b; Führs *et al.,* 2009). Furthermore, the concentrations of phenolics, which suppress

the NADH-peroxidase activity, are lowered in the apoplast and, as a consequence, H_2O_2

production is accelerated in the apoplast (Fecht-Christoffers *et al.,* 2006; Führs *et al.,* 2009).

The H_2O_2 consumption reaction by the guaiacol activity of PODs also proceeds by utilising

phenolics in the apoplast; subsequently, the intermediate phenolic oxidation product,

phenoxyl radicals (PhO ∙), oxidises Mn^{2+} to Mn^{3+} (Takahama, 2004). Because PhO ∙ is

regenerated to phenolics after the reaction with Mn^{2+}, the continuous reactions of

H_2O_2-production/consumption by the class III PODs accumulate the oxidised phenolics and

oxidised Mn in the apoplast (Kenten and Mann, 1957).

Although brownish spots are an indicator of Mn toxicity in terrestrial plants,

attenuation of plant growth by Mn toxicity is not always accompanied by the expression of

brown spots in leaves (Blamey *et al.,*1986, 2015; Horiguchi, 1987; Wissemeier and Horst, 1992; González *et al.*, 1998; Chen *et al.*, 2013; Tsunemitsu *et al.*, 2018). Under excess Mn accumulation in the symplast, leaves show lower photosynthetic activity and lower Chl content (Nable *et al.*, 1988; Macfie and Taylor, 1992; González and Lynch, 1997; Kitao *et al.*, 1997; Fernando and Lynch, 2015). Houtz *et al.* (1988) demonstrated a decrease in photosynthetic activity despite the absence of brown necrotic spots on leaves grown under high Mn concentrations, implying that a decrease in photosynthetic activity occurred independently of apoplastic Mn toxicity. These characteristics are termed symplastic Mn toxicity (Fernando *et al.*, 2016). The decreases in Chl concentration and photosynthetic activities have been hypothesised to be caused by photoinhibition by ROS; the decrease in photosystem I (PSI) content or the disturbance of Chl synthesis due to the inhibition of Fe absorption; Mn binding to Rubisco instead of Mg; or stomatal dysfunction (Clairmont *et al.*, 1986; Suresh *et al.*, 1987; Houtz *et al.*, 1988; McDaniel and Toman, 1994; Millaleo *et al.*, 2013). However, these hypotheses are often independently discussed at both *in vivo* and *in vitro* levels; therefore, the mechanisms by which symplastic Mn toxicity suppresses photosynthesis are less evident. Elucidating Mn toxicity including both its apoplastic and symplastic mechanisms would contribute to establishing a strategy for protecting terrestrial plants against Mn toxicity in agricultural land (Fernando and Lynch, 2015). Currently, the strategy for preventing Mn toxicity depends on two mechanisms. One is limiting Mn absorption and transport, and the other is Mn sequestration from the cytosol to the vacuole (Li *et al.*, 2019). These strategies are important in preventing Mn toxicity. However, identifying the expression mechanisms of symplastic Mn toxicity would be informative for manipulating Mn toxicity in terrestrial plants without limiting the Mn transportation system. This knowledge would be applicable to plant engineering processes such as Mn phytoremediation (Lei *et al.*, 2007).

In this study, we investigated the mechanisms of symplastic Mn toxicity to suppress photosynthetic activity in rice leaves. Rice shows high tolerance toward Mn toxicity, which make it easy to investigate the effect of excess Mn concentrations without severe necrosis

and chlorosis (Führs *et al.*, 2010). Additionally, complete genetic information is available for

investigating gene expression. Previous studies have indicated that photosynthetic electron

transport activities on the thylakoid membrane and Rubisco isolated from leaves grown

under high Mn concentrations were robust, although CO_2 assimilation was substantially

suppressed in leaves (Houtz *et al.*, 1988; Nable *et al.*, 1988; Chatterjee *et al.*, 1994). Based

on these observations, we hypothesised that chloroplast proteins are not the primary targets

of Mn toxicity, but factors that modulate CO_2 assimilation are targeted under Mn toxicity.

Here, we discuss how photosynthetic activity is limited under Mn toxicity conditions from a

wide range of physiological responses in plant cells.

MATERIALS AND METHODS

Plant materials and plant growth conditions

Oryza sativa L. 'Nipponbare' wild-type (WT) plants were grown in hydroponic culture, as reported in a previous study (Makino *et al.,* 1994). We employed two Mn concentration conditions by changing the $MnSO_4$ application to 0.5 μM (control conditions) and 200 μM (Mn-toxic conditions) (Tsunemitsu *et al.,* 2018). The pH of the hydroponic culture was adjusted to 5.2 using HCl, and the solution was replenished twice a week. The chamber was maintained at 60% relative humidity with a 14 h light (28 ^{o}C) and 10 h dark (25 ^{o}C) photoperiod. The light intensity was 500–600 μmol photons m^{-2} s^{-1}. All physiological, structural, and genetic analyses were performed for fully and newly expanded leaves after 70 d of germination, and the analyses were completed before heading.

Leaf photosynthesis and respiration measurements in rice plants

The gas exchange analysis, Chl fluorescence, and oxidised reaction centre Chl in PSI (P700^{+}) were simultaneously measured using a combined system of Li-6400 (Li-COR Inc., Lincoln, USA), Mini-PAM, and PAM-101 equipped with a dual-wavelength emitter-detector unit (ED-P700DW) (Heinz Walz GmbH, Effeltrich, Germany) with a cold halogen lamp. Ambient air (40 Pa CO_2 and 21 kPa O_2) and pure CO_2 gas were mixed to maintain the CO_2 concentration during measurements. The gases were saturated with water vapour at 18.0 ± 0.1 ^{o}C, and the leaf temperature was maintained at 28 ^{o}C. The Chl fluorescence parameters Y(II), Y(NPQ), and Y(NO) were calculated as described by Baker (2008), and a measuring light (0.1 μmol photons m^{-2} s^{-1}) and a saturated pulse (10,000 μmol photons m^{-2} s^{-1}, 600 ms) were employed to calculate the photosynthetic parameters of PSII. The oxidation-reduction state of P700 in PSI was determined according to the method described by Klughammer and Schreiber (1994). The maximum oxidation level of P700 (Pm) was obtained using a saturated pulse under far-red light illumination, and PSI photosynthetic parameters [Y(I), Y(ND), and Y(NA)] were calculated using the oxidation-reduction state of P700 as determined by a saturated pulse method.

The CO_2 emission rate was measured using a Li-6400 in the gaseous phase. The

O_2 absorption rate was measured using an O_2-electrode (CB1D; Hansatech Instruments

Ltd., King's Lynn, UK) in the aqueous phase at 25 ^oC (Hachiya *et al.,* 2010). For efficient

absorption of the chemical reagent into leaves in the aqueous phase, the leaf surface was

washed with 10% (v/v) dimethyl sulfoxide, and the leaves were finely cut up. The O_2

absorption   rate   was   measured   in   the   reaction   buffer   [50   mM

N-(2-hydroxyethyl)piperazine-N-2-ethanesulfonic acid (HEPES)-KOH (pH 6.6), 10 mM

2-(N-morpholino)ethanesulfonic acid, 0.2 mM $CaCl_2$, 50 mM sucrose]. To evaluate the

uncoupled respiration rate and alternative oxidase (AOX) and cytochrome *c* oxidase (COX)

activities in the mitochondria, 10 μM carbonyl cyanide m-chlorophenyl hydrazine, 2 mM

n-propyl gallate, and 2 mM KCN were added sequentially.

Quantification of leaf mineral elements and chlorophyll content

Leaf mineral elements were analysed according to Takagi *et al.* (2020). Briefly, the leaf

blades were dried at 70 ^oC and ground using a homogeniser. To determine the mineral

content, except for that of N, the dry matter was digested overnight in an acid mixture

(HNO_3:H_2SO_4:$HClO_4$ = 5:1:2). Subsequently, the solution was heated at 150 ^oC for 30 min in

a heat block. After cooling, the solutions were heated again at 200 ^oC for 1 h. After

decomposition, the mineral contents were quantified by inductively coupled plasma optical

emission spectroscopy (ICP-OES) (iCAP™ 7200; Thermo Fisher Scientific Inc., Waltham,

USA). To measure the total N concentration in the leaves, dried leaves were decomposed

using 60% (v/v) H_2SO_4 with 30% (v/v) H_2O_2. The N concentration was determined using

Nessler's reagent in a decomposed solution after the addition of 10% (w/v) potassium

sodium tartrate and 2.5 M NaOH, and the absorbance was measured at 420 nm (Takagi *et*

al., 2017). The leaf Chl content was quantified using fresh samples with *N,*

N-dimethylformamide, as described previously (Porra *et al.,* 1989).

Carbohydrates quantification in leaves

Leaves were sampled at the end of the day (1 h before the start of the dark period) and at

the end of the night (1 h before the start of the light period); subsequently, leaves were

frozen using liquid nitrogen. Glucose, sucrose, and starch were extracted from leaves

according to Morita *et al.* (2015) and quantified using the Enzytech^{TM} D-Glucose/Sucrose kit

(R-Biopharm AG, Darmstadt, Germany).

Quantification of leaf amino acid contents

The leaves were homogenised using 10 mM HCl with liquid nitrogen. After centrifugation

(15,000 × g, 4 ^{o}C, 5 min), the supernatant was applied to a centrifugation filter (Amicon Ultra

0.5 mL 3 K device; Merck KGaA, Darmstadt, Germany), and the fraction containing free

amino acids was obtained by centrifugation (15,000 × g, 4 ^{o}C, 10 min). The amino acid was

labelled using AccQ-Tag Ultra Derivatization Kit (Waters Co., Milford, USA) according to the

manufacture's instructions, and the free amino acids were separated using an HPLC system.

The HPLC system consisted of a 305 Piston Pump as the system controller and a 306 Pump

as a multi-pump application equipped with an 811D Dynamic Mixer and 805 Manometric

Module, using the Trilution LC software (Gilson, Villiers le Bel, France). The sample solution

was manually injected into the HPLC system. The derivative amino acids were separated on

a Sepax Bio-C18 (φ4.6 × 250 mm) (Sepax Technologies, Inc., Newark, USA) and eluted by

100% solvent A [5% (v/v) methanol in 20 mM CH_3COONa (pH 6.5)] for 4 min after injection

and by a linear gradient of 94% solvent A to 30% solvent B (100% acetonitrile) for another

65 min. The column was washed with 100 % solvent B for 20 min and equilibrated with

100% solvent A for 16 min. The flow rate was 0.6 mL min^{-1}, and the column temperature was

maintained at 25 ^{o}C by incubation in a hot pocket (Thermo Fisher Scientific Inc., Waltham,

USA). The peaks derived from derivatised amino acids were detected using UV/VIS-155

(Gilson, Villiers le Bel, France).

TCA cycle enzyme activities

The TCA cycle enzyme activities in the leaf blades were determined according to Noguchi

and Terashima (2006). Briefly, the leaves were homogenised with the extraction buffer [100

mM HEPES-KOH (pH 7.5), 10 mM KH_2PO_4, 0.5 mM EDTA-Na, 10 mM dithiothreitol, 0.05%

(v/v) TritonX-100, and 20% (v/v) glycerol] with polyvinylpyrrolidone. The reaction medium

was prepared for each enzyme according to the previous study, and the enzyme activity was

determined at 30 ^{o}C using UV-160A equipped with a temperature control system (Shimadzu,

Kyoto, Japan).

Structural analysis of stomata complex and leaf cross-section

For scanning electron microscopic analysis, the leaves were fixed with half-strength

Karnovsky's solution [50 mM phosphate buffer (pH 7.2), 2.5% (v/v) glutaraldehyde, 2% (v/v)

paraformaldehyde] with vacuum infiltration for 3 h. After fixation, the leaves were washed

with the phosphate buffer and dehydrated using a dilution series of ethanol, followed by

t-butanol. Leaves were dried using a t-butanol freeze drier (Vacuum Device Inc., Ibaraki,

Japan) and coated with Pt (JEC-300FC; JEOL Ltd., Tokyo, Japan). The stomatal complex

structure was monitored using a JSM-IT200 scanning electron microscope (JEOL Ltd.,

Tokyo, Japan). The size of the stomatal complex, comprising the paired guard cells, the

paired subsidiary cells, and the pore itself, and the stomatal density were analysed using

SEM Operation software (JEOL Ltd., Tokyo, Japan). Under each growth condition, the size

of the stomata complex and the stomatal density were analysed on an independent section

of the leaves as indicated in figure legends of six biologically independent plants. To analyse

the vertical and horizontal leaf sections, leaves were sectioned and fixed with

formalin-acetate-alcohol (FAA) [63% (v/v) ethanol, 5% (v/v) acetic acid, 5% (v/v) formalin]

with vacuum infiltration for 1 h, followed by incubation at 4 ^{o}C for 2 d. The leaves were

dehydrated using a dilution series of ethanol, followed by n-butanol. The dehydrated leaves

were embedded in paraffin and were sectioned at 10 μm thickness using an RX-860

microtome (Yamato Kohki Industrial Co., Ltd., Saitama. Japan). The FAA-fixed leaf sections

were stained with 0.05% (w/v) toluidine blue O. To investigate the intact leaf sections, leaves

were embedded in 50 mM phosphate buffer (pH 7.2) containing 5% (w/v) agar. The

agar-embedded leaves were sectioned at a thickness of 30 μm using DTL-1000 vibratome

(Dosaka EM Co., Ltd., Kyoto, Japan). Intact leaf sections and FAA-fixed leaf sections were

analysed using an optical microscope (BX53; Olympus Co., Japan).

Indole acetic acid quantification

The IAA content in leaves was quantified using a GC-MS system, according to Nghi *et al.*

(2021). Briefly, the leaves were homogenised in 70% (v/v) cold acetone. After centrifugation,

the supernatant was reduced to the aqueous phase by vacuum centrifugation equipped with

a cold trap. Subsequently, HCl (0.1 mM) was added to adjust the pH to 2.8, and diethyl ether

was added to dissolve IAA into the diethyl ether fraction. The diethyl ether fraction was dried

under vacuum and subsequently dissolved in *N*,*O*-bis(trimethylsilyl) tri-fluoroacetamide

(BSFTA) containing 1% trimethylchlorosilane. IAA was silylated at 70 ^{o}C for 1 h. The

silylated IAA was identified using a GC-MS-QP2010 SE (Shimadzu, Kyoto, Japan) with a

DB-WAX column. The conditions for GC-MS analysis were the same as those described

previously (Fujii *et al.*, 2010). The silylated IAA was identified by fragment ion 202 (*m/z*) and

molecular ion 319 (*m/z*) (Ljung *et al.*, 2005; Gutierrez *et al.*, 2009; Nghi *et al.*, 2021).

Gene expression analysis

To investigate gene expression in rice leaves, mRNA was isolated from the leaf blades

according to Suzuki *et al.* (2004). After mRNA isolation, cDNA was synthesised using

PrimeScript™ RT Reagent Kit (Takara Bio Inc., Shiga, Japan), and the mRNA content was

quantified by real-time PCR using the LightCycler® 96 System (Roche Diagnostics K.K.,   Table S1

Tokyo, Japan) with a KAPA SYBR FAST One-Step qRT-PCR Kit (Nippon Genetics Co., Ltd,

Tokyo, Japan). The primer sequences used for real-time PCR analyses are listed in

Supplementary Table S1.

Simulation of CO$_2$ fixation rate according to biochemical photosynthetic model

Rubisco limited CO$_2$ fixation rate was calculated from the equation of von Caemmerer and

Farquhar (1981):

$A = EV_c(Cc - \Gamma^*) / [Cc + K_c(1 + O / K_o)] - Rd$, (1)

Where E is the amount of Rubisco protein (assumed to be 5.3 µmol m^{-2}, from rice grown

with hydroponic solution same as in the present study described by Suganami et al. 2021),

V_c is the Rubisco activity of carboxylation, Cc is the chloroplastic CO_2 partial pressure, O is

the partial pressure of O_2 in the chloroplast (assumed to be the same as in the atmosphere,

21 kPa), K_c and K_o are the Michaelis-Menten constants for CO_2 and O_2, and Rd is the day

respiration (assumed to be 1.0 µmol m^{-2} s^{-1}). Γ^* is the CO_2 compensation point of

photosynthesis in the absence of Rd as defined as follows:

$\Gamma^* = 0.5V_oK_cO / V_cK_o$, (2)

where V_o is the Rubisco activity of oxygenation. Mg^{2+}-binding Rubisco kinetics were taken

from Makino et al., (1988). Mn^{2+}-binding Rubisco kinetics were calculated by multiplying the

average ratio of Mn^{2+}/Mg^{2+}-binding Rubisco kinetics and Mg^{2+}-binding Rubisco kinetics. The

average ratio of Mg^{2+}/Mn^{2+}-binding Rubisco kinetics were calculated by the data from Bloom

and Kameritsch (2017). Rubisco kinetics were summarized in Supplementary Table S2.   **Table S2**

Photorespiration rate were was calculated from the equation of von Caemmerer and

Farquhar (1981):

$F = 0.5EV_oO / (O + K_o(1 + Cc / K_c)$, (3)

Statistical analysis

All measurement data are expressed as means ± standard deviation (SD), or box plots with

bars indicating the 1.5 × interquartile range (IQR) and with squares indicating the mean

value of at least three independent biological analyses. Significant differences in

physiological and biological parameters between the control- and Mn-toxic conditions were

detected by the non-parametric Kruskal-Wallis test. All statistical analyses were performed

using Origin Pro 2020 (LightStone Corp., Tokyo, Japan).

RESULTS

Phenotypes of rice grown under Mn-toxic conditions

To study the effects of Mn toxicity, *Oryza sativa* L. 'Nipponbare' was grown under high Mn concentrations (200 µM; Mn-toxic conditions) in hydroponic culture for 70 d after germination. Under Mn-toxic conditions, the leaves showed pale green and brown-coloured sections, especially at the tips (Fig. 1A, B). These observations correspond with typical Mn toxicity symptoms in rice leaves (Vlamis and Williams, 1964; Chen *et al.*, 2013). Compared with the control conditions, Mn-toxic conditions decreased the total dry weight of rice, especially the dry matter of the leaf sheath and root (Fig. 1C, D).

Fig. 1

To study the effects of high Mn application on leaf mineral composition, the leaf mineral concentrations were quantified. The leaf Mn concentration was significantly increased under Mn-toxic conditions (Fig. 2). This concentration was comparable to that of WT rice, which showed growth inhibition when grown at a concentration of 1,000 µM Mn for 3 weeks (Sasaki *et al.*, 2011). The K, Ca, and Fe concentrations were similar under the control and Mn-toxic conditions (Fig. 2). In contrast, the Mg, Zn, and Cu concentrations increased under Mn-toxic conditions (Fig. 2). These results demonstrated decreased growth and the Mn toxicity symptoms accompanied by Mn accumulation in leaves grown under Mn-toxic conditions.

Fig. 2

Mn-toxic conditions suppressed CO_2 assimilation by limiting stomatal conductance in leaves

Next, the steady-state photosynthetic activities were measured under ambient air conditions (40 Pa CO_2 and 21 kPa O_2). The CO_2 assimilation rate was significantly decreased under the high light irradiance by the Mn-toxic treatments (Fig. 3A). The Mn-toxic treatments showed lower stomatal conductance (g_s) than did the control plants, and the change in g_s with increasing light intensity was also suppressed (Fig. 3B). These stomatal responses to the increase in Mn concentration were consistent with previous studies (Suresh *et al.*, 1987; Nable *et al.*, 1988; González and Lynch, 1997; Lidon *et al.*, 2004; Santos *et al.*, 2017). The

Fig. 3

internal CO_2 concentration in leaves (Ci) was lower under the Mn-toxic conditions than

under control conditions (Fig. 3C). These results indicated that, under Mn-toxic conditions,

the stomata limit CO_2 diffusion into leaves and CO_2 assimilation is suppressed, particularly

under high light conditions.

In addition to the CO_2 assimilation rate, the photosynthetic electron transport

activities in both PSII and PSI were analysed by chlorophyll fluorescence and

$P700^+$-dependent absorbance changes in the leaves. The quantum yield of PSII [Y(II)] was

lower under Mn-toxic conditions than under control conditions at the high light irradiance

(Fig. 3D). This result indicated that the electron transport activity in PSII was suppressed at

the high light irradiance, but compared with the CO_2 assimilation rate, the change in Y(II)

was marginal. The quantum yield of non-photochemical quenching [Y(NPQ)] increased

similarly with the increase in light intensity under both control and Mn-toxic conditions (Fig.

3E). The quantum yield of non-radiative energy loss [Y(NO)] was different between the

control and Mn-toxic conditions under low light irradiance; however, this difference was

masked under high light irradiance (Fig. 3F). These results showed that the photoprotective

mechanisms in PSII are robust, but the redox state in PSII can be perturbed under Mn

toxicity. The quantum yield of PSI [Y(I)] showed similar kinetics to that of Y(II), and no

significant differences were observed between the control and Mn toxic condition (Fig. 3G).

The quantum yield of non-photochemical quenching at the donor side of PSI [Y(ND)] was

higher under Mn-toxic conditions than under the control conditions at high light irradiance

(Fig. 3H). In contrast, the quantum yield of non-photochemical quenching at the acceptor

side of PSI [Y(NA)] was lower under Mn-toxic conditions than under the control conditions at

high light irradiance (Fig. 3I). These results indicated that the whole-chain photosynthetic

electron transport rate was less affected under Mn-toxic conditions than under the control

conditions; however, the photosynthetic electron transport reaction was limited to the donor

side of PSI, and PSI was more oxidised under Mn-toxic conditions than under the control

conditions.

The maximum quantum yield of PSII (Fv/Fm) was significantly, but marginally

decreased under Mn-toxic conditions (Fig. 2J). The total Chl content in leaves was also

decreased under Mn-toxic conditions (Fig. 2K). These results indicated that Mn toxicity

stimulates PSII photoinhibition. In contrast, the total N concentration was not significantly

different between the control and Mn-toxic conditions (Fig. 2L). Based on the linear function

between the N and Rubisco concentrations in the leaves (Nakano *et al.,* 1997), this result

indicated that the quantity of Rubisco was less affected by Mn toxicity.

Carbohydrate metabolism in leaves grown under Mn-toxic conditions

To examine whether the suppression of photosynthesis by excessive Mn accumulation

affects carbon acquisition during growth, the major carbohydrate concentrations were

quantified in the leaves. Leaves were sampled at both the end of the day and the end of the

night to visualise the CO_2 assimilation activity during the day (Takagi *et al.,* 2016*a*). At the

end of the day, Mn-toxicity treatments showed lower sucrose concentrations in the leaves   **Fig. 4**

than did the control plants (Fig. 4). In contrast, at the end of the night, the sucrose

concentration was similar under both control and Mn-toxic conditions (Fig. 4). The glucose

and starch concentrations were lower than the sucrose concentration under both control and

Mn-toxic conditions, and significant differences were not detected between the growth

conditions at the end of the day and the end of the night (Fig. 4). The sum of the sucrose,

glucose, and starch concentrations was lower in leaves grown under Mn-toxic conditions at

the end of the day, but the sum of these carbohydrate concentrations was similar between

the growth conditions at the end of the night (Fig. 4). These results indicated that carbon

acquisition is suppressed, corresponding to lower CO_2 assimilation activities, in leaves

grown under Mn-toxic conditions.

Effects of Mn toxicity on mitochondrial respiration in the leaves

The carbohydrates fixed during the day are metabolised at night and consumed at a

constant rate from the beginning of the night to the dawn (Moraes *et al.*, 2019). Therefore,

the difference in carbohydrate concentrations between the end of the day and end of the

night indicates the carbon catabolic activity during the night (Takagi *et al.*, 2016*a*). To

confirm whether carbon catabolism is suppressed under Mn-toxic conditions (Fig. 4), leaf

respiration was examined under control and Mn-toxic conditions. The respiratory CO_2

emission rate was decreased under Mn-toxic conditions (Fig. 5A). To examine the effect of

the change in respiration activities on carbon catabolism, the leaf amino acid content was

quantified during the night (Sulpice *et al.*, 2014). Subsequently, the Asp, Glu, and Gly   **Fig. 5**

contents were significantly decreased, and the Ser content was increased under Mn-toxic

conditions (Fig. 5B). These results indicated that carbon catabolism is affected by Mn-toxic

conditions during the night.

Next, we investigated the effects of Mn toxicity on the mitochondrial enzymes

involved in respiration. First, the respiratory electron transport activities were measured in

the presence of sucrose and mitochondrial electron transport inhibitors *in vivo* using an

aqueous phase O_2-electrode (Hachiya *et al.*, 2010). In the absence of inhibitors, the O_2

consumption rate did not differ between the control and Mn-toxic conditions (Fig. 5C). CCCP

dissipated mitochondrial membrane potential ($\Delta\psi_m$) and similarly stimulated O_2

consumption rate in both leaves grown under the control and Mn-toxic conditions (Fig. 5C).

The COX and AOX activities in the presence of CCCP and sucrose were determined by

adding n-propyl gallate and KCN. Subsequently, the control and Mn-toxic conditions showed

similar O_2 consumption rates depending on the COX and AOX activities (Fig. 5C). The effect

of Mn toxicity on mitochondrial respiratory electron transport activity was reproduced when

these activities were determined on a leaf fresh-weight basis (Supplementary Fig. S1).   **Fig. S1**

These results indicated that under sucrose feeding, the mitochondrial respiratory electron

transport activities were similar between the control and Mn-toxic conditions.

To understand the effects of Mn toxicity on the enzyme activities of the TCA-cycle,

the six types of TCA cycle enzyme activities were determined on both leaf area and

fresh-weight basis. NADP-isocitrate dehydrogenase, NAD-isocitrate dehydrogenase,   **Fig. S2**

NAD-malate dehydrogenase, and aconitase showed similar activities under the control and

Mn-toxic conditions (Fig. 5D-G, I, Supplementary Fig. S2). The NAD-malic enzyme activity

decreased under Mn-toxic conditions (Fig. 5H, Supplementary Fig. S2), whereas the

fumarase activity increased (Fig. 5I, Supplementary Fig. S2). These results indicated that

Mn toxicity did not completely suppress the enzyme activities of the TCA cycle, but

specifically stimulated fumarase and suppressed NAD-malic enzyme activities.

Difference in stomatal development in the leaves between the control and Mn-toxic

conditions

To examine whether the decrease in g_s under Mn-toxic conditions was related to the change

in stomatal structure and development, the stomatal complex on the adaxial side of a leaf

was monitored. The stomatal complex, consisting of guard and subsidiary cells, was located

linearly to the vein under the control conditions (Fig. 6A). However, under Mn-toxic   **Fig. 6**

conditions, the stomatal complex showed a more scattered distribution (Fig. 6B). The

stomatal density increased under Mn-toxic conditions (Fig. 6C). In contrast, the size of the

stomatal complex evaluated by the maximum major and minor axis was smaller under

Mn-toxic conditions than under control conditions (Fig. 6A, B, D, E). The changes in

stomatal size were consistent with a previous study evaluating rice stomata structure under

excess Mn application (Lidon, 2002). These results indicated that Mn-toxic conditions

change stomatal development in the leaves.

Anatomical changes within a leaf blade grown under Mn-toxic conditions

Following the change in stomatal development on the leaf epidermis, the leaf anatomy was

also examined. Figure 7A shows the horizontal and vertical cross-sections of the leaves

against the vein. Apparent internal air spaces were observed within leaves grown under   **Fig. 7**

control conditions. In contrast, cells were highly condensed in leaves grown under Mn-toxic

conditions, and the internal air space was hardly observed (Fig. 7A). These results showed

that in addition to stomatal complex, leaf anatomical development was also altered under

Mn-toxic conditions.

The vertical cross-section revealed that the bundle-sheath cells were brownish

under Mn-toxic conditions (Fig. 7A). To investigate whether the brown spots were artifacts

caused by the chemical fixing treatment, intact leaf sections were also studied.

Subsequently, we found that the brownish spots natively existed in the bundle-sheath cells,

both small and large bundles gown under Mn-toxic conditions (Fig. 7B, C). These results

indicated that under Mn-toxic conditions, rice leaves caused apoplastic Mn toxicity-like

symptoms in addition to symplastic Mn toxicity, and the symptoms are more evident in the

bundle-sheath cells in rice leaves.

Auxin content in the leaves and mRNA expression of auxin-responsive genes

Our findings demonstrated that excess Mn accumulation affects the developmental process

of leaves. IAA is an important hormone that determines cell fate and differentiation in plants

(Perrot-Rechenmann, 2010). Morgan et al. (1966) reported that Mn toxicity stimulates IAA

oxidation activity in cotton leaves, indicating that the auxin concentration could be lowered

by accelerated auxin degradation in leaves under excess Mn conditions. To examine this

possibility, the IAA concentration in leaves was quantified by GC-MS. Figure 8A shows the

Fig. 8

520   total ion chromatograms (TIC) with fragment ion chromatograms of the 202 (*m/z*) and 319

521   (*m/z*) of the IAA standard and leaf extract from the control and Mn-toxic conditions (Fig 8A)

522   (see Materials and Methods). Subsequently, this analysis revealed that the IAA

523   concentration was significantly decreased by 76% in the leaves under Mn-toxic conditions

524   (Fig. 8B).

525   To confirm that the decrease in IAA concentration affects physiological reactions

within the leaves, the IAA-responsive gene expression was investigated. *OsARF1*,

OsARF15, *OsARF16*, *OsAUX/IAA1,* and *OsAUX/IAA24* showed increases in their

expression level with an increase in IAA concentration in the leaves (Waller *et al.,* 2002; Jain

et al., 2006; Wang *et al.,* 2007). Among these genes, the expression levels of *OsARF1*,

OsARF16, and *OsAUX/IAA1* were significantly decreased under Mn-toxic conditions (Fig.

8C). *OsARF15* showed similar trends to other *ARF* genes, but the difference was not

significant between the growth conditions. In contrast, *OsAUX/IAA24,* and *OsACT1-1*

showed comparable expression levels between the control and Mn-toxic conditions (Fig.

8C). From these results, the auxin-responsive gene expression indicated a lower IAA

concentration in leaves under Mn-toxic conditions in concert with the results of GC-MS

although all genes were not responded to equally.

537   To regulate auxin concentration in plant cells, DIOXYGENASE FOR AUXIN

OXIDATION (DAO) undertakes IAA catabolism to maintain IAA homeostasis by oxidising

IAA to an inactive form, oxindole-3-acetic acid, in plant cells (Zhao *et al.,* 2013). Because

Mn-toxicity stimulates oxidative IAA degradation *in vitro* (Morgan *et al.,* 1966), we

hypothesised that *OsDAO* is upregulated following an increase in Mn concentration;

however, the transcriptional activation of *OsDAO* was suppressed under Mn-toxic conditions

(Fig. 8). The expression of *OsDAO* is activated under high IAA concentrations to maintain

IAA concentrations in plant tissues (Zhao *et al.,* 2013). Based on this homeostatic response

of *OsDAO*, the decrease in *OsDAO* expression under Mn-toxic conditions would be a

consequence of the lower IAA concentration in leaves to avoid further inactivation of IAA.

DISCUSSION

In acidic or waterlogged agricultural fields, Mn is easily released from the soil and causes

Mn toxicity in crop plants (Fernando *et al.*, 2016). In addition to the current agricultural

situations, the effects of climate change, such as the rises in the atmospheric temperature

and the frequency of flooding or acidic rainfall, can further accelerate Mn dissolution in soils

(Fernando and Lynch, 2015). Therefore, the mechanisms of Mn toxicity should be

elucidated to manipulate the future threat of increasing Mn availability on agricultural lands.

To address the mechanisms of symplastic Mn toxicity, we investigated the physiological

effects of high Mn application in rice. In this study, we propose that the disturbance of IAA

homeostasis is one of the critical causes of symplastic Mn-toxicity in rice leaves.

Mn toxicity disturbs IAA homeostasis and attenuates auxin signalling in the leaves

Our results showed that the IAA concentration was decreased in the leaves under Mn-toxic

conditions (Fig. 8). A previous study demonstrated that IAA oxidation activity increased in

leaf extracts grown under high Mn concentration conditions (Morgan *et al.,* 1966). Our

observations suggest that the increase in IAA oxidation activity functions *in vivo.* In addition

to the decrease in *ARF* and *AUX/IAA* gene expression levels in rice, the microRNA

expression profiles revealed that miRNA160 which targets *ARF* genes was downregulated,

and miRNA164 which targets the auxin-signal related F-box protein (TRANSPORT

INHIBITOR RESPONSE 1, TIR1) was upregulated in the leaves of common beans under

Mn toxicity (Valdés-López *et al.,* 2010). From these observations, we suggest that Mn

toxicity affects the auxin signalling cascade at the transcriptional level by decreasing the IAA

concentration in the leaves.

We suggest that the decrease in IAA concentration, together with the change in

auxin signalling activities, causes a change in leaf anatomy and stomatal

patterning/development under Mn-toxic conditions. When IAA deficiency is caused by the

overexpression of GRETCHEN HAGEN3 which conjugates IAA to amino acids or the anion

peroxidase which oxidatively inactivates IAA *in planta*, the leaf cell size becomes smaller

and the leaf anatomy shows more condensed cell structures (Lagrimini *et al.*, 1997*a, b*; Du *et*

al., 2012). Alternatively, the transgenic plants that inhibit auxin signalling or auxin polar

transport activities represent the IAA-deficient phenotype and show the abnormal leaf

anatomical development (Pérez-Pérez *et al.*, 2010; Guo *et al.*, 2013; Qi *et al.*, 2014;

Muñoz-Nores *et al.*, 2017). The aforementioned relationships among the IAA-deficient

phenotypes and the change in leaf structure caused by attenuating auxin signal transduction

are similar to those among the leaf phenotypes under Mn-toxic conditions (Figs. 3B, 6).

Moreover, stomatal development is regulated by auxin signalling (Balcerowicz and Hoecker,

2014). Exogenous IAA application decreased stomatal density in a concentration-depending

manner; whereas the attenuation of ARF-dependent auxin signal transduction increased it

(Le *et al.,* 2014; Zhang *et al.,* 2014). Based on these results, a decrease in IAA

concentrations could affect the change in leaf development and stomatal

patterning/development.

Generally, the terrestrial plants with smaller stomata and higher stomata density

show higher transpiration ability than those with larger stomata and lower density (Franks

and Beerling, 2009). This relationship does not apply to g_s and stomatal patterning (Figs. 3B,

6C); therefore, additional factors to suppress stomatal opening should exist under Mn

toxicity. IAA stimulates stomatal opening by activating plasma membrane H^+-ATPase (Jezek

and Blatt, 2017), and the exogenous application of IAA stimulates stomatal opening in the

leaves (Klein *et al.*, 2003). Therefore, the decrease in g_s could be caused by reducing the

IAA concentration in the leaves. However, the regulation of stomatal function is quite

complex, and the ROS and organic acid composition are also determinants of stomatal

movement (Jezek and Blatt, 2017). Considering the stimulation of oxidative stress under Mn

toxicity (see the following discussion), the decrease in IAA would not be the only cause of

stomatal dysfunction. Interestingly, fumarase activity increased under Mn toxicity (Fig. 5I).

Because the increase in malate/fumarate ratio is important for stomatal opening (Araújo *et*

al., 2011), the increased fumarase activity might be a counteraction toward stomatal

dysfunction under Mn toxicity.

617    Currently, we cannot determine how the IAA concentration is lowered under Mn

618 toxicity at the molecular level. Further research is required on a molecular genetic basis to

619 elucidate the key factors that cause IAA deficiency. We now consider the possibility that IAA

620 oxidation can be undertaken by unknown proteins such as class III PODs as reported

621 previously (Ray, 1958; Lagrimini *et al.*, 1997*a, b;* Tognetti *et al.*, 2012). Because IAA is

622 transported through the apoplast, the increase in POD activity in the apoplast might

623 influence IAA catabolism under Mn toxicity (Robert and Friml, 2009).

625 **Suppression of photosynthesis is caused by stomatal dysfunction in rice**

626 We suggest that the simultaneous prolonged limitation of the CO_2 assimilation reaction and

627 stimulation of photorespiration caused oxidative stress by ROS under Mn toxicity. Under Mn

628 toxicity, ROS-scavenging enzyme activities are upregulated in various terrestrial plants

629 (Gonzáletz *et al.*, 1988; Lei *et al.*, 2007; Santos *et al.*, 2017). This response indicates that

630 oxidative stress is stimulated by ROS in cells under Mn toxicity. Furthermore, Houtz *et al.*

631 (1988) showed an increase in the RuBP concentration in leaves under Mn toxicity. This

632 result indicated that the turnover of photosynthates is limited by Rubisco activity (Suzuki *et*

633 *al.*, 2012). Here, g_s was lowered and the CO_2 assimilation rate was limited under Mn-toxic

634 conditions (Fig. 2B). In contrast to the CO_2 assimilation rate, which showed a 39% decrease

635 at the highest light irradiance, Y(II) only decreased by 9% at the same light intensity (Fig. 3).

636 This difference would have been caused by the difference in the electron distribution

637 between carboxylation and oxygenase reaction by Rubisco (Takagi *et al.*, 2016*b*; Wada *et*

638 *al.*, 2019). Even when the CO_2 availability within chloroplasts is limited by stomatal closure,

639 O_2 can diffuse within the leaves; therefore, Rubisco drives photorespiration (Zivcak *et al.*,

640 2013). Under such conditions, and the *proton motive force* is built up to stimulate luminal

641 acidification of the thylakoid membranes owing to the lower ATP consumption rate in the

642 photorespiration reaction compared with that in the carboxylation reactions in the

643 Calvin-Benson cycle (Takagi *et al.*, 2016*b*, 2017). Luminal acidification is represented by an

644 increase in Y(ND) (Fig. 3), which is increased by limiting the electron flow from

plastoquinone to cytochrome b_6f depending on the luminal acidification (Kramer *et al.*, 1999),

under Mn-toxic conditions. These results indicated that Mn toxicity stimulates

photorespiration via stomatal closure. The increase in photorespiration activity can also be

supported by the change in the Gly/Ser ratio under Mn-toxic conditions (Fig. 5B). Wada *et al.*

(2019) reported that the limitation of the CO_2 assimilation rate by drought stress with a

decrease in g_s stimulates PSI photoinhibition rather than PSII photoinhibition by ROS. A

similar situation would occur under Mn toxicity, which is why the decrease in PSII activity is

modest, although PSI inhibition is accentuated under Mn-toxic conditions (Kitao *et al.*, 1997;

Lidon *et al.*, 2004; Millaleo *et al.*, 2013). Previous studies have shown that Mn toxicity

targets PSI by attenuating Fe absorption (Andressen *et al.*, 2018). However, the

concentration was maintained within a sufficient range for Fe nutrition in rice [from 70 to 300

µg g^{-1} (D.W.)] (Fageria and Stone, 2008). Furthermore, a decrease in the PSI content

caused an increase in Y(NA), instead of Y(ND) during steady-state photosynthesis (Brestic

et al., 2015). Hence, we suggest that the decrease in PSI content due to Mn toxicity would

be caused not only by the Fe status but also by PSI photoinhibition, as a consequence of

limited CO_2 assimilation. Other nutrients such as Ca, Mg and Zn were also reported to be

deficient under Mn toxicity (Broadley *et al.*, 2012). We did not observe a decrease in these

minerals in the leaves, but rather increased Mg, Cu, and Zn concentrations under Mn

toxicity (Fig. 2). Because Cu/Zn-type SOD activities are upregulated under Mn toxicity

(Gonzáletz *et al.*, 1988; Lei *et al.*, 2007; Santos *et al.*, 2017) and Mg supply rescues Mn

toxicity symptoms (Broadley *et al.*, 2012), the requirement of these minerals might be

upregulated.

Our results suggest that chloroplasts have homeostatic systems to evade

excessive Mn accumulation. *In vitro* studies have shown that excess free Mn stimulates

ROS production by PSII and causes severe oxidative damage to thylakoid lipids and PSII

(Panda *et al.,* 1987). Although the Mn concentration was high (Fig. 2), clear evidence of

severe damage to PSII was not detected here (Fig. 3J). Furthermore, Mn is to bind Rubisco,

and an *in vitro* study indicated that the substitution of Mg^{2+} with Mn^{2+} decreases the

specificity of Rubisco for CO_2 (Jordan and Ogren, 1983; Bloom and Kameritsch, 2017).

Previous studies have suggested that Mn binding to Rubisco causes a decrease in

photosynthesis under Mn toxicity (Houtz *et al.,* 1988; Kitao *et al.,* 1997). However, the

contribution of Mn-binding Rubisco to decrease CO_2 assimilation reactions remains

unknown under Mn-toxic conditions *in vivo* (Houtz *et al.,* 1988; Chatterjee *et al.,* 1994). To

address this question, we simulated the change in CO_2 fixation rate between Mg^{2+}- and

Mn^{2+}-binding Rubisco using photosynthetic biochemical model (von Caemmerer and

Farquhar, 1981). Calculated CO_2 assimilation rate using Mn^{2+}-binding Rubisco kinetics was

negative value below Cc = 50 Pa and greatly decreased compared to calculated CO_2

assimilation rate using Mg^{2+}-binding Rubisco kinetics (Supplementary Fig. S3). However,

such a drastic decrease in CO_2 assimilation rate was not observed (Fig. 3). Although

Mn^{2+}-binding Rubisco showed a higher photorespiration rate than Mg^{2+}-binding Rubisco,

this difference cannot compensate the difference in CO_2 assimilation rate. Moreover,

investigation of the relationship between g_s and the CO_2 assimilation rate did not show a

decrease in the CO_2 assimilation rate independent of the change in g_s (Supplementary Fig.

S4). These considerations reconciled the cause of the decrease in CO_2 assimilation as

decrease in g_s, and Rubisco is protected from excess Mn in chloroplasts. To protect

chloroplasts from excess free Mn accumulation, the homeostatic function of Mn^{2+} in the

chloroplasts should function. At this point, two mechanisms can be considered. The first

mechanism is binding Mn to the thylakoid membranes. Although this modulates membrane

structure of the thylakoid membranes, no inhibitory effect on photosynthetic activity was

observed (Lidon and Teixeira, 2000*a, b*). The binding of Mn to the thylakoid membrane

could be a storage system for Mn homeostasis. The second mechanism is the sequestration

or limiting Mn transport within chloroplasts. Indeed, Mn distributed to the chloroplasts was

decreased under Mn-toxic conditions in rice (Führs *et al.,* 2010). These homeostatic

systems would prevent the deleterious effects of Mn^{2+} within the chloroplasts.

Carbon catabolism due to reduced carbohydrates suppresses growth

A decrease in respiration rate under Mn toxicity has been reported in wheat and cotton

plants (Sirkar and Amin, 1974; Macfie and Taylor, 1992). We also observed a decrease in

leaf respiration, but not in sucrose feeding (Fig. 5). Moreover, no decreases in mitochondrial

enzyme activities, including TCA cycle and respiratory electron transport chain enzymes,

were observed, except for the NAD-malic enzyme (Fig. 5). The physiological importance of

the NAD-malic enzyme is highly limited in C3 plants, and a pronounced phenotype has not

been observed in NAD-malic enzyme mutants in *Arabidopsis* (Tronconi *et al.*, 2008). This

could be because of the lower contribution to the supply of pyruvate into the TCA cycle by

NAD-malic enzyme activity (Williams *et al.,* 2008). These observations suggested that the

significant decrease in NAD-malic enzyme activity could be independent of the growth

defect and decreased respiration rate under Mn-toxic conditions (Figs. 1, 5). However, the

change in amino acid composition could be influenced by a decrease in the NAD-malic

enzyme (Tronconi *et al.,* 2008). From these observations, the decreased respiration rate was

caused by a decrease in carbohydrate concentrations due to the suppression of

photosynthesis (Sulpice *et al.,* 2014). Carbohydrate catabolism during the night is a

determinant of plant growth (Takagi *et al.,* 2016*a*). We suggest that a decrease in the

capacity of carbon catabolism causes growth defects in rice under Mn toxicity. Interestingly,

Hibberd and Quick (2002) reported that similar to C4 plants, the NAD-malic enzyme activity

is high in the bundle-sheath cells in C3 plants. Here, we found that the bundle-sheath cells

turned brown under Mn-toxic conditions (Fig. 7B, C). Based on a previous study on

apoplastic Mn toxicity, brown spots contained oxidised phenol, oxidised Mn, and callose.

Fernando *et al.* (2016) showed that wheat leaves concentrate Mn in the bundle-sheath cell.

However, this specific localisation was not observed in soybean and canola, suggesting that

the bundle-sheath cell could function as storage organs for attenuating Mn toxicity in

monocots. The Mn accumulation in the bundle sheath cell could cause the suppression of

the NAD-malic enzyme activity in leaves. These results provide insight for elucidating the

function of bundle-sheath cells in C3 monocots.

CONCLUDING REMARKS

We proposed that the disturbance of IAA homeostasis is an initial cause of the inhibition of CO_2 assimilation and photoinhibition caused chlorosis in the leaves (Fig. 9). As a short-term effect, the decrease in IAA suppresses stomatal opening. Subsequently, the decrease in IAA concentration modifies leaf anatomy and stomatal development in newly emerged leaves as a long-term effect. Because leaf anatomy determines the CO_2 pathway for efficient CO_2 absorption by the chloroplasts, the lower internal air space could further lower the CO_2 fixation efficiency under Mn toxicity (Earles *et al.,* 2018). The decreased CO_2 assimilation rate and increased photorespiration rate decelerate carbon acquisition for catabolism; thus, vegetative growth is suppressed. To our knowledge, there are limited studies focusing on the relationship between auxins and Mn toxicity (Morgan *et al.,* 1966). Tsunemitsu *et al.* (2018) reported that the *osmtp11* mutant decreased grain fertility and yield in rice. *OsMTP11* undertakes Mn sequestration to the endoplasmic reticulum and is highly expressed in inflorescences, including the anthers, pistil, lemmas, paleas, and ovaries (Tsunemitsu *et al.,* 2018). Together with our conclusion, Mn toxicity may also affect grain fertilisation by disturbing IAA homeostasis. Not only the photosynthetic measurements but also the developmental analyses from leaf emergence to the reproductive phase would provide more insights into the mechanisms of the symplastic Mn-toxicity by focusing on IAA.

Fig. 9

SUPPLEMENTARY DATA

Figure S1; The sucrose-feeding respiration activities and mitochondrial respiratory electron transport activities depending on ATP synthase (uncoupled), COX, and AOX based on leaf fresh weight

Figure S2; The enzyme activities involved in the TCA cycle calculated on the fresh weight basis.

Figure S3; The simulated CO_2 assimilation in leaves containing Mg-binding Rubisco and Mn-binding Rubisco.

Figure S4; The relationship between g_s and CO_2 fixation rate under the control and the Mn-toxic conditions.

Table S1; The primer list for the gene expression analysis in rice leaf blade

Table S2; Mg^{2+}-binding Rubisco and Mn^{2+}-binding Rubisco kinetics.

ACKNOWLEDGEMENTS

The authors thank Editage (https://www.editage.jp) for English language editing of the manuscript. The authors also thank Prof. Yukio Ishikawa, Prof. Yutaka Okumoto, Prof. Takeo Yamakawa, and Associate Prof. Shuji Sano at Setsunan University for their kind technical support of this study. This work was supported by the Japan Society for the Promotion of Science (JSPS) research fellowship (JSPS KAKENHI Grant No. 18J00852 [DT] and KAKENHI Grant No.16H06379 [AM]).

Conflicts of interest:

The authors declare no conflicts of interest.

Author contributions:

Conceptualisation: DT; investigation: DT, KI, MS, TU, TF, YT, MK, and KO; original draft: DT; writing, review, and editing: DT, KI, MS, TU, TF, YT, MK, KO and AM; funding acquisition: DT and AM.

Data availability statements

All data supporting the findings of this study are available within the article and its supplementary figures and tables published online.

REFERENCES

Andresen E, Peiter E, Küpper H. 2018. Trace metal metabolism in plants. Journal of Experimental Botany **69,** 909-954.

Araújo WL, Nunes-Nesi A, Osorio S, *et al.* 2011. Antisense inhibition of the iron-sulphur subunit of succinate dehydrogenase enhances photosynthesis and growth in tomato via an organic acid–mediated effect on stomatal aperture. The Plant Cell **23,** 600-627.

Baker NR. 2008. Chlorophyll fluorescence: a probe of photosynthesis in vivo. Annual Review of Plant Biology **59,** 89-113.

Balcerowicz M, Hoecker U. 2014. Auxin–a novel regulator of stomata differentiation. Trends in Plant Science **19,** 747-749.

Blamey FPC, Joyce DC, Edwards DG, Asher CJ. 1986. Role of trichomes in sunflower tolerance to manganese toxicity. Plant and Soil **91,** 171-180.

Blamey FPC, Hernandez-Soriano MC, Cheng M, Tang C, Paterson DJ, Lombi E, Wang WH, Scheckel KG, Kopittke PM. 2015. Synchrotron-based techniques shed light on mechanisms of plant sensitivity and tolerance to high manganese in the root environment. Plant Physiology **169,** 2006-2020.

Bloom AJ, Kameritsch P. 2017. Relative association of Rubisco with manganese and magnesium as a regulatory mechanism in plants. Physiologia Plantarum **161,** 545-559.

Brestic M, Zivcak M, Kunderlikova K, Sytar O, Shao H, Kalaji HM, Allakhverdiev SI. 2015. Low PSI content limits the photoprotection of PSI and PSII in early growth stages of chlorophyll b-deficient wheat mutant lines. Photosynthesis Research **125,** 151-166.

Broadley M, Brown P, Cakmak I, Rengel Z, Zhao F. 2012. Function of nutrients: micronutrients. In *Marschner's mineral nutrition of higher plants* (eds Marschner, P.), pp. 135-189. Academic Press, London.

Chatterjee C, Nautiyal N, Agarwala SC. 1994. Influence of changes in manganese and magnesium supply on some aspects of wheat physiology. Soil Science and Plant Nutrition **40,** 191-197.

Chen Z, Fujii Y, Yamaji N, *et al.* 2013. Mn tolerance in rice is mediated by MTP8. 1, a member of the cation diffusion facilitator family. Journal of Experimental Botany **64,** 4375-4387.

Clairmont KB, Hagar WG, Davis EA. 1986. Manganese toxicity to chlorophyll synthesis in tobacco callus. Plant Physiology **80,** 291-293.

Delhaize E, Kataoka T, Hebb DM, White RG, Ryan PR. 2003. Genes encoding proteins of the cation diffusion facilitator family that confer manganese tolerance. The Plant Cell **15,** 1131-1142.

Du H, Wu N, Fu J, Wang S, Li X, Xiao J, Xiong L. 2012. A GH3 family member, OsGH3-2, modulates auxin and abscisic acid levels and differentially affects drought and cold tolerance in rice. Journal of Experimental Botany **63,** 6467-6480.

Earles JM, Theroux-Rancourt G, Roddy AB, Gilbert ME, McElrone AJ, Brodersen CR. 2018. Beyond porosity: 3D leaf intercellular airspace traits that impact mesophyll conductance. Plant Physiology **178,** 148-162.

Eroglu S, Meier B, von Wirén N, Peiter E. 2016. The vacuolar manganese transporter MTP8 determines tolerance to iron deficiency-induced chlorosis in *Arabidopsis*. Plant Physiology **170,**

1030-1045.

Fageria NK, Stone LF. 2008. Micronutrient deficiency problems in South America.

In *Micronutrient deficiencies in global crop production* (pp. 245-266). Springer, Dordrecht.

Fecht-Christoffers MM, Braun HP, Lemaitre-Guillier C, VanDorsselaer A, Horst WJ. 2003a.

Effect of manganese toxicity on the proteome of the leaf apoplast in cowpea. Plant

Physiology **133,** 1935-1946.

Fecht-Christoffers MM, Führs H, Braun HP, Horst WJ. 2006. The role of hydrogen

peroxide-producing and hydrogen peroxide-consuming peroxidases in the leaf apoplast of

cowpea in manganese tolerance. Plant Physiology **140,** 1451-1463.

Fecht-Christoffers MM, Maier P, Horst WJ. 2003b. Apoplastic peroxidases and ascorbate are

involved in manganese toxicity and tolerance of Vigna unguiculata. Physiologia Plantarum **117,**

237-244.

Fernando DR, Bakkaus EJ, Perrier N, Baker AJM, Woodrow IE, Batianoff GN, Collins RN.

2006. Manganese accumulation in the leaf mesophyll of four tree species: a PIXE/EDAX

localization study. New Phytologist **171,** 751-758.

Fernando DR, Lynch JP. 2015. Manganese phytotoxicity: new light on an old problem. Annals

of Botany **116,** 313-319.

Fernando DR, Moroni SJ, Scott BJ, Conyers MK, Lynch J, Marshall AT. 2016. Temperature

and light drive manganese accumulation and stress in crops across three major plant

families. Environmental and Experimental Botany **132,** 66-79.

Foy CD, Chaney RT, White MC. 1978. The physiology of metal toxicity in plants. Annual Review

of Plant Physiology **29,** 511-566.

Franks PJ, Beerling DJ. 2009. Maximum leaf conductance driven by CO2 effects on stomatal

size and density over geologic time. Proceedings of the National Academy of Sciences **106,**

10343-10347.

Führs H, Behrens C, Gallien S, Heintz D, Van Dorsselaer A, Braun HP, Horst WJ. 2010.

Physiological and proteomic characterization of manganese sensitivity and tolerance in rice

(*Oryza sativa*) in comparison with barley (*Hordeum vulgare*). Annals of Botany **105,** 1129-1140.

Führs H, Götze S, Specht A, Erban A, Gallien S, Heintz D, Van Dorsselaer A, Kopka J,

Braun H-P, Horst WJ. 2009. Characterization of leaf apoplastic peroxidases and metabolites in

Vigna unguiculata in response to toxic manganese supply and silicon. Journal of Experimental

Botany **60,** 1663-1678.

Fujii T, Nakano R, Takubo Y, Qian S, Yamakawa R, Ando T, Ishikawa Y. 2010. Female sex

pheromone of a lichen moth Eilema japonica (Arctiidae, Lithosiinae): components and control of

production. Journal of Insect Physiology **56,** 1986-1991.

González A, Lynch JP 1997. Effects of manganese toxicity on leaf CO_2 assimilation of

contrasting common bean genotypes. Physiologia Plantarum **101,** 872-880.

González A, Steffen K., Lynch JP 1998. Light and excess manganese: implications for

oxidative stress in common bean. Plant Physiology **118,** 493-504.

Gutierrez CK, Matsui GY, Lincoln DE, Lovell CR. 2009. Production of the phytohormone

indole-3-acetic acid by estuarine species of the genus Vibrio. Applied and Environmental

Microbiology **75,** 2253-2258.

Guo X, Lu W, Ma Y, Qin Q, Hou S. 2013. The BIG gene is required for auxin-mediated organ

growth in *Arabidopsis*. Planta **237,** 1135-1147.

Hachiya T, Watanabe CK, Boom C, Tholen D, Takahara K, Kawai-Yamada M, Uchimiya H,

Uesono Y, Terashima I. Noguchi K. 2010. Ammonium‐dependent respiratory increase is

dependent on the cytochrome pathway in Arabidopsis thaliana shoots. Plant, Cell &

Environment **33,** 1888-1897.

Hibberd JM, Quick WP. 2002. Characteristics of C4 photosynthesis in stems and petioles of C3

flowering plants. Nature **415,** 451-454.

Hirschi KD, Korenkov VD, Wilganowski NL, Wagner GJ. 2000. Expression of Arabidopsis

CAX2 in tobacco. Altered metal accumulation and increased manganese tolerance. Plant

Physiology **124,** 125-134.

Horiguchi T. 1987. Mechanism of manganese toxicity and tolerance of plants: II. Deposition of

oxidized manganese in plant tissues. Soil Science and Plant Nutrition **33,** 595-606.

Horst WJ, Fecht M, Naumann A, Wissemeier AH, Maier P. 1999. Physiology of manganese

toxicity and tolerance in *Vigna unguiculata* (L.) Walp. Journal of Plant Nutrition and Soil

Science **162,** 263-274.

Houtz RL, Nable RO, Cheniae GM. 1988. Evidence for effects on the in vivo activity of

ribulose-bisphosphate carboxylase/oxygenase during development of Mn toxicity in

tobacco. Plant Physiology **86,** 1143-1149.

Jain M, Kaur N, Garg R, Thakur JK, Tyagi AK, Khurana JP. 2006. Structure and expression

analysis of early auxin-responsive Aux/IAA gene family in rice (*Oryza sativa*). Functional &

Integrative Genomics **6,** 47-59.

Jezek M, Blatt MR. 2017. The membrane transport system of the guard cell and its integration

for stomatal dynamics. Plant Physiology **174,** 487-519.

Jordan DB, Ogren WL. 1983. Species variation in kinetic properties of ribulose 1,

5-bisphosphate carboxylase/oxygenase. Archives of Biochemistry and Biophysics **227,** 425-433.

Kenten RH, Mann PJG. 1957. Manganese oxidation in the pea plant (Pisum sativum L.) grown

under conditions of manganese toxicity. Biochemical Journal **65,** 179-185.

Kitao M, Lei TT, Koike T. 1997. Effects of manganese toxicity on photosynthesis of white birch

(Betula platyphylla var. japonica) seedlings. Physiologia Plantarum **101,** 249-256.

Klein M, Perfus-Barbeoch L, Frelet A, Gaedeke N, Reinhardt D, Mueller-Roeber B,

Martinoia E, Forestier C. 2003. The plant multidrug resistance ABC transporter AtMRP5 is

involved in guard cell hormonal signalling and water use. The Plant Journal **33,** 119-129.

Klughammer C, Schreiber U. 1994. Saturation pulse method for assessment of energy

conversion in PS I. Planta **192,** 261-268.

Kramer DM, Sacksteder CA, Cruz JA. 1999. How acidic is the lumen?. Photosynthesis

Research **60,** 151-163.

Lagrimini LM, Joly RJ, Dunlap JR, Liu TTY. 1997a. The consequence of peroxidase

overexpression in transgenic plants on root growth and development. Plant Molecular Biology **33,**

887-895.

Lagrimini LM, Gingas V, Finger F, Rothstein S, Liu TTY. 1997b. Characterization of antisense transformed plants deficient in the tobacco anionic peroxidase. Plant Physiology **114,** 1187-1196.

Le J, Liu XG, Yang KZ, *et al.* (2014). Auxin transport and activity regulate stomatal patterning and development. Nature Communications **5,** 1-8.

Lei Y, Korpelainen H, Li C. 2007. Physiological and biochemical responses to high Mn concentrations in two contrasting *Populus cathayana* populations. Chemosphere **68,** 686-694.

Li J, Jia Y, Dong R, Huang R, Liu P, Li X, Wang X, Liu G, Chen Z. 2019. Advances in the mechanisms of plant tolerance to manganese toxicity. International Journal of Molecular Sciences **20,** 5096.

Lidon FC. 2002. Rice plant structural changes by addition of excess manganese. Journal of Plant Nutrition **25,** 287-296.

Lidon FC, Barreiro MG, Ramalho JC. 2004. Manganese accumulation in rice: implications for photosynthetic functioning. Journal of Plant Physiology **161,** 1235-1244.

Lidon FC, Teixeira MG. 2000a. Oxy radicals production and control in the chloroplast of Mn-treated rice. Plant Science **152,** 7-15.

Lidon FC, Teixeira MG. 2000b. Rice tolerance to excess Mn: implications in the chloroplast lamellae and synthesis of a novel Mn protein. Plant Physiology and Biochemistry **38,** 969-978.

Ljung K, Hull AK, Celenza J, Yamada M, Estelle M, Normanly J, Sandberg G. 2005. Sites

1009 and regulation of auxin biosynthesis in Arabidopsis roots. The Plant Cell **17,** 1090-1104.

1010

1011 **Macfie SM, Taylor GJ.** 1992. The effects of excess manganese on photosynthetic rate and

1012 concentration of chlorophyll in *Triticum aestivum* grown in solution culture. Physiologia

1013 Plantarum **85,** 467-475.

1014

1015 **Makino A, Mae T, Ohira K.** 1988. Differences between wheat and rice in the enzymic properties

1016 of ribulose-1,5-bisphosphate carboxylase/oxygenase and the relationship to photosynthetic gas

1017 exchange. Planta, **174,** 30-38.

1018

1019 **Makino A, Nakano H, Mae T.** 1994. Responses of ribulose-1,5-bisphosphate carboxylase,

1020 cytochrome *f*, and sucrose synthesis enzymes in rice leaves to leaf nitrogen and their

1021 relationships to photosynthesis. Plant Physiology **105,** 173-179.

1022

1023 **McDaniel KL, Toman FR.** 1994. Short‐term effects of manganese toxicity on ribulose 1, 5

1024 bisphosphate carboxylase in tobacco chloroplasts. Journal of Plant Nutrition **17,** 523-536.

1025

1026 **McHargue JS.** 1922. The role of manganese in plants. Journal of the American Chemical

1027 Society **44,** 1592-1598.

1028

1029 **Millaleo R, Reyes-Díaz M, Alberdi M, Ivanov AG, Krol M, Hüner NPA.** 2013. Excess

1030 manganese differentially inhibits photosystem I versus II in Arabidopsis thaliana. Journal of

1031 Experimental Botany **64,** 343-354.

1032

1033 **Millaleo R, Reyes-Díaz M, Ivanov AG, Mora ML, Alberdi M.** 2010. Manganese as essential

1034 and toxic element for plants: transport, accumulation and resistance mechanisms. Journal of Soil

1035 Science and Plant Nutrition **10,** 470-481.

1036

1037 **Moraes TA, Mengin V, Annunziata MG, Encke B, Krohn N, Hoehne M, Stitt M.** 2019.

1038 Response of the circadian clock and diel starch turnover to one day of low light or low CO_2. Plant

1039 Physiology **179,** 1457-1478.

1040

1041 **Morgan PW, Joham HE, Amin JV.** 1966. Effect of manganese toxicity on the indoleacetic acid

1042 oxidase system of cotton. Plant Physiology **41,** 718-724.

1043

1044 **Morita R, Sugino M, Hatanaka T, Misoo S, Fukayama H.** 2015. CO_2-responsive CONSTANS,

1045 CONSTANS-like, and time of chlorophyll a/b binding protein Expression1 protein is a positive

1046 regulator of starch synthesis in vegetative organs of rice. Plant Physiology **167,** 1321-1331.

1047

1048 **Muñoz−Nortes T, Pérez−Pérez JM, Ponce MR, Candela H, Micol JL.** 2017. The ANGULATA 7

1049 gene encodes a DnaJ‐like zinc finger‐domain protein involved in chloroplast function and leaf

1050 development in Arabidopsis. The Plant Journal **89,** 870-884.

1051

1052 **Nable RO, Houtz RL, Cheniae GM.** 1988. Early inhibition of photosynthesis during development

1053 of Mn toxicity in tobacco. Plant Physiology **86,** 1136-1142.

1054

1055 **Nakano H, Makino A, Mae T.** 1997. The effect of elevated partial pressures of CO_2 on the

1056 relationship between photosynthetic capacity and N content in rice leaves. Plant Physiology **115,**

1057 191-198.

1058

1059 **Nghi KN, Tagliani A, Mariotti L, Weits DA, Perata P, Pucciariello C.** 2021. Auxin is required

1060 for the long coleoptile trait in japonica rice under submergence. New Phytologist **229,** 85-93.

1061

1062 **Noguchi Ko, Terashima I.** 2006. Responses of spinach leaf mitochondria to low N

1063 availability. Plant, Cell & Environment **29,** 710-719.

1064

1065 **Panda S, Mishra AK, Biswal UC.** 1987. Manganese induced peroxidation of thylakoid lipids and
1066 changes in chlorophyll-α fluorescence during aging of cell free chloroplasts in
1067 light. Phytochemistry **26,** 3217-3219.

1068

1069 **Pérez-Pérez JM, Candela H, Robles P, López-Torrejón G, Del Pozo JC, Micol JL.** 2010. A
1070 role for AUXIN RESISTANT3 in the coordination of leaf growth. Plant and Cell Physiology **51,**
1071 1661-1673.

1072

1073 **Porra RJ, Thompson WA, Kriedemann PE.** 1989. Determination of accurate extinction
1074 coefficients and simultaneous equations for assaying chlorophylls *a* and *b* extracted with four
1075 different solvents: verification of the concentration of chlorophyll standards by atomic absorption
1076 spectroscopy. Biochimica et Biophysica Acta-Bioenergetics **975,** 1104 384-394.

1077

1078 **Qi J, Wang Y, Yu T, Cunha A, Wu B, Vernoux T, Meyerowitz E, Jiao Y.** 2014. Auxin depletion
1079 from leaf primordia contributes to organ patterning. Proceedings of the National Academy of
1080 Sciences **111,** 18769-18774.

1081

1082 **Ray PM.** 1958. Destruction of auxin. Annual Review of Plant Physiology **9,** 81-118.

1083

1084 **Perrot-Rechenmann C.** 2010. Cellular responses to auxin: division versus expansion. Cold
1085 Spring Harbor Perspectives in Biology **2,** a001446.

1086

1087 **Robert HS, Friml J.** 2009. Auxin and other signals on the move in plants. Nature Chemical
1088 Biology **5,** 325-332.

1089

1090 **Santos EF, Santini JMK, Paixão AP, Júnior EF, Lavres J, Campos M, Dos Reis AR.** 2017.
1091 Physiological highlights of manganese toxicity symptoms in soybean plants: Mn toxicity
1092 responses. Plant Physiology and Biochemistry **113,** 6-19.

1093

1094   **Sasaki A, Yamaji N, Xia J, Ma JF.** 2011. OsYSL6 is involved in the detoxification of excess

1095   manganese in rice. Plant Physiology **157,** 1832-1840.

1096

1097   **Shao JF, Yamaji N, Shen RF, Ma JF.** 2017. The key to Mn homeostasis in plants: regulation of

1098   Mn transporters. Trends in Plant Science **22,** 215-224.

1099

1100   **Sirkar S, Amin JV.** 1974. The manganese toxicity of cotton. Plant Physiology **54,** 539-543.

1101

1102   **Suga M, Akita F, Hirata K,** *et al.* 2015. Native structure of photosystem II at 1.95 Å resolution

1103   viewed by femtosecond X-ray pulses. Nature **517,** 99-103.

1104

1105   **Suganami M, Suzuki Y, Tazoe Y, Yamori W, Makino A**. 2021. Co-overproducing Rubisco

1106   and Rubisco activase enhances photosynthesis in the optimal temperature range in

1107   rice. Plant Physiology, **185,** 108–119.

1108

1109   **Sulpice R, Flis A, Ivakov AA, Apelt F, Krohn N, Encke B, Abel C, Feil R, Lunn JE, Stitt M**.

1110   2014. Arabidopsis coordinates the diurnal regulation of carbon allocation and growth across a

1111   wide range of photoperiods. Molecular Plant **7,** 137-155.

1112

1113   **Suresh R, Foy CD, Weidner JR.** 1987. Effects of excess soil manganese on stomatal function in

1114   two soybean cultivars. Journal of Plant Nutrition **10,** 749-760.

1115

1116   **Suzuki Y, Fujimori T, Kanno K, Sasaki A, Ohashi Y, Makino A.** 2012. Metabolome analysis of

1117   photosynthesis and the related primary metabolites in the leaves of transgenic rice plants with

1118   increased or decreased Rubisco content. Plant, Cell & Environment **35,** 1369-1379.

1119

1120   **Suzuki Y, Kawazu T, Koyama H.** 2004. RNA isolation from siliques, dry seeds and other

1121   tissues of Arabidopsis thaliana. Biotechniques **37,** 542-544.

1122

1123   **Takagi D, Amako K, Hashiguchi M,** *et al.* 2017. Chloroplastic ATP synthase builds up a proton

1124   motive force preventing production of reactive oxygen species in photosystem I. The Plant

1125   Journal **91,** 306-324.

1126

1127   **Takagi D, Hashiguchi M, Sejima T, Makino A, Miyake C.** 2016b. Photorespiration provides the

1128   chance of cyclic electron flow to operate for the redox-regulation of P700 in photosynthetic

1129   electron transport system of sunflower leaves. Photosynthesis Research **129,** 279-290.

1130

1131   **Takagi D, Ifuku K, Ikeda KI, Inoue KI, Park P, Tamoi M, Inoue H, Sakamoto K, Saito R,**

1132   **Miyake C.** 2016a. Suppression of chloroplastic alkenal/one oxidoreductase represses the

1133   carbon catabolic pathway in Arabidopsis leaves during night. Plant Physiology **170,** 2024-2039.

1134

1135   **Takagi D, Miyagi A, Tazoe Y, Suganami M, Kawai–Yamada M, Ueda A, Suzuki Y, Noguchi K,**

1136   **Hirotsu N, Makino A.** 2020. Phosphorus toxicity disrupts Rubisco activation and reactive

1137   oxygen species defence systems by phytic acid accumulation in leaves. Plant, Cell &

1138   Environment **43,** 2033-2053.

1139

1140   **Takahama U.** 2004. Oxidation of vacuolar and apoplastic phenolic substrates by peroxidase:

1141   physiological significance of the oxidation reactions. Phytochemistry Reviews **3,** 207-219.

1142

1143   **Tognetti VB, Mühlenbock PER, Van Breusegem F.** 2012. Stress homeostasis–the redox and

1144   auxin perspective. Plant, Cell & Environment, **35,** 321-333.

1145

1146   **Tsunemitsu Y, Genga M, Okada T, Yamaji N, Ma JF, Miyazaki A, Kato S, Iwasaki K, Ueno D.**

1147   2018. A member of cation diffusion facilitator family, MTP11, is required for manganese tolerance

1148   and high fertility in rice. Planta **248,** 231-241.

1149

Tronconi MA, Fahnenstich H, Weehler MCG, Andreo CS, Flügge UI, Drincovich MF, Maurino VG. 2008. Arabidopsis NAD-malic enzyme functions as a homodimer and heterodimer and has a major impact on nocturnal metabolism. Plant Physiology **146,** 1540-1552.

Valdés−López O, Yang SS, Aparicio−Fabre R, Graham PH, Reyes JL, Vance CP, Hernández G. 2010. MicroRNA expression profile in common bean (Phaseolus vulgaris) under nutrient deficiency stresses and manganese toxicity. New Phytologist **187,** 805-818.

Vlamis J, Williams DE. 1964. Iron and manganese relations in rice and barley. Plant and Soil **20,** 221-231.

von Caemmerer S, Farquhar GD. 1981. Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. Planta **153,** 376-387

Wada S, Takagi D, Miyake C, Makino A, Suzuki Y. 2019. Responses of the photosynthetic electron transport reactions stimulate the oxidation of the reaction center chlorophyll of photosystem I, P700, under drought and high temperatures in rice. International Journal of Molecular Sciences **20,** 2068.

Walle, F, Furuya M, Nick P. 2002. OsARF1, an auxin response factor from rice, is auxin-regulated and classifies as a primary auxin responsive gene. Plant Molecular Biology 50, 415-425.

Wang D, Pei K, Fu Y, Sun Z, Li S, Liu H, Tang K, Han B, Tao Y. 2007. Genome-wide analysis of the auxin response factors (ARF) gene family in rice (*Oryza sativa*). Gene 394, 13-24.

Williams TC, Miguet L, Masakapalli SK, Kruger NJ, Sweetlove LJ, Ratcliffe RG. 2008.

Metabolic network fluxes in heterotrophic Arabidopsis cells: stability of the flux distribution under different oxygenation conditions. Plant Physiology 148, 704-718.

Wissemeier AH, Horst WJ. 1987. Callose deposition in leaves of cowpea (Vigna unguiculata [L.] Walp.) as a sensitive response to high Mn supply. Plant and Soil 102, 283-286.

Wissemeier AH, Horst WJ. 1992. Effect of light intensity on manganese toxicity symptoms and callose formation in cowpea (*Vigna unguiculata* (L.) Walp.). Plant and Soil 143, 299-309.

Zhang JY, He SB, Li L, Yang HQ. 2014. Auxin inhibits stomatal development through MONOPTEROS repression of a mobile peptide gene STOMAGEN in mesophyll. Proceedings of the National Academy of Sciences 111, E3015-E3023.

Zhao Z, Zhang Y, Liu X, *et al.* 2013. A role for a dioxygenase in auxin metabolism and reproductive development in rice. Developmental Cell 27, 113-122.

Zivcak M, Brestic M, Balatova Z, Drevenakova P, Olsovska K, Kalaji HM, Yang X, Allakhverdiev SI. 2013. Photosynthetic electron transport and specific photoprotective responses in wheat leaves under drought stress. Photosynthesis Research 117, 529-546.

 FIGURE LEGENDS

Figure 1

Phenotypes of rice plants grown under control and Mn-toxic conditions. The pictures of fully

expanded leaf blades grown under control and Mn-toxic conditions are shown in (A), and (B)

shows an enlarged picture of these. The white bars indicate lengths of 1 cm (A) and 1 mm

(B), respectively. (C) The dried weights of the leaf blade, leaf sheath, and root, respectively.

(D) The sum of the dried weights shown in (C). Data are shown as box plots (n = 9), black

squares indicate the mean value, and bars indicate the range of the maximum or minimum

data within a 1.5 × interquartile range (IQR). Green boxes indicate the results of the control

conditions, and red boxes indicate those of the Mn-toxic conditions. Asterisks show

significant differences between the control and Mn-toxic conditions (*: $p < 0.05$,

Kruskal–Wallis test).

1217

Figure 2

Mineral concentration in the rice leaf blade. Data are shown as box plots (control conditions

n = 8, Mn-toxic conditions n = 7), black squares indicate the mean value, and bars indicate

the range of the maximum or minimum data within a 1.5 × interquartile range (IQR). Green

boxes indicate the results of the control conditions, and red boxes indicate those of the

Mn-toxic conditions. Asterisks show significant differences between the control and Mn-toxic

conditions (*: $p < 0.05$, **: $p < 0.01$, Kruskal–Wallis test).

1225

Figure 3

Photosynthetic activities in leaves grown under control and the Mn-toxic conditions. The

CO_2 assimilation rate (A), stomatal conductance (g_s) (B), and intercellular CO_2 concentration

(C) are shown. For the PSII photosynthetic parameters, the quantum yields of PSII [Y(II)] (D),

non-photochemical quenching [Y(NPQ)] (E), and non-radiative decay [Y(NO)] (F) are shown.

The maximum quantum yield of PSII (Fv/Fm) is shown (J). For PSI photosynthetic

parameters, the quantum yields of PSI [Y(I)] (G), non-photochemical quenching at the donor

side [Y(ND)] (H), and non-photochemical quenching at the acceptor side [Y(NA)] (I) are shown. (K) and (L) show the total Chl and N concentrations in the leaves, respectively. Data of photosynthetic parameters are shown as means with standard deviation (SD) (n = 7). The results for Fv/Fm (n = 7) and total Chl (n = 7) and N (n = 9) concentrations are shown as box plots. Black squares indicate the mean value, and bars indicate the range of the maximum or minimum data within a 1.5 × interquartile range (IQR). Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, **: $p < 0.01$, Kruskal–Wallis test).

Figure 4

Carbohydrate content in leaf blades at the end of the day and end of the night. Data are shown as means with standard deviation (SD) (n = 3–4), and squares and diamonds show the distribution of the raw data. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, Kruskal–Wallis test).

Figure 5

Mitochondrial respiration activities and amino acid contents in leaf blades grown under control and Mn-toxic conditions. (A) Leaf respiration rate (n = 12–13) in the dark and (B) sucrose feeding respiration activities and mitochondrial respiratory electron transport activities depending on ATP synthase (uncoupled), COX, and AOX (n = 7). (C) Amino acid content in leaves sampled in the middle of the night (n = 6). (D–I) Enzyme activities involved in the TCA cycle (n = 13–14). Data are shown as box plots, black squares indicate the mean value, and bars indicate the range of the maximum or minimum data within a 1.5 × interquartile range (IQR). Green boxes indicate the results under control conditions, and red boxes indicate those under Mn-toxic conditions. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, **: $p < 0.01$, Kruskal–Wallis test).

Figure 6

Stomatal structures and characteristics in the rice leaf blade. Pictures of the stomata on the adaxial side of the leaf blade grown under control (A) and Mn-toxic (B) conditions. The upper pictures focus on the stomatal distributions, and the lower pictures focus on the single stomata complex. (C) Showed the stomatal density (n = 10–12). (D) and (E) showed the major and the minor axis of the stomatal complex, respectively (n = 53-63). Data are shown as box plots, black squares indicate the mean value, and bars indicate the range of the maximum or minimum data within a 1.5 × interquartile range (IQR). Green boxes indicate the results under control conditions, and red boxes indicate those under Mn-toxic conditions. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, ***: $p < 0.001$, Kruskal–Wallis test).

Figure 7

Characteristics of the leaf section grown under control and Mn-toxic conditions. (A) Fixed leaf sections sectioned from the horizontal and vertical side against the leaf vein. Fresh leaf sections of the small (B) and large (C) bundles. Black bars show the scales.

Figure 8

Leaf indole acetic acid (IAA) concentration and gene expression analysis relating to IAA and stomata development. (A) The result of GC-MS analysis targeting IAA, together with the chemical structures of IAA and its derivatives. The total ion chromatography (TIC) and chromatographs of $m/z = 202$ and $m/z = 319$ in the standard IAA solution and the leaf extract from plants under control and Mn-toxic conditions are shown. The retention time of the silylated IAA was 15.2 min. (B) The IAA concentrations in the leaf blade (n = 8). (D) The results of gene expression analysis involving IAA-responsive gene and stomatal patterning (n = 8). The quantified results are expressed on an *OsATC1-2* expression basis. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, **: $p < 0.01$, Kruskal–Wallis test).

Figure 9

Visual scheme of symplastic Mn toxicity to suppress CO_2 assimilation. When Mn is excessively accumulated in leaves, the leaf IAA concentration is lowered, which might cause the stimulation of IAA degradation activities under Mn toxicity (Morgan et al., 1996). As a short-term effect, the decrease in IAA concentration affects stomatal opening (Blatt & Thiel, 1994; Klein et al., 2003). Subsequently, as a long-term effect, the auxin-dependent signal transduction involving ARF transcriptional factor is perturbated in the leaves to cause an IAA-deficient phenotype, changing both stomatal and leaf anatomical structures (Le et al., 2014; Zhang et al., 2014). The change in stomatal function and leaf structure severely limits CO_2 diffusion to the chloroplasts, and CO_2 assimilation by Rubisco is inhibited. In contrast, the photorespiration reaction limits the photosynthetic electron activities to cause ROS production (Wada et al., 2019). The decreased sugar production efficiency in photosynthesis suppresses sugar catabolism. Therefore, growth is inhibited under Mn-toxic conditions.

Figure 1

Phenotypes of rice plants grown under control and Mn-toxic conditions. The pictures of fully expanded leaf blades grown under control and Mn-toxic conditions are shown in (A), and (B) shows an enlarged picture of these. The white bars indicate lengths of 1 cm (A) and 1 mm (B), respectively. (C) The dried weights of the leaf blade, leaf sheath, and root, respectively. (D) The sum of the dried weights shown in (C). Data are shown as box plots (n = 9), black squares indicate the mean value, and bars indicate the range of the maximum or minimum data within a 1.5× interquartile range (IQR). Green boxes indicate the results of the control conditions, and red boxes indicate those of the Mn-toxic conditions. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, Kruskal–Wallis test).

Figure 2

Mineral concentration in the rice leaf blade. Data are shown as box plots (control conditions n = 8, Mn-toxic conditions n = 7), black squares indicate the mean value, and bars indicate the range of the maximum or minimum data within a 1.5× interquartile range (IQR). Green boxes indicate the results of the control conditions, and red boxes indicate those of the Mn-toxic conditions. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, **: $p < 0.01$, Kruskal–Wallis test).

Figure 3

Photosynthetic activities in leaves grown under control and the Mn-toxic conditions. The CO_2 assimilation rate (A), stomatal conductance (g_s) (B), and intercellular CO_2 concentration (C) are shown. For the PSII photosynthetic parameters, the quantum yields of PSII [Y(II)] (D), non-photochemical quenching [Y(NPQ)] (E), and non-radiative decay [Y(NO)] (F) are shown. The maximum quantum yield of PSII (Fv/Fm) is shown (J). For PSI photosynthetic parameters, the quantum yields of PSI [Y(I)] (G), non-photochemical quenching at the donor side [Y(ND)] (H), and non-photochemical quenching at the acceptor side [Y(NA)] (I) are shown. (K) and (L) show the total Chl and N concentrations in the leaves, respectively. Data of photosynthetic parameters are shown as means with standard deviation (SD) (n = 7). The results for Fv/Fm (n = 7) and total Chl (n = 7) and N (n = 9) concentrations are shown as box plots. Black squares indicate the mean value, and bars indicate the range of the maximum or minimum data within a 1.5× interquartile range (IQR). Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, **: $p < 0.01$, Kruskal–Wallis test).

Figure 4

Carbohydrate content in leaf blades at the end of the day and end of the night. Data are shown as means with standard deviation (SD) (n = 3–4), and squares and diamonds show the distribution of the raw data. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, Kruskal–Wallis test).

Figure 5

Mitochondrial respiration activities and amino acid contents in leaf blades grown under control and Mn-toxic conditions. (A) Leaf respiration rate (n = 12–13) in the dark and (B) sucrose feeding respiration activities and mitochondrial respiratory electron transport activities depending on ATP synthase (uncoupled), COX, and AOX (n = 7). (C) Amino acid content in leaves sampled in the middle of the night (n = 6). (D–I) Enzyme activities involved in the TCA cycle (n = 13–14). Data are shown as box plots, black squares indicate the mean value, and bars indicate the range of the maximum or minimum data within a 1.5× interquartile range (IQR). Green boxes indicate the results under control conditions, and red boxes indicate those under Mn-toxic conditions. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, **: $p < 0.01$, Kruskal–Wallis test).

A; The control-conditions

B; The Mn-toxic conditions

Figure 6

Stomatal structures and characteristics in the rice leaf blade. Pictures of the stomata on the adaxial side of the leaf blade grown under control (A) and Mn-toxic (B) conditions. The upper pictures focus on the stomatal distributions, and the lower pictures focus on the single stomata complex. (C) Showed the stomatal density (n = 10–12). (D) and (E) showed the major and the minor axis of the stomatal complex, respectively (n = 53-63). Data are shown as box plots, black squares indicate the mean value, and bars indicate the range of the maximum or minimum data within a 1.5× interquartile range (IQR). Green boxes indicate the results under control conditions, and red boxes indicate those under Mn-toxic conditions. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, ***: $p < 0.001$, Kruskal–Wallis test).

Figure 7

Characteristics of the leaf section grown under control and Mn-toxic conditions. (A) Fixed leaf sections sectioned from the horizontal and vertical side against the leaf vein. Fresh leaf sections of the small (B) and large (C) bundles. Black bars show the scales.

Figure 8

Leaf indole acetic acid (IAA) concentration and gene expression analysis relating to IAA and stomata development. (A) The result of GC-MS analysis targeting IAA, together with the chemical structures of IAA and its derivatives. The total ion chromatography (TIC) and chromatographs of m/z = 202 and m/z = 319 in the standard IAA solution and the leaf extract from plants under control and Mn-toxic conditions are shown. The retention time of the silylated IAA was 15.2 min. (B) The IAA concentrations in the leaf blade (n = 8). (D) The results of gene expression analysis involving IAA-responsive gene and stomatal patterning (n = 8). The quantified results are expressed on an *OsATC1-2* expression basis. Asterisks show significant differences between the control and Mn-toxic conditions (*: $p < 0.05$, **: $p < 0.01$, Kruskal–Wallis test).

Figure 9

Visual scheme of symplastic Mn toxicity to suppress CO_2 assimilation. When Mn is excessively accumulated in leaves, the leaf IAA concentration is lowered, which might cause the stimulation of IAA degradation activities under Mn toxicity (Morgan *et al.,* 1996). As a short-term effect, the decrease in IAA concentration affects stomatal opening (Blatt and Thiel, 1994; Klein *et al.,* 2003). Subsequently, as a long-term effect, the auxin-dependent signal transduction involving ARF transcriptional factor is perturbated in the leaves to cause an IAA-deficient phenotype, changing both stomatal and leaf anatomical structures (Le *et al.*, 2014; Zhang *et al.,* 2014). The change in stomatal function and leaf structure severely limits CO_2 diffusion to the chloroplasts, and CO_2 assimilation by Rubisco is inhibited. In contrast, the photorespiration reaction limits the photosynthetic electron activities to cause ROS production (Wada *et al.,* 2019). The decreased sugar production efficiency in photosynthesis suppresses sugar catabolism. Therefore, growth is inhibited under Mn-toxic conditions.