A genomic and morphometric analysis of alpine bumblebees: ongoing reductions in

tongue length but no clear genetic component

Running title: Genome variation in alpine bumblebees

Matthew J. Christmas[1], Julia C. Jones[1,2], Anna Olsson[1], Ola Wallerman[1], Ignas Bunikis[3],

Marcin Kierczak[4], Kaitlyn M. Whitley[5,6], Isabel Sullivan[5,7], Jennifer C. Geib[5], Nicole E. Miller-

Struttmann[8], Matthew T. Webster[1]

1) Department of Medical Biochemistry and Microbiology, Science for Life Laboratory,

Uppsala University, Uppsala, Sweden

2) School of Biology and Environmental Science, University College Dublin, Dublin, Ireland

3) Department of Immunology, Genetics and Pathology, Science for Life Laboratory, Uppsala

University, Uppsala, Sweden

4) Department of Cell and Molecular Biology, National Bioinformatics Infrastructure

Sweden, Science for Life Laboratory, Uppsala University, Uppsala, Sweden

5) Department of Biology, Appalachian State University, Boone, North Carolina, USA

6) U.S. Department of Agriculture, Agriculture Research Service, Charleston, South Carolina,

USA

7) Marine Estuarine Environmental Sciences, University of Maryland, College Park Maryland

8) Biological Sciences Department, Webster University, St. Louis, Missouri, USA

Corresponding author: Matthew T. Webster

Address: Dept. Medical Biochemistry and Microbiology, Box 582, 75123 Uppsala, Sweden

Email: matthew.webster@imbim.uu.se

Tel: +46 18 471 4391

ABSTRACT

Over the last six decades, populations of the bumblebees *Bombus sylvicola* and *Bombus balteatus* in Colorado have experienced decreases in tongue length, a trait important for plant-pollinator mutualisms. It has been hypothesized that this observation reflects selection resulting from shifts in floral composition under climate change. Here we use morphometrics and population genomics to determine whether morphological change is ongoing, investigate the genetic basis of morphological variation, and analyze population structure in these populations. We generate a genome assembly of *B. balteatus*. We then analyze whole-genome sequencing data and morphometric measurements of 580 samples of both species from seven high-altitude localities. Out of 281 samples originally identified as *B. sylvicola*, 67 formed a separate genetic cluster comprising a newly-discovered cryptic species ("incognitus"). However, an absence of genetic structure within species suggests that gene flow is common between mountains. We find a significant decrease in tongue length between bees collected between 2012-2014 and in 2017, indicating that morphological shifts are ongoing. We did not discover any genetic associations with tongue length, but a SNP related to production of a proteolytic digestive enzyme is implicated in body size variation. We identify evidence of covariance between kinship and both tongue length and body size, which is suggestive of a genetic component of these traits, although it is possible that shared environmental effects between colonies are responsible. Our results provide evidence for ongoing modification of a morphological trait important for pollination and indicate that this trait likely has a complex genetic and environmental basis.

INTRODUCTION

The importance of pollinators for ecology and the maintenance of biodiversity is widely recognized (Goulson *et al.* 2015). Bumblebees are large-bodied, mainly cold-adapted species with broad significance for agricultural production and plant-pollinator networks in the wild (Goulson 2003). Many bumblebee species are experiencing range shifts or declines due to climate change or habitat loss (Goulson *et al.* 2008a; Cameron *et al.* 2011; Cameron & Sadd 2020; Soroye *et al.* 2020). In particular, a general worldwide trend for species to shift to higher elevations and latitudes has been documented, with losses occurring at a faster rate than expansions (Kerr *et al.* 2015; Marshall *et al.* 2020). Arctic and alpine species are

particularly threatened by these trends, and can be considered "canaries in the coal mine"

for detecting early effects (Elsen & Tingley 2015).

In addition to range shifts there is evidence that the morphology of certain populations of

bumblebees has shifted in recent decades. Miller-Struttmann et al. (2015) compared

morphometric measurements of contemporary samples and historical specimens of the

species *B. balteatus* and *B. sylvicola* from the same high-elevation alpine localities in

Colorado. These species historically comprise the majority of bumblebees in these locations

(Miller-Struttmann *et al.* 2015). They reported a significant decrease in tongue length in

both species in a period between 1966 and 2014. The cause of this was inferred to be a

general decrease in floral abundance resulting from warmer temperatures and dryer soils

resulting from climate change. These conditions could favor more generalist foraging

strategies, which several studies indicate are performed more efficiently by bees with

shorter tongues (Heinrich 2004; Goulson & Darvill 2004; Goulson *et al.* 2005, 2008b; Huang

et al. 2015). Continued monitoring of these populations is important to understand the

causes of these morphological shifts and their ecological consequences. It is unknown

whether the trend towards shorter tongues has continued in more recent years.

One explanation for the observed morphological shifts in *B. balteatus* and *B. sylvicola* in

Colorado is that natural selection for shorter tongues has occurred (Miller-Struttmann et al.,

2015). However, it also possible that the observed changes are a plastic response to

environmental change (Merilä & Hendry 2014). It is unknown whether this trait has a

genetic component that could be acted on by natural selection. Uncovering the size of the

genetic contribution to this trait and the number and identity of any genetic loci involved

would be an important step towards understanding the underlying cause of the

morphological changes. The feasibility of genome sequencing on a population scale has

enabled studies of quantitative genetics (Gienapp *et al.* 2017) and genome-wide association

studies (Santure & Garant 2018) in wild populations, allowing these goals to be pursued.

The genetic structure of populations is an important factor for interpretation of genotype-

phenotype correlations and the cause of morphological shifts. We previously presented a

genome assembly of *B. sylvicola* and analyzed genome-wide variation by resequencing 281

samples (Christmas *et al.* 2021). This identified the presence of a previously-undetected

cryptic species, which we gave the preliminary name *Bombus incognitus*. We refer to this

species hereafter simply as "incognitus" to reflect the fact that a formal taxonomic

description is not currently available. This species was identified morphologically as *B.*

sylvicola but forms a distinct genetic cluster. It is therefore possible that changes in tongue

length in the combined populations of these species could be brought about by changes in

the relative abundance of each species, although this has not previously been investigated.

Additional population structure within species could lead to a similar effect, particularly if

genetic structure associated with morphology exists. An understanding of genetic structure

and spatial connectivity between populations in different locations is also needed to predict

how they will evolve under climate change and to define the conservation value of

subpopulations (Pauls *et al.* 2013; Razgour *et al.* 2019).

The degree of gene flow in bumblebee populations is determined by the distances that

reproductive individuals (males and queens) travel in order to mate and establish new nests

(Heinrich 2004; Woodard *et al.* 2015). Genetic methods have been used to investigate

population structure of several species of bumblebees across a variety of habitats in the UK,

continental Europe and continental USA (Darvill *et al.* 2006; Ellis *et al.* 2006; Jha 2015;

Woodard *et al.* 2015; Koch *et al.* 2017; Jackson *et al.* 2018; Ghisbain *et al.* 2020). A general

finding is that populations of common species tend to exhibit very little structure in the

absence of geographical barriers, even at continental scales. However, genetic

differentiation can occur in rare or declining species, or when there are natural barriers

(Woodard *et al.* 2015). Populations restricted to high elevations have been shown to exhibit

higher genetic differentiation (Lozier *et al.* 2011, 2013). However, the degree of

fragmentation of populations of bumblebee species inhabiting high elevations in mountain

ranges is not fully understood.

Genetic methods can also be applied to estimate nest density and foraging distances of

bumblebees, by analyzing where genetically-related nestmates are caught. Such studies

indicate that maximum foraging distances range from less than 100 m to over 10 km (Darvill

et al. 2004; Knight *et al.* 2005; Jha & Kremen 2013; Geib *et al.* 2015; Woodard *et al.* 2015).

These variables show large variation between species and are strongly dependent on

landscape composition. Genome sequencing of multiple individuals has the potential to add

greater resolution to our understanding of population density, structure and dispersal.

In this study we present a high-quality annotated genome assembly of the bumblebee

species *B. balteatus* based on long-read Oxford Nanopore sequencing. We also conducted

whole-genome sequencing of 299 samples of this species sampled from seven high-altitude

localities spread across 150 km in Colorado. These localities have an average separation of

54 km between each other, and consist of alpine tundra separated by forest. We analyze

these data together with the previously-published population genomic dataset of 284

samples *B. sylvicola* and "incognitus" from the same localities (Christmas *et al.* 2021). We

use these data to assess whole-genome variation and its connection to geographical and

morphological variation in these populations, which were previously studied by (Miller-

Struttmann *et al.* 2015). We then analyzed morphological variation in tongue length and

body size in context of previous studies (Miller-Struttmann *et al.* 2015) to determine

whether the trend towards decreasing tongue length is ongoing. We also performed a

genome-wide analysis of the genetic basis of morphological variation using both genome-

wide association studies (GWAS) and genome-wide complex trait analysis (GCTA) (Yang *et*

al. 2011). The results are informative regarding the degree of connectivity between

bumblebee populations, the genetic basis of morphological variation, and the mechanisms

of morphological evolution.

Our study species have recently been subject to taxonomic revision. The species we refer to

here as *B. balteatus* was previously described as occurring in both North America and

Eurasia, but a recent study classified these geographically separated populations as two

different species: *B. balteatus* in Eurasia and *Bombus kirbiellus* in North America (Williams *et*

al. 2019). The species we refer to here as *B. sylvicola* was previously split into a North

American form (*B. sylvicola*) and a Eurasian form (*Bombus lapponicus*). However, a recent

analysis synonymized these two species as *B. lapponicus* (Martinet *et al.* 2019). We continue

to use the previous names here to avoid confusion with comparison to previous ecological

and evolutionary genetic studies.

MATERIALS AND METHODS

Sample collection

We collected samples identified as *B. balteatus* and *B. sylvicola* (Koch *et al.* 2014) during the summer of 2017. Bumblebees were collected on seven different mountains in Colorado, USA, with an average distance of 54 km between mountains (greatest distance: 134 km, Quail Mountain – Niwot Ridge; shortest distance: 10km, Mount Democrat – Pennsylvania Mountain). The existence of "incognitus" was unknown to us when the bees were sampled but a large fraction of bees identified as *B. sylvicola* were subsequently assigned to "incognitus" on the basis of genetic analysis (Christmas *et al.* 2021). Three of these mountains (Mount Evans (39.583883, -105.63845), Niwot Ridge (40.05945, -105.616667), and Pennsylvania Mountain (39.263383, -106.142733)) were also sampled previously by (Miller-Struttmann *et al.* 2015). Four additional mountains were sampled here (Boreas Mountain (39.419375, -105.95441), Mount Democrat (39.33721, -106.12581), Horseshoe Mountain (39.204605, -106.181085) and Quail Mountain (39.018158, -106.404877)) but were not sampled in the previous study. Each of the mountain locations comprised several neighboring sampling sites separated by short distances (mean = 1.1 km) and elevations (range: 3,473 – 4,012 m.a.s.l.) (full details in supplementary tables S1, S2).

We collected samples of foraging worker bees from each site using sweeping hand nets. We kept samples in Falcon tubes in cool boxes for transport. Species assignment was performed by observing standard morphological characters (Koch *et al.* 2014). We dissected each sample after placing it at -20°C for approximately 10 minutes. We stored thoraces in 95% ethanol prior to DNA extraction whereas the heads were stored for morphometric analysis.

Morphological measurements

We measured intertegular distance (ID) and tongue length on all of the samples of *B. sylvicola,* "incognitus" and *B. balteatus* using scaled photographs of individual bee heads processed by ImageJ (Kearns & Inouye 1993; Schneider *et al.* 2012). Following netting, bees were anesthetized by chilling them on ice or in a freezer (depending on field conditions). Once anesthetized, the bees were removed from the vial, gently pressed prone to the table with the thorax parallel to the camera lens using flat forceps, and photographed with a metric ruler for scaling purposes. The distance between the tegula was measured as a straight line between the edge of one tegula to the other tegula, running as near to

perpendicular to the tegula as possible. The head was then immediately removed from the

body and placed into an individual labeled vial and kept frozen until further characterization

in the lab. Following the protocol used in Miller-Struttmann *et al.* (2015), the prementum

and glossa were dissected from the head while the tissue was flexible, straightened and

aligned as possible without damaging the tissue, and mounted on acid-free paper. The

samples were then photographed with a ruler in the field of view under a microscope, and

the traits measured (i.e., prementum length and glossa length) via ImageJ (Schneider *et al.*

2012). We visually characterized the diagnostic traits visible on each head, such as the shape

of the malar space, the pile color between and antenna and above the ocelli, the size, color

and location of the ocelli relative to the supraorbital line, and the presence of black pile on

the scutellum. None of the morphological characters we measured could be used to

distinguish between "incognitus" and *B. sylvicola*.

Genome sequencing and assembly

We used the same sequencing strategy to produce high-quality genome assemblies for both

B. sylvicola and *B. balteatus*. The *B. sylvicola* assembly was presented previously (Christmas

et al. 2021). This involved combining data from Oxford Nanopore (ONT) and 10x Genomics

Chromium technologies. For both species, we used DNA extracted from male bees sampled

on Niwot Ridge and stored in 95% ethanol prior to DNA extraction. DNA from a single bee

was used for each assembly. We performed DNA extraction using a salt-isopropanol

extraction followed by size selection using magnetic bead purification to remove fragments

< 1 kbp. ONT sequencing was performed using the MinION instrument. For *B. sylvicola*, we

used two R9.4 flowcells and the RAD004 kit starting with 3-400 ng DNA per run. This

resulted in a total of 9.4 Gbp of sequence data, in 2.5 million reads with a mean length of

3.7 kbp (Christmas *et al.* 2021). For *B. balteatus*, we used 1μg DNA in a modified LSK-108

one-pot protocol. 1.75 μl NEB FFPE buffer, 1.75 μl NEB Ultra-II End Prep buffer and 1 μl FFPE

repair mix was added to 24 μl DNA and incubated at 20°C for 30 minutes. 2 μl End Prep

enzyme mix was then added for blunting, phosphorylation and A-tailing (20 min. 20°C, 10

min. 65°C, 10 min. 70°C). AMX (20 μl), ligation enhancer (1 μl) and Ultra-II ligase (40 μl) was

added for ligation at room temperature for 30 minutes. Clean-up and flowcell loading was

done following the standard ONT LSK-108 protocol and the library was sequenced on one

R9.4 flowcell. This resulted in 23.3 Gpb of sequence data, in 3.9 million reads with a mean

length of 5.9 kbp.

We assembled the ONT reads using the following process: We first used downpore with

default parameters for adaptor trimming and splitting chimeric reads. We then assembled

the trimmed reads using wtdbg2 (Ruan & Li 2020) on default settings. We then ran two

rounds of the consensus module Racon and contig improvements using medaka v.0.4,

removing contigs of length < 20 kbp. We used Illumina short-read data from the population

sequencing to perform two rounds of Pilon to improve sequence accuracy, particularly

around indels.

We also sequenced the same sample of each species using 10x Genomics Chromium. A 10x

GEM library was constructed according to the manufacturer's protocols. We quantitated

each library using qPCR and sequenced them together on a single lane of HiSeq 2500 using

the HiSeq Rapid SBS sequencing kit version 2, resulting in 150 bp paired-end sequences. The

linked reads were mapped to the assembly using Longranger v.2.1.4. We then ran Tigmint

v1.1.2 to correct errors in the assembly detected by the read mappings. We removed

contigs containing multiple mitochondrial genes identified using BLAST+ v2.9.0. We also

removed all contigs shorter than 10 kbp.

Genome analysis

BUSCO v3.0.2b (Simão *et al.* 2015) was used to assess the assembly completeness using the

hymenoptera_odb9 lineage set and species *B. impatiens*. We ran RepeatMasker v.4.1.0 on

each genome assembly to characterize genome repeat content, including interspersed

repeats and low complexity sequences. It has been shown that bumblebees have a stable 18

chromosome karyotype across all subgenera (Sun *et al.* 2020). We therefore performed

whole-genome synteny alignments between the *B. terrestris* chromosome-level genome

assembly (Sadd *et al.* 2015) (downloaded from NCBI; BioProject PRJNA45869) and our

assemblies using Satsuma v.3 (Grabherr *et al.* 2010) to arrange the contigs from our

assemblies onto 18 pseudochromosomes.

RNAseq and Annotation

We used the Nextflow pipelines available at https://github.com/NBISweden/pipelines-

nextflow for genome annotation. The pipelines, in turn, depend on two additional

annotation-specific toolkits, AGAT and GAAS, available

at https://github.com/NBISweden/AGAT and https://github.com/NBISweden/GAAS respecti

vely. Both repositories contain extensive documentation and installation instructions. As

reference proteins database, we used the SwissProt/UniProt (561,356 curated proteins,

downloaded 2019-11). The parameter files stated below are available at

https://github.com/matt-webster-lab/alpine_bumblebees.

We performed RNAseq on a single lane of Illumina HiSeq 2500 using samples from four body

parts from a single sample of *B. balteatus* (abdomen, head, legs and thorax), which

generated 42 Gbp of reads. We performed quality control of the data using FastQC and read

trimming using trimmomatic (v. 0.39, trimmomatic.params). We next generated a guided

assembly with Hisat2 and Stringtie using the AnnotationPreprocessing.nf (parameters as in

AnnotationPreprocessing_params.conf) TranscriptAssembly.nf Nextflow pipeline with

parameters as in the TranscriptAssembly_params.config file. In addition to the guided

assembly, we performed a de novo assembly for reads from each of the four tissues using

Trinity (v. 2.0.4, --seqType fq --SS_lib_type RF).

Next, we constructed a species-specific repeat library using RepeatModeler package (v.

open-1.0.8; -engine ncbi -pa 35). Since the de novo identified repeats may still include parts

of protein-coding genes, we perform an additional filtering step and remove from the

database the repeats that can be found in a comprehensive set of known proteins. We

remove all repeats that match to a pre-computed set non-transposable proteins from our

protein reference database. We used custom script (filter_repeats.bash) for filtering. To

identify the location of repeats from this library in the genomes, we used RepeatMasker

(open-4.0.9).

We used the MAKER package to compute gene builds. An evidence-guided gene build was

computed by MAKER (-fix-nucleotides) constructing gene models from these reference

proteins and from the transcript sequences aligned to the genome assembly (parameters in

evidence_build.maker_opts.ctl). We also used prepared an ab initio build using the

AbInitioTraining.nf pipeline and AbInitioTraining_params.config as the parameters file.

Both the evidence-guided and the ab initio builds were than merged using MAKER

(hybrid_build_maker_opts.ctl). Finally, we performed functional annotation using the

FunctionalAnnotation.nf pipeline with FunctionalAnnotation_params.config as the

parameter file.

Population sequencing, read mapping and variant calling

We extracted DNA from the thoraces of 299 worker bees of *B. balteatus* collected from

across the seven sampling sites using the Qiagen Blood and Tissue kit. We prepared dual-

indexed libraries using the Nextera Flex kit and performed sequencing on an Illumina HiSeq

X to produce 2 x 150 bp reads, using an average of 36 samples per lane. We mapped reads

to the *B. balteatus* reference using the mem algorithm in BWA (Li & Durbin 2009). We

performed sorting and indexing of the resultant bam files using samtools (Li *et al.* 2009) and

marked duplicate reads using Picard. We used the Genome Analysis Toolkit (GATK) to call

variants (McKenna *et al.* 2010). We first ran HaplotypeCaller using default parameters on

the bam file of each sample to generate a gVCF file for each sample. We then used

GenomicsDBImport and GenotypeGVCFs with default parameters to call variants for all *B.*

balteatus samples together. We applied a set of hard filters using the VariantFiltration tool

to filter for reliable SNPs using the following thresholds: QD < 2, FS > 60, MQ < 40,

MQRankSum < -12.5, ReadPosRankSum < -8 (see the GATK documentation for full

descriptions of each filter). Only biallelic SNPs were considered for downstream analysis.

Samples from *B. sylvicola* and "incognitus" samples were analyzed in the same way and are

presented in Christmas *et al.* (2021).

Relatedness and population structure

We estimated the pairwise kinship coefficient between all samples within each species using

the method described in (Manichaikul *et al.* 2010) implemented in the --relatedness2 option

of vcftools v0.1.16 (Danecek *et al.* 2011). Worker bees from the same bumblebee colony are

expected to have a relatedness of 0.75 under the assumption of monandry, which is

reasonable for the study species (Estoup *et al.* 1995). The kinship coefficient (φ) predicted

from this relationship is $\frac{3}{8}$ (the probability that two alleles sampled at random from two

individuals are identical by descent). We estimated the lower cutoff for the inference

criteria to infer this level of relatedness using powers of two as described in (Manichaikul *et*

al. 2010), providing an estimate of $\frac{3}{2^{7/2}}$.

We tested for correlations between relatedness and geographic distance for each species

using Mantel tests with the mantel function in the 'ecodist' package in R v4.0.2. Coordinates

for each sample were converted to geographic distances using the R package 'geodist' and

these were compared to the matrix of kinship coefficients described above. We used the

mantel.correlog function to produce mantel correlograms and assess whether correlations

were significant over different geographic distances. We performed these analyses after

excluding sisters from the datasets to ensure any patterns we observe were not driven by

an effect of sampling individuals from the same colony.

We thinned the VCF files to 1 SNP every 10 kbp using the --max-missing 1 and --thin flags in

vcftools v.0.1.16 to ensure no missing data and to more efficiently assess population

structure and relatedness. We calculated p-distance matrices for *B. balteatus* samples and

B. sylvicola and "incognitus" samples combined from the thinned vcf files using VCF2Dis

(https://github.com/BGI-shenzhen/VCF2Dis) and then constructed neighbor-joining trees

using the 'ape' package in R v4.0.2. We converted the thinned vcf files to plink format and

performed principal components analyses using the 'adegenet' package in R v4.0.2. We

performed PCAs on all samples as well as after removing one of each pair of sisters

identified via the kinship analysis to ensure that relatedness did not heavily influence any

clustering. We used admixture v1.3.0 (Alexander *et al.* 2009) to assess the most likely

number of genetic clusters within each species. We ran admixture for K=1 through to 10,

with 10 iterations per K value and used the calculated cross-validation (cv) errors to assess

the most suitable values of K, with low cv-error indicating higher support.

Statistical morphometric analysis

We combined the phenotypic measurements made in this study with those from previous

collections published in (Miller-Struttmann *et al.* 2015). These measurements are ID, glossa

length, prementum length, and total tongue length (equal to glossa length plus prementum

length). This gave us three time periods to compare amongst: '2017' (samples collected for

this study), '2012-2014', and '1966-1980' (both sets published in (Miller-Struttmann *et al.*

2015)). We performed statistical analysis of variation in glossa length, prementum length,

and ID in R v4.0.2. We investigated correlations between all traits for each species in the

2017 set using the 'cor.test' function, which calculates a Pearson's product moment

correlation coefficient.

Within our 2017 dataset, we compared glossa lengths between the three species by fitting

linear models of species against glossa length with ID included as a random effect to

account for body size using the lm function. We then calculated least-squares mean glossa

lengths from this model using the lsmeans function and tested for significant differences

using the Tukey method for p-value adjustment for multiple testing. We employed a similar

approach for comparing glossa lengths between our dataset and historical datasets for each

species. Here we used a linear mixed effect model framework with the 'lmer' function in the

lme4 package in R v4.0.2, where mountain and ID were included as random effects in each

model. Significance of each model was assessed using analysis of variance with the 'anova'

function. We calculated least-squares mean glossa lengths from each model using the

lsmeans function and tested for significant differences using the Tukey method for p-value

adjustment for multiple testing.

We analyzed variation in ID between species and geographic locations for our 2017 dataset

using analysis of variance ('aov' function) and assessed significance of comparisons using

Tukey's honestly significant difference (HSD) post hoc test ('TukeyHSD' function). For

comparisons of ID between years, we fitted linear mixed effect models and included

mountain as a random effect, then compared least-squares means from the models using

the Tukey method for p-value adjustment for multiple testing.

Genome-wide association study

We carried out mixed linear model-based association analyses, implemented in GCTA-

MLMA (Yang *et al.* 2011, 2014), to identify correlations between allele frequencies and trait

variation across the genome for *B. balteatus* and *B. sylvicola* samples separately

("incognitus" samples were left out of this analysis due to too few samples). First, we
created a genetic relationship matrix (GRM) from the genome wide SNPs using the --make-
grm function in GCTA. We removed one sample from each pair of sisters by setting a
relatedness cut-off of 0.67 using the '--grm-cutoff' flag. We ran the analyses for three traits
per species:ID, glossa length, and the residuals from a linear regression of glossa~ID to
control for body size. The GRM was included as a polygenic/random effect. The analysis
assumes phenotypes are normally distributed and so samples were removed in the tails of
the distributions to meet this assumption for all traits except ID in *B. sylvicola*, which was
normally distributed. For *B. balteatus*, this resulted in 279, 291, and 262 samples for the
glossa length, ID, and residuals analyses respectively. For *B. sylvicola* there were 176, 192,
and 152 samples for the glossa length, ID, and residuals analyses respectively. We assessed
divergence of p-values from neutral expectations using QQ-plots and carried out Bonferroni
correction of p-values using the 'p.adjust' function in R v4.0.2 to assess genome-wide
significance.
389
Genome-wide complex trait analysis
We used GCTA (Yang *et al.* 2011) to perform genome-wide complex trait analysis. We
estimated the phenotypic variance explained by the genome-wide SNPs using genome-
based restricted maximum likelihood (GREML). The analysis was performed separately for
each species on measures of ID, glossa length, and glossa length with ID included as a
covariate to control for body size. We ran each analysis twice, once using a GRM based on
all samples and a second time with the same GRM produced above where one from each
pair of sisters was removed. We calculated 95% confidence intervals around each estimate
by multiplying the standard error by 1.96. We used R functions provided in (Wang & Xu
2019) as well as the GCTA-GREML power calculator (Visscher *et al.* 2014) to perform power
analysis and estimate the effect sizes we can expect to detect with our population datasets.
401
RESULTS
Generation of a highly-contiguous genome assembly of *B. balteatus*
We generated a *de novo* genome assembly for the species *B. balteatus* using Oxford
Nanopore (ONT) long-read and 10x Chromium linked-read sequencing and compared this to
the assembly for *B. sylvicola* that we recently published (Christmas *et al.* 2021). Assembly

statistics for BBAL_1.0 (*B. balteatus*) and BSYL_1.0 (*B. sylvicola*) are presented in table 1,

which includes statistics for the assemblies of *B. terrestris* (BTER_1.0) and *B. impatiens*

(BIMP_2.2) for comparison. Total assembly sizes of 250.1 and 252.1 Mbp for *B. balteatus*

and *B. sylvicola* respectively are both above the average assembly size of *Bombus* genomes

assembled to date of 247.9 Mbp (22 genomes, range 229.8 – 282.1 Mbp) (Sadd *et al.* 2015;

Heraghty *et al.* 2020; Sun *et al.* 2020). Contig N50s of 8.60 and 3.02 for BBAL_1.0 and

BSYL_1.0 respectively are the longest and third longest contig N50s of all published *Bombus*

assemblies, demonstrating these to be highly contiguous assemblies. Megabase-scale contig

N50 are typical of long-read assemblies, such as the ones presented here and in (Heraghty

et al. 2020), and are significantly more contiguous than those based on short-read

sequencing technologies such as BTER_1.0 and BIMP_2.2.

High genome completeness and accuracy is also reflected in the BUSCO analysis, with scores

of 99.0% (98.7% single copy, 0.3% duplicated, 0.5% fragmented, 0.5% missing) and 98.2%

(97.9% single copy, 0.3% duplicated, 0.4% fragmented, 1.4% missing) reported for BBAL_1.0

and BSYL_1.0 respectively. Whole genome alignments to the *B. terrestris* assembly (Sadd *et*

al. 2015) resulted in 93.9% and 91.1% of the *B. balteatus* and *B. sylvicola* genomes being

placed on 18 pseudochromosomes. Our annotation pipeline annotated 11,711 and 11,585

genes in BBAL_1.0 and BSYL_1.0 respectively. These are highly comparable to the 11,874

genes in the *B. terrestris* gene set (v. 1.0), but lower than the 12,728 genes in the *B.*

impatiens gene set (v. 2.1). Gene annotation of the *Bombus* assemblies presented in

(Heraghty *et al.* 2020; Sun *et al.* 2020) also resulted in a greater number of genes annotated,

ranging from 13,325 to 16,970 genes. BUSCO analyses of the annotated genes in our

assemblies do suggest that we are missing some genes from our annotations (complete

BUSCO genes in BBAL_01 and BSYL_01 respectively: 86.6% and 86.9%). Repeat content of

our assemblies is comparable to that of other *Bombus* assemblies, with 17.11% of BBAL_1.0

and 17.85% of BSYL_1.0 identified as repetitive by RepeatMasker (full details of repeat

classes in supplementary table S3). These proportions are comparable to the repeat content

inferred in the *B. terrestris* and *B. impatiens* genomes.

Population resequencing indicates three species clusters, with little population structure

within species

The locations of sample collections, which comprise 580 samples from across seven

localities in the Rocky Mountains, CO, USA, are presented in Fig. 1 (full details in table S1,

S2). Illumina sequencing resulted in an average genome coverage of 13.1x per sample, with

4.5 and 3.5 million SNPs identified in *B. balteatus* and *B. sylvicola* samples respectively

(supplementary table S4). These samples were initially identified morphologically as *B.*

balteatus and *B. sylvicola*, and those identified as *B. sylvicola* have been presented

previously (Christmas *et al.* 2021). Neighbor-joining trees produced from samples of both

groups of samples indicate that the *B. balteatus* samples form a single cluster whereas the

B. sylvicola samples form two clusters (Fig 1). By comparison with an additional dataset

(Martinet *et al.* 2019) we previously inferred the largest cluster to be comprised of *B.*

sylvicola samples whereas the second cluster is comprised of the cryptic species

"incognitus" (Christmas *et al.* 2021). Among samples originally designated as *B. sylvicola,*

24% (n=67) are identified in the separate cluster of "incognitus" whereas the remaining 76%

(n=214) are *B. sylvicola. Bombus balteatus* (n=299) was the most commonly observed

species across all localities. The proportions vary between localities (table S1). "Incognitus"

samples were collected on six out of the seven mountains, whereas *B. balteatus* and *B.*

sylvicola were found on all seven.

We calculated a pairwise kinship coefficient between samples to infer degrees of

relatedness (Manichaikul *et al.* 2010; Danecek *et al.* 2011). We inferred a set of samples that

were likely to be nestmates (sisters) based on them having the same maternal and paternal

parents, which is likely to be the case for all workers from the same colony in these species,

under the realistic assumption that each colony is headed by a single monandrous female

queen (Estoup *et al.* 1995). Those identified as sisters from the same single-mated queen

from the same colony were always collected foraging on the same mountain and at the

same or immediately neighboring site (table S1). This suggests that foraging bees were

generally found in the immediate vicinity of their nests. Among the 299 *B. balteatus*

samples, we identified 43 samples from 17 colonies that had another nestmate in the

dataset, indicating that we sampled a total of 273 colonies. For the 214 *B. sylvicola*, we

identified 14 samples from 7 colonies that met this criterion, indicating a total of 207

colonies. For 67 "incognitus", we identified two samples from the same colony, indicating

that we sampled 66 colonies.

We investigated the presence of population genetic structure within each species using

several methods. Neighbor-joining trees (Fig. 1) do not reveal the presence of any

substructure related to geography for *B. balteatus* or "incognitus". For *B. sylvicola*, all of the

samples from Quail Mountain (the furthermost southwestern site) cluster together, but do

not separate from the rest on the tree. Principal component analysis further reflected the

extremely low level of geographical structure across all species (Fig. 2). One slight exception

to this trend is the Quail Mountain population of *B. sylvicola* which, as with the neighbor-

joining tree, shows a slight degree of clustering away from the other samples at the positive

extreme of PC1 (Fig. 2B), but they still overlap with other samples and PC1 explains only

1.17% of the variance in the data. Outlier clusters seen in the PCA plots of *B. balteatus* and

"incognitus" (Fig. 2A,C, circled) are composed of highly related individuals as shown in the

kinship analysis, suggesting that the effect of kinship overrides any signal of genetic

structure. After removing one of each pair of sisters, nearly all *B. balteatus* samples cluster

extremely tightly, with PC1 only explaining 0.8% of the variance (Fig. S1A). A similar pattern

is observed for *B. sylvicola*, with PC1 explaining 1.2% of the variance (Fig. S1B). For

"incognitus", two pairs of highly related individuals are separated from the main cluster on

PC1, but again only a small amount of the genetic variance is captured by PC1 (2.6%),

suggesting they are still highly similar (Fig. S1C).

We assessed connectivity between colonies on different mountains by analyzing the

association between kinship and geography using Mantel tests after removing sisters from

the datasets (Fig. 3). We found no significant relationship between geographic distance and

relatedness for *B. balteatus* (Mantel R = -0.03 (95% ci: -0.04 - -0.02), p value = 0.11; Fig.

3A,D). There were small but significant negative relationships for *B. sylvicola* (Mantel R = -

0.06 (95% c.i. -0.07 - -0.04), p value = 0.0001; Fig. 3B,E) and "incognitus" (Mantel R = -0.04

(95% ci: -0.06 - -0.03), p value = 0.029; Fig. 3C,F), demonstrating greater relatedness at

closer locations. In these species, Mantel correlograms indicate significant associations

between kinship and geographical distance only at shorter distances (from the same or

neighboring mountains), which are no longer significant at greater distances (Fig. 3E-F)).

These results indicate that the limited population structure is mainly driven by the

occurrence of a few highly-related samples from the same localities in the dataset.

Decrease in tongue length is ongoing in all species

We made morphological measurements for each sample (table S2). Intertegular distance (ID) is considered a proxy for body size (Cane 1987). We also measured the length of two parts of the tongue: glossa and prementum. These two tongue measures are correlated in both species (*B. balteatus* Pearson's r = 0.45, p < 0.001; *B. sylvicola* Pearson's r = 0.38, p < 0.001; Figs. S2,S3). Only glossa is considered further, as it is the tongue part with most relevance to feeding (Cariveau *et al.* 2016) and was measured by (Miller-Struttmann *et al.* 2015). Glossa is also significantly correlated with ID (*B. balteatus* Pearson's r = 0.41, p < 0.001; *B. sylvicola* Pearson's r = 0.26, p < 0.001) (Figs. S2,3).

A comparison of phenotypes between species shows that glossa length is substantially longer in *B. balteatus*, consistent with previous knowledge (ls-means = 4.72 mm, 3.34 mm, and 3.40 mm for *B. balteatus*, *B. sylvicola* and "incognitus" respectively; Tukey test, t ratio = 12.25 and 21.17, p < 0.001 for *B. balteatus* compared to *B. sylvicola* and "incognitus"*;* Fig. 4A). Average glossa length is extremely similar in *B. sylvicola* and "incognitus" (Tukey test, t ratio =0.50, p = 0.87), suggesting that this character does not distinguish this newly-discovered species from *B. sylvicola*. *Bombus balteatus* bees are significantly larger bees as indicated by a greater ID, and *B. sylvicola* is significantly larger than "incognitus" (mean ID = 4.00 mm, 3.83 mm, 3.58 mm for *B. balteatus*, *B. sylvicola* and "incognitus" respectively; ANOVA, F=28.75, p < 0.001, Tukey post-hoc test revealed differences are significant among all species at p < 0.001; Fig. S4A).

We next compared the distribution of glossa lengths in our samples with historical data. We combined *B. sylvicola* and "incognitus" for this analysis because they do not differ in glossa length on average and they were not distinguished in previous studies. We find an overall trend of decreasing glossa length in both *B. sylvicola*-incognitus and *B. balteatus* over time (Fig. 4B,C). This was reported previously between the collections from 1966-1980 and from 2012-2014 for both species (Miller-Struttmann *et al.* 2015). Here we show that this trend is continuing, with further reductions in glossa length between the 2012-2014 collections and our collections from 2017. These reductions are significant for both *B. sylvicola*-incognitus (ls-means: 2012-2014 = 3.51 mm, 2017 = 3.21 mm; Tukey test, t-ratio = 3.198, p = 0.004)

and *B. balteatus* (ls-means: 2012-2014 = 5.24 mm, 2017 = 4.84 mm; Tukey test, t-ratio =

2.79, p = 0.015). In contrast, there is no significant trend shown by ID (Fig. S4B,C), although

for *B. sylvicola*-incognitus, ID is significantly larger in both the 1966-80 and 2017 samples

compared to the 2012-2014 samples (ANOVA, F = 15.28, p < 0.001. Tukey post-hoc test, p <

0.001).

For samples collected in 2017, we compared the distribution of glossa lengths for each

species among the seven mountains where they were collected (Fig. S5). There is an overall

trend for glossa lengths to be longer in bees from Mount Evans than on other mountains in

all species. The distributions for *B. balteatus* and *B. sylvicola* show significant differences,

with samples from Mount Evans having significantly longer glossae than on Mount

Democrat, Horseshoe Mountain, and Pennsylvania Mountain in *B. balteatus* (ls-mean glossa

length on Mt. Evans = 5.34 mm, other mountains 4.51 – 4.73 mm; Tukey p < 0.01 in each

case; Fig. S5A) and significantly longer glossae than on all other mountains except Niwot

Ridge in *B. sylvicola* (ls-mean glossa length on Mt. Evans = 3.77 mm, other mountains 3.02 –

3.24 mm; Tukey p < 0.01 in each case; Fig. S5B). None of the trends for "incognitus" were

significant (Fig. S5C). Comparisons of ID between mountains revealed that *B. balteatus* bees

sampled on Niwot Ridge are significantly larger than those on Boreas Mountain, Mount

Democrat, Horseshoe Mountain, and Quail Mountain (Niwot Ridge mean ID = 4.25 mm,

other mountains 3.84 - 4.00 mm, ANOVA, F = 4.09, p < 0.001, Tukey post-hoc test p < 0.05 in

four comparisons; Fig. S6A). We did not identify any significant differences in ID of bees

across mountains for *B. sylvicola* (ANOVA, F = 1.83, p = 0.10) or "incognitus" (ANOVA, F =

1.62, p = 0.17; Fig. S6B,C).

Genome-wide association study identifies a SNP associated with a proxy for body size

We used a linear mixed-model analysis to search for correlations between allele frequencies

and quantitative trait variation in *B. sylvicola* and *B. balteatus.* We used the traits glossa

length, ID and glossa length corrected for ID (see Methods). One from each pair of sisters

identified in the kinship analysis was removed from all analyses to reduce the level of

relatedness in the sample set. We detected a genome-wide significant association between

ID and a SNP in intron 9 of the *CTRB1* (Chymotrypsinogen B1) gene on chromosome 10 in *B.*

sylvicola (p < 0.001 after Bonferroni correction; Fig. 5). The SNP is found at an allele

frequency of 0.14 across all populations, with 41 individuals heterozygous and nine

homozygous for the allele. The allele is present on all mountains, ranging in allele frequency

from 0.06 (Quail Mountain) to 0.24 (Niwot Ridge). The *CTRB1* gene is involved in production

of chymotrypsin, a proteolytic digestive enzyme (Giebel *et al.* 1971; Grogan & Hunt 1979;

Rawlings & Barrett 1994; Burgess *et al.* 1996; Matsuoka *et al.* 2015; Lazarević & Janković-

Tomanić 2015). Intron 9 is a large 18 Kbp intron and the significant SNP is located 2 Kbp

upstream of exon 10. The GWAS significance suggests it is of likely functional importance

and its location is suggestive of a regulatory element, such as an intron splice enhancer or

silencer.

Two further SNPs, located 4 bp apart on chromosome 11, also significantly associated with

ID (p = 0.039 after Bonferroni correction). However, they are not located in or close to any

genes according to our annotation, so any potential functional significance of these SNPs is

unknown. We did not find any significant associations between allele frequencies and ID in

B. balteatus (Fig. S7). Furthermore, no significant associations were found between allele

frequencies and glossa length, or glossa length corrected for ID in either species (Fig. S7).

This indicates that the trait has a polygenic or mainly environmental component. A power

analysis (Visscher *et al.* 2014) revealed that our datasets have sufficient power (>0.8) to

detect a QTL with a h^2 of ~0.1 but that loci with lower effect sizes would likely be

undetectable.

Limited evidence that morphological variation has a genetic component

We performed GCTA-GREML analyses (Yang *et al.* 2011) to determine the genetic

component of trait variation for *B. sylvicola* and *B. balteatus* (Fig. 6, supplementary table

S5). The "incognitus" samples were not included here due to smaller sample size. When

each species' entire dataset is considered, we detect a heritable component for both glossa

length and ID in both species, where the 95% confidence interval does not overlap zero. For

B. sylvicola, but not *B. balteatus*, this is also the case when ID is introduced as a covariate to

account for body size when considering glossa length. However, when we remove samples

that we infer to be workers from the same colony from the kinship analysis, we only detect

a heritable component with 95% confidence for ID in *B. balteatus* (all others then overlap

with zero). This indicates that a large proportion of the signal of covariance between genetic

and trait variance is due to the presence of sisters from the same colony. This could be a

confounding factor because environmental effects such as level of nutrition and colony

health are likely to covary with colony. However, such covariation could also reflect

genetics. The level of dependency of phenotype on genotype is therefore unclear from this

analysis.

The 95% confidence intervals for the estimate of heritability have a wide span and overlap

zero in many cases. The power to accurately determine heritability depends on several

factors, including sample size, the structure of relatedness in the samples, and the true

value of heritability. When samples are all completely unrelated, it has been estimated

that > 3,000 samples are required to achieve a SE below 0.1 (Visscher *et al.* 2014). For

datasets of unrelated samples of the size we have analyzed here, we estimate a power of

~0.06, or ~6% probability, to detect $h^2 > 0$ for a trait with a SNP-heritability of 0.5 and a type

1 error rate (α) of 0.05. Greater sample size or higher levels of relatedness within the

dataset would therefore increase accuracy of our heritability estimates.

DISCUSSION

In this study, we generated a high-quality reference genome and performed genome

sequencing of a population sample of the bumblebee species *B. balteatus*. Combining this

data with a previous dataset from *B. sylvicola* produced the most extensive population

genomics dataset for bumblebees to date, with 580 samples from these two species and the

newly-discovered "incognitus" collected from high-altitude locations in Colorado, USA.

These data were combined with morphometric measurements from each sample and used

to investigate genetic structure, morphological evolution and the genetic basis of

phenotypic traits in these populations. Our main findings are 1) no evidence for gene flow

between species, including the recently-inferred "incognitus", but lack of population

structure within species, indicating high levels of connectivity among subpopulations. 2) An

ongoing decrease in tongue (glossa) length compared to historical measurements,

previously inferred to reflect the effects of decreasing floral abundance on bumble bee

foraging (Miller-Struttmann *et al.* 2015). 3) An association between body size and variation

at a single SNP located in a gene related to production of a digestive enzyme in *B. sylvicola*.

4) A lack of conclusive evidence for a genetic component to tongue length in both species.

Our analysis is unable to determine the degree of heritability underlying these traits but is

consistent with a highly polygenic genetic component.

This investigation adds to the number of bumblebee species with reference genomes

available. The genomes of the common species *B. impatiens* and *B. terrestris* were

sequenced using short-read technologies (Sadd *et al.* 2015). Genome assemblies are also

available for representatives of all 15 *Bombus* subgenera (Sun *et al.* 2020) based on short-

read sequencing of jumping libraries and scaffolding using Hi-C, resulting in contig N50 of

325 kb. In addition to the two genomes produced here, the genomes of the North American

species *B. terricola*, *B. bifarius*, *B. vancouverensis*, and *B. vosnesenskii* have also been

produced using long-read sequencing (PacBio or ONT) and have megabase-scale contig N50s

(Kent *et al.* 2018; Heraghty *et al.* 2020).

Identification of the previously-unknown species "incognitus" demonstrates the power of

population-scale genome sequencing to discover new cryptic variation (Christmas *et al.*

2021). None of the morphological features that we studied could distinguish between *B.*

sylvicola and "incognitus". *Bombus sylvicola* did however have a significant tendency to be

collected at higher altitudes and have larger body size. However, importantly for this study,

there were no significant differences in glossa or prementum length between the two

species. The wider distribution of "incognitus" is unknown, as is the true composition of

populations described as *B. sylvicola* in other geographical locations. Existing collections of

B. sylvicola from Colorado are likely to be a mixture of both species.

We find no evidence for population structure among samples of any of the of the three

species studied here. This indicates an absence of long-term geographical barriers to gene

flow between the high-elevation localities included here. Furthermore, although we observe

elevated degrees of relatedness among foragers collected on the same mountain, there is

no significant tendency for elevated relatedness of foragers caught on nearby mountains,

indicating a lack of population structure at this scale. Foragers inferred to belong to the

same nest (i.e. with identical parents) were always found in the same or adjacent sites,

consistent with the limited foraging distance (on average < 110m) inferred previously by

Geib et al. (Geib *et al.* 2015) for these species.

Studies of several bumblebee species across continental USA, including *B. bifarius* and *B.*

vosnesenkii, also found in Colorado and across western North America, indicate weak

geographical differentiation even at distances over 1000 km (Lozier *et al.* 2011; Ghisbain *et*

al. 2020). However, the complex topography of mountain ranges likely modulates gene flow

between populations (Lozier *et al.* 2011, 2013). Populations of *B. bifarius* are inferred to be

more fragmented in the southern portions of their ranges, where they occur at higher

elevations (Jackson *et al.* 2018). The species studied here, *B. sylvicola* and *B. balteatus* are

generally restricted to high elevation localities above the tree line separated by forest

(Williams *et al.* 2014). However, our analysis indicates that this complex landscape has not

historically limited mating and dispersal between localities in the same mountain range. This

is consistent with previous studies of these species that found substantial genetic

differentiation between mountain ranges in the Pacific Northwest, but not within the same

mountain range (Koch *et al.* 2017; Whitley 2018).

Morphological change over time has already been documented in the species studied here,

with a significant reduction in tongue length observed since 1966 (Miller-Struttmann *et al.*

2015). During this time period, there has also been a shift in species composition, with

shorter-tongued species becoming more predominant. Previous collections of *B. sylvicola* in

this region are likely a mixture of *B. sylvicola* and "incognitus", as the existence of

"incognitus" was unknown. However, as *B. sylvicola* and "incognitus" do not differ in glossa

length, undetected changes in the relative abundance of these species over time are

unlikely to be responsible for the observed shifts in glossa length observed in samples that

are defined as *B. sylvicola.* This effect could however contribute to the observed variation in

body size over time as *B. sylvicola* have a significantly greater ID on average than

"incognitus" (Fig. S4).

A major finding that we present here is that the temporal shift toward shorter tongue

length is ongoing. In addition to the significant reduction previously observed over a period

of six decades up until 2014, we also observe a significant reduction over a much shorter

time period between 2012-14 and 2017 in both *B. balteatus* and *B. sylvicola*-incognitus.

Mean tongue length reportedly decreased on average by 0.61% annually in a period

between 1966 and 2014 for *B. sylvicola* (Miller-Struttmann *et al.* 2015). According to our

recent measurements, the corresponding decrease in 3-5 subsequent years has been

12.9 %, or 3.2% per year (compared to the 2012-14 average).

Decreases in tongue-length have been previously attributed to an advantage of shorter

tongues for foraging on a wide range of flowers. Such generalist foraging could be favored

due to a general decline of floral resources observed in connection with warmer summers

(Miller-Struttmann *et al.* 2015). While both long- and short-tongued bees are capable of

accessing nectar in plants with short flower tubes, short-tongued bees are more efficient at

foraging from them (Inouye 1980) and are also more efficient at foraging across a diversity

of flower tube depths (Plowright & Plowright 1997; Arbulo *et al.* 2011; Geib & Galen 2012).

Short-tongued bumble bees often forage from a wider suite of plants both in these alpine

bumble bee species (Miller-Struttmann *et al.* 2015) and others (Heinrich 2004; Goulson &

Darvill 2004; Goulson *et al.* 2008b; Huang *et al.* 2015). These considerations indicate that

shorter tongues are likely advantageous for generalist foraging. Although short-tongued

bees are unable to pollinate plants with long flower tubes, they are likely important for

supporting a diverse array of flowering plants.

It is unclear whether the tongue-length changes we observe are the result of natural

selection or phenotypic plasticity. There are now many examples of rapid evolution of traits

driven by natural selection due to climatic events in the wild, such as a shift in beak shape

over a period of a few years in Darwin's finches related to competition for food during

drought (Lamichhaney *et al.* 2016) and genomic shifts related to cold-tolerance in the green

anole lizard related to severe winter storms (Campbell-Staton *et al.* 2017). Human activity

may also promote rapid adaptation by natural selection, as demonstrated by the evolution

of longer beaks in great tits promoted by the use of bird feeders in the UK (Bosse *et al.*

2017). Global climate change is also expected to be a driver of rapid evolution, but it is often

difficult to establish whether trends of phenotypic change result from evolution or

phenotypic plasticity (Merilä & Hendry 2014). Long-term changes in breeding timing in great

tits (Charmantier *et al.* 2008) and body mass in red-billed gulls (Teplitsky *et al.* 2008) that

track temperature increases have been shown to result solely from phenotypic plasticity.

To address the causes of the changes in bumblebee tongue length observed here, we

attempted to uncover the genetic basis of morphological variation using genome-wide

association studies (GWAS) and genome-wide complex trait analysis (GCTA). No significant

associations were observed for tongue length in the genome of any of the species using

GWAS. However, a significant association with body size was identified in *B. sylvicola* at a

single SNP in the *CTRB1* (Chymotrypsinogen B1) gene. The lack of association at neighboring

SNPs could be attributable to high rates of meiotic recombination and low levels of linkage

disequilibrium in bumblebees (Kawakami *et al.* 2019). The *CTRB1* gene produces a precursor

of chymotrypsin, a key digestive enzyme in animals, which is likely important for pollen

digestion in bees (Giebel *et al.* 1971; Grogan & Hunt 1979; Rawlings & Barrett 1994; Burgess

et al. 1996; Matsuoka *et al.* 2015; Lazarević & Janković-Tomanić 2015). The associated SNP

is found in a large intron (18 Kbp) of this gene, 2 Kbp upstream of the nearest exon. We lack

any functional annotation for this SNP but its location and GWAS significance suggests it

may play a role in gene regulation. For example, if located within an intronic splicing

enhancer or silencer, it may result in variation in expression levels of different isoforms of

the CTRB1 enzyme being produced. Such changes could influence body size through an

effect on the efficiency of digestion.

We detected a significant signal of covariance of trait measurements with relatedness for

both tongue length and ID using GCTA when closely-related individuals were included in the

dataset. However, no significant signal of heritability in morphological traits was detected

when the dataset was purged of close relatives. The first result could potentially reflect a

genetic component, but it is not possible to disentangle this component from the shared

environmental effects due to close relatives sharing a nest. This analysis is therefore

underpowered to detect genetic effects due to low sample size and a general lack of related

individuals in the dataset. A possible explanation for our results is that the morphological

traits tongue length and ID have a highly polygenic genetic component, with multiple QTL of

small effect ($h^2 < 0.1$), such that GWAS and GCTA were underpowered to detect QTL and

estimate heritability. The size of the genetic component of variation in both traits is

therefore still unknown in each species.

It therefore remains unclear whether the observed changes in tongue length are caused by

natural selection acting on functional genetic variation, or whether they are the result of

phenotypic plasticity. There is large variation in body size between individuals in bumblebee

nests, indicating substantial phenotypic plasticity in this trait (Peat *et al.* 2005). This

variability could be explained by a correlation between larval position in the nest and size,

which is related to feeding intensity (Couvillon & Dornhaus 2009). There is evidence from

solitary bees of an effect of temperature on body size (Scaven & Rafferty 2013), although

such an effect is likely to be less pronounced in social insects such as bumblebees that

regulate the temperature of their nests. However, it is unclear whether tongue length also

exhibits plasticity, as this trait correlates with body size among nestmates (Peat *et al.* 2005),

indicating that plasticity in tongue length independent of body size is less pronounced.

However, our observation of significant differences in mean tongue length between

mountains could potentially reflect phenotypic plasticity.

In summary, we have shown that decreasing tongue length observed over the last six

decades is still ongoing in populations of the bumblebee species *B. sylvicola* and *B. balteatus*

(and the recently-reported "incognitus") at high-altitudes in Colorado. We have generated

the genomic resources to study these species on the genetic level in the form of highly-

contiguous genome assemblies and comprehensively characterized genome variation. Our

results indicate that the complex topology of the mountain landscape in Colorado does not

contain significant barriers to mating and dispersal of bumblebees between within this area.

We are unable to conclude whether the observed morphological shifts over time are the

result of genetic adaptation or phenotypic plasticity. Alpine ecosystems are particularly

sensitive to climate change and may suffer its effects before other geographical regions

(Elsen & Tingley 2015). Continued monitoring of changes in morphology in these and other

populations in these regions could be highly valuable for understanding how pollinators

respond to the effects of a warming climate, the degree to which rapid evolution by natural

selection or phenotypic plasticity are possible, and the effects of the changes on these

ecosystems.

DATA AND CODE AVAILABILITY

All sequence data presented here are available at NCBI under BioProject IDs PRJNA646847

and PRJNA704506. Scripts and annotation parameters used in the study are available at

GitHub (https://github.com/matt-webster-lab/alpine_bumblebees). VCF files containing

genotypic variation are available at the Dryad Digital Repository

(https://doi.org/10.5061/dryad.mgqnk990p).

ACKNOWLEDGEMENTS

This research was funded by Swedish Research Council Formas grant 2016-00535 and a

SciLifeLab National Biodiversity Project NP00046 to MTW. MK is financially supported by the

Knut and Alice Wallenberg Foundation as part of the National Bioinformatics Infrastructure

Sweden at SciLifeLab. The authors acknowledge support from the National Genomics

Infrastructure in Stockholm and Uppsala funded by Science for Life Laboratory, the Knut and

Alice Wallenberg Foundation and the Swedish Research Council. The computations and data

handling were enabled by resources provided by the Swedish National Infrastructure for

Computing (SNIC) at UPPMAX partially funded by the Swedish Research Council through

grant agreement no. 2018-05973. We thank the U.S. Forest Service, Dr. Rosemond Greinetz

and the Mountain Area Land Trust for providing access to field sites. We thank Jacqueline

Staab and Michael Osbourn who aided in specimen collection, identification and

preservation and Lisa Danback, Kate Schleicher, Ying Thao, Anna Grobelny and Breana Cook

who carried out phenotypic measurements.

REFERENCES

Alexander DH, Novembre J, Lange K (2009) Fast model-based estimation of ancestry in
unrelated individuals. *Genome Research*, **19**, 1655–1664.

Arbulo N, Santos E, Salvarrey S, Invernizzi C (2011) Proboscis Length and Resource
Utilization in Two Uruguayan Bumblebees: Bombus atratus Franklin and Bombus
bellicosus Smith (Hymenoptera: Apidae). *Neotropical Entomology*, **40**, 72–77.

Bosse M, Spurgin LG, Laine VN, Cole EF, Firth JA, Gienapp P, … Slate J (2017) Recent natural
selection causes adaptive evolution of an avian polygenic trait. *Science*, **358**, 365–
368.

Burgess EPJ, Malone LA, Christeller JT (1996) Effects of two proteinase inhibitors on the
digestive enzymes and survival of honey bees (Apis mellifera). *Journal of Insect
Physiology*, **42**, 823–828.

Cameron SA, Lozier JD, Strange JP, Koch JB, Cordes N, Solter LF, Griswold TL (2011) Patterns
of widespread decline in North American bumble bees. *Proceedings of the National
Academy of Sciences*, **108**, 662–667.

Cameron SA, Sadd BM (2020) Global Trends in Bumble Bee Health. *Annual Review of Entomology*, **65**, 209–232.

Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards SV (2017) Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole lizard. *Science*, **357**, 495–498.

Cane JH (1987) Estimation of Bee Size Using Intertegular Span (Apoidea). *Journal of the Kansas Entomological Society*, **60**, 145–147.

Cariveau DP, Nayak GK, Bartomeus I, Zientek J, Ascher JS, Gibbs J, Winfree R (2016) The Allometry of Bee Proboscis Length and Its Uses in Ecology. *PLOS ONE*, **11**, e0151482.

Charmantier A, McCleery RH, Cole LR, Perrins C, Kruuk LEB, Sheldon BC (2008) Adaptive Phenotypic Plasticity in Response to Climate Change in a Wild Bird Population. *Science*, **320**, 800–803.

Christmas MJ, Jones JC, Olsson A, Wallerman O, Bunikis I, Kierczak M, … Webster MT (2021) Genetic Barriers to Historical Gene Flow between Cryptic Species of Alpine Bumblebees Revealed by Comparative Population Genomics. *Molecular Biology and Evolution*.

Couvillon MJ, Dornhaus A (2009) Location, location, location: larvae position inside the nest is correlated with adult body size in worker bumble-bees (Bombus impatiens). *Proceedings of the Royal Society B: Biological Sciences*, **276**, 2411–2418.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, … Durbin R (2011) The variant call format and VCFtools. *Bioinformatics*, **27**, 2156–2158.

Darvill B, Ellis JS, Lye GC, Goulson D (2006) Population structure and inbreeding in a rare and declining bumblebee, Bombus muscorum (Hymenoptera: Apidae). *Molecular Ecology*, **15**, 601–611.

Darvill B, Knight ME, Goulson D (2004) Use of Genetic Markers to Quantify Bumblebee Foraging Range and Nest Density. *Oikos*, **107**, 471–478.

Ellis JS, Knight ME, Darvill B, Goulson D (2006) Extremely low effective population sizes, genetic structuring and reduced genetic diversity in a threatened bumblebee species, Bombus sylvarum (Hymenoptera: Apidae). *Molecular Ecology*, **15**, 4375–4386.

Elsen PR, Tingley MW (2015) Global mountain topography and the fate of montane species under climate change. *Nature Climate Change*, **5**, 772–776.

Estoup A, Scholl A, Pouvreau A, Solignac M (1995) Monoandry and polyandry in bumble bees (Hymenoptera; Bombinae) as evidenced by highly variable microsatellites. *Molecular Ecology*, **4**, 89–93.

Geib JC, Galen C (2012) Tracing impacts of partner abundance in facultative pollination mutualisms: from individuals to populations. *Ecology*, **93**, 1581–1592.

Geib JC, Strange JP, Galen C (2015) Bumble bee nest abundance, foraging distance, and host-plant reproduction: implications for management and conservation. *Ecological Applications*, **25**, 768–778.

Ghisbain G, Lozier JD, Rahman SR, Ezray BD, Tian L, Ulmer JM, … Hines HM (2020) Substantial genetic divergence and lack of recent gene flow support cryptic speciation in a colour polymorphic bumble bee (Bombus bifarius) species complex. *Systematic Entomology*, **45**, 635–652.

Giebel W, Zwilling R, Pfleiderer G (1971) The evolution of endopeptidases—XII. The proteolytic enzymes of the honeybee (Apis Mellifica L.): Purification and characterization of endopeptidases in the midgut of adult workers and comparative

studies on the endopeptidase-pattern of different castes and on different ontogenetic stages. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, **38**, 197–210.

Gienapp P, Fior S, Guillaume F, Lasky JR, Sork VL, Csilléry K (2017) Genomic Quantitative Genetics to Study Evolution in the Wild. *Trends in Ecology & Evolution*, **32**, 897–908.

Goulson D (2003) *Bumblebees: Their Behaviour and Ecology*. Oxford University Press.

Goulson D, Darvill B (2004) Niche overlap and diet breadth in bumblebees; are rare species more specialized in their choice of flowers? *Apidologie*, **35**, 55–63.

Goulson D, Hanley ME, Darvill B, Ellis JS, Knight ME (2005) Causes of rarity in bumblebees. *Biological Conservation*, **122**, 1–8.

Goulson D, Lye GC, Darvill B (2008a) Decline and conservation of bumble bees. *Annual Review of Entomology*, **53**, 191–208.

Goulson D, Lye GC, Darvill B (2008b) Diet breadth, coexistence and rarity in bumblebees. *Biodiversity and Conservation*, **17**, 3269–3288.

Goulson D, Nicholls E, Botías C, Rotheray EL (2015) Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science (New York, N.Y.)*, **347**, 1255957.

Grabherr MG, Russell P, Meyer M, Mauceli E, Alföldi J, Di Palma F, Lindblad-Toh K (2010) Genome-wide synteny through highly sensitive sequence alignment: Satsuma. *Bioinformatics*, **26**, 1145–1151.

Grogan DE, Hunt JH (1979) Pollen proteases: Their potential role in insect digestion. *Insect Biochemistry*, **9**, 309–313.

Heinrich B (2004) *Bumblebee Economics*. Harvard University Press, Cambridge.

Heraghty SD, Sutton JM, Pimsler ML, Fierst JL, Strange JP, Lozier JD (2020) De Novo Genome Assemblies for Three North American Bumble Bee Species: Bombus bifarius, Bombus vancouverensis, and Bombus vosnesenskii. *G3: Genes, Genomes, Genetics*, **10**, 2585–2592.

Huang J, An J, Wu J, Williams PH (2015) Extreme Food-Plant Specialisation in Megabombus Bumblebees as a Product of Long Tongues Combined with Short Nesting Seasons. *PLOS ONE*, **10**, e0132358.

Inouye DW (1980) The effect of proboscis and corolla tube lengths on patterns and rates of flower visitation by bumblebees. *Oecologia*, **45**, 197–201.

Jackson JM, Pimsler ML, Oyen KJ, Koch-Uhuad JB, Herndon JD, Strange JP, … Lozier JD (2018) Distance, elevation and environment as drivers of diversity and divergence in bumble bees across latitude and altitude. *Molecular Ecology*, **27**, 2926–2942.

Jha S (2015) Contemporary human-altered landscapes and oceanic barriers reduce bumble bee gene flow. *Molecular Ecology*, **24**, 993–1006.

Jha S, Kremen C (2013) Resource diversity and landscape-level homogeneity drive native bee foraging. *Proceedings of the National Academy of Sciences*, **110**, 555–558.

Kawakami T, Wallberg A, Olsson A, Wintermantel D, Miranda JR de, Allsopp M, … Webster MT (2019) Substantial Heritable Variation in Recombination Rate on Multiple Scales in Honeybees and Bumblebees. *Genetics*, genetics.302008.2019.

Kearns CA, Inouye DW (1993) *Techniques for Pollination Biologists*. University Press of Colorado, Boulder.

Kent CF, Dey A, Patel H, Tsvetkov N, Tiwari T, MacPhail VJ, … Zayed A (2018) Conservation Genomics of the Declining North American Bumblebee Bombus terricola Reveals Inbreeding and Selection on Immune Genes. *Frontiers in Genetics*, **9**.

919   Kerr JT, Pindar A, Galpern P, Packer L, Potts SG, Roberts SM, … Pantoja A (2015) CLIMATE
920   CHANGE. Climate change impacts on bumblebees converge across continents.
921   *Science (New York, N.Y.)*, **349**, 177–180.

922   Knight ME, Martin AP, Bishop S, Osborne JL, Hale RJ, Sanderson RA, Goulson D (2005) An
923   interspecific comparison of foraging range and nest density of four bumblebee
924   (Bombus) species. *Molecular Ecology*, **14**, 1811–1820.

925   Koch JB, Looney C, Sheppard WS, Strange JP (2017) Patterns of population genetic structure
926   and diversity across bumble bee communities in the Pacific Northwest. *Conservation*
927   *Genetics*, **18**, 507–520.

928   Koch J, Strange J, Williams P (2014) *Bumble Bees of the Western United States*. U.S. Forest
929   Service.

930   Lamichhaney S, Han F, Berglund J, Wang C, Almen MS, Webster MT, … Andersson L (2016) A
931   beak size locus in Darwins finches facilitated character displacement during a
932   drought. *Science*, **352**, 470–474.

933   Lazarević J, Janković-Tomanić M (2015) Dietary and phylogenetic correlates of digestive
934   trypsin activity in insect pests. *Entomologia Experimentalis et Applicata*, **157**, 123–
935   151.

936   Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler
937   transform. *Bioinformatics*, **25**, 1754–1760.

938   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, … Durbin R (2009) The Sequence
939   Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078–2079.

940   Lozier JD, Strange JP, Koch JB (2013) Landscape heterogeneity predicts gene flow in a
941   widespread polymorphic bumble bee, Bombus bifarius (Hymenoptera: Apidae).
942   *Conservation Genetics*, **14**, 1099–1110.

943   Lozier JD, Strange JP, Stewart IJ, Cameron SA (2011) Patterns of range-wide genetic variation
944   in six North American bumble bee (Apidae: Bombus) species. *Molecular Ecology*, **20**,
945   4870–4888.

946   Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen W-M (2010) Robust
947   relationship inference in genome-wide association studies. *Bioinformatics*, **26**, 2867–
948   2873.

949   Marshall L, Perdijk F, Dendoncker N, Kunin W, Roberts S, Biesmeijer JC (2020) Bumblebees
950   moving up: shifts in elevation ranges in the Pyrenees over 115 years. *Proceedings.*
951   *Biological Sciences*, **287**, 20202201.

952   Martinet B, Lecocq T, Brasero N, Gerard M, Urbanová K, Valterová I, … Rasmont P (2019)
953   Integrative taxonomy of an arctic bumblebee species complex highlights a new
954   cryptic species (Apidae: Bombus). *Zoological Journal of the Linnean Society*, **187**,
955   599–621.

956   Matsuoka T, Takasaki A, Mishima T, Kawashima T, Kanamaru Y, Nakamura T, Yabe T (2015)
957   Expression and characterization of honeybee, Apis mellifera, larva chymotrypsin-like
958   protease. *Apidologie*, **46**, 167–176.

959   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, … DePristo MA
960   (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
961   generation DNA sequencing data. *Genome research*, **20**, 1297–1303.

962   Merilä J, Hendry AP (2014) Climate change, adaptation, and phenotypic plasticity: the
963   problem and the evidence. *Evolutionary Applications*, **7**, 1–14.

Miller-Struttmann NE, Geib JC, Franklin JD, Kevan PG, Holdo RM, Ebert-May D, … Galen C (2015) Functional mismatch in a bumble bee pollination mutualism under climate change. *Science*, **349**, 1541–1544.

Pauls SU, Nowak C, Bálint M, Pfenninger M (2013) The impact of global climate change on genetic diversity within populations and species. *Molecular Ecology*, **22**, 925–946.

Peat J, Tucker J, Goulson D (2005) Does intraspecific size variation in bumblebees allow colonies to efficiently exploit different flowers? *Ecological Entomology*, **30**, 176–181.

Plowright CMS, Plowright RC (1997) THE ADVANTAGE OF SHORT TONGUES IN BUMBLE BEES (BOMBUS) — ANALYSES OF SPECIES DISTRIBUTIONS ACCORDING TO FLOWER COROLLA DEPTH, AND OF WORKING SPEEDS ON WHITE CLOVER. *The Canadian Entomologist*, **129**, 51–59.

Rawlings ND, Barrett AJ (1994) Families of serine peptidases. *Methods in Enzymology*, **244**, 19–61.

Razgour O, Forester B, Taggart JB, Bekaert M, Juste J, Ibáñez C, … Manel S (2019) Considering adaptive genetic variation in climate change vulnerability assessment reduces species range loss projections. *Proceedings of the National Academy of Sciences*, **116**, 10418–10423.

Ruan J, Li H (2020) Fast and accurate long-read assembly with wtdbg2. *Nature Methods*, **17**, 155–158.

Sadd BM, Barribeau SM, Bloch G, Graaf DC de, Dearden P, Elsik CG, … Worley KC (2015) The genomes of two key bumblebee species with primitive eusocial organization. *Genome Biology*, **16**, 76.

Santure AW, Garant D (2018) Wild GWAS—association mapping in natural populations. *Molecular Ecology Resources*, **18**, 729–738.

Scaven VL, Rafferty NE (2013) Physiological effects of climate warming on flowering plants and insect pollinators and potential consequences for their interactions. *Current Zoology*, **59**, 418–426.

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, **9**, 671–675.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM (2015) BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, **31**, 3210–3212.

Soroye P, Newbold T, Kerr J (2020) Climate change contributes to widespread declines among bumble bees across continents. *Science*, **367**, 685–688.

Sun C, Huang J, Wang Y, Zhao X, Su L, Thomas GWC, … Mueller RL (2020) Genus-wide characterization of bumblebee genomes provides insights into their evolution and variation in ecological and behavioral traits. *Molecular Biology and Evolution*.

Teplitsky C, Mills JA, Alho JS, Yarrall JW, Merilä J (2008) Bergmann's rule and climate change revisited: Disentangling environmental and genetic responses in a wild bird population. *Proceedings of the National Academy of Sciences*, **105**, 13492–13496.

Visscher PM, Hemani G, Vinkhuyzen AAE, Chen G-B, Lee SH, Wray NR, … Yang J (2014) Statistical Power to Detect Genetic (Co)Variance of Complex Traits Using SNP Data in Unrelated Samples. *PLOS Genetics*, **10**, e1004269.

Wang M, Xu S (2019) Statistical power in genome-wide association studies and quantitative trait locus mapping. *Heredity*, **123**, 287–306.

Whitley KM (2018) Genetic structure and gene flow barriers among populations of an alpine
bumble bee (Bombus balteatus) in the central Rocky Mountains. M.Sc. Thesis.
Appalachian State University.
Williams PH, Berezin MV, Cannings SG, Cederberg B, Ødegaard F, Rasmussen C, … Byvaltsev
AM (2019) The arctic and alpine bumblebees of the subgenus Alpinobombus revised
from integrative assessment of species' gene coalescents and morphology
(Hymenoptera, Apidae, Bombus). *Zootaxa*, **4625**, zootaxa.4625.1.1.
Williams P, Thorp R, Richardson L, Colla (2014) *Bumble Bees of North America*.
Woodard SH, Lozier JD, Goulson D, Williams PH, Strange JP, Jha S (2015) Molecular tools and
bumble bees: revealing hidden details of ecology and evolution in a model system.
Molecular Ecology, **24**, 2916–2936.
Yang J, Lee SH, Goddard ME, Visscher PM (2011) GCTA: A Tool for Genome-wide Complex
Trait Analysis. *The American Journal of Human Genetics*, **88**, 76–82.
Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL (2014) Advantages and pitfalls in the
application of mixed-model association methods. *Nature Genetics*, **46**, 100–106.

1024
1025
1026 **AUTHOR CONTRIBUTIONS**
1027 Designed research: MJC, NEM-S, JCG, MTW; Performed research: all authors; Analyzed data:
1028 MJC, JCJ, OW, IB, MK, MTW; Wrote the paper: MJC, MTW.
1029
Figure 1. Population sampling and genetic variation in three *Bombus* species. (A) Map
showing the sampling locations of *Bombus balteatus, B. sylvicola,* and "incognitus" on seven
mountains in the Rocky Mountains, CO, USA. Insert indicates location in the USA.
Neighbour-joining trees of (B) *B. balteatus* and (C) *B. sylvicola* and "incognitus" populations
sampled across the seven mountains, based on genome-wide single nucleotide
polymorphisms. Coloured tips indicate the mountain each sample was collected on, with
colours corresponding to those in (A).
1037
1038 **Figure 2. Principal component analysis of Rocky Mountain populations of three *Bombus***
1039 **species.** Plots show the first two principal components from principal component analyses
1040 of (A) *Bombus balteatus*, (B) *B. sylvicola* and (C) "incognitus" populations. Numbers in
1041 brackets on axes indicate the percentage variance explained by each principal component.
1042
Figure 3. Correlations between kinship and geographic distance. Geographic distance
between all pairs of samples plotted against kinship coefficient (a measure of relatedness)
between pairs are shown for (A) *Bombus balteatus*, (B) *B. sylvicola*, and (C) "incognitus",
where each point represents a pair. (D-F) Mantel correlograms showing the Mantel
correlation between kinship and geographic distance over 10 km distance classes for (D)
Bombus balteatus, (E) *B. sylvicola*, and (F) "incognitus". Filled in boxes indicate significant
correlations (p < 0.01). There were insufficient data points for calculating Mantel
correlations at distance classes > 60 km for *B. balteatus* and *B. sylvicola*. For "incognitus",
distance class 10-20 km as well as all distance classes > 50 km lacked sufficient data.
1052
1053 **Figure 4. Comparisons of glossa lengths between species, years, and locations.** (A) Least-
1054 squares mean (ls-mean) glossa lengths correcting for body size for the three *Bombus*
1055 species, based on measurements of the 2017 collections. Least-squares mean glossa

lengths, correcting for body size and sample location, per sampling period for (B) *B. balteatus* and (C) *B. sylvicola* and "incognitus" combined. In all plots, error bars represent ls-means ± standard error and significant differences between classes are indicated by asterisks ($p < 0.05$).

Figure 5. Genome-wide association Manhattan plot. (A) The $-\log_{10}$ p-values from an association test using a mixed linear model between allele frequencies and intertegular distance (ID) are plotted for 2,705,361 genome-wide SNPs against their genome positions over the 18 *Bombus balteatus* pseudochromosomes. Red dashed line indicates genome-wide significance after Bonferroni correction for $p = 0.01$. A single SNP on chromosome 10, coloured and circled in red, had a $-\log_{10}$ p-value above this threshold. Green dashed line indicates genome-wide significance after Bonferroni correction for $p = 0.05$. Two neighbouring SNPs located on chromosome 14, coloured green, had $-\log_{10}$ p-values above this threshold. (B) Zoomed-in plot of the significant SNP in chromosome 10 showing its location within an intron of the *CTRB1* gene. Exons of this gene are represented by purple boxes, introns are represented by dashed purple lines.

Figure 6. Estimates of SNP-heritability for two traits in *Bombus* species. The proportion of variance in glossa length and intertegular distance (ID) explained by all genome-wide SNPs (the SNP-based heritability) was calculated using the genome-base restricted maximum likelihood (GREML) method in GCTA. These were calculated separately for *B. sylvicola* (left panel) and *B. balteatus* (right panel). We included ID as a covariate with glossa length to control for body size. Error bars indicate 95% confidence intervals around estimates. Each analysis was carried out twice, once with all samples (red), and a second time with one from each pair of sisters removed (black), to assess the effect of including highly related samples in the analysis.

Table 1: Assembly metrics for the two genomes, with two other published *Bombus* genomes included for comparison

Species	*B. balteatus*	*B. sylvicola*	*B. terrestris*	*B. impatiens*
Assembly	BBAL_1.0	BSYL_1.0	Bter_1.0	BIMP_2.2
Size (Mbp)†	250.07	252.08	248.65 (236.38)	246.86 (241.98)
Scaffolds (n)	-	-	5,678	5,460
Contigs (n)	336	592	10,672	16,060
Contig N50 (Mbp)	8.60	3.02	0.08	0.06
Contig L50 (n)	12	28	890	54
GC (%)	37.64	38.23	37.51	37.76
Repeat content (%)	17.1	17.9	14.8	17.9
Complete hymenoptera BUSCO genes (%)*	99.0	98.2	96.9	98.3

*BUSCO analysis using the OrthoDB v. 10, Hymenoptera dataset.
† Numbers in brackets represent total ungapped length