1 **Genome sequencing and morphometric analysis of alpine bumblebees experiencing**

2 **climate change reveal ongoing phenotypic shifts and a lack of population structure**

4 Running title: Genome variation in alpine bumblebees
5

Matthew J. Christmas[1], Julia C. Jones[1,2], Anna Olsson[1], Ola Wallerman[1], Ignas Bunikis[3],

Marcin Kierczak[4], Kaitlyn M. Whitley[5,6], Isabel Sullivan[5,7], Jennifer C. Geib[5], Nicole E. Miller-

Struttmann[8], Matthew T. Webster[1]

1) Department of Medical Biochemistry and Microbiology, Science for Life Laboratory,

Uppsala University, Uppsala, Sweden

2) School of Biology and Environmental Science, University College Dublin, Dublin, Ireland

3) Department of Immunology, Genetics and Pathology, Science for Life Laboratory, Uppsala

University, Uppsala, Sweden

4) Department of Cell and Molecular Biology, National Bioinformatics Infrastructure

Sweden, Science for Life Laboratory, Uppsala University, Uppsala, Sweden

5) Department of Biology, Appalachian State University, Boone, North Carolina, USA

6) U.S. Department of Agriculture, Agriculture Research Service, Charleston, South Carolina,

USA

7) Marine Estuarine Environmental Sciences, University of Maryland, College Park Maryland

8) Biological Sciences Department, Webster University, St. Louis, Missouri, USA

Data submission: NCBI project IDs PRJNA646847, PRJNA704506

24
25

Corresponding author: Matthew T. Webster

Address: Dept. Medical Biochemistry and Microbiology, Box 582, 75123 Uppsala, Sweden

Email: matthew.webster@imbim.uu.se

Tel: +46 18 471 4391

ABSTRACT

Populations of the bumblebees *Bombus sylvicola* and *Bombus balteatus* in Colorado have

experienced decreases in tongue length, a trait important for plant-pollinator mutualisms, in

the last six decades. It is hypothesized that this reflects selection exerted by changes in

floral composition under climate change. Here we combine extensive morphometric and

population genomic data to investigate population structure, whether morphological

change is ongoing, and the genetic basis of morphological change. We generate highly-

contiguous genome assemblies of both species using long-read sequencing. We then

perform whole-genome sequencing and morphometric measurements of 580 samples of

these species from seven high-altitude localities. Out of 281 samples originally identified as

B. sylvicola, 67 formed a separate genetic cluster comprising the newly-discovered cryptic

species *B. incognitus*. However, there is very little additional genetic substructure,

suggesting that gene flow occurs readily between mountains. We find a significant decrease

in tongue length between bees collected between 2008-2014 and in 2017, indicating that

morphological shifts are ongoing. We did not discover any genetic associations with tongue

length, but a SNP related to production of a proteolytic digestive enzyme is implicated in

body size variation. We identify evidence of covariance between kinship and both tongue

length and body size, which is suggestive of a genetic component of these traits, although it

is not possible to rule out shared environmental effects between colonies. Our results

provide evidence for ongoing evolution of a morphological trait important for pollination

and indicate that this trait likely has a complex genetic and environmental basis.

Keywords: genome-wide complex trait analysis, plant-pollinator interaction, population

structure, genome assembly, population genomics, global warming.

INTRODUCTION

The importance of pollinators for ecology and the maintenance of biodiversity is widely

recognized (Goulson, Nicholls, Botías, & Rotheray, 2015). Bumblebees are large-bodied,

mainly cold-adapted species with broad significance for agricultural production and plant-

pollinator networks in the wild (Goulson, 2003). Many bumblebee species are experiencing

range shifts or declines due to climate change or habitat loss (Cameron et al., 2011;

Cameron & Sadd, 2020; Goulson, Lye, & Darvill, 2008; Soroye, Newbold, & Kerr, 2020). In

particular, a general worldwide trend for species to shift to higher elevations and latitudes

has been documented, with losses occurring at a faster rate than expansions (Kerr et al.,

2015; Marshall et al., 2020). Arctic and alpine species are particularly threatened by these

trends.

In addition to range shifts there is also evidence that the morphology of certain populations

of bumblebees has changed in recent decades. Miller-Struttmann et al. (Miller-Struttmann

et al., 2015) compared morphometric measurements of contemporary samples and

historical specimens of the species *Bombus balteatus* and *Bombus sylvicola* from the same

high-altitude localities in Colorado. These species historically comprise the majority of

bumblebees in these locations. They reported a significant decrease in tongue length in

both species in a period between 1966 and 2014. The cause of this was inferred to be a

general decrease in floral abundance resulting from warmer temperatures and dryer soils.

These conditions favor more generalist foraging that is performed more optimally by bees

with shorter tongues (Goulson, Hanley, Darvill, Ellis, & Knight, 2005). Continued monitoring

of these populations is important to understand the ongoing effects of climate change, but

it is unknown whether the trend toward shorter tongues has continued in more recent

years.

One explanation for the observed morphological shifts in *B. balteatus* and *B. sylvicola* in

Colorado is that natural selection for shorter tongues has occurred (Miller-Struttmann et al.,

2015). However, it is also possible that the observed changes are due to phenotypic

plasticity. It is unknown whether this trait has a strong genetic component that could be

acted on by natural selection. Uncovering the size of the genetic contribution to this trait

and the number and identity of genetic loci involved is an important step towards

understanding the underlying cause of the morphological changes.

The genetic structure of these populations is also relevant to how they evolve under climate

change (Pauls, Nowak, Bálint, & Pfenninger, 2013). It is also important to understand

population structure in order to interpret genotype-phenotype correlations and cause of

morphological shifts. Importantly, using population resequencing, we have identified the

presence of a previously-undetected cryptic species, which we called *Bombus incognitus*

(Christmas, submitted). This species is identified morphologically as *B. sylvicola* but forms a

distinct genetic cluster. It is therefore possible that morphological changes in the combined

populations of these species could be brought about by changes in the relative abundance

of each species, although this has not previously been investigated. Additional population

structure within species could lead to a similar effect, particularly if genetic structure

associated with morphology exists. An understanding of spatial connectivity between

populations in different locations is also needed to predict how they will evolve under

climate change and to define the conservation value of subpopulations (Pauls et al., 2013;

Razgour et al., 2019).

The aim of this study was to produce high-quality annotated genomes of the bumblebee

species *B. balteatus* and *B. sylvicola* and use them to assess whole-genome variation and its

connection to geographical and morphological variation in the populations of these species

previously studied by (Miller-Struttmann et al., 2015). We constructed genome assemblies

based on long-read Oxford Nanopore sequencing. We then performed population re-

sequencing of 580 samples identified morphologically as these species in Colorado in seven

high-altitude localities spread across 150 km. We studied levels of genetic variation,

geographical structure and patterns of relatedness between geographical areas in *B.*

balteatus, *B. sylvicola*, and the recently-revealed cryptic species *B. incognitus* (Christmas,

submitted), which was also present in our dataset. We then analyzed morphological

variation in tongue length and body size in context of previous studies (Miller-Struttmann et

al., 2015) in order to determine whether the trend towards decreasing tongue length is

ongoing. We also performed a genome-wide analysis of the genetic basis of morphological

variation using both genome-wide association studies (GWAS) and genome-wide complex

trait analysis (GCTA) (Yang, Lee, Goddard, & Visscher, 2011). The results are informative

regarding the degree of connectivity between bumblebee populations, the genetic basis of

morphological variation, and the mechanisms of morphological evolution. They also indicate

a continuing impact of climate change on the populations.

MATERIALS AND METHODS

Study system

The species *B. balteatus* and *B. sylvicola* belong to two subgenera of bumblebees

(Alpinobombus and Pyrobombus respectively) that are common in alpine and arctic

environments. Both species are closely related to populations in Eurasia that different

studies have considered either species or subspecies. *Bombus sylvicola* in North America is

closely related to *B. lapponicus* in Eurasia and a recent study classifies it as a subspecies of *B.*

lapponicus rather than a separate species (Martinet et al., 2019). Similarly, two subspecies

of *B. balteatus* have been described previously, restricted respectively to North America and

Eurasia, but recent study split these populations into two separate species: North American

B. kirbiellus and Eurasian *B. balteatus* (Williams et al., 2019). Here we maintain the names *B.*

balteatus and *B. sylvicola* of our species of interest for consistency with previous ecological

studies that we are directly comparing here (Geib, Strange, & Galen, 2015; Miller-

Struttmann et al., 2015) and ambiguities regarding species/subspecies status.

Sample collection

We collected samples identified as *B. balteatus* and *B. sylvicola* during the summer of 2017.

Bumblebees were collected on seven different mountains in Colorado, USA. The existence

of *B. incognitus* was unknown to us when the bees were sampled and a large fraction of

bees identified as *B. sylvicola* were subsequently assigned to *B. incognitus* on the basis of

genetic analysis (Christmas, submitted). Three of these mountains (Mount Evans

(39.583883, -105.63845), Niwot Ridge (40.05945, -105.616667), and Pennsylvania Mountain

(39.263383, -106.142733)) were also sampled previously by (Miller-Struttmann et al., 2015).

Four additional mountains were sampled here (Boreas Mountain (39.419375, -105.95441),

Mount Democrat (39.33721, -106.12581), Horseshoe Mountain (39.204605, -106.181085)

and Quail Mountain (39.018158, -106.404877)) but were not sampled in the previous study.

Each of the mountain locations comprised several neighboring sampling sites separated by

short distances and altitudes (full details in supplementary tables S1, S2).

We collected samples of foraging worker bees from each site using sweeping hand nets. We

kept samples in Falcon tubes in cool boxes for transport. Species assignment was performed

by observing standard morphological characters. We dissected each sample after placing it

at -20°C for approximately 10 minutes. We stored thoraces in 95% ethanol prior to DNA

extraction whereas the heads were stored for morphometric analysis.

Morphological measurements

Morphological measurements

We measured intertegular distance (ID) on all of the samples of *B. sylvicola, B. incognitus*

and *B. balteatus* using scaled photographs of individual bee heads processed by ImageJ

("ImageJ," n.d.). We characterized the diagnostic traits visible on each head, such as the

shape of the malar space, the pile color between and antenna and above the ocelli, the size,

color and location of the ocelli relative to the supraorbital line, and the presence of black

pile on the scutellum. We then dissected the prementum and glossa from the head, which

were then mounted and measured by photographing and analysis in ImageJ ("ImageJ," n.d.).

None of the morphological characters measured could be used to distinguish between *B.*

incognitus and *B. sylvicola*.

Genome sequencing and assembly

We used the same sequencing strategy to produce high-quality genome assemblies for both

B. sylvicola and *B. balteatus*. This involved combining data from Oxford Nanopore (ONT) and

10x Genomics Chromium technologies. In both cases, we used DNA extracted from male

bees sampled on Niwot Ridge and stored in 95% ethanol prior to DNA extraction. DNA from

a single bee was used for each assembly. We performed DNA extraction using a salt-

isopropanol extraction followed by size selection using magnetic bead purification to

remove fragments < 1 kbp. ONT sequencing was performed using the MinION instrument.

For *B. sylvicola*, we used two R9.4 flowcells and the RAD004 kit starting with 3-400 ng DNA

per run. This resulted in a total of 9.4 Gbp of sequence data, in 2.5 million reads with a

mean length of 3.7 kbp. For *B. balteatus*, we used 1µg DNA in a modified LSK-108 one-pot

protocol. 1.75 µl NEB FFPE buffer, 1.75 µl NEB Ultra-II End Prep buffer and 1 µl FFPE repair

mix was added to 24 µl DNA and incubated at 20°C for 30 minutes. 2 µl End Prep enzyme

mix was then added for blunting, phosphorylation and A-tailing (20 min. 20°C, 10 min. 65°C,

10 min. 70°C). AMX (20 µl), ligation enhancer (1µl) and Ultra-II ligase (40 µl) was added for

ligation at room temperature for 30 minutes. Clean-up and flowcell loading was done

following the standard ONT LSK-108 protocol and the library was sequenced on one R9.4

flowcell. This resulted in 23.3 Gpb of sequence data, in 3.9 million reads with a mean length

of 5.9 kbp.

We assembled the ONT reads using the following process: We first used downpore for

adaptor trimming and splitting chimeric reads. We then assembled the trimmed reads using

wtdbg2 (Ruan & Li, 2020) on default settings. We then ran two rounds of the consensus

module Racon and contig improvements using medaka v.0.4, removing contigs of length <

20 kbp. We used Illumina short-read data from the population sequencing to perform two

rounds of Pilon in order to improve sequence accuracy, particularly around indels.

We also sequenced the same sample of each species using 10x Genomics Chromium. A 10x

GEM library was constructed according to the manufacturer's protocols. We quantitated

each library using qPCR and sequenced them together on a single lane of HiSeq 2500 using

the HiSeq Rapid SBS sequencing kit version 2, resulting in 150 bp paired-end sequences. The

linked reads were mapped to the assembly using Longranger v.2.1.4. We then ran Tigmint

v1.1.2 to correct errors in the assembly detected by the read mappings. We removed

contigs containing multiple mitochondrial genes identified using BLAST+ v2.9.0. We also

removed all contigs shorter than 10 kbp.

Genome analysis

BUSCO v3.0.2b (Simão, Waterhouse, Ioannidis, Kriventseva, & Zdobnov, 2015) was used to

assess the assembly completeness using the hymenoptera_odb9 lineage set and species *B.*

impatiens. We ran RepeatMasker v.4.1.0 on each genome assembly to characterize genome

repeat content, including interspersed repeats and low complexity sequences. It has been

shown that bumblebees have a stable 18 chromosome karyotype across all subgenera (Sun

et al., 2020). We therefore performed whole-genome synteny alignments between the *B.*

terrestris chromosome-level genome assembly (Sadd et al., 2015) (downloaded from NCBI;

BioProject PRJNA45869) and our assemblies using Satsuma v.3 (Grabherr et al., 2010) to

arrange the contigs from our assemblies onto 18 pseudochromosomes.

RNAseq & Annotation

We performed RNAseq on a single lane of Illumina HiSeq 2500 using samples from four

tissues from both *B. syvicola* and *B. balteatus* (abdomen, head, legs and thorax). This

generated 44 Gbp of reads for *B. syvicola* and 42 Gbp for *B. balteatus.* We performed quality

control of the data using FastQC and read trimming using fastp (v. 0.20.0). We next

generated a guided assembly using Hisat2 and Stringtie (v. 2.0). We also performed a *de*

novo assembly for reads from each of the four tissues using Trinity (v. 2.0.4). We then

generated a species-specific repeat library using the RepeatModeller package. We used

RepeatMasker to identify the location of repeats from this library in the genomes.

We used the MAKER package to compute gene builds. As protein evidence, we downloaded

curated protein sequences from the Uniprot Swiss-prot database (561,356 proteins,

downloaded 11/2019). An evidence-guided gene build was computed by MAKER

constructing gene models from these reference proteins and from the transcript sequences

aligned to the genome assembly.

Population sequencing, read mapping and variant calling

We extracted DNA from the thoraces of worker bees of the species *B. sylvicola, B. incognitus*

and *B. balteatus* collected from across the seven sampling sites using the Qiagen Blood and

Tissue kit. We prepared libraries using the Nextera Flex kit and performed sequencing on

the Illumina HiSeq X to produce 2 x 150 bp reads. We mapped reads to the appropriate

reference genome (*B. sylvicola* for *B. sylvicola* and *B. incognitus* samples; *B. balteatus* for *B.*

balteatus samples) using the mem algorithm in BWA (H. Li & Durbin, 2009). We performed

sorting and indexing of the resultant bam files using samtools (Heng Li et al., 2009) and

marked duplicate reads using Picard. We used the Genome Analysis Toolkit (GATK)

according to best practices to call variants (McKenna et al., 2010). We first ran

HaplotypeCaller on the bam file of each sample to generate a gVCF file for each sample. We

then used GenomicsDBImport followed by GenotypeGVCFs to call variants for all *B.*

balteatus samples together and all *B. sylvicola* and *B. incognitus* samples together. We used

thresholds recommended by GATK best practices to filter SNPs using VariantFiltration tool.

Only biallelic SNPs were considered for downstream analysis.

Population structure and relatedness

We thinned the VCF files to 1 SNP every 10 kbp using the --thin flag in vcftools v.0.1.16 in

order to more efficiently assess population structure and relatedness. We calculated p-

distance matrices for *B. balteatus* samples and *B. sylvicola* and *B. incognitus* samples

combined from the thinned vcf files using VCF2Dis (https://github.com/BGI-

shenzhen/VCF2Dis) and then constructed neighbor-joining trees using the 'ape' package in R

v4.0.2. We converted the thinned vcf files to plink format and performed principal

components analyses using the 'adegenet' package in R v4.0.2. We used admixture v1.3.0

(Alexander, Novembre, & Lange, 2009) to assess the most likely number of genetic clusters

within each species. We ran admixture for K=1 through to 10, with 10 iterations per K value

and used the calculated cross-validation (cv) errors to assess the most suitable values of K,

with low cv-error indicating higher support.

We estimated the pairwise kinship coefficient between all samples using the method

described in (Manichaikul et al., 2010) implemented in the --relatedness2 option of vcftools

v0.1.16 (Danecek et al., 2011). Worker bees from the same bumblebee colony are expected

to have a relatedness of 0.75 under the assumption of monandry, which is reasonable for

the study species (Estoup, Scholl, Pouvreau, & Solignac, 1995). The kinship coefficient (φ)

predicted from this relationship is $\frac{3}{8}$ (the probability that two alleles sampled at random

from two individuals are identical by descent). We estimated the lower cutoff for the

inference criteria to infer this level of relatedness using powers of two as described in

(Manichaikul et al., 2010), providing an estimate of $\frac{3}{2^{7/2}}$.

Statistical morphometric analysis

We combined the phenotypic measurements made in this study with those from previous

collections published in (Miller-Struttmann et al., 2015). These measurements are ID, glossa

length, prementum length, and total tongue length (equal to glossa length plus prementum

length). This gave us three time periods to compare amongst: '2017' (samples collected for

this study), '2012-2014', and '1966-1980' (both sets published in (Miller-Struttmann et al.,

2015)). We performed statistical analysis of variation in glossa length, prementum length,

and ID in R v4.0.2. We investigated correlations between all traits for each species in the

2017 set using the 'cor.test' function, which calculates a Pearson's product moment

correlation coefficient. For glossa length, we calculated least-squares means (ls-means) to

account for body size (represented by ID) and compared between species, geographic

locations and years of sample collection using the 'lsmeans' function. We analyzed variation

in ID between species, geographic locations and years of sample collection using analysis of

variance ('aov' function) and assessed significance of comparisons using Tukey's honestly

significant difference (HSD) post hoc test ('TukeyHSD' function).

Genome-wide association study

We carried out mixed linear model-based association analyses, implemented in GCTA-

MLMA (Yang et al., 2011; Yang, Zaitlen, Goddard, Visscher, & Price, 2014), to identify

correlations between allele frequencies and trait variation across the genome for *B.*

balteatus and *B. sylvicola* samples separately (*B. incognitus* samples were left out of this

analysis due to too few samples). First, we created a genetic relationship matrix (GRM) from

the genome wide SNPs using the --make-grm function in GCTA. We removed one sample

from each pair of sisters by setting a relatedness cut-off of 0.67 using the '--grm-cutoff' flag.

We ran the analyses for three traits per species:ID, glossa length, and the residuals from a

linear regression of glossa~ID in order to control for body size. The GRM was included as a

polygenic/random effect. The analysis assumes phenotypes are normally distributed and so

samples were removed in the tails of the distributions to meet this assumption for all traits

except ID in *B. sylvicola*, which was normally distributed. We assessed divergence of p-

values from neutral expectations using QQ-plots and carried out Bonferroni correction of p-

values using the 'p.adjust' function in R v4.0.2 to assess genome-wide significance.

Genome-wide complex trait analysis

We used GCTA (Yang et al., 2011) to perform genome-wide complex trait analysis. We

estimated the phenotypic variance explained by the genome-wide SNPs using genome-

based restricted maximum likelihood (GREML). The analysis was performed separately for

each species on measures of ID, glossa length, and glossa length with ID included as a

covariate to control for body size. We ran each analysis twice, once using a GRM based on

all samples and a second time with the same GRM produced above where one from each

pair of sisters was removed. We calculated 95% confidence intervals around each estimate

by multiplying the standard error by 1.96. We used R functions provided in (Wang & Xu,

2019) as well as the GCTA-GREML power calculator (Visscher et al., 2014) to perform power

analysis and estimate the effect sizes we can expect to detect with our population datasets.

RESULTS

Generation of highly-contiguous genome assemblies of *B. sylvicola* and *B. balteatus*

We generated *de novo* genome assemblies for the species *B. sylvicola* and *B. balteatus* using

Oxford Nanopore (ONT) long-read and 10x Chromium linked-read sequencing. Assembly

statistics for BBAL_1.0 (*B. balteatus*) and BSYL_1.0 (*B. sylvicola*) are presented in table 1,

which includes statistics for the assemblies of *B. terrestris* (BTER_1.0) and *B. impatiens*

(BIMP_2.2) for comparison. Total assembly sizes of 250.1 and 252.1 Mbp for *B. balteatus*

and *B. sylvicola* respectively are both above the average assembly size of *Bombus* genomes

assembled to date of 247.9 Mbp (22 genomes, range 229.8 – 282.1 Mbp) (Heraghty et al.,

2020; Sadd et al., 2015; Sun et al., 2020). Contig N50s of 8.60 and 3.02 for BBAL_1.0 and

BSYL_1.0 respectively are the longest and third longest contig N50s of all published *Bombus*

assemblies, demonstrating these to be highly contiguous assemblies. Megabase-scale contig

N50 are typical of long-read assemblies, such as the ones presented here and in (Heraghty

et al., 2020), and are significantly more contiguous than those based on short-read

sequencing technologies such as BTER_1.0 and BIMP_2.2.

High genome completeness and accuracy is also reflected in the BUSCO analysis, with scores

of 99.0% (98.7% single copy, 0.3% duplicated, 0.5% fragmented, 0.5% missing) and 98.2%

(97.9% single copy, 0.3% duplicated, 0.4% fragmented, 1.4% missing) reported for BBAL_1.0

and BSYL_1.0 respectively. Whole genome alignments to the *B. terrestris* assembly (Sadd et

al., 2015) resulted in 93.9% and 91.1% of the *B. balteatus* and *B. sylvicola* genomes being

placed on 18 pseudochromosomes. Our annotation pipeline annotated 11,711 and 11,585

genes in BBAL_1.0 and BSYL_1.0 respectively. These are highly comparable to the 11,874

genes in the *B. terrestris* gene set (v. 1.0), but lower than the 12,728 genes in the *B.*

impatiens gene set (v. 2.1). Gene annotation of the *Bombus* assemblies presented in

(Heraghty et al., 2020; Sun et al., 2020) also resulted in a greater number of genes

annotated, ranging from 13,325 to 16,970 genes. BUSCO analyses of the annotated genes in

our assemblies do suggest that we are missing some genes from our annotations (complete

BUSCO genes in BBAL_01 and BSYL_01 respectively: 86.6% and 86.9%). Repeat content of

our assemblies is comparable to that of other *Bombus* assemblies, with 17.11% of BBAL_1.0

and 17.85% of BSYL_1.0 identified as repetitive by RepeatMasker (full details of repeat

classes in supplementary table S3). These proportions are comparable to the repeat content

inferred in the *B. terrestris* and *B. impatiens* genomes.

Population resequencing indicates three species clusters, with little population structure within species

The locations of sample collections, which comprise 580 samples from across seven localities in the Rocky Mountains, CO, USA, are presented in Fig. 1 (full details in table S1, S2). These samples were initially identified morphologically as *B. balteatus* and *B. sylvicola*. However, neigbor-joining trees produced from samples of both groups of samples indicate that the *B. balteatus* samples form a single cluster whereas the *B. sylvicola* samples form two clusters (Fig 1). By comparison with an additional dataset (Martinet et al., 2019) we previously inferred the largest cluster to be comprised of *B. sylvicola* samples whereas the second cluster is comprised of the cryptic species *B. incognitus* (Christmas, submitted). Among samples originally designated as *B. sylvicola,* 24% (n=67) are identified in the separate cluster of *B. incognitus* whereas the remaining 76% (n=214) are *B. sylvicola. Bombus balteatus* (n=299) was the most commonly observed species across all localities, followed by *B. sylvicola* (n=214) and *B. incognitus* (n=67). The proportions vary between localities (table S1). *Bombus incognitus* samples were collected on six out of the seven mountains, whereas *B. balteatus* and *B. sylvicola* were found on all seven.

We calculated a pairwise kinship coefficient between samples to infer degrees of relatedness (Danecek et al., 2011; Manichaikul et al., 2010). We inferred a set of samples that were likely to be nestmates. In haplodiploids such as bumblebees, females with the same maternal and paternal parents have a relatedness of 0.75. This is likely to be the case for all workers from the same colony in these species, under the realistic assumption that each colony is headed by a single monandrous female queen (Estoup et al., 1995). Those identified as sisters from the same single-mated queen from the same colony were always collected foraging on the same mountain and at the same or immediately neighboring site (Table S1). This suggests that foraging bees were generally found in the immediate vicinity of their nests.

We investigated the presence of population genetic structure within each species using several methods. Neighbor-joining trees (Fig. 1) do not reveal the presence of any substructure related to geography for *B. balteatus* or *B. incognitus*. For *B. sylvicola*, all of the

samples from Quail Mountain (the furthermost southwestern site) cluster together, but do

not separate from the rest on the tree. Principal component analysis further reflected the

extremely low level of geographical structure across all species (Fig. 2). Outlier clusters seen

in the PCA plots of *B. balteatus* and *B. incognitus* (Fig. 2A,C, circled) are composed solely of

the groups of sibling nestmates identified by the kinship analysis, suggesting that the effect

of kinship overrides any signal of genetic structure. One slight exception to this trend is the

Quail Mountain population of *B. sylvicola* which, as with the neighbor-joining tree, shows a

slight degree of clustering away from the other samples at the positive extreme of PC1

(Fig.2B), but they still overlap with other samples and PC1 explains only 1.17% of the

variance in the data.

We assessed connectivity between colonies on different mountains by analyzing the

association between kinship and geography using a Mantel test. Mantel correlograms

indicate a significant association between kinship and geographical distance at shorter

distances (from the same mountain), which is no longer significant at larger distances (Fig.

3). Hence there is a no significant signal of increased degrees of kinship on neighboring

mountains in our data.

Decrease in tongue length is ongoing in both species

We made morphological measurements for each sample (Table S2). Intertegular distance

(ID) is considered a proxy for body size (Cane, 1987). We also measured the length of two

parts of the tongue: glossa and prementum. These two tongue measures are correlated in

both species (*B. balteatus* Pearson's r = 0.45, p < 0.001; *B. sylvicola* Pearson's r = 0.38, p <

0.001; Figs. S1,S2). Only glossa is considered further, as it is the tongue part with most

relevance to feeding (Cariveau et al., 2016) and was measured by (Miller-Struttmann et al.,

2015). Glossa is also significantly correlated with ID (*B. balteatus* Pearson's r = 0.41, p <

0.001; *B. sylvicola* Pearson's r = 0.26, p < 0.001) (Figs. S1,2).

A comparison of phenotypes between species shows that glossa length is substantially

longer in *B. balteatus*, consistent with previous knowledge (ls-means = 4.72 mm, 3.34 mm,

and 3.40 mm for *B. balteatus*, *B. sylvicola* and *B. incognitus* respectively; Tukey test, t ratio =

12.25 and 21.17, p < 0.001 for *B. balteatus* compared to *B. sylvicola* and *B. incognitus*;

Fig.4A). Average glossa length is extremely similar in *B. sylvicola* and B. *incognitus* (Tukey

test, t ratio =0.50, p = 0.87), suggesting that this character does not distinguish this newly-

discovered species from *B. sylvicola*. *Bombus balteatus* bees are significantly larger bees as

indicated by a greater ID, and *B. sylvicola* is significantly larger than *B. incognitus* (mean ID =

4.00 mm, 3.83 mm, 3.58 mm for *B. balteatus*, *B. sylvicola* and *B. incognitus* respectively;

ANOVA, F=28.75, p < 0.001, Tukey post-hoc test revealed differences are significant among

all species at p < 0.001; Fig. S3A).

We next compared the distribution of glossa lengths in our samples with historical data. We

combined *B. sylvicola* and *B. incognitus* for this analysis because they do not differ in glossa

length on average, and in order to compare to historical data in which the two species were

not distinguished. We find an overall trend of decreasing glossa length in both *B. sylvicola-*

incognitus and *B. balteatus* over time (Fig. 4B,C). This was reported previously between the

collections from 1966-1980 and from 2012-2014 for both species (Miller-Struttmann et al.,

2015). Here we show that this trend is continuing, with further reductions in glossa length

between the 2012-2014 collections and our collections from 2017. This reduction is

significant for *B. sylvicola-incognitus* (ls-means: 2012-2014 = 3.72 mm, 2017 = 3.24 mm;

Tukey test, t-ratio = 3.785, p < 0.001) and suggestive for *B. balteatus* (ls-means: 2012-2014 =

5.45 mm, 2017 = 5.11 mm; Tukey test, t-ratio = 2.026, p = 0.11). In contrast, there is no

significant trend shown by ID (Fig. S3B,C), although for *B. sylvicola-incognitus,* ID is

significantly larger in both the 1966-80 and 2017 samples compared to the 2012-2014

samples (ANOVA, F = 15.28, p < 0.001. Tukey post-hoc test, p < 0.001).

The trend for a decrease in glossa length occurs independently on all mountains for which

data is available in both species, but the decrease between 2012-2014 samples and 2017

samples is only significant for *B. sylvicola-incognitus* on Pennsylvania mountain (ls-means:

2012-2014 = 3.58 mm, 2017 = 2.96 mm, Tukey test, t-ratio = 3.61, p = 0.002 ; Fig. 4D,E).

There are no discernable historical trends over the same period for ID, and fluctuations in

opposite directions are observed among mountains (Fig. S3D,E).

For samples collected in 2017, we compared the distribution of glossa lengths for each

species among the seven mountains where they were collected (Fig. S4). There is an overall

trend for glossa lengths to be longer in bees from Mount Evans than on other mountains in

all species. The distributions for *B. balteatus* and *B. sylvicola* show significant differences,

with samples from Mount Evans having significantly longer glossae than on Mount

Democrat, Horseshoe Mountain, and Pennsylvania Mountain in *B. balteatus* (ls-mean glossa

length on Mt. Evans = 5.34 mm, other mountains 4.51 – 4.73 mm; Tukey p < 0.01 in each

case; Fig. S4A) and significantly longer glossae than on all other mountains except Niwot

Ridge in *B. sylvicola* (ls-mean glossa length on Mt. Evans = 3.77 mm, other mountains 3.02 –

3.24 mm; Tukey p < 0.01 in each case; Fig. S4B). None of the trends for *B. incognitus* were

significant (Fig. S4C). Comparisons of ID between mountains revealed that *B. balteatus* bees

sampled on Niwot Ridge are significantly larger than those on Boreas Mountain, Mount

Democrat, Horseshoe Mountain, and Quail Mountain (Niwot Ridge mean ID = 4.25 mm,

other mountains 3.84 - 4.00 mm, ANOVA, F = 4.09, p < 0.001, Tukey post-hoc test p < 0.05 in

four comparisons; Fig. S5A). We did not identify any significant differences in ID of bees

across mountains for *B. sylvicola* (ANOVA, F = 1.83, p = 0.10) or *B. incognitus* (ANOVA, F =

1.62, p = 0.17; Fig. S5B,C).

Genome-wide association study identifies a SNP associated with a proxy for body size

We used a linear mixed-model analysis to search for correlations between allele frequencies

and quantitative trait variation in *B. sylvicola* and *B. balteatus.* We used the traits glossa

length, ID and glossa length corrected for ID (see Methods). One from each pair of sisters

identified in the kinship analysis was removed from all analyses to reduce the level of

relatedness in the sample set. We detected a genome-wide significant association between

ID and a SNP in an intron of the *CTRB1* (Chymotrypsinogen B1) gene on chromosome 10 in

B. sylvicola (p < 0.001 after Bonferroni correction; Figs. 5, S6F). This gene is involved in

production of chymotrypsin, a key proteolytic digestive enzyme in bees, involved in

digestion of pollen. It is highly plausible that variation in this gene affects the rate of growth

and development in bees through its role in digestion. Two further SNPs, located 4 bp apart

on chromosome 11, also significantly associated with ID (p = 0.039 after Bonferroni

correction). However, they are not located in or close to any genes according to our

annotation, so any potential functional significance of these SNPs is unknown.

We did not find any significant associations between allele frequencies and ID in *B.*

balteatus (Fig. S6). Furthermore, no significant associations were found between allele

frequencies and glossa length, or glossa length corrected for ID in either species (Fig. S6).

This indicates that the trait has a polygenic or mainly environmental component. A power

analysis (Visscher et al., 2014) revealed that our datasets have sufficient power (>0.8) to

detect a QTL with a h^2 of ~0.1 but that loci with lower effect sizes would likely be

undetectable.

Limited evidence that morphological variation has a genetic component

We performed GCTA-GREML analyses (Yang et al., 2011) to determine the genetic

component of trait variation for *B. sylvicola* and *B. balteatus* (Fig. 6, table S4). The *B.*

incognitus samples were not included here due to smaller sample size. When each species'

entire dataset is considered, we detect a heritable component for both glossa length and ID

in both species, where the 95% confidence interval does not overlap zero. For *B. sylvicola*,

but not *B. balteatus*, this is also the case when ID is introduced as a covariate to account for

body size when considering glossa length. However, when we remove samples that we infer

to be workers from the same colony from the kinship analysis, we only detect a heritable

component with 95% confidence for ID in *B. balteatus* (all others then overlap with zero).

This indicates that a large proportion of the signal of covariance between genetic and trait

variance is due to the presence of sisters from the same colony. This could be a confounding

factor because environmental effects such as level of nutrition and colony health are likely

to covary with colony. However, such covariation could also reflect genetics. The level of

dependency of phenotype on genotype is therefore unclear from this analysis.

In these analyses, 95% confidence intervals are calculated by multiplying the standard error

by 1.96. As such, to achieve 95% confidence intervals that do not span the entire range of

possible heritability (0-1), standard errors must be kept low by increasing sample size. For

unrelated individuals it has been estimated that > 3,000 samples are required to achieve a

SE below 0.1 (Visscher et al., 2014).With our datasets, we have a power of ~0.06, or a ~6%

probability, to detect h^2 > 0 for a trait with a SNP-heritability of 0.5 and a type 1 error rate

(α) of 0.05. The low power of our datasets has therefore strongly influenced the estimates

we achieved.

DISCUSSION

DISCUSSION

In this study, we generated high-quality reference genomes for two bumblebee species: *B.*

sylvicola and *B. balteatus*. We also produced the most extensive population genomics

dataset for bumblebees to date, with 583 samples from these two species and the newly-

discovered *B. incognitus* collected from high-altitude locations in Colorado, USA. These data

were combined with morphometric measurements from each sample and used to

investigate genetic structure, morphological evolution and the genetic basis of phenotypic

traits in these populations. Our main findings are 1) no evidence for gene flow between but

lack of population structure within species, indicating high levels of connectivity among

subpopulations. 2) An ongoing decrease in tongue (glossa) length compared to historical

measurements, which likely reflects the continuing effects of decreasing floral abundance

under climate change on bumble bee foraging (Miller-Struttmann et al., 2015). 3) An

association between body size and variation at a single SNP located in a gene related to

production of a digestive enzyme in *B. sylvicola*. 4) A lack of conclusive evidence for a

genetic component to tongue length in both species. Our analysis is unable to determine

the degree of heritability underlying these traits but is consistent with a highly polygenic

genetic component.

This investigation adds to the number of bumblebee species with reference genomes

available. The genomes of the common species *B. impatiens* and *B. terrestris* were

sequenced using short-read technologies (Sadd et al., 2015). The availability of genetic maps

for *B. terrestris* allowed placement of contigs on chromosomes. Genome assemblies are also

available for representatives of all 15 *Bombus* subgenera (Sun et al., 2020) based on short-

read sequencing of jumping libraries and scaffolding using Hi-C, resulting in contig N50 of

325 kb. In addition to the two genomes produced here, the genomes of the North American

species *B. terricola*, *B. bifarius*, *B. vancouverensis*, and *B. vosnesenskii* have also been

produced using long-read sequencing (PacBio or ONT) and have megabase-scale contig N50s

(Heraghty et al., 2020; Kent et al., 2018).

Identification of the previously-unknown species *B. incognitus* demonstrates the power of

population-scale genome sequencing to discover new cryptic variation (Christmas,

submitted). None of the morphological features that we studied could distinguish between *B. sylvicola* and *B. incognitus*. There were no significant differences in glossa or prementum length. *Bombus sylvicola* did however have a significant tendency to be collected at higher altitudes and have larger body size. The wider distribution of *B. incognitus* is unknown, as is the true composition of populations described as *B. sylvicola* in other geographical locations.

Another cryptic species, *B. interacti*, has been reported in Alaska, which is related to *B. sylvicola* from North America and *B. lapponicus* from Eurasia (Martinet et al., 2019). This species has subsequently been inferred to be synonymous with a previously-described species found in the far north of North America, *B. johanseni* (Sheffield, Oram, & Heron, 2020). However, this species clusters more closely with *B. monticola* based on genetic data, whereas *B. incognitus* displays closer affinity to *B. sylvicola* and *B. lapponicus* (Christmas, submitted; Martinet et al., 2019). The same study also classified *B. sylvicola* as the same species as *B. lapponicus*, which is found in Eurasia, due to a short genetic distance between them (Martinet et al., 2019). It is however clear that the North American *B. sylvicola* and the Eurasian *B. lapponicus* have experienced a prolonged period of geographic isolation, during which intrinsic barriers to gene flow may have emerged. Further population genomic studies of Pryobombus bumblebees from other geographic regions could potentially uncover new cryptic species and determine species relationships and population structure at larger scale.

We find no evidence for geographical barriers to gene flow between subpopulations of any of the three species studied here. Furthermore, although we observe elevated degrees of relatedness among foragers collected on the same mountain, there is no significant tendency for elevated relatedness of foragers caught on nearby mountains. These results indicate that populations of these species are effectively panmictic across the area surveyed, which has significance for their management. Foragers inferred to belong to the same nest (i.e. with identical parents) were always found in the same or adjacent sites, consistent with the short foraging distance (on average < 110m) inferred by (Geib et al., 2015) for these species. Previous studies have used microsatellites to determine genetic relatedness between individuals in order to survey colony density, foraging distance (Geib et al., 2015), and lineage survival (Carvell et al., 2017). Whole genome sequencing is now a

relatively inexpensive alternative, which, together with availability of a reference sequence

assembly, can provide an accurate measurement of kinship between samples.

Morphological change over time has already been documented in the species studied here,

with a significant reduction in tongue length observed since 1966 (Miller-Struttmann et al.,

2015). During this time period, there has also been a shift in species composition, with

shorter-tongued species becoming more predominant. Previous collections of *B. sylvicola* in

this region are likely a mixture of *B. sylvicola* and *B. incognitus*, as the existence of *B.*

incognitus was unknown. However, as *B. sylvicola* and *B. incognitus* do not differ in glossa

length, undetected changes in the relative abundance of these species over time are

unlikely to be responsible for the observed shifts in glossa length observed in samples that

are defined as *B. sylvicola.* This effect could however contribute to the observed variation in

body size over time as *B. sylvicola* have a significantly greater ID on average than *B.*

incognitus (Figure S3).

A major finding that we present here is that the temporal shift toward shorter tongue

length is ongoing. In addition to the significant reduction previously observed over a period

of six decades up until 2014, we also observe a significant reduction over a much shorter

time period between 2012-14 and 2017 in *B.sylvicola-incognitus*. Our results also indicate

that the shift has not only continued, but has accelerated compared to previous

measurements. Mean tongue length in *B. sylvicola* decreased on average by 0.61% annually

in a period between 1966 and 2014. According to our recent measurements, the

corresponding decrease in 3-5 subsequent years has been 12.9 %, or 3.2% per year

(compared to the 2012-14 average). These changes have been previously attributed to a

connection between shorter tongues and generalist foraging, a strategy which is favored at

lower floral density. It is suggested that these conditions have become more common in the

habitats of these species in the study area due to the effects of climate change (Miller-

Struttmann et al., 2015). We also observe significant differences in tongue length between

mountains, which could potentially reflect variation in these environmental factors between

mountains. In contrast to tongue length, ID has remained relatively constant despite some

fluctuations.

Our results are consistent with a hypothesis that tongue length changes are driven by

natural selection due to changes in flower abundance driven by climate change. There are

now many examples of rapid morphological evolution driven by natural selection related to

climatic events in the wild, such as a shift in beak shape over a period of a few years in

Darwin's finches related to competition for food during drought (Lamichhaney et al., 2016)

and genomic shifts related to cold-tolerance in the green anole lizard related to severe

winter storms (Campbell-Staton et al., 2017). Human activity may also promote rapid

adaptation by natural selection, as demonstrated by the evolution of longer beaks in great

tits promoted by the use of bird feeders in the UK (Bosse et al., 2017).

It is however unclear whether the observed changes in tongue length due to natural

selection acting on genetic variation. It is also possible that the changes in tongue length we

observe are due to plasticity. In order to address this question, we attempted to uncover

the genetic basis of morphological variation using genome-wide association studies (GWAS)

and genome-wide complex trait analysis (GCTA). No significant associations were observed

for tongue length in the genome of any of the species. However, a significant association

with body size was identified in *B. sylvicola* in the *CTRB1* (Chymotrypsinogen B1) gene. This

gene produces of a precursor of chymotrypsin, which is involved in proteolysis and likely

plays a role in digestion of pollen (Grogan & Hunt, 1979; Lazarević & Janković-Tomanić,

2015; Rawlings & Barrett, 1994). It is therefore highly plausible that variation in the activity

of this enzyme could influence body size, and intriguing that a variant of relatively large

effect is segregating in this natural population.

We detected a significant signal of covariance of trait measurements with relatedness using

GCTA when closely-related individuals were included in the dataset. However, no significant

signal of heritability in morphological traits was detected when the dataset was purged of

close relatives. The first result could potentially reflect a genetic component, but it is not

possible to disentangle this component from the shared environmental effects due to close

relatives sharing a nest. This analysis is therefore underpowered to detect genetic effects

due to low sample size and a general lack of related individuals in the dataset. We estimated

that a sample size 10-times larger would be necessary to estimate heritability with a

standard error below 0.1 (Visscher et al., 2014). For polygenic traits, such as body size and

tongue length, there are likely multiple QTL of small effect ($h^2 < 0.1$) throughout the genome

(Yang et al., 2010) and it is therefore not surprising that such loci were not detected. A study

of such a magnitude in a wild bumblebee population would require a huge sampling effort

that could have detrimental efforts on colony survival. A probable explanation for our

results is that the morphological traits have a highly polygenic genetic component, and

GWAS and GCTA were underpowered to detect QTL and estimate heritability. The size of

the genetic component of variation in both traits in each species is therefore still unknown.

In summary, we have shown here that morphological change under climate change is still

ongoing in alpine bumblebee populations at high-altitudes in Colorado. We generated the

genomic resources to study these species on the genetic level in the form of highly-

contiguous genome assemblies and comprehensively characterized genome variation. Our

results identified a previously-undescribed species and provided insights into population

structure and the genetic basis of morphological variation in these populations. Alpine

ecosystems are particularly sensitive to climate change and may suffer its effects before

other regions (Elsen & Tingley, 2015). Continued monitoring of changes in morphology and

species abundance of the populations under study here would be highly valuable to

understand how species elsewhere will adapt in future. Similar genomic studies of these

bumblebee subgenera in other mountain and artic locations would also be valuable in order

to fully characterize species diversity, including additional cryptic species, and to further

understand the ecological and genetic causes for morphological variation.

ACKNOWLEDGEMENTS

This research was funded by Swedish Research Council Formas grant 2016-00535 and a

SciLifeLab National Biodiversity Project NP00046 to MTW. MK is financially supported by the

Knut and Alice Wallenberg Foundation as part of the National Bioinformatics Infrastructure

Sweden at SciLifeLab. The authors acknowledge support from the National Genomics

Infrastructure in Stockholm and Uppsala funded by Science for Life Laboratory, the Knut and

Alice Wallenberg Foundation and the Swedish Research Council. The computations and data

handling were enabled by resources provided by the Swedish National Infrastructure for

Computing (SNIC) at UPPMAX partially funded by the Swedish Research Council through

grant agreement no. 2018-05973. U.S. Forest Service provided access to field sites. We

thank Jacqueline Staab and Michael Osbourn who aided in specimen collection,

identification and preservation and Lisa Danback, Kate Schleicher, Ying Thao, Anna Grobelny

and Breana Cook who carried out phenotypic measurements.

REFERENCES

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry
676 in unrelated individuals. *Genome Research*, *19*(9), 1655–1664. doi:
677 10.1101/gr.094052.109
678 Bosse, M., Spurgin, L. G., Laine, V. N., Cole, E. F., Firth, J. A., Gienapp, P., … Slate, J. (2017).
679 Recent natural selection causes adaptive evolution of an avian polygenic trait.
680 *Science*, *358*(6361), 365–368. doi: 10.1126/science.aal3298
681 Cameron, S. A., Lozier, J. D., Strange, J. P., Koch, J. B., Cordes, N., Solter, L. F., & Griswold, T.
682 L. (2011). Patterns of widespread decline in North American bumble bees.
683 *Proceedings of the National Academy of Sciences*, *108*(2), 662–667. doi:
684 10.1073/pnas.1014743108
685 Cameron, S. A., & Sadd, B. M. (2020). Global Trends in Bumble Bee Health. *Annual Review of*
686 *Entomology*, *65*, 209–232. doi: 10.1146/annurev-ento-011118-111847
687 Campbell-Staton, S. C., Cheviron, Z. A., Rochette, N., Catchen, J., Losos, J. B., & Edwards, S.
688 V. (2017). Winter storms drive rapid phenotypic, regulatory, and genomic shifts in
689 the green anole lizard. *Science*, *357*(6350), 495–498. doi: 10.1126/science.aam5512
690 Cane, J. H. (1987). Estimation of Bee Size Using Intertegular Span (Apoidea). *Journal of the*
691 *Kansas Entomological Society*, *60*(1), 145–147.
692 Cariveau, D. P., Nayak, G. K., Bartomeus, I., Zientek, J., Ascher, J. S., Gibbs, J., & Winfree, R.
693 (2016). The Allometry of Bee Proboscis Length and Its Uses in Ecology. *PLOS ONE*,
694 *11*(3), e0151482. doi: 10.1371/journal.pone.0151482
695 Carvell, C., Bourke, A. F. G., Dreier, S., Freeman, S. N., Hulmes, S., Jordan, W. C., … Heard, M.
696 S. (2017). Bumblebee family lineage survival is enhanced in high-quality landscapes.
697 *Nature*, *543*(7646), 547–549. doi: 10.1038/nature21709
698 Christmas, M. (submitted). *Genetic barriers to historical gene flow between cryptic species*
699 *of alpine bumblebees revealed by comparative population genomics*.
700 Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R.
701 (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. doi:
702 10.1093/bioinformatics/btr330
703 Elsen, P. R., & Tingley, M. W. (2015). Global mountain topography and the fate of montane
704 species under climate change. *Nature Climate Change*, *5*(8), 772–776. doi:
705 10.1038/nclimate2656
706 Estoup, A., Scholl, A., Pouvreau, A., & Solignac, M. (1995). Monoandry and polyandry in
707 bumble bees (Hymenoptera; Bombinae) as evidenced by highly variable
708 microsatellites. *Molecular Ecology*, *4*(1), 89–93. doi: 10.1111/j.1365-
709 294x.1995.tb00195.x
710 Geib, J. C., Strange, J. P., & Galen, C. (2015). Bumble bee nest abundance, foraging distance,
711 and host-plant reproduction: Implications for management and conservation.
712 *Ecological Applications*, *25*(3), 768–778. doi: https://doi.org/10.1890/14-0151.1

Goulson, D. (2003). *Bumblebees: Their Behaviour and Ecology*. Oxford University Press.

Goulson, D., Hanley, M. E., Darvill, B., Ellis, J. S., & Knight, M. E. (2005). Causes of rarity in
bumblebees. *Biological Conservation*, *122*(1), 1–8. doi: 10.1016/j.biocon.2004.06.017

Goulson, D., Lye, G. C., & Darvill, B. (2008). Decline and conservation of bumble bees.
Annual Review of Entomology, *53*, 191–208. doi:
10.1146/annurev.ento.53.103106.093454

Goulson, D., Nicholls, E., Botías, C., & Rotheray, E. L. (2015). Bee declines driven by
combined stress from parasites, pesticides, and lack of flowers. *Science (New York,*
N.Y.), *347*(6229), 1255957. doi: 10.1126/science.1255957

Grabherr, M. G., Russell, P., Meyer, M., Mauceli, E., Alföldi, J., Di Palma, F., & Lindblad-Toh,
K. (2010). Genome-wide synteny through highly sensitive sequence alignment:
Satsuma. *Bioinformatics*, *26*(9), 1145–1151. doi: 10.1093/bioinformatics/btq102

Grogan, D. E., & Hunt, J. H. (1979). Pollen proteases: Their potential role in insect digestion.
Insect Biochemistry, *9*(3), 309–313. doi: 10.1016/0020-1790(79)90011-8

Heraghty, S. D., Sutton, J. M., Pimsler, M. L., Fierst, J. L., Strange, J. P., & Lozier, J. D. (2020).
De Novo Genome Assemblies for Three North American Bumble Bee Species:
Bombus bifarius, Bombus vancouverensis, and Bombus vosnesenskii. *G3: Genes,*
Genomes, Genetics, *10*(8), 2585–2592. doi: 10.1534/g3.120.401437

ImageJ. (n.d.). Retrieved December 7, 2020, from https://imagej.nih.gov/ij/

Kent, C. F., Dey, A., Patel, H., Tsvetkov, N., Tiwari, T., MacPhail, V. J., … Zayed, A. (2018).
Conservation Genomics of the Declining North American Bumblebee Bombus
terricola Reveals Inbreeding and Selection on Immune Genes. *Frontiers in Genetics*,
9. doi: 10.3389/fgene.2018.00316

Kerr, J. T., Pindar, A., Galpern, P., Packer, L., Potts, S. G., Roberts, S. M., … Pantoja, A. (2015).
CLIMATE CHANGE. Climate change impacts on bumblebees converge across
continents. *Science (New York, N.Y.)*, *349*(6244), 177–180. doi:
10.1126/science.aaa7031

Lamichhaney, S., Han, F., Berglund, J., Wang, C., Almen, M. S., Webster, M. T., … Andersson,
L. (2016). A beak size locus in Darwins finches facilitated character displacement
during a drought. *Science*, *352*(6284), 470–474. doi: 10.1126/science.aad8786

Lazarević, J., & Janković-Tomanić, M. (2015). Dietary and phylogenetic correlates of
digestive trypsin activity in insect pests. *Entomologia Experimentalis et Applicata*,
157(2), 123–151. doi: 10.1111/eea.12349

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics*, *25*(14), 1754–1760. doi: 10.1093/bioinformatics/btp324

Li, Heng, Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … Durbin, R. (2009).
The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–
2079. doi: 10.1093/bioinformatics/btp352

Manichaikul, A., Mychaleckyj, J. C., Rich, S. S., Daly, K., Sale, M., & Chen, W.-M. (2010).
Robust relationship inference in genome-wide association studies. *Bioinformatics*,
26(22), 2867–2873. doi: 10.1093/bioinformatics/btq559

Marshall, L., Perdijk, F., Dendoncker, N., Kunin, W., Roberts, S., & Biesmeijer, J. C. (2020).
Bumblebees moving up: Shifts in elevation ranges in the Pyrenees over 115 years.
Proceedings. Biological Sciences, *287*(1938), 20202201. doi: 10.1098/rspb.2020.2201

Martinet, B., Lecocq, T., Brasero, N., Gerard, M., Urbanová, K., Valterová, I., … Rasmont, P.
(2019). Integrative taxonomy of an arctic bumblebee species complex highlights a

new cryptic species (Apidae: Bombus). *Zoological Journal of the Linnean Society*, *187*(3), 599–621. doi: 10.1093/zoolinnean/zlz041

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., … DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303. doi: 10.1101/gr.107524.110

Miller-Struttmann, N. E., Geib, J. C., Franklin, J. D., Kevan, P. G., Holdo, R. M., Ebert-May, D., … Galen, C. (2015). Functional mismatch in a bumble bee pollination mutualism under climate change. *Science*, *349*(6255), 1541–1544. doi: 10.1126/science.aab0868

Pauls, S. U., Nowak, C., Bálint, M., & Pfenninger, M. (2013). The impact of global climate change on genetic diversity within populations and species. *Molecular Ecology*, *22*(4), 925–946. doi: https://doi.org/10.1111/mec.12152

Rawlings, N. D., & Barrett, A. J. (1994). Families of serine peptidases. *Methods in Enzymology*, *244*, 19–61. doi: 10.1016/0076-6879(94)44004-2

Razgour, O., Forester, B., Taggart, J. B., Bekaert, M., Juste, J., Ibáñez, C., … Manel, S. (2019). Considering adaptive genetic variation in climate change vulnerability assessment reduces species range loss projections. *Proceedings of the National Academy of Sciences*, *116*(21), 10418–10423. doi: 10.1073/pnas.1820663116

Ruan, J., & Li, H. (2020). Fast and accurate long-read assembly with wtdbg2. *Nature Methods*, *17*(2), 155–158. doi: 10.1038/s41592-019-0669-3

Sadd, B. M., Barribeau, S. M., Bloch, G., Graaf, D. C. de, Dearden, P., Elsik, C. G., … Worley, K. C. (2015). The genomes of two key bumblebee species with primitive eusocial organization. *Genome Biology*, *16*(1), 76. doi: 10.1186/s13059-015-0623-3

Sheffield, C. S., Oram, R., & Heron, J. M. (2020). Bombus (Pyrobombus) johanseni Sladen, 1919, a valid North American bumble bee species, with a new synonymy and comparisons to other "red-banded" bumble bee species in North America (Hymenoptera, Apidae, Bombini). *ZooKeys*, *984*, 59–81. doi: 10.3897/zookeys.984.55816

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, *31*(19), 3210–3212. doi: 10.1093/bioinformatics/btv351

Soroye, P., Newbold, T., & Kerr, J. (2020). Climate change contributes to widespread declines among bumble bees across continents. *Science*, *367*(6478), 685–688. doi: 10.1126/science.aax8591

Sun, C., Huang, J., Wang, Y., Zhao, X., Su, L., Thomas, G. W. C., … Mueller, R. L. (2020). Genus-wide characterization of bumblebee genomes provides insights into their evolution and variation in ecological and behavioral traits. *Molecular Biology and Evolution*, (msaa240). doi: 10.1093/molbev/msaa240

Visscher, P. M., Hemani, G., Vinkhuyzen, A. A. E., Chen, G.-B., Lee, S. H., Wray, N. R., … Yang, J. (2014). Statistical Power to Detect Genetic (Co)Variance of Complex Traits Using SNP Data in Unrelated Samples. *PLOS Genetics*, *10*(4), e1004269. doi: 10.1371/journal.pgen.1004269

Wang, M., & Xu, S. (2019). Statistical power in genome-wide association studies and quantitative trait locus mapping. *Heredity*, *123*(3), 287–306. doi: 10.1038/s41437-019-0205-3

Williams, P. H., Berezin, M. V., Cannings, S. G., Cederberg, B., Ødegaard, F., Rasmussen, C., …
Byvaltsev, A. M. (2019). The arctic and alpine bumblebees of the subgenus
Alpinobombus revised from integrative assessment of species' gene coalescents and
morphology (Hymenoptera, Apidae, Bombus). *Zootaxa*, *4625*(1), zootaxa.4625.1.1.
doi: 10.11646/zootaxa.4625.1.1
Yang, J., Benyamin, B., McEvoy, B. P., Gordon, S., Henders, A. K., Nyholt, D. R., … Visscher, P.
M. (2010). Common SNPs explain a large proportion of the heritability for human
height. *Nature Genetics*, *42*(7), 565–569. doi: 10.1038/ng.608
Yang, J., Lee, S. H., Goddard, M. E., & Visscher, P. M. (2011). GCTA: A Tool for Genome-wide
Complex Trait Analysis. *The American Journal of Human Genetics*, *88*(1), 76–82. doi:
10.1016/j.ajhg.2010.11.011
Yang, J., Zaitlen, N. A., Goddard, M. E., Visscher, P. M., & Price, A. L. (2014). Advantages and
pitfalls in the application of mixed-model association methods. *Nature Genetics*,
46(2), 100–106. doi: 10.1038/ng.2876
819
820
821 **DATA ACCESSIBILTY**
822
823 All sequence data is associated with NCBI project IDs: PRJNA646847 and PRJNA704506
824
825 **AUTHOR CONTRIBUTIONS**
826
827 Designed research: MJC, NEM-S, JCG, MTW; Performed research: all authors; Analyzed data:
828 MJC, JCJ, OW, IB, MK, MTW; Wrote the paper: MJC, MTW.

Table 1: Assembly metrics for the two genomes, with two other published *Bombus* genomes included for comparison

Species	*B. balteatus*	*B. sylvicola*	*B. terrestris*	*B. impatiens*
Assembly	BBAL_1.0	BSYL_1.0	Bter_1.0	BIMP_2.2
Size (Mbp)†	250.07	252.08	248.65 (236.38)	246.86 (241.98)
Scaffolds (n)	-	-	5,678	5,460
Contigs (n)	336	592	10,672	16,060
Contig N50 (Mbp)	8.60	3.02	0.08	0.06
Contig L50 (n)	12	28	890	54
GC (%)	37.64	38.23	37.51	37.76
Repeat content (%)	17.1	17.9	14.8	17.9
Complete hymenoptera BUSCO genes (%)*	99.0	98.2	96.9	98.3

*BUSCO analysis using the OrthoDB v. 10, Hymenoptera dataset.
† Numbers in brackets represent total ungapped length

Figure 1. Population sampling and genetic variation in three *Bombus* species. (A) Map showing the sampling locations of *Bombus balteatus, B. sylvicola,* and *B. incognitus* on seven mountains in the Rocky Mountains, CO, USA. Insert indicates location in the USA. Neighbour-joining trees of (B) *B. balteatus* and (C) *B. sylvicola* and *B. incognitus* populations sampled across the seven mountains, based on genome-wide single nucleotide polymorphisms. Coloured tips indicate the mountain each sample was collected on, with colours corresponding to those in (A).

Figure 2. Principal component analysis of Rocky Mountain populations of three *Bombus* species. Plots show the first two principal components from principal component analyses of (A) *Bombus balteatus*, (B) *B. sylvicola* and (C) *B. incognitus* populations. Numbers in brackets on axes indicate the percentage variance explained by each principal component. Samples in clusters indicated by ellipses in (A) and (C) are full sisters.

Figure 3. Mantel correlograms showing correlations between kinship and geographic distance in different distance classes. The Mantel correlation between a kinship matrix and a geographic distance matrix (y-axis) is calculated over 15 distance classes (x-axis, in metres) for (A) *Bombus balteatus,* (B) *B. sylvicola*, and (C) *B. incognitus*. Filled in boxes indicate significant correlations (p < 0.01). In all three cases, there is a significant negative correlation in the first distance class (< 10 km), indicating the shorter the distance the higher the relatedness over short distances. This reflects the fact that nest mates forage over short distances. At greater distance classes, the correlations are mostly nonsignificant (open boxes). Distances over 60 km had insufficient data points for calculating Mantel correlations.

Figure 4. Comparisons of glossa lengths between species, years, and locations. (A) Least-squares mean (ls-mean) glossa lengths correcting for body size for the three *Bombus* species, based on measurements of the 2017 collections. Least-squares mean glossa lengths per sampling period for (B) *B. balteatus* and (C) *B. sylvicola* and *B. incognitus* combined based on measurements made from samples collected at the same locations in each time period. These locations are shown in (D) for *B. balteatus* and (E) for *B. sylvicola-incognitus*, where the samples have been separated out per location. In all plots, error bars represent ls-means ± standard error and significant differences between classes are indicated by asterisks (p < 0.05).

Figure 5. Genome-wide association Manhattan plot. The -log$_{10}$ p-values from an association test using a mixed linear model between allele frequencies and intertegular distance (ID) are plotted for 2,705,361 genome-wide SNPs against their genome positions over the 18 *Bombus* chromosomes. Red dashed line indicates genome-wide significance after Bonferroni correction for p = 0.01. A single SNP on chromosome 10, coloured red, had a -log$_{10}$ p-value above this threshold. Green dashed line indicates genome-wide significance after Bonferroni correction for p = 0.05. Two neighbouring SNPs located on chromosome 14, coloured green, had -log$_{10}$ p-values above this threshold.

Figure 6. Estimates of SNP-heritability for two traits in *Bombus* species. The proportion of variance in glossa length and intertegular distance (ID) explained by all genome-wide SNPs (the SNP-based heritability) was calculated using the genome-base restricted maximum likelihood (GREML) method in GCTA. These were calculated separately for *B. sylvicola* (left panel) and *B. balteatus* (right panel). We included ID as a covariate with glossa length to control for body size. Error bars indicate 95% confidence intervals around estimates. Each analysis was carried out twice, once with all samples (red), and a second time with one from each pair of sisters removed (black), in order to assess the effect of including highly related samples in the analysis.