A genomic and morphometric analysis of alpine bumblebees: ongoing reductions in

tongue length but no clear genetic component

Running title: Genome variation in alpine bumblebees

Matthew J. Christmas[1], Julia C. Jones[1,2], Anna Olsson[1], Ola Wallerman[1], Ignas Bunikis[3],

Marcin Kierczak[4], Kaitlyn M. Whitley[5,6], Isabel Sullivan[5,7], Jennifer C. Geib[5], Nicole E. Miller-

Struttmann[8], Matthew T. Webster[1]

1) Department of Medical Biochemistry and Microbiology, Science for Life Laboratory,

Uppsala University, Uppsala, Sweden

2) School of Biology and Environmental Science, University College Dublin, Dublin, Ireland

3) Department of Immunology, Genetics and Pathology, Science for Life Laboratory, Uppsala

University, Uppsala, Sweden

4) Department of Cell and Molecular Biology, National Bioinformatics Infrastructure

Sweden, Science for Life Laboratory, Uppsala University, Uppsala, Sweden

5) Department of Biology, Appalachian State University, Boone, North Carolina, USA

6) U.S. Department of Agriculture, Agriculture Research Service, Charleston, South Carolina,

USA

7) Marine Estuarine Environmental Sciences, University of Maryland, College Park Maryland

8) Biological Sciences Department, Webster University, St. Louis, Missouri, USA

Corresponding author: Matthew T. Webster

Address: Dept. Medical Biochemistry and Microbiology, Box 582, 75123 Uppsala, Sweden

Email: matthew.webster@imbim.uu.se

Tel: +46 18 471 4391

ABSTRACT

Over the last six decades, populations of the bumblebees *Bombus sylvicola* and *Bombus balteatus* in Colorado have experienced decreases in tongue length, a trait important for plant-pollinator mutualisms. It has been hypothesized that this observation reflects selection resulting from shifts in floral composition under climate change. Here we used morphometrics and population genomics to determine whether morphological change is ongoing, investigate the genetic basis of morphological variation, and analyze population structure in these populations. We generated a genome assembly of *B. balteatus*. We then analyzed whole-genome sequencing data and morphometric measurements of 580 samples of both species from seven high-altitude localities. Out of 281 samples originally identified as *B. sylvicola*, 67 formed a separate genetic cluster comprising a newly-discovered cryptic species ("incognitus"). However, an absence of genetic structure within species suggests that gene flow is common between mountains. We found a significant decrease in tongue length between bees collected between 2012-2014 and in 2017, indicating that morphological shifts are ongoing. We did not discover any genetic associations with tongue length, but a SNP related to production of a proteolytic digestive enzyme was implicated in body size variation. We identified evidence of covariance between kinship and both tongue length and body size, which is suggestive of a genetic component of these traits, although it is possible that shared environmental effects between colonies are responsible. Our results provide evidence for ongoing modification of a morphological trait important for pollination and indicate that this trait likely has a complex genetic and environmental basis.

INTRODUCTION

The importance of pollinators for ecology and the maintenance of biodiversity is widely recognized (Goulson, Nicholls, Botías, & Rotheray, 2015). Bumblebees are large-bodied, mainly cold-adapted species with broad significance for agricultural production and plant-pollinator networks in the wild (Goulson, 2003). Many bumblebee species are experiencing range shifts or declines due to climate change or habitat loss (Cameron et al., 2011; Cameron & Sadd, 2020; Goulson, Lye, & Darvill, 2008a; Soroye, Newbold, & Kerr, 2020). In particular, a general worldwide trend for species to shift to higher elevations and latitudes has been documented, with losses occurring at a faster rate than expansions (Kerr et al., 2015; Marshall et al., 2020). Arctic and alpine species are particularly threatened by these

trends, and can be considered "canaries in the coal mine" for detecting early effects (Elsen

& Tingley, 2015).

In addition to range shifts there is evidence that the morphology of certain populations of

bumblebees has shifted in recent decades. Miller-Struttmann et al. (2015) compared

morphometric measurements of contemporary samples and historical specimens of the

species *B. balteatus* and *B. sylvicola* from the same high-elevation alpine localities in

Colorado. These species historically comprise the majority of bumblebees in these locations

(Miller-Struttmann et al., 2015). They reported a significant decrease in tongue length in

both species in a period between 1966 and 2014. The cause of this was inferred to be a

general decrease in floral abundance resulting from warmer temperatures and drier soils

resulting from climate change. These conditions could favor more generalist foraging

strategies, which several studies indicate are performed more efficiently by bees with

shorter tongues (Goulson & Darvill, 2004; Goulson, Hanley, Darvill, Ellis, & Knight, 2005;

Goulson, Lye, & Darvill, 2008b; Heinrich, 2004; Huang, An, Wu, & Williams, 2015). Continued

monitoring of these populations is important to understand the causes of these

morphological shifts and their ecological consequences. It is unknown whether the trend

towards shorter tongues has continued in more recent years.

One explanation for the observed morphological shifts in *B. balteatus* and *B. sylvicola* in

Colorado is that natural selection for shorter tongues has occurred (Miller-Struttmann et al.,

2015). However, it also possible that the observed changes are a plastic response to

environmental change (Merilä & Hendry, 2014). It is unknown whether this trait has a

genetic component that could be acted on by natural selection. Uncovering the size of the

genetic contribution to this trait and the number and identity of any genetic loci involved

would be an important step towards understanding the underlying cause of the

morphological changes. The feasibility of genome sequencing on a population scale has

enabled studies of quantitative genetics (Gienapp et al., 2017) and genome-wide association

studies (Santure & Garant, 2018) in wild populations, allowing these goals to be pursued.

The genetic structure of populations is an important factor for interpretation of genotype-

phenotype correlations and the cause of morphological shifts. We previously presented a

genome assembly of *B. sylvicola* and analyzed genome-wide variation by resequencing 281

samples (Christmas et al., 2021). This identified the presence of a previously-undetected

cryptic species, which we gave the preliminary name *Bombus incognitus*. We refer to this

species hereafter simply as "incognitus" to reflect the fact that a formal taxonomic

description is not currently available. This species was identified morphologically as *B.*

sylvicola but forms a distinct genetic cluster. It is therefore possible that changes in tongue

length in the combined populations of these species could be brought about by changes in

the relative abundance of each species, although this has not previously been investigated.

Additional population structure within species could lead to a similar effect, particularly if

genetic structure associated with morphology exists. An understanding of genetic structure

and spatial connectivity between populations in different locations is also needed to predict

how they will evolve under climate change and to define the conservation value of

subpopulations (Pauls, Nowak, Bálint, & Pfenninger, 2013; Razgour et al., 2019).

The degree of gene flow in bumblebee populations is determined by the distances that

reproductive individuals (males and queens) travel in order to mate and establish new nests

(Heinrich, 2004; Woodard et al., 2015). Genetic methods have been used to investigate

population structure of several species of bumblebees across a variety of habitats in the UK,

continental Europe and continental USA (Darvill, Ellis, Lye, & Goulson, 2006; Ellis, Knight,

Darvill, & Goulson, 2006; Ghisbain et al., 2020; Jackson et al., 2018; Jha, 2015; Koch, Looney,

Sheppard, & Strange, 2017; Woodard et al., 2015). A general finding is that populations of

common species tend to exhibit very little structure in the absence of geographical barriers,

even at continental scales. However, genetic differentiation can occur in rare or declining

species, or when there are natural barriers (Woodard et al., 2015). Populations restricted to

high elevations have been shown to exhibit higher genetic differentiation (Lozier, Strange, &

Koch, 2013; Lozier, Strange, Stewart, & Cameron, 2011). However, the degree of

fragmentation of populations of bumblebee species inhabiting high elevations in mountain

ranges is not fully understood.

Genetic methods can also be applied to estimate nest density and foraging distances of

bumblebees, by analyzing where genetically-related nestmates are caught. Such studies

indicate that maximum foraging distances range from less than 100 m to over 10 km

(Darvill, Knight, & Goulson, 2004; Geib, Strange, & Galen, 2015; Jha & Kremen, 2013; Knight

et al., 2005; Woodard et al., 2015). These variables show large variation among species and

are strongly dependent on landscape composition. Genome sequencing of multiple

individuals has the potential to add greater resolution to our understanding of population

density, structure and dispersal.

In this study we present a high-quality annotated genome assembly of the bumblebee

species *B. balteatus* based on long-read Oxford Nanopore sequencing. We also conducted

whole-genome sequencing of 299 samples of this species sampled from seven high-altitude

localities spread across 150 km in Colorado. These localities have an average separation of

54 km between each other, and consist of alpine tundra separated by forest. We analyze

these data together with the previously-published population genomic dataset of 284

samples of *B. sylvicola* and "incognitus" from the same localities (Christmas et al., 2021). We

use these data to assess whole-genome variation and its connection to geographical and

morphological variation in these populations, which were previously studied by Miller-

Struttmann et al. (2015). We then analyzed morphological variation in tongue length and

body size in context of previous studies (Miller-Struttmann et al., 2015) to determine

whether the trend towards decreasing tongue length is ongoing. We also performed a

genome-wide analysis of the genetic basis of morphological variation using both genome-

wide association studies (GWAS) and genome-wide complex trait analysis (GCTA) (Yang, Lee,

Goddard, & Visscher, 2011). The results are informative regarding the degree of connectivity

among bumblebee populations, the genetic basis of morphological variation, and the

mechanisms of morphological evolution.

Our study species have recently been subject to taxonomic revision. The species we refer to

here as *B. balteatus* was previously described as occurring in both North America and

Eurasia, but a recent study classified these geographically separated populations as two

different species: *B. balteatus* in Eurasia and *Bombus kirbiellus* in North America (Williams et

al., 2019). The species we refer to here as *B. sylvicola* was previously split into a North

American form (*B. sylvicola*) and a Eurasian form (*Bombus lapponicus*). However, a recent

analysis synonymized these two species as *B. lapponicus* (Martinet et al., 2019). We

continue to use the previous names here to avoid confusion with comparison to previous

ecological and evolutionary genetic studies.

MATERIALS AND METHODS

Sample collection

We collected samples identified as *B. balteatus* and *B. sylvicola* (Koch, Strange, & Williams,

2014) during the summer of 2017. Bumblebees were collected on seven different

mountains in Colorado, USA, with an average distance of 54 km between mountains

(greatest distance: 134 km, Quail Mountain – Niwot Ridge; shortest distance: 10km, Mount

Democrat – Pennsylvania Mountain). The existence of "incognitus" was unknown to us

when the bees were sampled but a large fraction of bees identified as *B. sylvicola* were

subsequently assigned to "incognitus" on the basis of genetic analysis (Christmas et al.,

2021). Three of these mountains (Mount Evans, Niwot Ridge, and Pennsylvania Mountain)

were also sampled previously by Miller-Struttmann et al. (2015). Four additional mountains

were sampled here (Boreas Mountain, Mount Democrat, Horseshoe Mountain and Quail

Mountain) but were not sampled in the previous study. Each of the mountain locations

comprised several neighboring sampling sites separated by short distances (mean = 1.1 km)

and elevations (range: 3,473 – 4,012 m.a.s.l.) (full details in supplementary tables S1, S2).

We collected samples of foraging worker bees from each site using sweeping hand nets. We

kept samples in Falcon tubes in cool boxes for transport. Species assignment was performed

by observing standard morphological characters (Koch et al., 2014). We dissected each bee

after placing it at -20°C for approximately 10 minutes.

Morphological measurements

We measured intertegular distance and tongue length on all of the samples of *B. sylvicola,*

"incognitus" and *B. balteatus* using scaled photographs of individual bee heads processed by

ImageJ (Kearns & Inouye, 1993; Schneider, Rasband, & Eliceiri, 2012). Following netting,

bees were anesthetized by chilling them on ice or in a freezer (depending on field

conditions). Once anesthetized, the bees were removed from the vial, gently pressed prone

to the table with the thorax parallel to the camera lens using flat forceps, and photographed

with a metric ruler for scaling purposes. The distance between the tegulae was measured as

a straight line between the edge of one tegula to the other tegula, running as near to

perpendicular to the tegula as possible. The head was then immediately removed from the

body and placed into an individual labeled vial and kept frozen until further characterization

in the lab. Following the protocol used in Miller-Struttmann *et al.* (2015), the prementum

and glossa were dissected from the head while the tissue was flexible, straightened and

aligned as possible without damaging the tissue, and mounted on acid-free paper. The

samples were then photographed with a ruler in the field of view under a microscope, and

the traits measured (i.e., prementum length and glossa length) via ImageJ (Schneider et al.,

2012). After photographing, we stored thoraces in 95% ethanol prior to DNA extraction. We

visually characterized the diagnostic traits visible on each head, such as the shape of the

malar space, the pile color between and antenna and above the ocelli, the size, color and

location of the ocelli relative to the supraorbital line, and the presence of black pile on the

scutellum. None of the morphological characters we measured could be used to distinguish

between "incognitus" and *B. sylvicola*.

Genome sequencing and assembly

We used the same sequencing strategy to produce high-quality genome assemblies for both

B. sylvicola and *B. balteatus*. The *B. sylvicola* assembly was presented previously (Christmas

et al., 2021). This involved combining data from Oxford Nanopore (ONT) and 10x Genomics

Chromium technologies. For both species, we used DNA extracted from male bees sampled

on Niwot Ridge and stored in 95% ethanol prior to DNA extraction. DNA from a single bee

was used for each assembly. We performed DNA extraction using a salt-isopropanol

extraction followed by size selection using magnetic bead purification to remove fragments

< 1 kbp. ONT sequencing was performed using the MinION instrument. For *B. sylvicola*, we

used two R9.4 flowcells and the RAD004 kit starting with 3-400 ng DNA per run. This

resulted in a total of 9.4 Gbp of sequence data, in 2.5 million reads with a mean length of

3.7 kbp (Christmas et al., 2021). For *B. balteatus*, we used 1µg DNA in a modified LSK-108

one-pot protocol. 1.75 µl NEB FFPE buffer, 1.75 µl NEB Ultra-II End Prep buffer and 1 µl FFPE

repair mix was added to 24 µl DNA and incubated at 20°C for 30 minutes. 2 µl End Prep

enzyme mix was then added for blunting, phosphorylation and A-tailing (20 min. 20°C, 10

min. 65°C, 10 min. 70°C). AMX (20 µl), ligation enhancer (1 µl) and Ultra-II ligase (40 µl) was

added for ligation at room temperature for 30 minutes. Clean-up and flowcell loading was

done following the standard ONT LSK-108 protocol and the library was sequenced on one

R9.4 flowcell. This resulted in 23.3 Gpb of sequence data, in 3.9 million reads with a mean

length of 5.9 kbp.

We assembled the ONT reads using the following process: We first used downpore

(Teutenberg, 2018/2020) with default parameters for adaptor trimming and splitting

chimeric reads. We then assembled the trimmed reads using wtdbg2 (Ruan & Li, 2020) on

default settings. We then ran two rounds of the consensus module Racon (Vaser, Sović,

Nagarajan, & Šikić, 2017) and contig improvements using medaka v.0.4 (*Medaka*,

2017/2021), removing contigs of length < 20 kbp. We used Illumina short-read data from

the population sequencing to perform two rounds of Pilon (Walker et al., 2014) to improve

sequence accuracy, particularly around indels.

We also sequenced the same sample of each species using 10x Genomics Chromium. A 10x

GEM library was constructed according to the manufacturer's protocols. We normalized

each library using qPCR and sequenced them together on a single lane of HiSeq 2500 using

the HiSeq Rapid SBS sequencing kit version 2, resulting in 150 bp paired-end sequences. The

linked reads were mapped to the assembly using Longranger v.2.1.4

(*10XGenomics/Longranger*, 2018/2021). We then ran Tigmint v1.1.2 (Jackman et al., 2018)

to correct errors in the assembly detected by the read mappings. We identified contigs that

contained mitochondrial genes, and were therefore likely fragments of the mitochondrial

genome, by running a BLAST search *of B. impatiens* mitochondrial genes across the

assembly using BLAST+ v2.9.0 (Camacho et al., 2009). Any contigs containing two or more

mitochondrial genes located within the expected distance of each other based on their

locations on the mitochondrial genome were removed from the assembly to ensure the

final assembly did not contain partially assembled mitochondrial genome sequence. We also

removed all contigs shorter than 10 kbp.

Genome analysis

BUSCO v3.0.2b (Simão, Waterhouse, Ioannidis, Kriventseva, & Zdobnov, 2015) was used to

assess the assembly completeness using the hymenoptera_odb9 lineage set and species *B.*

impatiens. We ran RepeatMasker v.4.1.0 (Smit, Hubley, & Green, 2015) on each genome

assembly to characterize genome repeat content, including interspersed repeats and low

complexity sequences. Chromosome-level assemblies of species from five diverse

bumblebee subgenera indicate that non-parasitic clades have a stable 18 chromosome

karyotype (Sun et al., 2020). We therefore performed whole-genome synteny alignments

between the *B. terrestris* chromosome-level genome assembly (Sadd et al., 2015)

(downloaded from NCBI; BioProject PRJNA45869) and our assemblies using Satsuma v.3

(Grabherr et al., 2010) to arrange the contigs from our assemblies onto 18

pseudochromosomes.

RNAseq and Annotation

We used the Nextflow pipelines available at https://github.com/NBISweden/pipelines-

nextflow for genome annotation. The pipelines, in turn, depend on two additional

annotation-specific toolkits, AGAT and GAAS, available

at https://github.com/NBISweden/AGAT and https://github.com/NBISweden/GAAS

respectively. Both repositories contain extensive documentation and installation

instructions. As reference proteins database, we used the SwissProt/UniProt (561,356

curated proteins, downloaded 2019-11). The parameter files stated below are available at

https://github.com/matt-webster-lab/alpine_bumblebees.

We performed RNAseq on a single lane of Illumina HiSeq 2500 using samples from four body

parts from a single sample of *B. balteatus* (abdomen, head, legs and thorax), which

generated 42 Gbp of reads. We performed quality control of the data using FastQC

(Andrews, 2017/2021) and read trimming using trimmomatic (v. 0.39,

trimmomatic.params). We next generated a guided assembly with Hisat2 (Kim, Paggi, Park,

Bennett, & Salzberg, 2019) and Stringtie (Pertea et al., 2015) using the

AnnotationPreprocessing.nf (parameters as in AnnotationPreprocessing_params.conf)

TranscriptAssembly.nf Nextflow pipeline with parameters as in the

TranscriptAssembly_params.config file. In addition to the guided assembly, we performed a

de novo assembly for reads from each of the four tissues using Trinity v. 2.0.4 (Grabherr et

al., 2011) (--seqType fq --SS_lib_type RF).

Next, we constructed a species-specific repeat library using RepeatModeler package (v.

open-1.0.8; -engine ncbi -pa 35). Since the de novo identified repeats may still include parts

of protein-coding genes, we perform an additional filtering step and remove from the

database the repeats that can be found in a comprehensive set of known proteins. We

remove all repeats that match to a pre-computed set non-transposable proteins from our

protein reference database. We used custom script (filter_repeats.bash) for filtering. To

identify the location of repeats from this library in the genomes, we used RepeatMasker

(open-4.0.9).

We used the MAKER (Cantarel et al., 2008) package to compute gene builds. An evidence-

guided gene build was computed by MAKER (-fix-nucleotides) constructing gene models

from these reference proteins and from the transcript sequences aligned to the genome

assembly (parameters in evidence_build.maker_opts.ctl). We also prepared an *ab initio*

build using the AbInitioTraining.nf pipeline and AbInitioTraining_params.config as the

parameters file. Both the evidence-guided and the ab initio builds were than merged using

MAKER (hybrid_build_maker_opts.ctl). Finally, we performed functional annotation using

the FunctionalAnnotation.nf pipeline with FunctionalAnnotation_params.config as the

parameter file.

Population sequencing, read mapping and variant calling

We extracted DNA from the thoraces of 299 worker bees of *B. balteatus* collected from

across the seven sampling sites using the Qiagen Blood and Tissue kit. We prepared dual-

indexed libraries using the Nextera Flex kit and performed sequencing on an Illumina HiSeq

X to produce 2 x 150 bp reads, using an average of 36 samples per lane. We mapped reads

to the *B. balteatus* reference using the mem algorithm in BWA (Li & Durbin, 2009). We

performed sorting and indexing of the resultant bam files using samtools (Li et al., 2009) and

marked duplicate reads using Picard. We used the Genome Analysis Toolkit (GATK) to call

variants (McKenna et al., 2010). We first ran HaplotypeCaller using default parameters on

the bam file of each sample to generate a gVCF file for each sample. We then used

GenomicsDBImport and GenotypeGVCFs with default parameters to call variants for all *B.*

balteatus samples together. We applied a set of hard filters using the VariantFiltration tool

to filter for reliable SNPs using the following thresholds: $QD < 2$, $FS > 60$, $MQ < 40$,

MQRankSum < -12.5, ReadPosRankSum < -8 (see the GATK documentation for full

descriptions of each filter). Only biallelic SNPs were considered for downstream analysis.

Samples from *B. sylvicola* and "incognitus" samples were analyzed in the same way and are

presented in Christmas *et al.* (2021).

Relatedness and population structure

We estimated the pairwise kinship coefficient between all samples within each species using

the method described in (Manichaikul et al., 2010) implemented in the --relatedness2

option of vcftools v0.1.16 (Danecek et al., 2011). Worker bees from the same bumblebee

colony are expected to have a relatedness of 0.75 under the assumption of monandry,

which is reasonable for the study species (Estoup, Scholl, Pouvreau, & Solignac, 1995). The

kinship coefficient (φ) predicted from this relationship is $\frac{3}{8}$ (the probability that two alleles

sampled at random from two individuals are identical by descent). We estimated the lower

cutoff for the inference criteria to infer this level of relatedness using powers of two as

described in (Manichaikul et al., 2010), providing an estimate of $\frac{3}{2^{7/2}}$.

We tested for correlations between relatedness and geographic distance for each species

using Mantel tests with the mantel function in the 'ecodist' package in R v4.0.2. Coordinates

for each sample were converted to geographic distances using the R package 'geodist' and

these were compared to the matrix of kinship coefficients described above. We used the

mantel.correlog function to produce mantel correlograms and assess whether correlations

were significant over different geographic distances. We performed these analyses after

excluding sisters from the datasets to ensure any patterns we observe were not driven by

an effect of sampling individuals from the same colony.

We thinned the VCF files to 1 SNP every 10 kbp using the --max-missing 1 and --thin flags in

vcftools v.0.1.16 to ensure no missing data and to assess population structure and

relatedness more efficiently. We calculated p-distance matrices for *B. balteatus* samples and

B. sylvicola and "incognitus" samples combined from the thinned vcf files using VCF2Dis

(https://github.com/BGI-shenzhen/VCF2Dis) and then constructed neighbor-joining trees

using the 'ape' package in R v4.0.2. We converted the thinned vcf files to plink format and

performed principal components analyses using the 'adegenet' package in R v4.0.2. We

performed PCAs on all samples as well after removing one of each pair of sisters identified

via the kinship analysis to ensure that relatedness did not heavily influence any clustering.

We used admixture v1.3.0 (Alexander, Novembre, & Lange, 2009) to assess the most likely

number of genetic clusters within each species. We ran admixture for K=1 through to 10,

with 10 iterations per K value and used the calculated cross-validation (cv) errors to assess

the most suitable values of K, with low cv-error indicating higher support.

Statistical morphometric analysis

We combined the phenotypic measurements made in this study with those from previous

collections published in Miller-Struttmann et al. (2015). These measurements are

intertegular distance, glossa length, prementum length, and total tongue length (equal to

glossa length plus prementum length). This gave us three time periods to compare amongst:

'2017' (samples collected for this study), '2012-2014', and '1966-1980' (both sets published

in Miller-Struttmann et al., 2015). We performed statistical analysis of variation in glossa

length, prementum length, and intertegular distance in R v4.0.2. We investigated

correlations between all traits for each species in the 2017 set using the 'cor.test' function,

which calculates a Pearson's product moment correlation coefficient.

Within our 2017 dataset, we compared glossa lengths between the three species by fitting

linear models of species against glossa length with intertegular distance included as a

random effect to account for body size using the lm function. We then calculated least-

squares mean glossa lengths from this model using the lsmeans function and tested for

significant differences using the Tukey method for p-value adjustment for multiple testing.

We employed a similar approach for comparing glossa lengths between our dataset and

historical datasets for each species. Here we used a linear mixed effect model framework

with the 'lmer' function in the lme4 package in R v4.0.2, where mountain and intertegular

distance were included as random effects in each model. Significance of each model was

assessed using analysis of variance with the 'anova' function. We calculated least-squares

mean glossa lengths from each model using the lsmeans function and tested for significant

differences using the Tukey method for p-value adjustment for multiple testing.

We analyzed variation in intertegular distance between species and geographic locations for

our 2017 dataset using analysis of variance ('aov' function) and assessed significance of

comparisons using Tukey's honestly significant difference (HSD) post hoc test ('TukeyHSD'

function). For comparisons of intertegular distance between years, we fitted linear mixed

effect models and included mountain as a random effect, then compared least-squares

means from the models using the Tukey method for p-value adjustment for multiple testing.

Genome-wide association study

We carried out mixed linear model-based association analyses, implemented in GCTA-

MLMA (Yang et al., 2011; Yang, Zaitlen, Goddard, Visscher, & Price, 2014), to identify

correlations between allele frequencies and trait variation across the genome for *B.*

balteatus and *B. sylvicola* samples separately ("incognitus" samples were left out of this

analysis due to too few samples). First, we created a genetic relationship matrix (GRM) from

the genome-wide SNPs using the --make-grm function in GCTA. We removed one sample

from each pair of sisters by setting a relatedness cut-off of 0.67 using the '--grm-cutoff' flag.

We ran the analyses for three traits per species: intertegular distance (ID), glossa length

(GL), and the residuals from a linear regression of GL~ID to control for body size. The GRM

was included as a polygenic/random effect. The analysis assumes phenotypes are normally

distributed and so samples were removed in the tails of the distributions to meet this

assumption for all traits except intertegular distance in *B. sylvicola*, which was normally

distributed. For *B. balteatus*, this resulted in 279, 291, and 262 samples for the glossa

length, intertegular distance, and residuals analyses respectively. For *B. sylvicola* there were

176, 192, and 152 samples for the glossa length, intertegular distance, and residuals

analyses respectively. We assessed divergence of p-values from neutral expectations using

QQ-plots and carried out Bonferroni correction of p-values using the 'p.adjust' function in R

v4.0.2 to assess genome-wide significance.

Genome-wide complex trait analysis

We used GCTA (Yang et al., 2011) to perform genome-wide complex trait analysis. We

estimated the phenotypic variance explained by the genome-wide SNPs using genome-

based restricted maximum likelihood (GREML). The analysis was performed separately for

each species on measures of intertegular distance, glossa length, and glossa length with

intergular distance included as a covariate to control for body size. We ran each analysis

twice, once using a GRM based on all samples and a second time with the same GRM

produced above where one from each pair of sisters was removed. We calculated 95%

confidence intervals around each estimate by multiplying the standard error by 1.96. We

used R functions provided in (Wang & Xu, 2019) as well as the GCTA-GREML power

calculator (Visscher et al., 2014) to perform power analysis and estimate the effect sizes we

can expect to detect with our population datasets.

RESULTS

Generation of a highly-contiguous genome assembly of *B. balteatus*

We generated a *de novo* genome assembly for the species *B. balteatus* using Oxford

Nanopore (ONT) long-read and 10x Chromium linked-read sequencing and compared this to

the assembly for *B. sylvicola* that we recently published (Christmas et al., 2021). Assembly

statistics for BBAL_1.0 (*B. balteatus*) and BSYL_1.0 (*B. sylvicola*) are presented in table 1,

which includes statistics for the assemblies of *B. terrestris* (BTER_1.0) and *B. impatiens*

(BIMP_2.2) for comparison. Total assembly sizes of 250.1 and 252.1 Mbp for *B. balteatus*

and *B. sylvicola* respectively are both above the average assembly size of *Bombus* genomes

assembled to date of 247.9 Mbp (22 genomes, range 229.8 – 282.1 Mbp) (Heraghty et al.,

2020; Sadd et al., 2015; Sun et al., 2020). Contig N50s of 8.60 and 3.02 for BBAL_1.0 and

BSYL_1.0 respectively are the longest and third longest contig N50s of all published *Bombus*

assemblies, demonstrating these to be highly contiguous assemblies. Megabase-scale contig

N50 are typical of long-read assemblies, such as the ones presented here and in Heraghty et

al. (2020), and are significantly more contiguous than those based on short-read sequencing

technologies such as BTER_1.0 and BIMP_2.2.

High genome completeness and accuracy is also reflected in the BUSCO analysis, with scores

of 99.0% (98.7% single copy, 0.3% duplicated, 0.5% fragmented, 0.5% missing) and 98.2%

(97.9% single copy, 0.3% duplicated, 0.4% fragmented, 1.4% missing) reported for BBAL_1.0

and BSYL_1.0 respectively. Whole genome alignments to the *B. terrestris* assembly (Sadd et

al., 2015) resulted in 93.9% and 91.1% of the *B. balteatus* and *B. sylvicola* genomes being

placed on 18 pseudochromosomes. Our annotation pipeline annotated 11,711 and 11,585

genes in BBAL_1.0 and BSYL_1.0 respectively. These are highly comparable to the 11,874

genes in the *B. terrestris* gene set (v. 1.0), but lower than the 12,728 genes in the *B. impatiens* gene set (v. 2.1). Gene annotation of the *Bombus* assemblies presented in (Heraghty et al., 2020; Sun et al., 2020) also resulted in a greater number of genes annotated, ranging from 13,325 to 16,970 genes. BUSCO analyses of the annotated genes in our assemblies do suggest that we are missing some genes from our annotations (complete BUSCO genes in BBAL_01 and BSYL_01 respectively: 86.6% and 86.9%). Repeat content of our assemblies is comparable to that of other *Bombus* assemblies, with 17.11% of BBAL_1.0 and 17.85% of BSYL_1.0 identified as repetitive by RepeatMasker (full details of repeat classes in supplementary table S3). These proportions are comparable to the repeat content inferred in the *B. terrestris* and *B. impatiens* genomes.

Population resequencing indicates three species clusters, with little population structure within species

The locations of sample collections, which comprise 580 samples from across seven localities in the Rocky Mountains, CO, USA, are presented in Fig. 1 (full details in table S1, S2). Illumina sequencing resulted in an average genome coverage of 13.1x per sample, with 4.5 and 3.5 million SNPs identified in *B. balteatus* and *B. sylvicola* samples respectively (supplementary table S4). Neighbor-joining trees produced from samples of both groups of samples indicate that the *B. balteatus* samples form a single cluster whereas the *B. sylvicola* samples form two clusters (Fig 1). We previously inferred the largest cluster to be comprised of *B. sylvicola* samples whereas the second cluster is comprised of the cryptic species "incognitus" (Christmas et al., 2021). Among samples originally designated as *B. sylvicola,* 24% (n=67) are identified as "incognitus" whereas the remaining 76% (n=214) are *B. sylvicola. Bombus balteatus* (n=299) was the most commonly observed species across all localities. The proportions vary between localities (table S1). "Incognitus" samples were collected on six out of the seven mountains, whereas *B. balteatus* and *B. sylvicola* were found on all seven.

We calculated a pairwise kinship coefficient between samples to infer degrees of relatedness (Danecek et al., 2011; Manichaikul et al., 2010). We inferred a set of samples that were likely to be nestmates (sisters) based on them having the same maternal and paternal parents, under the realistic assumption that each colony is headed by a single

monandrous female queen (Estoup et al., 1995). Those identified as sisters were always

collected foraging on the same mountain and at the same or immediately neighboring site

(table S1). This suggests that foraging bees were generally found in the immediate vicinity of

their nests. Among the 299 *B. balteatus* samples, we identified 43 samples from 17 colonies

that had another nestmate in the dataset, indicating that we sampled a total of 273

colonies. For the 214 *B. sylvicola*, we identified 14 samples from 7 colonies that met this

criterion, indicating a total of 207 colonies. For 67 "incognitus", we identified two samples

from the same colony, indicating that we sampled 66 colonies.

We investigated the presence of population genetic structure within each species using

several methods. Neighbor-joining trees (Fig. 1) do not reveal the presence of any

substructure related to geography for *B. balteatus* or "incognitus". For *B. sylvicola*, all of the

samples from Quail Mountain (the furthermost southwestern site) cluster together, but do

not separate from the rest on the tree. Principal component analysis further reflected the

extremely low level of geographical structure across all species (Fig. 2). One slight exception

to this trend is the Quail Mountain population of *B. sylvicola* which, shows a slight degree of

separation from the other samples at the positive extreme of PC1 (Fig. 2B), although PC1

explains only 1.17% of the variance in the data. Outlier clusters seen in the PCA plots of *B.*

balteatus and "incognitus" (Fig. 2A,C, circled) are composed of highly related individuals as

shown in the kinship analysis, suggesting that the effect of kinship overrides any signal of

genetic structure. After removing one of each pair of sisters, samples of the same species

cluster tightly together and very little variance is explained by PC1 (Fig S1).

We assessed connectivity between colonies on different mountains by analyzing the

association between kinship and geography using Mantel tests after removing sisters from

the datasets (Fig. 3). We found no significant relationship between geographic distance and

relatedness for *B. balteatus* (Mantel R = -0.03 (95% ci: -0.04 - -0.02), p value = 0.11; Fig.

3A,D). There were small but significant negative relationships for *B. sylvicola* (Mantel R = -

0.06 (95% c.i. -0.07 - -0.04), p value = 0.0001; Fig. 3B,E) and "incognitus" (Mantel R = -0.04

(95% ci: -0.06 - -0.03), p value = 0.029; Fig. 3C,F), demonstrating greater relatedness at

closer locations. In these species, Mantel correlograms indicate significant associations

between kinship and geographical distance only at shorter distances (from the same or

neighboring mountains), which are no longer significant at greater distances (Fig. 3E-F).

These results indicate that the limited population structure is mainly driven by the

occurrence of a few highly-related samples from the same localities in the dataset.

Decrease in tongue length is ongoing in all species

We made morphological measurements for each sample (table S2). Intertegular distance is

considered a proxy for body size (Cane, 1987). We also measured the length of two parts of

the tongue: glossa and prementum. These two tongue measures are correlated in all three

species (*B. balteatus,* Pearson's r = 0.45, p < 0.001; *B. sylvicola,* Pearson's r = 0.38, p < 0.001;

incognitus, Pearson's r = 0.53, p < 0.001, Figs. S2-4). Only glossa is considered further, as it

was measured by Miller-Struttmann et al. (2015) and is the tongue part with most relevance

to feeding (Cariveau et al., 2016). Glossa is also significantly correlated with intertegular

distance (*B. balteatus,* Pearson's r = 0.41, p < 0.001; *B. sylvicola,* Pearson's r = 0.26, p <

0.001; incognitus, Pearson's r = 0.56, p < 0.001, Figs. S2-4).

A comparison of phenotypes between species shows that glossa length is substantially

longer in *B. balteatus*, consistent with previous knowledge (ls-means = 4.72 mm, 3.34 mm,

and 3.40 mm for *B. balteatus*, *B. sylvicola* and "incognitus" respectively; Tukey test, t ratio =

12.25 and 21.17, p < 0.001 for *B. balteatus* compared to *B. sylvicola* and "incognitus"*;* Fig.

4A). Average glossa length is extremely similar in *B. sylvicola* and "incognitus" (Tukey test, t

ratio =0.50, p = 0.87), suggesting that this character does not distinguish this newly-

discovered species from *B. sylvicola*. *Bombus balteatus* bees are significantly larger bees as

indicated by a greater intertegular distance, and *B. sylvicola* is significantly larger than

"incognitus" (mean intertegular distance = 4.00 mm, 3.83 mm, 3.58 mm for *B. balteatus*, *B.*

sylvicola and "incognitus" respectively; ANOVA, F=28.75, p < 0.001, Tukey post-hoc test

revealed differences are significant among all species at p < 0.001; Fig. S5A).

We next compared the distribution of glossa lengths in our samples with historical data. We

combined *B. sylvicola* and "incognitus" for this analysis because they do not differ in glossa

length on average and they were not distinguished in previous studies. We find an overall

trend of decreasing glossa length in both *B. sylvicola*-incognitus and *B. balteatus* over time

(Fig. 4B,C). This was reported previously between the collections from 1966-1980 and from

2012-2014 for both species (Miller-Struttmann et al., 2015). Here we show that this trend is

continuing, with further reductions in glossa length between the 2012-2014 collections and

our collections from 2017. These reductions are significant for both *B. sylvicola*-incognitus

(ls-means: 2012-2014 = 3.51 mm, 2017 = 3.21 mm; Tukey test, t-ratio = 3.198, p = 0.004)

and *B. balteatus* (ls-means: 2012-2014 = 5.24 mm, 2017 = 4.84 mm; Tukey test, t-ratio =

2.79, p = 0.015). In contrast, there is no significant trend shown by intertegular distance (Fig.

S5B,C), although for *B. sylvicola*-incognitus, intertegular distance is significantly larger in

both the 1966-80 and 2017 samples compared to the 2012-2014 samples (ANOVA, F =

15.28, p < 0.001. Tukey post-hoc test, p < 0.001).

For samples collected in 2017, we compared the distribution of glossa lengths for each

species among the seven mountains where they were collected (Fig. S6). There is an overall

trend for glossa lengths to be longer in bees from Mount Evans than on other mountains in

all species. The distributions for *B. balteatus* and *B. sylvicola* show significant differences,

with samples from Mount Evans having significantly longer glossae than on Mount

Democrat, Horseshoe Mountain, and Pennsylvania Mountain in *B. balteatus* (ls-mean glossa

length on Mt. Evans = 5.34 mm, other mountains 4.51 – 4.73 mm; Tukey p < 0.01 in each

case; Fig. S6A) and significantly longer glossae than on all other mountains except Niwot

Ridge in *B. sylvicola* (ls-mean glossa length on Mt. Evans = 3.77 mm, other mountains 3.02 –

3.24 mm; Tukey p < 0.01 in each case; Fig. S6B). A similar but non-significant trend could

also be seen for "incognitus" (Fig. S6C). Comparisons of intertegular distance between

mountains revealed that *B. balteatus* bees sampled on Niwot Ridge are significantly larger

than those on Boreas Mountain, Mount Democrat, Horseshoe Mountain, and Quail

Mountain (Niwot Ridge mean intertegular distance = 4.25 mm, other mountains 3.84 - 4.00

mm, ANOVA, F = 4.09, p < 0.001, Tukey post-hoc test p < 0.05 in four comparisons; Fig. S7A).

We did not identify any significant differences in intertegular distance of bees across

mountains for *B. sylvicola* (ANOVA, F = 1.83, p = 0.10) or "incognitus" (ANOVA, F = 1.62, p =

0.17; Fig. S7B,C).

Genome-wide association study identifies a SNP associated with a proxy for body size

We used a linear mixed-model analysis to search for correlations between allele frequencies

and quantitative trait variation in *B. sylvicola* and *B. balteatus.* We used the traits glossa

length, intertegular distance and glossa length corrected for intertegular distance (see

Methods). One from each pair of sisters identified in the kinship analysis was removed from

all analyses to reduce the level of relatedness in the sample set. We detected a genome-

wide significant association between intertegular distance and a SNP in intron 9 of the

CTRB1 (Chymotrypsinogen B1) gene on chromosome 10 in *B. sylvicola* (p < 0.001 after

Bonferroni correction; Fig. 5). The SNP is found at an allele frequency of 0.14 across all

populations, with 41 individuals heterozygous and nine homozygous for the allele. The allele

is present on all mountains, ranging in allele frequency from 0.06 (Quail Mountain) to 0.24

(Niwot Ridge). The *CTRB1* gene is involved in production of chymotrypsin, a proteolytic

digestive enzyme (Burgess, Malone, & Christeller, 1996; Giebel, Zwilling, & Pfleiderer, 1971;

Grogan & Hunt, 1979; Lazarević & Janković-Tomanić, 2015; Matsuoka et al., 2015; Rawlings

& Barrett, 1994). Intron 9 is a large 18 Kbp intron and the significant SNP is located 2 Kbp

upstream of exon 10. The GWAS significance suggests it is of likely functional importance

and its location is suggestive of a regulatory element, such as an intron splice enhancer or

silencer.

Two further SNPs, located 4 bp apart on chromosome 11, also significantly associated with

intertegular distance (p = 0.039 after Bonferroni correction). However, they are not located

in or close to any genes according to our annotation, so any potential functional significance

of these SNPs is unknown. We did not find any significant associations between allele

frequencies and intertegular distance in *B. balteatus* (Fig. S8). Furthermore, no significant

associations were found between allele frequencies and glossa length, or glossa length

corrected for intertegular distance in either species (Fig. S8). This indicates that the trait has

a polygenic or mainly environmental component. A power analysis (Visscher et al., 2014)

revealed that our datasets have sufficient power (>0.8) to detect a QTL with a h^2 of ~0.1 but

that loci with lower effect sizes would likely be undetectable.

Limited evidence that morphological variation has a genetic component

We performed GCTA-GREML analyses (Yang et al., 2011) to determine the genetic

component of trait variation for *B. sylvicola* and *B. balteatus* (Fig. 6, supplementary table

S5). The "incognitus" samples were not included here due to smaller sample size. When

each species' entire dataset is considered, we detect a heritable component for both glossa

length and intertegular distance in both species, where the 95% confidence interval does

not overlap zero. For *B. sylvicola*, but not *B. balteatus*, this is also the case when intertegular

distance is introduced as a covariate to account for body size when considering glossa

length. However, when we remove samples that we infer to be workers from the same

colony from the kinship analysis, we only detect a heritable component with 95%

confidence for intertegular distance in *B. balteatus* (all others then overlap with zero). This

indicates that a large proportion of the signal of covariance between genetic and trait

variance is due to the presence of sisters from the same colony. This could be a confounding

factor because environmental effects such as level of nutrition and colony health are likely

to covary with colony. However, such covariation could also reflect genetics. The level of

dependency of phenotype on genotype is therefore unclear from this analysis.

The 95% confidence intervals for the estimate of heritability have a wide span and overlap

zero in many cases. The power to determine heritability accurately depends on several

factors, including sample size, the structure of relatedness in the samples, and the true

value of heritability. When samples are all completely unrelated, it has been estimated

that > 3,000 samples are required to achieve a SE below 0.1 (Visscher et al., 2014). For

datasets of unrelated samples of the size we have analyzed here, we estimate a power of

~0.06, or ~6% probability, to detect $h^2 > 0$ for a trait with a SNP-heritability of 0.5 and a type

1 error rate (α) of 0.05. Greater sample size or higher levels of relatedness within the

dataset would therefore increase accuracy of our heritability estimates.

DISCUSSION

We generated a high-quality reference genome and performed genome sequencing of a

population sample of the bumblebee species *B. balteatus*. Combining these data with a

previous dataset from *B. sylvicola* produced the most extensive population genomics

dataset for bumblebees to date, with 580 samples from these two species and the newly-

discovered "incognitus" collected from high-altitude locations in Colorado, USA. These data

were combined with morphometric measurements from each sample and used to

investigate genetic structure, morphological evolution and the genetic basis of phenotypic

traits in these populations. Our main findings are 1) no evidence for gene flow between

species, including the recently-inferred "incognitus", but lack of population structure within

species, indicating high levels of connectivity among subpopulations. 2) An ongoing

decrease in tongue (glossa) length compared to historical measurements, previously

inferred to reflect the effects of decreasing floral abundance on bumble bee foraging

(Miller-Struttmann et al., 2015). 3) An association between body size and variation at a

single SNP located in a gene related to production of a digestive enzyme in *B. sylvicola*. 4) A

lack of conclusive evidence for a genetic component to tongue length in both species. Our

analysis is unable to determine the degree of heritability underlying tongue length or body

size but is consistent with these traits having a polygenic genetic component.

This investigation adds to the number of bumblebee species with reference genomes

available. The genomes of the common species *B. impatiens* and *B. terrestris* were

sequenced using short-read technologies (Sadd et al., 2015). Genome assemblies are also

available for representatives of all 15 *Bombus* subgenera (Sun et al., 2020) based on short-

read sequencing of jumping libraries and scaffolding using Hi-C, resulting in contig N50 of

325 kb. In addition to the two genomes produced here, the genomes of the North American

species *B. terricola*, *B. bifarius*, *B. vancouverensis*, and *B. vosnesenskii* have also been

produced using long-read sequencing (PacBio or ONT) and have megabase-scale contig N50s

(Heraghty et al., 2020; Kent et al., 2018).

Identification of the previously-unknown species "incognitus" demonstrates the power of

population-scale genome sequencing to discover new cryptic variation (Christmas et al.,

2021). None of the morphological features that we studied could distinguish between *B.

sylvicola* and "incognitus". *Bombus sylvicola* did however have a significant tendency to be

collected at higher altitudes and have larger body size. The wider distribution of

"incognitus" is unknown, as is the true composition of populations described as *B. sylvicola*

in other geographical locations. Existing collections of *B. sylvicola* from Colorado are likely to

be a mixture of both species. However, importantly for this study, there were no significant

differences in glossa or prementum length between the two species.

We find no evidence for population structure among samples of any of the three species

studied here. This indicates an absence of long-term geographical barriers to gene flow

among the high-elevation localities included here. Furthermore, although we observe

elevated degrees of relatedness among foragers collected on the same mountain, there is

no significant tendency for elevated relatedness of foragers caught on nearby mountains,

indicating a lack of population structure at this scale. Foragers inferred to belong to the

same nest (i.e. with identical parents) were always found in the same or adjacent sites,

consistent with the limited foraging distance (on average < 110m) inferred previously by

Geib et al. (2015) for these species.

Studies of several bumblebee species across continental USA, including *B. bifarius* and *B.*

vosnesenkii, also found in Colorado and across western North America, indicate weak

geographical differentiation even at distances over 1000 km (Ghisbain et al., 2020; Lozier et

al., 2011). However, the complex topography of mountain ranges likely modulates gene

flow among populations (Lozier et al., 2013, 2011). Populations of *B. bifarius* are inferred to

be more fragmented in the southern portions of their ranges, where they occur at higher

elevations (Jackson et al., 2018). The species studied here, *B. sylvicola* and *B. balteatus,* are

generally restricted to high-elevation localities above the tree line separated by forest

(Williams, Thorp, Richardson, & Colla, 2014). However, our analysis indicates that this

complex landscape has not historically limited mating and dispersal between localities in the

same mountain range. This is consistent with previous studies of these species that found

substantial genetic differentiation between mountain ranges in the Pacific Northwest, but

not within the same mountain range (Koch et al., 2017; Whitley, 2018).

Morphological change over time has already been documented in the species studied here,

with a significant reduction in tongue length observed since 1966 (Miller-Struttmann et al.,

2015). During this time period, there has also been a shift in species composition, with

shorter-tongued species becoming more predominant. Previous collections of *B. sylvicola* in

this region are likely a mixture of *B. sylvicola* and "incognitus", as the existence of

"incognitus" was unknown. However, as *B. sylvicola* and "incognitus" do not differ in glossa

length, undetected changes in the relative abundance of these species over time are

unlikely to be responsible for the observed shifts in glossa length observed in samples that

are defined as *B. sylvicola.* This effect could, however, potentially contribute to the

observed variation in body size over time as *B. sylvicola* have a significantly greater

intertegular distance on average than "incognitus" (Fig. S5).

A major finding that we present here is that the temporal shift toward shorter tongue

length is ongoing. In addition to the significant reduction previously observed over a period

of six decades up until 2014, we also observe a significant reduction over a much shorter

time period between 2012-14 and 2017 in both *B. balteatus* and *B. sylvicola*-incognitus.

Mean tongue length reportedly decreased on average by 0.61% annually in a period

between 1966 and 2014 for *B. sylvicola* (Miller-Struttmann et al., 2015). According to our

recent measurements, the corresponding decrease in 3-5 subsequent years has been

12.9 %, or 3.2% per year (compared to the 2012-14 average).

Decreases in tongue-length have been previously attributed to an advantage of shorter

tongues for foraging on a wide range of flowers. Such generalist foraging could be favored

due to a general decline of floral resources observed in connection with warmer summers

(Miller-Struttmann et al., 2015). While both long- and short-tongued bees are capable of

accessing nectar in plants with short flower tubes, short-tongued bees are more efficient at

foraging from them (Inouye, 1980) and are also more efficient at foraging across a diversity

of flower tube depths (Arbulo, Santos, Salvarrey, & Invernizzi, 2011; Geib & Galen, 2012;

Plowright & Plowright, 1997). Short-tongued bumble bees often forage from a wider suite of

plants both in these alpine bumble bee species (Miller-Struttmann et al., 2015) and others

(Goulson & Darvill, 2004; Goulson et al., 2008b; Heinrich, 2004; Huang et al., 2015). These

considerations indicate that shorter tongues are likely advantageous for generalist foraging.

Although short-tongued bees are unable to pollinate plants with long flower tubes, they are

likely important for supporting a diverse array of flowering plants.

It is unclear whether the tongue-length changes we observe are the result of natural

selection or phenotypic plasticity. There are now many examples of rapid evolution of traits

driven by natural selection due to climatic events in the wild, such as a shift in beak shape

over a period of a few years in Darwin's finches related to competition for food during

drought (Lamichhaney et al., 2016) and genomic shifts related to cold-tolerance in the green

anole lizard related to severe winter storms (Campbell-Staton et al., 2017). Human activity

may also promote rapid adaptation by natural selection, as demonstrated by the evolution

of longer beaks in great tits promoted by the use of bird feeders in the UK (Bosse et al.,

2017). Global climate change is also expected to be a driver of rapid evolution, but it is often

difficult to establish whether trends of phenotypic change result from evolution or

phenotypic plasticity (Merilä & Hendry, 2014). Long-term changes in breeding timing in

great tits (Charmantier et al., 2008) and body mass in red-billed gulls (Teplitsky, Mills, Alho,

Yarrall, & Merilä, 2008) that track temperature increases have been shown to result solely

from phenotypic plasticity.

To address the causes of the changes in bumblebee tongue length observed here, we

attempted to uncover the genetic basis of morphological variation using genome-wide

association studies (GWAS) and genome-wide complex trait analysis (GCTA). No significant

associations were observed for tongue length in the genome of any of the species using

GWAS. However, a significant association with body size was identified in *B. sylvicola* at a

single SNP in the *CTRB1* (Chymotrypsinogen B1) gene. The lack of association at neighboring

SNPs could be attributable to high rates of meiotic recombination and low levels of linkage

disequilibrium in bumblebees (Kawakami et al., 2019). The *CTRB1* gene produces a

precursor of chymotrypsin, a key digestive enzyme in animals, which is likely important for

pollen digestion in bees (Burgess et al., 1996; Giebel et al., 1971; Grogan & Hunt, 1979;

Lazarević & Janković-Tomanić, 2015; Matsuoka et al., 2015; Rawlings & Barrett, 1994). The

associated SNP is found in a large intron (18 Kbp) of this gene, 2 Kbp upstream of the

nearest exon. We lack any functional annotation for this SNP but its location and GWAS

significance suggests it may play a role in gene regulation. For example, if located within an

intronic splicing enhancer or silencer, it may result in variation in expression levels of

different isoforms of the CTRB1 enzyme being produced. Such changes could influence body

size through an effect on the efficiency of digestion.

We detected a significant signal of covariance of trait measurements with relatedness for

both tongue length and intertegular distance using GCTA when closely-related individuals

were included in the dataset. However, no significant signal of heritability in morphological

traits was detected when the dataset was purged of close relatives. The first result could

potentially reflect a genetic component, but it is not possible to disentangle this component

from the shared environmental effects due to close relatives sharing a nest. This analysis is

therefore underpowered to detect genetic effects due to low sample size and a general lack

of related individuals in the dataset. A possible explanation for our results is that the

morphological traits tongue length and intertegular distance have a highly polygenic genetic

component, with multiple QTL of small effect ($h^2 < 0.1$), such that GWAS and GCTA were

underpowered to detect QTL and estimate heritability. The size of the genetic component of

variation in both traits is therefore still unknown in each species. These results contrast with

findings regarding the genetic control of bumblebee coloration, in which variation at single

loci control intraspecific variation in color of abdominal segments (Rahman, Cnaani, Kinch,

Grishin, & Hines, 2021; Tian et al., 2019).

It therefore remains unclear whether the observed changes in tongue length are caused by

natural selection acting on functional genetic variation, or whether they are the result of

phenotypic plasticity. There is large variation in body size between individuals in bumblebee

nests, indicating substantial phenotypic plasticity in this trait (Peat, Tucker, & Goulson,

2005). This variability could be explained by a correlation between larval position in the nest

and size, which is related to feeding intensity (Couvillon & Dornhaus, 2009). There is

evidence from solitary bees of an effect of temperature on body size (Scaven & Rafferty,

2013), although such an effect is likely to be less pronounced in social insects such as

bumblebees that regulate the temperature of their nests. However, it is unclear whether

tongue length also exhibits plasticity, as this trait correlates with body size among

nestmates (Peat et al., 2005), indicating that plasticity in tongue length independent of body

size is less pronounced. However, our observation of significant differences in mean tongue

length between mountains could potentially reflect phenotypic plasticity.

In summary, we have shown that decreasing tongue length observed over the last six

decades is still ongoing in populations of the bumblebee species *B. sylvicola* and *B. balteatus*

(and the recently-reported "incognitus") at high altitudes in Colorado. We have generated

the genomic resources to study these species on the genetic level in the form of highly-

contiguous genome assemblies and comprehensively characterized genome variation. Our

results indicate that the complex topology of the mountain landscape in Colorado does not

contain significant barriers to mating and dispersal of bumblebees in the study area. We are

unable to conclude whether the observed morphological shifts over time are the result of

genetic adaptation or phenotypic plasticity. Alpine ecosystems are particularly sensitive to

climate change and may suffer its effects before other geographical regions (Elsen &

Tingley, 2015). Continued monitoring of changes in morphology in these and other

populations in these regions could be highly valuable for understanding how pollinators

respond to the effects of a warming climate, the degree to which rapid evolution by natural

selection or phenotypic plasticity are possible, and the effects of the changes on these

ecosystems.

DATA AND CODE AVAILABILITY

All sequence data presented here are available at NCBI under BioProject IDs PRJNA646847

and PRJNA704506. Scripts and annotation parameters used in the study are available at

GitHub (https://github.com/matt-webster-lab/alpine_bumblebees). VCF files containing

genotypic variation are available at the Dryad Digital Repository

(https://doi.org/10.5061/dryad.mgqnk990p).

ACKNOWLEDGEMENTS

This research was funded by Swedish Research Council Formas grant 2016-00535 and a

SciLifeLab National Biodiversity Project NP00046 to MTW. MK is financially supported by the

Knut and Alice Wallenberg Foundation as part of the National Bioinformatics Infrastructure

Sweden at SciLifeLab. The authors acknowledge support from the National Genomics

Infrastructure in Stockholm and Uppsala funded by Science for Life Laboratory, the Knut and

Alice Wallenberg Foundation and the Swedish Research Council. The computations and data

handling were enabled by resources provided by the Swedish National Infrastructure for

Computing (SNIC) at UPPMAX partially funded by the Swedish Research Council through

grant agreement no. 2018-05973. We thank the Mountain Research Station (U.C. Boulder),

Mt. Evans Research Station (University of Denver), the U.S. Forest Service, Dr. Rosemond

Greinetz and the Mountain Area Land Trust for providing assistance and access to field sites.

We thank Jacqueline Staab and Michael Osbourn who aided in specimen collection,

identification and preservation and Lisa Danback, Kate Schleicher, Ying Thao, Anna Grobelny

and Breana Cook who carried out phenotypic measurements.

REFERENCES

10XGenomics/longranger [Python]. (2021). 10x Genomics. Retrieved from
https://github.com/10XGenomics/longranger (Original work published 2018)
Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry
in unrelated individuals. *Genome Research*, *19*(9), 1655–1664. doi:
10.1101/gr.094052.109
Andrews, S. (2021). *FastQC* [Java]. Retrieved from https://github.com/s-andrews/FastQC
(Original work published 2017)
Arbulo, N., Santos, E., Salvarrey, S., & Invernizzi, C. (2011). Proboscis Length and Resource
Utilization in Two Uruguayan Bumblebees: *Bombus atratus* Franklin and *Bombus*
bellicosus Smith (Hymenoptera: Apidae). *Neotropical Entomology*, *40*, 72–77. doi:
10.1590/S1519-566X2011000100010
Bosse, M., Spurgin, L. G., Laine, V. N., Cole, E. F., Firth, J. A., Gienapp, P., … Slate, J. (2017).
Recent natural selection causes adaptive evolution of an avian polygenic trait.
Science, *358*(6361), 365–368. doi: 10.1126/science.aal3298
Burgess, E. P. J., Malone, L. A., & Christeller, J. T. (1996). Effects of two proteinase inhibitors
on the digestive enzymes and survival of honey bees (*Apis mellifera*). *Journal of*
Insect Physiology, *42*(9), 823–828. doi: 10.1016/0022-1910(96)00045-5
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T.
L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*, 421. doi:
10.1186/1471-2105-10-421
Cameron, S. A., Lozier, J. D., Strange, J. P., Koch, J. B., Cordes, N., Solter, L. F., & Griswold, T.
L. (2011). Patterns of widespread decline in North American bumble bees.
Proceedings of the National Academy of Sciences, *108*(2), 662–667. doi:
10.1073/pnas.1014743108
Cameron, S. A., & Sadd, B. M. (2020). Global Trends in Bumble Bee Health. *Annual Review of*
Entomology, *65*, 209–232. doi: 10.1146/annurev-ento-011118-111847
Campbell-Staton, S. C., Cheviron, Z. A., Rochette, N., Catchen, J., Losos, J. B., & Edwards, S.
V. (2017). Winter storms drive rapid phenotypic, regulatory, and genomic shifts in
the green anole lizard. *Science*, *357*(6350), 495–498. doi: 10.1126/science.aam5512
Cane, J. H. (1987). Estimation of Bee Size Using Intertegular Span (Apoidea). *Journal of the*
Kansas Entomological Society, *60*(1), 145–147.
Cantarel, B. L., Korf, I., Robb, S. M. C., Parra, G., Ross, E., Moore, B., … Yandell, M. (2008).
MAKER: An easy-to-use annotation pipeline designed for emerging model organism
genomes. *Genome Research*, *18*(1), 188–196. doi: 10.1101/gr.6743907
Cariveau, D. P., Nayak, G. K., Bartomeus, I., Zientek, J., Ascher, J. S., Gibbs, J., & Winfree, R.
(2016). The Allometry of Bee Proboscis Length and Its Uses in Ecology. *PLOS ONE*,
11(3), e0151482. doi: 10.1371/journal.pone.0151482
Charmantier, A., McCleery, R. H., Cole, L. R., Perrins, C., Kruuk, L. E. B., & Sheldon, B. C.
(2008). Adaptive Phenotypic Plasticity in Response to Climate Change in a Wild Bird
Population. *Science*, *320*(5877), 800–803. doi: 10.1126/science.1157174
Christmas, M. J., Jones, J. C., Olsson, A., Wallerman, O., Bunikis, I., Kierczak, M., … Webster,
M. T. (2021). Genetic Barriers to Historical Gene Flow between Cryptic Species of
Alpine Bumblebees Revealed by Comparative Population Genomics. *Molecular*
Biology and Evolution, (msab086). doi: 10.1093/molbev/msab086
Couvillon, M. J., & Dornhaus, A. (2009). Location, location, location: Larvae position inside
the nest is correlated with adult body size in worker bumble-bees (*Bombus*

impatiens). *Proceedings of the Royal Society B: Biological Sciences*, *276*(1666), 2411–2418. doi: 10.1098/rspb.2009.0172

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. doi: 10.1093/bioinformatics/btr330

Darvill, B., Ellis, J. S., Lye, G. C., & Goulson, D. (2006). Population structure and inbreeding in a rare and declining bumblebee, *Bombus muscorum* (Hymenoptera: Apidae). *Molecular Ecology*, *15*(3), 601–611. doi: 10.1111/j.1365-294X.2006.02797.x

Darvill, B., Knight, M. E., & Goulson, D. (2004). Use of Genetic Markers to Quantify Bumblebee Foraging Range and Nest Density. *Oikos*, *107*(3), 471–478. JSTOR. Retrieved from JSTOR.

Ellis, J. S., Knight, M. E., Darvill, B., & Goulson, D. (2006). Extremely low effective population sizes, genetic structuring and reduced genetic diversity in a threatened bumblebee species, *Bombus sylvarum* (Hymenoptera: Apidae). *Molecular Ecology*, *15*(14), 4375–4386. doi: 10.1111/j.1365-294X.2006.03121.x

Elsen, P. R., & Tingley, M. W. (2015). Global mountain topography and the fate of montane species under climate change. *Nature Climate Change*, *5*(8), 772–776. doi: 10.1038/nclimate2656

Estoup, A., Scholl, A., Pouvreau, A., & Solignac, M. (1995). Monoandry and polyandry in bumble bees (Hymenoptera; Bombinae) as evidenced by highly variable microsatellites. *Molecular Ecology*, *4*(1), 89–93. doi: 10.1111/j.1365-294x.1995.tb00195.x

Geib, J. C., & Galen, C. (2012). Tracing impacts of partner abundance in facultative pollination mutualisms: From individuals to populations. *Ecology*, *93*(7), 1581–1592. doi: 10.1890/11-1271.1

Geib, J. C., Strange, J. P., & Galen, C. (2015). Bumble bee nest abundance, foraging distance, and host-plant reproduction: Implications for management and conservation. *Ecological Applications*, *25*(3), 768–778. doi: https://doi.org/10.1890/14-0151.1

Ghisbain, G., Lozier, J. D., Rahman, S. R., Ezray, B. D., Tian, L., Ulmer, J. M., … Hines, H. M. (2020). Substantial genetic divergence and lack of recent gene flow support cryptic speciation in a colour polymorphic bumble bee (*Bombus bifarius*) species complex. *Systematic Entomology*, *45*(3), 635–652. doi: 10.1111/syen.12419

Giebel, W., Zwilling, R., & Pfleiderer, G. (1971). The evolution of endopeptidases—XII. The proteolytic enzymes of the honeybee (*Apis Mellifica* L.): Purification and characterization of endopeptidases in the midgut of adult workers and comparative studies on the endopeptidase-pattern of different castes and on different ontogenetic stages. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, *38*(1), 197–210. doi: 10.1016/0305-0491(71)90297-5

Gienapp, P., Fior, S., Guillaume, F., Lasky, J. R., Sork, V. L., & Csilléry, K. (2017). Genomic Quantitative Genetics to Study Evolution in the Wild. *Trends in Ecology & Evolution*, *32*(12), 897–908. doi: 10.1016/j.tree.2017.09.004

Goulson, D. (2003). *Bumblebees: Their Behaviour and Ecology*. Oxford University Press.

Goulson, D., & Darvill, B. (2004). Niche overlap and diet breadth in bumblebees; are rare species more specialized in their choice of flowers? *Apidologie*, *35*(1), 55–63. doi: 10.1051/apido:2003062

Goulson, D., Hanley, M. E., Darvill, B., Ellis, J. S., & Knight, M. E. (2005). Causes of rarity in bumblebees. *Biological Conservation*, *122*(1), 1–8. doi: 10.1016/j.biocon.2004.06.017

Goulson, D., Lye, G. C., & Darvill, B. (2008a). Decline and conservation of bumble bees.
Annual Review of Entomology, *53*, 191–208. doi:
10.1146/annurev.ento.53.103106.093454
Goulson, D., Lye, G. C., & Darvill, B. (2008b). Diet breadth, coexistence and rarity in
bumblebees. *Biodiversity and Conservation*, *17*(13), 3269–3288. doi:
10.1007/s10531-008-9428-y
Goulson, D., Nicholls, E., Botías, C., & Rotheray, E. L. (2015). Bee declines driven by
combined stress from parasites, pesticides, and lack of flowers. *Science (New York,*
N.Y.), *347*(6229), 1255957. doi: 10.1126/science.1255957
Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., … Regev, A.
(2011). Full-length transcriptome assembly from RNA-Seq data without a reference
genome. *Nature Biotechnology*, *29*(7), 644–652. doi: 10.1038/nbt.1883
Grabherr, M. G., Russell, P., Meyer, M., Mauceli, E., Alföldi, J., Di Palma, F., & Lindblad-Toh,
K. (2010). Genome-wide synteny through highly sensitive sequence alignment:
Satsuma. *Bioinformatics*, *26*(9), 1145–1151. doi: 10.1093/bioinformatics/btq102
Grogan, D. E., & Hunt, J. H. (1979). Pollen proteases: Their potential role in insect digestion.
Insect Biochemistry, *9*(3), 309–313. doi: 10.1016/0020-1790(79)90011-8
Heinrich, B. (2004). *Bumblebee Economics*. Cambridge: Harvard University Press. Retrieved
from https://www.hup.harvard.edu/catalog.php?isbn=9780674016392
Heraghty, S. D., Sutton, J. M., Pimsler, M. L., Fierst, J. L., Strange, J. P., & Lozier, J. D. (2020).
De Novo Genome Assemblies for Three North American Bumble Bee Species:
Bombus bifarius, *Bombus vancouverensis*, and *Bombus vosnesenskii*. *G3: Genes,*
Genomes, Genetics, *10*(8), 2585–2592. doi: 10.1534/g3.120.401437
Huang, J., An, J., Wu, J., & Williams, P. H. (2015). Extreme Food-Plant Specialisation in
Megabombus Bumblebees as a Product of Long Tongues Combined with Short
Nesting Seasons. *PLOS ONE*, *10*(8), e0132358. doi: 10.1371/journal.pone.0132358
Inouye, D. W. (1980). The effect of proboscis and corolla tube lengths on patterns and rates
of flower visitation by bumblebees. *Oecologia*, *45*(2), 197–201. doi:
10.1007/BF00346460
Jackman, S. D., Coombe, L., Chu, J., Warren, R. L., Vandervalk, B. P., Yeo, S., … Birol, I. (2018).
Tigmint: Correcting assembly errors using linked reads from large molecules. *BMC*
Bioinformatics, *19*(1), 393. doi: 10.1186/s12859-018-2425-6
Jackson, J. M., Pimsler, M. L., Oyen, K. J., Koch-Uhuad, J. B., Herndon, J. D., Strange, J. P., …
Lozier, J. D. (2018). Distance, elevation and environment as drivers of diversity and
divergence in bumble bees across latitude and altitude. *Molecular Ecology*, *27*(14),
2926–2942. doi: 10.1111/mec.14735
Jha, S. (2015). Contemporary human-altered landscapes and oceanic barriers reduce
bumble bee gene flow. *Molecular Ecology*, *24*(5), 993–1006. doi: 10.1111/mec.13090
Jha, S., & Kremen, C. (2013). Resource diversity and landscape-level homogeneity drive
native bee foraging. *Proceedings of the National Academy of Sciences*, *110*(2), 555–
558.
Kawakami, T., Wallberg, A., Olsson, A., Wintermantel, D., Miranda, J. R. de, Allsopp, M., …
Webster, M. T. (2019). Substantial Heritable Variation in Recombination Rate on
Multiple Scales in Honeybees and Bumblebees. *Genetics*, genetics.302008.2019. doi:
10.1534/genetics.119.302008
Kearns, C. A., & Inouye, D. W. (1993). *Techniques for Pollination Biologists*. Boulder:
University Press of Colorado.

Kent, C. F., Dey, A., Patel, H., Tsvetkov, N., Tiwari, T., MacPhail, V. J., … Zayed, A. (2018).
Conservation Genomics of the Declining North American Bumblebee *Bombus*
terricola Reveals Inbreeding and Selection on Immune Genes. *Frontiers in Genetics*,
9. doi: 10.3389/fgene.2018.00316

Kerr, J. T., Pindar, A., Galpern, P., Packer, L., Potts, S. G., Roberts, S. M., … Pantoja, A. (2015).
CLIMATE CHANGE. Climate change impacts on bumblebees converge across
continents. *Science (New York, N.Y.)*, *349*(6244), 177–180. doi:
10.1126/science.aaa7031

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*,
37(8), 907–915. doi: 10.1038/s41587-019-0201-4

Knight, M. E., Martin, A. P., Bishop, S., Osborne, J. L., Hale, R. J., Sanderson, R. A., & Goulson,
D. (2005). An interspecific comparison of foraging range and nest density of four
bumblebee (Bombus) species. *Molecular Ecology*, *14*(6), 1811–1820. doi:
10.1111/j.1365-294X.2005.02540.x

Koch, J. B., Looney, C., Sheppard, W. S., & Strange, J. P. (2017). Patterns of population
genetic structure and diversity across bumble bee communities in the Pacific
Northwest. *Conservation Genetics*, *18*(3), 507–520. doi: 10.1007/s10592-017-0944-8

Koch, J. B., Strange, J. P., & Williams, P. H. (2014). *Bumble Bees of the Western United*
States. U.S. Forest Service.

Lamichhaney, S., Han, F., Berglund, J., Wang, C., Almen, M. S., Webster, M. T., … Andersson,
L. (2016). A beak size locus in Darwins finches facilitated character displacement
during a drought. *Science*, *352*(6284), 470–474. doi: 10.1126/science.aad8786

Lazarević, J., & Janković-Tomanić, M. (2015). Dietary and phylogenetic correlates of
digestive trypsin activity in insect pests. *Entomologia Experimentalis et Applicata*,
157(2), 123–151. doi: 10.1111/eea.12349

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics*, *25*(14), 1754–1760. doi: 10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … Durbin, R. (2009). The
Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079.
doi: 10.1093/bioinformatics/btp352

Lozier, J. D., Strange, J. P., & Koch, J. B. (2013). Landscape heterogeneity predicts gene flow
in a widespread polymorphic bumble bee, *Bombus bifarius* (Hymenoptera: Apidae).
Conservation Genetics, *14*(5), 1099–1110. doi: 10.1007/s10592-013-0498-3

Lozier, J. D., Strange, J. P., Stewart, I. J., & Cameron, S. A. (2011). Patterns of range-wide
genetic variation in six North American bumble bee (Apidae: Bombus) species.
Molecular Ecology, *20*(23), 4870–4888. doi: 10.1111/j.1365-294X.2011.05314.x

Manichaikul, A., Mychaleckyj, J. C., Rich, S. S., Daly, K., Sale, M., & Chen, W.-M. (2010).
Robust relationship inference in genome-wide association studies. *Bioinformatics*,
26(22), 2867–2873. doi: 10.1093/bioinformatics/btq559

Marshall, L., Perdijk, F., Dendoncker, N., Kunin, W., Roberts, S., & Biesmeijer, J. C. (2020).
Bumblebees moving up: Shifts in elevation ranges in the Pyrenees over 115 years.
Proceedings. Biological Sciences, *287*(1938), 20202201. doi: 10.1098/rspb.2020.2201

Martinet, B., Lecocq, T., Brasero, N., Gerard, M., Urbanová, K., Valterová, I., … Rasmont, P.
(2019). Integrative taxonomy of an arctic bumblebee species complex highlights a
new cryptic species (Apidae: Bombus). *Zoological Journal of the Linnean Society*,
187(3), 599–621. doi: 10.1093/zoolinnean/zlz041

Matsuoka, T., Takasaki, A., Mishima, T., Kawashima, T., Kanamaru, Y., Nakamura, T., & Yabe, T. (2015). Expression and characterization of honeybee, *Apis mellifera*, larva chymotrypsin-like protease. *Apidologie*, *46*(2), 167–176. doi: 10.1007/s13592-014-0313-2

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., … DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303. doi: 10.1101/gr.107524.110

Medaka [Python]. (2021). Oxford Nanopore Technologies. Retrieved from https://github.com/nanoporetech/medaka (Original work published 2017)

Merilä, J., & Hendry, A. P. (2014). Climate change, adaptation, and phenotypic plasticity: The problem and the evidence. *Evolutionary Applications*, *7*(1), 1–14. doi: 10.1111/eva.12137

Miller-Struttmann, N. E., Geib, J. C., Franklin, J. D., Kevan, P. G., Holdo, R. M., Ebert-May, D., … Galen, C. (2015). Functional mismatch in a bumble bee pollination mutualism under climate change. *Science*, *349*(6255), 1541–1544. doi: 10.1126/science.aab0868

Pauls, S. U., Nowak, C., Bálint, M., & Pfenninger, M. (2013). The impact of global climate change on genetic diversity within populations and species. *Molecular Ecology*, *22*(4), 925–946. doi: https://doi.org/10.1111/mec.12152

Peat, J., Tucker, J., & Goulson, D. (2005). Does intraspecific size variation in bumblebees allow colonies to efficiently exploit different flowers? *Ecological Entomology*, *30*(2), 176–181. doi: 10.1111/j.0307-6946.2005.00676.x

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., & Salzberg, S. L. (2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology*, *33*(3), 290–295. doi: 10.1038/nbt.3122

Plowright, C. M. S., & Plowright, R. C. (1997). The advantage of short tongues in bumble bees (Bombus)—Analyses of species distributions according to flower corolla depth, and of working speeds on white clover. *The Canadian Entomologist*, *129*(1), 51–59. doi: 10.4039/Ent12951-1

Rahman, S. R., Cnaani, J., Kinch, L. N., Grishin, N. V., & Hines, H. M. (2021). A combined RAD-Seq and WGS approach reveals the genomic basis of yellow color variation in bumble bee Bombus terrestris. *Scientific Reports*, *11*(1), 7996. doi: 10.1038/s41598-021-87194-y

Rawlings, N. D., & Barrett, A. J. (1994). Families of serine peptidases. *Methods in Enzymology*, *244*, 19–61. doi: 10.1016/0076-6879(94)44004-2

Razgour, O., Forester, B., Taggart, J. B., Bekaert, M., Juste, J., Ibáñez, C., … Manel, S. (2019). Considering adaptive genetic variation in climate change vulnerability assessment reduces species range loss projections. *Proceedings of the National Academy of Sciences*, *116*(21), 10418–10423. doi: 10.1073/pnas.1820663116

Ruan, J., & Li, H. (2020). Fast and accurate long-read assembly with wtdbg2. *Nature Methods*, *17*(2), 155–158. doi: 10.1038/s41592-019-0669-3

Sadd, B. M., Barribeau, S. M., Bloch, G., Graaf, D. C. de, Dearden, P., Elsik, C. G., … Worley, K. C. (2015). The genomes of two key bumblebee species with primitive eusocial organization. *Genome Biology*, *16*(1), 76. doi: 10.1186/s13059-015-0623-3

Santure, A. W., & Garant, D. (2018). Wild GWAS—association mapping in natural
populations. *Molecular Ecology Resources*, *18*(4), 729–738. doi: 10.1111/1755-
0998.12901
Scaven, V. L., & Rafferty, N. E. (2013). Physiological effects of climate warming on flowering
plants and insect pollinators and potential consequences for their interactions.
Current Zoology, *59*(3), 418–426.
Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of
image analysis. *Nature Methods*, *9*(7), 671–675. doi: 10.1038/nmeth.2089
Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).
BUSCO: Assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics*, *31*(19), 3210–3212. doi: 10.1093/bioinformatics/btv351
Smit, A., Hubley, R., & Green, P. (2015). *RepeatMasker Open-4.0. 2013-2015*
<http://www.repeatmasker.org>.
Soroye, P., Newbold, T., & Kerr, J. (2020). Climate change contributes to widespread
declines among bumble bees across continents. *Science*, *367*(6478), 685–688. doi:
10.1126/science.aax8591
Sun, C., Huang, J., Wang, Y., Zhao, X., Su, L., Thomas, G. W. C., … Mueller, R. L. (2020).
Genus-wide characterization of bumblebee genomes provides insights into their
evolution and variation in ecological and behavioral traits. *Molecular Biology and
Evolution*, (msaa240). doi: 10.1093/molbev/msaa240
Teplitsky, C., Mills, J. A., Alho, J. S., Yarrall, J. W., & Merilä, J. (2008). Bergmann's rule and
climate change revisited: Disentangling environmental and genetic responses in a
wild bird population. *Proceedings of the National Academy of Sciences*, *105*(36),
13492–13496.
Teutenberg, J. (2020). *Downpore* [Go]. Retrieved from
https://github.com/jteutenberg/downpore (Original work published 2018)
Tian, L., Rahman, S. R., Ezray, B. D., Franzini, L., Strange, J. P., Lhomme, P., & Hines, H. M.
(2019). A homeotic shift late in development drives mimetic color variation in a
bumble bee. *Proceedings of the National Academy of Sciences*, *116*(24), 11857–
11865. doi: 10.1073/pnas.1900365116
Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. (2017). Fast and accurate de novo genome
assembly from long uncorrected reads. *Genome Research*, *27*(5), 737–746. doi:
10.1101/gr.214270.116
Visscher, P. M., Hemani, G., Vinkhuyzen, A. A. E., Chen, G.-B., Lee, S. H., Wray, N. R., … Yang,
J. (2014). Statistical Power to Detect Genetic (Co)Variance of Complex Traits Using
SNP Data in Unrelated Samples. *PLOS Genetics*, *10*(4), e1004269. doi:
10.1371/journal.pgen.1004269
Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., … Earl, A. M.
(2014). Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection and
Genome Assembly Improvement. *PLOS ONE*, *9*(11), e112963. doi:
10.1371/journal.pone.0112963
Wang, M., & Xu, S. (2019). Statistical power in genome-wide association studies and
quantitative trait locus mapping. *Heredity*, *123*(3), 287–306. doi: 10.1038/s41437-
019-0205-3
Whitley, K. M. (2018). *Genetic structure and gene flow barriers among populations of an
alpine bumble bee (Bombus balteatus) in the central Rocky Mountains* (M.Sc.).
Appalachian State University.

Williams, P. H., Berezin, M. V., Cannings, S. G., Cederberg, B., Ødegaard, F., Rasmussen, C., …
Byvaltsev, A. M. (2019). The arctic and alpine bumblebees of the subgenus
Alpinobombus revised from integrative assessment of species' gene coalescents and
morphology (Hymenoptera, Apidae, Bombus). *Zootaxa*, *4625*(1), zootaxa.4625.1.1.
doi: 10.11646/zootaxa.4625.1.1
Williams, P., Thorp, R., Richardson, L., & Colla. (2014). *Bumble Bees of North America*.
Princeton University Press.
Woodard, S. H., Lozier, J. D., Goulson, D., Williams, P. H., Strange, J. P., & Jha, S. (2015).
Molecular tools and bumble bees: Revealing hidden details of ecology and evolution
in a model system. *Molecular Ecology*, *24*(12), 2916–2936. doi: 10.1111/mec.13198
Yang, J., Lee, S. H., Goddard, M. E., & Visscher, P. M. (2011). GCTA: A Tool for Genome-wide
Complex Trait Analysis. *The American Journal of Human Genetics*, *88*(1), 76–82. doi:
10.1016/j.ajhg.2010.11.011
Yang, J., Zaitlen, N. A., Goddard, M. E., Visscher, P. M., & Price, A. L. (2014). Advantages and
pitfalls in the application of mixed-model association methods. *Nature Genetics*,
46(2), 100–106. doi: 10.1038/ng.2876

1119
1120

1121 **AUTHOR CONTRIBUTIONS**

1122 Designed research: MJC, NEM-S, JCG, MTW; Performed research: all authors; Analyzed data:
1123 MJC, JCJ, OW, IB, MK, MTW; Wrote the paper: MJC, MTW.

1124

Figure 1. Population sampling and genetic variation in three *Bombus* species. (A) Map
showing the sampling locations of *Bombus balteatus, B. sylvicola,* and "incognitus" on seven
mountains in the Rocky Mountains, CO, USA. Insert indicates location in the USA.
Neighbour-joining trees of (B) *B. balteatus* and (C) *B. sylvicola* and "incognitus" populations
sampled across the seven mountains, based on genome-wide single nucleotide
polymorphisms. Coloured tips indicate the mountain each sample was collected on, with
colours corresponding to those in (A).
1132

1133 **Figure 2. Principal component analysis of Rocky Mountain populations of three *Bombus*
1134 species.** Plots show the first two principal components from principal component analyses
1135 of (A) *Bombus balteatus*, (B) *B. sylvicola* and (C) "incognitus" populations. Numbers in
1136 brackets on axes indicate the percentage variance explained by each principal component.
1137

Figure 3. Correlations between kinship and geographic distance. Geographic distance
between all pairs of samples plotted against kinship coefficient (a measure of relatedness)
between pairs are shown for (A) *Bombus balteatus*, (B) *B. sylvicola*, and (C) "incognitus",
where each point represents a pair. (D-F) Mantel correlograms showing the Mantel
correlation between kinship and geographic distance over 10 km distance classes for (D)
Bombus balteatus, (E) *B. sylvicola*, and (F) "incognitus". Filled in boxes indicate significant
correlations ($p < 0.01$). There were insufficient data points for calculating Mantel
correlations at distance classes > 60 km for *B. balteatus* and *B. sylvicola*. For "incognitus",
distance class 10-20 km as well as all distance classes > 50 km lacked sufficient data.
1147

1148 **Figure 4. Comparisons of glossa lengths between species, years, and locations.** (A) Least-
1149 squares mean (ls-mean) glossa lengths correcting for body size for the three *Bombus*

species, based on measurements of the 2017 collections. Least-squares mean glossa lengths, correcting for body size and sample location, per sampling period for (B) *B. balteatus* and (C) *B. sylvicola* and "incognitus" combined. In all plots, error bars represent ls-means $\pm$ standard error and significant differences between classes are indicated by asterisks ($p < 0.05$).

Figure 5. Genome-wide association Manhattan plot. (A) The -$\log_{10}$ p-values from an association test using a mixed linear model between allele frequencies and intertegular distance (ID) are plotted for 2,705,361 genome-wide SNPs against their genome positions over the 18 *Bombus balteatus* pseudochromosomes. Red dashed line indicates genome-wide significance after Bonferroni correction for $p = 0.01$. A single SNP on chromosome 10, coloured and circled in red, had a -$\log_{10}$ p-value above this threshold. Green dashed line indicates genome-wide significance after Bonferroni correction for $p = 0.05$. Two neighbouring SNPs located on chromosome 14, coloured green, had -$\log_{10}$ p-values above this threshold. (B) Zoomed-in plot of the significant SNP in chromosome 10 showing its location within an intron of the *CTRB1* gene. Exons of this gene are represented by purple boxes, introns are represented by dashed purple lines.

Figure 6. Estimates of SNP-heritability for two traits in *Bombus* species. The proportion of variance in glossa length and intertegular distance (ID) explained by all genome-wide SNPs (the SNP-based heritability) was calculated using the genome-base restricted maximum likelihood (GREML) method in GCTA. These were calculated separately for *B. sylvicola* (left panel) and *B. balteatus* (right panel). We included intertegular distance as a covariate with glossa length to control for body size. Error bars indicate 95% confidence intervals around estimates. Each analysis was carried out twice, once with all samples (red), and a second time with one from each pair of sisters removed (black), to assess the effect of including highly related samples in the analysis.

Table 1: Assembly metrics for *B. balteatus*, with three other published *Bombus* genomes included for comparison

Species	*B. balteatus*	*B. sylvicola*	*B. terrestris*	*B. impatiens*
Assembly	BBAL_1.0	BSYL_1.0	Bter_1.0	BIMP_2.2
Size (Mbp)†	250.07	252.08	248.65 (236.38)	246.86 (241.98)
Scaffolds (n)	-	-	5,678	5,460
Contigs (n)	336	592	10,672	16,060
Contig N50 (Mbp)	8.60	3.02	0.08	0.06
Contig L50 (n)	12	28	890	54
GC (%)	37.64	38.23	37.51	37.76
Repeat content (%)	17.1	17.9	14.8	17.9
Complete hymenoptera BUSCO genes (%)*	99.0	98.2	96.9	98.3

*BUSCO analysis using the OrthoDB v. 10, Hymenoptera dataset.

† Numbers in brackets represent total ungapped length

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6