***Curtobacterium glycinis* sp. nov. from *Glycine max*, *Curtobacterium gossypii* sp.**

nov. from *Gossypium hirsutum* and *Curtobacterium oryzae* sp. nov. from *Oryza sativa*,

three new *Curtobacterium* species and endophytes from agricultural crops

Sarah Seaton[1], Jacqueline Lemaire[1], Patrik Inderbitzin[1], Victoria Knight-Connoni[1],

James F. White[2], Martha E. Trujillo[3]

[1]Indigo Ag, Inc., 500 Rutherford Avenue, Boston, MA 02129, United States

[2]Department of Plant Biology, Rutgers University, New Brunswick, NJ 08901,

United States

[3]Departamento de Microbiología y Genética, Campus Miguel de Unamuno,

University of Salamanca, Salamanca, Spain

ABSTRACT

Three new *Curtobacterium* species from healthy tissues of agricultural crop plants

in the United States are reported. They are *Curtobacterium glycinis* sp. nov. from soybean

in Missouri, *Curtobacterium gossypii* from cotton in Puerto Rico and corn in Missouri, and

Curtobacterium oryzae sp. nov. from rice in Texas.

Keywords: novel species, taxonomy, bacteria, prokaryotes, endophytes, agriculture,

Actinobacteria, *Microbacteriaceae*

INTRODUCTION

Bacterial endophytes of plants have many beneficial interactions with their hosts

and play a crucial role in promoting plant health (Finkel et al. 2017; Schenk et al. 2012).

Microbiomes of the major agricultural crops including Cotton (*Gossypium hirsutum* L.),

rice (*Oryza sativa* L.) and soybean (*Glycine max* (L.) Merr.) have been characterized

(Edwards et al. 2015; Liu et al. 2019; Longley et al. 2020; Qiao et al. 2017; Roman-Reyna

et al. 2020; Ullah et al. 2019), and beneficial endophytes have been detected (Bertani et al.

2016; de Almeida Lopes et al. 2018; Zhou et al. 2018). Plant-associated microorganisms

are increasingly used for biotechnological applications, including biological control of

plant pathogens, plant growth promotion, or isolation of active compounds (Ryan et al.

2008; Glick 2012; Bouizgarne 2013; Dey et al. 2014).

Curtobacterium is a genus of *Actinobacteria* in the family *Microbacteriaceae*

comprising eight validly published species (Parte 2018). Several of the species have been

isolated from agricultural plants. These include *Curtobacterium albidum* shown to

counteract salt stress in rice (Vimal et al. 2019), *Curtobacterium flaccumfaciens* a

pathogen of bean, beat, tulip and poinsetta (Collins and Jones 1983), *Curtobacterium*

herbarum from grasses (Behrendt et al. 2002), and *Curtobacterium plantarum* from corn

and soybean seed (Dunleavy 1989).

In this study, three *Curtobacterium* strains were isolated from healthy plant tissues

in the United States. Strain OG107 from soybean in Missouri, strain VK105 from cotton in

Puerto Rico, and strain SS108 from rice in Texas. The strains were characterized using

molecular and phenotypic tests to determine their taxonomic placement. Our results

indicate that the strains represent three new species of *Curtobacterium*, and we propose the

names *Curtobacterium glycinis* sp. nov. (OG107), *Curtobacterium gossypii* sp. nov.

(VK105) and *Curtobacterium oryzae* sp. nov. (SS108). These strains are taxonomically

separated from the plant pathogens found within this genus.

METHODS

Isolation. Strains OG107 and SS108 were collected from the roots of healthy field-

grown *Glycine max* in Missouri and *Oryza sativa* seedlings in Texas, United States,

respectively. Plant tissue was washed with a mild detergent to remove particulates,

surface-sterilized with bleach (1% v/v sodium hypochlorite) and ethanol (70% v/v), and

homogenized. Serial dilutions of tissue homogenate were plated on a panel of media types

for endophyte cultivation. Strain OG107, a small (0.7 mm diameter), pale yellow colony,

arose on R2A agar after 5 days of incubation at 24°C, and strain SS108 arose as a non-

descript colony on Actinomycete Isolation Agar. Both strains were streaked to purity and

stored in glycerol (20% v/v) at -80°C until subjected to further testing.

Strain VK105 was isolated as described in Irizarry and White (2017). Briefly, seeds

of wild, non-cultivated *G. hirsutum* plants were collected roadside in Guayama, Puerto

Rico. Seeds were inoculated on Potato Dextrose Agar (PDA) and incubated at room

temperature (25°C). Strain VK105 arose as an irregular yellow-orange pigmented colony.

The colony was streaked to purity and stocked in 25% glycerol at −80°C until further

analysis.

Motility. Strains were tested for flagellar-dependent swimming and swarming
motility on R2A plates solidified with 0.3% and 0.6% agar, respectively. Three
independent colonies were inoculated onto R2A broth and grown for 36 hr at 24°C. Broth
cultures were normalized to an OD600 of 0.1, and 1.5 µl of culture was spotted directly
onto the surface of the motility agar. The diameter of colony expansion was measured for
5 days.

Carbon source utilization. Substrate utilization was assessed using Biolog GenIII
Microplates (Catalogue No. 1030) (Biolog Inc., Hayward, CA). Each bacterium was
inoculated in duplicate plates using Protocol A described by the manufacturer, with the
exception that plates were incubated at 30°C for 36 hr. Respiration leading to reduction of
the tetrazolium indicator was measured by absorbance at 590 nm.

Biochemical analyses. Catalase activity was evaluated by immediate effervescence
after the application of 3% (v/v) hydrogen peroxide solution via the tube method, a
positive reaction was indicated by the production of bubbles. *Staphylococcus aureus*
NCIMB 12702 and *Streptococcus pyogenes* ATCC 19615 were used as positive and
negative controls, respectively. Oxidase activity was evaluated via the oxidation of Kovács
oxidase reagent, 1% (w/v) tetra-methyl-p-phenylenediamine dihydrochloride in water, via
the filter-paper spot method. A positive reaction was indicated when the microorganism's
color changed to dark purple. *Pseudomonas aeruginosa* NCIMB 12469 and *Escherichia*
coli ATCC 25922 were used as positive and negative controls, respectively.

Phylogenetic and genomic analyses. DNA was extracted from pure cultures using
the Omega Mag-Bind Universal Pathogen Kit according to manufacturer's protocol with a
final elution volume of 60µl (Omega Biotek Inc., Norcross, GA). DNA samples were
quantified using Qubit fluorometer (ThermoFisher Scientific, Waltham, MA) and
normalized to 100 ng. DNA was prepped using Nextera DNA Flex Library Prep kit
according to manufacturer's instructions (Illumina Inc., San Diego, CA). DNA libraries
were quantified via qPCR using KAPA Library Quantification kit (Roche Sequencing and
Life Science, Wilmington, MA) and combined in equimolar concentrations into one 24-
sample pool. Libraries were sequenced on a MiSeq using pair-end reads (2x200bp). Reads
were trimmed of adapters and low-quality bases using Cutadapt (version 1.9.1) and
assembled into contigs using MEGAHIT (version 1.1.2) (Li et al. 2015). Reads were
mapped to contigs using Bowtie2 (version 2.3.4) (Langmead and Salzberg 2012), and
contigs were assembled into scaffolds using BESST (2.2.8) (Sahlin et al. 2014).

Average nucleotide identity analyses were performed using the pyani ANIm

algorithm (Richter and Rosselló-Móra 2009) implemented in the MUMmer package

(Kurtz et al. 2004) retrieved from https://github.com/widdowquinn/pyani.

16S rRNA gene sequences were extracted from genome assemblies using barrnap

(Seemann 2019) and submitted to GenBank. Phylogenetic analyses based on the 16S

rRNA gene were performed using FastTree (Price et al. 2010) with a General Time

Reversible substitution model. Taxon sampling for each species is described in the

respective phylogenetic tree figure legend.

Average nucleotide identity (ANI) analyses between genome assemblies were

performed using the pyani ANIm algorithm (Richter and Rosselló-Móra 2009)

implemented in the MUMmer package (Kurtz et al. 2004) retrieved from

https://github.com/widdowquinn/pyani.

Geographic distribution and host range of novel species were inferred by ANI to

assemblies from unidentified species from GenBank (Ciufo et al. 2018) and the Indigo

internal collection. An ANI threshold of ≥95% indicated conspecificity (Chun et al. 2018;

Richter and Rosselló-Móra 2009). Distribution maps were generated in R version 4.0.4 (R

Core Team 2021) with tigris 1.5 (Walker 2021), sf 1.0-2 (Pebesma 2018), ggplot2 3.3.2

(Wickham 2016), cowplot 1.1.1 (Wilke 2020) and other packages.

RESULTS

Phylogenetic and genomic analyses

Curtobacterium glycinis sp. nov. strain OG107

Strain OG107 shared 98.9% 16S rRNA gene sequence identity with

Curtobacterium citreum DSM 20528[T] and less with the remaining *Curtobacterium*

species. A phylogenetic tree using FastTree (Price et al. 2010) confirmed the affiliation of

strain OG107 with the genus *Curtobacterium*. OG107 was most closely related to *C.*

herbarum P 420/07[T] with 84% bootstrap support (Figure 1). The top average nucleotide

identity (ANI) value of OG107 was 89.2% with *C. luteum* ATCC 15830[T]. This value was

well below the threshold for species demarcation (Richter and Rosselló-Móra 2009; Chun

et al. 2018) providing further genomic support that strain OG107 represents a new

genomic species of *Curtobacterium*.

Curtobacterium gossypii sp. nov. strain VK105

Strain VK105 shared 99.8% 16S rRNA gene sequence identity with

Curtobacterium citreum DSM 20528[T] and less with the remaining *Curtobacterium*

species. A phylogenetic tree using FastTree (Price et al. 2010) confirmed the affiliation of

strain VK105 with the genus *Curtobacterium*. VK105 formed a monophyletic group with

the species *C. albidum* and *C. citreum* supported by high bootstrap support (Figure 1) and

was equally related to either of the two species. Average nucleotide identity (ANI) values

of *C. albidum* DSM 20512[T] and *C. citreum* DSM 20528[T] to VK105 were both 87.7%.

These values are well below the threshold for species demarcation (Richter and Rosselló-

Móra 2009; Chun et al. 2018) providing further genomic support that strain VK105

represents a new genomic species of *Curtobacterium.*

Curtobacterium oryzae sp. nov. strain SS108

Strain SS108 shared 99.6% 16S rRNA gene sequence identity with *Curtobacterium*

luteum strain DSM 20542[T] and less with the remaining *Curtobacterium* species. A

phylogenetic tree using FastTree (Price et al. 2010) confirmed the affiliation of strain

SS108 with the genus *Curtobacterium*. SS108 was most closely related to *C. luteum* with

79% bootstrap support (Figure 1). The top average nucleotide identity (ANI) value of

SS108 was 89.2% with *C. luteum* DSM 20542[T]. This value was well below the threshold

for species demarcation (Richter and Rosselló-Móra 2009; Chun et al. 2018) providing

further genomic support that strain SS108 represents a new genomic species of

Curtobacterium.

Geographic distribution and host range

Geographic distribution and host range of the novel species was inferred by

comparison to congeneric genome assemblies from the Indigo internal collection. Hits

included the Indigo strain *Curtobacterium gossypii* strain JL114 (ANI: 99.99%; query

coverage: 99.33%) collected from healthy *Zea mays* plants in Missouri. Known geographic

distributions of the novel species from culturing is illustrated in Figure 2, Figure 3 and

Figure 4, and substrates are compiled in Table 1.

Morphology, physiology and biochemical characteristics

Curtobacterium glycinis sp. nov. strain OG107

Strain OG107 stained Gram-positive. Cells had a slightly curved rod shape (0.2-0.4

µm in width and 1-3 µm in length); V and Y shaped forms were observed (Figure 5). They

were aerobic, non-sporulating and showed motility. On nutrient agar, colonies were cream,

entire, punctiform and convex. After 48 hr, good growth is observed on TSA, R2A and NA

at 22 and 30°C. Growth was poor to moderate on the same media and incubation times at

37°C. Growth is observed when media are supplemented with 4% NaCl (w/v). Strain OG

107 was positive for catalase, but not for oxidase or urease. Being aerobic, the strain has an

oxidative metabolism and according to the Biolog system, it is capable of using a range of

substrates as carbon sources that include sugars, alcohol sugars and organic acids.

Compared to its closest phylogenetic neighbor, *C. citreum* JCM 1345[T], OG17 uses N-

Acetyl-D-Galactosamine, D-arabitol, gentobiose, *myo*-inositol, mannitol and turanose as

carbon sources. These tests help to differentiate between the two strains. A detailed

description is given in Table 2 and the species description.

Curtobacterium gossypii sp. nov. strain VK105

The cells of strain VK105 stained Gram-positive, formed short irregular rods (0.4-0.6 µm

in width and 2-4 µm in length), with cells diving by bending (Figure 6). Y shaped-forms

were sometimes visible. Cells were non-motile and did not produce spores. Strain VK105

showed good aerobic growth on R2A and TSA agars. On both media, yellow to orange

colonies were observed after 2 days incubation at 22 and 30°C. At 7 days, colonies were

circular, smooth, glistening and slightly convex. The new strain produced catalase but not

oxidase. Strain VK105 has an oxidative metabolism and uses a range of substrates as

carbon sources that are also useful for to differentiate against its closest phylogenetic

neighbors, *Curtocabterium citreum* and *C. albidum*, namely D-arabitol, L-fucose, D-

glucuronic acid, and *myo*-inositol. These and other physiological characteristics are

presented in Table 2 and the species description.

Curtobacterium oryzae sp. nov. strain SS108

Cells stained-Gram positive, are non-motile, non-spore-forming and rod shaped or

slightly curved. In many cases, two cells remained together and formed V shapes (Figure

7). Average cell size ranged from 0.2-0.4 µm in width and 1.5 x 2 µm in length. Colonies

of strain SS108 were light yellow, entire, round and raised when grown on nutrient agar.

After 2 days, good growth was observed on TSA and R2A agars at 22, 30 and 37°C. On

nutrient agar, good growth was obtained at 30°C after 2 days and was moderate at 22 and

37°C. However, after 7 days, good growth was also seen at 22°C. Aerobic growth.

Tolerates 4% NaCl (w/v). Enzyme activity was detected for catalase, but not for oxidase or

urease. Several carbon sources are useful to differentiate between the new strain and *C.*

luteum JCM 1480^T and include the utilization of N-Acetyl-D-Galactosamine, D-arabitol,

gentobiose, *myo*-inositol and D-mannitol among others (Table 2). Other phenotypic

characteristics are given in the species description.

DESCRIPTION OF CURTOBACTERIUM GLYCINIS SP. NOV.

Curtobacterium glycinis (gly.ci'nis. N.L. fem. gen. n. *glycinis* of *Glycine max*, the

soybean, referring to the origin of the type strain)

Cells are Gram-stain positive, non-spore forming, motile and rod-shaped with V

and Y shapes produced. Aerobic and chemoorganotrophic. Colonies are circular, smooth,

glistening and slightly convex. On R2A and nutrient agar, colonies are cream color and

entire, punctiform and convex. Growth is abundant at 22 and 30°C but moderate at 37°C.

Grows in the presence of 4% NaCl. Growth is observed between pH 5-7. Positive for

catalase; negative for oxidase and urease. The following substrates are used as carbon

sources: N-acetyl-D-glucosamine, N-acetyl-D-galactosamine, D-arabitol, D-cellobiose, L-

fucose, gentiobiose, *myo*-inositol, α-D-lactose, D-mannitol, D-melibiose, β-methyl-D-

glucoside, D-raffinose, turanose, D-gluconic acid, D-glucuronic acid, L-alanine, L-

glutamic acid (weak), L-rhamnose, D-Glucose-6-PO4.

The type strain is resistant to nalidixic acid and aztreonam.

The type strain OG 17^T was isolated from the roots of healthy field-grown *Glycine*

max in Missouri, USA.

DESCRIPTION OF CURTOBACTERIUM GOSSYPII SP. NOV.

*Curtobacterium gossypii (*gos.sy′pi.i. N.L. gen. n. *gossypii* of *Gossypium,* the

generic name of cotton, referring to the origin of the type strain)

Cells are Gram-stain positive, non-motile, non-spore-forming, coccoid or rod-

shaped. Aerobic and chemoorganotrophic. Yellow to orange colonies on R2A and TSA

agars. Growth on these media is observed after 48 h. Catalase is produced but not oxidase.

Grows in the presence of 1 - 4 % NaCl; weak at 8% and at pH 5-7. The following
substrates are used as sole carbon sources: dextrin, D-maltose, D-trehalose, D-cellobiose,
gentiobiose, sucrose, D-turanose, stachyose, D-raffinose, α-D-lactose, D-melibiose, β-
methylD-glucoside, D-salicin, N-acetyl-D-glucosamine, α-D-glucose, D-mannose, D-
fructose, D-galactose, inosine, sodium lactate, D-mannitol, D-arabitol, *myo*-inositol,
glycerol, L-aspartic acid, L-glutamic acid, pectin, D-arabitol, D-gluconic acid, D-
glucuronic acid and glucuronamide.

The type strain is resistant to nalidixic acid and aztreonam.

The type strain, VK105[T], was isolated from seeds of wild cotton collected in Puerto
Rico, USA.

DESCRIPTION OF CURTOBACTERIUM ORYZAE SP. NOV.

Curtobacterium oryzae (o.ry′zae. N.L. gen. n. *oryzae* of rice, referring to the origin of the
type strain)

Cells are Gram-stain positive, non-motile, non-spore-forming and rod shaped or
slightly curved; V shapes are formed. Aerobic and chemoorganotrophic. Colonies are light
yellow, entire, round and raised on nutrient agar. Good growth is obtained at 22, 30 and
37°C on the media tested. Grows on media supplemented with 4% NaCl and at pH 5-7.
Positive for catalase, but negative for oxidase or urease. The following substrates are used
as carbon sources: N-Acetyl-D-Glucosamine, N-Acetyl-D-Galactosamine, N-Acetyl
Neuraminic Acid, D-Arabitol, D-Cellobiose, L-Fucose, Gentiobiose, myo-Inositol, α-D-
Lactose, D-Mannitol, D-Melibiose, β-Methyl-D-Glucoside, D-Raffinose, D-Sorbitol,
Turanose, D-Gluconic Acid, D-Glucuronic Acid, L-Alanine, L-Glutamic Acid, L-
Rhamnose, D-Glucose-6-PO4 and Acetic Acid.

The type strain is resistant to nalidixic acid and aztreonam.

The type strain, SS108[T], was isolated from *Oryza sativa* seedlings collected in
Texas, USA.

ACKNOWLEDGEMENTS

We would like to thank Professor Aharon Oren, The Hebrew University of

Jerusalem, for checking Latin species names. Support was provided by USDA-NIFA

Multistate Project W4147 and the New Jersey Agricultural Experiment Station.

LITERATURE CITED

Aizawa, T., Ve, N. B., Kimoto, K., Iwabuchi, N., Sumida, H., Hasegawa, I., et al. 2007.
Curtobacterium ammoniigenes sp. nov., an ammonia-producing bacterium isolated from
plants inhabiting acidic swamps in actual acid sulfate soil areas of Vietnam. Int. J. Syst.
Evol. Microbiol. 57:1447–1452.
de Almeida Lopes, K. B., Carpentieri-Pipolo, V., Fira, D., Balatti, P. A., López, S. M. Y.,
Oro, T. H., et al. 2018. Screening of bacterial endophytes as potential biocontrol agents
against soybean diseases. J. Appl. Microbiol. 125:1466–1481.
Behrendt, U., Ulrich, A., Schumann, P., Naumann, D., and Suzuki, K. 2002. Diversity of
grass-associated Microbacteriaceae isolated from the phyllosphere and litter layer after
mulching the sward; polyphasic characterization of *Subtercola pratensis* sp. nov.,
Curtobacterium herbarum sp. nov. and *Plantibacter flavus* gen. nov., sp. nov. Int. J. Syst.
Evol. Microbiol. 52:1441–1454.
Bertani, I., Abbruscato, P., Piffanelli, P., Subramoni, S., and Venturi, V. 2016. Rice
bacterial endophytes: isolation of a collection, identification of beneficial strains and
microbiome analysis. Environ. Microbiol. Rep. 8:388–398.
Bouizgarne, B. 2013. Bacteria for plant growth promotion and disease management. In
Bacteria in Agrobiology: Disease Management, Springer, p. 15–47.
Chun, J., Oren, A., Ventosa, A., Christensen, H., Arahal, D. R., da Costa, M. S., et al.
2018. Proposed minimal standards for the use of genome data for the taxonomy of
prokaryotes. Int. J. Syst. Evol. Microbiol. 68:461–466.
Ciufo, S., Kannan, S., Sharma, S., Badretdin, A., Clark, K., Turner, S., et al. 2018. Using
average nucleotide identity to improve taxonomic assignments in prokaryotic genomes at
the NCBI. Int. J. Syst. Evol. Microbiol. 68:2386–2392.
Collins, M., and Jones, D. 1983. Reclassification of *Corynebacterium flaccumfaciens*,
Corynebacterium betae, *Corynebacterium oortii* and *Corynebacterium poinsettiae* in the
genus *Curtobacterium*, as *Curtobacterium flaccumfaciens* comb. nov. Microbiology.
129:3545–3548.
Dey, R., Pal, K. K., and Tilak, K. 2014. Plant growth promoting rhizobacteria in crop
protection and challenges. In *Future Challenges in Crop Protection Against Fungal
Pathogens*, Springer, p. 31–58.
Dunleavy, J. 1989. *Curtobacterium plantarum* sp. nov. is ubiquitous in plant leaves and is
seed transmitted in soybean and corn. Int. J. Syst. Evol. Microbiol. 39:240–249.
Edwards, J., Johnson, C., Santos-Medellín, C., Lurie, E., Podishetty, N. K., Bhatnagar, S.,
et al. 2015. Structure, variation, and assembly of the root-associated microbiomes of rice.
Proc. Natl. Acad. Sci. 112:E911–E920.
Finkel, O. M., Castrillo, G., Paredes, S. H., González, I. S., and Dangl, J. L. 2017.
Understanding and exploiting plant beneficial microbes. Curr. Opin. Plant Biol. 38:155–
163.
Glick, B. R. 2012. Plant growth-promoting bacteria: mechanisms and applications.
Scientifica. 2012.

Irizarry, I., and White, J. 2017. Application of bacteria from non-cultivated plants to promote growth, alter root architecture and alleviate salt stress of cotton. J. Appl. Microbiol. 122:1110–1120.

Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C., et al. 2004. Versatile and open software for comparing large genomes. Genome Biol. 5:R12.

Langmead, B., and Salzberg, S. L. 2012. Fast gapped-read alignment with bowtie 2. Nat Methods. 9 Available at: https://doi.org/10.1038/nmeth.1923.

Li, D., Liu, C.-M., Luo, R., Sadakane, K., and Lam, T.-W. 2015. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics. 31:1674–1676.

Liu, F., Hewezi, T., Lebeis, S. L., Pantalone, V., Grewal, P. S., and Staton, M. E. 2019. Soil indigenous microbiome and plant genotypes cooperatively modify soybean rhizosphere microbiome assembly. BMC Microbiol. 19:1–19.

Longley, R., Noel, Z. A., Benucci, G. M. N., Chilvers, M., Trail, F., and Bonito, G. 2020. Crop management impacts the soybean (*Glycine max*) microbiome. Front. Microbiol. 11:1116.

Parte, A. C. 2018. LPSN-List of prokaryotic names with standing in nomenclature (bacterio.net), 20 years on. Int. J. Syst. Evol. Microbiol. 68:1825–1829.

Pebesma, E. 2018. Simple features for R: standardized support for spatial vector data. R J. 10:439–446.

Price, M. N., Dehal, P. S., and Arkin, A. P. 2010. FastTree 2 - approximately maximum-likelihood trees for large alignments. PLoS ONE. 5:e9490.

Qiao, Q., Wang, F., Zhang, J., Chen, Y., Zhang, C., Liu, G., et al. 2017. The variation in the rhizosphere microbiome of cotton with soil type, genotype and developmental stage. Sci. Rep. 7:1–10.

R Core Team. 2021. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. Available at: https://www.r-project.org/.

Richter, M., and Rosselló-Móra, R. 2009. Shifting the genomic gold standard for the prokaryotic species definition. Proc. Natl. Acad. Sci. 106:19126–19131.

Roman-Reyna, V., Pinili, D., Borja, F. N., Quibod, I. L., Groen, S. C., Alexandrov, N., et al. 2020. Characterization of the leaf microbiome from whole-genome sequencing data of the 3000 rice genomes project. Rice. 13:1–8.

Ryan, R. P., Germaine, K., Franks, A., Ryan, D. J., and Dowling, D. N. 2008. Bacterial endophytes: recent developments and applications. FEMS Microbiol. Lett. 278:1–9.

Sahlin, K., Vezzi, F., Nystedt, B., Lundeberg, J., and Arvestad, L. 2014. BESST-efficient scaffolding of large fragmented assemblies. BMC Bioinformatics. 15:281.

Schenk, P. M., Carvalhais, L. C., and Kazan, K. 2012. Unraveling plant-microbe interactions: can multi-species transcriptomics help? Trends Biotechnol. 30:177–184.

Seemann, T. 2019. *barrnap 0.9: rapid ribosomal RNA prediction*. Available at: https://github.com/tseemann/barrnap.

Ullah, A., Akbar, A., Luo, Q., Khan, A. H., Manghwar, H., Shaban, M., et al. 2019. Microbiome diversity in cotton rhizosphere under normal and drought conditions. Microb. Ecol. 77:429–439.

Vimal, S. R., Patel, V. K., and Singh, J. S. 2019. Plant growth promoting *Curtobacterium albidum* strain SRV4: an agriculturally important microbe to alleviate salinity stress in paddy plants. Ecol. Indic. 105:553–562.

Walker, K. 2021. *tigris: Load Census TIGER/Line Shapefiles*. Available at: https://CRAN.R-project.org/package=tigris.

Wickham, H. 2016. *ggplot2: elegant graphics for data analysis*. New York: Springer
Verlag. Available at: https://ggplot2.tidyverse.org.
Wilke, C. O. 2020. *cowplot: streamlined plot theme and plot annotations for "ggplot2."*
Available at: https://CRAN.R-project.org/package=cowplot.
Zhou, W., Wheeler, T. A., Starr, J. L., Valencia, C. U., and Sword, G. A. 2018. A fungal
endophyte defensive symbiosis affects plant-nematode interactions in cotton. Plant Soil.
422:251–266.
369

 FIGURES AND TABLES

 Figure 1. Phylogenetic 16S rRNA gene tree of *Curtobacterium glycinis* sp. nov.

 strain OG107, *Curtobacterium gossypii* sp. nov. strain VK105, *Curtobacterium oryzae* sp.

 nov. strain SS108 and relatives generated using FastTree (Price et al. 2010). All validly

 published *Curtobacterium* species were included in the tree, the tree is rooted with

 Frondihabitans australicus. New species are in bold. Strain identifiers and GenBank

 accession numbers follow species names, T stands for 'type'. Support values above 70%

 are given by the branches. *Curtobacterium glycinis* is most closely related to *C. herbarum*

 with 84% support, *Curtobacterium gossypii* to *C. albidum* and *C. citreum* with 82%

 support and *Curtobacterium oryzae* to *C. luteum* with 79% support. Branch lengths are

 proportional to the changes along the branches, a scale bar is provided.

Figure 2. Geographic distribution of *Curtobacterium glycinis* sp. nov. based on culturing.

Dark blue indicates state of origin for type strain.

Figure 3. Geographic distribution of *Curtobacterium gossypii* sp. nov. based on culturing.

Dark blue indicates state of origin for type strain, light blue additional strain. Inset: Puerto

Rico (not to scale).

Figure 4. Geographic distribution of *Curtobacterium oryzae* sp. nov. based on culturing.

Dark blue indicates state of origin for type strain.

Figure 5. Morphology of *Curtobacterium glycinis* sp. nov. strain OG107 depicted following Gram stain using bright field microscopy. Bar = 10 µm.

Figure 6. Morphology of *Curtobacterium gossypii* sp. nov. strain VK105 depicted

following Gram stain using bright field microscopy. Bar = 10 µm.

Figure 7. Morphology of *Curtobacterium oryzae* sp. nov. strain SS108 depicted following

Gram stain using bright field microscopy. Bar = 10 µm.

TABLES

Table 1. Substrates of novel species based on culturing.

Species	Substrate[1]
Curtobacterium glycinis	*Glycine max*
Curtobacterium gossypii	*Gossypium hirsutum, Zea mays*
Curtobacterium oryzae	*Oryza sativa*

430 [1] No disease symptoms reported for strains isolated from plants.

Table 2. Physiological characteristics of newly described *Curtobacterium glycinis*, *C. gossypii* and *C. oryzae* species and related *Curtobacterium* type strains. All strains were positive for the assimilation of dextrin, D-fructose, D-galactose, α-D-glucose, D-mannose and sucrose (data not shown).

	C. gossypii VK105	*C. oryzae* OG107	*C. glycinis* SS108	*C. citreum* JCM 1345^T	*C. pusillum* JCM 1350^T	*C. luteum* JCM 1480^T	*C. albidum* NBRC 15078^T	*C. herbarum* DSM 14013^T
General characteristics:								
Colony pigment		light yellow	ivory	yellow	yellow	yellow	ivory	orange
Motility	-	w	w	+	+	+	-	+
Oxidase	-	-	-	-	-	-	-	-
Catalase	+	+	+	+	+	+	+	+
Urease	-	-	-	-	-	-	-	nr
Growth with 4% NaCl	+	+	+	nr	nr	nr	nr	nr
Growth with 8% NaCl	+	-	-	nr	nr	nr	nr	nr
Utilization of:								
N-Acetyl-D-Glucosamine	+	+	+	+	-	+	w	nr
N-Acetyl-D-Galactosamine	-	+	+	-	-	-	-	nr
N-Acetyl Neuraminic Acid	-	+	-	nr	nr	nr	nr	nr
D-Arabitol	+	+	+	-	w	-	-	nr
D-Cellobiose	+	+	+	+	-	+	+	nr
L-Fucose	w	+	+	+	-	+	-	nr
Gentiobiose	+	+	+	-	+	-	+	nr
myo-Inositol	+	+	+	-	-	-	-	nr
α-D-Lactose	+	+	+	+	-	+	+	nr
D-Mannitol	+	+	+	-	+	-	w	+
D-Melibiose	+	+	+	+	+	+	w	+
β-Methyl-D-Glucoside	+	+	+	-	+	-	w	nr
D-Raffinose	+	+	+	-	-	-	+	nr

D-Sorbitol	w	+	-	-	+	-	+	+
Turanose	+	+	+	-	+	-	+	nr
D-Gluconic Acid	+	+	+	-	+	-	+	nr
D-Glucuronic Acid	+	+	+	-	-	-	-	nr
L-Alanine	+	+	+	-	-	-	w	nr
L-Glutamic Acid	+	+	w	+	-	-	-	nr
L-Rhamnose	w	+	+	-	-	-	-	+
D-Glucose-6-PO4	-	+	+	-	-	-	-	nr
Acetic Acid	w	+	-	-	-	-	-	nr

+, Positive; w, Weakly positive; -, Negative; nr, Not reported.

Data from this study, Aizawa et al. (2007) and Behrendt et al. (2002).