1   ***Curtobacterium glycinis* sp. nov. from *Glycine max*, *Curtobacterium gossypii* sp.**
2   **nov. from *Gossypium hirsutum* and *Curtobacterium oryzae* sp. nov. from *Oryza sativa*,**
3   **three new *Curtobacterium* species and endophytes from agricultural crops**
4

5   Sarah Seaton[1], Jacqueline Lemaire[1], Patrik Inderbitzin[1], Victoria Knight-Connoni[1],
6   James F. White[2], Martha E. Trujillo[3]
7

8   [1]Indigo Ag, Inc., 500 Rutherford Avenue, Boston, MA 02129, United States
9   [2]Department of Plant Biology, Rutgers University, New Brunswick, NJ 08901,
10   United States
11   [3]Departamento de Microbiología y Genética, Campus Miguel de Unamuno,
12   University of Salamanca, Salamanca, Spain

13
14

15   **ABSTRACT**
16   Three new *Curtobacterium* species from healthy tissues of agricultural crop plants
17   in the United States are reported. They are *Curtobacterium glycinis* sp. nov. from soybean
18   in Missouri, *Curtobacterium gossypii* from cotton in Puerto Rico and *Curtobacterium*
19   *oryzae* sp. nov. from rice in Texas.
20

21   Keywords: novel species, taxonomy, bacteria, prokaryotes, endophytes, agriculture,
22   *Actinobacteria*, *Microbacteriaceae*

23
24

INTRODUCTION
Bacterial endophytes of plants have many beneficial interactions with their hosts
and play a crucial role in promoting plant health (Finkel et al. 2017; Schenk et al. 2012).
Microbiomes of the major agricultural crops including Cotton (*Gossypium hirsutum* L.),
rice (*Oryza sativa* L.) and soybean (*Glycine max* (L.) Merr.) have been characterized
(Edwards et al. 2015; Liu et al. 2019; Longley et al. 2020; Qiao et al. 2017; Roman-Reyna
et al. 2020; Ullah et al. 2019), and beneficial endophytes have been detected (Bertani et al.
2016; de Almeida Lopes et al. 2018; Zhou et al. 2018). Plant-associated microorganisms
are increasingly used for biotechnological applications, including biological control of

plant pathogens, plant growth promotion, or isolation of active compounds (Ryan et al.

2008; Glick 2012; Bouizgarne 2013; Dey et al. 2014).

Curtobacterium is a genus of *Actinobacteria* in the family *Microbacteriaceae*

comprising eight validly published species (Parte 2018). Several of the species have been

isolated from agricultural plants. These include *Curtobacterium albidum* shown to

counteract salt stress in rice (Vimal et al. 2019), *Curtobacterium flaccumfaciens* a

pathogen of bean, beat, tulip and poinsetta (Collins and Jones 1983), *Curtobacterium*

herbarum from grasses (Behrendt et al. 2002), and *Curtobacterium plantarum* from corn

and soybean seed (Dunleavy 1989).

In this study, three *Curtobacterium* strains were isolated from healthy plant tissues

in the United States. Strain OG107 from soybean in Missouri, strain VK105 from cotton in

Puerto Rico, and strain SS108 from rice in Texas. The strains were characterized using

molecular and phenotypic tests to determine their taxonomic placement. Our results

indicate that the strains represent three new species of *Curtobacterium*, and we propose the

names *Curtobacterium glycinis* sp. nov. (OG107), *Curtobacterium gossypii* sp. nov.

(VK105) and *Curtobacterium oryzae* sp. nov. (SS108). These strains are taxonomically

separated from the plant pathogens found within this genus.

METHODS

Isolation. Strains OG107 and SS108 were collected from the roots of healthy field-

grown *Glycine max* in Missouri and *Oryza sativa* seedlings in Texas, United States,

respectively. Plant tissue was washed with a mild detergent to remove particulates,

surface-sterilized with bleach (1% v/v sodium hypochlorite) and ethanol (70% v/v), and

homogenized. Serial dilutions of tissue homogenate were plated on a panel of media types

for endophyte cultivation. Strain OG107, a small (0.7 mm diameter), pale yellow colony,

arose on R2A agar after 5 days of incubation at 24°C, and strain SS108 arose as a non-

descript colony on Actinomycete Isolation Agar. Both strains were streaked to purity and

stored in glycerol (20% v/v) at -80°C until subjected to further testing.

Strain VK105 was isolated as described in Irizarry and White (2017). Briefly, seeds

of wild, non-cultivated *G. hirsutum* plants were collected roadside in Guayama, Puerto

Rico. Seeds were inoculated on Potato Dextrose Agar (PDA) and incubated at room

temperature (25°C). Strain VK105 arose as an irregular yellow-orange pigmented colony.

The colony was streaked to purity and stocked in 25% glycerol at −80°C until further

analysis.

Motility. Strains were tested for flagellar-dependent swimming and swarming

motility on R2A plates solidified with 0.3% and 0.6% agar, respectively. Three

independent colonies were inoculated onto R2A broth and grown for 36 hr at 24°C. Broth

cultures were normalized to an OD600 of 0.1, and 1.5 µl of culture was spotted directly

onto the surface of the motility agar. The diameter of colony expansion was measured for

5 days.

Carbon source utilization. Substrate utilization was assessed using Biolog GenIII

Microplates (Catalogue No. 1030) (Biolog Inc., Hayward, CA). Each bacterium was

inoculated in duplicate plates using Protocol A described by the manufacturer, with the

exception that plates were incubated at 30°C for 36 hr. Respiration leading to reduction of

the tetrazolium indicator was measured by absorbance at 590 nm.

Biochemical analyses. Catalase activity was evaluated by immediate effervescence

after the application of 3% (v/v) hydrogen peroxide solution via the tube method, a

positive reaction was indicated by the production of bubbles. *Staphylococcus aureus*

NCIMB 12702 and *Streptococcus pyogenes* ATCC 19615 were used as positive and

negative controls, respectively. Oxidase activity was evaluated via the oxidation of Kovács

oxidase reagent, 1% (w/v) tetra-methyl-p-phenylenediamine dihydrochloride in water, via

the filter-paper spot method. A positive reaction was indicated when the microorganism's

color changed to dark purple. *Pseudomonas aeruginosa* NCIMB 12469 and *Escherichia*

coli ATCC 25922 were used as positive and negative controls, respectively.

Phylogenetic and genomic analyses. DNA was extracted from pure cultures using

the Omega Mag-Bind Universal Pathogen Kit according to manufacturer's protocol with a

final elution volume of 60µl (Omega Biotek Inc., Norcross, GA). DNA samples were

quantified using Qubit fluorometer (ThermoFisher Scientific, Waltham, MA) and

normalized to 100 ng. DNA was prepped using Nextera DNA Flex Library Prep kit

according to manufacturer's instructions (Illumina Inc., San Diego, CA). DNA libraries

were quantified via qPCR using KAPA Library Quantification kit (Roche Sequencing and

Life Science, Wilmington, MA) and combined in equimolar concentrations into one 24-

sample pool. Libraries were sequenced on a MiSeq using pair-end reads (2x200bp). Reads

were trimmed of adapters and low-quality bases using Cutadapt (version 1.9.1) and

assembled into contigs using MEGAHIT (version 1.1.2) (Li et al. 2015). Reads were

mapped to contigs using Bowtie2 (version 2.3.4) (Langmead and Salzberg 2012), and

contigs were assembled into scaffolds using BESST (2.2.8) (Sahlin et al. 2014).

Average nucleotide identity analyses were performed using the pyani ANIm

algorithm (Richter and Rosselló-Móra 2009) implemented in the MUMmer package

(Kurtz et al. 2004) retrieved from https://github.com/widdowquinn/pyani.

16S rRNA gene sequences were extracted from genome assemblies using barrnap

(Seemann 2019) and submitted to GenBank. Phylogenetic analyses based on the 16S

rRNA gene were performed using FastTree (Price et al. 2010) with a General Time

Reversible substitution model. Taxon sampling for each species is described in the

respective phylogenetic tree figure legend.

RESULTS

Phylogenetic and genomic analyses

Curtobacterium glycinis sp. nov. strain OG107

Strain OG107 shared 98.9% 16S rRNA gene sequence identity with

Curtobacterium citreum DSM 20528[T] and less with the remaining *Curtobacterium*

species. A phylogenetic tree using FastTree (Price et al. 2010) confirmed the affiliation of

strain OG107 with the genus *Curtobacterium*. OG107 was most closely related to *C.*

herbarum P 420/07[T] with 84% bootstrap support (Figure 1). The top average nucleotide

identity (ANI) value of OG107 was 89.2% with *C. luteum* ATCC 15830[T]. This value was

well below the threshold for species demarcation (Richter and Rosselló-Móra 2009; Chun

et al. 2018) providing further genomic support that strain OG107 represents a new

genomic species of *Curtobacterium*.

Curtobacterium gossypii sp. nov. strain VK105

Strain VK105 shared 99.8% 16S rRNA gene sequence identity with

Curtobacterium citreum DSM 20528[T] and less with the remaining *Curtobacterium*

species. A phylogenetic tree using FastTree (Price et al. 2010) confirmed the affiliation of

strain VK105 with the genus *Curtobacterium*. VK105 formed a monophyletic group with

the species *C. albidum* and *C. citreum* supported by high bootstrap support (Figure 1) and

was equally related to either of the two species. Average nucleotide identity (ANI) values

of *C. albidum* DSM 20512[T] and *C. citreum* DSM 20528[T] to VK105 were both 87.7%.

These values are well below the threshold for species demarcation (Richter and Rosselló-

Móra 2009; Chun et al. 2018) providing further genomic support that strain VK105

represents a new genomic species of *Curtobacterium*.

135    *Curtobacterium oryzae* sp. nov. strain SS108

136    Strain SS108 shared 99.6% 16S rRNA gene sequence identity with *Curtobacterium*

137 *luteum* strain DSM 20542^T and less with the remaining *Curtobacterium* species. A

138 phylogenetic tree using FastTree (Price et al. 2010) confirmed the affiliation of strain

139 SS108 with the genus *Curtobacterium*. SS108 was most closely related to *C. luteum* with

140 79% bootstrap support (Figure 1). The top average nucleotide identity (ANI) value of

141 SS108 was 89.2% with *C. luteum* DSM 20542^T. This value was well below the threshold

142 for species demarcation (Richter and Rosselló-Móra 2009; Chun et al. 2018) providing

143 further genomic support that strain SS108 represents a new genomic species of

144 *Curtobacterium.*

146    **Morphology, physiology and biochemical characteristics**

147    *Curtobacterium glycinis* sp. nov. strain OG107

148    Strain OG107 stained Gram-positive. Cells had a slightly curved rod shape (0.2-0.4

149 µm in width and 1-3 µm in length); V and Y shaped forms were observed (Figure 2). They

150 were aerobic, non-sporulating and showed motility. On nutrient agar, colonies were cream,

151 entire, punctiform and convex. After 48 hr, good growth is observed on TSA, R2A and NA

152 at 22 and 30°C. Growth was poor to moderate on the same media and incubation times at

153 37°C. Growth is observed when media are supplemented with 4% NaCl (w/v). Strain OG

154 107 was positive for catalase, but not for oxidase or urease. Being aerobic, the strain has an

155 oxidative metabolism and according to the Biolog system, it is capable of using a range of

156 substrates as carbon sources that include sugars, alcohol sugars and organic acids.

157 Compared to its closest phylogenetic neighbor, *C. citreum* JCM 1345^T, OG17 uses N-

158 Acetyl-D-Galactosamine, D-arabitol, gentobiose, *myo*-inositol, mannitol and turanose as

159 carbon sources. These tests help to differentiate between the two strains. A detailed

160 description is given in Table 1 and the species description.

161    *Curtobacterium gossypii* sp. nov. strain VK105

162 The cells of strain VK105 stained Gram-positive, formed short irregular rods (0.4-0.6 µm

163 in width and 2-4 µm in length), with cells diving by bending (Figure 3). Y shaped-forms

164 were sometimes visible. Cells were non-motile and did not produce spores. Strain VK105

165 showed good aerobic growth on R2A and TSA agars. On both media, yellow to orange

166 colonies were observed after 2 days incubation at 22 and 30°C. At 7 days, colonies were

167 circular, smooth, glistening and slightly convex. The new strain produced catalase but not

168 oxidase. Strain VK105 has an oxidative metabolism and uses a range of substrates as

carbon sources that are also useful for to differentiate against its closest phylogenetic

neighbors, *Curtocabterium citreum* and *C. albidum*, namely D-arabitol, L-fucose, D-

glucuronic acid, and *myo*-inositol. These and other physiological characteristics are

presented in Table 1 and the species description.

Curtobacterium oryzae sp. nov. strain SS108

Cells stained-Gram positive, are non-motile, non-spore-forming and rod shaped or

slightly curved. In many cases, two cells remained together and formed V shapes (Figure

4). Average cell size ranged from 0.2-0.4 µm in width and 1.5 x 2 µm in length. Colonies

of strain SS108 were light yellow, entire, round and raised when grown on nutrient agar.

After 2 days, good growth was observed on TSA and R2A agars at 22, 30 and 37°C. On

nutrient agar, good growth was obtained at 30°C after 2 days and was moderate at 22 and

37°C. However, after 7 days, good growth was also seen at 22°C. Aerobic growth.

Tolerates 4% NaCl (w/v). Enzyme activity was detected for catalase, but not for oxidase or

urease. Several carbon sources are useful to differentiate between the new strain and *C.*

luteum JCM 1480^T and include the utilization of N-Acetyl-D-Galactosamine, D-arabitol,

gentobiose, *myo*-inositol and D-mannitol among others (Table 1). Other phenotypic

characteristics are given in the species description.

DESCRIPTION OF CURTOBACTERIUM GLYCINIS SP. NOV.

Curtobacterium glycinis (gly.ci'nis. N.L. fem. gen. n. *glycinis* of *Glycine max*, the

soybean, referring to the origin of the type strain)

Cells are Gram-stain positive, non-spore forming, motile and rod-shaped with V

and Y shapes produced. Aerobic and chemoorganotrophic. Colonies are circular, smooth,

glistening and slightly convex. On R2A and nutrient agar, colonies are cream color and

entire, punctiform and convex. Growth is abundant at 22 and 30°C but moderate at 37°C.

Grows in the presence of 4% NaCl. Growth is observed between pH 5-7. Positive for

catalase; negative for oxidase and urease. The following substrates are used as carbon

sources: N-acetyl-D-glucosamine, N-acetyl-D-galactosamine, D-arabitol, D-cellobiose, L-

fucose, gentiobiose, *myo*-inositol, α-D-lactose, D-mannitol, D-melibiose, β-methyl-D-

glucoside, D-raffinose, turanose, D-gluconic acid, D-glucuronic acid, L-alanine, L-
glutamic acid (weak), L-rhamnose, D-Glucose-6-PO4.

The type strain is resistant to nalidixic acid and aztreonam.

The type strain OG 17^T was isolated from the roots of healthy field-grown *Glycine*
max in Missouri, USA.

DESCRIPTION OF CURTOBACTERIUM GOSSYPII SP. NOV.

*Curtobacterium gossypii (*gos.sy′pi.i. N.L. gen. n. *gossypii* of *Gossipium,* the
generic name of cotton, referring to the origin of the type strain)

Cells are Gram-stain positive, non-motile, non-spore-forming, coccoid or rod-
shaped. Aerobic and chemoorganotrophic. Yellow to orange colonies on R2A and TSA
agars. Growth on these media is observed after 48 h. Catalase is produced but not oxidase.
Grows in the presence of 1 - 4 % NaCl; weak at 8% and at pH 5-7. The following
substrates are used as sole carbon sources: dextrin, D-maltose, D-trehalose, D-cellobiose,
gentiobiose, sucrose, D-turanose, stachyose, D-raffinose, α-D-lactose, D-melibiose, β-
methylD-glucoside, D-salicin, N-acetyl-D-glucosamine, α-D-glucose, D-mannose, D-
fructose, D-galactose, inosine, sodium lactate, D-mannitol, D-arabitol, *myo*-inositol,
glycerol, L-aspartic acid, L-glutamic acid, pectin, D-arabitol, D-gluconic acid, D-
glucuronic acid and glucuronamide.

The type strain is resistant to nalidixic acid and aztreonam.

The type strain, VK105^T, was isolated from seeds of wild cotton collected in Puerto
Rico, USA.

DESCRIPTION OF CURTOBACTERIUM ORYZAE SP. NOV.

Curtobacterium oryzae (o.ry′zae. N.L. gen. n. *oryzae* of rice, referring to the origin of the
type strain)

Cells are Gram-stain positive, non-motile, non-spore-forming and rod shaped or slightly curved; V shapes are formed. Aerobic and chemoorganotrophic. Colonies are light yellow, entire, round and raised on nutrient agar. Good growth is obtained at 22, 30 and 37°C on the media tested. Grows on media supplemented with 4% NaCl and at pH 5-7. Positive for catalase, but negative for oxidase or urease. The following substrates are used as carbon sources: N-Acetyl-D-Glucosamine, N-Acetyl-D-Galactosamine, N-Acetyl Neuraminic Acid, D-Arabitol, D-Cellobiose, L-Fucose, Gentiobiose, myo-Inositol, α-D-Lactose, D-Mannitol, D-Melibiose, β-Methyl-D-Glucoside, D-Raffinose, D-Sorbitol, Turanose, D-Gluconic Acid, D-Glucuronic Acid, L-Alanine, L-Glutamic Acid, L-Rhamnose, D-Glucose-6-PO4 and Acetic Acid.

The type strain is resistant to nalidixic acid and aztreonam.

The type strain, SS108^T, was isolated from *Oryza sativa* seedlings collected in Texas, USA.

ACKNOWLEDGEMENTS

We would like to thank Professor Aharon Oren, The Hebrew University of Jerusalem, for checking Latin species names. Support was provided by USDA-NIFA Multistate Project W4147 and the New Jersey Agricultural Experiment Station.

LITERATURE CITED

Aizawa, T., Ve, N. B., Kimoto, K., Iwabuchi, N., Sumida, H., Hasegawa, I., et al. 2007. *Curtobacterium ammoniigenes* sp. nov., an ammonia-producing bacterium isolated from plants inhabiting acidic swamps in actual acid sulfate soil areas of Vietnam. Int. J. Syst. Evol. Microbiol. 57:1447–1452.

de Almeida Lopes, K. B., Carpentieri-Pipolo, V., Fira, D., Balatti, P. A., López, S. M. Y., Oro, T. H., et al. 2018. Screening of bacterial endophytes as potential biocontrol agents against soybean diseases. J. Appl. Microbiol. 125:1466–1481.

Behrendt, U., Ulrich, A., Schumann, P., Naumann, D., and Suzuki, K. 2002. Diversity of grass-associated Microbacteriaceae isolated from the phyllosphere and litter

layer after mulching the sward; polyphasic characterization of *Subtercola pratensis* sp.

nov., *Curtobacterium herbarum* sp. nov. and *Plantibacter flavus* gen. nov., sp. nov. Int. J.

Syst. Evol. Microbiol. 52:1441–1454.

Bertani, I., Abbruscato, P., Piffanelli, P., Subramoni, S., and Venturi, V. 2016. Rice

bacterial endophytes: isolation of a collection, identification of beneficial strains and

microbiome analysis. Environ. Microbiol. Rep. 8:388–398.

Bouizgarne, B. 2013. Bacteria for plant growth promotion and disease

management. In *Bacteria in Agrobiology: Disease Management*, Springer, p. 15–47.

Chun, J., Oren, A., Ventosa, A., Christensen, H., Arahal, D. R., da Costa, M. S., et

al. 2018. Proposed minimal standards for the use of genome data for the taxonomy of

prokaryotes. Int. J. Syst. Evol. Microbiol. 68:461–466.

Collins, M., and Jones, D. 1983. Reclassification of *Corynebacterium*

flaccumfaciens, *Corynebacterium betae*, *Corynebacterium oortii* and *Corynebacterium*

poinsettiae in the genus *Curtobacterium*, as *Curtobacterium flaccumfaciens* comb. nov.

Microbiology. 129:3545–3548.

Dey, R., Pal, K. K., and Tilak, K. 2014. Plant growth promoting rhizobacteria in

crop protection and challenges. In *Future Challenges in Crop Protection Against Fungal*

Pathogens, Springer, p. 31–58.

Dunleavy, J. 1989. *Curtobacterium plantarum* sp. nov. is ubiquitous in plant leaves

and is seed transmitted in soybean and corn. Int. J. Syst. Evol. Microbiol. 39:240–249.

Edwards, J., Johnson, C., Santos-Medellín, C., Lurie, E., Podishetty, N. K.,

Bhatnagar, S., et al. 2015. Structure, variation, and assembly of the root-associated

microbiomes of rice. Proc. Natl. Acad. Sci. 112:E911–E920.

Finkel, O. M., Castrillo, G., Paredes, S. H., González, I. S., and Dangl, J. L. 2017.

Understanding and exploiting plant beneficial microbes. Curr. Opin. Plant Biol. 38:155–

163.

Glick, B. R. 2012. Plant growth-promoting bacteria: mechanisms and applications.

Scientifica. 2012.

Irizarry, I., and White, J. 2017. Application of bacteria from non-cultivated plants

to promote growth, alter root architecture and alleviate salt stress of cotton. J. Appl.

Microbiol. 122:1110–1120.

Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C.,

et al. 2004. Versatile and open software for comparing large genomes. Genome Biol.

5:R12.

Langmead, B., and Salzberg, S. L. 2012. Fast gapped-read alignment with bowtie

2. Nat Methods. 9 Available at: https://doi.org/10.1038/nmeth.1923.

Li, D., Liu, C.-M., Luo, R., Sadakane, K., and Lam, T.-W. 2015. MEGAHIT: an

ultra-fast single-node solution for large and complex metagenomics assembly via succinct

de Bruijn graph. Bioinformatics. 31:1674–1676.

Liu, F., Hewezi, T., Lebeis, S. L., Pantalone, V., Grewal, P. S., and Staton, M. E.

2019. Soil indigenous microbiome and plant genotypes cooperatively modify soybean

rhizosphere microbiome assembly. BMC Microbiol. 19:1–19.

Longley, R., Noel, Z. A., Benucci, G. M. N., Chilvers, M., Trail, F., and Bonito, G.

2020. Crop management impacts the soybean (*Glycine max*) microbiome. Front.

Microbiol. 11:1116.

Parte, A. C. 2018. LPSN-List of prokaryotic names with standing in nomenclature

(bacterio.net), 20 years on. Int. J. Syst. Evol. Microbiol. 68:1825–1829.

311  Price, M. N., Dehal, P. S., and Arkin, A. P. 2010. FastTree 2 - approximately

312 maximum-likelihood trees for large alignments. PLoS ONE. 5:e9490.

313  Qiao, Q., Wang, F., Zhang, Jingxia, Chen, Y., Zhang, C., Liu, G., et al. 2017. The

314 variation in the rhizosphere microbiome of cotton with soil type, genotype and

315 developmental stage. Sci. Rep. 7:1–10.

316  Richter, M., and Rosselló-Móra, R. 2009. Shifting the genomic gold standard for

317 the prokaryotic species definition. Proc. Natl. Acad. Sci. 106:19126–19131.

318  Roman-Reyna, V., Pinili, D., Borja, F. N., Quibod, I. L., Groen, S. C., Alexandrov,

319 N., et al. 2020. Characterization of the leaf microbiome from whole-genome sequencing

320 data of the 3000 rice genomes project. Rice. 13:1–8.

321  Ryan, R. P., Germaine, K., Franks, A., Ryan, D. J., and Dowling, D. N. 2008.

322 Bacterial endophytes: recent developments and applications. FEMS Microbiol. Lett.

323 278:1–9.

324  Sahlin, K., Vezzi, F., Nystedt, B., Lundeberg, J., and Arvestad, L. 2014. BESST-

325 efficient scaffolding of large fragmented assemblies. BMC Bioinformatics. 15:281.

326  Schenk, P. M., Carvalhais, L. C., and Kazan, K. 2012. Unraveling plant-microbe

327 interactions: can multi-species transcriptomics help? Trends Biotechnol. 30:177–184.

328  Seemann, T. 2019. *barrnap 0.9: rapid ribosomal RNA prediction*. Available at:

329 https://github.com/tseemann/barrnap.

330  Ullah, A., Akbar, A., Luo, Q., Khan, A. H., Manghwar, H., Shaban, M., et al. 2019.

331 Microbiome diversity in cotton rhizosphere under normal and drought conditions. Microb.

332 Ecol. 77:429–439.

333  Vimal, S. R., Patel, V. K., and Singh, J. S. 2019. Plant growth promoting

334 *Curtobacterium albidum* strain SRV4: an agriculturally important microbe to alleviate

335 salinity stress in paddy plants. Ecol. Indic. 105:553–562.

336   Zhou, W., Wheeler, T. A., Starr, J. L., Valencia, C. U., and Sword, G. A. 2018. A

337   fungal endophyte defensive symbiosis affects plant-nematode interactions in cotton. Plant

338   Soil. 422:251–266.

TABLES

Table 1. Physiological characteristics of newly described *Curtobacterium glycinis*, *C. gossypii* and *C. oryzae* species and related *Curtobacterium* type strains. All strains were positive for the assimilation of dextrin, D-fructose, D-galactose, α-D-glucose, D-mannose and sucrose (data not shown).

	C. gossypii VK105	*C. oryzae* OG107	*C. glycinis* SS108	*C. citreum* JCM 1345[T]	*C. pusillum* JCM 1350[T]	*C. luteum* JCM 1480[T]	*C. albidum* NBRC 15078[T]	*C. herbarum* DSM 14013[T]
General characteristics:								
Colony pigment		light yellow	ivory	yellow	yellow	yellow	ivory	orange
Motility	-	w	w	+	+	+	-	+
Oxidase	-	-	-	-	-	-	-	-
Catalase	+	+	+	+	+	+	+	+
Urease	-	-	-	-	-	-	-	nr
Growth with 4% NaCl	+	+	+	nr	nr	nr	nr	nr
Growth with 8% NaCl	+	-	-	nr	nr	nr	nr	nr
Utilization of:								
N-Acetyl-D-Glucosamine	+	+	+	+	-	+	w	nr
N-Acetyl-D-Galactosamine	-	+	+	-	-	-	-	nr
N-Acetyl Neuraminic Acid	-	+	-	nr	nr	nr	nr	nr
D-Arabitol	+	+	+	-	w	-	-	nr
D-Cellobiose	+	+	+	+	-	+	+	nr
L-Fucose	w	+	+	+	-	+	-	nr
Gentiobiose	+	+	+	-	+	-	+	nr
myo-Inositol	+	+	+	-	-	-	-	nr
α-D-Lactose	+	+	+	+	-	+	+	nr
D-Mannitol	+	+	+	-	+	-	w	+
D-Melibiose	+	+	+	+	+	+	w	+
β-Methyl-D-Glucoside	+	+	+	-	+	-	w	nr

D-Raffinose	+	+	+	-	-	-	+	nr
D-Sorbitol	w	+	-	-	+	-	+	+
Turanose	+	+	+	-	+	-	+	nr
D-Gluconic Acid	+	+	+	-	+	-	+	nr
D-Glucuronic Acid	+	+	+	-	-	-	-	nr
L-Alanine	+	+	+	-	-	-	w	nr
L-Glutamic Acid	+	+	w	+	-	-	-	nr
L-Rhamnose	w	+	+	-	-	-	-	+
D-Glucose-6-PO4	-	+	+	-	-	-	-	nr
Acetic Acid	w	+	-	-	-	-	-	nr

+, Positive; w, Weakly positive; -, Negative; nr, Not reported.

Data from this study, Aizawa et al. (2007) and Behrendt et al. (2002).

FIGURES

├──┤ 0.001

Figure 1. Phylogenetic 16S rRNA gene tree of *Curtobacterium glycinis* sp. nov.

strain OG107, *Curtobacterium gossypii* sp. nov. strain VK105, *Curtobacterium oryzae* sp.

nov. strain SS108 and relatives generated using FastTree (Price et al. 2010). All validly

published *Curtobacterium* species were included in the tree, the tree is rooted with

Frondihabitans australicus. New species are in bold. Strain identifiers and GenBank

accession numbers follow species names, T stands for 'type'. Support values above 70%

are given by the branches. *Curtobacterium glycinis* is most closely related to *C. herbarum*

with 84% support, *Curtobacterium gossypii* to *C. albidum* and *C. citreum* with 82%

support and *Curtobacterium oryzae* to *C. luteum* with 79% support. Branch lengths are

proportional to the changes along the branches, a scale bar is provided.

Figure 2. Morphology of *Curtobacterium glycinis* sp. nov. strain OG107 depicted following Gram stain using bright field microscopy. Bar = 10 µm.

389 **Figure 3.** Morphology of *Curtobacterium gossypii* sp. nov. strain VK105 depicted

390 following Gram stain using bright field microscopy. Bar = 10 μm.

392
393
394   **Figure 4.** Morphology of *Curtobacterium oryzae* sp. nov. strain SS108 depicted following
395   Gram stain using bright field microscopy. Bar = 10 μm.