Predicting tomato field-yield using continuous monitoring of young tomato water

status

Sanbon Chaka Gosa[1*], Amit Koch[1*], Itamar Shenhar[1], Joseph Hirschberg[2], Dani Zamir[1,]

Menachem Moshelion[1]

[1] The R.H. Smith Institute of Plant Sciences and Genetics in Agriculture, The R.H. Smith Faculty

of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot, 76100

Israel

[2] Department of Genetics, Alexander Silberman Institute of Life Sciences, The Hebrew

University of Jerusalem, Jerusalem 9190401, Israel

* Equal contribution

Abstract

To address the challenge of predicting tomato yields in the field, we used whole-plant

functional phenotyping to evaluate water relations under well-irrigated and drought

conditions. The genotypes tested are known to exhibit variability in their yields in wet and

dry fields. The examined lines included two lines with recessive mutations that affect

carotenoid biosynthesis, zeta z^{2083} and tangerine t^{3406}, both isogenic to the processing

tomato variety M82. The two mutant lines were reciprocally grafted onto M82 and multiple

physiological characteristics were measured continuously, as well as before, during and

after drought treatment in the greenhouse. A comparative analysis of greenhouse and field

yields showed that the whole-canopy stomatal conductance (g_{sc}) in the morning and

cumulative transpiration (CT) were strongly correlated with field measurements of total

yield (TY: $r^2 = 0.9$ and 0.77, respectively) and plant vegetative weight (PW: $r^2 = 0.6$ and

0.94, respectively). Furthermore, the minimum CT during drought and the rate of recovery

when irrigation was resumed were both found to predict resilience.

Keywords: drought tolerance, functional genomic mapping, functional phenotyping,

physiological trait, time-series measurements, tomato, yield prediction, yield-prediction models

Introduction

Water stress is the main factor limiting crop yields worldwide (Tilkat *et al.*, 2012; Halperin

et al., 2017; Polania *et al.*, 2017). Despite intense research over the last decades, drought

tolerance is still a poorly understood complex trait (Anderegg *et al.*, 2013). Accurate yield

prediction is important for national food security and global food production (Horie *et al.*,

1992) and it also aids policymaking. From the research and development perspective, yield

prediction tools would enable breeders to reduce the time and cost required to select the best

parent lines and test new hybrids under different environmental conditions (Han et al., 2020;

Khaki & Wang, 2019; Weber et al., 2012). Finally, reliable yield prediction would benefit

the growers who are the end-users of newly developed, improved varieties, aiding their crop

management and helping them to make wise economic decisions (Horie *et al.*, 1992).

However, early growth-stage prediction of crop yields is a challenging task, in general, and

is even more challenging under water stress. Several yield-prediction models have been

developed, some of which consider yield as a function of genotype (G) and environment (E),

and treat the interaction between the two (G×E) as a noise (DeLacy *et al.*, 1996; Heslot *et*

al., 2014). Some other models address G×E interactions using multiplicative models (Crossa

and Cornelius, 1997), factor analytic (FA) models and linear mixed models to cluster

environments and genotypes and detect their interactions (Burgueño *et al.*, 2008, 2011;

Montesinos-López *et al.*, 2018). A recently developed yield-prediction model, which is based

on a deep neural network fed with weather and soil-condition data for 2,247 sites and yield

data for 2,267 maize hybrids) was found to accurately predict yields. The developers of that system concluded that environmental factors had a stronger effect on the crop yield than genotype did (Khaki & Wang, 2019). Thus, early-season yield prediction may require a large amount of data from the soil-plant-atmosphere continuum (SPAC). Plant physiological traits that are most relevant to productivity and are very responsive to environmental conditions are expected to serve as important yield predictors (Gosa *et al.*, 2019).

Recent advances in crop physiology show that under drought conditions, quantitative physiological traits such as stomatal conductance (Bahar *et al.*, 2009), osmotic adjustment, accumulation and remobilization of stem reserves and photosynthetic efficiency are strongly correlated with yield (Richards *et al.*, 2002; Aisawi *et al.*, 2015; Teodoro *et al.*, 2019). Nevertheless, most of the available models do not include key plant physiological traits, such as g_{sc} and photosynthesis, which contribute to crop productivity (Roche, 2015; Faralli and Lawson, 2020). These traits are among the primary and most sensitive responses of the plant to the changing environment (Pirasteh-Anosheh *et al.*, 2016) and this dynamic behavior helps to optimize the plant's response to changing environmental conditions and probably also helps to maximize yield.

Therefore, we hypothesized that having a set of high-resolution and continuous data for many key-physiological traits, measured under different environmental conditions at an early growth stage, could improve our ability to predict the yields of particular genotypes under field conditions. To profile physiological traits that reliably contribute to the yield-prediction model, we used different genotypes known for carotenoid biosynthesis mutations and yield reduction and compared them with a control genotype (see Materials and Methods). Preliminary field trials that we conducted showed that these mutant lines exhibit pleiotropic mutations of the zeta and tangerine genes.

Materials and Methods

Plant material and the grafting procedure

Tomato cv. M82 seeds (Eshed and Zamir, 1995), the recessive mutant *zeta z^{2083}* (ZET) described by Koch et al. (2021) and the *tangerine t^{3406}* (TAN) mutant described in (Isaacson *et al.*, 2002; Kachanovsky *et al.*, 2012) were used. These mutant lines have mutations that affect two of the four enzymes reported to convert phytoene into phytoene desaturase (PDS), ζ-carotene desaturase (ZDS), zeta isomerase (ZISO) and carotene isomerase (CRTISO; (Hirschberg, 2001); (Enfissi *et al.*, 2017). Fruits of the *tangerine* mutant are orange and accumulate prolycopene (7,9,7′,9′- tetra-cis-lycopene) instead of the all-translycopene that is normally synthesized in wild-type fruit, as a result of a mutation in the carotenoid cis-trans isomerase (Kachanovsky *et al.*, 2012). In terms of phenotype, the *tangerine* mutant has yellowish young leaves, light green foliage and pale flowers. The *zeta* leaf phenotype resembles that of *tangerine*, but the fruit of *zeta* mutants are yellow-orange in color, due to a reduction in lycopene that is accompanied by the accumulation of ζ-carotene, phytofluene and phytoene. To increase the phenotypic variation in yield, we made seven grafting combinations, namely, four hetero graftings in which M82 was reciprocally grafted with ZET and TAN and three self-grafts for each genotype.

Open-field experiments

The results presented here are from work that was done in two consecutive growing seasons, 2018 and 2019, at the Western Galilee Experimental Station in Akko, Israel. In those trials, we used a low planting density of one plant per m^2. In 2018, the experiment involved individual plants in a completely randomized design in blocks, with a minimum of 15 replicates per block. In 2019, the experiment was conducted in plots of 10 plants per 5 m^2, arranged in a randomized block design. The seedlings were grown in greenhouse for 35 days

and then transplanted into the field at the beginning of April; wet and dry trials were

conducted. Both wet and dry fields started the growing season at field capacity, which

represents the maximum amount of water that the soil could hold. In the wet treatment, 320

m^3 of water were applied per 1000 m^2 of field throughout the growing season, according to

the irrigation protocols commonly used in the area. In the limited-irrigation (drought)

treatment, we stopped irrigation 3 weeks after planting, so only 30 m^3 of water were applied

per 1000 m^2 of field. There was no rain during the experimental period, so the drought

scenario was managed entirely via irrigation.

Measurements of yield and yield components

The experiments were harvested when nearly 100% ripened. Plant vegetative weight (PW, g m^{-2}) was

determined by weighing only the vegetative tissue (after harvesting the fruits) without the roots. Total

fruit yield (TY, g m^{-2}) per plant or plot included both the red and a few green fruits. Mean fruit weight

(FW) was estimated from a random sample of 20 fruits per plant or plot. The concentration of total

soluble solids (Brix %) was measured using a digital refractometer and a random sample of 10 fruits

per plant or 20 fruits per plot. The sugar output per plant or plot was calculated as the product of Brix

and TY.

Pigment extraction and analysis

118　Fresh samples of root and flower tissues (approximately 50 to 100 mg) were harvested and

119　immediately frozen in liquid nitrogen. Carotenoids were extracted and quantified according

120　to protocols described by (Karniel *et al.*, 2020).

Greenhouse experiment using the physiological-phenotyping platform

122　A greenhouse experiment was conducted in parallel with a field experiment from mid-April

123　to mid-May in 2018. The grafted and well-established seedlings were transplanted into 4-L

124　pots filled with potting soil (Bental 11, Tuff Marom Golan, Israel). Plants were grown under

125　semi-controlled greenhouse conditions with naturally fluctuating light (see Fig. 1A). Whole-

plant, continuous physiological measurements were taken using a high-throughput, telemetric, gravimetric-based phenotyping system (Plantarry 3.0 system; Plant-DiTech, Israel) in the greenhouse of the I-CORE Center for Functional Phenotyping (http://departments.agri.huji.ac.il/plantscience/icore.phpon), as described in (Dalal *et al.*, 2020).

The set-up included 72 highly sensitive, temperature-compensated load cells, which were used as weighing lysimeters. Each unit was connected to its own controller, which collected data and controlled the irrigation to each plant separately. A pot containing a single plant was placed on each load cell. (For more details, see the "Experimental set-up" section.) The data were analyzed using SPAC-analytics (Plant-Ditech), a web-based software program that allowed us to view and analyze the real-time data collected from the Plantarray system.

Experimental set-up

The experimental set-up was generally similar to that described by (Dalal *et al.*, 2020), with some modifications. Briefly, before the start of the experiment, all load-cell units were calibrated for accuracy and drift level under constant load weights (1 kg and 5 kg). Each pot was placed into a Plantarray plastic drainage container on a lysimeter. The containers fit the pot size, to enable the accurate return to pot capacity after irrigation and to prevent evaporation. The container had orifices on its side walls that were located at different heights, to allow for different water levels after the drainage of excess water following irrigation. Evaporation from the soil surface was prevented by a cover with a circle cut out at its center through which the plant grew.

Each pot was irrigated with a multi-outlet dripper assembly that was pushed into the soil to ensure that the medium in the pot was uniformly wetted at the end of the free-drainage period following each irrigation event. Irrigation events were programmed to take place

during the night in four consecutive pulses. The amount of water left in the drainage

containers underneath the pots at the end of the irrigation events was intended to provide

water to the well-irrigated plants beyond the water volume at pot capacity. The associated

monotonic weight loss over the course of the daytime hours was essential for the calculation

of the different physiological traits using the data-analysis algorithms (see inset in Fig. 1B).

Measurement of quantitative physiological traits

The following water-relations kinetics and quantitative physiological traits of the plants were

determined simultaneously, following protocols and equations (Halperin *et al.*, 2017)

implemented in the SPAC-analytics software for daily transpiration, transpiration rate,

normalized transpiration (E) and WUE. Cumulative transpiration (CT) was calculated as the

sum of daily transpiration for all 29 days of the experiment for each plant. The other

physiological traits used in this experiment are described in (Halperin *et al.*, 2017). The

estimated plant weight at the beginning of the experiment was calculated as the difference

between the total system weight and the sum of the tare weight of the pot + the drainage

container, the weight of the soil at pot capacity and the weight of the water in the drainage

container at the end of the free drainage. The plant weight at the end of a growth period

(calculated plant weight) was calculated as the sum of the initial plant weight and the product

of the multiplication of the cumulative transpiration during the period by the WUE. The

latter, determined as the ratio between the daily weight gain and the transpiration during that

day, was calculated automatically each day by the SPAC-analytics software. The plant's

recovery from drought was described by the rate at which the plant gained weight following

the resumption of irrigation (recovery stage).

Data presentation and statistical analysis

We used the JMP® ver. 14 statistical packages (SAS Institute, Cary, NC, USA) for our

statistical analyses. Levene's test was used to examine the homogeneity of variance among

the treatments. Differences between the genotypes were examined using Tukey HSD. Each

analysis involved a set significance level of $P \leq 0.05$ or 20 fruits per plot.

Pair-wise Pearson correlations between traits under greenhouse conditions and the yield and

yield components measured in the open field (i.e., plant vegetative weight, red yield, green

yield, Brix yield and total yield) were calculated using the genotype's mean performance.

Results

Field-based plant weight and total yield

The yield components plant vegetative weight (PW), total yield (TY), and green yield (GY)

were tested under well-irrigated and dry conditions in the 2018 and 2019 growing seasons.

Comparing two key traits TY and PW we found similar performances of the genotypes across

years in 2018 and 2019.

Plant vegetative weight (PW): In the well-irrigated field, the M82 self-grafted plants

(M82_scion/M82_rootstock) had a significantly higher PW than the TAN/TAN and

ZET/ZET plants. Under the dry condition, no significant difference was observed between

the M82 and TAN self-grafted plants; whereas the plant vegetative weights of the ZET/ZET

plants were significantly lower (Fig. 2A, B) under both well-irrigated and dry conditions.

Total Yield (TY): Under well-irrigated conditions, the TY of the different self-grafted M82

was significantly different from both mutants across both years. The total yield of M82/M82

was significantly higher than the yields of the other self-grafted plants, TAN/TAN was a

medium yielder and ZET/ZET had the lowest yield of all of the self-grafted plants across

both years. Under the drought condition, the total yield of M82/M82 remained higher than

those of the other two genotypes, which were not different from each other (Fig. 2C and D,

respectively). However, the TY under the drought condition was less than half of that

observed under the well-irrigated condition. To increase the phenotypic variation in yield,

we used a reciprocal-grafting approach, in which seven combinations of the three tomato

cultivars resulted different gradient of yield performance under wet and dry conditions (Fig.

2E and F, respectively). TY increased more than 2-fold when TAN and ZET scions were

grafted onto M82 rootstock, especially under dry conditions.

Early-stage physiological traits measured in the greenhouse.

To identify physiological traits of young tomato plants that might serve as good predictors

of yield in the field, we profiled multiple physiological traits using continuous data collected

on a minute time-scale, such as whole-canopy stomatal conductance (g_{sc}); continuous data

collected on a daily time-scale, such as transpiration throughout the experimental period as a

cumulative transpiration (CT); and single-point measurements such as growth rate and plant

net weight (see Table 1 and Supplementary Datasets 1 and 2).

The continuous measurement data show that the traits varied with the environment.

For example, as shown in Figure 3, the whole-canopy conductance measured every 3 min

over the whole day fluctuated over the course of the day in response to the environment. To

better understand this trait, we divided the day into three periods: morning, mid-day and late

afternoon. We found that stomatal conductance was relatively high at morning time (Fig. 3,

marked in green), declined between mid-day and late afternoon to some point, and then

increased again during the late afternoon. We also performed a correlation analysis using the

averaged 3-min measurements, in which each time regime was a single point, and correlated

that measurement data with field-based yield and biomass data.

Correlation of greenhouse physiological traits with yield and yield components in the field

Data from the functional-phenotyping system were composed of continuous soil-plant-atmosphere measurements, with each data point representing the trait at a certain time point. In contrast, field data are normally composed of a single-point measurement that represents the plant's absolute performance throughout the season (e.g., total fruit yield or plant vegetative weight). When we compared time-series, cumulative and single-point physiological traits (measured traits) of young tomato plants with their field-based yield-related traits (TY, PW, RF, GF and Brix,), we found only a few traits that were highly correlated with each other (Table 1), out of about 95 bivariate combinations (see Fig. 4, Supplementary Figs. S3 and S4, and Supplementary Dataset S2). Here, we present a few physiological traits for which the greenhouse data was strongly correlated with the field data and for which we observed low *p*-values (e.g., the highly correlated traits in Table 1).

Time-series data are highly dynamic because of the plant's continuous response to environmental changes (e.g., stomatal conductance, Fig.3; transpiration rate, Supplementary Dataset S1). Therefore, some data points were strongly correlated with yield (e.g., g_{sc} in the morning, Table 1) while others were weakly correlated with yield (e.g., g_{sc} at mid-day; Table 1). Looking at cumulative physiological data or single-point traits, both presented as a single value (e.g., CT, growth rate, plant net weight), eliminated the need to select a specific time point and revealed highly significant and positive correlations between CT and yield and most of the yield components under well-irrigated conditions (Fig. 4A–D). Similarly, the CT of drought-treated plants after recovery in the greenhouse was positively correlated with yield and with most yield components, but poorly correlated with green yield (Fig. 4C). A similar positive correlation between CT and yield in the field was observed in 2019 (Supplementary Figs. S5 and S6).

Cumulative transpiration as an indicator of resilience and yield performance

The rate of plants' recovery from drought stress (i.e., drought resilience) is an important trait. To evaluate this resilience, we measured the CT for the first week after recovery from drought. We then compared that CT data with CT data from two other periods during the experiment: the pre-drought period and the drought period (Fig. 5A). While the CT over the pre-drought treatment showed a similar positive correlation with that of the entire well-irrigated experiment (Fig. 5B), we found a strong negative correlation between total yield and CT and under drought conditions (Fig. 5C). We also observed a strong positive correlation between CT and TY during the recovery period (Fig. 5D), even though the actual total yield of the drought-treated plants was half that of the plants grown under the well-irrigated condition.

Discussion

Physiological traits (e.g. photosynthesis or stomatal conductance) are key contributors to plant productivity and yield (Roche, 2015; Faralli and Lawson, 2020). However, existing methods of measuring these traits are mostly manual and thus are limited to a single point on a single leaf at a time (Iseki and Olaleye, 2020). As these physiological traits are very sensitive to ambient conditions (especially light and VPD; (Gosa *et al.*, 2019), conventional manual measurements fail to capture the temporal and spatial dynamic interactions between the genotype and the environment. This could be misleading for yield prediction, as plants respond differently to dynamic growing conditions (Gosa *et al.*, 2019). Hence, the integration of manual physiological measurements into breeding programs is limited, most likely due to their low-throughput nature and the large degree of variation within and between temporal and spatial measurements.

In this study, we used continuous measurements of physiological traits to assess

whether those traits could serve as early predictors of plant responses to environmental

conditions. We used a high-throughput, physiological phenotyping platform, with a high

resolution of 3-min intervals, to capture plant responses to the environment. In our

experiments, we captured a detailed profile of each plant's performance. However,

correlating this detailed profile with actual field performance was another challenge (e.g., the

choice of the measurement points to be used in the analysis). A good example of this

challenge is demonstrated in Figure 3, which shows how continuous g_{sc} measurements were

more or less correlated with yield performance at different hours of the day (Table 1). In line

with our finding, a recent research reported by (Du *et al.*, 2020) high stomatal conductance

and photosynthesis was seen in rice tiller in the early morning, which could be related to

relatively low VPD and sufficient light for photosynthesis (Gosa *et al.*, 2019), and it was

suggested that early morning might be a "golden hour" for plants (Gosa *et al.*, 2019). Here,

we demonstrate that this morning g_{sc} peak is strongly correlated with TY and PW in the field.

Moreover, the fact that our midday g_{sc} data was less strongly correlated with field

performance is in agreement with the common practice of measuring g_{sc} between 10:00 and

14:00 (Gooch, 2011; Roberts *et al.*, 2017). The weak correlation between midday g_{sc} and

yield could be related to the dynamic patterns of daily whole-plant water-use efficiency

suggested by (Gosa *et al.*, 2019). Nevertheless, the identification of the best time to measure

each trait and/or weather condition would require the use of new data-analysis tools, such as

AI-based algorithms, to create a more comprehensive understanding of our large amount of

data. Moreover, a better understanding the genetic mechanism governing the morning peak

could contribute to the improvement of crop productivity through breeding, in addition to

yield prediction, as plants use water very efficiently at that time of day. It is also important

to examine many genotypes. For example, the tomato introgression line (IL) population

(Eshed and Zamir, 1995) for which we have multiple years of field data, to verify whether

these morning peaks are present in all genotypes. This would improve our understanding of

the genetic mechanism for this important trait.

The relationship between transpiration and net carbon assimilation or dry weight has

been well studied (de Wit, 1958; Tolk and Howell, 2009). The reason for this correlation is

most likely due to the fact that CO_2 enters via the same open stomata through which the plant

transpires. Indeed, we found a positive correlation between CT and yield. Yet, this correlation

was weaker than the correlation between morning g_{sc} and TY ($r^2 = 0.9$ versus $r^2 = 0.77$ and

$p = 0.004$ versus $p = 0.04$, respectively), suggesting that the correlation between CT and CO_2

absorption might be affected by other environmental factors, such as VPD, radiation and

humidity, which are all known to affect stomatal conductance (Zhang *et al.*, 2018). On the

other hand, CT is a stable, single-point measurement that is relatively simple to measure, yet

integrates the overall responses of plants to the environment throughout the experimental

period. Nevertheless, these correlations should be examined in other plant species, as

different vegetative stages, reproductive systems, growth, and development patterns may

involve different yield-related predictive traits.

Another goal of this study was to evaluate stress-related traits that could predict yield.

Under water-deficient conditions, the plant undergoes several changes ranging from

molecular and cellular changes to changes at the whole-plant level. This reprogramming of

metabolic pathways and physiological response patterns enables the plant to better cope with

drought stress (Jiang and Huang, 2001; Claeys and Inzé, 2013). Many of the physiological

responses to stress [e.g., reduced stomatal conductance, damage to the photosynthetic parts,

reduced chlorophyll content; (Fu and Huang, 2001)] have dramatic effects on plant

productivity. Under stressful conditions, plants enter a protective or survival mode (Claeys

and Inzé, 2013) at the expense of their productive mode (Moshelion, 2020). Here, we found

that CT was strongly and positively correlated with TY under well-irrigated conditions, but negatively correlated with yield under stressful conditions (Fig. 5C). This reversal reflects the productive-survival transition mode of the plant (Moshelion, 2020). Namely, a plant that can maximize its transpiration under well-irrigated conditions and swiftly minimize it under stressful conditions is more likely to produce more yield by the end of the season, if it recovers quickly after the stress ends. This is clearly shown in Figure 5B: M82/M82 and TAN/M82 had higher levels of transpiration pre-stress, but swiftly reduced their transpiration during the stress period (Fig. 5C) and went back to their high levels of transpiration after recovery (Fig. 5D), which might have led them to have higher yields than the other lines. Thus, this transition mode is important for distinguishing plants' stress-response (protective) mode from their normal growth response (productive mode). An additional important phase of the plant-stress response is the plant's post-stress performance, often called resilience.

Resilience to water limitations, specifically the plant's ability to resume growth and gain yield after water resumption following drought stress, was acknowledged by (Fang and Xiong, 2015). Resilience is considered to be a key trait for crop improvement for water stress (Dalal *et al.*, 2019). Although it has not received much attention for some time (Chen *et al.*, 2016), this trait has been prioritized in some breeding programs (Chapuis *et al.*, 2012). In this study, we found that the CT of all of the treatment periods together (pre-treatment, drought and the recovery period) and the CT of only the recovery period each had a strong, positive relationship with TY (Figs. 4B, 5D), suggesting the importance of this trait for stress-response profiling.

Although the current study did not cover stress-response mechanisms in detail, it is interesting to note that ZET and TAN are prone to slow recovery rates (Fig. 5A, D), in terms of cumulative and daily transpiration, which probably led to their low yields. Their lower CT levels may be related to their root systems, since both ZET and TAN grafted as scions on

M82 performed a lot better than ZET and TAN when M82 was used as the scion. This

behavior could be accounted for by the higher carotenoid concentrations in ZET roots, as

compared to M82 roots. The roots of M82 had undetectable levels of carotenoids; whereas

the roots of *tangerine* and *zeta* contained the same composition of carotenoids typically found

in their fruits and flowers. The low concentration of carotenoids in wild-type roots can be

explained, in part, by the synthesis of ABA and strigolactones in the tomato roots, as

reviewed in (Saeed *et al.*, 2017). In these mutants, carotenoid synthesis is blocked, so

intermediate products accumulate. This blockage is very effective in the roots due to their

lack of exposure to light, whereas exposure to light in the leaves partially compensates for

the lack of carotenoid isomerase CRTISO and ZISO(Isaacson *et al.*, 2002; Fantini *et al.*,

2013). However, this cannot explain the lower yields of ZET and TAN under the well-

irrigated condition. The relatively low yield of TAN plants under the well-irrigated condition

might be linked to the lower concentrations of carotenoids, such as violaxanthin and

neoxanthin, in their flowers, as compared to M82 (Supplemental Fig. S2). Violaxanthin and

neoxanthin are the precursors for ABA synthesis (Galpaz *et al.*, 2008), which suggests that

ABA might have been involved in reducing the yields of these mutants. However, this

hypothesis needs to be tested in future research.

In conclusion, continuous measurements of dynamic traits such as g_{sc} provide a

dataset that is rich, yet also very challenging to analyze. In our current study, we confirmed

that early-morning g_{sc} is an important physiological trait that can predict yield performance.

Understanding the genetic mechanism underlying early-morning g_{sc} could be a potential

avenue for breeding programs aimed at developing lines that will perform well under water-

deficit conditions. Furthermore, future data-science tools are likely to improve our

understanding of the mechanisms involved and allow us to use these dynamic traits in yield-

prediction models. On the other hand, the relatively simple trait of CT of young tomato was

proven to be a good predictor of plant biomass and yield performance. The inclusion of CT

in yield models is expected to improve the accuracy and consistency of those models, which

should facilitate the selection of complex traits for water-stress conditions.

It is important to note that various crops may present different response profiles, as

well as different levels of susceptibility to a particular type of stress, depending on their

biochemical, physiological and phonological stage. For example, for more than five decades,

we have known that maize crops are most sensitive to drought stress that occurs at flowering

time (Claassen and Shaw, 1970; Deikman *et al.*, 2012). Therefore, it is important to conduct

screening and follow-up testing using stress applied around different phenological and

developmental stages (e.g., flowering time) of each crop of interest.

In addition to yield prediction and crop improvement (i.e., at pre-breeding stages),

high-resolution, continuous physiological data could further be exploited to help bridge the

genotype-phenotype gap, by combining the functional-genomics approach with a high-

resolution time axis on a QTL map. This combined approach may help to identify time-

dependent QTLs for dynamic physiological traits such as g_{sc} and help us to understand the

genetic mechanisms that underlie those dynamic traits.

Table and Figure Legends

Table 1. Correlations between the physiological traits of young tomato plants in the greenhouse and their field-based yield and biomass under well-irrigated conditions; means of each genotype were used for the correlation. The greenhouse measurements were categorized as continuous (i.e., whole-canopy stomatal conductance, g_{sc},), cumulative or single-point measurements. All measurements were taken 1 week before the stress treatment started. r^2 and p-values indicate the range of weak to strong correlations.

Fig. 1. Atmospheric conditions and experimental progress are represented as the fluctuations in pot weight over the course of the experiment in the greenhouse. **(A)** Daily vapor pressure deficit (VPD) and photosynthetically active radiation (PAR) during the 29 consecutive days of the experiment. **(B)** Continuous weight measurements of all the plants during the 29 days of the experiment. Each line represents one plant/pot. The decreasing slope of the lines during the day indicates that the system lost weight as the plants transpired. The three sharp peaks during the nighttime show system weight gain during irrigation events.

Fig. 2. Total yield and plant weight among three reciprocal-grafted tomatoes grown in the field. Boxplot showing the differences in **(A)** fresh weights of self-grafted plants under the well-irrigated condition and **(B)** fresh weights of self-grafted plants under the limited-irrigation condition. **(C)** Total fruit yield under the well-irrigated condition. **(D)** Total fruit yield under the limited-irrigation condition. **(E)** Total fruit yield of reciprocal-graft plants under the well-irrigated condition and **(F)** total fruit yield of reciprocal-graft plants under the limited-irrigation condition. Data from 2018 are indicated in grey (with small letters) and data from the 2019 experiments are indicated in red (with capital letters). Different letters indicate significantly different means, according to Tukey's Honest Significant Difference test ($p < 0.05$). Box edges represent the upper and lower quantile with the median value

418 shown as a bold line in the middle of the box. Whiskers represent 1.5 times the quantile of

419 the data.

Fig. 3. Daily pattern of whole-canopy stomatal conductance ($gsc(g_{water}-1g_{plant}-1min$) presented as an example of continuous whole plant physiological measurement. Well-irrigated M82 tomato plants were used. The line is an average of three days. Data are shown as means ($\pm$ SE, n=10).

Fig. 4. Correlations between yield components and cumulative transpiration of different tomato genotypes. **(A)** Plant vegetative weight in the field, **(B)** total fruit yield, **(C)** green yield and **(D)** red yield. Measurements taken at harvest time were correlated with the CT throughout the 29 days of the greenhouse experiment. Symbols are the means $\pm$ SE of traits for each genotype under the limited-irrigation condition (circles) and the well-irrigated condition (square box). Vertical SE ($n = 12$–15), Horizontal SE ($n = 8$–10).

Fig. 5. The differential contribution of transpiration periods to yield prediction. **(A)** Mean $\pm$ SE daily transpiration continuously measured during the whole experimental period for all genotypes. We examined the relative contributions of the three phases for yield prediction: well-irrigated (green box), drought treatment (orange), and recovery from drought (light green). CT was measured and correlated with TY for each period: **(B)** pre-treatment, **(C)** drought period and **(D)** recovery period. Vertical SE ($n = 12$–15) field-based data; horizontal SE ($n = 8$–10) greenhouse-based data.

Supplementary Fig. S1. Fruit color and leaf characteristics of M82 and the two mutant lines. **(A, a)** Fruit and leaves of M82. **(B, b)** *Zeta* mutant's fruit and yellowish leaves characteristic. **(C, c)** The *tangerine* mutant's fruit and leaf characteristics.

Supplementary Fig. S2. Carotenoid concentrations in the **(A)** flowers and **(B)** roots of wild-type (M82), TAN and ZET plants and their reciprocal combinations (μg gFW^{-1}). Roots of wild-type tomato contained undetectable level of carotenoids; whereas roots of *tangerine* and *zeta* mutants contained the same compositions of carotenoids typically found in their fruit and flowers. These results indicate that active carotenoid biosynthesis takes place in roots. In the tomato mutants, carotenoid biosynthesis was blocked and, therefore, intermediate products accumulated. This blockage was very effective in the roots due to their lack of exposure to light, which compensates the lack of CRTISO and ZISO carotenoid isomerase in leaves(Isaacson *et al.*, 2002).

Supplementary Fig. S3. Scatter-plot matrices for traits under the well-irrigated condition in 2018. The figure depicts the matrices of scatter plots and Pearson's correlation coefficients among the field-measured yield data and yield components correlated with the greenhouse-based traits of 7 different young tomato plants under the well-irrigated condition. The windows show Pearson's correlation coefficients (*r*) and bivariate scatter plot matrices with a density ellipse

Supplementary Fig. S4. Scatter-plot matrices for traits under the water-deficit condition in 2018. The figure depicts the relationships between field-measured yield and yield components, and the greenhouse-based traits of 7 different young tomato plants that were exposed to drought. The windows show Pearson's correlation coefficients (*r*) and bivariate scatter-plot matrices with a density ellipse

Supplementary **Figure 5: Scatter plot matrix for traits under well irrigated conditions in 2019.** The relationships between fields measured yield and its components and greenhouse-based traits of 7 different young tomato plants under irrigated condition. The windows are Pearson's correlation coefficients (r) and bivariate scatter plots matrix with density ellipse.

Supplementary Fig. S6. Scatter-plot matrices of correlations among studied physiological

yield traits under the water-deficient conditions in 2019. The figure depicts the relationships

between field-measured yield and yield components, and the greenhouse-based traits of 7

different young tomato plants under drought condition. The windows show Pearson's

correlation coefficients (*r*) and bivariate scatter-plot matrices with a density ellipse.

Acknowledgements

The work was funded by the Israel Ministry of Agriculture and Rural Development

(Eugene Kandel Knowledge centers) as part of the "Root of the Matter" – The root zone

knowledge center for leveraging modern agriculture to MM. This research was also

supported by the ISF-NSFC joint research program (grant No. 2436/18) and the United

States–Israel Binational Science Foundation (BSF Grant # 2015100) to MM. D.Z. was

funded by a TOMRES grant (H2020 #727929)

Author contributions

Original research plan conceived by MM, DZ and JH, carried out by SG, AK, and IS. SG, and

AK analyzed the data and wrote the MS draft. All authors contributed to the writing,

reviewing, and editing of the manuscript, and have read and approved the final version.

References

Aisawi KAB, Reynolds MP, Singh RP, Foulkes MJ. 2015. The physiological basis of the genetic progress in yield potential of CIMMYT spring wheat cultivars from 1966 to 2009. Crop Science **55**, 1749–1764.

Anderegg LDL, Anderegg WRL, Berry JA. 2013. Not all droughts are created equal: Translating meteorological drought into woody plant mortality. Tree Physiology **33**, 701–712.

Bahar B, Yildirim M, Barutcular C. 2009. Relationships between stomatal conductance and yield components in spring durum wheat under Mediterranean conditions. Notulae Botanicae Horti Agrobotanici Cluj-Napoca **37**, 45–48.

Burgueño J, Crossa J, Cornelius PL, Yang RC. 2008. Using factor analytic models for joining environments and genotypes without crossover genotype x environment interaction. Crop Science.

Burgueño J, Crossa J, Cotes JM, Vicente FS, Das B. 2011. Prediction assessment of linear mixed models for multienvironment trials. Crop Science.

Chapuis R, Delluc C, Debeuf R, Tardieu F, Welcker C. 2012. Resiliences to water deficit in a phenotyping platform and in the field: How related are they in maize? European Journal of Agronomy.

Chen D, Wang S, Cao B, Cao D, Leng G, Li H, Yin L, Shan L, Deng X. 2016. Genotypic variation in growth and physiological response to drought stress and re-watering reveals the critical role of recovery in drought adaptation in maize seedlings. Frontiers in Plant Science.

Claassen MM, Shaw RH. 1970. Water Deficit Effects on Corn. II. Grain Components 1 .

508 Agronomy Journal.

509 **Claeys H, Inzé D**. 2013. The agony of choice: How plants balance growth and survival

510 under water-limiting conditions. Plant Physiology **162**, 1768–1779.

511 **Crossa J, Cornelius PL**. 1997. Sites regression and shifted multiplicative model clustering

512 of cultivar trial sites under heterogeneity of error variances. Crop Science.

513 **Dalal A, Bourstein R, Haish N, Shenhar I, Wallach R, Moshelion M**. 2019. Dynamic

514 physiological phenotyping of drought-stressed pepper plants treated with "productivity-

515 enhancing" and "survivability-enhancing" biostimulants. Frontiers in Plant Science.

516 **Dalal A, Shenhar I, Bourstein R, Mayo A, Grunwald Y, Averbuch N, Attia Z, Wallach**

517 **R, Moshelion M**. 2020. A telemetric, gravimetric platform for real-time physiological

518 phenotyping of plant–environment interactions. Journal of Visualized Experiments.

519 **Deikman J, Petracek M, Heard JE**. 2012. Drought tolerance through biotechnology:

520 Improving translation from the laboratory to farmers' fields. Current Opinion in

521 Biotechnology **23**, 243–250.

522 **DeLacy IH, Basford KE, Cooper M, Bull JK, McLaren CG**. 1996. Analysis of

523 multienvironment trials-an historical perspective. Plant adaptation and crop improvement.

524 **Du T, Meng P, Huang J, Peng S, Xiong D**. 2020. Fast photosynthesis measurements for

525 phenotyping photosynthetic capacity of rice. Plant Methods.

526 **Enfissi EMA, Nogueira M, Bramley PM, Fraser PD**. 2017. The regulation of carotenoid

527 formation in tomato fruit. The Plant Journal **89**, 774–788.

528 **Eshed Y, Zamir D**. 1995. An introgression line population of Lycopersicon pennellii in the

529 cultivated tomato enables the identification and fine mapping of yield- associated QTL.

530 Genetics.

Fang Y, Xiong L. 2015. General mechanisms of drought response and their application in
drought resistance improvement in plants. Cellular and Molecular Life Sciences.

Fantini E, Falcone G, Frusciante S, Giliberto L, Giuliano G. 2013. Dissection of tomato
lycopene biosynthesis through virus-induced gene silencing. Plant Physiology **163**, 986–
998.

Faralli M, Lawson T. 2020. Natural genetic variation in photosynthesis: an untapped
resource to increase crop yield potential? Plant Journal.

Fu J, Huang B. 2001. Involvement of antioxidants and lipid peroxidation in the adaptation
of two cool-season grasses to localized drought stress. Environmental and Experimental
Botany.

Galpaz N, Wang Q, Menda N, Zamir D, Hirschberg J. 2008. Abscisic acid deficiency in
the tomato mutant high-pigment 3 leading to increased plastid number and higher fruit
lycopene content. Plant Journal.

Gooch JW. 2011. *Centimeter*.

Gosa SC, Lupo Y, Moshelion M. 2019. Quantitative and comparative analysis of whole-
plant performance for functional physiological traits phenotyping: New tools to support
pre-breeding and plant stress physiology studies. Plant Science.

Halperin O, Gebremedhin A, Wallach R, Moshelion M. 2017. High-throughput
physiological phenotyping and screening system for the characterization of
plant???environment interactions. Plant Journal **89**, 839–850.

Han J, Zhang Z, Cao J, Luo Y, Zhang L, Li Z, Zhang J. 2020. Prediction of Winter
Wheat Yield Based on Multi-Source Data and Machine Learning in China.

Heslot N, Akdemir D, Sorrells ME, Jannink JL. 2014. Integrating environmental

covariates and crop modeling into the genomic selection framework to predict genotype by

environment interactions. Theoretical and Applied Genetics.

Hirschberg J. 2001. Carotenoid biosynthesis in flowering plants. Current Opinion in Plant

Biology **4**, 210–218.

Horie T, Yajima M, Nakagawa H. 1992. Yield forecasting. Agricultural Systems.

Isaacson T, Ronen G, Zamir D, Hirschberg J. 2002. Cloning of tangerine from tomato

reveals a Carotenoid isomerase essential for the production of β-carotene and xanthophylls

in plants. Plant Cell **14**, 333–342.

Iseki K, Olaleye O. 2020. A new indicator of leaf stomatal conductance based on thermal

imaging for field grown cowpea. Plant Production Science **23**, 136–147.

Jiang Y, Huang B. 2001. Drought and heat stress injury to two cool-season turfgrasses in

relation to antioxidant metabolism and lipid peroxidation. Crop Science.

Kachanovsky DE, Filler S, Isaacson T, Hirschberg J. 2012. Epistasis in tomato color

mutations involves regulation of phytoene synthase 1 expression by cis-carotenoids.

Proceedings of the National Academy of Sciences **109**, 19021–19026.

Karniel U, Koch A, Zamir D, Hirschberg J. 2020. Development of zeaxanthin-rich

tomato fruit through genetic manipulations of carotenoid biosynthesis. Plant Biotechnology

Journal.

Khaki S, Wang L. Crop Yield Prediction Using Deep Neural Networks. Frontiers in Plant

Science **10**.

Montesinos-López OA, Montesinos-López A, Crossa J, Montesinos-López JC, Mota-

Sanchez D, Estrada-González F, Gillberg J, Singh R, Mondal S, Juliana P. 2018.

Prediction of multiple-trait and multiple-environment genomic data using recommender

systems. G3: Genes, Genomes, Genetics.

Moshelion M. 2020. The dichotomy of yield and draught resistance. EMBO reports, 1–7.

Pirasteh-Anosheh H, Saed-Moucheshi A, Pakniyat H, Pessarakli M. 2016. Stomatal

responses to drought stress. Water Stress and Crop Plants: A Sustainable Approach.

Polania J, Rao I, Cajiao C, Grajales M, Rivera M, Velasquez C, Raatz B, Beebe S.

2017. Shoot and Root Traits Contribute to Drought Resistance in Recombinant Inbred

Lines of MD 23-24 x SEA 5 of Common Bean. Frontiers in Plant Science **8**, 296.

Richards RA, Rebetzke GJ, Condon AG, Van Herwaarden AF. 2002. Breeding

opportunities for increasing the efficiency of water use and crop yield in temperate cereals.

Crop Science **42**, 111–121.

Roberts SL, Dun X, Doddrell RDS, Mindos T, Drake LK, Onaitis MW, Florio F,

Quattrini A, Antonio D, Parkinson DB. 2017. 1 . 1. , 1–19.

Roche D. 2015. Stomatal Conductance Is Essential for Higher Yield Potential of C3 Crops.

Critical Reviews in Plant Sciences.

Saeed W, Naseem S, Ali Z. 2017. Strigolactones biosynthesis and their role in abiotic

stress resilience in plants: A critical review. Frontiers in Plant Science.

Teodoro LPR, Bhering LL, Gomes BEL, Campos CNS, Baio FHR, Gava R, da Silva

CA, Teodoro PE. 2019. Understanding the combining ability for physiological traits in

soybean. PLoS ONE **14**, 1–13.

Tilkat E, Akdemir H, Onay A, Tilkat EA. 2012. *Crop Production for Agricultural*

Improvement.

Tolk JA, Howell TA. 2009. Transpiration and yield relationships of grain sorghum grown

in a field environment. Agronomy Journal **101**, 657–662.

Weber VS, Araus JL, Cairns JE, Sanchez C, Melchinger AE, Orsini E. 2012.

Prediction of grain yield using reflectance spectra of canopy and leaves in maize plants

grown under different water regimes. Field Crops Research.

de Wit C. 1958. Transpiration and crop yields. Versl. Landbouwk. Onderz. 64. 6., 18–20.

Zhang J, Jiang H, Song X, Jin J, Zhang X. 2018. The responses of plant leaf CO_2/H_2O

exchange and water use efficiency to drought: A meta-analysis. Sustainability

(Switzerland) **10**, 1–13.

Fig. 1. Atmospheric conditions and experimental progress are represented as the fluctuations in pot weight over the course of the experiment in the greenhouse. **(A)** Daily vapor pressure deficit (VPD) and photosynthetically active radiation (PAR) during the 29 consecutive days of the experiment. **(B)** Continuous weight measurements of all the plants during the 29 days of the experiment. Each line represents one plant/pot. The decreasing slope of the lines during the day indicates that the system lost weight as the plants transpired. The three sharp peaks during the nighttime show system weight gain during irrigation events.

Fig. 2. Total yield and plant weight among three reciprocal-grafted tomatoes grown in the field. Boxplot showing the differences in (**A**) fresh weights of self-grafted plants under the well-irrigated condition and (**B**) fresh weights of self-grafted plants under the limited-irrigation condition. (**C**) Total fruit yield under the well-irrigated condition. (**D**) Total fruit yield under the limited-irrigation condition. (**E**) Total fruit yield of reciprocal-graft plants under the well-irrigated condition and (**F**) total fruit yield of reciprocal-graft plants under the limited-irrigation condition. Data from 2018 are indicated in grey (with small letters) and data from the 2019 experiments are indicated in red (with capital letters). Different letters indicate significantly different means, according to Tukey's Honest Significant Difference test ($p < 0.05$). Box edges represent the upper and lower quantile with the median value shown as a bold line in the middle of the box. Whiskers represent 1.5 times the quantile of the data.

Fig. 3. Daily pattern of whole-canopy stomatal conductance (gsc(g_{water}-1g_{plant}-1min) presented as an example of continuous whole plant physiological measurement. Well-irrigated M82 tomato plants were used. The line is an average of three days. Data are shown as means (± SE, n=10).

			Field-Based Measurements			
			Total Yield		Plant fresh weight	
			R^2 range	p-Value ranges	R^2 range	p-Value range
Greenhouse-Based Measurements	Continuous	$g_{sc(7:00-10:00)}$	0.55 to 0.90	0.19 to 0.004	0.2 to 0.60	0.62 to 0.14
		$g_{sc(10:00-13:00)}$	0.45 to 0.72	0.304 to 0.067	0.44 to 0.89	0.32 to 0.006
		$g_{sc(13:00-16:00)}$	0.34 to 0.71	0.447 to 0.073	0.74 to 0.93	0.054 to 0.002
	Single point	Cumulative transpiration	0.77	0.04	0.94	0.001
		Growth rate	0.62	0.12	0.89	0.0065
		Plant net weight	0.70	0.076	0.79	0.038

Table 1. Correlations between the physiological traits of young tomato plants in the greenhouse and their field-based yield and biomass under well-irrigated conditions; means of each genotype were used for the correlation. The greenhouse measurements were categorized as continuous (i.e., whole-canopy stomatal conductance, g_{sc}), cumulative or single-point measurements. All measurements were taken 1 week before the stress treatment started. r^2 and *p*-values indicate the range of weak to strong correlations.

Fig. 4. Correlations between yield components and cumulative transpiration of different tomato genotypes. **(A)** Plant vegetative weight in the field, **(B)** total fruit yield, **(C)** green yield and **(D)** red yield. Measurements taken at harvest time were correlated with the CT throughout the 29 days of the greenhouse experiment. Symbols are the means ± SE of traits for each genotype under the limited-irrigation condition (circles) and the well-irrigated condition (square box). Vertical SE (*n* = 12–15), Horizontal SE (*n* = 8–10).

Fig. 5. The differential contribution of transpiration periods to yield prediction. (**A**) Mean ± SE daily transpiration continuously measured during the whole experimental period for all genotypes. We examined the relative contributions of the three phases for yield prediction: well-irrigated (green box), drought treatment (orange), and recovery from drought (light green). CT was measured and correlated with TY for each period: (**B**) pre-treatment, (**C**) drought period and (**D**) recovery period. Vertical SE ($n = 12$–15) field-based data; horizontal SE ($n = 8$–10) greenhouse-based data.

Supplementary figures

Supplementary Fig. S1. Fruit color and leaf characteristics of M82 and the two mutant lines. **(A, a)** Fruit and leaves of M82. **(B, b)** *Zeta* mutant's fruit and yellowish leaves characteristic. **(C, c)** The *tangerine* mutant's fruit and leaf characteristics.

Supplementary Fig. S2. Carotenoid concentrations in the **(A)** flowers and **(B)** roots of wild-type (M82), TAN and ZET plants and their reciprocal combinations (µg gFW⁻¹). Roots of wild-type tomato contained undetectable level of carotenoids; whereas roots of *tangerine* and *zeta* mutants contained the same compositions of carotenoids typically found in their fruit and flowers. These results indicate that active carotenoid biosynthesis takes place in roots. In the tomato mutants, carotenoid biosynthesis was blocked and, therefore, intermediate products accumulated. This blockage was very effective in the roots due to their lack of exposure to light, which compensates the lack of CRTISO and ZISO carotenoid isomerase in leaves(Isaacson *et al.*, 2002).

Supplementary Fig. S3. Scatter-plot matrices for traits under the well-irrigated condition in 2018. The figure depicts the matrices of scatter plots and Pearson's correlation coefficients among the field-measured yield data and yield components correlated with the greenhouse-based traits of 7 different young tomato plants under the well-irrigated condition. The windows show Pearson's correlation coefficients (r) and bivariate scatter plot matrices with a density ellipse

Fig.S3

Fig.S3

Fig.S3

Supplementary Fig. S4. Scatter-plot matrices for traits under the water-deficit condition in 2018. The figure depicts the relationships between field-measured yield and yield components, and the greenhouse-based traits of 7 different young tomato plants that were exposed to drought. The windows show Pearson's correlation coefficients (*r*) and bivariate scatter-plot matrices with a density ellipse

Fig.S4

Fig.S4

Fig.S4

Supplementary **Figure 4 continued...**

Fig.S4

Supplementary **Figure 5: Scatter plot matrix for traits under well irrigated conditions in 2019.** The relationships between fields measured yield and its components and greenhouse-based traits of 7 different young tomato plants under irrigated condition. The windows are Pearson's correlation coefficients (r) and bivariate scatter plots matrix with density ellipse.

Fig.S5

Fig.S5

Fig.S5

Supplementary Fig. S6. Scatter-plot matrices of correlations among studied physiological yield traits under the water-deficient conditions in 2019. The figure depicts the relationships between field-measured yield and yield components, and the greenhouse-based traits of 7 different young tomato plants under drought condition. The windows show Pearson's correlation coefficients (*r*) and bivariate scatter-plot matrices with a density ellipse.

Fig.S6