Title page

Title: Differential responses of amphibians and reptiles to land-use change in the biodiversity hotspot of north-eastern Madagascar

Running Title: Herpetofauna response to land-use change in Madagascar

Authors' name: Thio Rosin Fulgence[1,2,3], Dominic Andreas Martin[3,4], Romual Randriamanantena[5], Ronick Botra[5], Erosiniot Befidimanana[1], Kristina Osen[6], Annemarie Wurz[7], Holger Kreft[3,8], Aristide Andrianarimisa[2], Fanomezana Mihaja Ratsoavina[2]

Authors' affiliations: [1] Natural and Environmental Sciences, Regional University Centre of the SAVA Region (CURSA), Antalaha, Madagascar

[2] Zoology and Animal Biodiversity, Faculty of Sciences, University of Antananarivo, Madagascar

[3] Biodiversity, Macroecology and Biogeography, University of Goettingen, Goettingen, Germany

[4] Wyss Academy for Nature, University of Bern, Bern, Switzerland

[5] Sciences of life and Environmental Department, Faculty of Sciences, University of Antsiranana, Madagascar

[6] Tropical Silviculture and Forest Ecology, University of Goettingen, Goettingen, Germany

[7] Agroecology, University of Goettingen, Goettingen, Germany

[8] Centre for Biodiversity and Sustainable Land Use (CBL), University of Goettingen, Goettingen, Germany

Corresponding author: Thio Rosin Fulgence; Natural and Environmental Sciences, Regional University Centre of the SAVA Region (CURSA), Antalaha, Madagascar;

Zoology and Animal Biodiversity, Faculty of Sciences, University of Antananarivo, Antananarivo, Madagascar,

29   Biodiversity, Macroecology and Biogeography, University of
30   Goettingen, Goettingen, Germany

31   Email: thiorosinf@yahoo.fr / Telephone: +261 32 029 5174

Abstract

Large expanses of tropical rainforest have been converted into agricultural landscapes
cultivated by smallholder farmers. This is also the case in north-eastern Madagascar; a
region that retains significant proportions of forest cover despite slash-and-burn hill rice
cultivation and vanilla agroforestry expansion. The region is also a global hotspot for
herpetofauna diversity, but how amphibians and reptiles are affected by land-use change
remains largely unknown. Using a space-for-time study design, we compared species
diversity and community composition across seven prevalent land uses: unburned (old-
growth forest, forest fragment, and forest-derived vanilla agroforest) and burned (fallow-
derived vanilla agroforest, woody fallow, and herbaceous fallow) land-use types, and rice
paddy. We conducted six comprehensive, time-standardized searches across at least ten
replicates of each land-use type and applied genetic barcoding to confirm species
identification. We documented an exceptional diversity of herpetofauna (119 species; 91%
endemic). Plot-level amphibian species richness was significantly higher in old-growth forest
than in all other land-use types. Plot-level reptile species richness was significantly higher in
unburned land-use types compared to burned land-use types. For both amphibians and
reptiles, the less-disturbed land-use types showed more uneven communities and the
species composition in old-growth forest differed significantly from all other land-use types.
Amphibians had higher forest dependency (38% of species occurred exclusively in old-
growth forest) than reptiles (26%). Our analyses thus revealed that the two groups respond
differently to land-use change: we found less pronounced losses of reptile species richness
especially in unburned agricultural habitats, suggesting that reptiles are less susceptible to
land-use change than amphibians, possibly due to their ability to cope with hotter and drier
microclimates. Overall, old-growth forest harboured a unique diversity, but some species also
thrived in vanilla agroforestry systems, especially if these were forest-derived. This highlights
the importance of conserving old-growth forests and non-burned land-use types within
agricultural landscapes.

Key words: agroecology, agroforestry, community ecology, herpetofauna, human-
dominated landscape, land-use history, slash-and-burn cultivation, vanilla, forest
dependency

1.Introduction

Demand for agricultural goods is on the rise due to a growing world population, rising per-capita consumption, and changing diets (Tilman et al. 2011). This situation is leading to both an expansion of croplands into natural areas and an intensification within existing production systems (Tscharntke et al. 2012). Most agricultural expansion in the tropics happens at the expense of forests and leads to an increase in forest fragmentation (Hansen et al. 2020). This expansion affects all aspects of biodiversity: research has documented losses in species richness (Scales and Marsden 2008), shifts in species composition (Newbold et al. 2016), reductions of functional diversity (Matuoka et al. 2020), and erosion of phylogenetic diversity (Li et al. 2020). Importantly, the response of biodiversity to land-use change differs among species groups (Newbold et al. 2014) and world regions (Williams and Newbold 2020), but certain groups and regions are understudied compared to others. While vertebrates are comparatively well studied, we know less about the responses of amphibians and reptiles than we know about mammals and birds (Newbold et al. 2014). Similarly, we know less about what happens in the more biodiverse tropics compared to temperate regions (Gardner et al. 2009). In sum, we know that land-use change is the main driver of biodiversity decline globally (Powers and Jetz 2019), but how the diversity of certain understudied taxa differs between land-use types remains an open question, especially in the tropics, where high land-use pressure and biodiversity coincide (Newbold et al. 2020).

Tropical agricultural landscapes contribute to food security but also provide opportunities for nature conservation (Perfecto and Vandermeer 2010). The quality of the agro-ecosystem for biodiversity conservation depends on the farming systems. Agricultural landscapes dominated by large-scale industrial monocultures have lower conservation value than diverse mosaics of forest fragments, agroforestry systems and more intensively farmed annual crop fields (Mendenhall et al. 2016; Murray and Nowakowski 2021). For example, Ricciardi et al. (2021) reported that small-scale agricultural landscapes host more biodiversity than large-scale ones and De Palma et al. (2015) show that low-intensity agriculture can maintain relatively diverse bee communities. Furthermore, forest patches within tropical small-scale agricultural landscapes increase the variety of ecological niches and benefit species with small ranges in particular (Gray et al. 2016). Besides the value for biodiversity, small-scale land-use mosaics can also provide essential ecosystem services and livelihoods for rural people, making such landscapes work for humans and nature (Kremen and Merenlender 2018). Most research investigating the value of tropical agricultural landscapes for biodiversity and humans was conducted in the Neotropics (Mendenhall et al. 2016) while the conservation value of Afrotropical agricultural landscapes is less understood (Powers et al. 2011). Furthermore, it remains largely unclear how different forms of forest conversion – that is with the use of fire (slash-and-burn shifting cultivation) or without the use of fire (forest

degradation and forest-derived agroforestry) – affect amphibian and reptile diversity in resulting land-use types.

Here, we investigate how herpetofauna taxonomic diversity and species composition differ across prevalent land-use types in Madagascar, providing insights into the vulnerability of amphibians and reptiles to various forms of land-use change. Madagascar has lost 44% of forest cover since the 1950s, mainly due to the transformation of natural ecosystems to agricultural lands (Vieilledent et al. 2018). Along with unsustainable extraction rates (Whitehurst et al. 2009), this has resulted in more than half of evaluated Malagasy vertebrate species being at risk of extinction (IUCN 2019). Outstanding levels of endemism (Goodman and Benstead 2005) and ongoing threats qualify Madagascar as a global biodiversity hotspot (Myers et al. 2000). While the forests and protected areas of the island are increasingly well surveyed, the biodiversity in the agricultural landscapes is less known (Irwin et al. 2010). Focusing on amphibians and reptiles is particularly relevant since Madagascar has a diverse and highly endemic herpetofauna: amphibian species diversity is currently estimated at around 370 (AmphibiaWeb 2020) and almost all species are endemic (Goodman and Benstead 2005). Reptile diversity stands at around 440 species (Uetz et al. 2021) with 91% endemism (Goodman and Benstead 2005). Due to cryptic taxonomic complexes, many species still await discovery or description, suggesting that total species richness numbers will increase further (Vieites et al. 2009). Globally, herpetofauna is sensitive to various anthropogenic threats (Vallan 2000) including chytrid fungi, environmental pollution, collection for the pet trade, climate change, and conversion of forest habitat into agricultural lands (Hof et al. 2011). All of these factors are also threatening Malagasy herpetofauna, with deforestation being the top pressure (Cordier et al. 2021).

Our study focuses on north-eastern Madagascar. The area retains more forest cover than other parts of the country (Vieilledent et al. 2018) and is a global priority area for amphibian research (Nori et al. 2018). Besides being known for its remarkable biodiversity, north-eastern Madagascar is also a global centre for vanilla cultivation (Hänke et al. 2018). The price boom of the spice between 2012 and 2019 has triggered an expansion of vanilla agroforests (Llopis et al. 2019), and roughly 80% of rural households in the study region farm vanilla (Hänke et al. 2018). How the expansion of vanilla agroforestry is impacting biodiversity is not well known, despite that the majority of global vanilla cultivation takes place across three biodiversity hotspots (Madagascar, Indonesia and Mexico; Myers et al. 2000, FAO 2020). Investigating how vanilla agroforestry compares to other land-uses in terms of biodiversity conservation thus represents a research gap. Besides farming vanilla, the rural population in north-eastern Madagascar also practices slash-and-burn shifting cultivation for hill rice production. Valleys and plains in the study region are commonly occupied by irrigated rice paddies, forming the backbone of staple crop supply. We thus

compared the taxonomic diversity and community composition of amphibians and reptiles between six land-use types within the small-scale agricultural landscape and compared this to continuous old-growth forest inside Marojejy National Park. Within the small-scale agricultural landscape, we sampled vanilla agroforest of contrasting land-use history (forest- and fallow-derived vanilla agroforests), as well as forest fragment, herbaceous fallow, woody fallow and rice paddy. Following Cordier et al. (2021) and Gonzalez and González-Trujillo (2021), we hypothesised that taxonomic diversity would be highest in old-growth forest (for both amphibians and reptiles) and expected losses and shifts in species composition after conversion, particularly in those cases where land-use conversion involves the use of fire for slash-and-burn shifting cultivation.

2.Methods

2.1. Study region and study design

We conducted our study in the SAVA region in north-eastern Madagascar (Fig. 1A and B) where forests outside protected areas are now highly fragmented (Vieilledent et al. 2018) and the landscape is dominated by smallholder agriculture.

We collected data on circular plots with a 25 m radius at low to mid-altitude (7-819 m above sea level; mean = 192 m±207 m) surrounding ten villages and in Marojejy National Park, a UNESCO World Heritage Site. In each village, we selected seven plots: three vanilla agroforests (forest-derived and/or fallow-derived vanilla agroforest), one forest fragment, one herbaceous fallow, one woody fallow, and one rice paddy. In the ten villages, we first selected 30 vanilla agroforests along a canopy cover gradient. After consultation with the agroforest owner and a visual confirmation on the plot, we found that twenty vanilla agroforests were fallow-derived while ten agroforests were forest-derived. This approach allowed us to understand the response of amphibians and reptiles to the land-use history of vanilla agroforests. Additionally, we chose ten plots at two sites (five plots each) inside Marojejy National Park. All land-use types, except fallow-derived vanilla agroforests, were replicated ten times. The average minimum distance between one plot and the next closest plot was 719 m± 438 m, while the smallest minimum distance between two plots was 260 m. In total, we surveyed 80 plots (Fig. 1C and D).

2.2. Sampled land-use types

We selected 10 *old-growth forest* plots at two sites with five plots each. One of the sites has experienced some selective logging in the past but is now well protected (touristic zone in Manantenina valley; site *Tourist* on Fig. 1), the other site suffers from ongoing occasional selective logging and trapping (Bangoabe; site *East* on Fig. 1), but we chose plots that did not show signs of recent disturbance. The old-growth forest plots are a minimum of 300 m from the National Park boundary. In the study region, isolated *forest fragments* occur around

villages and represent remnants of the continuous forest cover that existed in the region prior
to the large-scale deforestation that began in the early 20[th] century (Gade 1996). The ten
fragments have not been burned in historic times. The forest fragments are all used for the
extraction of timber and non-timber forest products. *Herbaceous fallows* occur after shifting
slash-and-burn hill rice cultivation (locally referred to as *tavy*; Styger et al. 2007) and are
sometimes used for grazing. The herbaceous fallow plots in this study had last burned at the
end of 2016, one year before the onset of data collection in 2017. *Woody fallows* represent
successional stages of herbaceous fallows, containing shrubs and small trees. Woody
fallows are also occasionally grazed by cows. The woody fallows in our study had last burnt
4-16 years before the onset of data collection in 2017, according to the reports of land-
owners. The climbing vanilla orchid (*Vanilla planifolia)* is farmed in agroforestry systems with
two distinct land-use histories (following Martin et al. 2020): *forest-derived vanilla*
agroforests, where vanilla is directly planted inside the forest after removing understory trees
and shrubs and managing tall trees for shade (not involving burning). In *fallow-derived vanilla*
agroforests, vanilla is planted on fallow land which resulted from shifting slash-and-burn hill
rice cultivation in the past. In these fallow-derived vanilla agroforests, farmers allow natural
regeneration of trees or plant trees to provide shade or as support structures for the vanilla
vines. Lastly, we studied irrigated *rice paddies* that occur in valley bottoms and flood plains
throughout the whole study region. Rice is planted and harvested between one to three times
per year. The rice paddies chosen for our study had wider-than-average banks to facilitate
movement and sampling within the plots.

2.3. Sampling, data collection, and field identification

To collect data in the villages, we organized two sampling campaigns during the driest period
of the year (October to December 2017 and late August to December 2018) and one
campaign during the wettest period (Mid-January to early April 2018). In Marojejy National
Park, we also organized two sampling campaigns during the driest period (late August –
early September 2018 and December 2018) and one during the wettest period (February
2019). During each campaign, we visited each plot once during the day (08:00 - 17:00) and
once at night (18:30-23:00). Overall, we did six visits per plot, three times during the day and
three times at night.

We collected data on amphibian and reptile communities during time-standardized searches
(Kadlec et al. 2012). During the survey, we systematically searched each plot in a zig-zag
pattern (Kadlec et al. 2012). Each search was standardized to 45 minutes of searching time
by two observers. In sum, we thus conducted 270 minutes of searching time on each plot,
summing up to 408 hours of searching time across all plots. To detect individuals hiding
under rocks, in leaf axils, tree barks, tree holes, leaf-litter, or deadwood, we actively
inspected those microhabitats by lifting removable objects to check underneath.

To identify individuals to species level during fieldwork, we used morphological
characteristics and referred to keys and visualisations available in 'A Field Guide to
Amphibians and Reptiles of Madagascar' (Glaw and Vences 2007) and additional literature
(Rakotoarison et al. 2017; Ratsoavina et al. 2019). For those individuals which we could not
identify with confidence in the field with the available literature, we extracted tissue samples
for follow-up DNA analysis and/or collected the specimen. To denote those individuals, we
used 'cf.' (confer or compare with) and 'sp. aff.' (affinis) in conjuncture with the genus name
when we found slightly different morphological characters compared to the literature that we
used or compared to a given identified species. We used sp. (sp1 or sp2) in conjuncture with
the genus name when we could not find any given identified species to compare with the
encountered individual. The sp. denotation was thus used to differentiate several unknown
species belonging to the same genus and we consider them as morphospecies but used
DNA barcodes to confirm species identification (see below). Following the field identification,
we resumed the searching time; thus abundance and diversity of amphibians and reptiles
were independent of the time spent on the plot. Throughout this manuscript, we refer to each
encountered individual as an 'encounter' rather than an individual as we cannot exclude the
possibility of having encountered the same individual at more than one sampling event.

2.4. Species identification with DNA barcodes
We collected muscle or toe clips as tissue samples of individuals in case of non-reliable
morphological characteristics based identification. We preserved tissue samples in
Eppendorf tubes with 90% ethanol, stored and analysed at the Evolutionary Biology
laboratory of Prof. Miguel Vences at the University of Braunschweig, Germany. We
preserved specimens in 70% ethanol for amphibians and 70% and deposited them in the
collection of the Regional University Centre of the SAVA region (CURSA), Antalaha,
Madagascar. We extracted genomic DNA following the standard single-tube salt extraction
protocol (Bruford et al. 1992). To do so, we cut small pieces from the collected tissue and
proceeded to protein digestion following the protocol of Cacciali et al. (2019). We then
passed the extracted DNA to a PCR (Polymerase Chain Reaction) thermocycler, in which we
amplified DNA fragments of two mitochondrial genes, "16S" (Vences et al. 2005) and "COI"
with the primers 16SAL (5´ - CGCCTGTTTATCAAAAACAT - 3´) and 16SBH (5´ -
CCGGTCTGAACTCAGATCACGT - 3´) from Palumbi et al. (1991). Amplification was set for
45 seconds with 94-96 °C as denaturation, for 45 seconds with 52 °C as hybridization and for
90 seconds with 72°C as elongation. The whole process was repeated for 35 cycles. DNA
sequences are generated by sequencer. We first cleaned the obtained sequences with the
software CodonCode Aligner version 8.0.2 (Codon Code Corporation, Centerville, MA, US).
Then, we processed newly generated sequences for additional quality control with a Blast
search (Altschul et al. 1990) which queries similar DNA sequences through GenBank

database to provide information on closest taxa or potential contamination. Finally, we built a
phylogenetic tree using MEGA-X (Kumar et al. 2018) to reveal clusters of each newly
generated sequence, allowing us to reliably identify individuals to species level. We
submitted all newly determined sequences to GenBank under accession numbers MZ019097
- MZ019430. In this study, the DNA barcoding differentiated between 43 species of
amphibians and 24 species of reptiles. Among them were 12 candidates for new species of
which eight were documented for the first time (marked as sp. CaNEW) and four were
already found by other researchers (marked as sp. Ca) but have not yet been described. The
new species were based on a comparison of obtained DNA sequences to available
sequences with the same extraction DNA method conducted in the same lab. In case the
DNA barcoding method did not yield results due to short sequences or contaminations, we
kept the field identification for those species (for 15 species, eight for amphibians and seven
for reptiles).

We attributed the red list status to evaluated species following IUCN (2019). We determined
endemicity according to AmphibiaWeb (2020) for amphibians and Uetz et al. (2021) for
reptiles. Because most amphibian and reptile genera are completely endemic to
Madagascar, we could attribute endemicity also to morphospecies.

2.5. Data analysis and visualisation: species diversities and encounters
During the data analysis, we conducted all statistical using R version 3.5.1 (R Core Team
2019) and R package 'ggplot2' (Wickham 2016) for visualisation. To compare observed
species richness and species encounters on plot-level among land-use types, we computed
a generalized linear model with species richness as a response, land-use type as the
explanatory variable, and village (respectively old-growth forest site) as a random factor and
with Poisson family for count data as distribution. We then ran the *glht* function of the R-
package 'multcomp' (Hothorn et al. 2008) applying a Tukey all-pair-comparison with
Bonferroni correction. We then used the same approach to compare observed species
between unburned plots (old-growth forest, forest fragment, forest-derived vanilla) and
burned plots (herbaceous fallow, woody fallow, fallow-derived vanilla).

We used the encounter data to compute sample-size-based extrapolation curves (Hsieh et
al. 2016) with the 'iNEXT' package to assess the diversity (species richness, Shannon, and
Simpson diversity) across land-use types using the Hill number framework (Chao et al.
2014). We calculated species richness, Shannon diversity and Simpson diversity (Hsieh et
al. 2016). Chao and Jost (2012) define q=0 (0D) as species richness, i.e. the effective
number of species in the community, giving equal weight to frequent and infrequent species;
q=1 (1D) as Shannon diversity, giving more weight to more frequently observed species and
q=2 (2D), as Simpson diversity interpreted as the effective number of abundant species. This

approach is now well established given key advantages over traditional Shannon and
Simpson diversity indices (Roswell et al. 2021).

To display the total species diversity in each land-use type, we sub-sampled 10 plots of the
20 fallow-derived vanilla agroforests. To do so, we randomly selected one fallow-derived
vanilla agroforest from each village. As one of the villages lacks fallow-derived vanilla
agroforests (village Andramanolotra, see Fig.1), this resulted in 9 agroforests. We proceeded
to select one additional plot from all remaining fallow-derived vanilla agroforests, enabling a
fair comparison of total species diversity across 10 plots of each land-use type.

To compare the evenness of communities between land-use types, we additionally plotted
the observed and extrapolated diversity across hill numbers for each land-use type, allowing
for a direct comparison of slopes between land-use types (Chao and Jost 2015; Roswell et
al. 2021).

2.6. Data analysis and visualization: species composition
To evaluate the differences in species community composition between land-use types,
between burned and unburned plots, and between villages, we used the *metaMDS* function
of the R-package 'vegan'(with 1000 permutations, Oksanen et al. 2020). We used non-
metric dimensional scaling (NMDS) of Bray-Curtis dissimilarities to visualize the dissimilarity
of species composition in two dimensions. Furthermore, to test the differences between land-
use types, we used PERMANOVA (Permutational multivariate analysis of variance)
implemented in the *adonis* function of the 'vegan' package (Oksanen et al. 2020) and
computed pairwise differences using the *pairwise.adonis* function with Bonferroni correction
of the '*pairwise.adonis*' package (Martinez 2020). Since PERMANOVA may confound
location and dispersion, we used PERMDISP (Permutational analysis of multivariate
dispersions) to see if PERMANOVA results may have been influenced by dispersion
differences between groups (Anderson and Walsh 2013). To do so, we used the function
betadisper and *permutest* and performed pairwise comparisons using the *TukeyHSD*
function. To analyse the degree of forest dependency (Rembold et al. 2017), we plotted the
proportion of encounters for each species across land-use types. To visualize the forest
dependency, we used the *barplot* function of R graphics version 3.5.1.

3. Results

3.1. Encounters and observed species richness
In total, we made 6215 encounters and observed 119 species of amphibians and reptiles.
The 3694 amphibian encounters belong to 58 species, 15 genera, and four families (see SI
1). The most species-rich genera of amphibians are *Boophis* (11 species), *Stumpffia* (10
species) and *Gephyromantis* (eight species). We found all but one species to be endemic

according to AmphibiaWeb (2020). Among the observed amphibian species, 22 (37%) could
not be identified to species level in the field. The 12 of them were recognized as candidates
for new species based on genetic barcoding, another eight of them were counted as
morphospecies based on criteria mentioned in the methods section and two were conferred
to identified species. Among the encountered amphibian species, seven are listed in the
'threatened' category (IUCN 2019). Among the threatened species, we recorded six
vulnerable and one endangered species.

The 2521 reptile encounters represent 61 species, 28 genera, and five families (see SI 1).
The most species-rich reptile genera were *Phelsuma* (11 species) and *Uroplatus* (six
species). We found 83% of reptile species to be endemic (Uetz et al. 2021). Amongst
observed reptile species, 15 could not be identified to species level in the field, seven of
them were counted as morphospecies and eight were conferred to identified species. Based
on the IUCN red list (IUCN 2019), seven encountered reptile species are listed as
'threatened'. Among the threatened species, we recorded four endangered and one critically
endangered species.

Plot-level observed amphibian species richness differed significantly among land-use types
($F_{6, 73}$ =19.59, p-value < 0.001) and when comparing all burned to all unburned land-use
types ($F_{1, 67}$ =15.89, p-value < 0.001; Figure SI 6 and table SI 7). A Tukey post-hoc test
revealed significant pairwise differences between pairs of land-use types (Fig. 2A and SI 2):
Old-growth forests plots had significantly higher observed amphibian species richness while
rice paddies plots had significantly lower compared to other land-use types (Fig. 2A). The
other land-use types had similar amphibian species richness with no significant differences
(see SI 2). Plot level amphibian encounters also differed significantly between land-use types
($F_{6, 72}$ =13.01, p-value < 0.001; Figure SI 9 and table SI 10 and SI 12).

Plot-level observed reptile species richness also varied significantly among land-use types
($F_{6, 73}$ =18.55, p-value < 0.001) and when comparing all burned to all unburned land-use
types ($F_{1, 67}$ =25.60, p-value < 0.001; Figure SI 6 and table SI 8). Tukey post-hoc tests
revealed no differences in observed species richness between old-growth forest and forest
fragment (p-value = 0.88) and between old-growth forest and forest-derived vanilla
agroforest (p-value = 0.77). These three land-use types are recorded with the highest
observed species richness, with 10 reptile species on average per plot. Rice paddy had the
lowest reptile species richness but there were no significant differences (p-value = 0.19)
compared to herbaceous fallow. Reptile species richness was significantly higher in forest-
derived vanilla agroforest than in fallow-derived vanilla agroforest (p-value = 0.01; Fig. 2B
and SI 2). Plot-level reptile encounters also differed significantly between land-use types ($F_{6,}$
$_{73}$ =14, p-value < 0.001; Figure SI 9 and table SI 11 and SI 12).

3.2. Species accumulation curves, diversity, evenness, and estimated species richness

Encounter-based accumulation curves for amphibians revealed the highest species richness, Shannon and Simpson diversity in old-growth forest and the lowest in rice paddy. In most land-use types, observed amphibian species richness curves flattened off, except for old-growth forest and forest fragment. The overlap of the 95% confidence interval of extrapolated amphibian richness for old-growth forest and forest fragment indicates no differences in species richness. Regarding the accumulated amphibian species diversity for Shannon and Simpson diversity, old-growth forest varied significantly compared to all other land-use types. Shannon and Simpson diversity also did not differ significantly between forest fragment and forest-derived vanilla agroforest, respectively, compared to burned land uses, except to rice paddy (Fig. 3, Table 1).

Encounter-based accumulation curves for reptiles revealed the highest species richness and Shannon diversity in old-growth forest and the lowest in herbaceous fallow. Observed species richness curves flattened off in all land-use types except woody fallow and fallow-derived vanilla agroforest. We also found an overlap of the 95% confidence intervals of extrapolated species richness as well as Shannon and Simpson diversity within unburned and burned land-use types except for Shannon and Simpson diversity in burned land-use types (Fig. 3, Table 1).

The species diversity dropped more strongly across hill number orders (q=0 to q=2) in amphibians than in reptiles after old-growth forest transformation (Fig. 3, SI 3 & SI 5), highlighting that amphibian communities were more uneven than reptile communities (SI 5). Furthermore, the comparison shows that old-growth forest communities of both amphibians and reptiles were most uneven (SI 5).

3.3. Species composition and forest dependency

The composition of amphibian communities differed significantly across land-use types (PERMANOVA: R^2=0.50, p-value<.001, Df=6), partly driven by differences in group dispersion (PERMDISP: F=4.400, p-value=0.004, Df=6, Table SI 18 – SI 19). Pairwise comparisons showed that amphibian communities in old-growth forest and rice paddy were significantly different from those of other land-use types. Forest fragments differed significantly from fallow-derived vanilla, woody fallow and herbaceous fallow, but no significant differences were observed between forest fragment and forest-derived vanilla. We found no significant differences between forest-derived vanilla, fallow-derived vanilla, woody fallow, and herbaceous fallow for amphibian communities (Fig. 5A, see SI 4). We compared also the amphibian communities between unburned and burned land-use types and found significant differences (PERMANOVA: R^2=0.193, p-value<.001, Df=1, Fig. SI 13 and Table SI

14), again partly drive by dispersion (PERMDISP: F=34.718, p-value=.001, Df=1, Table SI

16). Similarly, amphibian communities differed between villages and Marojejy National Park

communities (PERMANOVA: R^2= 0.416, p-value<.001, Df=10, Fig. SI 21 and Table SI 22

and SI 24), in this case not driven by differences in group dispersion (PERMDISP: F=1.254,

p-value=0.283, Df=11, Table SI 25). Here, communities were very similar between villages

(Fig SI 21).

Reptile species composition showed significant differences among land-use types

(PERMANOVA: R^2=0.40, p-value<.001, Df=6) and non-significant differences within-group

dispersion (PERMDISP: F=1.870, p-value=0.088, Df=6, SI 20) indicating the homogeneity of

group dispersion. Pairwise comparisons showed that the reptile communities in old-growth

forest and forest fragment were significantly different from all other land-use types. The

communities in other land-use types showed no significant differences (Fig. 5B, SI 4). Old-

growth forest showed unique communities of reptiles and amphibians. The reptile species

composition between unburned and burned land-use types showed significant differences

(PERMANOVA: R^2=0.210, p-value=.001, Df=1, Table SI 15) not driven by differences in

group dispersion (PERMDISP: F=3.918, p-value=0.057, Df=1, Table SI 17). The reptile

communities between villages and Marojejy National Park showed significant differences

(PERMANOVA: R^2=0.389, p-value<.001, Df=10, Table SI 23 and SI 24) not driven by

differences in group dispersion (PERMDISP: F=0.835, p-value=0.626, Df=11, Table SI 26).

Finally, the analyses of forest dependency revealed that 22 amphibian species (38%) were

found exclusively in old-growth forest, 11 amphibian species (19%) exclusively in forest

fragment, three amphibian species (5%) exclusively in forest-derived vanilla agroforest, and

four amphibian species (7%) exclusively in fallow-derived vanilla agroforest (Fig. 4A).

We observed 16 reptile species (26%) exclusively in old-growth forest, three reptile species

(5%) exclusively in forest fragment, four reptile species (7%) exclusively in forest-derived

vanilla agroforest, and one reptile species (2%) exclusively in fallow-derived vanilla

agroforest (Fig. 4B). No species were exclusively found in woody fallow, herbaceous fallow

and rice paddy for both groups.

418 4.Discussion

Based on a rigorous study design, extensive field sampling, and DNA-based species

identification, we provide a comprehensive assessment of the response of amphibian and

reptile species diversity to land-use change in the biodiversity hotspot of north-eastern

Madagascar. Overall, we document a highly diverse herpetofauna with 58 amphibian and 61

reptile species and a high proportion of endemic species, 98% for amphibians and 83% for

reptiles.

At a plot-level, species richness for both amphibians and reptiles was very high, with up to 12
and 14 species, respectively. For both groups, old-growth forest was significantly different
from all other land-use types in terms of total species richness and community composition.
Rice paddy for amphibian and herbaceous fallow for reptile harboured the lowest species
richness. Reptile species diversity varied significantly between forest-derived and fallow-
derived vanilla agroforests whereas amphibian species showed no difference. For woody
fallow, herbaceous fallow and rice paddy, no unique species occurred in any of these land-
use types. Importantly, we found that amphibians and reptiles responded differently to land-
use history. After any kind of old-growth forest conversion, whether through burning or not,
amphibian species communities were decimated and showed a high shift in community
structure compared to old-growth forest. In reptile communities, the forest alteration through
slash-and-burn showed a strong species loss as well, but changes were less pronounced if
old-growth forest was transformed to forest fragment or forest-derived vanilla agroforest
since these transformations refrain from using fire.

4.1. Outstanding diversity of amphibians and reptiles in north-eastern Madagascar
The diversity of amphibians and reptiles documented here was exceptional, both within
Madagascar and compared to other tropical biodiversity hotspots. The few available studies
on the response of amphibian communities to land-use change in Madagascar reported
lower overall values, with 32 and 28 species, respectively (Vallan 2000; Ndriantsoa et al.
2017). Andreone et al. (2000) found 42 amphibian and 23 reptile species in north-eastern
Malagasy rainforest. Despite differences in methodology and study sites, our study provides
additional evidence that the north-east of Madagascar is one of the most species-rich regions
for herpetofauna, as previously suggested (Brown et al. 2016). The documented diversity
also exceeds values found in other tropical biodiversity hotspots (Cordier et al. 2021; Murray
and Nowakowski 2021) – for example, Urbina-Cardona et al. (2006) found 21 amphibian and
33 reptile species in region of Veracruz, Mexico, and Kurz et al. (2014) found 25 amphibian
and 20 reptile species in region of north-eastern Costa Rica (versus 58 amphibian and 61
reptile species in this study). The high species diversity may partly be driven by our extensive
sampling effort, in terms of number of plots (total n=80), diversity of different land-use types
(seven), and search effort (total of 270 min searching time by two observers per plot), and is
likely also influenced by the use of genetic samples for species identification.

4.2. Response of amphibian diversity to land-use change
We found a strong negative response of amphibian species richness to any anthropogenic
land uses. Old-growth forest had a significantly higher observed amphibian species richness
with 38% species exclusively occurring there. Besides, the amphibian species composition is
unique compared to all other land-use types, driven by a high share belonging to the
Microhylidae family and *Spinomantis*, *Guibemantis*, and *Mantella* genera (Fig. 4A). Only

forest fragments harboured comparably high values to old-growth forest for accumulated
amphibian species richness. Subsequently, we infer that amphibians are very sensitive to
habitat change. This result is in line with findings from Ndriantsoa et al. (2017) in disturbed
habitats of eastern Madagascar which documented strongly impoverished frog communities
in secondary vegetation and rice fields compared to forests. Forest fragments are thus
valuable for Malagasy amphibians to maintain diversity within the agricultural landscape, yet
forest fragments cannot substitute large and continuous old-growth forest (Vallan 2000).

The other land-use types (fallows that form part of the slush-and-burn cycle, vanilla
agroforests, and rice paddies) are of minor importance for amphibian conservation, given the
low species diversity that consists mainly of common and widely ranged species.
Nonetheless, amphibians could play an important functional role in these habitats:
abundance of amphibians is high throughout, reflected by high number of encounters,
particularly in rice paddies (see Fig. 3B and SI 9). As such, they may be an important food
source for other taxa or could provide pest control services in crops (Hocking et al. 2014).

The strong negative response of amphibians to deforestation could be explained by the fact
that numerous species rely on moist environments to avoid dehydration (Watling and Braga
2015), especially species living in evergreen forest habitat (Hof et al. 2011). Given that
selective logging, forest fragmentation and deforestation severely change microclimate
conditions (Ewers and Banks-Leite 2013), many species may struggle to cope. Furthermore,
amphibians have different habitat requirements throughout their life cycle stages (Hof et al.
2011; Hocking et al. 2014). Such heterogeneity is often lost in concert with the simplification
in forest structure (Wanger et al. 2009), making new habitats unsuitable for many species.

4.3. Response of reptile diversity to land-use change
We found a strong effect of land-use history on reptile diversity. Unburned land-use types
had a significantly higher average species richness, more uneven communities, more unique
species, and a distinct species composition compared to burned land-use types and rice
paddy. The distinct species composition in unburned land-use types was caused by an
increased abundance compared to other land-use types of species belonging to the
Chamaeleonidae family and *Uroplatus* genus (Fig. 4B). This is, to our knowledge, the first
study shedding light on the response of reptiles to land-use change in Madagascar's humid
eastern landscape. Studies from the drier southern region (Scott et al. 2006; Gardner et al.
2016; Nopper et al. 2018) show that Malagasy reptile species react less strongly to habitat
change than other taxa, especially if diurnal (Nopper et al. 2018).

Interestingly, the less pronounced effects of land-use change on reptiles compared to
amphibians in tropical landscapes has also been demonstrated by others (Wanger 2010;
Kurz et al. 2014) and was found in a review by Palacios et al. (2013). This may be due to the

high thermotolerance in reptiles (Thompson and Donnelly 2018) thanks to a fairly impermeable skin covered with scales. This structure prevents excessive evaporative water loss and adapts reptiles to various microclimates (Watling and Braga 2015). Furthermore, egg development is not heavily impacted by temperature rise, which reduces the size of hatchlings and accelerates incubation (Packard et al. 1982). Reptile lifestyle and traits may thus enable reptiles to adapt to more open, hotter, and drier environments (Morin 2005), making them less vulnerable to land-use change.

4.4. Land-use history of vanilla agroforests affects reptiles but not amphibians

In our study, we have separated forest-derived vanilla from fallow-derived vanilla agroforests, thereby explicitly accounting for land-use history (Martin et al. 2020). For amphibians, we found no differences between the two agroforestry systems across metrics. Reptile communities, on the other hand, were significantly more diverse in forest-derived vanilla agroforests on plot level, more species-rich overall, more uneven, and compositionally different compared to fallow-derived vanilla. Reptile communities observed in forest-derived vanilla agroforests are thus more similar to old-growth forest and forest fragment. The reptile communities recorded in fallow-derived vanilla agroforests were comparable to fallow land. Both kinds of agroforest thus resembled the land-use types they were derived from. An increase in accumulated amphibian species richness is also demonstrated from fallow-derived vanilla agroforest over fallow land, which may emphasize the rehabilitation opportunity of fallow land through agroforestry (Osen et al. 2021). These findings are in line with the prediction from a recent study of Martin et al. (2020) which found that the land-use history of agroforestry systems matters for biodiversity.

We further hypothesize, that the strong importance of forest-derived vanilla agroforests for reptiles may be in part driven by leaf-litter depth, which is known to positively influence reptile diversity and abundance (Urbina-Cardona et al. 2006). According to the same study, these microenvironmental variables also showed positive effects on amphibians and thus cannot explain why there are no differences between the two kinds of agroforests for amphibians. Elsewhere, the loss of canopy cover had negative impacts on amphibian and reptile species richness (Scott et al. 2006). Since canopy cover differed significantly between forest-derived vanilla agroforest (higher) and fallow-derived vanilla agroforest (lower) in the same study site (Osen et al. 2021), this is calling for further investigation combining habitat characteristics with species traits (Oliveira et al. 2017) to elucidate the drivers of change in both species groups.

4.5. Conservation implications

The strong negative response of amphibians to old-growth forest modification and the high old-growth forest dependency of amphibians and reptiles calls for the effective protection of

the last remaining old-growth forests. Additionally, conserving forest fragments within the agricultural landscape will be important to many reptile and amphibian species that are absent from other land-use types. This is particularly important given that numerous species are micro-endemics (Brown et al. 2014), meaning that they may only consist of few fragmented populations (Vieites et al. 2009). Management strategies are thus needed to safeguard the long-term existence of these fragments and to improve the connectivity between fragments. The protection of large continuous forests throughout the region is also important under a changing climate (Hof et al. 2011) and in light of emerging threats, such as the chytrid fungi (Kolby and Skerratt 2015) and the recent spread of the invasive Asian common toad (*Duttaphrynus melanostictus,* Soorae et al. 2020). These conservation needs are underscored by the exceptional diversity of reptiles and amphibians in north-eastern Madagascar as well as by the high proportion of endemic species.

Our findings from the north-eastern Madagascar also have broader relevance to herpetofauna conservation in other tropical biodiversity hotspots. The differential response of amphibians and reptiles to land-use change shows that conservationists should not treat herpetofauna as a homogenous group when devising conservation programs. Furthermore, the distinct differences in reptile communities between plots of contrasting land-use history shows that past land-use should also not be overlooked, suggesting that the maintenance of remnant forest fragments and forest-derived agroforests is important. Lastly, we highlight the importance of unburned compared to burned land-use types in the agricultural matrix for the conservation of reptiles.

6. Acknowledgements

We are grateful to all *chef de fokontany*, landowners, and Madagascar National Parks for granting us access to sites and information. We thank Prof. Miguel Vences and his laboratory staff for invaluable support with the DNA barcoding, Saskia Dröge for preparing panel C of Fig. 1, James Herrera for feedback, and Prof. Arne Mooers, Philippe Fernandez-Fournier, and an anonymous reviewer for their constructive reviews. We also thank 'CodonCode grants' for offering us a free two-year license under the CodonCode Aligner License Grant Program. We collected data under research permits N°100/17/MEEF/SG/DGF/DSAP/SCB.Re, N°163/17/MEEF/SG/DGF/DSAP/SCB.Re, N°18/18/MEEF/SG/DGF/DSAP/SCB.Re and N°254/18/MEEF/SG/DGF/DSAP/SCB.Re granted by the Ministry of Environment and Sustainable Development, Antananarivo. We transported DNA samples domestically under the transport permit N° 34/17-MEEF/SG/DREEF/SAVA/SRF granted by the Regional Office of the Ministry of Environment and Sustainable Development, Sambava. We exported the samples from Madagascar under CITES permits N°323C-EA05/MG18 and N°180C-EA03/MG19 granted by the Ministry of

Environment and Sustainable Development, Antananarivo, and imported the samples to
Germany under CITES permits DE-E-03377/18, DE-E-03378/18, DE-E-02422/19, DE-E-
02423/19, DE-E-02424/19, and DE-E-02425/19 granted by the German Federal Agency for
Nature Conservation, Bonn. This study was financially supported by the Niedersächsisches
Vorab of Volkswagen Foundation as part of the research project 'Diversity Turn in Land Use
Science' (Grant number 11-76251-99-35/13 (ZN3119)) and by the German Academic Ex-
Change Service (DAAD) within the 'Partnerships for Supporting Biodiversity in Developing
Countries' initiative (Project Nr. 57449386).

7. Conflict of interest

579 No conflict of interest to declare.

8. Author's contributions

581 TRF, DAM, KO, AW, HK, AA, and FMR designed the study. TRF, EB, RB, and RR collected
582 amphibian and reptile data under the lead of TRF. TRF and DAM analysed the data. TRF
583 and DAM wrote the first manuscript draft. All authors revised the manuscript.

9. Data availability

585 Data are available from Zenodo: http://doi.org/10.5281/zenodo.4548955

10. Reference

AmphibiaWeb. 2020. List of amphibian of Madagascar. Retrieve from:
https://amphibiaweb.org/cgi/amphib_query?rel-isocc=like&orderbyaw=Order&where-
isocc=Madagascar. Accessed December 14, 2020.

Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and D.J. Lipman. 1990. Basic local
alignement search tool. J. Mol. Biol. 215:403–410.

Anderson, M.J., and D.C.I. Walsh. 2013. PERMANOVA, ANOSIM, and the Mantel test in the
face of heterogeneous dispersions: What null hypothesis are you testing? Ecol. Monogr.
83:557–574. DOI: https://doi.org/10.1890/12-2010.1

Andreone, F., J.E. Randrianirina, P.D. Jenkins, and G. Aprea. 2000. Species diversity of
Amphibia, Reptilia and Lipotyphla (Mammalia) at Ambolokopatrika, a rainforest between
the Anjanaharibe-Sud and Marojejy massifs, NE Madagascar. Biodivers. Conserv.
9:1587–1622. DOI: https://doi.org/10.1023/A:1026559728808

Brown, J.L., N. Sillero, F. Glaw, P. Bora, D.R. Vieites, and M. Vences. 2016. Spatial
biodiversity patterns of Madagascar's amphibians and reptiles. PLoS One 11:1–26. DOI:
https://doi.org/10.1371/journal.pone.0144076

Brown, J.L., A. Cameron, A.D. Yoder, and M. Vences. 2014. A necessarily complex model to
explain the biogeography of the amphibians and reptiles of Madagascar. Nat. Commun.
5. DOI: https://doi.org/10.1038/ncomms6046

Bruford, M.W., Hanotte, O., Brookfield, J.F.Y. & Burke, T. 1992. Single-locus and multilocus
DNA fingerprinting. pp. 225-269. In: Hoelzel, A.R. 1 Ed. Molecular genetic analysis of
populations: A practical approach. Oxford: IRL Press, 325 p

Cacciali, P., E. Buongermini, and G. Köhler. 2019. Barcoding analysis of Paraguayan squamata. Diversity 11. DOI: https://doi.org/10.3390/d11090152

Chao, A., N.J. Gotelli, T.C. Hsieh, E.L. Sander, K.H. Ma, R.K. Colwell, and A.M. Ellison. 2014. Rarefaction and extrapolation with Hill numbers: A framework for sampling and estimation in species diversity studies. Ecol. Monogr. 84:45–67. DOI: https://doi.org/10.1890/13-0133.1

Chao, A., and L. Jost. 2012. Coverage-based rarefaction and extrapolation: Standardizing samples by completeness rather than size. Ecology 93:2533–2547. DOI: https://doi.org/10.1890/11-1952.1

Chao, A., and L. Jost. 2015. Estimating diversity and entropy profiles via discovery rates of new species. Methods Ecol. Evol. 6:873–882. DOI: https://doi.org/10.1111/2041-210X.12349

Cordier, J.M., R. Aguilar, J.N. Lescano, G.C. Leynaud, A. Bonino, D. Miloch, R. Loyola, and J. Nori. 2021. A global assessment of amphibian and reptile responses to land-use changes. Biol. Conserv. 253. DOI: https://doi.org/10.1016/j.biocon.2020.108863

Dröge, S., D.A. Martin, R. Andriafanomezantsoa, … H. Kreft. 2021. Listening to a changing landscape: Acoustic indices reflect bird species richness and plot-scale vegetation structure across different land-use types in north-eastern Madagascar. Ecol. Indic. 120:106929. DOI: https://doi.org/10.1016/j.ecolind.2020.106929

Ewers, R.M., and C. Banks-Leite. 2013. Fragmentation Impairs the Microclimate Buffering Effect of Tropical Forests. PLoS One 8. DOI: https://doi.org/10.1371/journal.pone.0058093

FAO. 2020. FAOSTAT. Food and Agriculture Organisation of the United Nations. http://www.fao.org/faostat/en/#home.

Gade, D.W. 1996. Deforestation and its effects in Highland Madagascar. Mt. Res. Dev. 16:101–116. DOI: https://doi.org/10.2307/3674005

Gardner, C.J., L.D. Jasper, C. Eonintsoa, J.J. Duchene, and Z.G. Davies. 2016. The impact of natural resource use on bird and reptile communities within multiple-use protected areas: evidence from sub-arid Southern Madagascar. Biodivers. Conserv. 25:1773–1793. DOI: https://doi.org/10.1007/s10531-016-1160-4

Gardner, T.A., J. Barlow, R. Chazdon, R.M. Ewers, C.A. Harvey, C.A. Peres, and N.S. Sodhi. 2009. Prospects for tropical forest biodiversity in a human-modified world. Ecol. Lett. 12:561–582. DOI: https://doi.org/10.1111/j.1461-0248.2009.01294.x

Glaw, F., and M. Vences. 2007. Ny Toro-hay momba ny Amphibia sy ny Reptilia an'i Madagasikara. 3rd editio. (M. Vences and V. Glaw, eds.). Cologne. 529 pp.

Gonzalez, T., and J.D. González-trujillo. 2021. Differential effects of fire on the occupancy of small mammals in neotropical savanna-gallery forests. DOI: https://doi.org/10.1016/j.pecon.2021.03.005

Goodman, S.M., and J.P. Benstead. 2005. Updated estimates of biotic diversity and endemism for Madagascar. Oryx 39:73–77. DOI: https://doi.org/10.1017/S0030605305000128

Gray, C.L., S.L.L. Hill, T. Newbold, … J.P.W. Scharlemann. 2016. Local biodiversity is higher inside than outside terrestrial protected areas worldwide. Nat. Commun. 7. DOI: https://doi.org/10.1038/ncomms12306

Hänke, H., J. Niens, K. Osen, V. Uruena, and S. Annette. 2018. Socio-economic, land use and value chain perspectives on vanilla farming in the SAVA Region (north-estern

Madagascar): The Diversity Turn Baseline Study (DTBS). Econstor 1–137.

Hansen, M.C., L. Wang, X.P. Song, A. Tyukavina, S. Turubanova, P. V. Potapov, and S. V.
Stehman. 2020. The fate of tropical forest fragments. Sci. Adv. 6:1–10. DOI:
https://doi.org/10.1126/sciadv.aax8574

Hocking, D.J., K.J. Babbitt, and D.J. Hocking. 2014. Amphibian contributions to ecosystem
services. Herpetol. Conserv. Biol. 9:1–17.

Hof, C., M.B. Araújo, W. Jetz, and C. Rahbek. 2011. Additive threats from pathogens, climate
and land-use change for global amphibian diversity. Nature 480:516–519. DOI:
https://doi.org/10.1038/nature10650

Hothorn, T., F. Bretz, and P. Westfall. 2008. Simultaneous inference in general parametric
models. Biometrical J. 50:346–363. DOI: https://doi.org/10.1002/bimj.200810425

Hsieh, T.C., K.H. Ma, and A. Chao. 2016. iNEXT: an R package for rarefaction and
extrapolation of species diversity (Hill numbers). Methods Ecol. Evol. 7:1451–1456.
DOI: https://doi.org/10.1111/2041-210X.12613

IUCN (2019). The IUCN red list of threatened species. Version 2019-1. IUCN, Gland,
Switzerland. Available from http://www.iucnredlist.org (accessed March 2019).

Irwin, M.T., P.C. Wright, C. Birkinshaw, … J.U. Ganzhorn. 2010. Patterns of species change
in anthropogenically disturbed forests of Madagascar. Biol. Conserv. 143:2351–2362.
DOI: https://doi.org/10.1016/j.biocon.2010.01.023

Kadlec, T., R. Tropek, and M. Konvicka. 2012. Timed surveys and transect walks as
comparable methods for monitoring butterflies in small plots. J. Insect Conserv. 16:275–
280. DOI: https://doi.org/10.1007/s10841-011-9414-7

Kolby, J.E., and L.F. Skerratt. 2015. Amphibian chytrid fungus in Madagascar neither shows
widespread presence nor signs of certain establishment. PLoS One 10:1–6. DOI:
https://doi.org/10.1371/journal.pone.0139172

Kremen, C., and A.M. Merenlender. 2018. Landscapes that work for biodiversity and people.
Science (80-.). 362. DOI: https://doi.org/10.1126/science.aau6020

Kumar, S., G. Stecher, M. Li, C. Knyaz, and K. Tamura. 2018. MEGA X: Molecular
evolutionary genetics analysis across computing platforms. Mol. Biol. Evol. 35:1547–
1549. DOI: https://doi.org/10.1093/molbev/msy096

Kurz, D.J., A.J. Nowakowski, M.W. Tingley, M.A. Donnelly, and D.S. Wilcove. 2014. Forest-
land use complementarity modifies community structure of a tropical herpetofauna. Biol.
Conserv. 170:246–255. DOI: https://doi.org/10.1016/j.biocon.2013.12.027

Li, D., J.D. Olden, J.L. Lockwood, S. Record, M.L. McKinney, and B. Baiser. 2020. Changes
in taxonomic and phylogenetic diversity in the Anthropocene: Changes in biodiversity.
Proc.   R.   Soc.   B   Biol.   Sci.   287.   DOI:
https://doi.org/10.1098/rspb.2020.0777rspb20200777

Llopis, J.C., P.C. Harimalala, R. Bär, A. Heinimann, Z.H. Rabemananjara, and J.G.
Zaehringer. 2019. Effects of protected area establishment and cash crop price dynamics
on land use transitions 1990–2017 in north-eastern Madagascar. J. Land Use Sci.
14:52–80. DOI: https://doi.org/10.1080/1747423X.2019.1625979

Martin, D.A., R. Andriafanomezantsoa, S. Dröge, K. Osen, E. Rakotomalala, A. Wurz, A.
Andrianarimisa, and H. Kreft. 2021. Bird diversity and endemism along a land-use
gradient in Madagascar: The conservation value of vanilla agroforests. Biotropica
53:179–190. DOI: https://doi.org/10.1111/btp.12859

Martin, D.A., K. Osen, I. Grass, D. Hölscher, T. Tscharntke, A. Wurz, and H. Kreft. 2020.

Land-use history determines ecosystem services and conservation value in tropical
agroforestry. Conserv. Lett. 13:1–12. DOI: https://doi.org/10.1111/conl.12740

Martinez Arbizu, P. pairwise.Adonis: Pairwise Multilevel Comparison Using Adonis. R
Package Version 0.0.1. Available online:
https://github.com/pmartinezarbizu/pairwiseAdonis (accessed on 6 December 2020).

Matuoka, M.A., M. Benchimol, J.M. de Almeida-Rocha, and J.C. Morante-Filho. 2020. Effects
of anthropogenic disturbances on bird functional diversity: A global meta-analysis. Ecol.
Indic. 116:106471. DOI: https://doi.org/10.1016/j.ecolind.2020.106471

Mendenhall, C.D., A. Shields-Estrada, A.J. Krishnaswami, and G.C. Daily. 2016. Quantifying
and sustaining biodiversity in tropical agricultural landscapes. Proc. Natl. Acad. Sci. U.
S. A. 113:14544–14551. DOI: https://doi.org/10.1073/pnas.1604981113

Morin, K.C. 2005. Herpetofaunal responses to prescibed fire in upland pine communities of
Northeast Florida. PhD Thesis, University of Florida. 77 pp.

Murray, A.H., and A.J. Nowakowski. 2021. Climate and land-use change severity alter trait-
based responses to habitat conversion. DOI: https://doi.org/10.1111/geb.13237

Myers, N., R.A. Mittermeler, C.G. Mittermeler, G.A.B. Da Fonseca, and J. Kent. 2000.
Biodiversity hotspots for conservation priorities. Nature 403:853–858. DOI:
https://doi.org/10.1038/35002501

Ndriantsoa, S.H., J.C. Riemann, N. Raminosoa, M.O. Rödel, and J.S. Glos. 2017. Amphibian
diversity in the matrix of a fragmented landscape around Ranomafana in Madagascar
depends on matrix quality. Trop. Conserv. Sci. 10. DOI:
https://doi.org/10.1177/1940082916686065

Newbold, T., L.N. Hudson, S.L.L. Hill, … A. Purvis. 2016. Global patterns of terrestrial
assemblage turnover within and among land uses. Ecography (Cop.). 39:1151–1163.
DOI: https://doi.org/10.1111/ecog.01932

Newbold, T., L.N. Hudson, H.R.P. Phillips, … A. Purvis. 2014. A global model of the
response of tropical and sub-tropical forest biodiversity to anthropogenic pressures.
Proc. R. Soc. B Biol. Sci. 281. DOI: https://doi.org/10.1098/rspb.2014.1371

Newbold, T., P. Oppenheimer, A. Etard, and J.J. Williams. 2020. Tropical and Mediterranean
biodiversity is disproportionately sensitive to land-use and climate change. Nat. Ecol.
Evol. 4:1630–1638. DOI: https://doi.org/10.1038/s41559-020-01303-0

Nopper, J., J.C. Riemann, K. Brinkmann, M.O. Rödel, and J.U. Ganzhorn. 2018. Differences
in land cover – biodiversity relationships complicate the assignment of conservation
values in human-used landscapes. Ecol. Indic. 90:112–119. DOI:
https://doi.org/10.1016/j.ecolind.2018.02.004

Nori, J., F. Villalobos, and R. Loyola. 2018. Global priority areas for amphibian research. J.
Biogeogr. 45:2588–2594. DOI: https://doi.org/10.1111/jbi.13435

Oksanen, J., F.G. Blanchet, M. Friendly, … E. Szoecs. 2020. Community Ecology Package.
1–299.

Oliveira, B.F., V.A. São-Pedro, G. Santos-Barrera, C. Penone, and G.C. Costa. 2017.
AmphiBIO, a global database for amphibian ecological traits. Sci. Data 4:1–7. DOI:
https://doi.org/10.1038/sdata.2017.123

Osen, K., M.R. Soazafy, D.A. Martin, A. Wurz, A. März, H.L.T. Ranarijaona, and D. Hölscher.
2021. Land-use history determines stand structure and tree diversity in vanilla
agroforests of northeastern Madagascar. Appl. Veg. Sci. 1–14. DOI:
https://doi.org/10.1111/avsc.12563

Packard, M.J., G.C. Packard, and T.J. Boardman. 1982. Structure of Eggshells and Water
Relations of Reptilian Eggs. 38. 136–155 pp.

Palacios, C.P., B. Agüero, and J.A. Simonetti. 2013. Agroforestry systems as habitat for
herpetofauna: Is there supporting evidence? Agrofor. Syst. 87:517–523. DOI:
https://doi.org/10.1007/s10457-012-9571-z

De Palma, A., M. Kuhlmann, S.P.M. Roberts, S.G. Potts, L. Börger, L.N. Hudson, I. Lysenko,
T. Newbold, and A. Purvis. 2015. Ecological traits affect the sensitivity of bees to land-
use pressures in European agricultural landscapes. J. Appl. Ecol. 52:1567–1577. DOI:
https://doi.org/10.1111/1365-2664.12524

Palumbi, S., S. Romano, W.O. Mcmillan, and G. Grabowski. 1991. the Simple Fool ' S Guide
To Pcr. October 96822:1–45.

Perfecto, I., and J. Vandermeer. 2010. The agroecological matrix as alternative to the land-
sparing/agriculture intensification model. Proc. Natl. Acad. Sci. U. S. A. 107:5786–5791.
DOI: https://doi.org/10.1073/pnas.0905455107

Powers, J.S., M.D. Corre, T.E. Twine, and E. Veldkamp. 2011. Geographic bias of field
observations of soil carbon stocks with tropical land-Use changes precludes spatial
extrapolation. Proc. Natl. Acad. Sci. U. S. A. 108:6318–6322. DOI:
https://doi.org/10.1073/pnas.1016774108

Powers, R.P., and W. Jetz. 2019. Global habitat loss and extinction risk of terrestrial
vertebrates under future land-use-change scenarios. Nat. Clim. Chang. 9:323–329. DOI:
https://doi.org/10.1038/s41558-019-0406-z

R Core Team.2019. R: A language and environment for statistical computing. **R** Foundation
for Statistical Computing, Vienna, Austria.

Rakotoarison, A., M.D. Scherz, F. Glaw, … M. Gonçalo. 2017. Describing the smaller
majority: Integrative taxonomy reveals twenty-six new species of tiny microhylid frogs (
genus Stumpffia) from Madagascar Describing the smaller majority: integrative
taxonomy reveals twenty-six new species of tiny microhylid fro.

Ratsoavina, F.M., A.P. Raselimanana, M.D. Scherz, A. Rakotoarison, J.H. Razafindraibe, F.
Glaw, and M. Vences. 2019. Finaritra! A splendid new leaf-tailed gecko (Uroplatus)
species from Marojejy National Park in north-eastern Madagascar. Zootaxa 4545:563–
577. DOI: https://doi.org/10.11646/zootaxa.4545.4.7

Rembold, K., H. Mangopo, S.S. Tjitrosoedirdjo, and H. Kreft. 2017. Plant diversity, forest
dependency, and alien plant invasions in tropical agricultural landscapes. Biol. Conserv.
213:234–242. DOI: https://doi.org/10.1016/j.biocon.2017.07.020

Ricciardi, V., Z. Mehrabi, H. Wittman, D. James, and N. Ramankutty. 2021. Higher yields and
more biodiversity on smaller farms. Nat. Sustain. 1–7. DOI:
https://doi.org/10.1038/s41893-021-00699-2

Roswell, M., J. Dushoff, and R. Winfree. 2021. A conceptual guide to measuring species
diversity. Oikos 130:321–338. DOI: https://doi.org/10.1111/oik.07202

Scales, B.R., and S.J. Marsden. 2008. Biodiversity in small-scale tropical agroforests: A
review of species richness and abundance shifts and the factors influencing them.
Environ. Conserv. 35:160–172. DOI: https://doi.org/10.1017/S0376892908004840

Scott, D.M., D. Brown, S. Mahood, B. Denton, A. Silburn, and F. Rakotondraparany. 2006.
The impacts of forest clearance on lizard, small mammal and bird communities in the
arid spiny forest, southern Madagascar. Biol. Conserv. 127:72–87. DOI:
https://doi.org/10.1016/j.biocon.2005.07.014

Soorae, P.S., G.J. Frankham, and A.A. Mohamed. 2020. The first record of the asian
common toad Duttaphrynus melanostictus schneider, 1799 in Abu Dhabi, United Arab
Emirates. BioInvasions Rec. 9:434–443. DOI: https://doi.org/10.3391/bir.2020.9.2.29

Styger, E., H.M. Rakotondramasy, M.J. Pfeffer, E.C.M. Fernandes, and D.M. Bates. 2007.
Influence of slash-and-burn farming practices on fallow succession and land
degradation in the rainforest region of Madagascar. Agric. Ecosyst. Environ. 119:257–
269. DOI: https://doi.org/10.1016/j.agee.2006.07.012

Thompson, M.E., and M.A. Donnelly. 2018. Effects of Secondary Forest Succession on
Amphibians and Reptiles: A Review and Meta-analysis. Copeia 106:10–19. DOI:
https://doi.org/10.1643/ch-17-654

Tilman, D., C. Balzer, J. Hill, and B.L. Befort. 2011. Global food demand and the sustainable
intensification of agriculture. Proc. Natl. Acad. Sci. U. S. A. 108:20260–20264. DOI:
https://doi.org/10.1073/pnas.1116437108

Tscharntke, T., Y. Clough, T.C. Wanger, L. Jackson, I. Motzke, I. Perfecto, J. Vandermeer,
and A. Whitbread. 2012. Global food security, biodiversity conservation and the future of
agricultural intensification. Biol. Conserv. 151:53–59. DOI:
https://doi.org/10.1016/j.biocon.2012.01.068

Uetz, P., Freed, P, Aguilar, R. & Hošek, J. (eds.) (2021) The Reptile Database,
http://www.reptile-database.org, accessed [August 25, 2021]

Urbina-Cardona, J.N., M. Olivares-Pérez, and V.H. Reynoso. 2006. Herpetofauna diversity
and microenvironment correlates across a pasture-edge-interior ecotone in tropical
rainforest fragments in the Los Tuxtlas Biosphere Reserve of Veracruz, Mexico. Biol.
Conserv. 132:61–75. DOI: https://doi.org/10.1016/j.biocon.2006.03.014

Vallan, D. 2000. Influence of forest fragmentation on amphibian diversity in the nature
reserve of Ambohitantely, highland Madagascar. Biol. Conserv. 96:31–43. DOI:
https://doi.org/10.1016/S0006-3207(00)00041-0

Vences, M., M. Thomas, R.M. Bonett, and D.R. Vieites. 2005. Deciphering amphibian
diversity through DNA barcoding: Chances and challenges. Philos. Trans. R. Soc. B
Biol. Sci. 360:1859–1868. DOI: https://doi.org/10.1098/rstb.2005.1717

Vieilledent, G., C. Grinand, F.A. Rakotomalala, R. Ranaivosoa, J.R. Rakotoarijaona, T.F.
Allnutt, and F. Achard. 2018. Combining global tree cover loss data with historical
national forest cover maps to look at six decades of deforestation and forest
fragmentation in Madagascar. Biol. Conserv. 222:189–197. DOI:
https://doi.org/10.1016/j.biocon.2018.04.008

Vieites, D.R., K.C. Wollenberg, F. Andreone, J. Köhler, F. Glaw, and M. Vences. 2009. Vast
underestimation of Madagascar's biodiversity evidenced by an integrative amphibian
inventory. Proc. Natl. Acad. Sci. U. S. A. 106:8267–8272. DOI:
https://doi.org/10.1073/pnas.0810821106

Wanger, T.C. 2010. Land-use change , tropical biodiversity , and ecosystem services –
Southeast Asian amphibians and reptiles in focus. Pp. 1–147 in PhD Thesis University
of Adelaide.

Wanger, T.C., A. Saro, D.T. Iskandar, B.W. Brook, N.S. Sodhi, Y. Clough, and T. Tscharntke.
2009. Conservation value of cacao agroforestry for amphibians and reptiles in South-
East Asia: Combining correlative models with follow-up field experiments. J. Appl. Ecol.
46:823–832. DOI: https://doi.org/10.1111/j.1365-2664.2009.01663.x

Watling, J.I., and L. Braga. 2015. Desiccation resistance explains amphibian distributions in a
fragmented tropical forest landscape. Landsc. Ecol. 30:1449–1459. DOI:
https://doi.org/10.1007/s10980-015-0198-0

Whitehurst, A.S., J.O. Sexton, and L. Dollar. 2009. Land cover change in western
Madagascar's dry deciduous forests: a comparison of forest changes in and around
Kirindy Mite National Park. Oryx 43:275. DOI:
https://doi.org/10.1017/S0030605309001756

Wickham, H. 2016. Programming with ggplot2. 241–253. DOI: https://doi.org/10.1007/978-3-
319-24277-4_12

Williams, J.J., and T. Newbold. 2020. Local climatic changes affect biodiversity responses to
land use: A review. Divers. Distrib. 26:76–92. DOI: https://doi.org/10.1111/ddi.12999

Figure. 1 Overview of the study region and the study design. **A** SAVA region in north-eastern
Madagascar; **B** study area within SAVA region; **C** study design showing the distribution of 80
plots in ten villages and at two sites inside Marojejy National Park; and **D** overview of the
studied land-use types and the typical land-use transformation trajectory from old-growth
forest to forest fragments and agricultural land-uses. Panel D modified after Martin et al.
(2021) and Panel C modified after Dröge et al. (2021). Rice paddy is not part of the main
land-use trajectory.

Figure. 2 Plot-level observed amphibian (A) and reptile (B) species richness across seven
land-use types (replicated 10 times each except fallow-derived vanilla, which is replicated 20
times) in north-eastern Madagascar. The dots represent the species richness per plot in each
land-use type. The black horizontal line in the box shows the median. Land-use types with
letters in common did not differ significantly based on pairwise comparisons that controlled
for inflated false-positive errors using the Tukey HSD approach (Numeric results in SI:
amphibians & reptiles SI 2). See icon reference in supplementary information (Free Icons
Library).

865
Figure. 3: Encounter-based accumulation curves showing interpolated (solid line) and extrapolated (dotted line) diversities for amphibians (first and

second column of panel from left to right) and reptiles (third and fourth panel) in north-eastern Madagascar. Unburned (first and third panel) and burned

(second and fourth panel) land-use types are separated. The species richness represented by q = 0 (first row of panel from above), Shannon diversity,

q=1 (second row of panel) and Simpson diversity q=2 (third row of panel) with 95% confidence intervals (shaded areas) for the amphibian and the reptile

data of seven land-use types. The solid dots, triangles and diamonds represent the reference samples, i.e. the observed number of encounters and

species   richness.   See   icon   reference   in   supplementary   information   (Free   Icons   Library).

Figure. 4: Species composition across seven land-use types in north-eastern Madagascar. All 58 amphibian species with 3694 encounters (A) and
all 61 reptile species (B) with 2521 encounters are displayed by forest dependency rank. 38% of amphibian species and 26% of reptile species
exclusively occurred in old-growth forest, despite that old-growth forest is only accounting for 12.5% of the total plots. See icon reference in
supplementary information (Free Icons Library).

Figure. 5: Species composition across seven land-use types in north-eastern Madagascar.
Non-metric dimensional scaling (NMDS) showing community dissimilarity of amphibian (A)
and reptile (B) communities. Old growth forest shows unique species composition for both
amphibians and reptiles See icon reference in supplementary information (Free Icons
Library).

Table 1: Estimated and observed amphibian and reptile species richness (q = 0) for all land-
use types and separated per land-use type showing the observed and extrapolated total
species richness. Each land-use type is represented by 10 plots; for fallow-derived vanilla,
the 10 plots are down-sampled from 20 plots. Extrapolated species richness is based on
5000 encounters and includes the lower and upper 95% confidence interval in italics below
(into bracket). See SI for results of q = 1 and q =2 (amphibians and reptiles: table SI 3).

Land- use type / Species Group (measure)	All 80 plots	Old-growth forest	Forest fragment	Forest-derived Vanilla agroforest	Fallow-derived Vanilla agroforest	Woody fallow	Herbaceous fallow	Rice paddy
Amphibian (observed)	58	32	26	14	16	8	6	4
Amphibian (extrapolated)	NA	60.0 *(20.4 - 99.6)*	46.1 *(12.4 - 79.8)*	18.0 *(9.1 - 26.9)*	25.0 *(7.0 - 42.9)*	9.0 *(7.4 - 10.6)*	6.5 *(3.9 - 9.1)*	5.0 *(4.3 - 5.6)*
Reptile (observed)	61	34	30	30	18	19	8	11
Reptile (extrapolated)	NA	42.0 *(20.4 - 63.5)*	40.1 *(10.2 - 69.9)*	35.1 *(18.2 - 52.4)*	26.9 *(8.2 - 45.7)*	46.6 *(11.5 - 81.7)*	8.0 *(6.8 - 9.2)*	18.9 *(4.3 - 33.4)*
Total (observed)	119	66	56	44	34	27	14	15

890