Long-term patterns of an interconnected core marine microbiota

Anders K. Krabberød[1*], Ina M. Deutschmann[2], Marit F. M. Bjorbækmo[1], Vanessa Balagué[2], Caterina R. Giner[2], Isabel Ferrera[2,3], Esther Garcés[2], Ramon Massana[2], Josep M. Gasol[2, 4], Ramiro Logares[2,1*]

[1] University of Oslo, Department of Biosciences, Section for Genetics and Evolutionary Biology (Evogene), Blindernv. 31, N-0316 Oslo, Norway

[2] Institute of Marine Sciences (ICM), CSIC, Passeig Marítim de la Barceloneta, Barcelona, Spain

[3] Centro Oceanográfico de Málaga, Instituto Español de Oceanografía, 29640 Fuengirola, Málaga, Spain

[4] Centre for Marine Ecosystems Research, School of Sciences, Edith Cowan University, Joondalup, WA, Australia

Corresponding authors:

*Anders K. Krabberød, University of Oslo, Department of Biosciences, Section for Genetics and Evolutionary Biology (Evogene), Blindernv. 31, N-0316 Oslo, Norway
Email: a.k.krabberod@ibv.uio.no

*Ramiro Logares, Institute of Marine Sciences (ICM), CSIC, Passeig Marítim de la Barceloneta, Barcelona, Spain
Email: ramiro.logares@icm.csic.es

Running title: Patterns in a core marine microbiota

Manuscript for: Microbiome

ABSTRACT

Background

Ocean microbes constitute ~70% of the marine biomass, are responsible for ~50% of the Earth's primary production, and are crucial for global biogeochemical cycles. Marine microbiotas include core taxa that are usually key for ecosystem function. Despite their importance, core marine microbes are relatively unknown, which reflects the lack of consensus on how to identify them. So far, most core microbiotas have been defined based on species occurrence and abundance. Yet, species interactions are also important to identify core microbes, as communities include interacting species. Here, we investigate interconnected bacteria and small protists of the core pelagic microbiota populating a long-term marine-coastal observatory in the Mediterranean Sea over a decade.

Results

Core microbes were defined as those present in >30% of the monthly samples over 10 years, with the strongest associations. The core microbiota included 259 Operational Taxonomic Units (OTUs) including 182 bacteria, 77 protists, and 1,411 strong and mostly positive (~95%) associations. Core bacteria tended to be associated with other bacteria, while core protists tended to be associated with bacteria. The richness and abundance of core OTUs varied annually, decreasing in stratified warmers waters and increasing in colder mixed waters. Most core OTUs had a preference for one season, mostly winter, which featured subnetworks with the highest connectivity. Groups of highly associated taxa tended to include protists and bacteria with predominance in the same season, particularly winter. A group of 13 highly-connected hub-OTUs, with potentially important ecological roles dominated in winter and spring. Similarly, 18 connector OTUs with a low degree but high centrality were mostly associated with summer or autumn and may represent transitions between seasonal communities.

Conclusions

We found a relatively small and dynamic interconnected core microbiota in a model temperate marine-coastal site, with potential interactions being more deterministic in winter than in other seasons. These core microbes would be essential for the functioning of this ecosystem over the year. Other non-core taxa may also carry out important functions but would be redundant and non-essential. Our work contributes toth e understanding of the dynamics and potential interactions of core microbes possibly sustaining ocean ecosystem function.

Keywords: bacteria, protists, ocean, time-series, seasonality, networks, associations

BACKGROUND

Ecosystems are composed of interacting units embedded in and influenced by their physicochemical environment. Ecosystem function can be broadly defined as the biological, geochemical, and physical processes that occur within it. These processes will likely change or halt if specific organisms or gene-functions are removed, driving the ecosystem towards a new state or its collapse. It is hypothesized that ecological redundancy guarantees continuous ecosystem function, as multiple species could carry out the same or similar function [1]. And while the amount of functional redundancy in microbial ecosystems is a matter of debate [2, 3] it has also been observed that microbiotas in comparable habitats tend to share "core" species that are hypothesized to be fundamental for ecosystem function [4]. These core organisms and the functions they carry out might not be easily replaced.

Identifying the core microbiota is not straightforward as there are different ways of defining a core depending on the habitats and the questions being addressed [4]. One often-used approach is to identify species that tend to be recurrently present across spatiotemporal scales. This definition might not be sufficient, however, since communities are made up of interacting species [5]. A more appropriate definition of a core, therefore, needs to incorporate ecological interactions fundamental for the community in the location under study [4, 5]. This is particularly important in studies using DNA to investigate microbial communities, as a fraction of the detected taxa could be dormant, dead, or transient [6-8]. In the interaction-based definition taxa that do not appear to be interacting are excluded from the core [4].

Core microbiotas based on common presence have been widely studied in terrestrial animals, in particular humans [9] or cattle [10], as well in marine animals, in particular corals [11, 12] and sponges [13, 14]. Core microbiotas in non-host-associated

systems, such as soils or the ocean, have been investigated to a lesser extent. In soils,

for example, a global analysis identified a core group of 241 ubiquitous and dominant

bacterial taxa with more or less invariant abundances and unclear habitat preferences

[15]. In the tropical and subtropical global-ocean, a total of 68 bacteria and 57

picoeukaryotic operational taxonomic units (OTUs) have been identified that could be

part of the core surface microbiota, as they were present in >80% of the globally-

distributed samples [16].

Analyses of ocean time-series have also pointed to the existence of core

microbiotas. For example, Gilbert et al. [17] investigated the microbiota of the English

Channel for 6 years and found 12 abundant OTUs that were detected throughout the

entire dataset (72 time-points), totaling ~35% of the sequence abundance. Potentially

core bacterial OTUs were detected in the SPOT time-series (southern California), in a

study covering 10 years of monthly samples in the euphotic zone [18]. These

potentially-core bacterial OTUs were present in >75% of the months, represented ~7%

(25-28 OTUs depending on depth) of the total richness, and had a high (>10%) relative

abundance [18].

These studies have provided substantial insights on core marine microbiotas,

although they typically define them in terms of species occurrence or abundance over

spatiotemporal scales, rather than on potential interactions. As in other ecosystems,

microbial interactions are essential for the functioning of the ocean ecosystem, where

they guarantee the transfer of carbon and energy to upper trophic levels, as well as the

recycling of carbon and nutrients [19]. Despite their importance, most microbial

interactions in the ocean remain unknown [20]. A recent literature survey spanning the

last 150 years indicated that we have documented a minor fraction of protist interactions

in the ocean [21] and most likely, the same is true if not worse for bacteria.

During the last decade, association networks have been used to bridge this knowledge gap. Association networks are based on correlations between species' abundances and they may reflect microbial interactions [22]. Contemporaneous positive correlations may point to interactions such as symbiosis, or similar niche preferences, while negative correlations may suggest predation, competition, or opposite niche preferences [23]. So far, network analyses have produced hypotheses on microbial interactions at the level of individual species across diverse ecosystems [22, 24, 25], a few of which have been experimentally validated [26]. In addition, networks can help detect species that have relatively more associations to other species ("hubs"), or species that connect different subgroups within a network, and which therefore may have important roles in the ecosystem. Groups of highly associated species in the network ("modules") may represent niches [27, 28], and the amount of these modules may increase with increasing environmental selection [22]. Networks can also produce ecological insight at the community level, since their architecture can reflect community processes, such as selection [27].

Network analyses have been particularly useful for the investigation of microbial interactions in the ocean [25, 29]. A surface global-ocean network analysis of prokaryotes and single-celled eukaryotes indicated that ~72% of the associations between microbes were positive and that most associations were between single-celled eukaryotes belonging to different organismal size-fractions [26]. Other studies using networks have indicated a limited number of associations between marine microbes and abiotic environmental variables [17, 18, 23, 26, 30-32], suggesting that microbial interactions have an important role in driving community turnover [32]. Despite the important insights these studies have provided, most of them share the limitation that

they do not disentangle whether microbial associations may represent ecological

interactions or environmental preferences [22].

Even though association networks based on long-term species dynamics may

allow a more accurate delineation of core marine microbiotas, few studies have

identified them in this manner. Consequently, we have a limited understanding of the

interconnected set of organisms that may be key for ocean ecosystem function. Here

we identify and investigate the core microbiota occurring in the marine-coastal Blanes

Bay Microbial Observatory (Northwestern Mediterranean Sea) over 10 years. We

delineated the core microbiota stringently, using potential interactions based on species

abundances. We also made an effort to disentangle environmental effects in association

networks by identifying and removing species associations that are a consequence of

shared environmental preference and not interactions between the species [33]. We

analyzed bacteria and protists from the pico- (0.2-3 μm) and nanoplankton (3-20 μm)

organismal size fractions, which show a strong seasonality in this location [34-36].

Taxa relative abundances were estimated by sequencing the 16S and 18S rRNA-gene

and delineating OTUs as Amplicon Sequence Variants (ASVs). Specifically, we ask:

What taxa constitute the interconnected core microbiota and what are the main patterns

of this assemblage over 10 years? Does the core microbiota feature seasonal sub-groups

of highly associated species? What degree of association do bacteria and microbial

eukaryotes have and do they show comparable connectivity? Can we identify core

OTUs with central positions in the network that could have important ecological roles?

RESULTS

Composition and dynamics of the resident microbiota

Based on the data set containing 2,926 OTUs, (1,561 bacteria and 1,365 microbial eukaryotes) we first defined the resident OTUs as the bacteria and microbial eukaryotes present in >30% of the samples, which equals 36 out of 120 months (not necessarily consecutive). This threshold was selected as it includes seasonal OTUs that would be present recurrently in at least one season. The residents consisted of 709 OTUs: 354 Bacteria (~54% relative read abundance) and 355 Eukaryotic OTUs (~46% relative read abundance) [**Table 1,** see methods for calculation of relative read abundance]. The most abundant resident bacteria OTUs belonged to Oxyphotobacteria (mostly *Synechococcus*; ~15% of total relative read abundance), Alphaproteobacteria (mostly SAR11 Clade Ia [~9%, and clade II [~4%]), and Gammaproteobacteria (mainly SAR86; ~2%). The most abundant resident protist OTUs belonged to Dinophyceae (predominantly an unclassified dinoflagellate lineage [~7%], Syndiniales Group I Clade 1 [~7%] and *Gyrodinium* [~4%]), Chlorophyta (mostly *Micromonas* [~3%] and *Bathycoccus* [~2%]), Ochrophyta (predominantly Mediophyceae [~2%] and *Chaetoceros* [~1%]) and Cryptophyceae (mainly a Cryptomonadales lineage [~2%]) [**Figure 3, Table S1**].

Table 1. Description of the datasets

	OTUs	OTUs (%)	Sequence abundance (%)[*]
All OTUs[1]	2,926	100	*100*
Bacteria	1,561	53.3	*50.7*
Protists	1,365	46.7	*49.3*
Resident microbiota[2]	709	100	100 (*85*)
Bacteria	354	49.9	53.6
Protists	355	50.1	46.4
Core microbiota[3]	**259**	**100**	**64.5** (*54*)
Bacteria	182	70.3	46.3
Protists	77	29.7	18.2
Picoplankton	109	42.1	32.4
Nanoplankton	150	57.9	32.1
Protists			
Heterotroph	5	1.9	0.3
Photoautotroph	37	14.3	11.8

Parasite	21	8.1	3.5
Mixotroph	3	1.2	0.7
Symbiont	1	0.4	0.1
Unknown	11	4.3	2.0
Bacteria			
Photoautotroph (cyanobacteria)	19	7.3	19.3
Non-photoautotroph[4]	163	62.5	26.8
Seasonal preference core OTUs			
Winter	156	60.2	21.8
Spring	24	9.3	16.4
Summer	44	17.0	8.2
Autumn	30	11.6	13.7
No seasonality	5	1.9	4.5
Seasonal subnetworks			
Winter	156	60.2	21.8
Spring	19	7.3	13.7
Summer	41	15.8	6.6
Autumn	26	10.0	12.9

[1] Number of OTUs in the full dataset that were left after quality control and rarefaction, which were present in at least 10% of the samples (i.e. 12 months, not necessarily consecutive).

[2] OTUs present in at least 30% of the samples (i.e. 36 months, not necessarily consecutive) [=Resident microbiota].

[3] OTUs included in the core network (core microbiota) with significant correlations (p&q <0.001), local similarity scores >|0.7| and Spearman correlations >|0.7|, being present in at least 30% of the samples.

[4] Includes non-photoautotrophic lifestyles (i.e., chemoautotrophs, photoheterotrophs, chemoheterotrophs, etc.).

* In *Italics* the abundances relative to all OTUs are indicated. All other values in normal text indicate abundances relative to OTUs in the resident microbiota.

The resident microbiota, including both protists and bacteria, showed seasonal variation over 10 years, with communities from the same season but different years tending to group (**Figure 1C and D**). The structure of the resident microbiota correlated to specific environmental variables during winter (nutrients, Total photosynthetic nanoflagellates [PNF; 2-5µm size], and small PNF [2µm]), spring (Total Chlorophyll a [Chla]), summer (daylength, temperature, Secchi disk depth and, the cell abundances of *Synechococcus,* Heterotrophic prokaryotes [HP] and Heterotrophic nanoflagellates [HNF, 2-5µm]) and autumn (salinity) [**Figure 1C**]. The environmental variables most relevant for explaining the variance of the resident microbiota were determined by stepwise model selection and distance-based redundancy analyses (dbRDA) [**Figure 1D**], leading to a dbRDA constrained and unconstrained variation of 41% and 59% respectively (**Figure 1D**). The selected variables were predominantly aligned with the axis summer (daylength, temperature, and the cell abundance of *Synechococcus* and HP) - winter (SiO$_2$, small PNF [**Figure 1D**]. This dbRDA axis had the highest eigenvalue, explaining ~55% of the constrained variation (**Figure 1D**). Even though

the measured environmental variables did not explain the majority of the variation of

the resident microbiota, they could account for a substantial fraction. This was further

supported by Adonis analyses, which indicated that the measured environmental

variables could explain ~45% of the resident microbiota variance, with temperature and

daylength having a predominant role by accounting for 30% of this variance (15%

each).

We then investigated whether temperature and daylength could determine the

main niches. We found that ~70% and ~68% of the OTUs in the resident microbiota

had niche preferences associated with temperature or daylength respectively (**Figure**

1E-F; Note that several OTUs preferring Spring or Autumn are not expected to be

detected with this approach, as their preferred temperature or daylength may not differ

significantly from the randomized mean). In total, 371 OTUs from the resident

microbiota had both a temperature and a daylength niche preference that departed

significantly from the randomization mean (**Figure 1E-F**). These 371 OTUs

represented ~52% of all OTUs in the resident microbiota, corresponding to ~90% of

the sequence abundance. In particular, 248 OTUs had a weighted mean for both

temperature and daylength below the randomization mean (corresponding to

winter/autumn), while 116 OTUs had a weighted mean above the randomization mean

for both variables (corresponding to summer/spring). Interestingly, 7 OTUs displayed

a weighted mean above and below the randomized mean for temperature and daylength

respectively (corresponding to autumn or spring).

Figure 1. The Blanes Bay Microbial Observatory and the variation of its resident microbiota and measured environmental variables over ten years. A) Location of the Blanes Bay Microbial Observatory. **B)** All possible correlations between the measured environmental variables including the richness and abundance of resident OTUs (NB: only 709 resident OTUs are considered, see **Table1**). Only significant Pearson correlation coefficients are shown ($p<0.01$). The p-values were corrected for multiple inference (Holm's method). **C)** Unconstrained

ordination (NMDS based on Bray Curtis dissimilarities) of communities including resident OTUs

only, to which environmental variables were fitted. Only variables with a significant fit are shown

(P<0.05). Arrows indicate the direction of the gradient and their length represents the strength

of the correlation between resident OTUs and a particular environmental variable. The color of

the samples (circles) indicates the season to which they belong. The bottom-left arrow indicates

the direction of the seasonal change. PNF = photosynthetic nanoflagellates. **D)** Constrained

ordination (Distance-based redundancy analyses, dbRDA, using Bray Curtis dissimilarities)

including only the most relevant variables after stepwise model selection using permutation tests.

Each axis (i.e., dbRDA1 and dbRDA2) indicates the amount of variance it explains according to

the associated eigenvalues. The color of the samples (circles) indicates the season to which they

belong. Arrows indicate the direction of the gradient and their length represents the strength of

the correlation between resident OTUs and a particular environmental variable. The bottom-left

arrow indicates the direction of the seasonal change. E-F) Resident OTUs displaying different

niche preferences (blueish areas) in terms of the two most important abiotic variables:

Temperature E) and Daylength F). The red dots indicate the randomization mean, and the orange

curves represent the confidence limits. Black dots indicate individual OTUs for which temperature

or daylength preferences are significantly (p<0.05) higher or lower than a random distribution

over 10 years. At least two assemblages with different niches become evident: one preferring

higher temperature and longer days (summer/spring), and another one preferring lower

temperature and shorter days (winter/autumn). Note that several OTUs associated to Spring or

Autumn are not expected to be detected with this approach, as their preferred temperature or

daylength may not differ significantly from the randomized mean.

Core network

To determine the core microbiota that incorporates possible interactions, we

constructed an association network based on the resident OTUs and removed all OTUs

that were not involved in strong and significant associations with any other OTUs.

Specifically, we kept only the associations (edges in the network) with Local similarity

score $|LS| > 0.7$, a false discovery rate adjusted p-value < 0.001 and Spearman $|r| > 0.7$.

In addition, we removed all associations that seemed to be caused by environmental

preferences of OTUs (see Methods). The core network consisted of 1,411 significant

and strong correlations (**Figure 2A**) and was substantially smaller than the network

based on the resident OTUs without stringent cut-offs (**Supp. Figure 1A**, removed

edges in **Supp. Figure 1B**). The core network includes only the strongest microbial

associations that are inferred during a decade and, according to our definition,

determines the core microbiota. The associations in the core microbiota may represent

proxies for species interactions since steps have been taken to remove associations that

are driven by environmental factors.

In the core network, most associations were positive (~95%), pointing to the

dominance of co-existence or symbiotic associations (**Table 2, Figure 2A**). The core

network had "small world" properties [37], with a small average path length (i.e.

number of nodes between any pair of nodes through the shortest path) and a relatively

high clustering coefficient, showing that nodes tend to be connected to other nodes,

forming tightly knit groups, more than what it would be expected by chance (**Table 3**).

Since node degree was not correlated with OTU abundance (**Supp. Figure 2**), the

associations between OTUs are not caused by a high sequence abundance alone, as the

most abundant OTUs did not tend to be the most connected.

Figure 2. Core microbiota resulting from 10 years of monthly pico- and nanoplankton relative abundances. A) Core network including bacteria and microbial eukaryotic OTUs that occur ≥ 30% of the time during the studied decade (i.e. resident microbiota), with highly significant and strong associations (P<0.001 and Q<0.001, absolute local similarity score |LS| >

0.7, Spearman correlation |ρ|>0.7), where detected environmentally-driven edges were

removed. The color of the edges (links) indicates whether the association is positive (grey) or

negative (red). The shape of nodes indicates bacteria (rhomboid) or microbial eukaryotes (circle),

and the color of nodes represents species seasonal preferences, determined using the indicator

value (*indval*, p<0.05). Node size indicates OTU relative abundance. B) Core network as a Circos

plot, indicating the high-rank taxonomy of the core OTUs. Since 95% of the associations are

positive (see Table 2), we do not indicate whether an edge is positive or negative.

The core network displayed a winter cluster, while no clear clusters could be

defined for the other seasons (**Figure 2A**). Of the 15 environmental variables analyzed,

only 3 were found to be significantly correlated with core OTUs: *daylength*, showing

strong correlations with 33 OTUs, *temperature*, correlated with 14 OTUs, and

Chlorophyll a, correlated with 1 OTU (**Figure 2A**). Therefore, the analysis of the core

network also points to the importance of temperature and daylength in the decade-long

seasonal dynamics of the studied microbial ecosystem. It is also coherent with the

Adonis and ordination analyses (**Figure 1C-B**). However, the associations between

these environmental parameters with taxa represented only 4% of all the associations

(**Figure 2B**).

Table 2. Core associations. See **Figure 2**.

	Association # (edges)	Co-occurrences (positive)	Co-exclusions (negative)
All	1,411	1,341 (95.0%)	70 (5.0%)
Within Picoplankton	378	353 (93.3%)	25 (6.6%)
Within Nanoplankton	791	748 (94.6%)	43 (5.4%)
Picoplankton-Nanoplankton	242	240 (99.2%)	2 (0.8%)

325   Of the 709 OTUs from the resident microbiota (**Figure 3**), only 259 OTUs

326 (35%) were left in the core network (182 bacteria (~70%) and 77 microbial eukaryotic

327 OTUs (~30%); **Table 1, Figure 2**). The monthly taxonomic composition of the resident

328 microbiota differed from that of the core (**Figure 3**). The core OTUs accounted for

329 ~64% of the relative read abundance of the resident microbiota (**Table 1**). The core

OTUs had annual variation in terms of richness and abundance over the 10 years for both the pico- and nanoplankton, with microbial eukaryotes decreasing markedly in OTU richness and relative read abundance in the warmer seasons, and increasing during colder periods (**Figure 3**).

Figure 3. The monthly variation in the resident and core microbiotas over 10 years. *Upper panels:* The resident microbiota is defined as those eukaryotes and bacteria that occur in at least

30% of the samples over 10 years. The relative OTU abundance (left panel) and number of OTUs

(right panel) for different domains and taxonomic levels in the resident microbiota are shown.

Note that the relative abundance of Bacteria vs. Eukaryotes does not necessarily reflect

organismal abundances on the sampling site, but the amplicon relative abundance after PCR.

Relative abundances were calculated for each year and aggregated over the corresponding

months along the 10 years for the resident microbiota, then split into size fractions (NB: relative

abundance for both domains and size fraction sums up to 1 for each month across ten years).

Lower panels: Core microbiota over 10 years. The relative abundances of core OTUs reflect the

remaining proportions after removing all the OTUs that were not strongly associated when

building networks. Relative OTU abundance (left panel) and number of OTUs (right panel) for

different domains and taxonomic levels among the core OTUs.

The most abundant bacteria (**Figure 3**; **Supplementary Table S2**) among the

core OTUs were Oxyphotobacteria (mostly *Synechococcus*), total abundance ~14% of

the resident microbiota, followed by Alphaproteobacteria, with SAR11 clades Ia and II

representing ~9% and ~2% respectively. The most abundant microbial eukaryotic

groups were *Micromonas*, *Bathycoccus*, Dinophyceae, and Cryptomonadales (each

~2%) [**Figure 3**; **Supplementary Table S3**]. In terms of diversity and abundance,

bacterial   non-phototrophs   (including   chemoautotrophs,   photoheterotrophs,

chemoheterotrophs) were the most prevalent in the core microbiota, representing ~62%

of the OTUs and a quarter of the total relative read abundance (**Table 1**). In turn,

protistan heterotrophs represented a minor fraction of the diversity and relative

abundance (**Table 1**). Bacteria photoautotrophs were relatively more abundant than

their protistan counterparts but less diverse (**Table 1**). Protistan parasites represented

~8% of the OTUs and ~3% of the abundance, while the remaining protistan lifestyles

had a minor relevance in the core microbiota (**Table 1**).

Intra- and cross-domain core associations

Bacteria tended to be associated with other bacteria (**Table 3 & 4**; **Figure 2B**), with

Bacteria-Bacteria associations making up ~54% of all associations, while Protist-Protist

associations accounted for 11% (**Table 4**). The connectivity of the bacterial

subnetworks was higher (mean degree ~10) than the protist counterparts (mean degree

~6), regardless of whether these networks included exclusively bacteria, protists, or

both (**Table 3**).

In particular, there was a substantial number of associations between Alpha-

and Gammaproteobacteria, between Alphaproteobacteria and Acidiimicrobia as well as

among Alphaproteobacteria OTUs (**Figure 2B**). Eukaryotic OTUs did not show a

similar trend with associations between OTUs of the same taxonomic ranks (**Figure**

2B). In terms of cross-domain associations, Alphaproteobacteria OTUs had several

associations with most major protistan groups (i.e. dinoflagellates, diatoms,

cryptophytes, Mamiellophyceae, and Syndiniales) [**Figure 2B**].

Core associations within the pico- and within the nanoplankton

While the pico- and nano-size fractions indicate different lifestyles in bacteria (free-

living or particle-attached), they indicate different cell sizes in protists, and this could

be reflected in association networks. Nanoplankton sub-networks were larger and more

connected than picoplankton counterparts (**Figure 4**, **Table 3**). This pattern was

observed in both sub-networks considering associations from the same or both size

fractions (**Table 3**). Nanoplankton sub-networks had a higher average degree (~10)

than picoplankton sub-networks (~7; Wilcoxon p<0.05), while not differing much in other network statistics (**Table 3**). Most associations in the pico- and nanoplankton were positive (>93%), while the associations between OTUs from different size fractions represented only ~17% of the total, being ~99% positive (**Table 2**).

In the pico- or nanoplankton sub-networks that include OTUs from the same size fraction, the number of bacterial core OTUs was higher than the protistan counterparts (103 bacterial vs. 47 protistan OTUs in the nanoplankton, and 79 bacterial vs. 30 protistan OTUs in the picoplankton) (**Figure 4**, **Table 3**). Still, core OTUs in both the pico- and nanoplankton had comparable sequence abundances: ~27% of the resident microbiota in each size fraction. Within the picoplankton, 64% of the associations were between bacteria, 8% between eukaryotes, and 25% between eukaryotes and bacteria (**Table 4**). In turn, in the nanoplankton, 50% of the edges were between bacteria, 14% between eukaryotes, and 31% between eukaryotes and bacteria (**Table 4**). Overall, the BBMO pico- and nanoplankton sub-networks differed in size, connectivity, and taxonomic composition, while they were similar in terms of positive connections and relative sequence abundance.

Table 3. Core network and sub-networks statistics.

Network	Nodes (#OTUs)	Edges	Di.	De.	Average degree	Average path length	Average clustering coefficient	Largest clique (#)	Mod.
Core network	262 (259)	1,411	11	0.04	10.7	3.45	0.52	13 (4)	0.19
Random core network	262	1,411	5	0.04	10.7	2.60	0.03	3(199)	0.13
Picoplankton all [1]	161 (160)*	620*	10	0.05	7.7	3.13	0.55	10(1)	0.22
Picoplankton only [2]	110 (109)	378	9	0.06	6.9	3.15	0.51	9(4)	0.29
Nanoplankton all [3]	197 (194)*	1,033*	10	0.05	10.5	3.18	0.57	13(4)	0.15
Nanoplankton only[4]	153 (150)	791	10	0.07	10.3	3.21	0.56	13(4)	0.17
Bacteria all [5]	233 (230)**	1,236**	10	0.04	10.6	3.34	0.52	11(3)	0.19
Bacteria only [6]	185 (182)	803	10	0.05	8.7	3.50	0.51	10(1)	0.31
Protists all [7]	147 (145)**	608**	5	0.06	8.3	2.40	0.48	8(2)	0.10
Protist only [8]	80 (77)	175	5	0.05	4.4	2.54	0.54	7(1)	0.32

NB: Networks and sub-networks include OTUs and environmental factors. Di=Network diameter. De=Network density. Largest clique = size of the largest clique(s) in the network, and in brackets, the number of them. Mod = Network modularity inferred using edge betweenness. [1]All associations where picoplankton OTUs are involved (including nanoplankton); [2]Associations between picoplankton OTU only; [3]All associations where nanoplankton OTUs are involved (including picoplankton); [4]Associations between nanoplankton OTU only; [5]All associations where bacterial OTUs are involved (including protists); [6]Associations between bacterial OTU only; [7]All associations where protist OTUs are involved (including bacteria); [8]Associations between protist OTU only. * Includes nodes and edges shared between pico- and nanoplankton. ** Includes nodes and edges shared between bacteria and protists.

Figure 4. Pico- and nanoplankton core sub-networks. The shape of the nodes indicates

bacteria (rhomboid) or microbial eukaryotes (circle), and the color of nodes represents species

seasonal preferences, determined using the indicator value (p<0.05). The color of the edges

419 indicates if the association is positive (grey) or negative (red). Node size indicates OTU relative

420 abundance from the core microbiota.

422 **Table 4.** Core associations within and between taxonomic domains and size fractions.

Network	Association type[1]	# Associations
Core network	Total	1,411
	Bacteria - Bacteria	767 (54%)
	Bacteria - Protist	433 (31%)
	Protist - Protist	161 (11%)
	Environmental factor - Bacteria	36 (3%)
	Environmental factor - Protist	14 (1%)
Picoplankton subnetwork	Total	378
	Bacteria - Bacteria	241 (64%)
	Bacteria - Protist	94 (25%)
	Protist - Protist	31 (8%)
	Environmental factor - Bacteria	12 (3%
	Environmental factor - Protist	0 (0%)
Nanoplankton subnetwork	Total	791
	Bacteria - Bacteria	394 (50%)
	Bacteria - Protist	246 (31%)
	Protist - Protist	113 (14%)
	Environmental factor - Bacteria	24 (3%)
	Environmental factor - Protist	14 (2%)

423 [1] "Bacteria – Bacteria" indicates associations between two bacterial OTUs. "Protist – Protist" are associations between two unicellular eukaryotes
424 and "Bacteria – Protist" are associations between one eukaryote and one bacterial OTU. "Environmental factor – Protist" and "Environmental
425 factor – Bacteria" are associations between an environmental factor and a eukaryotic or bacterial OTU.
426

Network seasonality

The indicator value (IndVal) was used to infer the seasonal preference of core OTUs.

Most of the core OTUs (98%; 254 out of 259 OTUs) showed a clear preference for one

of the four seasons, pointing to a marked seasonality in the core microbiota (**Figure 4;**

Table 5; Tables S4 & S5). Winter had the highest quantity of core OTUs and the

highest network connectivity (average degree ~13), compared to the other seasons

(average degrees ~2 – ~6) [**Figure 4**; **Table 5**]. The average path length was larger in

the core network compared to a random network of the same size (**Table 3**). Yet, all

sub-networks associated with size fractions and seasons (**Table 5**) had shorter path

lengths than the random network,

that nodes tended to be connected within seasons and size fractions. This was also

supported by an increase in network density when comparing the core network (**Table**

3) and the core network subdivided into seasons (**Table 5**), against the core network

subdivided into both seasons and size fractions (**Table 5**). The five OTUs that did not

show any seasonal preference, among them SAR11 Clades Ia & II, showed high to

moderate abundances but had a low number of associations to other OTUs (**Tables S4,**

S5, S6). Thus, network connectivity in the BBMO appears to be heterogeneous over

time, peaking in winter and remaining low in the other seasons.

Table 5: Subnetworks including core OTUs displaying seasonal preference.
447

	Sub-network	Number of OTUs	Edges	Di.	De.	Average degree	Average path length	Average clustering coefficient	Largest clique (#)	Mod.
All	Winter	156	1,175	7	0.10	15.1	2.62	0.54	13(4)	0.19
	Spring	19	16	4	0.09	1.7	1.56	0.44	4(1)	0.75
	Summer	41	56	7	0.07	2.7	2.90	0.49	6(1)	0.53
	Autumn	26	25	3	0.08	1.9	1.59	0.46	4(2)	0.73
Pico	Winter	63	286	6	0.15	9.1	2.35	0.53	9(4)	0.10
	Spring	8	5	3	0.18	1.2	1.50	0.00	2(5)	0.56
	Summer	25	36	5	0.12	2.9	2.20	0.41	6(1)	0.23
	Autumn	5	3	2	0.30	1.2	1.25	0.00	2(3)	0.44
Nano	Winter	92	658	6	0.16	14.3	2.40	0.61	13(4)	0.04
	Spring	11	11	4	0.20	2.0	1.59	0.57	4(1)	0.56
	Summer	13	17	3	0.22	2.6	1.70	0.65	4(1)	0.50
	Autumn	17	18	3	0.13	2.1	1.35	0.56	4(2)	0.60

448
449 NB: Subnetworks include OTUs only. Di=Network diameter. De=Network density. Largest clique = size of the largest clique(s) in the network, and in brackets, the number of them. Mod = Network modularity inferred using edge betweenness.
450

Groups of highly associated OTUs

Within the core network, we identified groups that were more connected to each other

than to the rest of the network (called modules). These groups of OTUs may indicate

recurring associations that are likely important for the stability of ecosystem function.

We identified 12 modules in both the pico- and nanoplankton subnetworks (**Table S7**).

Modules tended to include OTUs from the same season (**Table S8**), with main modules

(i.e. MCODE score >4) including OTUs predominantly associated with winter,

summer, and autumn (**Figure 5**). Overall, winter modules prevailed (5 out of 7) among

the main modules (**Figure 5**), while modules with scores ≤ 4 did not tend to be

associated with a specific season (**Table S8**). Two main winter modules had members

that were negatively correlated to temperature and daylength (**Figure 5**; Modules 1 and

4, nanoplankton).

The total relative sequence abundance of core OTUs included in modules was

~24% (proportional to the resident microbiota), while the total abundance of individual

modules ranged between ~6% and ~0.3% (**Table S7**). In turn, the relative abundance

of core OTUs included in modules ranged between 0.01% and ~2% (**Table S8**). In most

modules, a few OTUs tended to dominate the abundance, although there were

exceptions, such as module 4 of the picoplankton, where all SAR11 members featured

abundances >1% (**Table S8**). In addition, several OTUs within modules had relatively

low abundances (**Table S8**), supporting modules as a real feature of the network and

not just the agglomeration of abundant taxa.

Central OTUs

Biological networks typically contain nodes (i.e. OTUs) that hold more "central"

positions in the network than others [22]. Even though the ecological role of these hub

and connector OTUs is unclear, it is acknowledged that they could reflect taxa with

important ecological functions [22]. There is no universal definition for hub or

connector OTUs, yet, in this work, we have used stringent thresholds to determine them

ad hoc (see Methods). We have identified 13 hub-OTUs that were associated with

winter or spring (**Table 6**). Hubs did not include highly abundant OTUs, such as

Synechococcus or SAR11 (**Table 6**), but instead, they included several OTUs with

moderate-low abundance (<1%) and high degree (ranging between 26-60) [**Table 6**].

For example, the Gammaproteobacteria OTU bn_000226 had a relative abundance of

0.04% and a degree of 60 (**Table 6**). Hubs included other moderately abundant OTUs,

such as the eukaryotic picoalgae *Bathycoccus,* which was abundant in winter, as well

as an unidentified dinoflagellate (**Table 6**).

We identified a total of 18 connector OTUs (featuring relatively low degree and

high centrality), which were predominantly associated with summer (5 out of 18) or

autumn (6 out of 18), contrasting with hub OTUs, which were associated mostly with

winter and spring (**Table 6**). Connectors may be linked to the seasonal transition

between main community states (**Figure 1 C & D**) and included several abundant

OTUs belonging to *Synechococcus* and SAR11 (**Table 6**). In particular, the SAR11

OTU bp_000007 displayed a relatively high abundance (1.4%), but a degree of 3

(relatively low) and a betweenness centrality of 0.6 (relatively high). In contrast, two

protist OTUs displayed low-moderate abundances (ep_00269, Chrysophyceae,

abundance 0.04% and en_00161, Syndiniales, abundance 0.4%), low degree <4, but a

high betweenness centrality (>0.8; **Table 6**).

Table 6. Central OTUs

OTU	Class	Lowest rank taxonomy	Relative Abundance (%)[1]	Degree	Betweenness Centrality	Closeness Centrality	Season
Hubs							
en_00092	Mamiellophyceae	*Bathycoccus*	0.51	42	0.04	0.42	Winter
en_00119	Dinophyceae	-	0.41	50	0.03	0.42	Winter
bp_000037	Alphaproteobacteria	Parvibaculales_OCS116	0.31	45	0.08	0.43	Winter
bp_000039	Gammaproteobacteria	SUP05_cluster	0.28	29	0.12	0.41	Spring
bn_000039	Gammaproteobacteria	SUP05_cluster	0.21	42	0.17	0.44	Spring
bn_000037	Alphaproteobacteria	Parvibaculales_OCS116	0.20	40	0.05	0.42	Spring
bp_000059	Gammaproteobacteria	SAR86	0.20	24	0.09	0.40	Spring
ep_00070	Cryptophyceae	Cryptomonadales_X	0.13	40	0.04	0.42	Winter
bn_000059	Gammaproteobacteria	SAR86	0.12	24	0.03	0.40	Spring
bn_000102	Alphaproteobacteria	Nisaeaceae_OM75	0.09	26	0.03	0.38	Winter
bp_000193	Alphaproteobacteria	-	0.06	37	0.03	0.40	Winter
bn_000170	Acidimicrobiia	Sva0996_marine_group	0.06	59	0.06	0.44	Winter
bn_000226	Gammaproteobacteria	HOC36	0.04	60	0.06	0.43	Winter
Connectors							
bp_000001	Oxyphotobacteria	*Synechococcus* (CC9902)	3.79	5	0.05	0.30	Autumn
bp_000002	Alphaproteobacteria	SAR11 Clade_Ia	2.26	2	0.40	0.56	Spring
bp_000004	Alphaproteobacteria	SAR11 Clade_Ia	2.02	3	0.15	0.63	NA
bp_000007	Alphaproteobacteria	SAR11 Clade_Ia	1.38	3	0.60	0.71	NA
bp_000008	Alphaproteobacteria	SAR11 Clade_Ia	1.15	3	0.15	0.63	NA
bn_000008	Alphaproteobacteria	SAR11 Clade_Ia	0.68	5	0.03	0.27	Winter

en_ 00059	Chlorodendrophyceae	*Tetraselmis*	0.66	4	0.05	0.26	Summer
bn_ 000020	Oxyphotobacteria	-	0.56	3	0.60	0.67	Autumn
en_ 00161	Syndiniales	Syndiniales-Group-I-Clade-4_X	0.42	4	0.80	0.75	Autumn
bn_ 000018	Oxyphotobacteria	*Prochlorococcus* MIT9313	0.41	5	0.04	0.24	Winter
bn_ 000054	Alphaproteobacteria	Puniceispirillales_SAR116	0.11	4	0.14	0.40	Autumn
bn_ 000062	Alphaproteobacteria	Puniceispirillales_SAR116	0.08	3	0.55	0.50	Autumn
bn_ 000077	Rhodothermia	*Balneola*	0.07	3	0.17	0.32	Summer
bn_ 000112	Gammaproteobacteria	KI89A	0.06	4	0.53	0.48	Summer
bn_ 000156	Alphaproteobacteria	Parvibaculales_PS1	0.05	4	0.14	0.40	Summer
bn_ 000281	Bacteroidia	Sphingobacteriales_NS11-12	0.05	5	0.16	0.44	Autumn
bn_ 000221	Alphaproteobacteria	Puniceispirillales_SAR116	0.04	5	0.05	0.30	Winter
ep_ 00269	Chrysophyceae	Clade-I_X	0.04	2	1.00	1.00	Summer

500   [1] Proportional to the resident microbiota

Figure 5. Main modules in the core network. Modules with MCODE score >4 are shown for picoplankton (upper panel) and nanoplankton (lower panel). For each module, the MCODE score and relative amplicon abundance of the taxa included in it (as % of the resident microbiota) are indicated. In addition, the numbers of edges and OTUs within the modules are shown as edges/OTUs; this quotient estimates the average number of edges per OTU within the different modules. The edges represent correlations with |LS| > 0.7, |ρ|>0.7, P<0.001 and Q<0.001. The color of the edges indicates positive (grey) or negative (red) associations. The shape of nodes

indicates bacteria (rhomboid) or microbial eukaryotes (circle), and the color of nodes represents species seasonal preferences, determined using the indicator value (p<0.05). pb = Proteobacteria

DISCUSSION

Identifying the most important microbes for the functioning of the ocean ecosystem is a challenge, which can be addressed by delineating core microbiotas [4]. Recognizing the most abundant and widespread microbes in the ocean is a step towards knowing the core microbiota. However, this does not take into account the importance that both microbial interactions and microbes with moderate or low abundance may have for the functioning of ecosystems [4, 29, 38]. Considering potential interactions when delineating core microbiotas may not only allow identifying moderate/low abundance taxa that may have important roles in the community but could also allow excluding taxa that are present in several locations but that may not have an important role for community function (e.g., dormant cells or cells being dispersed [8]). Here, we have delineated and analyzed the core microbiota of a coastal ecosystem-based on 10 years of occurrence data considering possible interactions.

To detect the core microbiota, we first identified the resident OTUs, that is, those that occur >30% of the time (i.e. >36 out of 120 months) over a decade. This threshold was selected as it allows for seasonal OTUs that would be present recurrently in at least one season. Analysis of the resident OTU dynamics indicated a clear seasonality (**Figure 1 C-D**), and that the measured environmental factors could explain ~45% of the resident microbiota variance. The main environmental drivers were temperature and daylength, which is consistent with previous works from the same time-series (BBMO) [34, 39, 40]. These values are lower than what has been reported

for bacteria in the English Channel, where daylength explains ~65% of community

variance [17], and higher than what has been reported for entire communities in the

time-series SPOT (California, 31%) [41] or SOLA (the Mediterranean Sea, ~130 km

from BBMO; 7-12%) [42]. Daylength may be more important in the English Channel

as it has a more pronounced annual variation than at BBMO, whereas the measured

differences could reflect a higher coupling of the resident OTUs with environmental

variation in BBMO than in SOLA or SPOT. SOLA is characterized by the occasional

winter storms that bring nutrients from the sediments to the water column as well as by

the freshwater inputs from nearby rivers during flash floods [43], and this could

partially explain the differences with BBMO. The importance of daylength and

temperature for community dynamics was reflected by niche analyses, which identified

two main niches associated with summer and winter at the BBMO, to which ~50% of

the resident OTUs were associated (**Figure 1 E-F**). Other resident OTUs likely have

spring and fall niches as indicated by **Figure 1 C-D**, yet these niches cannot be detected

with the used null model analysis, as their preferred temperatures or daylengths will not

depart significantly from the randomized mean.

Based on the resident OTUs, we built networks to define the core microbiota.

We identified a total of 259 core OTUs (182 bacteria and 77 protists) that represented

64% of the abundance of the resident microbiota and that showed seasonal variation.

We could only find supporting evidence from the literature (PIDA database) [21] for

85 associations of the core (6 %), indicating that most of them still need to be validated

with direct observation or experimentally. This is not surprising, as the most studied

hosts in PIDA are protists from the micro-plankton (>20 µm cell size), which are mostly

absent from our pico- and nanoplankton networks. Also, PIDA does not cover Bacteria-

Bacteria associations. Nevertheless, the detected core OTUs from BBMO represent a

fraction of the core microbiota at this site, since larger microbial size fractions were not

sampled. Including these larger size fractions would expand the composition of the core

and could unveil additional patterns. For example, in a global ocean network including

size fractions >20 µm cell size, protists or small multicellular eukaryotes dominated the

interactome [26].

566   Alpha-/Gammaproteobacteria, Bateroidia, Acidimicrobiia were the main

bacterial groups in the core, including also common marine taxa, such as

Synechococcus or SAR11. The main protists in the core included Syndiniales

(parasites), Dinoflagellates, Mammiellales (*Micromonas* and *Bathycoccus*), and

diatoms. These taxa are likely the most important in sustaining ecosystem function at

BBMO, and probably have similar importance in other coastal areas. Other studies have

reported important roles in marine association networks for SAR11 and *Synechococcus*

[31, 44]. Syndiniales, Haptophytes, and Dinoflagellates dominated networks in terms

of the number of nodes and edges at SPOT, while Mamiellales (*Micromonas &*

Bathycoccus) and diatoms also had relevant roles [41]. Syndiniales, Dinoflagellates,

and Diatoms were also predominant in global ocean networks, which is coherent with

our results [26].

578   Bacteria-Bacteria associations were the most abundant (54%) in the core

BBMO microbiota, followed by Bacteria-Protists (31%) and Protist-Protist (11%)

associations. Associations tended to occur among bacteria or protists, rather than

between them, in the English Channel time-series [17]. However, the study used

microscopy to determine protist community composition, while it used 16S-rRNA gene

data for analyzing bacteria communities and this might explain the limited number of

connections between protists and bacteria. Most associations occurred among protists

in a global-ocean network that included a broad range of microbial size-fractions [26].

This suggests that time-series analyses including larger size-fractions may determine a

higher proportion of associations among protists, which may turn out to be prevalent.

The core network had "small world" properties (that is, high clustering

coefficient and relatively short path lengths) [37] when compared to randomized

networks (**Table 3**) or particular subnetworks from size fractions or specific seasons

(**Table 5**). The small-world topology is characteristic of many different types of

networks [45], including marine microbial temporal or spatial networks [23, 26, 30,

31]. Some of our network statistics were similar to those obtained at SPOT [23, 30], in

particular the averages of degree, clustering coefficient, and path length (**Table 3**).

Furthermore, the BBMO network had an average path length similar to a global ocean

network [26] and also, similarly to this network, the node degree of the BBMO core

members was independent of their relative abundances, showing that the associations

between core OTUs were not merely a consequence of high prevalence and abundance.

The BBMO core network had a clustering coefficient that was ten times larger

than that of an Erdős–Rényi random network of the same size (**Table 3**), which agrees

with what was observed at SPOT [23, 30]. The large proportion of positive associations

in BBMO networks (~95%) was in agreement with results from other temporal [23, 41]

or large-scale spatial [26] microbiota analyses, where positive associations were also

predominant (~70-98%), although these values include taxa that are not necessarily part

of the core. This suggests that interactions such as syntrophy or symbiotic associations

are more important than competition in marine microbial systems and that these types

of associations may underpin marine ecosystem function. These findings are also

coherent with a recent large-scale literature survey that found that ~47% of the validated

associations between protists and bacteria are symbiotic [21]. Nevertheless, it is also

possible that common sampling strategies and methodological approaches do not detect

a substantial fraction of negative associations. For example, while positive correlations

in taxa abundance pointing to positive interactions may be easier to detect, negative

associations may be missed due to plummeting species abundances that would prevent

establishing significant correlations, or to a delay between the increase and decrease in

abundance of interacting taxa that are not synchronized with sampling time. Future

studies adapting the sampling scheme to the timing of interactions (e.g., daily or weekly

sampling) and the use of other approaches apart from taxa abundances, such as analyses

of single-cell genomic data to determine protistan predation, or controlled experiments,

will likely generate new insights on negative microbial interactions.

The relatively high clustering coefficient of the core network (compared to a

random network) and its short path length indicate that most OTUs are connected

through < 3 intermediary OTUs. It has been shown that a large proportion of strong

positive associations, as in the BBMO core network, may destabilize communities due

to positive feedbacks between species [46]. When a species decreases in abundance as

a response to environmental variation, it may pull others with it, generating a cascade

effect propagated by the many positive associations in the network. Accordingly, the

change of abundance in specific OTUs in one section of the network could affect OTUs

in other network sections not necessarily affected directly by the environmental

variation. This cascade effect may help to explain a paradox: environmental variables

affect the structure of marine microbial communities and consequently association

networks. Yet, our and others' results [17, 18, 23, 26, 30-32] have reported a limited

number of associations between environmental variables and network nodes (OTUs).

Environmental heterogeneity might affect network structure by acting on a small subset

of nodes (OTUs), which would then influence other nodes through cascading

interactions facilitated by the highly interconnected nature of the networks as well as

positive feedbacks promoted by the high proportion of positive associations [46].

If OTUs susceptible to environmental variation are also highly connected, then

their effect on the entire network structure may be larger. In line with this, we found

that the connectivity of OTUs associated with environmental variables at BBMO (49

OTUs out of 259) had a mean degree of ~25 (SD ~14), while for all the 259 OTUs of

the core network, the mean degree was ~11 (SD ~13). The seasonal dynamics of the

BBMO microbiota may partially be driven by a subset of OTUs that vary with

environmental factors (e.g. temperature, daylength). These may exert a destabilizing

influence over the entire community over time, promoting the annual turnover of

communities and networks.

Most core OTUs (98%) showed a clear preference for one season. Interestingly,

the distribution of core OTUs among the seasons was uneven, with 61% of these OTUs

showing a winter preference. Network connectivity at BBMO was correspondingly

heterogeneous between seasons, peaking in winter and remaining low in the other

seasons. Specifically, the winter subnetwork included ~92% of the seasonal edges. This

indicates that winter associations are not only specific (i.e. they do not tend to change

partners), but they also have a relatively high recurrence (otherwise, winter networks

would be smaller). A higher similarity between winter communities when compared to

other seasons was also indicated by our ordination analyses of the resident OTUs

(**Figure 1**), as well as by studies of the entire protist community at BBMO [34] or whole

community analyses at SPOT [23].

The structure of communities is determined by the interplay of selection,

dispersal, speciation, and ecological drift [47]. Our results indicate that selection, a

deterministic process, is stronger in winter, leading to winter sub-communities that tend

to be more similar between each other than to communities from other seasons. Given

that we have removed edges associated with the measured environmental variables, we

do not expect that the identified edges between winter OTUs represent selection

associated to these variables (e.g. low temperature). Consequently, winter edges may

represent associations linked to unmeasured variables or ecological interactions that

may be more likely to develop during winter due to stronger environmental selection.

Due to weaker selection in other seasons species occurrence would display less

recurrent (or more random) patterns, preventing specific associations to be formed. This

also suggests that ecological redundancy changes over time, and is lower in winter

compared to the other seasons (even though the number of OTUs is larger in winter).

A reduction in redundancy may also promote strong ecological interactions in winter.

The existence of subsets of species that interact more often between themselves

than with other species (modules), is characteristic of biological networks, and can

contribute to overall network stability [48, 49]. Modules can represent divergent

selection, niches, the clustering of evolutionary closely related species or co-

evolutionary units [50, 51]. Modules in the core BBMO network (total 12) included

positive associations between diverse taxa, and could represent divergent selection,

driven by unmeasured environmental variables, or examples of syntrophic or symbiotic

interactions between microbes from different taxonomic groups.

Most BBMO modules included diverse lifestyles (heterotrophs, mixotrophs,

phototrophs, parasites), similar to what has been observed at SPOT [41]. Yet, a number

of modules appeared to be predominantly heterotrophic or autotrophic (**Table S8**).

Some modules included OTUs from the same species, such as Module 4 in the

picoplankton, which included several SAR11 Clade I OTUs, and Module 7 of the

nanoplankton, which included several *Synechococcus* OTUs. These modules could

reflect similar niches, associated with unmeasured variables, or the dependence on

metabolites produced by other organisms (auxotrophy). There is evidence of

auxotrophy for both SAR11 (e.g. thiamin, glycine)[52-54] and *Synechococcus* (e.g.

cobalamin) [55]. Recently it has been observed in co-culture experiments that

Prochlorococcus may fulfill some metabolic requirements of SAR11, promoting the

growth of the latter in a commensal relationship [56]. In our analyses of the BBMO

core microbiota, we did not find strong associations between SAR11 and

Prochlorococcus or the more abundant relative, *Synechococcus*. Yet, SAR11 formed

strong associations with a plethora of taxa with which could potentially have

commensal relationships.

The overall importance of the observed modules was indicated by the total

abundance of their constituent OTUs (24% of the reads compared to the resident

microbiota). Most of the modules at BBMO were associated with a single season,

suggesting that they reflect seasonal niches. Since these modules were inferred over 10

years, they represent recurrent network features. Chafee et al. [57] also identified

season-specific modules in a 2-year time series in the North Sea (Helgoland), including

samples taken weekly or bi-weekly. These modules were much larger than ours, and

they may also include environmentally-driven edges. Nevertheless, the Helgoland

modules seem to be driven by eutrophic (spring & summer) vs. oligotrophic (autumn

& winter) conditions in this location. In contrast, the BBMO modules, displayed weaker

correlations with nutrients and seem to be influenced by temperature and daylength

(**Figure 5**). Differences in the sampling scheme between Helgoland and BBMO

((bi)weekly vs. monthly) as well as between both locations (different seas and latitudes,

affecting temperature and daylength) may explain these differences.

Keystone species have a high influence in ecosystems relative to their

abundance [58]. Network analyses may help to identify them [24, 59], yet, there is no

clear consensus of what network features are the best unequivocal indicator of keystone

species [60-62]. Therefore, we focused on identifying central OTUs (hubs or

connectors) that may be important for ecosystem function [22, 24] and could represent

keystone species. We identified 13 hubs in the BBMO core network with moderate-low

abundances (<1%) and high degree (26-60) that were associated with winter or spring.

These moderate-low abundance OTUs may affect nutrient cycling directly [63] or

indirectly, by affecting other OTUs with higher abundance. The putative stronger

selection exerted by low temperatures and short daylengths during winter and early

spring, as compared to summer and autumn, may lead to a higher species recurrence

[34], larger networks, and possibly, more hubs. An OTU of the abundant picoalgae

Bathycoccus (en_00092) was identified as a winter hub, which is consistent with

reported *Bathycoccus* abundance peaks in late winter (February-March) in both BBMO

[64] and the nearby station SOLA [42]. This *Bathycoccus* hub may be associated with

diverse taxa, such as prokaryotes that may benefit from algal exudates [65] or even via

mixotrophy [66]. In agreement with this, out of the 42 associations of this hub OTU,

25 were with bacteria and the rest with protists.

In contrast to hubs, connector OTUs were predominantly associated with

warmer waters, that is, summer and autumn, and may represent transitions in

community states. This was consistent with the associations observed in an abundant

Synechococcus connector OTU (bp_000001, **Table 6**). This OTU was predominant in

summer-autumn, in agreement with previous BBMO reports [36, 67], but it was

associated with other OTUs from spring (negative association with bp_000017), winter

(negative association with bp_000039), summer (positive association with bp_000087,

bp_000012) and autumn (positive association with bp_000022), thus likely holding a

central position in the network. Another abundant spring connector OTU (SAR11 Clade

Ia, bp_000002), featured only two connections to spring (positive association with

bp_000007) and summer (positive association with bp_000046) OTUs.

CONCLUSION

Our decade-long analysis of the dynamics of a microbiota populating a time-series in

the Mediterranean Sea allowed us to determine the interconnected core microbiota,

which likely includes several microbes that are important for the functioning of this

coastal ecosystem. We found a relatively small core microbiota that displayed seasonal

variation, with a heterogeneous distribution of associations over different seasons,

indicating different degrees of recurrence and selection strength over the year. Future

analyses of other core marine microbiotas will determine how universal are the patterns

found in BBMO. These studies will be crucial to determine potential long-term effects

of climate change on the architecture of the interaction networks that underpin the

functioning of the ocean ecosystem.

METHODS

Study site and sampling

Surface water (~1 m depth) was sampled monthly from January 2004 to December

2013 at the Blanes Bay Microbial Observatory (BBMO) in the Northwestern

Mediterranean Sea (41º40'N, 2º48'E) [**Figure 1A**]. The BBMO is an oligotrophic

coastal site ~1 km offshore with ~20 m depth and with limited riverine or human

influence [36]. Seawater was pre-filtered with a 200 μm nylon mesh and then

transported to the laboratory in 20 L plastic carboys and processed within 2 hours.

Microbial plankton from about 6 L of the pre-filtered seawater was separated into two

size fractions: picoplankton (0.2-3 μm) and nanoplankton fraction (3-20 μm). To achieve this, the seawater was first filtered through a 20 μm nylon mesh using a peristaltic pump. Then the nanoplankton (3-20 μm) was captured on a 3 μm pore-size polycarbonate filter. Subsequently, a 0.2 μm pore-size Sterivex unit (Millipore, Durapore) was used to capture the picoplankton (0.2-3 μm). Sterivex units and 3 μm filters were stored at -80 ºC until further processed. The sequential filtering process aimed to capture free-living bacteria and picoeukaryotes in the 0.2-3 μm size fraction (picoplankton), and particle/protist-attached bacteria or nanoeukaryotes in the 3-20 μm fraction (nanoplankton). The 3μm filter was replaced if clogging was detected; DNA from all 3μm filters from the same sample were extracted together.

A total of 15 contextual abiotic and biotic variables were considered for each sampling point: Daylength (hours of light), Temperature (°C), Turbidity (estimated as Secchi disk depth [m]), Salinity, Total Chlorophyll a [Chla] (μg/l), PO_4^{3-} (μM), NH_4^+ (μM), NO_2^- (μM), NO_3^- (μM), SiO_2 (μM), abundances of Heterotrophic prokaryotes [HP] (cells/ml), *Synechococcus* (cells/ml), Total photosynthetic nanoflagellates [PNF; 2-5μm size] (cells/ml), small PNF (2μm ; cells/ml) and, Heterotrophic nanoflagellates [HNF] (cells/ml) [**Figure 1B**]. Water temperature and salinity were sampled *in situ* with a SAIV A/S SD204 CTD. Inorganic nutrients (NO_3^-, NO_2^-, NH_4^+, PO_4^{3-}, SiO_2) were measured using an Alliance Evolution II autoanalyzer [68]. Cell counts were done by flow cytometry (heterotrophic prokaryotes, *Synechococcus*) or epifluorescence microscopy (PNF, small PNF and HNF). See Gasol *et al.* [36] for specific details on how other variables were measured. Environmental variables were z-score standardized before running statistical analysis.

DNA extraction, sequencing, and metabarcoding

DNA was extracted from the filters using a standard phenol-chloroform protocol [69],

purified in Amicon Units (Millipore), and quantified and qualitatively checked with a

NanoDrop 1000 Spectrophotometer (Thermo Fisher Scientific). Eukaryotic PCR

amplicons were generated for the V4 region of the 18S rDNA (~380 bp), using the

primer pair TAReukFWD1 and TAReukREV3 [70]. The primers Bakt_341F [71] and

Bakt_806RB [72] were used to amplify the V4 region of the 16S rDNA. PCR

amplification and amplicon sequencing were carried out at the Research and Testing

Laboratory (http://rtlgenomics.com/) on the *Illumina* MiSeq platform (2x250 bp paired-

end sequencing). DNA sequences and metadata are publicly available at the European

Nucleotide Archive (http://www.ebi.ac.uk/ena; accession numbers PRJEB23788 for

18S rRNA genes & PRJEB38773 for 16S rRNA genes).

A total of 29,952,108 and 16,940,406 paired-end *Illumina* reads were produced

for microbial eukaryotes and prokaryotes respectively. Adapters and primers were

removed with Cutadapt v1.16 [73]. DADA2 v1.10.1 [74] was used for quality control,

trimming, and inference of Operational Taxonomic Units (OTUs) as Amplicon

Sequence Variants (ASVs). For both microbial eukaryotes and prokaryotes, the

Maximum number of expected errors (MaxEE) was set to 2 and 4 for the forward and

reverse reads respectively. No ambiguous bases (Ns) were allowed. Microbial

eukaryotic sequences were trimmed to 220 bp (forward) and 190 bp (reverse), while

prokaryotic sequences were trimmed to 225 bp (both forward and reverse reads). A

total of 28,876 and 19,604 OTUs were inferred for microbial eukaryotes and

prokaryotes respectively.

OTUs were assigned taxonomy using the naïve Bayesian classifier method [75]

together with the SILVA version 132 [76] database as implemented in DADA2.

Eukaryotic OTUs were also BLASTed [77] against the Protist Ribosomal Reference

database (PR^2, version 4.10.0; [78]). When the taxonomic assignments for the

eukaryotes disagreed between SILVA and PR^2, the conflict was resolved manually by

inspecting a pairwise alignment of the OTU and the closest hits from the two databases.

OTUs assigned to Metazoa, Streptophyta, nucleomorph, chloroplast, and mitochondria

were removed before further analysis. Archaea were removed from downstream

analyses as the used primers are not optimal for recovering this domain [79].

Each sample (corresponding to a specific gene, size fraction, and timepoint) was

subsampled with the *rrarefy* function from the R package *Vegan* [80] to 4,907 reads,

corresponding to the number of reads in the sample with the lowest sequencing depth,

to normalize for different sequencing depth between samples. OTUs present in <10%

of the samples were removed. After quality control and rarefaction, the number of

OTUs was 2,926 (1,561 bacteria, and 1,365 microeukaryotes; **Table 1**).

Due to a suboptimal sequencing of the amplicons, we did not use nanoplankton

samples of bacteria and protists from the period May 2010 to July 2012 (27 samples)

as well as March 2004 and February 2005. OTU read abundance for samples with

missing values were estimated using seasonally aware missing value imputation by

weighted moving average for time series as implemented in the R package *imputeTS*

[81].

Cell/particle dislodging or filter clogging during the sequential filtration process

may affect the taxonomic diversity observed in the different size fractions, with

nanoplankton DNA leaking into the picoplankton fraction, or picoplankton DNA

getting stuck in the nanoplankton fraction. To minimize the effects of cell/particle

dislodging or filter clogging on the diversity recovered from the different size fractions,

we calculated the sequence-abundance ratio for OTUs appearing in both pico- and

nano-plankton fractions. When the ratio exceeded 2:1, we removed the OTU from the

size fraction with the lowest number of reads. After subsampling and filtering the OTU

tables were joined for each time point, and since the samples had been normalized to

the same sequencing depth, we calculated the relative read abundance for the OTUs for

each year and aggregated over the corresponding months along the 10 years for the

resident microbiota. This means that the relative abundance for both domains and size

fractions sums up to 1 for each month across ten years.

Resident microbiota

We defined *ad hoc* the resident microbiota as the set of OTUs present in >30% of the

samples over 10 years (that is, present in >36 months, not necessarily consecutive).

This value was chosen as it allows for seasonal OTUs, which may only be present 3-4

months each year, and still be considered as part of the resident microbiota. The

residents included 355 eukaryotic and 354 bacteria OTUs (**Table 1**), and excluded a

substantial amount of rare OTUs, which can cause spurious correlations during network

construction due to sparsity [i.e. too many zeros] [22]. The relative abundance of the

taxonomic groups included in the resident microbiota was fairly stable from year to

year (**Figure 3**).

Environmental variation and resident OTUs

All possible correlations among the measured environmental variables and resident

OTU richness and abundance were computed in R and plotted with the package

corrplot. Only significant Pearson correlation coefficients were considered ($p<0.01$),

and the p-values were corrected for multiple inference (Holm's method) using the

function *rcorr.adjust* from the R package *RcmdrMisc*. Unconstrained ordination

analyses were carried out using NMDS based on Bray Curtis dissimilarities between

samples including resident OTUs only. Environmental variables were fitted to the

NMDS using the function *envfit* from the R package *Vegan* [80]. Only variables

displaying a significant correlation (p<0.05) were considered. Constrained ordination

was performed using distance-based redundancy analyses (dbRDA) in *Vegan*,

considering Bray Curtis dissimilarities between samples including resident OTUs only.

The most relevant variables for constrained ordination were selected by stepwise model

selection using 200 permutations, as implemented in *ordistep* (*Vegan*). Ordinations

were plotted using the R package *ggplot2* and *ggord*. The amount of community

variance explained by the different environmental variables was calculated with *Adonis*

(Vegan) using 999 permutations. Resident OTUs displaying niche preference in terms

of Temperature and Daylength, the most important environmental variables, were

determined using the function *niche.val* from the R package *EcolUtils* with 1,000

permutations.

Delineation of seasons

Seasons were defined following Gasol *et al.* [36] with a small modification: months

with water temperature (at the sampling time) >17 °C and daylength >14 h d^{-1} were

considered to be summer. Months with water temperature <17 °C and < 11 h d^{-1} of

daylength were considered to be winter. Months with water temperature >17°C and

daylength <14 h d^{-1} were considered as autumn, while months with water temperature

<17°C and > 11 h d^{-1} of daylength were considered to be spring. The indicator value

[82] was calculated using the R package *labdsv* [83] to infer OTU seasonal preference.

Core microbiota delineated using networks

OTUs from the resident microbiota together with the 15 environmental variables were

used to construct association networks using extended Local Similarity Analysis

(eLSA) [84-86]. eLSA was run on the OTU table with subsampled reads with default

normalization: a z-score transformation using the median and median absolute

deviation. P-value estimations were run under a mixed model that performs a random

permutation test of a co-occurrence only if the theoretical p-values for the comparison

are <0.05. Bonferroni false discovery rate (q) was calculated for all edges based on the

p-values using the *p.adjust* package in R.

To detect environmentally-driven associations between OTUs induced by the

measured environmental variables we used the program EnDED [87].

Environmentally-driven associations indicate similar or different environmental

preferences between OTUs and not ecological interactions. In short, EnDED evaluates

associations between two OTUs that are both connected to the same environmental

variable based on a combination of four methods: *Sign Pattern*, *Overlap, Interaction*

Information, and *Data Processing Inequality.* These methods use the sign (positive or

negative) and the duration of the association, the relative abundance of OTUs as well

as environmental parameters to determine if an association is environmentally-driven.

If the four methods agreed that an association was environmentally-driven, then it was

removed from the network. The initial number of edges was 199,937, of which 180,345

were OTU-OTU edges that were at least in one triplet with an environmental parameter.

In total 65,280 (~33%) edges in the network were identified as indirect by EnDED and

removed. Afterward, only edges representing the strongest associations (i.e., absolute

local similarity score $|LS| > 0.7$, Spearman correlation $|\rho| > 0.7$, P<0.001 and Q<0.001)

were retained for downstream analysis and are hereafter referred to as "core

associations". Those OTUs participating in core associations were defined as core

OTUs, although their involvement in ecological interactions need further experimental validation. Both core associations and core OTUs constitute the "core network", which also represents the core microbiota (both "core network" and "core microbiota" are used indistinctively). The core network was randomized using the Erdős–Rényi model [88], using 262 nodes and 1,411 edges.

For the core network, we calculated: 1) *Density:* quantifies the proportion of actual network connections out of the total number of possible connections, 2) *Transitivity* or *Clustering coefficient:* measures the probability that nodes connected to a node are also connected, forming tight clusters, 3) *Average path length:* mean number of steps (edges) along the shortest paths for all possible pairs of nodes in the network (a low average path length indicates that most species in the network are connected through a few intermediate species), 4) *Degree:* number of associations per node, 5) *Betweenness centrality:* measures how often an OTU (node) appears on the shortest paths between other OTUs in the network, 6) *Closeness centrality:* indicates how close a node is to all other nodes in a network, 7) *Cliques:* refers to sets of interconnected nodes where all possible connections are realized, 8) *Modularity:* measures the division of a given network into modules (that is, groups of OTUs that are highly interconnected between themselves).

The Degree, Betweenness centrality and Closeness centrality were used to identify central OTUs using *ad hoc* definitions. "Hub" OTUs were those with a score above the average for the three statistics and were normally among the top 25% in each score [22, 62, 89]. Specifically, hub OTUs featured a degree >24, Betweenness centrality >0.03 and Closeness centrality >0.3. Similarly, "connector" OTUs were defined as those featuring a relatively low degree and high centrality and could be seen as elements that connect different regions of a network or modules [50]. Connector

OTUs featured a degree <5, Betweenness centrality > 0.03 and Closeness centrality >0.2. Network statistics were calculated with *igraph* in R [90] , Gephi [91] and Cytoscape v3.6.1 [92]. Visualizations were made in Cytoscape v3.6.1. Modules in the core network were identified with MCODE [93].

DECLARATIONS

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

DNA sequences and metadata are publicly available at the European Nucleotide Archive (http://www.ebi.ac.uk/ena; accession numbers PRJEB23788 [18S rRNA genes] & PRJEB38773 [16S rRNA genes]).

Competing interests

The authors declare that they have no competing interests

Funding

RL was supported by a Ramón y Cajal fellowship (RYC-2013-12554, MINECO, Spain). IMD was supported by an ITN-SINGEK fellowship (ESR2-EU-H2020-MSCA-

ITN-2015, Grant Agreement 675752 [ESR2] to RL). This work was supported by the projects INTERACTOMICS (CTM2015-69936-P, MINECO, Spain to RL), MicroEcoSystems (240904, RCN, Norway to RL), MINIME (PID2019-105775RB-I00, AEI, Spain, to RL), ALLFLAGS (CTM2016-75083-R, MINECO to RM), MIAU (RTI2018-101025-B-I00, to JMG) and DEVOTES (grant agreement n° 308392, European Union to EG). It was further supported by Grup Consolidat de Recerca 2017SGR/1568 (Generalitat de Catalunya).

Authors' contributions

AKK & RL designed the study. JMG, RM organized sampling. VB, CRG & IF collected samples, extracted the DNA, and organized its sequencing. AKK, RL & ID analyzed the data, while JMG, RM, IF, CRG & EG, provided contextual ecological or environmental pre-processed data. AKK, MFMB & RL interpreted the results. AKK & RL wrote the manuscript. All authors contributed substantially to manuscript revisions. All authors read and approved the final manuscript.

Acknowledgements

We thank all members of the Blanes Bay Microbial Observatory sampling and analyses team. Bioinformatics analyses were performed at the MARBITS platform of the Institut de Ciències del Mar (ICM; http://marbits.icm.csic.es).

REFERENCES

1. Gitay H, Wilson JB, Lee WG. Species Redundancy: A Redundant Concept? Journal of Ecology. 1996; 84(1):121-124.
2. Louca S, Parfrey LW, Doebeli M. Decoupling function and taxonomy in the global ocean microbiome. Science. 2016; 353:1272-1277.

3.   Galand PE, Pereira O, Hochart C, Auguet JC, Debroas D. A strong link between
marine microbial community composition and function challenges the idea of
functional redundancy. ISME J. 2018; 12(10):2470-2478.

4.   Shade A, Handelsman J. Beyond the Venn diagram: the hunt for a core
microbiome. Environ Microbiol. 2012; 14(1):4-12.

5.   Little AEF, Robinson CJ, Peterson SB, Raffa KF, Handelsman J. Rules of
Engagement: Interspecies Interactions that Regulate Microbial Communities.
Annual Review of Microbiology. 2008; 62(1):375-401.

6.   Lennon JT, Jones SE. Microbial seed banks: the ecological and evolutionary
implications of dormancy. Nat Rev Microbiol. 2011; 9(2):119-130.

7.   Singer E, Wagner M, Woyke T. Capturing the genetic makeup of the active
microbiome in situ. ISME J. 2017; 11(9):1949-1963.

8.   Mestre M, Höfer J. The Microbial Conveyor Belt: Connecting the Globe
through Dispersion and Dormancy. Trends Microbiol. 2021; in press.

9.   Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JI.
The Human Microbiome Project. Nature. 2007; 449(7164):804-810.

10.   Wirth R, Kádár G, Kakuk B, Maróti G, Bagi Z, Szilágyi Á, Rákhely G, Horváth
J, Kovács KL. The Planktonic Core Microbiome and Core Functions in the
Cattle Rumen by Next Generation Sequencing. Front Microbiol. 2018; 9(2285).

11.   Rubio-Portillo E, Kersting DK, Linares C, Ramos-Esplá AA, Antón J.
Biogeographic Differences in the Microbiome and Pathobiome of the Coral
Cladocora caespitosa in the Western Mediterranean Sea. Front Microbiol.
2018; 9(22).

12.   Sweet MJ, Bulling MT. On the Importance of the Microbiome and Pathobiome
in Coral Health and Disease. Frontiers in Marine Science. 2017; 4(9).

13.   Lurgi M, Thomas T, Wemheuer B, Webster NS, Montoya JM. Modularity and
predicted functions of the global sponge-microbiome network. Nature
communications. 2019; 10(1):992.

14.   Björk JR, O'Hara RB, Ribes M, Coma R, Montoya JM. The dynamic core
microbiome: Structure, dynamics and stability. bioRxiv. 2018.

15.   Delgado-Baquerizo M, Oliverio AM, Brewer TE, Benavent-González A,
Eldridge DJ, Bardgett RD, Maestre FT, Singh BK, Fierer N. A global atlas of
the dominant bacteria found in soil. Science. 2018; 359:320-325.

16.   Logares R, Deutschmann IM, Junger PC, Giner CR, Krabberød AK, Schmidt
TSB, Rubinat-Ripoll L, Mestre M, Salazar G, Ruiz-González C *et al*.
Disentangling the mechanisms shaping the surface ocean microbiota.
Microbiome. 2020; 8(1):55.

17.   Gilbert JA, Steele JA, Caporaso JG, Steinbruck L, Reeder J, Temperton B, Huse
S, McHardy AC, Knight R, Joint I *et al*. Defining seasonal marine microbial
community dynamics. ISME J. 2012; 6(2):298-308.

18.   Chow CE, Sachdeva R, Cram JA, Steele JA, Needham DM, Patel A, Parada
AE, Fuhrman JA. Temporal variability and coherence of euphotic zone bacterial
communities over a decade in the Southern California Bight. ISME J. 2013;
7(12):2259-2273.

19.   Worden AZ, Follows MJ, Giovannoni SJ, Wilken S, Zimmerman AE, Keeling
PJ. Rethinking the marine carbon cycle: factoring in the multifarious lifestyles
of microbes. Science. 2015; 347(6223):1257594.

20.   Krabberød AK, Bjorbækmo MFM, Shalchian-Tabrizi K, Logares R. Exploring
the oceanic microeukaryotic interactome with metaomics approaches. Aquat
Microb Ecol. 2017; 79(1):1-12.

21. Bjorbaekmo MFM, Evenstad A, Rosaeg LL, Krabberod AK, Logares R. The
planktonic protist interactome: where do we stand after a century of research?
ISME J. 2020; 14(2):544-559.
22. Röttjers L, Faust K. From hairballs to hypotheses - biological insights from
microbial networks. FEMS Microbiol Rev. 2018; 42(6):761-780.
23. Chow CE, Kim DY, Sachdeva R, Caron DA, Fuhrman JA. Top-down controls
on bacterial community structure: microbial network analysis of bacteria, T4-
like viruses and protists. ISME J. 2014.
24. Layeghifard M, Hwang DM, Guttman DS. Disentangling Interactions in the
Microbiome: A Network Perspective. Trends Microbiol. 2017; 25(3):217-228.
25. Fuhrman JA, Cram JA, Needham DM. Marine microbial community dynamics
and their ecological interpretation. Nat Rev Microbiol. 2015; 13(3):133-146.
26. Lima-Mendez G, Faust K, Henry N, Decelle J, Colin S, Carcillo F, Chaffron S,
Ignacio-Espinosa JC, Roux S, Vincent F *et al*. Determinants of community
structure in the global plankton interactome. Science. 2015;
348(6237):1262073.
27. Ponisio LC, Valdovinos FS, Allhoff KT, Gaiarsa MP, Barner A, Guimarães PR,
Hembry DH, Morrison B, Gillespie R. A Network Perspective for Community
Assembly. Frontiers in Ecology and Evolution. 2019; 7(103).
28. Chaffron S, Rehrauer H, Pernthaler J, von Mering C. A global network of
coexisting microbes from environmental and whole-genome sequence data.
Genome Res. 2010; 20(7):947-959.
29. Krabberod AK, Bjorbaekmo MFM, Shalchian-Tabrizi K, Logares R. Exploring
the oceanic microeukaryotic interactome with metaomics approaches. Aquat
Microb Ecol. 2017; 79(1):1-12.
30. Cram JA, Xia LC, Needham DM, Sachdeva R, Sun F, Fuhrman JA. Cross-depth
analysis of marine bacterial networks suggests downward propagation of
temporal changes. ISME J. 2015; 9(12):2573-2586.
31. Steele JA, Countway PD, Xia L, Vigil PD, Beman JM, Kim DY, Chow CE,
Sachdeva R, Jones AC, Schwalbach MS *et al*. Marine bacterial, archaeal and
protistan association networks reveal ecological linkages. ISME J. 2011;
5(9):1414-1425.
32. Needham DM, Fuhrman JA. Pronounced daily succession of phytoplankton,
archaea and bacteria following a spring bloom. Nat Microbiol. 2016;
1(4):16005.
33. Deutschmann IM, Lima-Mendez G, Krabberød AK, Raes J, Vallina SM, Faust
K, Logares R. Disentangling environmental effects in microbial association
networks. preprint. 2021.
34. Giner CR, Balague V, Krabberod AK, Ferrera I, Rene A, Garces E, Gasol JM,
Logares R, Massana R. Quantifying long-term recurrence in planktonic
microbial eukaryotes. Mol Ecol. 2019; 28(5):923-935.
35. Alonso-Saez L, Balague V, Sa EL, Sanchez O, Gonzalez JM, Pinhassi J,
Massana R, Pernthaler J, Pedros-Alio C, Gasol JM. Seasonality in bacterial
diversity in north-west Mediterranean coastal waters: assessment through clone
libraries, fingerprinting and FISH. FEMS Microbiol Ecol. 2007; 60(1):98-112.
36. Gasol JM, Cardelus C, Moran XAG, Balague V, Forn I, Marrase C, Massana
R, Pedros-Alio C, Sala MM, Simo R *et al*. Seasonal patterns in phytoplankton
photosynthetic parameters and primary production at a coastal NW
Mediterranean site. Sci Mar. 2016; 80(S1):63-77.

37. Watts DJ, Strogatz SH. Collective dynamics of 'small-world' networks. Nature.
1998; 393(6684):440-442.
38. Pedrós-Alió C. The rare bacterial biosphere. Ann Rev Mar Sci. 2012; 4:449-
466.
39. Mestre M, Höfer J, Sala MM, Gasol JM. Seasonal Variation of Bacterial
Diversity Along the Marine Particulate Matter Continuum. Front Microbiol.
2020; 11(1590).
40. Auladell A, Sánchez P, Sánchez O, Gasol JM, Ferrera I. Long-term seasonal
and interannual variability of marine aerobic anoxygenic photoheterotrophic
bacteria. ISME J. 2019; 13(8):1975-1987.
41. Berdjeb L, Parada A, Needham DM, Fuhrman JA. Short-term dynamics and
interactions of marine protist communities during the spring-summer transition.
ISME J. 2018; 12:1907–1917.
42. Lambert S, Tragin M, Lozano JC, Ghiglione JF, Vaulot D, Bouget FY, Galand
PE. Rhythmicity of coastal marine picoeukaryotes, bacteria and archaea despite
irregular environmental perturbations. ISME J. 2019; 13(2):388-401.
43. Charles F, Lantoine F, Brugel S, Chrétiennot-Dinet M-J, Quiroga I, Rivière B.
Seasonal survey of the phytoplankton biomass, composition and production in
a littoral NW Mediterranean site, with special emphasis on the picoplanktonic
contribution. Estuarine, Coastal and Shelf Science. 2005; 65(1):199-212.
44. Milici M, Deng Z-L, Tomasch J, Decelle J, Wos-Oxley ML, Wang H, Jáuregui
R, Plumeier I, Giebel H-A, Badewien TH *et al*. Co-occurrence Analysis of
Microbial Taxa in the Atlantic Ocean Reveals High Connectivity in the Free-
Living Bacterioplankton. Front Microbiol. 2016; 7(649).
45. Newman M. Networks: Oxford University Press; 2018.
46. Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: Networks,
competition, and stability. Science. 2015; 350:663-666.
47. Vellend M. The theory of ecological communities. Princeton: Princeton
University Press; 2016.
48. Stouffer DB, Bascompte J. Compartmentalization increases food-web
persistence. Proceedings of the National Academy of Sciences. 2011; 108:3648-
3652.
49. Krause AE, Frank KA, Mason DM, Ulanowicz RE, Taylor WW. Compartments
revealed in food-web structure. Nature. 2003; 426(6964):282-285.
50. Olesen JM, Bascompte J, Dupont YL, Jordano P. The modularity of pollination
networks. Proceedings of the National Academy of Sciences. 2007; 104:19891-
19896.
51. Medeiros LP, Garcia G, Thompson JN, Guimarães PR. The geographic mosaic
of coevolution in mutualistic networks. Proceedings of the National Academy
of Sciences. 2018; 115:12017-12022.
52. Tripp HJ, Schwalbach MS, Meyer MM, Kitner JB, Breaker RR, Giovannoni SJ.
Unique glycine-activated riboswitch linked to glycine–serine auxotrophy in
SAR11. Environ Microbiol. 2009; 11(1):230-238.
53. Carini P, Steindler L, Beszteri S, Giovannoni SJ. Nutrient requirements for
growth of the extreme oligotroph 'Candidatus Pelagibacter ubique'
HTCC1062 on a defined medium. ISME J. 2013; 7(3):592-602.
54. Carini P, Campbell EO, Morré J, Sañudo-Wilhelmy SA, Cameron Thrash J,
Bennett SE, Temperton B, Begley T, Giovannoni SJ. Discovery of a SAR11
growth requirement for thiamin's pyrimidine precursor and its distribution in
the Sargasso Sea. ISME J. 2014; 8(8):1727-1738.

55. Włodarczyk A, Selão TT, Norling B, Nixon PJ. Newly discovered *Synechococcus* sp. PCC 11901 is a robust cyanobacterial strain for high biomass production. Communications Biology. 2020; 3(1):215.

56. Becker JW, Hogle SL, Rosendo K, Chisholm SW. Co-culture and biogeography of *Prochlorococcus* and SAR11. ISME J. 2019; 13(6):1506-1519.

57. Chafee M, Fernàndez-Guerra A, Buttigieg PL, Gerdts G, Eren AM, Teeling H, Amann RI. Recurrent patterns of microdiversity in a temperate coastal marine environment. ISME J. 2018; 12(1):237-252.

58. Paine RT. A Note on Trophic Complexity and Community Stability. The American Naturalist. 1969; 103(929):91-93.

59. Banerjee S, Schlaeppi K, van der Heijden MGA. Keystone taxa as drivers of microbiome structure and functioning. Nat Rev Microbiol. 2018; 16:567-576.

60. Berry D, Widder S. Deciphering microbial interactions and detecting keystone species with co-occurrence networks. Front Microbiol. 2014; 5(MAY):219.

61. Freilich MA, Wieters E, Broitman BR, Marquet PA, Navarrete SA. Species co-occurrence networks: Can they reveal trophic and non-trophic interactions in ecological communities? Ecology. 2018; 99(3):690-699.

62. Banerjee S, Kirkby CA, Schmutter D, Bissett A, Kirkegaard JA, Richardson AE. Network analysis reveals functional redundancy and keystone taxa amongst bacterial and fungal communities during organic matter decomposition in an arable soil. Soil Biology and Biochemistry. 2016; 97:188-198.

63. Pester M, Bittner N, Deevong P, Wagner M, Loy A. A 'rare biosphere' microorganism contributes to sulfate reduction in a peatland. ISME J. 2010; 4(12):1591-1602.

64. Zhu F, Massana R, Not F, Marie D, Vaulot D. Mapping of picoeucaryotes in marine ecosystems with quantitative PCR of the 18S rRNA gene. FEMS Microbiol Ecol. 2005; 52(1):79-92.

65. Seymour JR, Amin SA, Raina JB, Stocker R. Zooming in on the phycosphere: the ecological interface for phytoplankton-bacteria relationships. Nat Microbiol. 2017; 2:17065.

66. Farnelid HM, Turk-Kubo KA, Zehr JP. Identification of Associations between Bacterioplankton and Photosynthetic Picoeukaryotes in Coastal Waters. Front Microbiol. 2016; 7(339).

67. Auladell A, Barberán A, Logares R, Garcés E, Gasol JM, Ferrera I. Seasonal niche differentiation between evolutionary closely related marine bacteria. bioRxiv. 2020.

68. Grasshoff K, Kremling K, Ehrhardt M. Methods of Seawater Analysis: Third, Completely Revised and Extended Edition; 2007.

69. Massana R, Castresana J, Balague V, Guillou L, Romari K, Groisillier A, Valentin K, Pedros-Alio C. Phylogenetic and ecological analysis of novel marine stramenopiles. Appl Environ Microbiol. 2004; 70(6):3528-3534.

70. Stoeck T, Bass D, Nebel M, Christen R, Jones MD, Breiner HW, Richards TA. Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. Mol Ecol. 2010; 19 Suppl 1(SUPPL. 1):21-31.

71. Herlemann DPR, Labrenz M, Jürgens K, Bertilsson S, Waniek JJ, Andersson AF. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. ISME J. 2011; 5(10):1571-1579.

| 1181 | 72. | Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb Ecol. 2015; 75(2):129-137. |

1181   72.   Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU
1182   rRNA 806R gene primer greatly increases detection of SAR11
1183   bacterioplankton. Aquat Microb Ecol. 2015; 75(2):129-137.

1184   73.   Martin M. Cutadapt removes adapter sequences from high-throughput
1185   sequencing reads. EMBnet Journal. 2011; 17(1):10-12.

1186   74.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP.
1187   DADA2: High-resolution sample inference from Illumina amplicon data. Nat
1188   Methods. 2016; 13(7):581-583.

1189   75.   Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid
1190   assignment of rRNA sequences into the new bacterial taxonomy. Appl Environ
1191   Microbiol. 2007; 73(16):5261-5267.

1192   76.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J,
1193   Glockner FO. The SILVA ribosomal RNA gene database project: improved
1194   data processing and web-based tools. Nucleic Acids Res. 2013; 41(Database
1195   issue):D590-596.

1196   77.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment
1197   search tool. Journal of molecular biology. 1990; 215(3):403-410.

1198   78.   Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, Boutte C, Burgaud
1199   G, de Vargas C, Decelle J *et al*. The Protist Ribosomal Reference database
1200   (PR2): a catalog of unicellular eukaryote small sub-unit rRNA sequences with
1201   curated taxonomy. Nucleic Acids Res. 2013; 41(Database issue):D597-604.

1202   79.   McNichol J, Berube PM, Biller SJ, Fuhrman JA. Evaluating and Improving
1203   SSU rRNA PCR Primer Coverage via Metagenomes from Global Ocean
1204   Surveys. bioRxiv. 2020.

1205   80.   Oksanen J, Guillaume Blanchet FFM, Kindt R, Legendre P, McGlinn D,
1206   Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH *et al*. vegan:
1207   Community Ecology Package. R package. In.; 2016.

1208   81.   Moritz S. imputeTS: Time Series Missing Value Imputation. In.; 2017.

1209   82.   Dufrêne M, Legendre P. Species assemblages and indicator species: The need
1210   for a flexible asymmetrical approach. Ecological Monographs. 1997.

1211   83.   Roberts DW. labdsv: Ordination and Multivariate Analysis for Ecology. R
1212   package version 1.8-0. In.; 2016.

1213   84.   Ruan Q, Dutta D, Schwalbach MS, Steele JA, Fuhrman JA, Sun F. Local
1214   similarity analysis reveals unique associations among marine bacterioplankton
1215   species and environmental factors. Bioinformatics. 2006; 22(20):2532-2538.

1216   85.   Xia LC, Ai D, Cram JA, Liang X, Fuhrman JA, Sun F. Statistical significance
1217   approximation in local trend analysis of high-throughput time-series data using
1218   the theory of Markov chains. BMC Bioinformatics. 2015; 16(1):301.

1219   86.   Xia LC, Ai D, Cram J, Fuhrman JA, Sun F. Efficient statistical significance
1220   approximation for local similarity analysis of high-throughput time series data.
1221   Bioinformatics. 2013; 29(2):230-237.

1222   87.   Deutschmann IM, Lima-Mendez G, Krabberød AK, Raes J, Vallina SM, Faust
1223   K, Logares R. Disentangling environmental effects in microbial association
1224   networks. ResearchSquare. 2020.

1225   88.   Erdős P, Rényi A. On random graphs. Publicationes Mathematicae. 1959;
1226   6:290–297.

1227   89.   Banerjee S, Baah-Acheamfour M, Carlyle CN, Bissett A, Richardson AE,
1228   Siddique T, Bork EW, Chang SX. Determinants of bacterial communities in
1229   Canadian agroforestry systems. Environ Microbiol. 2016; 18(6):1805-1816.

1230   90.   Yu G, Chen Y-s, Guo Y-c. Design of integrated system for heterogeneous
1231   network query terminal. Journal of Computer Applications. 2009; 29(8):2191-
1232   2193.
1233   91.   Bastian M, Heymann S, Jacomy M. Gephi: an open source software for
1234   exploring and manipulating networks. In: *International AAAI Conference on*
1235   *Weblogs and Social Media*. 2009.
1236   92.   Smoot ME, Ono K, Ruscheinski J, Wang PL, Ideker T. Cytoscape 2.8: New
1237   features for data integration and network visualization. Bioinformatics. 2011.
1238   93.   Bader GD, Hogue CW. An automated method for finding molecular complexes
1239   in large protein interaction networks. BMC Bioinformatics. 2003; 4(1):2.
1240
1241

Supplementary Figures

Supplementary Figure 1. Panel A shows the full network constructed with the resident microbiota (that is, OTUs present in >30% of the samples over 10 years; Table 1). **Panel B** displays network elements that were removed as they did not fulfill the cut-offs (that is, highly significant correlations (P & Q <0.001), local similarity scores >|0.7| and Spearman correlations >|0.7|).

1253

1254
1255

1256
1257 **Supplementary Figure 2.** OTU relative abundance vs. degree shows no relationship
1258 in the core network.