Transcriptomic analysis of temporal shifts in berry development

between two grapevine cultivars of the Pinot family reveals potential

genes controlling ripening time

Jens Theine[1], Daniela Holtgräwe[1], Katja Herzog[2], Florian Schwander[2], Anna Kicherer[2], Ludger

Hausmann[2], Prisca Viehöver[1], Reinhard Töpfer[2], Bernd Weisshaar[1]*

[1] Bielefeld University, Genetics and Genomics of Plants, Faculty of Biology & Center for

Biotechnology, Bielefeld

[2] Julius Kühn-Institute, Institute for Grapevine Breeding Geilweilerhof, Siebeldingen

*corresponding author:
BW Bernd Weisshaar (0000-0002-7635-3473)
14
JT Jens Theine (0000-0002-6042-373X)
DH Daniela Holtgräwe (0000-0002-1062-4576)
KH Katja Herzog
FS Florian Schwander
AK Anna Kicherer
LH Ludger Hausmann (0000-0002-4046-9626)
PV Prisca Viehöver
RT Reinhard Töpfer (0000-0003-1569-2495)
23

Abstract

Background

Grapevine cultivars of the Pinot family represent clonally propagated mutants with major

phenotypic and physiological differences, such as different colour or shifted ripening time, as

well as changes in important viticultural traits. Specifically, the cultivars 'Pinot Noir' (PN) and

'Pinot Noir Precoce' (PNP, early ripening) flower at the same time, but vary in the beginning

of berry ripening (veraison) and, consequently, harvest time. In addition to genotype,

seasonal climatic conditions (i.e. high temperatures) also affect ripening times. To reveal

possible regulatory genes that affect the timing of veraison onset, we investigated

differences in gene expression profiles between PN and PNP throughout berry development

with a closely meshed time series and over two separate years.

Results

The difference in the duration of berry formation between PN and PNP was quantified to be

approximately two weeks under the growth conditions applied, using plant material with a

proven PN and PNP clonal relationship. Clusters of co-expressed genes and differentially

expressed genes (DEGs) were detected which reflect the shift in the timing of veraison onset.

Functional annotation of these DEGs fit to observed phenotypic and physiological changes

during berry development. In total, we observed 3,342 DEGs in 2014 and 2,745 DEGs in 2017

between PN and PNP, with 1,923 DEGs across both years. Among these, 388 DEGs were

identified as veraison-specific and 12 were considered as berry ripening time regulatory

candidates. The expression profiles revealed two candidate genes for ripening time control

which we designated *VviRTIC1* and *VviRTIC2* (VIT_210s0071g01145 and

VIT_200s0366g00020, respectively). These genes likely contribute the phenotypic

differences observed between PN and PNP.

Conclusions

Many of the 1,923 DEGs show highly similar expression profiles in both cultivars if the

patterns are aligned according to developmental stage. In our work, putative genes

differentially expressed between PNP and PN which could control ripening time as well as

veraison-specific genes were identified. We point out connections of these genes to

molecular events during berry development and discuss potential candidate genes which

may control ripening time. Two of these candidates were observed to be differentially

expressed in the early berry development phase. Several down-regulated genes during berry

ripening are annotated as auxin response factors / ARFs. Conceivably, general changes in

auxin signaling may cause the earlier ripening phenotype of PNP.

Key words

Vitis vinifera, Pinot Noir, Pinot Noir Precoce, grapevine, berry ripening, fruit development,

differential gene expression, transcriptome profiling, ripening time control, veraison

Background

Vitis vinifera subsp. vinifera (grapevine) belongs to the family *Vitaceae*. With 6,000 to 11,000

cultivars, it is one of the most important perennial crops worldwide [1]. Grapevine fruit

development can be divided into two physiological phases, berry formation and berry

ripening. Veraison refers to the transition from berry formation to berry ripening, and each

of the two phases is represented by a sigmoidal growth curve of development [2]. The

progress through development is described by stages referred to as "BBCH stages" (acronym

derived from the names of the coordinating institutions involved in stage definition) that

have been defined for several crops including grapevine [3, 4]. The first physiological phase is

described as berry formation (berry initiation and growth with cell divisions) and lasts from

the end of flowering (BBCH71) until ~60 days later when the majority of berries are touching

each other (BBCH79). The developmental stage of veraison (BBCH81) is the end of berry

formation and the start of berry ripening [2]. Phenotypically, veraison is the developmental

switch when the berries start to soften, accompanied by the onset of accumulation of

phenylpropanoids. In red grapevine cultivars, veraison is also indicated by a colour change of

the berries that is caused by the beginning of accumulation of anthocyanins, a major class of

phenylpropanoids. Members of the well-studied protein superfamily of R2R3-MYB

transcription factors (TFs) are considered to be mainly accountable for controlling

anthocyanin accumulation [5-7]. Berry ripening starts at veraison and continues until harvest

(BBCH89), this phase includes cell enlargement, sugar accumulation and acidity decline.

Timing of veraison has also been studied at the level of genetic loci and genomic regions that

control this trait. Since it is a quantitative trait influenced by several to many genetic loci,

quantitative trait locus (QTL) analyses have been performed. These studies detected a major

QTL for timing of the onset of veraison on chromosome 16, combined with a minor QTL on

chromosome 18 [8]. By integrating a number of QTL studies, several meta-QTLs connected

to genetic control of veraison time were detected, with the most relevant located on

chromosome 14, 16, and 18 [9].

Anthropogenic climate change is resulting in successively earlier ripening of grapes with a

significant impact on berry quality and consequently the expected flavours of a desired wine

style [10]. In addition, the time of veraison and harvest of a given cultivar may differ greatly,

driven by regional and/or year-specific differences in weather conditions. Obviously, this

calls for a better molecular understanding of the control of ripening time in grapevine.

Comparison of different grapevine cultivars grown at the same environmental conditions

often uncovers differences in the duration of berry formation, timing of veraison, duration of

berry ripening, and ripening time in general. However, the underlying genetic factors are

mostly unknown. Previous studies have elucidated how ripening time is affected by internal

and external factors. For example, the effect of phytohormones on berry ripening has been

widely studied [1]. In general, fruit growth is discussed to be controlled by several

phytohormones which play essential roles to trigger or delay ripening processes [11].

Grapevine is a non-climacteric fruit and effects of abscisic acid (ABA) have been investigated

in many studies as ABA is considered to trigger ripening [12-14]. Furthermore, it was shown

that ABA is involved in controlling leaf senescence [15], responses to drought [16] and

pathogen defense [17]. In grapevine, although not as central as in climacteric fruits like

tomato (*Solanum lycopersicum*), the phytohormone ethylene is involved in the control of

berry ripening [1, 13, 18, 19], while auxin has been shown to induce a delay of ripening [20,

21].

Fruit development of both, dry and fleshy fruits, has been studied very intensively for the

obvious reason that fruits are central to human nutrition [22, 23]. The main model system

for studies on fleshy fruits is tomato, because of established genetics and molecular biology,

access to mutants, and well advanced transgenic approaches to gene function identification

[24, 25]. Berry development of grapevines has also been studied intensively [1, 26] and often

at the level of the transcriptome. In quite some of the studies, predominantly late berry

development stages were sampled to bring the development stage of veraison into the focus

[9, 27-29]. In addition, whole berry development was studied with coarse time point

distribution [30-34].

To monitor gene expression changes at a high resolution throughout grapevine berry

development, starting from flowering until berries are matured, we sampled a

comprehensive time series from two Pinot cultivars across two years. The samples were

collected from the grapevine cultivar 'Pinot Noir' (PN) and the comparably earlier ripening

cultivar 'Pinot Noir Precoce' (PNP) that is expected to be closely related to PN. The cultivar

PNP is listed in the *Vitis* International Variety Catalogue (*V*IVC; [35]) and described to flower

at the same time as PN but to reach veraison significantly earlier than PN [36]. Quantitative

data for transcript levels, interpreted as values for gene expression, were generated by RNA-

Seq. We studied the general course of gene expression patterns throughout berry

development in both years and cultivars, and identified a number of differentially expressed

genes (DGEs) between PN and PNP prior to veraison. These DEGs can be considered as

important candidates for either delaying or pushing forward berry development. Our main

aim was the identification of genes controlling the speed of development, to offer an entry

point into characterization of the relevant molecular functions in grapevine, and to facilitate

future breeding strategies that address traits relevant to, and affected by, climate change.

Results

Phenotypical comparison between two Pinot cultivars

To study ripening shifts, we used samples of two closely related grapevine cultivars. The

cultivar PNP is an earlier ripening clonal variant of its ancestor PN. Clonal relation of PN and

PNP was confirmed by a set of 24 SSR markers that all displayed the identical allele status for

both cultivars (Additional file 1: Table S1). To confirm and validate the phenotypic

differences between PN and PNP, detailed BBCH developmental stages were determined

and documented (Figure 1). PN and PNP display similar phenotypic properties during

development and flower (BBCH65) at the same time. However, veraison (BBCH81) is shifted

to ~2 weeks earlier for PNP, and similar shifts were observed in four different years

(Table 1). In addition, Figure 1A shows an overview across the time points at which samples

were taken. The phenotypic differences between PN and PNP are illustrated in images of

developing berries taken between onset of berry formation and veraison (Figure 1B and

Additional file 1: Table S2 and Table S3). Veraison (BBCH81) is visible as the onset of

anthocyanin accumulation and is detected ~2 weeks later in PN compared to PNP.

Figure 1: Phenotypical observations and sampling scheme. (A) Sampling time points and days after onset of flowering (DAF) are indicated in red. The developmental stage observed is shown in the BBCH stages [3, 4]. DAF zero (0) is set at BBCH61 (onset of flowering, 10% of flowerhoods fallen [3]). Berry development is depicted schematically and categorized into the phases flowering (yellow), berry formation (green), and berry ripening (purple) for both cultivars. The junction between green and purple indicates veraison (BBCH81). To orient for

time of year, numbered days after January 1st are shown. (B) Images of grape bunches and

developing berries taken in 2014 are shown to document the differences between PN and

PNP. Images were taken 35, 41, 49 and 56 DAF. Scale bar: 50 mm.

Table 1: Observed flowering- and berry development shifts between the cultivars PNP and

PN in 2014, 2015, 2016 and 2017 at the Geilweilerhof vineyards, Siebeldingen, Germany (in

days after January 1st).

Year	Cultivar	Start of flowering (BBCH61)	End of flowering period / Start of berry formation (BBCH71)	End of berry formation / veraison (BBCH81)	Flowering time [Δ days]	Berry formation time [Δ days]
2014	PNP	155	162	196	0	19
2014	PN	155	162	215		
2015	PNP	159	166	201	7	14
2015	PN	159	173	222		
2016	PNP	171	180	215	0	14
2016	PN	171	180	229		
2017	PNP	151	163	200	0	11
2017	PN	151	163	211		

Global view of gene expression patterns

We harvested triplicate samples in 2014 and 2017 from flowering until after veraison (for

time points see Figure 1 and Additional file 1: Tables S2 and S3), individual harvests are

referred to below as subsamples, and analyzed them by RNA-Seq. After preprocessing the

raw data (see Methods), reads derived from each subsample were mapped to the reference

sequence from PN40024 and analyzed with respect to the CRIBI V2.1 annotation dataset. For

2014, approximately 19.7 million reads per subsample were obtained from each of the 72

libraries. An overall alignment rate of 79 % to the grape reference genome sequence was

reached. For 2017, approximately 43.5 million reads per subsample were obtained from

each of the 78 libraries. From these, an overall alignment rate of 92 % to the reference was

calculated. Expression values were initially detected as Transcripts Per Kilobase Million

(TPM) and averaged over the three subsamples for each sample. Considering both years

separately, a total of 28,692 genes were detected as expressed in both cultivars and in both

years. In contrast, 2,152 CRIBI V2.1 genes were found to be not expressed.

The correlation between gene expression data, determined as TPM values per sample, of the

datasets from both years over all genes was r = 0.5095 (Pearson correlation coefficient) for

PN and r = 0.6557 for PNP, respectively. For PN and PNP, 10,205 and 16,226 genes,

respectively, expression values were significantly correlated (p-value < 0.05) between the

years 2014 and 2017. A list of the correlation strength of the eight time points with the same

BBCH stage is provided in Additional file 1: Table S4.

To visualize global trends and similarity of the gene expression values obtained from all

subsamples, a Principal Component Analysis (PCA) of both years was performed with

keeping the subsamples separate. The first component PC1 explains 57% of the variance,

while the second component PC2 explains 17% (Figure 2). Almost all data points of the

subsamples (triplicates within a sample) from both years cluster near to each other. The data

follow a track of time in a nearly consecutive and continuous way. Main actors, which

influence most of the variance in the data, were genes related to cell wall modification,

secondary metabolism, wounding-response and hormone signaling. The top 500 genes

responsible for most of the variance in PC1 and PC2 are listed in Additional file 1: Table S5

and Table S6, together with functional information for each listed gene from

MapMan/Mercator and RefSeq.

Figure 2: Principle Component Analysis (PCA) of gene expression values from all subsamples.
Each data point represents a single subsample of the triplicates for each time point of both
years (2014 and 2017 as indicated by [DAF]_14 and [DAF]_17 with the colour code) and for
both cultivars (PN as triangles, PNP as circles).

Cluster analysis for identification of co-expressed genes

Gene expression time series profiles combined data from the subsamples/triplicates for each

time point across two years for both culrivars. Expression profiles were compared using the

clustering tool CLUST. The goal was the characterization of similarity and/or differences in

gene expression among years and cultivars throughout berry development. Over all four

datasets, 13 PN/PNP clusters of genes with similar gene expression patterns (C1-C13) were

obtained (Additional file 2: Figure S1A). In these clusters, 3,316 (12.2 %) of the 27,139 genes

expressed during berry development were found co-expressed among both years and

cultivars. It should be noted that CLUST uses criteria to define expressed and not expressed

genes that differ from the ones applied above (see Methods). The observed expression

profiles differ clearly between the clusters, which was in part the result of the restricted

number of clusters that CLUST extracts. Manual inspection of the clusters revealed little

deviation of individual gene expression profiles within each individual cluster for a given year

or cultivar. All cluster gene memberships, also those for the additional cultivar-specific

cluster analyses (see below), are available in Additional file 1: Table S7-S9.

The PN/PNP clusters C2, C5, C6 and C12 (C12_PN/PNP selected as example, see Figure 3)

reveal a small but detectable difference in the gene expression profile between both

sampled years, but are almost identical for both cultivars. The PN/PNP clusters C1, C7 and

C11 (C1_PN/PNP selected as example, see Figure 3) show similar expression profiles over the

two years, but stand out by shifted expression peaks that distinguish PN and PNP.

To characterize the clusters with respect to potential functions of the co-expressed genes

included in a given cluster, GO term enrichment for biological processes was calculated. The

full list of enriched GO terms for all clusters is listed in Additional file 1: Table S10-S12. Two

examples for GO terms appearing with highly significant incidence were 'response to

oxidative stress' in cluster C11_PN/PNP (term GO:0051276) and 'regulation of defense

response' in cluster C5_PN/PNP (term GO:0031347).

Two additional cluster analyses were performed, one for the PN data from both years

(Additional file 2: Figure S1B) and one for the PNP data from both years (Additional file 2:

Figure S1C). These analyses revealed a high abundance of genes from the expansin gene

family with similar expression profiles in clusters C0_PNP and C6_PN. The cluster C16_PN

was found to have a highly significant enrichment for 'vegetative to reproductive phase

transition' (GO:0010228). Cluster-gene memberships for the cultivar-specific clustering are

available in Additional file 1: Table S8-9, and the corresponding GO term enrichment is

summarized in Additional file 1: Table S11-12.

Figure 3: Two selected gene expression profile clusters with either a cultivar-specific

difference (C1_PN/PNP) or a weather/field condition-specific difference (C12_PN/PNP) after

clustering all data (both cultivars and both years). Strength of gene expression (quantile

normalization) was plotted over the time course of berry development. Sampling time points

are detailed in Figure 1 and were restricted to those eight equivalent time points at which

the cultivars display the same BBCH stage (Additional file 1: Table S4). For all PN/PNP

clusters see Additional file 2: Figure S1A. IDs of genes that make up the clusters are listed in

Additional file 1: Table S7.

Analyses of differentially expressed genes

The gene expression time series throughout berry development was analyzed for

differentially expressed genes (DEGs) between the two cultivars PN and PNP with DESeq2.

Genes with significantly differential expression were selected by using the filters adjusted

p-value (PADJ) < 0.05 and log2fold change (LFC) > 2. The results are summarized in Table 2

and are detailed at the gene level per time point compared in Additional file 1: Table S13.

Table 2: Filtering steps applied for selecting DEGs, and the number of DEGs that were carried

on after each selection step. For details see Methods.

	PN/PNP 2014 [DEGs]	PN/PNP 2017 [DEGs]
adjusted p-value (PADJ) < 0.05: (counted over all sample pairs)	8,206	4,419
log2fold change (LFC) > 2: (counted over all sample pairs)	6,629	4,298
down- / up-regulated (in PNP vs. PN):	3,293 / 3,336	2,130 / 2,168
unique: (non-redundant within time series)	3,342	2,745
intersection: (detected in both years)	1,923	
excluded due to intersection:	*1,419*	*822*
veraison-specific genes: (detected within BBCH79-81 of PNP)	388	
potentially regulatory: (detected within BBCH61-79 of PNP)	12	

In total, 8,206 and 4,419 DEGs were identified with PADJ greater than 0.05 for 2014 and

2017, respectively. Almost twice as many DEGs were initially detected in 2014 samples

compared to 2017. By applying the filter for an at least 2-fold difference in expression level

(LFC > 2), the number of significant DEGs decreased, mainly for the PN/PNP time series from

2014. Only few DEGs between PN and PNP were observed during flowering (BBCH61 to 69)

at the beginning of both time series (see Figure 4). Within the berry formation phase

(BBCH71 to 79), the number of DEGs detected increased towards veraison (BBCH81) as both

genotypes increasingly varied in physiological stages. The highest number of DEGs was

observed in parallel to the time-shifted veraison of PNP relative to PN. This time point was

also the most phenotypically different between the two cultivars (see Figures 1 and 4). A set

of veraison-specific genes was defined by selecting the DEGs from time points DAF35 and

DAF41 from 2014 that were also observed to coincide with this phenotype at DAF42 and

DAF49 in the 2017 gene expression data. These criteria identified 388 veraison-specific

DEGs. This set of 388 genes was compared to results from similar studies and found to be in

agreement (e.g. 81% [27] and 52% [28]; IDs of the 388 genes, the genes that match results

from the other studies and their functional annotation, are included in Additional file 1:

Table S14). During the subsequent phase of berry ripening (BBCH81 to 89, after around

DAF56 in 2014 and around DAF60 in 2017), the number of DEGs detected decreased.

We developed a visualization for the numbers of DEGs detected and the changes with

respect to which genes are newly appearing as differentially expressed at a given time point

(sample pair PN/PNP) in the time series (Figure 4). Groups of newly appearing DEGs are

containing only few genes early in berry formation, while numbers increase at veraison of

PNP. After veraison of PN, the number of DEGs decreased. If DEGs appearing in several time

points are counted only once, 3,342 and 2,745 unique DEGs (different genes) are detected

from 2014 and 2017, respectively (compare Table 2).

Figure 4: Visualization of the number of DEGs detected between PN and PNP in a logarithmic

scale ($\log_{10}$). Results for 2014 are shown in purple, those for 2017 in orange. The time series

from the two years were aligned at veraison of PNP; the timeline is given as days after onset

of flowering (DAF). DEGs are counted for a pair of PN/PNP samples for each time point

individually, the number above each column mentions the number of DEGs detected at the

respective time point. Groups of newly appearing DEGs relative to an earlier time point are

indicated by a new colour shade in the column (bar) for each time point. For members of a

given group of DEGs, the attributed colour shade is kept for the subsequent time points

(columns/samples). The pictures between the two column series display BBCH65 (full bloom,

50% of flowerhoods fallen [3]) and BBCH81 (veraison) of PNP and PN.

To further increase the reliability, reproducibility, and relevance of the selected DEGs, the

intersection between the DEGs identified in the two years studied was computed. In total,

1,923 unique DEGs were obtained (Table 2). To reveal DEGs potentially involved in the

control of timing of ripening, i.e. genes that might be involved in the trait that mainly

distinguishes PN and PNP, only intersecting DEGs which appeared at time points before

veraison in PNP were picked. This resulted in a list of 12 DEGs that may control ripening

time. It should be noted that these putative regulatory DEGs are supposed to be relevant

before the set of veraison-specific genes implements the phenotypic changes at veraison.

The full list of DEGs, their identity and annotation information as well as their fit to the

selection criteria on the way from all (raw) DEGs to potentially regulatory DEGs are detailed

in Additional file 1: Table S13. IDs of the 12 putative ripening time control genes, the genes

that match results from related studies (7 DEGs [27], 4 DEGs [34] and 3 DEGs [28]) and their

functional annotation are included in Additional file 1: Table S14, the most relevant data are

summarized in Table 3.

Functional classification of DEGs

To complement the gene lists with functional information from grapevine that might

potentially be informative for berry development, the 1,923 intersecting DEGs were

analyzed with respect to enrichment of genes that have been assigned to biological

pathways already established for grapevine (see Methods). For 46 of the 247 defined

grapevine pathways, significant enrichment (permuted p-value <0.1) was detected. The most

reliable predictions (permuted p-value <0.001) for pathways that might be relevant were

photosynthesis antenna proteins (vv10196; 9 DEGs); nitrogen metabolism (vv10910, 19

DEGs); phenylpropanoid biosynthesis (vv10940, 66 DEGs), tyrosine metabolism (vv10350, 33

DEGs); transport electron carriers (vv50105, 18 DEGs); phenylalanine metabolism (vv10360,

33 DEGs); brassinosteroid biosynthesis (vv10905, 8 DEGs) and flavonoid biosynthesis

(vv10941, 30 DEGs). The enrichment results are provided in (Additional file 1: Table S15). The

same analysis was also carried out for the 12 putative ripening time control (Additional file 1:

Table S16) and the 388 veraison-specific DEGs (Additional file 1: Table S17).

A check of the 1,923 intersecting DEGs revealed that 141 TF genes are included. Of these, 48

DEGs were clearly up- and 93 down-regulated at their first appearance in the time series.

The full list of TF encoding genes that were higher expressed in PNP (up-regulated), or lower

expressed in PNP (down-regulated), compared to PN, is shown in Additional file 1: Table S18.

For a more detailed view on the expression patterns of selected TF encoding genes, we

generated for the TF gene family with the highest abundance among the 141 TF genes,

namely the R2R3-MYB-type TFs with 22 cases in the MapMan functional assignment, an

expression heatmap (Additional file 2: Figure S2). As a result, *VviMYB24*

(VIT_214s0066g01090), which is related to At3g27810/*AtMYB21*, At5g40350/*AtMYB24* and

At3g01530/*AtMYB57* according to TAIR/PhyloGenes, was identified as an early appearing

DEG that showed its highest expression level at flowering (BBCH61). Prominent *R2R3-MYB*

genes known to be relevant for anthocyanin accumulation like *VviMYBA1*

(VIT_202s0033g00410), *VviMYBA2* (VIT_202s0033g00390), *VviMYBA3*

(VIT_202s0033g00450) and *VviMYBA8* (VIT_202s0033g00380) were detected as expressed

starting from veraison (BBCH81) in both cultivars and with a time shift towards earlier

expression in PNP. An additional *R2R3-MYB* gene with a similar expression pattern is

VviMYB15 (VIT_205s0049g01020). Other *R2R3-MYB* genes are expressed early during berry

formation, these include *VviMYBF1* (VIT_207s0005g01210, related to

At2g47460/*AtMYB12*/*AtPFG1*) as well as *VviMYBPA5* (VIT_209s0002g01400) and *VviMYBPA7*

(VIT_204s0008g01800, both related to At5g35550/*AtMYB123*/*AtTT2*). According to their

related expression patterns visualized in the heatmaps (Additional file 2: Figure S2), the

R2R3-MYB genes fall into three groups that roughly fit to the three phases marked in Figure

1B, namely flowering, berry formation, and berry ripening (see discussion).

Putative candidates for ripening time control genes

As mentioned above, DEGs detected in both years at time points before veraison of PNP

were selected and considered as putative genes that control ripening time (Table 3,

Additional file 1: Table S13). The VitisNet enrichment analyses performed for these 12

candidates resulted in 2 pathways that showed significant (permuted p-value < 0.05)

enrichment with two genes in the pathway: auxin signaling (vv30003 with *VviEXPA5*

(VIT_206s0004g00070) and *VviEXPA14* (VIT_213s0067g02930)) and cell wall (vv40006 with

VviPL1 (VIT_205s0051g00590) and *VviGRIP28* (VIT_216s0022g00960)); see Additional file 1:

Table S16).

A detailed check of the data presented in Figure 4, together with results for these putatively

ripening time control DEGs, resulted in the identification of two DEGs that stand out from

the whole list of DEGs. Both genes almost completely lack expression in the early ripening

cultivar PNP while there is clear expression in PN. Therefore, these two genes were detected

as DEGs throughout the whole time series in both years. The first of the two, designated

VviRTIC1 for "Ripening Time Control" (VIT_210s0071g01145, encoding a protein similar to

"protein of unknown function DUF789"), is expressed during flowering (BBCH61 - 65) and is

more or less continuously down-regulated over time in PN (Figure 5A). The second of the

two, designated *VviRTIC2* (VIT_200s0366g00020, encoding a protein similar to "cysteine-rich

receptor-like protein kinase"), displays expression in PN during berry formation as well as

during berry ripening with a peak before veraison in 2017 (Figure 5B). The expression

patterns of the 10 remaining DEGs of the putative ripening time control gene set are shown

in Additional file 2: Figure S3.

Data on the set of 12 putative ripening time controlling genes are collected in Table 3, with a

focus on relative up- or downregulation of expression before veraison of PNP. Also, a short

description of the differences of the expression patterns before, at and after veraison of PN

and PNP is included.

Table 3: Collection of features of the set of 12 genes classified as potential regulators based

on their differential expression before veraison of PNP. The detailed expression patterns are

shown in Figure 5 and Additional file 2: Figure S3. Details on annotation are listed in

Additional file 1: Table S13.

gene ID	gene symbol	functional annotation (transferred via BLASTp)	relative transcriptional change	
			pre veraison of PNP	before, at or after veraison of PN & PNP
VIT_205s0051g00590	VviPL1	pectate lyase 8	upregulated in PNP	peaks before veraison, goes down after veraison
VIT_205s0077g01980	-	uncharacterized protein LOC100248252	upregulated in PNP	peaks before veraison, stays up after veraison
VIT_206s0004g00070	VviEXPA5	expansin A10	upregulated in PNP	upregulated in PN, strong peak before veraison
VIT_206s0009g02560	VviPME10	pectinesterase 2	upregulated in PNP	off in PN, may come up later after veraison
VIT_209s0018g01490	-	methanol O-anthraniloyltransferase	upregulated in PNP	goes up before veraison, further up after veraison
VIT_210s0071g01145	VviRTIC1	plant protein with Domain of Unknown Function 789	upregulated in PN	off in PNP
VIT_213s0067g02930	VviEXPA14	expansin A8	upregulated in PNP	upregulated in PN, peak before veraison, down after
VIT_216s0013g00880	VviOLE5	oleosin 1	upregulated in PNP and PN	peaks before veraison, low consistency between years
VIT_216s0022g00960	VviGRIP28	ripening-related protein-like precursor	upregulated in PNP	goes up before veraison, stays up after veraison
VIT_216s0098g01170	VviHDZ28	homeobox-leucine zipper protein ATHB-12	upregulated in PN	up after flowering, goes down long before veraison
VIT_200s0366g00020	VviRTIC2	cysteine-rich receptor-like protein kinase 10	upregulated in PN	off in PNP
VIT_200s0956g00020	VviLEC1	nuclear transcription factor Y subunit B-6	upregulated in PNP and PN	peaks before/at verais., low consistency between years

The expression patterns derived from RNA-Seq confirm each other and also allow digital

quantification of low transcript accumulation levels. Nevertheless, three genes were chosen

for confirmation via qRT-PCR, namely *VviRTIC1*, *VviRTIC2* and *VviERF027* (VIT

216s0100g00400). *VviERF027* was included to cover a gene that displays, in different

samples of the time series, differential expression as well as equally high expression in PN

and PNP. The results obtained by qRT-PCR are fully congruent with the data from RNA-Seq

(see Additional file 2: Figure S4). Thus, differential gene activity in a developmental pattern

and in a genotype specific way has been detected by RNA-Seq as well as qRT-PCR.

Figure 5: Expression patterns of *VviRTIC1* (VIT_210s0071g01145) in (A) and *VviRTIC2*

(VIT_200s0366g00020) in (B) from RNA-Seq data of PN (blue) and PNP (red). Error bars

display the standard deviation of triplicates. Left, expression profile from 2014. Right,

expression profile from 2017. The y-axis represents the read counts from the output of

DESeq2. The x-axis represents the development stages in days after onset of flowering (DAF).

Discussion

One of the first detectable mentions of the cultivar 'Pinot Precoce' in connection with the

synonym '(German) Früh Burgunder Traube, translated: Early Burgundy Grape' (PNP) is in

the French book "Ampelographie retrospective" [37]. PNP is considered to be closely related

to 'Pinot noir' (PN) grapes and, here, we have confirmed the clonal relationship of PN and

PNP by 24 well distributed genomic SSR markers. Although this does not prove that PN is the

ancestor, it is very likely that PNP was derived from the cultivar PN by somatic mutation as

suggested earlier [36]. We used these two isogenic cultivars, that are distinguished by a clear

duration of berry formation phenotype, to analyze changes in gene expression throughout

berry development. The aim was to identify candidate genes that control the speed of berry

development and veraison timing. Samples from inflorescences as well as from forming and

ripening berries were collected from the onset of flowering until after PN and PNP veraison

in 2014 and 2017. These samples were subjected to RNA-Seq analyses in two well resolved

time series.

Phenotypic differences between the cultivars PN and PNP

The data from 4 years of careful assessment of the BBCH developmental stages of PN and

PNP at the same location validate earlier observations from viticulture [36] that lead to

establishment of PNP as a distinct grapevine cultivar in north European wine growing

countries. Berry formation lasts about two weeks less in PNP, is clearly accelerated

compared to PN and results in PNP entering veraison approximately two weeks earlier than

PN (Figure 1A, Table 1). It is reasonable to assume that this acceleration affects berry

formation throughout, from immediately after fruit set until veraison. Functionally, this

hypothesis implies that the genes that are responsible for the control of timing of berry

development, and for the establishment of the phenotypic difference between PN and PNP,

should be acting already very early in berry development, starting at least shortly after

flowering and at or even before BBCH61 to BBCH79. At the end of ripening (harvest), the

berries of PNP reach high sugar content earlier within the season when compared to PN.

General validation of the RNA-Seq dataset

To estimate overall data quality, the expression profiles obtained from PN and PNP were

correlated for the two sampled years, 2014 and 2017. Pearson correlation was moderate,

but this is expected considering the conditions of the free field environment. Exposure of the

vines to external factors like biotic or abiotic stressors, including weather conditions that

differ significantly between the years, also affect the transcriptome which reduces the level

of correlation [38]. In a PCA, almost all data points lie on the same intended track, and

biological replicates (subsamples) from both years are located close to each other. The main

actors, which predominantly influence the variance in the dataset, are genes related to cell

wall modification, secondary metabolism, wounding and hormone signaling. These gene

categories fit locical expectations since berry development is known to (i) be controlled by

hormones, (ii) require new cell walls, and (iii) be accompanied by accumulation of

secondary/specialized metabolites [24, 26]. These initial results validated the quality of the

dataset and indicated clearly that sampling of biologically closely related material for the

subsamples/triplicates was successful.

Co-expression analysis shows similar gene expression clusters between cultivars and years

To further validate the data with respect to comparability as well as reproducibility between the two years, related gene expression profiles were identified among all genes by clustering the data from the four different time series. Generally, clusters of the same genes with similar expression patters over time were observed for both cultivars and both years. Also, the cluster analyses for gene expression patterns among the years 2014 & 2017 in only PN and in only PNP, confirmed comparability of the gene expression patterns obtained in these two years. Detailed inspection revealed clusters representing expression profiles (and clusters of genes) with and without an environmental influence. Especially the cultivar-specific clusters C1_PN/PNP, C7_PN/PNP and C11_PN/PNP stand out. Comparison of the expression profiles for PN to those of PNP in these clusters identified a similar pattern that is moved to a different time in PNP. These findings are in agreement with the shifted ripening time phenology of the two cultivars discussed above. In contrast, the clusters C2_PN/PNP, C5_PN/PNP, C6_PN/PNP and C12_PN/PNP display more pattern similarity among the two years than among the two cultivars. Thus, the genes in these four clusters may display dependence on environmental factors in their expression patterns, potentially due to differences in the weather conditions between the two years studied. Strong environmental effects on gene expression patterns have also been described for grapevine berry development at 11 different environments (vineyards) from northern Italy [39]. The remaining other PN/PNP clusters C0, C3, C4, C8, C9 and C10 display highly similar expression profiles ofer al four conditions. The genes included in these clusters are probably less affected by environmental factors and/or the genotypic difference between PN and PNP. We conclude that our RNA-Seq results and expression level comparisons between two years are based on valid data.

When the genomic location of the DEGs is analyzed, a genome region on chromosome 16

comes into focus. In this region, 54 of the DEGs from the set of 1,923 intersecting DEGs

(Table 2) are located. Of these, 28 encode stilbene synthases [40] that are all up-regulated

after veraison of PNP (BBCH83). Stilbenes are a group of phenylpropanoid compounds (that

includes resveratrol) which are detected in many plants, which often accumulate in response

to biotic and abiotic stresses, and which are formed as a basic structuce by the key enzyme

stilbene synthase. The genome region fits to a major QTL (Ver1) for "timing of the onset of

veraison" on chromosome 16 [8]. It remains to be determined if this aggregation of DEGs is

by chance. Potentially, the observation is biased by co-regulation of a large number of

closely linked stilbene synthase genes.

Differentially expressed genes throughout berry development and identification of

veraison-specific genes

Differential gene expression analysis and subsequent filtering revealed 1,923 DEGs between

PNP and PN. DEG detection was based on a comparison of samples taken from the two

cultivars at very similar DAF. As expected for the characterized phenotype, PNP reaches

veraison when PN is still in the phase of berry formation. Consequently, the strong increase

in the number of detected DEGs shortly before and at veraison of PNP results from the

different developmental stage of PNP compared to the lagging PN. Subsequently, when also

PN enters veraison, the number of DEGs declines (note that Figure 4 uses a logarithmic

scale). A list of 388 genes that show up in both years with a veraison-specific expression

pattern was extracted and compared to published results. Interestingly, about 81.5% of the

388 PN/PNP veraison-specific genes were also described in the 4,351 differentially expressed

genes between the table grape cultivar '8612-CTR' (wild type) and its early ripening bud

mutation '8612-ERN' [27]. Also, analyses of berries from the cultivars 'Cabernet Sauvignon'

and 'Pinot Noir' by RNA-Seq identified a gene set of 5,404 genes marking the onset of berry

ripening [28]. This set covers 51.5% of the 388 PN/PNP veraison-specific genes (Additional

file 1: Table S14). Several "switch genes" which are supposed to encode key regulators of the

developmental transition at veraison [34, 41] are also included in the 388 veraison-specific

gene set (Additional file 1: Table S14). We conclude that the PN/PNP veraison-specific set of

388 genes represents a core set of genes that are relevant for executing the switch from

berry formation to berry ripening. As a relatively small gene set was detected that still

displays high overlap to those found by other studies that addressed similar biological

questions, indicates that the specific experimental setup and implemented filters used here

are appropriate to remove unrelated genes. Here, the comparison of "wildtype to mutant"

RNA-Seq results in isogenic background between PN and PNP, reduced environmentally

controlled transcriptome differences by sampling in the same vineyard/location, and dense

time course sampling together with high RNA-Seq read coverage allowed good resolution

power.

In order to check for potentially co-expressed genes within the veraison-specific gene set,

the memberships of these genes in the PN/PNP cluster analysis were investigated. A total of

48 veraison-specific DEGs were detected in cluster C6_PN/PNP (contains 914 genes). These

48 genes include several prominent ripening-related genes like *VviGRIP61*

(VIT_201s0011g05110), *VviMYBA8*, *VviMRIP1* (VIT_205s0049g00760, [42]), *VviGRIP4*

(VIT_205s0049g00520) and *VviGRIP28*. The ~20 *VviGRIP* genes were previously detected by

differential cDNA screening as ripening-induced genes in grape [43]. Another relevant cluster

is C5_PN/PNP (contains 263 genes) which includes 37 of the 388 veraison-specific DEGs.

Among these are *VviMYBA1*, *VviMYBA2*, *VviMYB15* and *VviGRIP22* (VIT_206s0004g02560).

The two clusters C5_PN/PNP and C6_PN/PNP show quite similar patterns (Additional file 2:

Figure S1A). It was, at first, not obvious which difference has forced CLUST to put a given

gene in either C5_PN/PNP or C6_PN/PNP. However, a comparison of the expression patterns

of *VviMYBA2* (in C5_PN/PNP) and *VviMYBA8* (in C6_PN/PNP; see Additional file 2: Figure S2

for a heatmap) shows that there are borderline cases regarding assignment to either

C5_PN/PNP or C6_PN/PNP.

In total, 22 genes encoding R2R3-MYB TFs were found among the 1,923 intersecting DEGs.

Based on the timing of expression in PN and PNP, the 22 *R2R3-MYB* genes can be classified

into three groups (Additional file 2: Figure S2). The first group is represented by *VviMYB24*

which is expressed during early flowering (BBCH61) but switched off already at the transition

from flowering to berry formation (BBCH71). *VviMYB24* is potentially orthologous to a group

of three *A. thaliana R2R3-MYB* genes (*AtMYB21/24/57*) that are expressed in flowers and

which function redundantly to regulate stamen development in the context of jasmonate

action [44]. It is tempting to speculate that *VviMYB24* has a similar function in grape.

The second group covers about 15 *R2R3-MYB* genes that are expressed during berry

formation and pre-veraison (BBCH71 to 77). This group includes *VviMYBF1* which regulates

flavonol biosynthesis [45], and *VviMYBPA5* as well as *VviMYBPA7* which belong to the clade

of *AtTT2*-related genes that control proanthocyanidin (PA, flavan-3-ol) biosynthesis [7, 46-

48]. The other *R2R3-MYB* genes in this group are less well characterized although there are

functions described for some of them, e.g. *VviMYBC2-L3* (VIT_214s0006g01620) as repressor

of specific branches of the phenylpropanoid pathway [49].

The third group of *R2R3-MYB* genes is active starting at veraison (after BBCH81) and covers

about six genes. Among them are the anthocyanin accumulation controlling genes,

VviMYBA1, VviMYBA2, VviMYBA3 and *VviMYBA8* for which there is good evidence that they

trigger anthocyanin biosynthesis [50]. Since PN and PNP are red berry cultivars, activity of

the TF genes that direct anthocyanin accumulation is expected. In addition, this group

includes *VviMYB14* and *VviMYB15* that are supposed to regulate the stilbene biosynthetic

pathway [51]. With regard to the heatmaps (Additional file 2: Figure S2) and the analyses of

the DEGs in this study in general, it should be noted that while the resolution within the

developmental program and time is quite good, our data do not resolve the exact location of

gene expression. Therefore, it remains to be determined if the expression detected is

derived from berry skin, flesh, the seed or other tissues/cells.

Putative ripening time control genes acting early in berry development

To focus on genes that are contributing to the acceleration of berry formation in PNP, and/or

to the control of timing of veraison, we selected DEGs detected at time points prior to

veraison of PNP (Table 2, Figure 4). This resulted in a set of only 12 genes that are potentially

involved in the regulation of ripening time (Table 3). According to our hypothesis that the

genes relevant for acceleration of berry formation in PNP, which cause the earlier onset of

ripening in PNP, should be acting from at least shortly after flowering, we designated this set

of genes as "ripening time controlling". However, genes that encode components of the

respective regulatory networks and target genes of regulators including secondarily affected

DEGs are surely included as well [52]. The 12 putative ripening time control genes, the DEGs

detected before veraison of PNP, encode proteins related to auxin action, pectin processing

enzymes related to cell wall modification, TFs from the HD-Zip as well as NF-Y/LEC families, a

cysteine-rich receptor-like protein kinase, an oleosin, and proteins with domains of unknown

function.

The expression patters of this set of differentially expressed genes, and also the complete

dataset of DEGs detected (Table 2), was screened for genes that were higher expressed in

PNP than in PN. However, although there are many genes that display earlier upregulation of

expression in PNP than in PN in the context of earlier veraison, none of the 12 "early

differential" putative ripening time control genes was significantly higher expressed in PNP

than in PN already at or early after flowering (Table 3). Also, among the other DEGs, no gene

with such a differential expression pattern was detected. Such early gene activation in PNP

compared to PN could hint at a dominant regulator that promotes faster ripening, but the

data are more in favor of loss of an inhibitor of fast ripening.

The two genes assigned to auxin signaling by VitisNet (vv30003) encode expansins (*VviEXPA5*

and *VviEXPA14*, [53]). Expansins are known to be involved in fruit ripening through cell wall

expansion and cell enlargement [54]. Auxin can delay the onset of veraison and ripening

processes in grapevine [19-21]. Since reduced expression of genes from the auxin signaling

pathway may indicate reduced auxin action due to lower auxin levels, the accelerated entry

of PNP into veraison might be initiated by reduced auxin levels. Additionally, the genes

VviPL1 (pectate lyase 1 [55]), *VviPME10* (pectin methylesterase 10, VIT_206s0009g02560)

VviGRIP28 (encoding a pectin methylesterase inhibitor precursor-like protein) are also

related to cell wall processes, indicating that cell wall modification is an important target

process also prior to veraison [54]. The gene *VviGRIP28* was also detected within a veraison-

specific meta-QTL designated ver/ph16.1 [9]. It remains to be determined if this correlation

has a functional basis.

The two genes in the set of 12 that encode TFs are *VviHDZ28* (VIT_216s0098g01170, [56])

and *VviLEC1* (VIT_200s0956g00020, [57]). The *V. vinifera* gene VIT_216s0098g01170 that has

been designated *VviHDZ28* has also been considered as a homolog of *AtHB12* (At3g61890),

but it seems that *VviHDZ07* (VIT_202s0025g02590) and *VviHDZ27* (VIT_215s0048g02870) are

more similar to *AtHB12*. In these cases, which lack clearly assignable homologs, transfer of

functional information reaches its limits and might be restricted to concluding that *VviHDZ28*

is important for organ development in *Vitis*. The gene *VviLEC1* is one of three genes in *V.*

vinifera which are homologs of *AtLEC1* (At1g21970, NF-YB9) and *AtL1L* (LEC1-like,

At1g21970, NF-YB6). LEC1 and L1L are central regulators of embryo and endosperm

development. They control, among other processes, embryo morphogenesis and

accumulation of storage reserve [58]. It is tempting to speculate that the reason for the

detection of *VviLEC1* among the 12 putative ripening time control genes is that also seed

development needs to be accelerated in PNP compared to PN. This would explain earlier and

higher expression of *VviLEC1* in PNP compared to PN as observed (Additional file 2: Figure

S3G). Consequently, *VviOLE5* (VIT_216s0013g00880, encoding an oleosin involved in oil body

formation [59]) would fit into the picture as relevant for lipid storage during seed

development. According to the proposed enzyme function as an alcohol acyltransferase by

the protein encoded by VIT_209s0018g01490 involved in volatile ester formation [60], this

gene could play a similar role. For the gene VIT_205s0077g01980 no functional annotation is

available (uncharacterized protein), although homologs exist throughout the Magnoliophyta.

605   Candidates for causal genes explaining the difference between PN and PNP

Among the 12 putative ripening time control genes, of which 10 are discussed above, two

are especially interesting. Detailed analyses of the full set of DEGs, visualized in Figure 4,

resulted in the identification of *VviRTIC1* and *VviRTIC2*, that could possibly be centrally

involved in the accelerated berry development and earlier beginning of ripening in PNP

compared to PN. The special feature of the expression patters of the two genes (Figure 5,

Table 3) is that both are differentially expressed already at the first time point analyzed

which was selected to hit the BBCH stage 61 (flowering before full bloom, DAF zero (0)).

Also, both genes are only barely expressed in PNP in both years studied, while expression in

PN is high at almost all time points. *VviRTIC1* is annotated to encode a protein containing a

domain of unknown function (DUF789), while *VviRTIC2* is annotated to encode a "cysteine-

rich receptor-like protein kinase". The best BLASTp hit to *A. thaliana* protein sequences

indicates that it is related to At4g23180/AtCRK10, but a closer inspection shows that

similarity to At4g05200/AtCRK25, At4g23160/AtCRK8 and At4g23140/AtCRK6 is almost as

high. This ambiguity, and also the fact that the *V. vinifera* genome contains several genes

related to *VviRTIC2* (e.g. VIT_210s0071g01200, VIT_202s0087g01020 or

VIT_203s0017g01550 as listed by PhyloGenes), complicates transfer of functional

information. CRKs are a subgroup of plant receptor-like kinases [61] and are encoded by a

family of 44 genes in *A. thaliana*. In a systematic analysis of the functions of *A. thaliana* CRKs,

evidence was collected for involvement in the control of plant development, biotic and

abiotic stress responses, photosynthesis as well as stomatal regulation [62]. This systematic

phenotypic screen of a large set of T-DNA insertion mutants revealed distinct phenotypes for

various of the CRK genes, but assignment of a molecular function to individual CRKs beyond

recognition of unknown ligands and signal transmission by phosphorylation remains a large

challenge.

As pointed out above, it is very possible that the genes we have identified are part of a

genetic pathway that controls timing of berry development in *V. vinifera*, but that we have

hit genes in a downstream part of this pathway. The relevance of the two candidate genes in

the causal genetic difference between PN and PNP remains to be determined. Phase-

separated genome sequences of the cultivars will be required to resolve the genome

structure of both alleles of *VviRTIC1 and 2* the genes in PN and PNP for an informative

comparison. In future studies, we will address this question, for example by long read DNA

sequencing.

Conclusions

This study detected 1,923 DEGs between the Pinot cultivars PN and PNP. The two clonal

cultivars display a phenotypic difference in berry development timing where PNP reaches

from full bloom to veraison faster than PN. We defined 388 DEGs as veraison-specific and 12

DEGs as putatively controlling ripening time. The relatively small number of veraison-specific

genes displays a very high overlap with results published for similar studies (see Additional

file 1: Table S14) and could be used for studying a phytohormone network that is acting

similarly in PN and PNP, but accelerated in PNP. Additionally, the ripening time control genes

identified here might offer access to a set of genes putatively important for triggering or

delaying the start of berry ripening in grapevine. Further investigations are needed to

elucidate structural differences in the genomes, the function of the observed DEGs, and their

role in shifting the onset of ripening in grapevine.

Material and Methods

Plant material and analysis of clonal relation

The grapevine (*Vitis vinifera* subsp. *vinifera* L.) cultivar PNP (Pinot Precoce Noir, VIVC No.

9280) is early ripening and has been described to be related to the cultivar PN (Pinot Noir,

VIVC No. 9279) [35] that ripens later than PNP. To prove the clonal relation, DNA from both

cultivars was genotyped utilizing 24 polymorphic SSR markers (VVS2, VVMD7, VVMD5,

VVMD32, VVMD28, VVMD27, VVMD25, VVMD24, VVMD21, VVIV67, VVIV37, VVIQ52,

VVIP60, VVIP31, VVIN73, VVIN16, VVIH54, VVIB01, VrZAG83, VrZAG79, VRZAG67, VrZAG62,

VMC4F3.1, VMC1B11) as described [63]. The two cultivars used have been identified as

accession DEU098_VIVC9280_Pinot_Precoce_Noir_DEU098-2008-076 and

DEU098_VIVC9279_Pinot_Noir_DEU098-2008-075, respectively. The tissue used for harvest

is indicated below and in Figure 1. Both cultivars do not belong to an endangered species

and were obtained and are grown in accordance with German legislation.

Phenotypical characterization and sampling of plant material

Plant material was harvested from PN and PNP grapevines trained in trellis. The plants are

growing at the vineyards of JKI Geilweilerhof located at Siebeldingen, Germany (N

49°21.747, E 8°04.678). The grapevine plants were planted with an interrow distance of 2.0

m and spacing of 1.0 m in north-south direction. Inflorescences, developing and ripening

berry samples of PNP and PN for RNA extraction were collected in two years with three

independent biological replicates (subsamples) each. Sampling took place at systematic time

points (12 time points in 2014, 13 time points in 2017), and at approx. 8 a.m. each day. In

2014, harvesting took place regularly every 7 days with only two exceptions (one day

deviation, DAF 13 and DAF 27). In 2017, harvesting was adapted to BBCH stages (Figure 1A).

The timeline in both years is described as days after onset of flowering (DAF), with onset of

flowering defined as the day at which 10% of the individual flowers have lost their caps

(BBCH61 [3]). For each subsample within the triplicates, material from two neighboring

grapevines was selected. Grapevine plants were weekly phenotyped according to BBCH

stage [3, 4]. Phenotyping was performed repeatedly to ensure sampling from vines of the

same development stage (e.g. percentage of open flowers during flowering, or berry

development stage) to reach uniform subsamples. The phenotypical observations were

summarized in Additional file 1: Table S2 and S3. From these, the durations of flowering,

berry formation and berry ripening as well as the resulting shifts between the cultivars were

calculated (Table 1). Furthermore, images from berry developmental stages of both cultivars

were taken in 2014 for 35, 41, 49 and 56 DAF. The sampled material was directly frozen in

liquid nitrogen and stored at -70°C until RNA extraction.

RNA extraction and cDNA library construction

Biological replicates, i.e. the subsamples, were ground separately under liquid nitrogen.

Total RNA was extracted using an RNA Isolation Kit (Sigma-Aldrich Spectrum™ Plant Total

RNA) according to suppliers' instructions. Quality control, determination of RIN numbers [64]

and estimation of the concentrations of all RNA samples was done on a Bioanalyzer 2100

(Agilent) using RNA Nano 6000 Chips. For RNA-Seq, 500 ng total RNA per subsample were

used to prepare sequencing libraries according to the Illumina TruSeq RNA Sample

Preparation v2 Guide. For subsamples from 2014 and 2017, 72 and 78 libraries were

constructed and sequenced, respectively. Enrichment of poly-A containing mRNA was

performed twice, using poly-T oligos attached to magnetic beads included in the Illumina kit.

During the second elution of the poly-A+ RNA, the RNA was fragmented and primed for

cDNA synthesis. After cDNA synthesis, the fragments were end-repaired and A-tailing was

performed. Multiple indexing adapters were ligated to the ends of the cDNA fragments and

the adapter ligated fragments were enriched by 10 cycles of PCR. After quality check using

Bioanalyzer 2100 HS-Chips (Agilent) and exact quantification by Quant-iT PicoGreen dsDNA

assay on a FLUOstar Optima Plate Reader (BMG LABTECH), the libraries were pooled

equimolarly.

RNA-Seq

Single end (SE) sequencing of the pooled barcoded libraries from 2014 was performed on an

Illumina HiSeq1500 in High Output mode generating 100 nt reads. For samples from 2017,

sequencing was done using an Illumina NextSeq500 generating 83 nt SE reads; two runs

were performed with the same pool of barcoded libraries from 2017.

Processing of RNA-Seq read data

Raw reads were trimmed with Trimmomatic (version 0.36) [65]. For raw reads from the year

2014, the following settings were used: LEADING:10 TRAILING:10 SLIDINGWINDOW:4:15

MINLEN:50. In addition, a collection of all available Illumina adapter sequences was supplied

to remove matches within the parameter 2:30:10. For raw reads from the year 2017,

trimming settings were set to LEADING:6 TRAILING:6 SLIDING WINDOW:4:15 MINLEN:36. All

trimmed reads were quality-checked via FastQC (version 0.11.8) [66]. Thus, possible adapter

sequences and low-quality bases were removed. All trimmed reads passing QC were mapped

to the reference genome sequence PN40024 (version 12Xv2) [67] using the graph-based

alignment tool HISAT2 (version 2.1.0) [68, 69] with no additional soft clipping. Afterwards, all

tagged genes (structural gene annotation: CRIBI v2.1) were counted as raw read counts with

FeatureCounts (Bioconductor package Rsubread version 3.8 [70]). To estimate transcript

abundance as a measure for gene expression, counts for Transcripts Per Kilobase Million

(TPM, [71]) were determined.

Basic gene expression analyses

TPM counts from the various samples were used for manual gene expression inspection, for

determination of the number of expressed and not expressed genes, and to calculate the

correlation between gene expression values from both years. Genes with a TPM value > 0

added up over all samples from one year were classified as expressed, conversely genes with

a TPM value = 0 added up over all samples as not expressed. A custom python script was

applied utilizing the function pearsonr from SciPy python package (v. 1.2.3) [72], which

calculates the Pearson correlation coefficient and the p-value for all year-to-year

comparisons. Expression data pairs for TPM counts per gene from both sampled years,

averaged over the three subsamples of each sample, were used. To test correlations and

relationships between expression values from the two years, where samples were harvested

with slightly different sampling patterns (see Figure 1A), eight equivalent time points with

the same BBCH stages between the cultivars of each year were selected (see Additional

file 1: Table S4).

Principal component analysis

To explore data similarity, a Principal Component Analysis (PCA) was calculated over all gene

expression values from both years and cultivars for all subsamples. All data points were

normalized using variance stabilizing transformation function 'vst' from the R package

DESeq2 (v. 1.12.4) [73]. Subsequently, the principal components were generated using

'prcomp' from the R package 'stats' (v. 3.5.2) [74]. The resulting PCA object, displaying the

main components PC1 and PC2, was plotted and exported. Additionally, genes with the

highest variance contribution to PC1 and PC2 were extracted separately.

Functional annotation of genes

Transfer of annotation information from other plant species, mainly *A. thaliana,* was

calculated using MapMan's sequence annotation tool Mercator (v. 3.6) [75, 76]. Additionally,

all open reading frame (CDS from *V. vinifera*/grapevine genes) sequences were aligned to

the non-redundant protein sequence data base RefSeq [77] with the basic local alignment

tool for proteins BLASTp [78] (e-value ≤ 0.001). Short descriptions of gene functions were

extracted and added to the gene lists in Additional file 1: Table S5, S6, S13, S14.

GO term enrichment for biological processes was calculated via the R package 'topGO' (v.

2.38.1) [79]. Subsequently, statistical reliability was calculated using Fishers exact test. All

Gene IDs and their corresponding GO terms were extracted from the CRIBI database

(http://genomes.cribi.unipd.it/DATA/V2/annotation/bl2go.annot_with_GO_description.txt).

All results of the GO term enrichment are deposited in Additional file 1: Table S10-12.

Cluster analysis

To reveal co-expressed genes over all four datasets, the tool CLUST (v. 1.10.8) was used with

default parameters [80]. As input, raw read counts from eight time points were used. These

time points were selected to cover the same BBCH stages of PN and PNP from the years

2014 and 2017 (Additional file 1: Table S4). First, all data were pre-processed as described in

the CLUST manual. Values from corresponding subsamples (triplicates) were combined and

averaged. To filter out uninformative (very low) gene expression values, an additional filter

was applied: genes not reaching a sample expression value > 1 in at least three conditions

and in at least one cultivar from one year were discarded (-fil-v 1 -fil-c 3 -fil-d 1). Afterwards,

the data were quantile normalized according to the RNA-Seq defaults of CLUST. Genes

showing a flat expression profile were filtered out by applying the default settings [80].

Differential gene expression analyses

For analyses of differentially expressed genes, DESeq2 (v. 1.12.4; R Bioconductor) was

employed. To test if gene expression differs significantly between two samples, the

likelihood ratio test nbinomLRT, included in the DESeq2 package, was used. As input, raw

read counts from all time points were used. Normalization factors and dispersion estimates

were used as described [73]. The output table contained all differentially expressed genes

(DEGs) and the corresponding values for baseMean, log2FoldChange (LFC), lfcSE (LFC

standard error), stat (difference in deviation between the reduced model and the full

model), p-value and PADJ (adjusted p-value). To focus on significantly differentially

expressed genes from the DESeq2 analyses, cut-off filters PADJ ≤ 0.05 and LFC > 2 were

applied.

Confirmation of differential gene expression by qRT-PCR

To verify the RNA-Seq results, four time points from 2017 were selected (DAF 0, DAF 28,

DAF57 and DAF77) for qRT-PCR. Synthesis of cDNA from the RNA subsamples was carried

out with First Strand cDNA Synthesis Kit (ProtoScript® II; NEB) according to the

manufacturer's instructions. The qRT-PCR assay was performed using Luna Universal qPCR

Master Mix (NEB) with a total volume of 20 µl. Sequences of the primers used are listed in

Additional file 1: Table S23. Reaction products/amplicons were detected based on SYBR

green via a CFX96 Real Time PCR Detection System (Bio-Rad). For each time point, three

biological and three technical replicates (i.e., each subsample in triplicate) were measured.

Cycling conditions after initial denaturation 2 min at 95°C: denaturation 5 sec at 95°C,

annealing/extension 30 sec at 60°C, cycled 35 times. For QC, each reaction was controlled by

product melt analysis (65°C - 95°C). As negative controls, no template control (NTC) and no

reverse transcriptase control (-RT) were measured as well using three technical replicates.

The polyubiquitin gene *VviUbiquitin1* (VIT_216s0098g01190) was used for normalization

[46]. Measurements were analyzed using the CFX Maestro V.4.1.2433.1219 (Bio-Rad) by

normalization via the relative quantitative ΔΔCt method.

Selection of gene sets potentially relevant for ripening time and comparison with

literature data

In order to identify gene sets from the DEGs relevant for control and implementation of

ripening, an intersection between the DEGs detected at all time points between both years

was built. To determine a subset of putatively ripening time control genes, the intersection

between both years covering the development stages BBCH61 (onset of flowering) to

BBCH79 (one developmental BBCH stage before veraison) was used (time points 2014:

DAF 0-35; 2017: DAF 0-42). Furthermore, a set of veraison specific genes was defined from

the DEGs detected at the intersection of development stages BBCH79 (one developmental

BBCH stage before veraison) to BBCH81 (onset of ripening / veraison; time points 2014:

DAF35-41; 2017: DAF42-49). To test for biological relevance of the subsets, all DEGs were

screened to their occurrence in similar relevant studies [9, 27, 28, 30, 34, 41, 81, 82].

Visualization of gene numbers newly appearing as differentially expressed

To visualize appearance of DEGs over time, a stacked bar plot script was set up using the R

package 'plotly' (v. 4.9.2.1) [83]. Each bar represents the amount of DEGs of a given time

point or condition. In order to track groups of DEGs newly appearing at a given time point

throughout the following time points, the colour shade representing the group of DEGs

remains the same.

Pathway enrichment analysis

To search for possible targets in known pathways of grapevine, a pathway enrichment

analysis using the tool VitisPathways [84] was performed. To achieve a reliable enrichment,

1000 permutations, a Fisher's exact test of p < 0.05 and permuted p-value < 0.1 were set.

Thus, all significant enriched pathway genes and their relations can be displayed in VitisNet

[85], a specific molecular network for grapevine.

Heatmaps

As an extension to assignment of genes to biosynthesis pathways, the genes were also

filtered for annotation as coding for transcription factors (TFs). This filter was based on the

annotation information transferred from Mercator and RefSeq (see above). To look at the

entire family of *R2R3-MYB* TF genes, the list of *MYB* genes identified via MapMan was

extended by additional grapevine *R2R3-MYB* gene family members that have been

characterized [6, 7]. The *R2R3-MYB* genes detected among the intersecting DEGs were

displayed in heatmaps addressing the four individual time series (2 cultivars, 2 years) using

the R package 'pheatmap' (v. 2.1.3) [86]. Predictions for phylogenetic relationships were

deduced from PhyloGenes v. 2.2 [87].

Abbreviations

ABA   abscisic acid

DAF   days after onset of flowering

DEGs   differentially expressed genes

PNP   Pinot Noir Precoce

PN   Pinot Noir

LFC   log2fold change

PADJ   adjusted p-value

PCA   principle component analysis

TF   transcription factor

TPM   transcripts per kilobase million

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The FASTQ files containing all RNA-Seq reads (PN2014, PNP2014, PN2017 and PNP2017)

have been deposited at the European Nucleotide Archive (ENA) according to the INTEGRAPE

guidelines under the accession numbers PRJEB39262, PRJEB39261, PRJEB39264 and

PRJEB39263, respectively (see Additional file 1: Tables S19 to S22). All scripts developed for

this study are available on GitHub [https://github.com/bpucker;

https://github.com/jenthein].

Competing interests

The authors declare that they have no competing interests

Funding

This project was funded by BMBF/PtJ (FKZ 031A3496).

Authors' contributions

Sampling and phenotyping were done at the JKI (Julius Kühn-Institute, Institute for

Grapevine Breeding Geilweilerhof, Siebeldingen, Germany). Sequencing and data analysis

were performed at Bielefeld University, Faculty of Biology & Center for Biotechnology

(CeBiTec). Organization, supervision at Bielefeld University and parts of sampling were done

by DH. KH, AK and FS coordinated weekly sampling, phenotyping, and image capture for

documentation. Logistic work was done by KH. The design of the experiments was set up by

FS, DH and KH. RNA isolation and cDNA synthesis were carried out by DH and PV. PV

accomplished library preparation and sequencing. Quantitative RT-PCR was performed by JT

and DH. RT and KH supervised the work at JKI Geilweilerhof. LH performed the SSR analysis.

BW supervised the work at Bielefeld University. FS, KH, RT and BW acquired project funding

and wrote the project proposal. All bioinformatic data analyses, creation of figures, tables

and writing of the manuscript were performed by JT with the help of DH. JT and BW drafted

the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

We would like to thank Boas Pucker for providing several python scripts for handling large

datasets as well as helpful comments on the manuscript, Katharina Frey-Sielemann for help

visualizing the DEG distributions and Wiebke Halpape for providing the GO annotation

enquiry. We also thank Kyle J. Lauersen for higly valuable language editing, and Willy Keller

for the excellent technical support. We are very grateful to the Bioinformatics Resource

Facility support team of the CeBiTec for providing computing infrastructure and excellent

technical support. This publication is based in part upon contributions from COST Action CA

17111 INTEGRAPE, supported by COST (European Cooperation in Science and Technology);

the authors thank BMBF/PtJ for funding in the context of IPAS (project acronym novisys).

The authors from UniBi also wish to thank the other members of the Chair of Genetics and

Genomics of Plants for their support. We would also like to thank the Open Access

Publication Funds of Bielefeld University for funding the publication charge.

Supplementary material

Additional file 1: Table S1: Clonal relation of PN and PNP confirmed by 24 SSR markers

Additional file 1: Table S2 - Sampling and phenotypical observations (BBCH) for 2014

Additional file 1: Table S3 - Sampling and phenotypical observations (BBCH) for 2017

Additional file 1: Table S4 - Pearsonr correlations of gene expression values over time

between both years

Additional file 1: Table S5 - Top 500 genes influencing principal component PC1

Additional file 1: Table S6 - Top 500 genes influencing principal component PC2

Additional file 1: Table S7 - Cluster memberships of clusters obtained from the

clusteranalysis PN/PNP

Additional file 1: Table S8 - Cluster memberships of clusters obtained from the cluster

analysis PN

Additional file 1: Table S9 - Cluster memberships of clusters obtained from the cluster

analysis PNP

Additional file 1: Table S10 - GO term enrichment for the clusters from the cluster analysis

of PN/PNP

Additional file 1: Table S11 - GO term enrichment for the clusters from the cluster analysis

of PN

Additional file 1: Table S12 - GO term enrichment for the clusters from the cluster analysis

of PNP

Additional file 1: Table S13 - Differentially expressed genes between PN and PNP in both

years and filtering steps

Additional file 1: Table S14 - Overlap of 12 putatively ripening time control DEGs (time

points: DAF 0-35 from 2014 intersecting with DAF 0-42 from 2017) and 388 veraison-specific

DEGs (time points: DAF 35+41 from 2014 intersecting with DAF 42+49 from 2017) with

previous transcriptomic grapevine studies.

Additional file 1: Table S15 - VitisNet enrichment for intersecting DEGs

Additional file 1: Table S16 - VitisNet enrichment for regulatory DEGs

Additional file 1: Table S17 - VitisNet enrichment for veraison specific DEGs

Additional file 1: Table S18 - Up- & Down-regulated TFs among the DEGs

Additional file 1: Table S19 - ENA sample identifier and metadata linked to the study for PNP

RNA-Seq reads from 2014 (PRJEB39262)

Additional file 1: Table S20 - ENA sample identifier and metadata linked to the study for PN

RNA-Seq reads from 2014 (PRJEB39261)

Additional file 1: Table S21 - ENA sample identifier and metadata linked to the study for PNP

RNA-Seq reads from 2017 (PRJEB39264)

Additional file 1: Table S22 - ENA sample identifier and metadata linked to the study for PN

RNA-Seq reads from 2017 (PRJEB39263)

Additional file 1: Table S23 - Sequences of primers used for qRT-PCR

Additional file 2: Figure S1 - Cluster analyses of PN/PNP in both years (A), PN in both years

(B) and PNP in both years (C)

Additional file 2: Figure S2 - Heatmap of all MYB-TFs gene expression among the DEGs for

PN and PNP in both years

Additional file 2: Figure S3 - Expression patterns of genes from the ripening-regulatory gene

set

Additional file 2: Figure S4 - Confirmation of expression data from RNA-Seq by qRT-PCR

References

1. Kuhn N, Guan L, Dai ZW, Wu BH, Lauvergeat V, Gomès E, Li SH, Godoy F, Arce-Johnson P, Delrot S: **Berry ripening: recently heard through the grapevine.** *Journal of Experimental Botany* 2014, **65**(16):4543-4559.

2. Coombe BG: **Research on development and ripening of the grape berry.** *American Journal of Enology and Viticulture* 1992, **43**(1):101-110.

3. Lorenz DH, Eichhorn KW, Bleiholder H, Klose R, Meier U, Weber E: **Phenological growth stages of the grapevine (Vitis vinifera L. ssp. vinifera) - Codes and descrptions according to the extended BBCH scale.** *Australian Journal of Grape and Wine Research* 1995, **1**(2):100-103.

4. Bleiholder H, Weber E, Feller C, Hess M, Wicke H, Meier U, van den Boom T, Lancashire PD, Buhr L, Hack H *et al*: **Growth stages of mono- and dicotyledonous plants - BBCH Monograph**: Federal Biological Research Centre for Agriculture and Forestry; 2001.

5. Dubos C, Stracke R, Grotewold E, Weisshaar B, Martin C, Lepiniec L: **MYB transcription factors in Arabidopsis.** *Trends in Plant Science* 2010, **15**(10):573-581.

6. Matus JT, Aquea F, Arce-Johnson P: **Analysis of the grape MYB R2R3 subfamily reveals expanded wine quality-related clades and conserved gene structure organization across Vitis and Arabidopsis genomes.** *BMC Plant Biology* 2008, **8**:83.

7. Wong DCJ, Schlechter R, Vannozzi A, Holl J, Hmmam I, Bogs J, Tornielli GB, Castellarin SD, Matus JT: **A systems-oriented analysis of the grapevine R2R3-MYB transcription factor family uncovers new insights into the regulation of stilbene accumulation.** *DNA Research* 2016, **23**(5):451-466.

8. Zyprian E, Ochßner I, Schwander F, Šimon S, Hausmann L, Bonow-Rex M, Moreno-Sanz P, Grando MS, Wiedemann-Merdinoglu S, Merdinoglu D *et al*: **Quantitative trait loci affecting pathogen resistance and ripening of grapevines.** *Molecular Genetics and Genomics* 2016, **291**(4):1573-1594.

9. Delfino P, Zenoni S, Imanifard Z, Tornielli GB, Bellin D: **Selection of candidate genes controlling veraison time in grapevine through integration of meta-QTL and transcriptomic data.** *BMC Genomics* 2019, **20**(1):739.

10. Delrot S, Grimplet J, Carbonell-Bejerano P, Schwandner A, Bert P-F, Bavaresco L, Costa LD, Di Gaspero G, Duchêne E, Hausmann L *et al*: **Genetic and Genomic Approaches for Adaptation of Grapevine to Climate Change.** In: *Genomic Designing of Climate-Smart Fruit Crops.* Edited by Kole C. Cham (Switzerland): Springer; 2020: 157-270.

11. Fenn MA, Giovannoni JJ: **Phytohormones in Fruit Development and Maturation.** *The Plant Journal* 2020.

12. Osorio S, Scossa F, Fernie AR: **Molecular regulation of fruit ripening.** *Frontiers in Plant Science* 2013, **4**:198.

13. Sun L, Zhang M, Ren J, Qi J, Zhang G, Leng P: **Reciprocity between abscisic acid and ethylene at the onset of berry ripening and after harvest.** *BMC Plant Biology* 2010, **10**:257.

14.   Pilati S, Bagagli G, Sonego P, Moretto M, Brazzale D, Castorina G, Simoni L, Tonelli C, Guella G,
Engelen K *et al*: **Abscisic Acid Is a Major Regulator of Grape Berry Ripening Onset: New**
Insights into ABA Signaling Network. *Frontiers in Plant Science* 2017, **8**:1093.

15.   Zhu Z, Li G, Yan C, Liu L, Zhang Q, Han Z, Li B: **DRL1, Encoding A NAC Transcription Factor, Is**
Involved in Leaf Senescence in Grapevine. *International Journal of Molecular Sciences* 2019,
20(11):2678.

16.   Rossdeutsch L, Edwards E, Cookson SJ, ., Barrieu F, Gambetta GA, Delrot S, Ollat N: **ABA-**
mediated responses to water deficit separate grapevine genotypes by their genetic
background. *BMC Plant Biology* 2016, **16**(91):s12870-12016-10778-12874.

17.   Coelho J, Almeida-Trapp M, Pimentel D, Soares F, Reis P, Rego C, Mithöfer A, Fortes AM: **The**
study of hormonal metabolism of Trincadeira and Syrah cultivars indicates new roles of
salicylic acid, jasmonates, ABA and IAA during grape ripening and upon infection with
Botrytis cinerea. *Plant Science* 2019, **283**:266-277.

18.   Chervin C, El-Kereamy A, Roustan J-P, Latché A, Lamon J, Bouzayen M: **Ethylene seems**
required for the berry development and ripening in grape, a non-climacteric friut. *Plant*
Science 2004, **167**(6):1301-1305.

19.   Böttcher C, Burbidge CA, Boss PK, Davies C: **Interactions between ethylene and auxin are**
crucial to the control of grape (Vitis vinifera L.) berry ripening. *BMC Plant Biology* 2013,
13:222.

20.   Böttcher C, Harvey K, Forde CG, Boss PK, Davies C: **Auxin treatment of pre-veraison grape**
(Vitis vinifera L.) berries both delays ripening and increases the synchronicity of sugar
accumulation. *Australian Journal of Grape and Wine Research* 2011, **17**(1):1-8.

21.   Ziliotto F, Corso M, Rizzini FM, Rasori A, Botton A, Bonghi C: **Grape berry ripening delay**
induced by a pre-véraison NAA treatment is paralleled by a shift in the expression pattern of
auxin- and ethylene-related genes. *BMC Plant Biology* 2012, **12**:185.

22.   Seymour GB, Ostergaard L, Chapman NH, Knapp S, Martin C: **Fruit Development and Ripening.**
Annual Review of Plant Biology 2013, **64**:219-241.

23.   Giovannoni J, Nguyen C, Ampofo B, Zhong S, Fei Z: **The Epigenome and Transcriptional**
Dynamics of Fruit Ripening. *Annual Review of Plant Biology* 2017, **68**:61-84.

24.   Karlova R, Chapman N, David K, Angenent GC, Seymour GB, de Maagd RA: **Transcriptional**
control of fleshy fruit development and ripening. *J Exp Bot* 2014, **65**(16):4527-4541.

25.   Wang R, Tavano E, Lammers M, Martinelli AP, Angenent GC, de Maagd RA: **Re-evaluation of**
transcription factor function in tomato fruit development and ripening with CRISPR/Cas9-
mutagenesis. *Scientic Reports* 2019, **9**(1):1696.

26.   Serrano A, Espinoza C, Armijo G, Inostroza-Blancheteau C, Poblete E, Meyer-Regueiro C, Arce
A, Parada F, Santibanez C, Arce-Johnson P: **Omics Approaches for Understanding Grapevine**
Berry Development: Regulatory Networks Associated with Endogenous Processes and
Environmental Responses. *Frontiers in Plant Science* 2017, **8**:1486.

27.   Ma Q, Yang J: **Transcriptome profiling and identification of the functional genes involved in**
berry development and ripening in Vitis vinifera. *Gene* 2019, **680**:84-96.

28.   Fasoli M, Richter CL, Zenoni S, Bertini E, Vitulo N, Dal Santo S, Dokoozlian N, Pezzotti M,
Tornielli GB: **Timing and Order of the Molecular Events Marking the Onset of Berry Ripening**
in Grapevine. *Plant Physiology* 2018, **178**(3):1187-1206.

29.   Sun R, He F, Lan Y, Xing R, Liu R, Pan Q, Wang J, Duan C: **Transcriptome comparison of**
Cabernet Sauvignon grape berries from two regions with distinct climate. *Journal of Plant*
Physiology 2015, **178**:43-54.

30.   Zenoni S, Ferrarini A, Giacomelli E, Xumerle L, Fasoli M, Malerba G, Bellin D, Pezzotti M,
Delledonne M: **Characterization of transcriptional complexity during berry development in**
Vitis vinifera using RNA-Seq. *Plant Physiology* 2010, **152**(4):1787-1795.

31.   Xu L, Yue Q, Xiang G, Bian F, Yao Y: **Melatonin promotes ripening of grape berry via increasing**
the levels of ABA, H2O2, and particularly ethylene. *Horticulture Research* 2018, **5**:41.

32.   Griesser M, Martinez SC, Eitle MW, Warth B, Andre CM, Schuhmacher R, Forneck A: **The**
ripening disorder berry shrivel affects anthocyanin biosynthesis and sugar metabolism in
Zweigelt grape berries. *Planta* 2018, **247**(2):471-481.

33.   Zarrouk O, Brunetti C, Egipto R, Pinheiro C, Genebra T, Gori A, Lopes CM, Tattini M, Chaves
MM: **Grape Ripening Is Regulated by Deficit Irrigation/Elevated Temperatures According to**
Cluster Position in the Canopy. *Frontiers in Plant Science* 2016, **7**:1640.

34.   Massonnet M, Fasoli M, Tornielli GB, Altieri M, Sandri M, Zuccolotto P, Paci P, Gardiman M,
Zenoni S, Pezzotti M: **Ripening Transcriptomic Program in Red and White Grapevine Varieties**
Correlates with Berry Skin Anthocyanin Accumulation. *Plant Physiology* 2017, **174**(4):2376-
2396.

35.   Maul E, Sudharma KN, Ganesch A, Brühl U, Hundemer M, Kecke S, Mahler-Ries A, Marx G,
Schreiber T, Walk M *et al*: **30 Years VIVC - Vitis International Variety Catalogue**
(www.vivc.de). In: *XI International Conference on Grapevine Breeding and Genetics; Yanqing,*
Beijing, China. 2014.

36.   Robinson J, Harding J, Vouillamoz J: **Wine Grapes: A complete guide to 1,368 vine varieties,**
including their origins and flavours: Penguin Books; 2012.

37.   Roy-Chevrier J: **Ampelographie retrospective**. Paris; 1900.

38.   Pucker B, Schwandner A, Becker S, Hausmann L, Viehover P, Topfer R, Weisshaar B, Holtgrawe
D: **RNA-Seq Time Series of Vitis vinifera Bud Development Reveals Correlation of Expression**
Patterns with the Local Temperature Profile. *Plants* 2020, **9**(11):1548.

39.   Dal Santo S, Tornielli GB, Zenoni S, Fasoli M, Farina L, Anesi A, Guzzo F, Delledonne M, Pezzotti
M: **The plasticity of the grapevine berry transcriptome**. *Genome Biology* 2013, **14**(6):R54.

40.   Vannozzi A, Dry IB, Fasoli M, Zenoni S, Lucchin M: **Genome-wide analysis of the grapevine**
stilbene synthase multigenic family: genomic organization and expression profiles upon
biotic and abiotic stresses. *BMC Plant Biology* 2012, **12**:130.

41.   Palumbo MC, Zenoni S, Fasoli M, Massonnet M, Farina L, Castiglione F, Pezzotti M, Paci P:
Integrated network analysis identifies fight-club nodes as a class of hubs encompassing key
putative switch genes that induce major transcriptome reprogramming during grapevine
development. *The Plant Cell* 2014, **26**(12):4617-4635.

42. Burger AL, Zwiegelaar JP, Botha FC: **Characterisation of the gene encoding the Merlot**
ripening-induced protein 1 (mrip1): evidence that this putative protein is a distinct member
of the plant proline-rich protein family. *Plant Science* 2004, **167**(5):1075-1089.

43. Davies C, Robinson SP: **Differential screening indicates a dramatic change in mRNA profiles**
during grape berry ripening. Cloning and characterization of cDNAs encoding putative cell
wall and stress response proteins. *Plant Physiology* 2000, **122**(3):803-812.

44. Qi T, Huang H, Song S, Xie D: **Regulation of Jasmonate-Mediated Stamen Development and**
Seed Production by a bHLH-MYB Complex in Arabidopsis. *The Plant Cell* 2015, **27**(6):1620-
1633.

45. Czemmel S, Stracke R, Weisshaar B, Cordon N, Harris NN, Walker AR, Robinson SP, Bogs J: **The**
grapevine R2R3-MYB transcription factor VvMYBF1 regulates flavonol synthesis in
developing grape berries. *Plant Physiology* 2009, **151**(3):1513-1530.

46. Bogs J, Downey MO, Harvey JS, Ashton AR, Tanner GJ, Robinson SP: **Proanthocyanidin**
Synthesis and Expression of Genes Encoding Leucoanthocyanidin Reductase and
Anthocyanidin Reductase in Developing Grape Berries and Grapevine Leaves. *Plant*
Physiology 2005, **139**(2):652-663.

47. Bogs J, Jaffe FW, Takos AM, Walker AR, Robinson SP: **The grapevine transcription factor**
VvMYBPA1 regulates proanthocyanidin synthesis during fruit development. *Plant Physiology*
2007, **143**(3):1347-1361.

48. Koyama K, Numata M, Nakajima I, Goto-Yamamoto N, Matsumura H, Tanaka N: **Functional**
characterization of a new grapevine MYB transcription factor and regulation of
proanthocyanidin biosynthesis in grapes. *Journal of Experimental Botany* 2014, **65**(15):4433-
4449.

49. Cavallini E, Matus JT, Finezzo L, Zenoni S, Loyola R, Guzzo F, Schlechter R, Ageorges A, Arce-
Johnson P, Tornielli GB: **The phenylpropanoid pathway is controlled at different branches by**
a set of R2R3-MYB C2 repressors in grapevine. *Plant Physiology* 2015, **167**(4):1448-1470.

50. Matus JT, Cavallini E, Loyola R, Holl J, Finezzo L, Dal Santo S, Vialet S, Commisso M, Roman F,
Schubert A *et al*: **A group of grapevine MYBA transcription factors located in chromosome 14**
control anthocyanin synthesis in vegetative organs with different specificities compared with
the berry color locus. *The Plant Journal* 2017, **91**(2):220-236.

51. Vannozzi A, Wong DCJ, Holl J, Hmmam I, Matus JT, Bogs J, Ziegler T, Dry I, Barcaccia G, Lucchin
M: **Combinatorial Regulation of Stilbene Synthase Genes by WRKY and MYB Transcription**
Factors in Grapevine (Vitis vinifera L.). *Plant and Cell Physiology* 2018, **59**(5):1043-1059.

52. Forlani S, Masiero S, Mizzotti C: **Fruit ripening: the role of hormones, cell wall modifications,**
and their relationship with pathogens. *Journal of Experimental Botany* 2019, **70**(11):2993-
3006.

53. Dal Santo S, Vannozzi A, Tornielli GB, Fasoli M, Venturini L, Pezzotti M, Zenoni S: **Genome-wide**
analysis of the expansin gene superfamily reveals grapevine-specific structural and
functional characteristics. *PLoS One* 2013, **8**(4):e62206.

54. Du M, Spalding EP, Gray WM: **Rapid Auxin-Mediated Cell Expansion**. *Annual Review of Plant*
Biology 2020, **71**:379-402.

55. Guillaumie S, Fouquet R, Kappel C, Camps C, Terrier N, Moncomble D, Dunlevy JD, Davies C,
Boss PK, Delrot S: **Transcriptional analysis of late ripening stages of grapevine berry**. *BMC*
Plant Biology 2011, **11**:165.

56. Li Z, Zhang C, Guo Y, Niu W, Wang Y, Xu Y: **Evolution and expression analysis reveal the**
potential role of the HD-Zip gene family in regulation of embryo abortion in grapes (Vitis
vinifera L.). *BMC Genomics* 2017, **18**(1):744.

57. Cui M, Wang W, Guo F, Fan X, Guan L, Zheng T, Zhu X, Jia H, Fang J, Wang C *et al*:
Characterization and temporal–spatial expression analysis of LEC1 gene in the development
of seedless berries in grape induced by gibberellin. *Plant Growth Regulation* 2020, **90**(3):585-
596.

58. Jo L, Pelletier JM, Harada JJ: **Central role of the LEAFY COTYLEDON1 transcription factor in**
seed development. *Journal of Integrative Plant Biology* 2019, **61**(5):564-580.

59. Liu Q, Sun Y, Su W, Yang J, Liu X, Wang Y, Wang F, Li H, Li X: **Species-specific size expansion and**
molecular evolution of the oleosins in angiosperms. *Gene* 2012, **509**(2):247-257.

60. Qian X, Liu Y, Zhang G, Yan A, Wang H, Wang X, Pan Q, Xu H, Sun L, Zhu B: **Alcohol**
acyltransferase gene and ester precursors differentiate composition of volatile esters in
three interspecific hybrids of Vitis labruscaxV. Vinifera during berry development period.
Food Chemistry 2019, **295**:234-246.

61. Liang X, Zhou JM: **Receptor-Like Cytoplasmic Kinases: Central Players in Plant Receptor**
Kinase-Mediated Signaling. *Annual Review of Plant Biology* 2018, **69**:267-299.

62. Bourdais G, Burdiak P, Gauthier A, Nitsch L, Salojarvi J, Rayapuram C, Idanheimo N, Hunter K,
Kimura S, Merilo E *et al*: **Large-Scale Phenomics Identifies Primary and Fine-Tuning Roles for**
CRKs in Responses Related to Oxidative Stress. *PLoS Genetics* 2015, **11**(7):e1005373.

63. Dallakyan M, Zyprian E, Maul E, Yesayan A, Hovhannisyan N: **Genetic characterization of rape**
varieties in Armenia. *Vitis* 2015, **54**:23-26.

64. Schroeder A, Mueller O, Stocker S, Salowsky R, Leiber M, Gassmann M, Lightfoot S, Menzel W,
Granzow M, Ragg T: **The RIN: an RNA integrity number for assigning integrity values to RNA**
measurements. *BMC Molecular Biology* 2006, **7**:3.

65. Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina sequence data**.
Bioinformatics 2014, **30**(15):2114-2120.

66. **FastQC, a quality control tool for high throughput sequence data**.
[http://www.bioinformatics.babraham.ac.uk/projects/fastqc/]

67. Jaillon O, Aury JM, Noel B, Policriti A, Clepet C, Casagrande A, Choisne N, Aubourg S, Vitulo N,
Jubin C *et al*: **The grapevine genome sequence suggests ancestral hexaploidization in major**
angiosperm phyla. *Nature* 2007, **449**(7161):463-467.

68. Kim D, Langmead B, Salzberg SL: **HISAT: a fast spliced aligner with low memory requirements**.
Nature Methods 2015, **12**(4):357-360.

69. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL: **Graph-based genome alignment and**
genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* 2019, **37**(8):907-915.

70.   Liao Y, Smyth GK, Shi W: **featureCounts: an efficient general purpose program for assigning**
sequence reads to genomic features. *Bioinformatics* 2014, **30**(7):923-930.

71.   Zhao S, Ye Z, Stanton R: **Misuse of RPKM or TPM normalization when comparing across**
samples and sequencing protocols. *RNA* 2020, **26**(8):903-909.

72.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E,
Peterson P, Weckesser W, Bright J *et al*: **SciPy 1.0: fundamental algorithms for scientific**
computing in Python. *Nature Methods* 2020, **17**(3):261-272.

73.   Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for RNA-**
seq data with DESeq2. *Genome Biology* 2014, **15**(12):550.

74.   **R: A language and environment for statistical computing** [https://www.R-project.org/]

75.   Thimm O, Bläsing O, Gibon Y, Nagel A, Meyer S, Krüger P, Selbig J, Müller LA, Rhee SY, Stitt M:
MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic
pathways and other biological processes. *The Plant Journal* 2004, **37**(6):914-939.

76.   Lohse M, Nagel A, Herter T, May P, Schroda M, Zrenner R, Tohge T, Fernie AR, Stitt M, Usadel
B: **Mercator: a fast and simple web server for genome scale functional annotation of plant**
sequence data. *Plant, Cell & Environment* 2014, **37**(5):1250-1258.

77.   Pruitt KD, Tatusova T, Maglott DR: **NCBI reference sequences (RefSeq): a curated non-**
redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Research*
2007, **35**(Database issue):D61-65.

78.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search tool.** *Journal*
of Molecular Biology 1990, **215**(3):403-410.

79.   Alexa A, Rahnenfuhrer J, Lengauer T: **Improved scoring of functional groups from gene**
expression data by decorrelating GO graph structure. *Bioinformatics* 2006, **22**(13):1600-1607.

80.   Abu-Jamous B, Kelly S: **Clust: automatic extraction of optimal co-expressed gene clusters**
from gene expression data. *Genome Biology* 2018, **19**:172.

81.   Zenoni S, D'Incà E, Tornielli GB: **Genetic dissection of grape berry ripening control: defining a**
role for NAC transcription factors. *Acta Horticulturae* 2019(1248):387-402.

82.   Wei L, Cao Y, Cheng J, Xiang J, Shen B, Wu J: **Comparative transcriptome analyses of a table**
grape 'Summer Black' and its early-ripening mutant 'Tiangong Moyu' identify candidate
genes potentially involved in berry development and ripening. *Journal of Plant Interactions*
2020, **15**(1):213-222.

83.   Sievert C: **Interactive Web-Based Data Visualization with R, plotly, and shiny**: CRC Press;
2020.

84.   Osier MV: **VitisPathway: gene pathway analysis for V. vinifera.** *Vitis* 2016, **55**:129-133.

85.   Grimplet J, Van Hemert J, Carbonell-Bejerano P, Díaz-Riquelme J, Dickerson J, Fennell A,
Pezzotti M, Martínez-Zapater JM: **Comparative analysis of grapevine whole-genome gene**
predictions, functional annotation, categorization and integration of the predicted gene
sequences. *BMC Research Notes* 2012, **5**:213.

86.   **pheatmap: Pretty Heatmaps. R package version 1.0.12** [https://CRAN.R-
project.org/package=pheatmap]

87.   Zhang P, Berardini TZ, Ebert D, Li Q, Mi H, Muruganujan A, Prithvi T, Reiser L, Sawant S, Thomas
PD *et al*: **PhyloGenes: An online phylogenetics and functional genomics resource for plant**
gene function inference. *Plant Direct* 2020, **4**(12):e00293.

1198