Phycobilisome breakdown effector NblD is required to maintain the cellular

amino acid composition during nitrogen starvation

Vanessa Krauspe,[a] Stefan Timm,[b] Martin Hagemann,[b] Wolfgang R. Hess,[a,#]

[a]Genetics and Experimental Bioinformatics, Institute of Biology 3, Faculty of Biology,

University of Freiburg, 79104 Freiburg, Germany

[b]Plant Physiology Department, Institute of Biosciences, University of Rostock,

Rostock, 18059 Rostock, Germany;

Running Head: NblD affects metabolite pools

[#]Correspondence: Wolfgang R. Hess, wolfgang.hess@biologie.uni-freiburg.de

Keywords: amino acids, cyanobacteria, nitrogen starvation, photosynthesis,

phycobilisomes, *Synechocystis* sp. PCC 6803, small proteins, stress response

Abstract

Small proteins are critically involved in the acclimation response of photosynthetic cyanobacteria to nitrogen starvation. NblD is the 66-amino-acid effector of nitrogen-limitation-induced phycobilisome breakdown, which is believed to replenish the cellular amino acid pools. To address the physiological functions of NblD, the concentrations of amino acids, intermediates of the arginine catabolism pathway and several organic acids were measured during the response to nitrogen starvation in the cyanobacterium *Synechocystis* sp. PCC 6803 wild type and in an *nblD* deletion strain. A characteristic signature of metabolite pool composition was identified, which shows that NblD-mediated phycobilisome degradation is required to maintain the cellular amino acid and organic acid pools during nitrogen starvation. Specific deviations from the wild type suggest wider-reaching effects that also affect such processes as redox homeostasis via glutathione and tetrapyrrole biosynthesis, both of which are linked to the strongly decreased glutamate pool, and transcriptional reprogramming via an enhanced concentration of 2-oxoglutarate, the metabolite co-regulator of the NtcA transcription factor. The essential role played by NblD in metabolic homeostasis is consistent with the widespread occurrence of NblD throughout the cyanobacterial radiation and the previously observed strong positive selection for the *nblD* gene under fluctuating nitrogen supply.

Importance

Cyanobacteria play important roles in the global carbon and nitrogen cycles. In their natural environment, these organisms are exposed to fluctuating nutrient conditions. Nitrogen starvation induces a coordinated nitrogen-saving program that includes the

breakdown of nitrogen-rich photosynthetic pigments, particularly phycobiliproteins.

The small protein NblD was recently identified as an effector of phycobilisome

breakdown in cyanobacteria. In this study, we demonstrate that the NblD-mediated

degradation of phycobiliproteins is needed to sustain cellular pools of soluble amino

acids and other crucial metabolites. The essential role played by NblD in metabolic

homeostasis explains why genes encoding this small protein are conserved in

almost all members of cyanobacterial radiation.

Introduction

The recent introduction of the antibiotic retapamulin as an inhibitor of the initiation of translation into ribosome-profiling studies in bacteria has revolutionized the view on the bacterial small proteome suggesting that a high number of previously unexplored small open reading frames (smORFs) exists in bacteria (1, 2). Small proteins, often defined as those containing 50 or fewer amino acids are too small to have own enzymatic activities and this applies also to the group of "larger" small proteins that range from 50 to 100 amino acids. Nevertheless, small proteins have been shown to fulfil multiple important functions, for example as regulatory subunits of multi-protein complexes, as modulators of enzymatic activities of larger proteins or as toxins (for a recent review see Hemm et al. (3)).

A large number of small proteins have been characterized in photosynthetic cyanobacteria due to the extensive work on characterizing the photosynthetic apparatus. Nineteen small proteins with fewer than 50 amino acids are directly involved in photosynthesis. These proteins constitute subunits of photosystem I (encoded by genes *psaM, psaJ* and *psaI* (4)), of photosystem II (genes *psbM, psbT* (*ycf8*), *psbI, psbL, psbJ, psbY, psbX, psb30* (*ycf12*), *psbN, psbF, psbK* (5, 6)), participate in photosynthetic electron transport (Cytb_6f complex; *petL, petN, petM, petG* (7–9)) or in photosynthetic complex I (*ndhP*, *ndhQ* (10, 11)). Other previously characterized small proteins in cyanobacteria have accessory functions to photosynthesis such as the small CAB like proteins HliA, HliB, HliC and HliD (12–14). The cytochrome b_6f complex subunit VIII, encoded by *petN* is with 29 amino acids the shortest annotated photosynthetic protein conserved in cyanobacteria (15). Additional smORFs were computationally predicted in the model cyanobacterium *Synechocystis* sp. PCC 6803 (hereafter designated *Synechocystis* 6803) (16). The

44 amino acids small protein AcnSP was characterized as regulator of the

tricarboxylic acid (TCA) cycle enzyme aconitase (17).

Another set of small proteins were found to be involved in the acclimation response

of the photosynthetic machinery to nitrogen (N) starvation, including NblA1, NblA2

and NblD, proteins of 62, 60 and 66 residues in *Synechocystis* 6803 (18, 19).

Nondiazotrophic cyanobacteria, such as *Synechocystis* 6803, depend on combined

nitrogen (N) sources in their environment. These bacteria favor ammonia (NH_3 or

NH_4^+) but can also utilize nitrate (NO_3^-) as an inorganic N source (20, 21). Under N-

limited conditions, cells can start an acclimation program, manifested as chlorosis,

and finally enter a dormant state (22–26). During chlorosis, N-rich pigments are

degraded, predominantly the giant light-harvesting pigment-protein complexes, the

phycobilisomes (22, 23, 27). The excess carbon (C) availability during chlorosis

induces the accumulation of C reserves, particularly glycogen (28). The tricarboxylic

acid (TCA) cycle intermediate 2-oxoglutarate (2-OG), which is primarily used as a C

skeleton for ammonium incorporation by the glutamine synthetase/glutamine 2-OG

aminotransferase (GS/GOGAT) cycle (29, 30), has been shown to serve as a cellular

signal metabolite to report the relative N/C status (31–35). If ammonia assimilation

via the GS/GOGAT cycle is slowed due to N deficiency, 2-OG is no longer converted

and accumulates in the cyanobacterial cell. As a result, 2-OG can bind to the signal

transduction protein PII (36, 37), thereby releasing PipX from the PipX/PII complex

(33, 35). Free PipX can subsequently associate with and activate the global nitrogen

transcription regulator NtcA together with 2-OG. In *Synechocystis* 6803, NtcA directly

controls 79 different genes after 4 h of N starvation (38). Depending on the location

of the respective binding motifs, NtcA can either activate genes, especially those

involved in nitrogen uptake and assimilation, or repress genes, as observed for the

GS inhibition factors *gifA* and *gifB* (39). Two of the NtcA-activated genes in

Synechocystis 6803 are *nblA1* and *nblA2* (*ssl0452* and *ssl0453*) (38), which encode

small proteins that are critically important in N starvation-induced phycobilisome

breakdown (18, 40–42).

Phycobilisome breakdown in N-starved cyanobacteria is a tightly controlled process

that is believed to fill cellular amino acid pools, compensating for the limited *de novo*

nitrogen assimilation via the GS-GOGAT cycle. NblA1/A2 play essential roles in the

coordinated process of phycobilisome degradation during N deprivation because

corresponding mutants show a so-called non-bleaching phenotype (*nbl*), i.e., they

retain their phycobilisomes under N-limiting conditions (18, 40). NblA proteins

function as adaptors for the Clp protease by connecting the phycobiliproteins and the

ClpC chaperone (41–44). Phycobilisome degradation can be generally divided into a

rapid early phase, in which almost 50% of phycocyanin is degraded within 5 h,

followed by a more sedate phase until only approximately 10% of the initial

phycocyanin content is attained after 24 h of N deficiency (45). The degradation of

the phycobilisome starts at the distal phycocyanin rods and extends to the

allophycocyanin core, which is in proximity to the photosystems situated in the

thylakoid membrane (44, 46). Previous studies showed that the inability to degrade

phycobilisomes in the Δ*nblA1/A2* mutant affected the composition of amino acid

pools during N starvation in a specific way in *Synechocystis* 6803 (47). These

authors demonstrated that the pool sizes of 12 amino acids were NblA1/A2-

dependent, while others were not.

NblD has been recently identified as another small protein determined to be

important for phycobilisome turnover because the pigmentation of Δ*nblD* mutants

was only marginally reduced under N starvation (19) resembling the non-bleaching

phenotype of Δ*nblA* mutants. NblD specifically binds to the phycocyanin beta subunit

(CpcB) of phycobilisomes at an early time point of nitrogen depletion (19). However,

the precise biochemical function of NblD remains to be determined. In this study, we

compared the metabolic composition of *Synechocystis* 6803 wild-type cells (WT)

with a mutant deficient for NblD (Δ*nblD*) over a time course of N starvation. A

characteristic signature in the composition of these metabolite pools was observed

that supports the essential role played by NblD in the maintenance of cellular amino

acid homeostasis during N starvation. Other deviations from the WT suggest wider-

reaching effects, also affecting transcriptional reprogramming via an unusually high

concentration of 2-OG, the metabolite co-regulator of the NtcA transcription factor.

Results

Global responses of an *nblD* deletion mutant to nitrogen starvation.

In the present study, the amino acid pools and amounts of other important

metabolites were compared in the *Synechocystis* 6803 wild type (WT) to an *nblD*

deletion mutant (Δ*nblD*). In the absence of a nitrogen source, cells of the Δ*nblD* strain

are bleaching significantly less than the WT (19). Accordingly, we hypothesized that

the soluble amino acid pools of Δ*nblD* cells may be exhausted more quickly than

those of WT cells. This assumption was tested by metabolomics based on liquid

chromatography mass spectrometry (LC-MS/MS) of three biologically independent

samples for each WT and Δ*nblD*, taken before (+N condition), and 3 h or 24 h after

the transfer into N-free BG11 medium. In total, 28 compounds, including almost all

amino acids (Asn and Cys were below the detection limit), intermediates of the

arginine catabolism pathway, and several organic acids, were quantified (see Table

S1 for the LC-MS/MS raw data).

Cells obtained from the +N condition and 3 h after nitrogen removal showed no

difference in pigmentation. After 24 h of N starvation, the WT samples showed

pronounced bleaching, while the Δ*nblD* mutant bleached much less (Fig. 1A and B).

Comparing the spectra with previously published spectra suggests that the decrease

in the 630 nm absorption maximum of phycobilisomes in Δ*nblD* under N starvation is

greater than in the Δ*nblA1/nblA2* double mutant (18, 42), although clearly not as

much as in the WT under the same condition. The degradation of phycobilisomes

contributed to transiently higher levels of total soluble amino acids in *Synechocystis*

6803 WT during N starvation (Fig. 2A, Table S2). The level of free amino acids

increased from approximately 261 ng OD_{750}^{-1} ml^{-1} (± 32.4) under nitrogen-replete

conditions (+N) by 65% on average after 3 h and approximately 27% (in relation to

the starting value) after 24 h of N limitation (-N). In contrast, the Δ*nblD* mutant

contained 335 ng OD_{750}^{-1} ml^{-1} (± 43.3) of free amino acids in the +N condition and

remained at a similar level after 3 h. After 24 h after nitrogen removal, the amino acid

content decreased dramatically by 70% to 75% in relation to the initial content in

Δ*nblD* (Fig. 2A). Compared to the amino acid pool, the total amounts of other

quantified metabolites, primarily organic acids, did not differ significantly between

cells of the WT and Δ*nblD* (Fig. 2B), at least in the +N condition and after 24 h. The

higher total amount after 3 h is mostly due to increased 2-OG and citrate

concentrations in the mutant (see below).

Changes in the composition of amino acid pools differ between WT and Δ*nblD*.

Despite the overall trend of a decreasing total amino acid pool in mutant Δ*nblD* compared to WT, the direction of changes in the amounts of individual amino acids were not unidirectional (Fig. 3). According to their differential changes between WT and Δ*nblD* during the N starvation period, these changes could be categorized into four groups. Group 1 comprised amino acids showing no major differences between WT and Δ*nblD;* group 2 included those amino acids that appeared different already at the +N condition, while group 3 exhibited differences between WT and Δ*nblD* early, after 3 h, or group 4 later, after 24 h of N starvation (Fig. 3).

Amino acids of group 1 showed similar trends in both WT and Δ*nblD* during nitrogen limitation, as was observed for Ala, Arg, Gln and His. The strongest deviation in the response to N starvation between the two strains was observed for Glu. The level of Glu did not decrease during the 24 h N starvation period in WT but decreased in mutant Δ*nblD* to approximately 17% of the initial value after 24 h of N starvation (Fig. 3). Glu is by far the most abundant soluble amino acid and serves as an amino group donor in the majority of transaminase reactions. Hence, the depletion of the Glu pool should have notable effects on the entire metabolism. Indeed, the lack of Glu extended into the pool of reduced glutathione (GSH), which is the tripeptide γ-L-glutamyl-L-cysteinylglycine; the initial increase observed in the WT was lower in cells of the mutant Δ*nblD* and led after 24 h into N limitation to a significantly reduced amount (Fig. 3). GSH plays an important role as a reducing agent in all living cells and was found to be essential in the acclimation of *Synechocystis* 6803 to environmental and redox perturbations (48); hence, its depletion probably should have an impact on redox homeostasis in the mutant Δ*nblD*. The amino acids of group 3, such as Gly, Pro, Ser, Thr and Val, exhibited transiently increased levels

after 3 h in WT, which was not the case or was less pronounced in the mutant (Fig.

3). The branched chain amino acids Ile, Leu and Val share the same biosynthetic

pathway, whereas Gly and Ser are closely linked to photorespiratory metabolism and

can be interconverted by serine hydroxymethyltransferase (49). Phe and Thr

exhibited increased levels in WT cells after 24 h of N starvation, while they returned

to initial levels in Δ*nblD,* accumulating at values of less than 50% of the WT. Finally,

the aromatic amino acid Tyr from group 2 accumulated nearly 100-fold more in WT

cells after 24 h of N starvation, whereas it remained at low levels in Δ*nblD* (Fig. 3).

This finding is consistent with a previous analysis of *Synechocystis* 6803 WT after 24

h N starvation (47). Tyr represents approximately 5% of the amino acids stored in

the phycobilisome (Fig. 4). The strong accumulation of Tyr in WT but not in Δ*nblD*

suggests that it originated directly from phycobilisome degradation, assuming that in

contrast to other amino acids, Tyr is not readily incorporated into the metabolism of

N-starved cells.

Changes in other N-containing metabolites and organic acids

In addition to proteinogenic amino acids, many additional metabolites could be

quantified. For example, ornithine, a key intermediate in the arginine catabolism

pathway, was clearly depleted in Δ*nblD* upon N starvation compared to WT (Fig. 5).

This points to the arginine catabolism pathway as another mechanism for

stabilization of the Glu pool. In this pathway, the arginine-guanidine-removing

enzyme (AgrE) together with proline oxidase (PutA) can produce Glu from the Arg

molecules released from phycocyanin via ornithine (50–52). Therefore, the

diminished ornithine pool adds another important element to the changed nitrogen

homeostasis in cells of the Δ*nblD* strain. This assumption was also supported by

changes in the 2-OG pool, which was lower in WT than in the mutant under all

conditions, indicating a relatively more N-starved state of $\Delta nblD$ cells. Moreover, we

observed that the absence of NblD also affected the pool of 3-phosphglyceric acid

(3PGA), the stable CO_2 fixation product of RubisCO, which was decreased under all

conditions. Excess 3PGA can be used to produce glycogen, which is highly

accumulating in N-starved cells during chlorosis (53). Otherwise, some 3PGA can be

diverted via phosphoglycerate mutase into the direction of lower glycolysis and the

TCA cycle, which is primarily regulated by PirC (54). In contrast to the decreased

flow into the oxidative branch, the reductive branch of the TCA cycle seemed to be

less affected by the absence of NblD, since the pool of malate was similar in cells of

the mutant and the WT (Fig. 5). However, succinate was clearly lowered in $\Delta nblD$

after 3 h under N limitation, probably due to altered consumption of succinate by the

respiratory chain. In the mutant cells, the succinate pool stabilized at a similar

quantity as was observed in WT cells under prolonged N depletion. Taken together,

the metabolic changes observed in this study enabled us to conclude that the

absence of NblD had a broad impact on primary metabolism.

Discussion

Comparison of the changes in Δ*nblD* with the previously characterized

Δ*nblA1/A2* mutant.

NblD is involved in the disassembly of phycobilisomes by targeting CpcB during

acclimation to N starvation (19). In contrast to the WT, the $\Delta nblD$ mutant maintains

the majority of its pigments under this condition resembling the non-bleaching

phenotype of the previously reported $\Delta nblA$ strains and suggesting that

phycobilisomes are retained in these mutants to a large extent during the considered

time frame (see Fig. 1). In case of the Δ*nblA1/A2* mutant, amino acid pools were

analyzed after 24 h in the presence or absence of nitrogen. The authors categorized

the changes in the amino acid pool sizes into NblA1/A2-dependent and -independent

responses. Most amino acids (Gln, Glu, Gly, Ile, Leu, Met, Phe, Pro, Ser, Thr, Tyr,

Val and also GSH) fell into the NblA1/A2-dependent category. In the present study,

we analyzed the effects of the absence of NblD on the amino acid and organic acid

pools in the Δ*nblD* deletion mutant after 3 h and 24 h of N starvation. The observed

differences between Δ*nblD* and WT in the amino acid pools were similar to those

described for the Δ*nblA1/A2* mutant (47) but were not identical, despite the fact that

both types of proteins play a role in phycobilisome degradation. Five amino acids

(Ile, Leu, Ser, Thr, Val) had after 24 h of N starvation higher levels in WT but lower

levels in the Δ*nblA1/A2* mutant compared to the control in N-replete media. While we

observed the same changes for these 5 amino acids after 24 h N starvation, we

noticed that their pools were peaking even higher in WT in the sample taken after 3

h. In the study by Kiyota et al. (47), the WT had been analyzed at even higher

temporal resolution (after 1, 3, 6, 12 and 24 h of N starvation) and they reported that

these 5 amino acids transiently overshot at the 3 h value, hence our data are

consistent. However, we found a transient accumulation of Ile, Leu, Ser, Thr, and Val

in cells of the Δ*nblD* mutant like in the WT in the 3 h sample. Hence, our data allow

to differentiate between an early NblD-independent response for these 5 amino acids

and a later response that was NblD-dependent. This finding suggests that the early,

very pronounced response was independent from amino acid release from the

phycobilisomes, whereas they were mobilized at the later time point, which both the

Δ*nblD* and Δ*nblA1/A2* mutants failed to do.

The levels of four other amino acids (Pro, Gly, Glu, Tyr) and GSH behaved very

similar in the Δ*nblA1/A2* and Δ*nblD* mutants. However, for three amino acids (Gln,

Met and Phe) we observed differences between the different non-bleaching mutants.

Gln levels steadily declined in our study, both in WT and Δ*nblD* over the time of the

experiment; hence, it appeared as an NblD-independent response (Fig. 3). For Met

and Phe, similar or higher concentrations had been reported in the WT but lower in

Δ*nblA1/A2* upon N starvation, with Met even falling below the detection limit after 24

h (47). While we observed the same effect for the WT, in contrast the Δ*nblD* mutant

exhibited transiently higher levels of Met and Phe after 3 h, and about the same

levels after 24 h as at the beginning.

In the work by Kiyota et al. (47) some amino acids were identified as NblA1/A2-

independent (Ala, Asn, Asp, Lys and Trp). The changes in the pools sizes of these

amino acids were also NblD-independent in our work. We can add Arg and His to

this category, which were below detection limit in the work on NblA, while we could

not measure Asn. Moreover, the decline in Gln levels appeared as an NblD-

independent response, different from the findings on NblA.

Collectively, the overall responses of Δ*nblA1/A2* and Δ*nblD* mutants were similar but

some deviations appeared. For example, our investigation of the 3 h time point

allowed to conclude that some amino acid changes are independent from amino acid

releases via the phycobilisome degradation. Consequently, though NblA1/NblA2-

and NblD-assisted phycobilisome degradation represents the most obvious protein

degradation process due to visible pigment changes, faster changes in the proteome

likely contribute to the dynamics of amino acid changes especially during the early

phase of acclimation to N starvation.

Phycobilisome proteins and amino acid pools during nitrogen starvation.

Degradation of phycocyanin rods and their proteolytic recycling are assumed to

contribute extensively to the soluble amino acid pool in the cell, making those amino

acids available to be incorporated into other proteins to acclimate towards nutrient

limitation. From the phycobilisome core, up to six rod antennas radiate, each of

which is built by three hexameric discs connected with linker polypeptides CpcG and

CpcC (55). One hexameric phycocyanin disc contains six subunits of each CpcA and

CpcB.

Therefore, the most abundant phycobilisome proteins are CpcA and CpcB, which are

rich in Ala, Gly, Leu, Val and Tyr. Assuming full phycobilisome turnover, including

also all linker proteins and allophycocyanin core, it would provide mostly Ala (~14%),

Leu, Gly (both ~8%), Ser, Val, Arg, and Asp (contributing 6-7% each to

phycobilisome composition). Accordingly, the pools of these amino acids were

expected to increase early during N-starvation-induced phycobilisome degradation in

the WT but not in Δ*nblD*. Such an increase was clearly observed for Tyr, whereas

the changes in the amounts of the other amino acids differed less pronounced

between WT and Δ*nblD* cells. The amounts of Gly, Leu, Ser, and Val were always

higher in the WT than in the mutant under -N conditions. However, the amount of

Ala, the most frequent amino acid in the phycobilisome showed a similar pool

change in WT and mutant cells, suggesting its increase resulted from synthesis

instead. Ala biosynthesis is derived from the transamination of pyruvate and Glu by

alanine dehydrogenase (*sll1682*) or by serine-pyruvate aminotransferase using

pyruvate and Ser. Therefore, Ala is directly linked to the metabolic pathways of Ser,

Gly and Thr, as well as to primary C metabolism, via pyruvate. Indeed, in

Synechococcus elongatus PCC 7942, alanine dehydrogenase was found to be

required for correct phycobilisome degradation during N starvation, resulting in lower

nblA transcription if the gene for alanine dehydrogenase was deleted (56).

As mentioned above, besides Tyr the largest difference between WT and mutant

ΔnblD was observed for the pool of Glu after 24 h of N starvation. Glu is can be

regarded as the central amino acid from several perspectives. It is certainly the most

relevant amino group donor for transaminations, consistent with its abundance,

which is far higher than any other amino acid. In addition, Glu serves as the

precursor of tetrapyrrole synthesis, leading to cytochromes and especially

chlorophylls in cyanobacteria, green algae and plants upon its activation by binding

to its cognate tRNA^{glu} (57–59). This C5 pathway is the sole biosynthetic pathway for

tetrapyrrole biogenesis in *Synechocystis* 6803 (60, 61). Glu has an approximate 5%

share in the composition of the phycobilisome proteins (Fig. 4). Therefore, it is likely

that amino groups of abundant amino acids that become available upon

phycobilisome degradation are utilized predominantly to stabilize the Glu pool via

transaminase reactions. Initially, stabilization of the Glu pool may occur through

glutamate synthase from Gln and 2-OG, which is also consistent with the relatively

lower amounts of 2-OG in WT compared to ΔnblD cells (Fig. 5), then via arginine

catabolism from Arg. A corresponding reduction in Gln and Arg pools is seen at 3 h

and at 24 h into N starvation in both WT and ΔnblD. At the later time point,

additionally Asp, Gly, Leu, possibly Ala and Val, might be a source of amino groups

as these amino acids are (i) relatively abundant, (ii) relatively frequent in the

phycobilisome and (iii) their pools initially increased in the WT after shift to N

starvation, followed by a particularly sharp reduction from 3 h to 24 h. In addition, the

development of pool sizes for these amino acids showed marked differences

between the WT and ΔnblD.

Our results support the hypothesis that amino acid pools are regulated to a large

extent independently of phycobilisome degradation by NblA1/A2 and NblD, because

the cells maintain distinct amino acid pool homeostasis involving transaminase

reactions, primarily using the amino groups from the released phycobilisome

constituents. Furthermore, such amino acids as Ala, Asp and Lys might originate

from the breakdown of several other proteins (47). Hence, even though

NblA1/NblA2- and NblD-assisted phycobilisome degradation represents the most

obvious protein degradation process due to visible pigment changes, additional

changes in the proteome likely contribute to the recycling of amino acids during

acclimation to N starvation.

Effects on the organic acids pools.

The striking differences between WT and Δ*nblD* observed for Glu at 24 h of N

starvation may extend also further, into the amounts of certain organic acids,

especially succinate. At 3 h -N, succinate levels were very low, whereas Glu levels

were high; at 24 -N, succinate levels were relatively high, whereas Glu levels were

very low. There are four possibilities for the relatively high succinate levels after 24 h

of N starvation:

(i) it may appear from an enhanced flux into the reductive branch of the TCA

cycle working in anabolic direction in both strains, because the malate content

also increased over time;

(ii) an enhanced closure of the "open" cyanobacterial TCA cycle due to the

shunt, which converts 2-OG into succinate via 2-oxoglutarate decarboxylase

and succinic semialdehyde dehydrogenase reactions (62) would provide higher

succinate amounts;

(iii) Succinate could also originate from the gamma-aminobutyric acid shunt

closing the "open" TCA cycle in which glutamate is converted into gamma-

aminobutyric acid and then via succinic semialdehyde to succinate (63);

(iv) The succinate consumption by respiration via succinate dehydrogenase

could have become reduced.

The flux through the alternative TCA shunts has been regarded to be rather low

under light conditions (62, 63) as applied here, therefore, the possibility (ii) appears

rather unlikely. Possibility (iv) is unlikely as respiration does not decrease but

increases upon N limitation (24). Consequently, the most likely scenario is that the

succinate accumulation after 3 h N starvation in WT cells resulted from increased

flow through the arginine catabolism pathway from phycobiliproteins (variant iii) via

Glu, while the succinate accumulation after 24 h in the WT and especially

pronounced in ΔnblD originates from the reductive branch of the TCA cycle (i). This

scenario is supported by the N starvation-induced expression (53, 64) of *slr1022*

encoding an enzyme that functions also as γ-aminobutyrate aminotransferase (63),

a key enzyme in the gamma-aminobutyric acid shunt pathway interconverting Glu to

succinate.

Connecting metabolic changes and the regulation of gene expression.

Several of the metabolites showing marked changes between WT and mutant ΔnblD

possess signaling functions connecting metabolism and the regulation of gene

expression. For example, Glu is used as a signaling molecule in green algae,

mosses and higher plants sensed by the PII protein (65). Moreover, in several Gram-

negative bacteria, Gln serves as a signaling molecule for N availability (reviewed by

Forchhammer, 2013 (66)). More recently, it was observed that Gln also exhibits a

signaling function in *Synechocystis* 6803 and other cyanobacteria via interaction with

the GlnA riboswitch governing the expression of *gifB* mRNA (67).

In contrast to Gram-negative bacteria, cyanobacteria use the TCA cycle intermediate

2-OG to sense the C/N status. 2-OG is the regulatory co-factor of NtcA and of PII,

where the 2-OG accumulation transduces the low N signal leading to the activation

of NtcA (36, 37). Conversely, under high N levels 2-OG is converted via the

GS/GOGAT cycle to Glu, thereby the low 2-OG level diminishes the activity of NtcA.

In addition, the decrease in 2-OG is characteristic for C limitation and activates the

transcription factor NdhR to upregulate genes involved in the cyanobacterial

inorganic C-concentrating mechanism (68, 69).

Hence, the increase in 2-OG in the WT observed during N starvation should

represent a pronounced signal for activation of NtcA, and indeed, the transcriptional

units within the NtcA regulon become strongly differentially regulated during this

condition (19). Notably, there is an NtcA motif in the promotor region of the *nblD*

gene itself (16), consistent with its role in N-limited conditions. However, compared

to WT, we observed that 2-OG increased even more in Δ*nblD* and at every sampling

point, including the +N condition (Fig. 5). Because of the function of 2-OG as the

main cellular indicator metabolite to report C/N homeostasis (31–35), increased 2-

OG levels therefore clearly indicated a reduced ammonia assimilating capacity of

mutant Δ*nblD* already under N replete conditions. The likely reduced use of 2-OG in

the GS/GOGAT cycle was supported by the observation that the levels of citrate and

aconitate, precursors of 2-OG in the oxidative branch in the open cyanobacterial

TCA cycle, were lowered in Δ*nblD* (Fig. 5). If the flux into the GS/GOGAT cycle

would be similar, this decrease should extend into the 2-OG pool; however, its

increase indicates reduced 2-OG utilization in mutant cells, whereas 2-OG was

extensively used in WT to replenish the Glu pool.

It could be expected that the higher 2-OG level in $\Delta nblD$ than in the WT cells would

contribute to the stimulation of the NtcA regulon, which was observed in $\Delta nblD$

during the previous transcriptome analysis. Indeed, 3 h after the induction of N

starvation, the mRNA levels of NtcA-regulated *nblA1A2* genes were more strongly

induced compared to the WT (log$_2$ fold changes of +4.28 and +4.09 in $\Delta nblD$ and

+3.71 and +3.47 in the WT), while *gifA* and *gifB* were more strongly reduced (for

both genes, log$_2$ fold change of -4.7 in $\Delta nblD$ and -2.4 in the WT control) (19). By

comparison, 2-OG concentrations were around 3.14 mM and 1.16 mM in $\Delta nblD$ and

WT, respectively (Table S3). Hence, a higher 2-OG concentration corresponded

roughly to higher mRNA levels 3 h after the induction of N starvation. However, in

the nitrogen-replete condition we measured an ~100 fold higher level of 2-OG in

$\Delta nblD$ compared to WT but the *nblA1A2* mRNA levels were only marginally higher.

This result strongly suggests that the high accumulation of 2-OG alone is not

sufficient to induce the complete N starvation response and an additional signal can

be postulated. Furthermore, the high accumulation of 2-OG in $\Delta nblD$ already in the

+N condition can be taken as evidence that NblD, in addition to its prominent

function during N starvation, also plays some role under N-replete conditions. This

function of NblD could be related to the remodeling of phycobilisome composition,

size and number per thylakoid membrane or to the turnover of phycobilisomes

occurring under N-replete conditions.

Materials and Methods

Cultivation conditions and sampling for metabolite analysis

Synechocystis 6803 PCC-M was used as the WT and background for the construction of mutants. *NblD* knockout mutants were generated by homologous replacement of the *nblD* coding sequence with a kanamycin resistance cassette (*nptII*) (19).

Strains were maintained in copper-free BG11 supplemented with 20 mM TES PUFFERAN®, pH 7.5, under continuous white light of 50 µmol photons m^{-2} s^{-1} at 30°C.

Wild-type and $\Delta nblD$ cultures were grown to OD ~0.8 in triplicate without antibiotics before cells were pelleted by centrifugation (3,210 × *g*, 8 minutes, room temperature), washed three times in BG11 without N and resuspended in BG11 lacking a N source. After three and 24 h of incubation in N-limited conditions, the OD was recorded, and 2 ml of each culture was collected by centrifugation (9,000 × *g*, 1 minute, room temperature) and immersed in liquid nitrogen. As controls, samples were also obtained before N removal (+N). For extraction, 630 µl methanol containing carnitine (internal standard, 1 µg per extraction) was added, mixed for one minute and incubated for 5 minutes in a sonic water bath. After another 15 minutes of shaking at room temperature, 400 µl of chloroform was added, and the sample was incubated at 37°C for 5 minutes. Next, 800 µl ROTISOLV® LC-MS grade H_2O (Roth) was added, and the sample was mixed thoroughly. Precipitation was enabled by storage at -20°C for at least 2 h, and phase separation was subsequently achieved by centrifugation for 5 minutes at room temperature (16,800 × *g*). The upper polar phase was collected and dried in a speed vac for 30 minutes followed by lyophilization overnight. A mock sample without any culture was subjected to the same treatments as the other samples.

Metabolite analysis

Absolute metabolite contents were quantified on a high-performance liquid

chromatograph mass spectrometer LCMS-8050 system (Shimadzu) as described

previously (70). Dried samples were dissolved in 200 µl LCMS grade water and

filtered through 0.2-mm filters (Omnifix-F, Braun, Germany), and 4 µl of the cleared

supernatant was separated on a pentafluorophenylpropyl column (Supelco

Discovery HS FS, 3 mm, 150 3 2.1 mm).

The compounds were identified and quantified using the multiple reaction monitoring

(MRM) values given in the LC-MS/MS method package and the LabSolutions

software package (Shimadzu). Authentic standard substances (Merck) at varying

concentrations were used for calibration, and peak areas were normalized to signals

of the internal standard (carnitine). The data were further normalized to the OD_{750}

measured for each sample and corrected by signals obtained from the mock sample.

The raw data and estimations as absolute values in nanogram $\times$ ml^{-1} $\times$ OD_{750}^{-1} are

given in Table S1, statistical evaluations in Table S4. These data were converted to

molar concentrations of metabolites per total cell volume assuming an average cell

diameter of 3 µm and 10^7 cells per milliliter at $OD_{750} = 1$ (Table S3).

Funding

We appreciate the support by the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) to WRH through the graduate school MeInBio -

322977937/GRK2344 and to WRH and MH through the priority program "Small

Proteins in Prokaryotes, an Unexplored World" SPP 2002 (grants DFG HE2544/12-2

and DFG HA2002/22-2). The LC-MS equipment at the University of Rostock was

financed through the Hochschulbauförderungsgesetz program (GZ: INST 264/125-1

FUGG to MH).

Conflict of interest statement

The authors declare that they have no conflicts of interest.

Author Contributions

WRH designed the project and secured funding. VK constructed and cultivated

strains, MH and ST measured metabolite concentrations. VK, MH and WRH wrote

the manuscript with input from all authors.

References

1. Meydan S, Marks J, Klepacki D, Sharma V, Baranov PV, Firth AE, Margus T, Kefi A, Vázquez-Laslop N, Mankin AS. 2019. Retapamulin-assisted ribosome profiling reveals the alternative bacterial proteome. Mol Cell 74:481-493.e6.

2. Weaver J, Mohammad F, Buskirk AR, Storz G. 2019. Identifying small proteins by ribosome profiling with stalled initiation complexes. mBio 10.

3. Hemm MR, Weaver J, Storz G. 2020. *Escherichia coli* small proteome. EcoSal Plus 9.

4. Fromme P, Melkozernov A, Jordan P, Krauss N. 2003. Structure and function of photosystem I: interaction with its soluble electron carriers and external antenna systems. FEBS Lett 555:40–44.

5. Guskov A, Kern J, Gabdulkhakov A, Broser M, Zouni A, Saenger W. 2009. Cyanobacterial photosystem II at 2.9-A resolution and the role of quinones, lipids, channels and chloride. Nat Struct Mol Biol 16:334–342.

6. Kashino Y, Lauber WM, Carroll JA, Wang Q, Whitmarsh J, Satoh K, Pakrasi HB. 2002. Proteomic analysis of a highly active photosystem II preparation from the cyanobacterium *Synechocystis* sp. PCC 6803 reveals the presence of novel polypeptides. Biochemistry 41:8004–8012.

7. Baniulis D, Yamashita E, Whitelegge JP, Zatsman AI, Hendrich MP, Hasan SS, Ryan CM, Cramer WA. 2009. Structure-function, stability, and chemical modification of the cyanobacterial cytochrome b_6f complex from *Nostoc* sp. PCC 7120. J Biol Chem 284:9861–9869.

8. Allen JF. 2004. Cytochrome b_6f: structure for signalling and vectorial metabolism. Trends Plant Sci 9:130–137.

9. Schneider D, Volkmer T, Rögner M. 2007. PetG and PetN, but not PetL, are essential subunits of the cytochrome b_6f complex from *Synechocystis* PCC 6803. Res Microbiol 158:45–50.

10. Nowaczyk MM, Wulfhorst H, Ryan CM, Souda P, Zhang H, Cramer WA, Whitelegge JP. 2011. NdhP and NdhQ: two novel small subunits of the cyanobacterial NDH-1 complex. Biochemistry 50:1121–1124.

11. Schwarz D, Schubert H, Georg J, Hess WR, Hagemann M. 2013. The gene *sml0013* of *Synechocystis* sp. strain PCC 6803 encodes for a novel subunit of

the NAD(P)H oxidoreductase or complex I that is ubiquitously distributed among
Cyanobacteria. Plant Physiol 163:1191–1202.

12. Knoppová J, Sobotka R, Tichy M, Yu J, Konik P, Halada P, Nixon PJ, Komenda
J. 2014. Discovery of a chlorophyll binding protein complex involved in the early
steps of photosystem II assembly in *Synechocystis*. Plant Cell 26:1200–1212.

13. Komenda J, Sobotka R. 2016. Cyanobacterial high-light-inducible proteins--
Protectors of chlorophyll-protein synthesis and assembly. Biochim Biophys Acta
1857:288–295.

14. Kufryk G, Hernandez-Prieto MA, Kieselbach T, Miranda H, Vermaas W, Funk C.
2008. Association of small CAB-like proteins (SCPs) of *Synechocystis* sp. PCC
6803 with photosystem II. Photosyn Res 95:135–145.

15. Hobbs EC, Fontaine F, Yin X, Storz G. 2011. An expanding universe of small
proteins. Curr Opin Microbiol 14:167–173.

16. Baumgartner D, Kopf M, Klähn S, Steglich C, Hess WR. 2016. Small proteins in
cyanobacteria provide a paradigm for the functional analysis of the bacterial
micro-proteome. BMC Microbiol 16:285.

17. de Alvarenga LV, Hess WR, Hagemann M. 2020. AcnSP - a novel small protein
regulator of aconitase activity in the cyanobacterium *Synechocystis* sp. PCC
6803. Front Microbiol 11:1445.

18. Baier K, Nicklisch S, Grundner C, Reinecke J, Lockau W. 2001. Expression of
two *nblA*-homologous genes is required for phycobilisome degradation in
nitrogen-starved *Synechocystis* sp. PCC6803. FEMS Microbiol Lett 195:35–39.

19. Krauspe V, Fahrner M, Spät P, Steglich C, Frankenberg-Dinkel N, Maček B,
Schilling O, Hess WR. 2021. Discovery of a small protein factor involved in the
coordinated degradation of phycobilisomes in cyanobacteria. Proc Natl Acad Sci
USA 118:e2012277118.

20. Muro-Pastor MI, Reyes JC, Florencio FJ. 2005. Ammonium assimilation in
cyanobacteria. Photosynth Res 83:135–150.

21. Herrero A, Muro-Pastor AM, Flores E. 2001. Nitrogen control in cyanobacteria. J
Bacteriol 183:411–425.

22. Allen MM, Smith AJ. 1969. Nitrogen chlorosis in blue-green algae. Archiv für
Mikrobiologie 69:114–120.

23. Görl M, Sauer J, Baier T, Forchhammer K. 1998. Nitrogen-starvation-induced
chlorosis in *Synechococcus* PCC 7942: adaptation to long-term survival.
Microbiology 144:2449–2458.

24. Krasikov V, Aguirre von Wobeser E, Dekker HL, Huisman J, Matthijs HC. 2012.
Time-series resolution of gradual nitrogen starvation and its impact on
photosynthesis in the cyanobacterium *Synechocystis* PCC 6803. Physiol plant
145:426–439.

25. Sauer J, Schreiber U, Schmid R, Völker U, Forchhammer K. 2001. Nitrogen
starvation-induced chlorosis in *Synechococcus* PCC 7942. Low-level
photosynthesis as a mechanism of long-lerm survival. Plant Physiol126:233–
243.

26. Spät P, Klotz A, Rexroth S, Maček B, Forchhammer K. 2018. Chlorosis as a
developmental program in cyanobacteria: The proteomic fundament for survival
and awakening. Mol Cell Proteomics 17:1650–1669.

27. Collier JL, Grossman AR. 1992. Chlorosis induced by nutrient deprivation in
Synechococcus sp. strain PCC 7942: not all bleaching is the same. J Bacteriol
174:4718–4726.

28. Gründel M, Scheunemann R, Lockau W, Zilliges Y. 2012. Impaired glycogen
synthesis causes metabolic overflow reactions and affects stress responses in
the cyanobacterium *Synechocystis* sp. PCC 6803. Microbiology, 158:3032–
3043.

29. Meeks JC, Wolk CP, Lockau W, Schilling N, Shaffer PW, Chien WS. 1978.
Pathways of assimilation of $^{[13N]}N_2$ and $^{13}NH_4^+$ by cyanobacteria with and without
heterocysts. J Bacteriol 134:125–130.

30. Florencio FJ, García-Domínguez M, Martín-Figueroa E, Crespo JL, Navarro F,
Muro-Pastor MI, Reyes JC. 1998. Ammonium assimilation in cyanobacteria.
The regulation of the GS-GOGAT pathway, p. 3607–3612. *In* Garab, G (ed.),
Photosynthesis: Mechanisms and Effects: Volume I–V: Proceedings of the XIth
International Congress on Photosynthesis, Budapest, Hungary, August 17–22,
1998. Springer Netherlands, Dordrecht.

31. Vázquez-Bermúdez MF, Herrero A, Flores E. 2002. 2-Oxoglutarate increases
the binding affinity of the NtcA (nitrogen control) transcription factor for the
Synechococcus glnA promoter. FEBS Lett 512:71–74.

32. Tanigawa R, Shirokane M, Maeda S -i., Omata T, Tanaka K, Takahashi H.
2002. Transcriptional activation of NtcA-dependent promoters of
Synechococcus sp. PCC 7942 by 2-oxoglutarate *in vitro*. Proc Natl Acad Sci
USA 99:4251–4255.

33. Espinosa J, Forchhammer K, Burillo S, Contreras A. 2006. Interaction network
in cyanobacterial nitrogen regulation: PipX, a protein that interacts in a 2-
oxoglutarate dependent manner with PII and NtcA. Mol Microbiol 61:457–469.

34. Espinosa J, Rodriguez-Mateos F, Salinas P, Lanza VF, Dixon R, de la Cruz F,
Contreras A. 2014. PipX, the coactivator of NtcA, is a global regulator in
cyanobacteria. Proc Natl Acad Sci USA 111:E2423–E2430.

35. Forcada-Nadal A, Llácer JL, Contreras A, Marco-Marín C, Rubio V. 2018. The
PII-NAGK-PipX-NtcA regulatory axis of cyanobacteria: A tale of changing
partners, allosteric effectors and non-covalent interactions. Front Mol Biosci
5:91.

36. Forchhammer K. 2004. Global carbon/nitrogen control by PII signal transduction
in cyanobacteria: from signals to targets. FEMS Microbiol Rev 28:319–333.

37. Forchhammer K. 2008. PII signal transducers: novel functional and structural
insights. Trends Microbiol 16:65–72.

38. Giner-Lamia J, Robles-Rengel R, Hernández-Prieto MA, Muro-Pastor MI,
Florencio FJ, Futschik ME. 2017. Identification of the direct regulon of NtcA
during early acclimation to nitrogen starvation in the cyanobacterium
Synechocystis sp. PCC 6803. Nucleic Acids Res 45:11800–11820.

39. García-Domínguez M, Reyes JC, Florencio FJ. 2000. NtcA represses
transcription of *gifA* and *gifB*, genes that encode inhibitors of glutamine
synthetase type I from *Synechocystis* sp. PCC 6803. Mol Microbiol 35:1192–
1201.

40. Collier JL, Grossman AR. 1994. A small polypeptide triggers complete
degradation of light-harvesting phycobiliproteins in nutrient-deprived
cyanobacteria. EMBO J 13:1039–1047.

41. Karradt A, Sobanski J, Mattow J, Lockau W, Baier K. 2008. NblA, a key protein
of phycobilisome degradation, interacts with ClpC, a HSP100 chaperone partner
of a cyanobacterial Clp protease. J Biol Chem 283:32394–32403.

42. Baier A, Winkler W, Korte T, Lockau W, Karradt A. 2014. Degradation of
phycobilisomes in *Synechocystis* sp. PCC6803: evidence for essential formation
of an NblA1/NblA2 heterodimer and its codegradation by a Clp protease
complex. J Biol Chem 289:11755–11766.

43. Nguyen AY, Bricker WP, Zhang H, Weisz DA, Gross ML, Pakrasi HB. 2017. The
proteolysis adaptor, NblA, binds to the N-terminus of β-phycocyanin:
Implications for the mechanism of phycobilisome degradation. Photosyn Res
132:95–106.

44. Sendersky E, Kozer N, Levi M, Moizik M, Garini Y, Shav-Tal Y, Schwarz R.
2015. The proteolysis adaptor, NblA, is essential for degradation of the core
pigment of the cyanobacterial light-harvesting complex. Plant J 83:845–852.

45. Elmorjani K, Herdman M. 1987. Metabolic control of phycocyanin degradation in
the cyanobacterium *Synechocystis* PCC 6803: a glucose effect. Microbiology
133:1685–1694.

46. Hu P-P, Hou J-Y, Xu Y-L, Niu N-N, Zhao C, Lu L, Zhou M, Scheer H, Zhao K-H.
2020. The role of lyases, NblA and NblB proteins and bilin chromophore transfer
in restructuring the cyanobacterial light-harvesting complex. Plant J 102:529–
540.

47. Kiyota H, Hirai MY, Ikeuchi M. 2014. NblA1/A2-dependent homeostasis of
amino acid pools during nitrogen starvation in *Synechocystis* sp. PCC 6803.
Metabolites 4:517–531.

48. Cameron JC, Pakrasi HB. 2010. Essential role of glutathione in acclimation to
environmental and redox perturbations in the cyanobacterium *Synechocystis* sp.
PCC 6803. Plant Physiol 154:1672–1685.

49. Eisenhut M, Kahlon S, Hasse D, Ewald R, Lieman-Hurwitz J, Ogawa T, Ruth W,
Bauwe H, Kaplan A, Hagemann M. 2006. The plant-like C2 glycolate cycle and
the bacterial-like glycerate pathway cooperate in phosphoglycolate metabolism
in cyanobacteria. Plant Physiol 142:333–342.

50. Quintero MJ, Muro-Pastor AM, Herrero A, Flores E. 2000. Arginine catabolism
in the cyanobacterium *Synechocystis* sp. strain PCC 6803 involves the urea
cycle and arginase pathway. J Bacteriol 182:1008–1015.

51. Zhang H, Yang C. 2019. Arginine and nitrogen mobilization in cyanobacteria.
Mol Microbiol 111:863–867.

52. Zhang H, Liu Y, Nie X, Liu L, Hua Q, Zhao G-P, Yang C. 2018. The
cyanobacterial ornithine-ammonia cycle involves an arginine dihydrolase. Nat
Chem Biol 14:575–581.

53. Klotz A, Georg J, Bučinská L, Watanabe S, Reimann V, Januszewski W,
Sobotka R, Jendrossek D, Hess WR, Forchhammer K. 2016. Awakening of a
dormant cyanobacterium from nitrogen chlorosis reveals a genetically
determined program. Curr Biol 26:2862–2872.

54. Orthwein T, Scholl J, Spät P, Lucius S, Koch M, Macek B, Hagemann M,
Forchhammer K. 2021. The novel PII-interactor PirC identifies
phosphoglycerate mutase as key control point of carbon storage metabolism in
cyanobacteria. Proc Natl Acad Sci USA 118:e2019988118.

55. Arteni AA, Ajlani G, Boekema EJ. 2009. Structural organisation of
phycobilisomes from *Synechocystis* sp. strain PCC6803 and their interaction
with the membrane. Biochim Biophys Acta 1787:272–279.

56. Lahmi R, Sendersky E, Perelman A, Hagemann M, Forchhammer K, Schwarz
R. 2006. Alanine dehydrogenase activity is required for adequate progression of
phycobilisome degradation during nitrogen starvation in *Synechococcus*
elongatus PCC 7942. J Bacteriol 188:5258–5265.

57. Wang W-Y, Gough SP, Kannangara CG. 1981. Biosynthesis of Δ-
aminolevulinate in greening barley leaves IV. Isolation of three soluble enzymes
required for the conversion of glutamate to Δ-aminolevulinate. Carlsberg Res
Commun 46:243.

58. Kannangara CG, Gough SP, Oliver RP, Rasmussen SK. 1984. Biosynthesis of
Δ-aminolevulinate in greening barley leaves VI. Activation of glutamate by
ligation to RNA. Carlsberg Res Commun 49:417.

59. Schön A, Krupp G, Gough S, Berry-Lowe S, Kannangara CG, Söll D. 1986. The
RNA required in the first step of chlorophyll biosynthesis is a chloroplast
glutamate tRNA. Nature 322:281–284.

60. O'Neill GP, Peterson DM, Schön A, Chen MW, Söll D. 1988. Formation of the
chlorophyll precursor delta-aminolevulinic acid in cyanobacteria requires
aminoacylation of a tRNAGlu species. J Bacteriol 170:3810–3816.

61. Rieble S, Beale SI. 1988. Transformation of glutamate to delta-aminolevulinic
acid by soluble extracts of *Synechocystis* sp. PCC 6803 and other oxygenic
prokaryotes. J Biol Chem 263:8864–8871.

62. Zhang S, Bryant DA. 2011. The tricarboxylic acid cycle in cyanobacteria.
Science 334:1551–1553.

63. Xiong W, Brune D, Vermaas WFJ. 2014. The γ-aminobutyric acid shunt
contributes to closing the tricarboxylic acid cycle in *Synechocystis* sp. PCC
6803. Mol Microbiol 93:786–796.

64. Kopf M, Klähn S, Scholz I, Matthiessen JKF, Hess WR, Voß B. 2014.
Comparative analysis of the primary transcriptome of *Synechocystis* sp. PCC
6803. DNA Res 21:527–539.

65. Chellamuthu V-R, Ermilova E, Lapina T, Lüddecke J, Minaeva E, Herrmann C,
Hartmann MD, Forchhammer K. 2014. A widespread glutamine-sensing
mechanism in the plant kingdom. Cell 159:1188–1199.

66. Forchhammer K. 2007. Glutamine signalling in bacteria. Front Biosci 12:10–
2741.

67. Klähn S, Bolay P, Wright PR, Atilho RM, Brewer KI, Hagemann M, Breaker RR,
Hess WR. 2018. A glutamine riboswitch is a key element for the regulation of
glutamine synthetase in cyanobacteria. Nucleic Acids Res 46:10082–10094.

68. Jiang Y-L, Wang X-P, Sun H, Han S-J, Li W-F, Cui N, Lin G-M, Zhang J-Y,
Cheng W, Cao D-D, Zhang Z-Y, Zhang C-C, Chen Y, Zhou C-Z. 2018.
Coordinating carbon and nitrogen metabolic signaling through the
cyanobacterial global repressor NdhR. Proc Natl Acad Sci USA 115:403–408.

69. Klähn S, Orf I, Schwarz D, Matthiessen JKF, Kopka J, Hess WR, Hagemann M.
2015. Integrated transcriptomic and metabolomic characterization of the low-
carbon response using an *ndhR* mutant of *Synechocystis* sp. PCC 6803. Plant
Physiol 169:1540.

70. Reinholdt O, Schwab S, Zhang Y, Reichheld J-P, Fernie AR, Hagemann M,
Timm S. 2019. Redox-regulation of photorespiration through mitochondrial
thioredoxin o1. Plant Physiol 181:442–457.

Figures

Figure 1. Response of the *Synechocystis* 6803 WT and Δ*nblD* mutant during the
response to N starvation. **A.** Phenotype under nitrogen-replete (+N) conditions and
after 24 h of nitrogen depletion (-N). **B.** Spectrometric measurements of cultures
normalized to OD_{750}.

Figure 2. **A.** Total soluble amino acid content and **B.** total soluble metabolite content

(non-proteinogenic amino acids and important intermediates) of wild-type (blue) and

Δ*nblD* (red) cultures in nitrogen-replete (+N) and nitrogen-depleted BG11 (-N) after 3

h and 24 h. For each time point and strain, triplicates were measured, and

significance ($p < 0.05$) was calculated by t-test; n.s. = not significant. The raw data

are given in Table S2.

Figure 3. Development of amino acid content (in mM for Glu and µM for all other) in WT (blue) and ΔnblD mutant (red) in nitrogen replete (+N) and nitrogen deplete (-N) medium after 3 h and 24 h. Amino acids are grouped together according to the observed differences between wild type and mutant as follows: Difference already observed in +N conditions; no difference between strains; early difference (after 3 h); later differences (24 h). Significance was calculated with a two-sample t-test with unequal variance (Welch's t-test; $p < 0.05$ = *; $p < 0.01$ = ** and $p < 0.001$ = ***) for

756   the development in each strain and between the strains at corresponding time points

757   (see Table S4 for details).

Figure 4. Phycobilisome individual amino acid (aa) composition (in %) plotted against their increase or decrease observed in relation to the initial values measured in nitrogen sufficient conditions. **A, C.** WT after 3 h of N limitation; **B, D.** WT after 24 h of N limitation. **E**, **F.** Data for Δ*nblD* mutant after 3 and 24 h. Phycobilisome composition was calculated based on the structure description by Arteni et al. (55), assuming a common complex with 6 rods each with 3 hexameric phycocyanin discs and a tricylindrical allophycocyanin core. A linear trend line shows very low correlation between the variables, and a poor R-squared value indicates high variation of the data in relation to this line.

Figure 5. Comparison of metabolite content (in mM for Glu, citrate, 3PGA, malate and 2-OG, in µM for all other) in WT (blue) and Δ*nblD* deletion mutant (red) in nitrogen replete (+N) and nitrogen deplete (-N) medium after 3 h and 24 h shown alongside their biosynthetic pathways. Succinate was below detection limit (b.d.) in the Δ*nblD* sample under nitrogen-replete conditions. Significance was calculated with the two-sample t-test with unequal variance (Welch's t-test; $p < 0.05$ = *; $p < 0.01$ = ** and $p < 0.001$ = ***) for the development in each strain and between the strains at corresponding time points (details in Table S4).

Supplementary Tables

Table S1. LC-MS/MS data for 18 amino acids (Cys and Asn are missing because they were below detection limit), several organic acids and GSH. Retention time (r) was individually monitored with 2 standards (line 4 and 5) and the average peak area for 1 ng (line 7), which were used to determine the ng per sample. The carnitine standard (columns E, F, G, H) was used to generate the Factor IS (internal standard), which is implemented in the normalization of the peak area for each sample.

Table S2. Total amount of all amino acids and the other soluble metabolites investigated in this study. Average, standard deviation and significance calculated by two-sample t-test (Welch's t-test) are visualized in figure 2.

Table S3. Conversion in molar concentrations of metabolites in µM (Glu in mM) per total cell volume assuming an average cell diameter of 3 µm and 10^7 cells per milliliter at $OD_{750} = 1$.

Table S4. Welch's t-test (two-sided, inequal variation) for evaluating significant differences in the amino acids and organic acids between the strains and the development in the strains. Significant differences are marked in red ($p < 0.05$), boxed ($p < 0.01$) and bold italics ($p < 0.001$).