Combined computational and cellular screening identifies synergistic inhibition of SARS-CoV-2 by lenvatinib and remdesivir

Marie O. Pohl[a,*], Idoia Busnadiego[a,*], Francesco Marrafino[b, c], Lars Wiedmer[b], Annika Hunziker[a], Sonja Fernbach[a], Elena Omori[b], Benjamin G. Hale[a], Amedeo Caflisch[b,§] & Silke Stertz[a,§]

[a]Institute of Medical Virology, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland.

[b]Department of Biochemistry, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland.

[c]Department of Pharmacy, University of Salerno, Fisciano SA, 84084, Italy.

[*] These authors contributed equally.

[§] To whom correspondence should be addressed: caflisch@bioc.uzh.ch or
                                                    stertz.silke@virology.uzh.ch

ABSTRACT

Rapid repurposing of existing drugs as new therapeutics for COVID-19 has been an

important strategy in the management of disease severity during the ongoing SARS-

CoV-2 pandemic. Here, we screened by high-throughput docking 6,000 compounds

within the DrugBank library for their potential to bind and inhibit the SARS-CoV-2 3CL

main protease, a chymotrypsin-like enzyme that is essential for viral replication. For 19

candidate hits, parallel *in vitro* fluorescence-based protease-inhibition assays and Vero-

CCL81 cell-based SARS-CoV-2 replication-inhibition assays were performed. One hit,

diclazuril (an investigational anti-protozoal compound), was validated as a SARS-CoV-2

3CL main protease inhibitor *in vitro* (IC50 value of 29 µM) and modestly inhibited SARS-

CoV-2 replication in Vero-CCL81 cells. Another hit, lenvatinib (approved for use in

humans as an anti-cancer treatment), could not be validated as a SARS-CoV-2 3CL

main protease inhibitor *in vitro*, but serendipitously exhibited a striking functional synergy

with the approved nucleoside analogue remdesivir to inhibit SARS-CoV-2 replication in

Vero-CCL81 cells. Lenvatinib is a broadly-acting host receptor tyrosine kinase (RTK)

inhibitor, but the synergistic effect with remdesivir was not observed with other approved

RTK inhibitors (such as pazopanib or sunitinib), suggesting that the mechanism-of-

action is independent of host RTKs. Furthermore, time-of-addition studies revealed that

lenvatinib/remdesivir synergy probably targets SARS-CoV-2 replication subsequent to

host-cell entry. Our study shows that combining computational and cellular screening is

an efficient means to identify existing drugs with repurposing potential as antiviral

compounds. Future studies should aim at understanding and optimizing the

lenvatinib/remdesivir synergistic mechanism as a therapeutic option.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) entered the human

population in late 2019 and rapidly spread around the globe causing a devastating

pandemic with significant impacts on human health and the worldwide economy [1, 2].

Infection with SARS-CoV-2 results in a respiratory disease, COVID-19, that is mild in

most individuals or even asymptomatic. However, in vulnerable patients, in particular in

individuals with compromised immune systems, such as the elderly or

immunosuppressed, the virus can cause a severe life-threatening pneumonia [2-4].

Thus, large efforts have been undertaken to develop vaccines and antiviral drugs for the

prevention and treatment of COVID-19. While several vaccines have been developed

and licensed at unprecedented speeds, the development of antiviral drugs has not yet

been successful [5-8]. Remdesivir is the only antiviral drug that is currently approved

and used to treat COVID-19 patients in many countries, but its efficacy has been

questioned by recent results from a large-scale clinical trial [9, 10]. Thus, additional

efforts are urgently needed to fill this gap.

Early on in the pandemic, drug development approaches focused on the repurposing of

drugs approved for other conditions as a means to accelerate the process of bringing

drug candidates to market [11, 12]. One of the viral drug targets that has gained

particular interest is the SARS-CoV-2 protease, 3CL M^{pro}, a chymotrypsin-like protease

that cleaves the two polyproteins, pp1a and pp1ab, and is essential for viral replication

[13, 14]. 3CL M^{pro} is considered a promising target for drug development as its structure

has been solved and there is no homologous human protease [15, 16]. Moreover,

protease inhibitors have been developed successfully to treat HIV-1 and hepatitis C

virus infections [17, 18]. While several inhibitors of 3CL M^{pro} have already been

identified, no drug targeting 3CL M^{pro} has been approved yet [15, 19-22]. Here, we

computationally screened a library of approved drugs for novel 3CL M^{pro} inhibitors. The

identified drug candidates were then tested for their antiviral potential in an *in vitro* 3CL

M^{pro} protease-inhibition assay, and in a Vero cell-based virus replication assay using

authentic SARS-CoV-2. While two potential inhibitors (diclazuril and ramelteon)

identified *in silico* possessed modest antiviral activity, we found that lenvatinib, an

inhibitor of receptor tyrosine kinases such as vascular endothelial growth factor

receptors, displayed strong synergistic inhibition of SARS-CoV-2 with the approved drug

remdesivir.

79   **RESULTS**

To identify novel inhibitors of the SARS-CoV-2 3CL M^{pro}, we performed an *in silico*

screening campaign of the Drugbank library which contains approximately 6000

approved drugs [23]. The docking was carried out by the computer program SEED

(details in Methods section) using two holo structures of 3CL M^{pro} (PDB codes 5r82 and

5re9). The binding pocket was considered rigid during docking. The calculation of the

binding energy makes use of a force field-based methodology. Upon filtering and

ranking, the top 19 compounds were purchased for experimental validation (Fig. 1A, B).

An *in vitro* fluorescence-based protease-inhibition assay with recombinant 3CL M^{pro}

protease was employed to measure the potential inhibition of enzymatic activity by each

compound. The positive control, GC376, was used as reference compound [22]. An IC$_{50}$

(inhibitor concentration that results in 50% reduction of activity with respect to buffer-only

solution) value of 316 nM was measured for GC376, which is consistent with a reported

value of 190 nM [22]. First, single-dose measurements were carried out at 100 μM for

each compound (Fig. 1C). For only two compounds a substantial reduction (> 40% with

respect to DMSO-only) of 3CL M^{pro} activity was measured. In follow-up dose-response

experiments with these two compounds, diclazuril showed an IC_{50} of 29 μM, and

oprea1_409650 showed an IC_{50} of 100 μM (Fig. 1D, E). The binding mode of diclazuril

as predicted by docking is shown in Figure 1F.

In parallel, we tested the 19 predicted 3CL M^{pro} protease inhibitors for their potential to

inhibit SARS-CoV-2 replication in Vero CCL81 cells. Following pretreatment with the

different inhibitors for 2 h, Vero CCL81 cells were infected with SARS-CoV-2 isolate

SARS-CoV-2/human/Switzerland/ZH-395-UZH-IMV5/2020 at a multiplicity of infection

(MOI) of 0.01 PFU/cell and viral titers were determined at 24 h post-infection.

Remdesivir, a known inhibitor of SARS-CoV-2 [24], was included as a positive control. In

parallel, inhibitor-treated cells were left uninfected and cell viability was determined. With

the exception of compounds 9-aminocampthecin and 1-deazaadenosine, cell viability

was only minimally affected by treatment with the inhibitors (Fig. 2A). In line with

previous reports, SARS-CoV-2 replication was inhibited by remdesivir in a dose-

dependent manner, thus validating our screening assay (Fig. 2B). We also observed

significant inhibition of virus replication with three of the predicted protease inhibitors,

lenvatinib, ramelteon and diclazuril (Fig. 2B). Of note, diclazuril, which was the most

potent protease inhibitor in the *in vitro* protease assay, was also the strongest hit in the

screen for inhibition of SARS-CoV-2 replication. Oprea1, which also displayed inhibition

in the protease assay did not reduce viral replication significantly. While the inhibitory

effects of the three hit compounds were robust and reproducible across the replicates of

the SARS-CoV-2 screening assay, the degree of inhibition was only around 50-60% and

thus modest (Fig. 2C). In order to test whether the inhibitors would show greater antiviral

potential when combined with an established inhibitor of SARS-CoV-2, we tested them

in combination with low concentrations of remdesivir. While ramelteon and diclazuril did

not increase virus inhibition by remdesivir, the combination of remdesivir and lenvatinib

showed a very strong synergistic effect: the low concentration of remdesivir used

reduced viral titers approximately 3-fold, but in combination with lenvatinib, SARS-CoV-2

titers were reduced >1000-fold (Fig. 2D), which was not due to cytotoxic effects (Supp.

Fig. 1A). When the three hit compounds were tested in human Calu-3 cells for their

impact on SARS-CoV-2 replication, we did not detect any effect on cell viability (Supp.

Fig. 1B) but also no inhibition of virus growth, either by the inhibitors alone or in

combination with low doses of remdesivir (Supp. Fig. 1C). Thus, the striking synergistic

effect of lenvatinib in combination with remdesivir appears to be specific to Vero CCL81

cells.

As lenvatinib did not inhibit SARS-CoV-2 3CL M^{pro} in the *in vitro* protease-inhibitor assay

at the concentration used in cell culture-based assays (Fig. 1C), we speculated that its

function as an inhibitor of receptor tyrosine kinases (RTKs) might contribute to the

observed synergy with remdesivir. We therefore selected pazopanib and sunitinib as

additional RTK inhibitors to test for synergy with remdesivir, as they exhibit a similar

inhibition profile to lenvatinib, with all three shown to inhibit vascular endothelial growth

factor receptors (VEGFR1, VEGFR2 and VEGFR3), platelet-derived growth factor

receptor (PDGFR-α/-β), and c-Kit with comparable specificity and potency [25, 26].

When taking cell viability into consideration, only lenvatinib displayed a clear synergistic

effect with remdesivir (Fig. 2E, Supp. Fig. 1D) suggesting that the observed synergistic

inhibitory effect is not linked to inhibition of VEGFR-, PDGFR- or c-Kit induced signaling.

Next, we performed a time-of-addition experiment, in which we treated cells with

remdesivir, lenvatinib, or the combination of both inhibitors, at 1 h prior to infection, 1 h

post infection, or 4 h post infection. At 24 h post infection, supernatants were harvested

and infectious virus titers determined by plaque assay. As expected from an RNA

polymerase inhibitor, remdesivir retained its antiviral potential if given post viral entry:

treatment with remdesivir 1 h post infection resulted in a similar inhibition of virus

replication as treatment of cells prior to infection (Fig. 2F). In addition, treatment with

remdesivir at 4 h post infection still led to a substantial decrease in viral replication. The

synergistic effect of lenvatinib and remdesivir was highly similar for pretreatment and 1 h

post-treatment conditions, both resulting in potent inhibition of virus replication.

Furthermore, the 4 h post infection treatment condition resulted in a less potent, but still

substantial, decrease in viral titers by more than 10-fold (Fig. 2F). Thus, our data

suggest that the synergistic effect of lenvatinib and remdesivir could be due to more

potent inhibition of the viral transcription and/or replication processes rather than any

effects on earlier stages of the viral replication cycle.

156   **DISCUSSION**

Our approach of combining *in silico* and *in vitro* screenings of an approved drug library

for inhibitors of the SARS-CoV-2 3CL M^{pro} protease identified diclazuril as a low

micromolar inhibitor of the 3CL M^{pro} protease, and three drugs (diclazuril, ramelteon, and

lenvatinib) that reduced viral replication in Vero CCL81 cells. Moreover, lenvatinib

showed an unexpectedly strong synergy with remdesivir. This shows that the approach

used here can identify compounds with antiviral potential.

There are several caveats to the approach adopted in this study. Of note, we only tested

the compounds at a single 10 µM concentration, and thus an immediate option to see

stronger inhibition of virus replication would be to test higher concentrations of the

inhibitors. However, this might result in the identification of additional compounds with

antiviral potential but only in a non-physiologically achievable range of concentrations.

Thus, extensive compound modification and optimization for increased antiviral potential

would be required in subsequent studies and would slow immediate roll-out of a

repurposed drug. Given that our goal was to rapidly identify approved drugs as

treatment options for COVID-19 and circumvent the lengthy process of progressing

inhibitors to approval, we set the threshold at 10 µM and did not consider higher drug

concentrations. Options to improve the yield of our screening approach could be to use

an alternative structure of 3CL M^{pro} that was subsequently solved by room temperature

X-ray crystallography [27]. It reveals the ligand-free structure of the active site and the

conformation at near-physiological temperature, and was suggested to be a better target

for *in silico* screening for inhibitors of 3CL M^{pro} [27]. Moreover, additional libraries of

approved drugs could be included, for example IUPHAR/BPS

(https://www.guidetopharmacology.org/) and ChEMBL (https://www.ebi.ac.uk/chembl/),

although they overlap substantially with DrugBank.

A second caveat is the apparent cell-type specificity of synergy observed between

lenvatinib and remdesivir: this effect was clear in Vero CCL81cells, but not in human

Calu-3 cells. One possible explanation could be that lenvatinib inhibits a Vero cell-

specific target that is not inhibited in Calu-3 cells. This could either be due to differential

expression levels of the respective target or could be related to differences in target

sequence and structure between the human and the African green monkey version of

the target. As the conversion of remdesivir to its active metabolite (remdesivir

triphosphate) is less efficient in Vero CCL81 cells as compared to Calu-3 cells [24], the

synergistic effect of lenvatinib with remdesivir could be linked to the metabolic

conversion of remdesivir. If lenvatinib treatment led to a higher conversion rate, and thus

to higher levels of the active metabolite of remdesivir it could be hypothesized that in

Calu-3 cells, where the conversion of remdesivir to remdesivir triphosphate is already

efficient, an increase in conversion efficiency would not result in a boost of the antiviral

potential of remdesivir. However, it is thus far unclear how lenvatinib would affect the

metabolic conversion of remdesivir. Given that lenvatinib did not significantly inhibit the

SARS-CoV-2 3CL M^{pro} protease in the *in vitro* protease assay, the known targets of

lenvatinib (i.e. receptor tyrosine kinases (RTKs), such as VEGFR, PDGFR or c-Kit),

could be the relevant drug targets for inhibition of virus replication and synergy with

remdesivir. We did not observe synergy for two other RTK inhibitors with comparable

RTK specificity and potency, and thus the observed synergy is probably not mediated by

inhibition of VEGFRs, PDGFRs or c-Kit, which are targeted by all three inhibitors. Thus,

future work should aim to reveal the mechanism behind the synergistic action of

remdesivir and lenvatinib. Such studies on the Vero cell-specific synergy could reveal

broader insights into the metabolism and antiviral mechanism of nucleoside analogs and

thus aid the development of antiviral drugs.

207   **MATERIALS AND METHODS**

In silico screen for 3CL M[pro] inhibitors. The evaluation of the binding energy by the

program SEED makes use of a force field-based energy function. Protein and ligand

electrostatic desolvation penalties are approximated by a continuum dielectric treatment

and the generalized Born model [28-30]. The partial charges and vdW parameters for

the atoms in the protein and in the ligands were taken from the CHARMM36 all-atom

force field and the CHARMM general force field (CGenFF), respectively [31-33].

Importantly, CHARMM36 and CGenFF employ a consistent paradigm for the

determination of partial charges and vdW parameters. The dielectric discontinuity

surface was approximated by the molecular surface. The dielectric constant was

assigned a value of 2.0 and 78.5 for the volume occupied by the solute and solvent,

respectively. Default SEED parameters were used. The nearly 6000 approved and

experimental drugs in DrugBank with more than 17 non-hydrogen atoms were docked

by SEED. This filter on molecular sizes was employed to avoid the selection of very

small molecules which cannot provide sufficient affinity. SEED offers three methods for

the actual placement of the fragments: (1) polar docking in which fragments are

positioned and oriented such that at least one favorable hydrogen bond with a residue in

the binding site is formed, (2) apolar docking in which fragments are placed in the

hydrophobic regions of the binding site, and (3) a combination of (1) and (2). The latter

placement procedure was used. The docking was carried out on two holo structures of

3CL M[pro] (PDB codes 5r82 and 5re9) with high resolution. The residue Glu166 was

selected for the definition of the binding pocket. For PDB 5r82 no water molecules were

retained, while for PDB 5re9 one buried water molecule was retained. Two energy filters

were used: total binding energy < -10 kcal/mol and van der Waals efficiency < -1.0

kcal/mol per non-hydrogen atom. Hydrogen bonding penalty was calculated for each

compound, according to the method described by Zhao and Huang [34]. If newly formed

hydrogen bonds between the ligand and the protein are not in good geometry or

desolvated donors/acceptors are not involved into new hydrogen bonds, an enthalpic

loss arises due to uncompensated breakage of hydrogen bonds with waters. The

hydrogen bonding penalty index is calculated by checking both the solvation and

hydrogen bonding state of donors/acceptors at the protein-ligand interface. A value of 1

in penalty corresponds to about 1.5 kcal/mol in binding free energy. This penalty can

then be integrated into total binding energy values according to the formula SEED total

energy + hydrogen bonding penalty*1.5kcal/mol. Compounds were then ranked by

recalculated total energy and heavy atom count. Up to top five compounds for each

heavy atom count tier were selected and experimental phase drugs were removed,

focusing on approved drugs. The resulting compounds were visually inspected, favoring

compounds that established H-bonds, and 19 compounds in total were selected.

***In vitro* protease inhibition assay.** The in vitro inhibition assay was based on

fluorescence and was carried out according to the specifications of the vendor (Reaction

Biology, Catalogue #: Mpro). The assay measures the inhibitory activity of compounds

that compete with a known peptide substrate for purified 3CL Mpro. The peptide

contains a quencher and a donor group, thus fluorescence is quenched in an intact

peptide. Upon cleavage of the peptide by 3CL Mpro the donor fragment generates

fluorescence which is quantified by a luminometer. The fluorogenic substrate used is

[NH2-C(EDANS)VNSTQSGLRK(DABCYL)M-COOH]. The positive control is GC376

which was employed as reference compound. Single-dose screenings were performed

at 100 µM.

Compound screening for inhibitors of SARS-CoV-2. To test the selected compounds

from the docking analysis for antiviral activity against SARS-CoV-2, 1.2×10^4 Vero CCL81

cells were seeded in DMEM supplemented with 10% fetal calf serum (FCS), 100 U/ml of

penicillin, and 100 µg/ml of streptomycin in 96-well plates and incubated overnight at

37°C and 5% CO_2. For the infection and cell viability experiments, compound dilutions of

10 µM and a final DMSO concentration of 0.1% were used. As negative control, 0.1%

DMSO was used while 10, 5 and 2.5 µM remdesivir (at a final DMSO concentration of

0.1%) served as positive controls. The day after seeding, cells were washed with PBS

followed by incubation with the individual compounds at a concentration of 10 µM diluted

in Opti-MEM (Gibco) for 2 h at 37°C and 5% CO_2. Next, medium was removed and cells

were infected with SARS-CoV-2 (SARS-CoV-2/human/Switzerland/ZH-395-UZH-

IMV5/2020) at a multiplicity of infection (MOI) of 0.01 PFU/cell diluted in PBS

supplemented with 0.3% BSA, 1 mM Ca^{2+}/Mg^{2+}, 100 U/ml penicillin, and 100 µg/ml

streptomycin [35]. Cells were then incubated for 1 h at 37°C and 5% CO_2. Following

infection, the inoculum was removed and DMEM supplemented with 10 µM of the

respective compound (or the positive or negative control), 100 U/ml penicillin,

100 µg/ml streptomycin, 0.3% BSA, 20 mM HEPES, 0.1% FCS, and 0.5 µg/ml

TPCK-treated trypsin was added to the cells. After incubation for 24 h at 37°C and 5%

CO_2, cell supernatants were harvested and virus titer was determined by plaque assay.

In parallel to the infection experiments, cell viability following incubation of Vero CCL81

cells with the respective compounds was assessed using the same experimental set-up

as for the infection experiments but instead of infection with SARS-CoV-2, cells were

mock-infected with PBS supplemented with 0.3% BSA, 1 mM Ca^{2+}/Mg^{2+}, 100 U/ml

penicillin, and 100 µg/ml streptomycin for 1 h at 37°C and 5% CO_2. Cell viability was

determined 24 h after the addition of compound-containing DMEM using the CellTiterGlo

kit (Promega) and a luminescence plate reader (Perkin Elmer). The infection and cell

viability experiments for the compound screen were performed in duplicates in three

independent experiments.

For the analysis of the compound screen, all titers were normalized to the negative

control (0.1% DMSO) and log2-transformed. Next, the z-score was calculated by

dividing the difference of the normalized and transformed titer of a given sample and the

average normalized and transformed titer of the negative control by the standard

deviation of the normalized and transformed negative control [36]. The average z-score

of the three independent replicates for a given sample was used for hit selection.

Compounds associated with a z-score of smaller than -2 were considered as hits for

further analysis. Compounds associated with an average reduction in cell viability of

more than 20% relative to the negative control were not considered as hits.

Testing for synergistic inhibition of SARS-CoV-2. Vero CCL81 cells were seeded in

96 well plates at a density of 1.2×10^4 cells/well and incubated overnight. Cells were

washed once with PBS and incubated for 2 h at 37°C and 5% CO_2 with Opti-MEM

(Gibco) containing 5 or 2.5 μM remdesivir alone or in combination with 10 μM ramelteon

(PubChem ID 208902), lenvatinib (PubChem ID 9823820), diclazuril (PubChem ID

25273624), pazopanib (LC laboratories, P-6706) or sunitinib (Selleckchem, S1042) at a

final concentration of 0.1 % DMSO. DMSO 0.1 % was used as negative control. Cells

were   infected   with   SARS-CoV-2   (SARS-CoV-2/human/Switzerland/ZH-395-UZH-

IMV5/2020) at a MOI of 0.01 in PBS supplemented with 0.3% BSA, 1 mM Ca^{2+}/Mg^{2+},

100 U/ml penicillin, 100 μg/ml streptomycin. After 1 h at 37°C and 5% CO_2, cells were

washed once in PBS and the indicated compounds were added in Dulbecco's modified

Eagle's medium (DMEM; Gibco) supplemented with 100 U/ml penicillin, 100 μg/ml

streptomycin, 0.3% BSA, 20 mM HEPES, 0.1 % FCS and 0.5 μg/ml TPCK-treated

trypsin. Samples were taken at 48 h post infection and supernatants were stored at -

80°C prior to titer determination by plaque assay. In parallel to the infection experiments,

cell viability was assessed as indicated in the compound screen section above. All

experiments were performed in duplicates in three independent experiments.

Time of addition experiments. Vero CCL81 cells were seeded in 96 well plates at a

density of 1.2 x 10^4 cells/well. After incubation overnight, cells were washed once with

PBS and incubated for 1 h at 37°C and 5% CO_2 with Opti-MEM (Gibco) containing the

indicated inhibitors. DMSO 0.1 % was used as negative control. Cells were infected with

SARS-CoV-2 (SARS-CoV-2/human/Switzerland/ZH-395-UZH-IMV5/2020) at a MOI of

0.01 PFU/cell in PBS supplemented with 0.3% BSA, 1 mM Ca^{2+}/Mg^{2+}, 100 U/ml

penicillin, 100 μg/ml streptomycin. Following infection, the inoculum was removed and

DMEM supplemented with 100 U/ml penicillin, 100 μg/ml streptomycin, 0.3% BSA,

20 mM HEPES, 0.1% FCS, and 0.5 μg/ml TPCK-treated trypsin was added to the

cells. To test for time of addition effect, the indicated compounds were added either 1 or

4 h post infection instead of 1 h before infection in parallel samples. After incubation for

24 h at 37°C and 5% CO_2, cell supernatants were harvested, and virus titer was

determined by plaque assay. In parallel to the infection experiments, cell viability was

assessed as indicated above. All experiments were performed in duplicates in three

independent experiments.

CONFLICT OF INTEREST

The author(s) declare that there are no conflicts of interest.

ACKNOWLEDGMENTS

Work in the Stertz and Hale groups is supported by the Swiss National Science

Foundation (grant numbers 31003A_176170 (to SSt) and 31003A_182464 (to BGH)). The

Caflisch group is supported by an Excellence grant of the Swiss National Science

Foundation (310030B_189363).

FIGURE LEGENDS

Fig. 1 *In silico* screen reveals novel inhibitors of the SARS-CoV-2 3CL M^{pro}

protease. A) Schematic overview of the *in silico* screen by high-throughput docking for

repurposing known drugs. **B)** List of 19 compounds identified as potential inhibitors of

3CL M^{pro}. **C)** The 19 compounds were tested in an *in vitro* fluorescence-based assay at

a concentration of 100 µM for their potential to inhibit 3CL M^{pro}. Mean values from three

independent results are shown with error bars representing standard deviations. GC376

is the positive control. **D-E)** The two compounds that showed a reduction of enzymatic

activity of more than 40% (relative to DMSO solution) in the *in vitro* protease assay,

oprea1_409650 (D) and diclazuril (E), were tested at varying concentrations to obtain a

dose-response curve. Average values from three independent experiments are plotted

with error bars representing standard deviations. **F)** Predicted binding mode of diclazuril

(carbon atoms in magenta) to 3CL M^{pro} (ribbon model in cyan with the side chains of the

catalytic dyad in yellow). The hydrogen bonds between diclazuril and the backbone polar

groups of Glu166 are shown (dotted lines).

Fig. 2. Lenvatinib acts synergistically with remdesivir to inhibit SARS-CoV-2. A)

Effect of inhibitor-treatment on cell viability. Vero CCL81 cells were pre-treated for 2

h with 10 µM of a putative 3CL M^{pro} inhibitor or with DMSO 0.1% or RDV (10, 5, and 2.5

µM) as negative and positive control, respectively. Then, cells were mock-infected for 1

h followed by incubation in the presence of the respective inhibitors or controls. 24 h

after mock-infection cell viability was assessed. Shown are means and standard

deviation of cell viability normalized to the DMSO control (depicted as a square) of three

independent experiments each performed in duplicates. RDV-treated cells are depicted

as triangles. Compounds reducing cell viability more than 20% compared to the DMSO

control (marked in red) were excluded from the selection. **B) Effect of inhibitor-**

treatment on SARS-CoV-2 growth. Experimental set-up as in A) but cells were

infected with SARS-CoV2 (MOI=0.01) for 1 h followed by incubation in the presence of

the respective inhibitors or controls. After 24 h of incubation, cell supernatants were

harvested and viral growth was assessed by plaque assay. Virus titers of three

independent experiments, each performed in duplicate, were normalized to the DMSO

control, log2-transformed and converted into z-scores. Z-score of DMSO and RDV-

treated cells are depicted as square and triangle, respectively. Excluded inhibitors are

marked in red. A z-score of smaller than -2 was used for the selection of inhibitors for

follow-up. **C) Normalized virus titers for selected inhibitors.** Experimental set-up as

in B). Shown are virus titers normalized to DMSO control of the selected inhibitors from

B). Depicted are means and standard deviation of three independent experiments, each

performed in duplicates. **D) Effect of combination of selected inhibitors with RDV on**

SARS-CoV-2 growth. Vero CCL81 cells were pre-treated for 2 h with 10 μM of ramelteon, lenvatinib, or diclazuril combined with either 5 or 2.5 μM RDV. DMSO 0.1% or RDV (5 and 2.5μM) were used as negative and positive control, respectively. Following infection with SARS-CoV-2 (MOI=0.01) for 1 h, cells were incubated in the presence of the respective inhibitors or controls. After 48 h of incubation, cell supernatants were harvested and viral growth was assessed by plaque assay. Shown are mean virus titers and standard deviation of three independent experiments, each performed in duplicate. **E) Effect of VEGFR inhibitors and RDV on SARS-CoV-2 growth.** Vero CCL81 cells were pre-treated for 2 h with lenvatinib (10 μM), pazopanib (10 μM), or sunitinib (1 μM) in combination with RDV (5 μM). DMSO 0.1% or RDV (5 μM) were used as negative and positive controls, respectively. Following infection with SARS-CoV2 (MOI=0.01) for 1 h, cells were incubated in the presence of the respective inhibitors or controls. After 48 h of incubation, cell supernatants were harvested and viral growth was assessed by plaque assay. Shown are mean virus titers and standard deviation of three independent experiments, each performed in duplicate. **F) Effect of time of addition on the synergistic reduction of SARS-CoV-2 growth by lenvatinib and RDV.** Vero CCL81 cells were treated either 1 h before (-1h), 1 h after (+1h) or 4 h after (+4h) infection with SARS-CoV-2 (MOI=0.01) with 10 μM of lenvatinib combined with either 5 or 2.5 μM RDV. DMSO 0.1% or RDV (10, 5 and 2.5μM) were used as negative and positive control, respectively. 24 h post infection, cell supernatants were harvested and viral growth was assessed by plaque assay. Shown are mean virus titers and standard deviation of three independent experiments, each performed in duplicates. LoD, limit of detection.

REFERENCES

1.   **Dong E, Du H, Gardner L**. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis* 2020;20(5):533-534.

2.   **Zhu N, Zhang D, Wang W, Li X, Yang B et al.** A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* 2020;382(8):727-733.

3.   **Wu Z, McGoogan JM**. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA* 2020;323(13):1239-1242.

4.   **Huang C, Wang Y, Li X, Ren L, Zhao J et al.** Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395(10223):497-506.

5.   **Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A et al.** Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med* 2020;383(27):2603-2615.

6.   **Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S et al.** Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med* 2021;384(5):403-416.

7.   **Voysey M, Clemens SAC, Madhi SA, Weckx LY, Folegatti PM et al.** Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet* 2021;397(10269):99-111.

8.   **Krammer F**. SARS-CoV-2 vaccines in development. *Nature* 2020;586(7830):516-527.

9.   **Consortium WHOST, Pan H, Peto R, Henao-Restrepo AM, Preziosi MP et al.** Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results. *N Engl J Med* 2021;384(6):497-511.

10.   **Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS et al.** Remdesivir for the Treatment of Covid-19 - Final Report. *N Engl J Med* 2020;383(19):1813-1826.

11.   **Riva L, Yuan S, Yin X, Martin-Sancho L, Matsunaga N et al.** Discovery of SARS-CoV-2 antiviral drugs through large-scale compound repurposing. *Nature* 2020;586(7827):113-119.

12.   **Gordon DE, Hiatt J, Bouhaddou M, Rezelj VV, Ulferts S et al.** Comparative host-coronavirus protein interaction networks reveal pan-viral disease mechanisms. *Science* 2020;370(6521).

13.   **V'Kovski P, Kratzel A, Steiner S, Stalder H, Thiel V**. Coronavirus biology and replication: implications for SARS-CoV-2. *Nat Rev Microbiol* 2020.

14.   **Anand K, Ziebuhr J, Wadhwani P, Mesters JR, Hilgenfeld R**. Coronavirus main proteinase (3CLpro) structure: basis for design of anti-SARS drugs. *Science* 2003;300(5626):1763-1767.

15. **Zhang L, Lin D, Sun X, Curth U, Drosten C et al.** Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved alpha-ketoamide inhibitors. *Science* 2020;368(6489):409-412.

16. **Douangamath A, Fearon D, Gehrtz P, Krojer T, Lukacik P et al.** Crystallographic and electrophilic fragment screening of the SARS-CoV-2 main protease. *Nat Commun* 2020;11(1):5047.

17. **Farnik H, Zeuzem S**. New antiviral therapies in the management of HCV infection. *Antivir Ther* 2012;17(5):771-783.

18. **Wensing AM, van Maarseveen NM, Nijhuis M**. Fifteen years of HIV Protease Inhibitors: raising the barrier to resistance. *Antiviral Res* 2010;85(1):59-74.

19. **Drayman N, Jones KA, Azizi SA, Froggatt HM, Tan K et al.** Drug repurposing screen identifies masitinib as a 3CLpro inhibitor that blocks replication of SARS-CoV-2 in vitro. *bioRxiv* 2020.

20. **Hung HC, Ke YY, Huang SY, Huang PN, Kung YA et al.** Discovery of M Protease Inhibitors Encoded by SARS-CoV-2. *Antimicrob Agents Chemother* 2020;64(9).

21. **Fu L, Ye F, Feng Y, Yu F, Wang Q et al.** Both Boceprevir and GC376 efficaciously inhibit SARS-CoV-2 by targeting its main protease. *Nat Commun* 2020;11(1):4417.

22. **Vuong W, Khan MB, Fischer C, Arutyunova E, Lamer T et al.** Feline coronavirus drug inhibits the main protease of SARS-CoV-2 and blocks virus replication. *Nat Commun* 2020;11(1):4282.

23. **Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A et al.** DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Res* 2018;46(D1):D1074-D1082.

24. **Pruijssers AJ, George AS, Schafer A, Leist SR, Gralinksi LE et al.** Remdesivir Inhibits SARS-CoV-2 in Human Lung Cells and Chimeric SARS-CoV Expressing the SARS-CoV-2 RNA Polymerase in Mice. *Cell Rep* 2020;32(3):107940.

25. **Capozzi M, De Divitiis C, Ottaiano A, von Arx C, Scala S et al.** Lenvatinib, a molecule with versatile application: from preclinical evidence to future development in anti-cancer treatment. *Cancer Manag Res* 2019;11:3847-3860.

26. **Suyama K, Iwase H**. Lenvatinib: A Promising Molecular Targeted Agent for Multiple Cancers. *Cancer Control* 2018;25(1):1073274818789361.

27. **Kneller DW, Phillips G, O'Neill HM, Jedrzejczak R, Stols L et al.** Structural plasticity of SARS-CoV-2 3CL M(pro) active site cavity revealed by room temperature X-ray crystallography. *Nat Commun* 2020;11(1):3202.

28. **Majeux N, Scarsi M, Caflisch A**. Efficient electrostatic solvation model for protein-fragment docking. *Proteins* 2001;42(2):256-268.

29. **Majeux N, Scarsi M, Apostolakis J, Ehrhardt C, Caflisch A**. Exhaustive docking of molecular fragments with electrostatic solvation. *Proteins* 1999;37(1):88-105.

30.   **Scarsi M, Apostolakis J, Caflisch A**. Continuum electrostatic energies of
macromolecules in aqueous solutions. *Journal of Physical Chemistry A*
1997;101(43):8098-8106.

31.   **MacKerell AD, Bashford D, Bellott M, Dunbrack RL, Evanseck JD et al.** All-
atom empirical potential for molecular modeling and dynamics studies of proteins.
Journal of Physical Chemistry B 1998;102(18):3586-3616.

32.   **MacKerell AD, Jr., Feig M, Brooks CL, 3rd**. Improved treatment of the protein
backbone in empirical force fields. *J Am Chem Soc* 2004;126(3):698-699.

33.   **Vanommeslaeghe K, Hatcher E, Acharya C, Kundu S, Zhong S et al.**
CHARMM General Force Field: A Force Field for Drug-Like Molecules Compatible with
the CHARMM All-Atom Additive Biological Force Fields. *Journal of Computational*
Chemistry 2010;31(4):671-690.

34.   **Zhao HT, Huang DZ**. Hydrogen Bonding Penalty upon Ligand Binding. *Plos One*
2011;6(6).

35.   **Busnadiego I, Fernbach S, Pohl MO, Karakus U, Huber M et al.** Antiviral
Activity of Type I, II, and III Interferons Counterbalances ACE2 Inducibility and Restricts
SARS-CoV-2. *mBio* 2020;11(5).

36.   **Konig R, Chiang CY, Tu BP, Yan SF, DeJesus PD et al.** A probability-based
approach for the analysis of large-scale RNAi screens. *Nat Methods* 2007;4(10):847-
849.

Figure 1

Internal name	Pubchem ID	Name
LWI_3CL_001	5323	Sulfadimethoxine
LWI_3CL_002	5215	Sulfadiazine
FMA_3CL_001	10127622	Selumetinib
LWI_3CL_003	5328	Sulfamethizole
LWI_3CL_004	71157	Nefiracetam
LWI_3CL_005	5336	Sulfapyridine
FMA_3CL_002	6918485	Isavuconazole
FMA_3CL_003	208902	Ramelteon
FMA_3CL_004	10090485	Galunisertib
FMA_3CL_005	9823820	Lenvatinib
FMA_3CL_006	82148	Agomelatine
FMA_3CL_007	72402	9-Aminocamptothecin
FMA_3CL_008	447980	Oprea1_409650
FMA_3CL_009	213053	Cimicoxib
LWI_3CL_006	5326	Sulfameter
FMA_3CL_010	25273624	Diclazuril
FMA_3CL_011	159738	1-Deazaadenosine
FMA_3CL_012	115348	2-Iodomelatonin
LWI_3CL_007	5503	Tolazamide

Figure 2