Molecular basis for recognition of the Group A Carbohydrate backbone by the PlyC

streptococcal bacteriophage endolysin

Harley King[1,^], Sowmya Ajay Castro[2,^], Juan A. Bueren-Calabuig[3,^], Amol Arunrao

Pohane[4], Cynthia M. Scholte[1], Vincent A. Fischetti[5], Natalia Korotkova[4,6] ,Fabio

Zuccotto[3], Daniel C. Nelson[1] and Helge C. Dorfmueller[2,*]

[1]Institute for Bioscience and Biotechnology Research, University of Maryland, Rockville, MD,

USA

[2]Division of Molecular Microbiology, School of Life Sciences, University of Dundee, Dundee, UK

[3]Drug Discovery Unit, Wellcome Centre for Anti-Infectives Research, Division of Biological

Chemistry, School of Life Sciences, University of Dundee, Dundee, UK

[4]Department of Microbiology, Immunology and Molecular Genetics, University of Kentucky,

Lexington, Kentucky, USA

[5]Laboratory of Bacterial Pathogenesis and Immunology, The Rockefeller University, New York,

NY, USA

[6]Department of Molecular and Cellular Biochemistry, University of Kentucky, Lexington, KY, USA

[^]equal contribution

[*]Correspondence: Helge C. Dorfmueller, h.c.z.dorfmueller@dundee.ac.uk

Keywords: Streptococcus; rhamnose, polysaccharide, cell wall, bacteriophage, endolysin,

pyogenes

Abbreviations (alphabetical order)

Cell wall-binding domain (CBD), enzymatically active domain (EAD), Glucose (Glc), glycerol

phosphate (GroP), Glycosyl Hydrolase (GlyH), Group A Carbohydrate (GAC), Group A

Streptococcus (GAS), Group A-variant Streptococcus (GAVS), Group C Carbohydrate (GCC),

Group C Streptococcus (GCS), Group G Carbohydrate (GGC), *N*-acetyl-glucosamine (GlcNAc),

N-acetyl-galactosamine (GalNAc), N-acetyl-muramic acid (MurNAc), Peptidoglycan (PG),

polyrhamnose (pRha), Rhamnose (Rha), *S. dysgalactiae subsp. equisimilis* (SDSE), *S. mutans*

serotype c carbohydrate (SCC)

Abstract

Endolysins are peptidoglycan (PG) hydrolases that function as part of the bacteriophage (phage) lytic system to release progeny phage at the end of a replication cycle. Notably, endolysins alone can produce lysis without phage infection, which offers an attractive alternative to traditional antibiotics. Endolysins from phage that infect Gram-positive bacterial hosts contain at least one enzymatically active domain (EAD) responsible for hydrolysis of PG bonds and a cell wall binding domain (CBD) that binds a cell wall epitope, such as a surface carbohydrate, providing some degree of specificity for the endolysin. Whilst the EADs typically cluster into conserved mechanistic classes with well-defined active sites, relatively little is known about the nature of the CBDs and only a few binding epitopes for CBDs have been elucidated. The major cell wall components of many streptococci are the polysaccharides that contain the polyrhamnose (pRha) backbone modified with species-specific and serotype-specific glycosyl side chains. In this report, using molecular genetics, microscopy, flow cytometry and lytic activity assays, we demonstrate the interaction of PlyCB, the CBD subunit of the streptococcal PlyC endolysin, with the pRha backbone of the cell wall polysaccharides, Group A Carbohydrate (GAC) and serotype *c*-specific carbohydrate (SCC) expressed by the Group A Streptococcus and *Streptococcus mutans*, respectively. Molecular dynamics simulations reveal a previously unidentified binding pocket that is regulated by a gatekeeper residue and uncover that a previously reported inactive PlyC mutant is locked into a 'closed' conformation. Docking studies with the short GAC backbone oligosaccharides expose potential protein-carbohydrate interactions and are consistent with PlyCB binding to the unmodified pRha or pRha decorated with the GAC side chains.

Introduction

Endolysins are bacteriophage-encoded PG hydrolases that normally function from within the cell to lyse the bacterial host, releasing progeny phage and completing the phage lifecycle [1]. However, the lytic activity of endolysins can be harnessed for antimicrobial use due to their ability to equally lyse bacteria when applied exogenously, without infection by a parental phage. Due to their direct lytic action on target PG, endolysins are not affected by efflux pumps, alterations in metabolism, or other mechanisms of antibiotic resistance, making them ideal candidates for development against multi-drug resistant organisms [2-4]. Notably, at least three endolysins, some of which are active against methicillin-resistant *Staphylococcus aureus*, are currently being evaluated in human clinical trials for their antimicrobial activity (reviewed in [5]).

Most endolysins, and in particular those from phage that infect Gram-positive bacterial hosts, are comprised of modular domains. An enzymatically active domain (EAD) is generally found in the N-terminal region, while a cell wall-binding domain (CBD) is located in the C-terminal region [6]. As the name implies, the EAD is a catalytic domain that is responsible for cleaving specific bonds in the PG, the nature of which is dependent on the mechanistic class of the EAD. Occasionally, endolysins contain two EADs, although both are not necessarily active. The CBD binds at high affinity [7] to a cell wall-specific epitope and was suggested to dictate genus, species and serovar-specificity of the endolysin. The CBD targets may be surface carbohydrates, wall teichoic acids linked to the Gram-positive bacterial cell wall, or the PG itself [8].

The endolysin now known as PlyC is one of the first described endolysins and remains one of the most studied. In 1934, Alice Evans noted a "nascent lysis" activity derived from streptococcal phage lysates on streptococcal strains that were not sensitive to the phage itself [9]. By 1957, Krause had determined that the phage used by Evans was specific for Group C

Streptococci (GCS), but an "enzyme" produced by the phage could lyse Groups A, A-variant,

and C Streptococci (GAS, GAVS, and GCS, respectively) [10]. These findings were confirmed

by Maxted later the same year and extended to include Group E Streptococci (GES) [11]. In

2001, PlyC, then referred to as the streptococcal C_1 lysin, became the first endolysin to be used

therapeutically, protecting mice from GAS challenge in a nasopharyngeal model [2].

Subsequent studies revealed that PlyC is a structurally unconventional endolysin, which is not

encoded by a single gene as found for all other endolysins described to date. Rather, PlyC is a

nine-subunit holoenzyme encoded by two distinct genes, *plyCB* and *plyCA*, within a

polycistronic operon [12]. Eight PlyCB subunits self-assemble into a ring structure and form the

basis of the CBD that binds the streptococcal surface. A single PlyCA subunit contains two

distinct EADs separated by an extended α-helical linker region, which interfaces with the N-

terminal residues of the PlyCB octamer [13]. The PlyCA EADs consist of a glycosyl hydrolase

(GH) domain and a cysteine, histidine-dependent aminohydrolase/peptidase (CHAP) domain.

The very high lytic activity of PlyC relative to other endolysins is attributed to

synergy/cooperativity between the two EADs.

Although the EADs of PlyCA have been extensively characterized with respect to specificity of

PG bonds they cleave, active-site residues, and their synergistic activity, it is unclear how PlyCB

recognizes PG of GAS, GCS and GES. Similar to other Gram-positive bacteria, the plasma

membrane of streptococci is surrounded by thick cell wall that consists of a complex network of

PG with covalently attached polysaccharides. Rebecca Lancefield utilized the unique

immunogenicity of the surface carbohydrates in β-hemolytic streptococci to subsequently

separate them into serogroups [14]. *S. pyogenes* is categorized as Group A Streptococcus,

whilst *S. dysgalactiae subsp equisimilis* (SDSE) produce at least two types of carbohydrates

and are annotated as Group C and G Streptococci, respectively [15]. Pioneering work by many

researchers have revealed that the cell wall polysaccharides of GAS, GCS and *S. mutans*,

consist of a pRha backbone modified with species-specific and serotype-specific glycosyl side

chains (Fig. 1A) [16, 17]. In GAS, GCS and *S. mutans* serotype *c*, the polysaccharides termed

the Lancefield group A carbohydrate (GAC), group C carbohydrate (GCC), and SCC,

respectively, have a conserved repeating →3)α-Rha(1→2)α-Rha(1→ di-saccharide backbone

[16, 17]. The β-*N*-acetylglucosamine (GlcNAc) side chains are attached to the 3-position of the

α-1,2 linked rhamnose (Rha) in GAC [16, 18, 19]. The GCC side chains have two *N*-acetyl-

galactosamine (GalNAc) residues attached to the same hydroxyl of Rha [16, 18, 20]. SCC

carries the α-glucose (Glc) side chains attached to the 2-position of the α-1,3 linked Rha in SCC

[17]. Additionally, the side chains of GAC and SCC are decorated in parts with the glycerol

phosphate (GroP) moiety [21]. The GAC and SCC biosynthetic pathways are encoded by the

12-gene loci *gacABCDEFGHIJKL* and *sccABCDEFGHMNPQ*, respectively. The molecular

mechanisms of GAC and SCC biosynthesis have been the focus in a number of recent studies

[21-25].

Remarkably, extracted GAC is known to partially inhibit the lytic actions of PlyC [26].

Furthermore, a GAS mutant carrying the unmodified GAC (lacking the GlcNAc side chains)

displayed enhanced susceptibility to PlyC [25], implicating PlyCB in recognition of the pRha

backbone. Here, we identify the pRha backbone of GAC as the definite minimal binding epitope

for PlyCB. Molecular dynamics simulations and docking studies based on our findings further

establish a model of interaction between PlyCB and selective surface carbohydrates in

streptococci.

Materials and Methods

Bacterial strains, growth conditions and media

Streptococcus pyogenes strain D471 was propagated on solid media in plates containing Todd-

Hewitt broth supplemented with yeast extract (0.2%) (THY) and agar (1.4%) or in liquid THY as

139 described by Gera et al [27]. *S. mutans* wild type (Xc) and mutants were grown in Todd-Hewitt

140 broth with 1% yeast extract. All cultures were grown without antibiotics and without aeration at

141 37°C. *E. coli* genotypes DH5α (NEB, cat. No. C2988J), DH10α (Thermo Fisher, cat No.

142 18297010), BL21 (NEB, cat No. C2530H) and Origami 2 (Millipore Sigma, cat. No. 71346) were

143 used for routine plasmid propagation or protein expression and grown in Lysogeny Broth (LB)

144 medium supplemented with either 50 µg/ml kanamycin, 100 µg/ml ampicillin or 35 µg/ml

145 chloramphenicol as needed.

147 Bacterial strains *E. coli* CS2775 were transformed with pRGP1 plasmid [28], *gacABCDEFG* [23]

148 to produce pRha or empty plasmid control (pHD0131). The bacterial cells were grown overnight

149 in LB containing erythromycin (150 µg/ml) at 37°C and used next day for whole cell Western

150 blots and FACS and microscopy analysis.

152 **Recombinant expression and purification of PlyCB$_{WT}$ and PlyCB$_{R66E}$**

153 PlyCB$_{WT}$ (GenBank ID: NC_004814.1:7517-7735) was expressed from a pBAD24 vector and

154 purified from BL21 cells as previously described [12]. In brief, the culture was grown in LB and

155 induced with 0.25% L-arabinose at OD$_{600}$ ~1.2-1.4 (Alfa Aesar, cat no. A11921). Cultures were

156 grown at 37°C with shaking at 180 RPM for 3-4 hours, centrifuged at 4500 x g and resuspended

157 in phosphate-buffered saline (PBS). Lysis was performed by a French press (1800 psi).

158 Benzonase (Millipore Sigma, cat. No. 70746-3) was added and the lysate incubated at room

159 temperature with rotation for 20-30 min. The lysate was centrifuged at 20,000 x g for 20 min and

160 the cleared lysate was passed through a 0.45-µm filter and loaded onto a XK-26/20 column

161 (Cytiva) with 30-35 ml ceramic hydroxyapatite (Bio-Rad, cat no. 1582000). PlyCB$_{WT}$ was eluted

162 from the column with three column volumes of 1 M sodium phosphate buffer (pH 7.2). Protein

163 was subsequently dialyzed in PBS, 10% glycerol and stored at -80°C until use. Protein

164 purification of PlyCB$_{R66E}$ was performed as PlyCB$_{WT}$ [13]. The fluorescent labeling of PlyCB$_{WT}$

and PlyCB$_{R66E}$ was performed using the manufacturer's recommended guidelines (Thermo

Fisher, cat. No. A20174).

Purification of GAC

GAS were grown overnight in THY media at 37˚C. Cultures were centrifuged at 4,500 x g.

Pellets were washed and resuspended in 40 ml distilled water per each original liter of media

used and combined in an 800 μl beaker. 22.5 ml 4N sodium nitrite (5 ml per liter of culture) was

added to the beaker in addition to 22.5 ml glacial acetic acid (5 ml per liter of culture). An orbital

shaker was used to gently mix the beaker for 15 min in a hood. The mixture was centrifuged in

500 ml bottles at 8000 x g for 15 min. The supernatant was decanted to a new beaker and

neutralized with 1M sodium hydroxide. The total solution, about 300 ml, was filtered with a 0.45-

micron filter assembly. 50-50 ml aliquots were deposited in a 3.5 kDa membrane and dialyzed in

a 4-liter beaker overnight with water. The following day, the solution was concentrated using an

Amicon 400 ml stirred cell (model 8400) filter assembly with a 76 mm diameter Ultracel® 5 kDa

ultrafiltration disc (Millipore Sigma, cat. No. PLCC07610) for 2 hours with 60 psi. The ~10 ml

concentrate was loaded onto an S-100 column for final purification. Fractions were assayed for

Rha, lyophilized and stored at 4°C until use.

Calculation of pRha concentration

A modified protocol by Edgar *et al.* [21] based on a protocol from DuBois *et al.* [29] was used to

determine Rha concentration in purified GAC. Briefly, anthrone reagent was prepared by

dissolving 0.2% w/w anthrone in H_2SO_4. Eighty microliters of aqueous samples or standards

containing either GAC or L-Rha at known concentrations were added to a 1.5 ml microfuge

tube. To this same tube, 320 μL of the anthrone reagent was added. Samples were boiled at

98°C for 10 min in a heat block. Samples were cooled to room temperature, transferred to a

quartz plate, and the absorbance at 580 nm was recorded using a spectrophotometer. Rha

concentration was interpolated using an L-Rha standard curve.

Precipitation of PlyCB with GAC

PlyCB$_{WT}$ and PlyCB$_{R66E}$ samples were defrosted from storage at -80°C. Lyophilized GAC was

resuspended in PBS. Both proteins and GAC were added to a 3.5 kDa dialysis membrane and

dialyzed overnight in PBS. Protein concentrations were determined using a NanoDrop

spectrophotometer (Thermo Fisher ND-2000) at 280 nm and were diluted to 5 mg/ml. The GAC

was also assayed and diluted with PBS to 1.6 mg/ml. One-hundred microliters of proteins and

100 µl of GAC or PBS were mixed in a 250 µl quartz plate and allowed to incubate without

shaking at room temperature. Visible precipitate formed in samples in 5-8 minutes. After

recording the precipitate at 340 nm using a Spectramax® M5 (Molecular Devices)

spectrophotometer, the total sample volume was transferred to a 1.5 ml microfuge tube.

Samples were centrifuged at 14,000 x g to pellet the precipitate and supernatants were

transferred to new 1.5 ml microfuge tubes. Two-hundred microliters of 8M urea was added to

the pellet. Pellets were resuspended and 5 µl of either pellet or supernatant were added to 40 µl

water with 8 µl 6x Laemmli buffer with DTT. Samples were boiled at 98°C for 8 min, and then

12.5 µl were loaded onto a 7.5% SDS-PAGE and run for 32 min at 200V. Proteins were

visualized using Coomassie stain.

Lysis assay

A turbidity reduction assay was used to ascertain strain sensitivity to PlyC. This assay was

performed as previously described [30], except PlyC was used at 2 µM. Eight technical

replicates were performed.

Sensitivity of streptococcal species to PlyC-mediated lysis was analyzed using a wide range of

clinical isolates: 1) GAS isolates: M1$_{UK}$, WT 5448 strain [M1], delta*gacI* 5448 strain [d*gacI*]; 2)

GCS isolate: stC74A.0; 3) SDSE_*gac* isolates: stG245.0, stG652.0. stG485.0; GGS: stG6.0,

stG485.0 and 4) *S. mutans* serotype c were used as negative controls. Briefly, all streptococcal

strains were grown in THY at 37˚C overnight in 5% CO_2, except for *S. mutans*, which was grown

in THB media. Next day, the bacterial cells were inoculated in 1:100 fresh media and grown until

mid-logarithmic phase (OD_{600} 1.0). The cells were washed in PBS and resuspended to an OD_{600}

of 2.0. In a 96-well plate, to a 100 µl of bacterial cells, 100 µl of PlyC [1 µg/ml] was added and

immediately read at an absorbance of OD_{600}. The obtained values were standardized by

subtracting from the background values. The data is plotted using GraphPad Prism version 9.

SDS-PAGE and blotting analysis

PlyCB$_{WT}$ binding to recombinant *E. coli* expressing pRha was conducted using blot analysis.

Briefly, the lysate from the overnight cultures was analyzed in 20% tricine gels. SDS-PAGE and

protein transfers were performed according to manufactures instructions, Atto Ae-6050 Mini Gel

chamber and Novex protein separation from Thermo Fisher, respectively. The PVDF

membranes were blocked with 5% non-fat dry milk with Tris-Buffered Saline, 0.1% Tween® 20

detergent prior to incubation with PlyCB$_{WT}$ labelled with Alexa Fluor® 647 (1:1000) for one hour

at room temperature. Goat anti-rabbit GAC antibodies conjugated with IRDye® 800CW were

used as a positive control (abcam ab216773). The resulting blots were imaged using the Licor

Odyssey FC Imaging System. All the blots were processed in parallel under the same

conditions.

Microscopy

Microscopic analysis of *E. coli* bacteria was performed using cells that were grown overnight in

LB supplemented with antibiotics at 37˚C and diluted 1:100 the next day and regrown until

OD_{600} reached 0.5. The cells were washed twice with PBS for 5 min at 10,000 rpm and stained

with 1:1000 dilution of PlyCB^{AF647} and left for 20 minutes on ice in the darkness. Prior to fixing

the cells with 4% paraformaldehyde, the cells were washed again with PBS. The fixed cells

were mounted on 1% agarose coated microscopic slides and viewed under the CY5 channel on

a fluorescent Deltavision widefield microscope.

FACS analysis

E. coli cells were grown overnight as described above and diluted 1:100 the next day, grown at

37˚C and used for the assay at OD_{600} = 0.5. The cells were washed with PBS and probed with

PlyCB^{AF647} or PlyCB$_{R66E}$^{AF645} at 1:1000 dilution at 1:1000 dilution. Anti-GAC antibodies

conjugated with FITC (ABIN238144, antibodies-online, titer 1:50) were used as a positive

control. The samples were incubated for 20 minutes on ice at dark conditions. The cells were

washed twice with PBS at 14,000 rpm for 5 minutes and fixed with 4% paraformaldehyde. BD

LSRFortessa Flow Cytometry software was used to analyse the samples and the data

interpretation was conducted with FlowJo^{TM} software v10.6.2.

PlyC hydrolysis of sacculi

S. mutans wild-type (WT) and the mutant strain sacculi were obtained by the SDS-boiling

procedure [25] followed by four washes each with 1 M NaCl and distilled water. The sacculi

were resuspended in PBS to OD_{600} of 1.0 and incubated with PlyC (5 µg/ml) in a 96-well plate.

The lysis was monitored after 10, 20, 30, 40, 50 and 60 min as a decrease in OD_{600}. Results

were reported as fold change in OD_{600} of the sacculi incubated with PlyC vs the sacculi

incubated without PlyC.

Molecular Dynamics

Monomers of PlyCB$_{WT}$ (PDB 4F87) and the PlyCB$_{R66E}$ mutant (PDB 4ZRZ) were simulated by

Molecular Dynamics (MD). The systems for MD simulations were prepared with the utility LEaP,

which is integrated in the suite of programs AMBER 16 [31]. The ff14SB force field [32] was

used. The N- and C-termini of the proteins were capped with an acetyl (ACE) and methylated

amino group (NME), respectively. Each simulated system was immersed in a water box (TIP3P

water model) and neutralized by adding the appropriate number of counterions. This was

followed by steepest-descent energy minimization to remove steric clashes. MD simulations

were performed using the pmemd.cuda module of AMBER 16. The cut-off distance for the non-

bonded interactions was set to 10Å. The periodic boundary conditions were used. Electrostatic

interactions were treated using the smooth particle mesh Ewald method [33]. The SHAKE

algorithm was applied to all bonds involving hydrogen atoms, and a time step of 2 fs was used

throughout [34]. Each energy minimized system was heated to 300K, equilibrated for 10 ns, and

further simulated for 2 µs without any restraints. Protein structures and MD trajectories were

visually inspected and analyzed using the molecular visualization programs PyMOL [35] and

VMD [36].

Binding cavity identification and druggability assessment

The site recognition software SiteMap, implemented in the Schrodinger suite of programs [37],

was used to investigate the cavities on the PlyCB wild type (PDB 4F87) and R66E mutant (PDB

4ZRZ) proteins surface, in terms of physicochemical properties (hydrophobic/hydrophilic

nature), volume, exposure, and enclosure. Based on those properties, an overall "SiteScore"

was generated providing an estimate value of the druggability of the pocket. Using the default

settings, scores of >= 0.8 define the limit between drug-binding and non-drug-binding cavities.

Molecular docking

A low energy 3D conformation of Rha di- and tri-saccharides was generated using LigPrep in

the Schrödinger platform [37]. The binding of the ligands in the Y28 gated cavity identified by

MD in the wild-type structure was evaluated by molecular docking. The program Glide [37], part

of the Schrodinger platform, was used in extra-precision mode with post-docking minimization.

No distance or hydrogen-bond constraints were applied. A 20Å cubic box centered on the

centroid of the residues Leu9, Phe10, His11, Thr12, Ser17, Tyr20 and Ile34 was used to

generate the docking grid. The default settings were modified to increase the conformational

sampling of rings by increasing the energy window to 6.0 kcal/mol.

Results & Discussion
301
Pathogenic streptococci producing the GAC are susceptible to PlyC

A major component of the GAS cell wall is the GAC, building approximately 50% of the cell wall

by weight [38]. The GAC is universally conserved amongst all isolated GAS strains on the basis

of the gene cluster sequence conservation [39]. A number of *S. dysgalactiae subsp. equisimilis*

(SDSE) isolates, naturally belonging to Group G Streptococci (GGS), have been reported to

have undergone homologous recombination and replaced their Group G Carbohydrate (GGC)

gene cluster in parts with the GAC gene cluster [40-42]. We therefore expanded the previously

reported PlyC streptococci cell lysis assay used by Nelson *et al.* [2] to investigate those new

isolates named SDSE_*gac*. We also tested if PlyC was able to lyse a selection of GAS

serotypes including a newly emerged isolate M1$_{UK}$ [43], and included negative controls GGS

isolates and *S. mutans* serotype C (Fig. 1B). In agreement with the published literature, all

tested GAS serotypes are susceptible to PlyC lysis and the two GGS isolates are resistant to

PlyC. The GGC does not contain the GAC, GCC and SCC pRha backbone with →3)α-

Rha(1→2)α-Rha(1→ di-saccharide repeats, but an alternating Rha-GalNAc carbohydrate [44]

(Fig. 1A). All three SDSE_*gac* isolates tested have inherited parts of the *gac* gene cluster and

produce the GAC instead of the GGC [41]. Strikingly, the SDSE_*gac* strains are all sensitive to

PlyC treatment. The fact that PlyC is able to lyse SDSE_*gac* strains that express the GAC, but

PlyC does not lyse the isogenic GGS strains producing the GGC strongly suggests that the

GAC is a critical component of PlyC recognition and subsequent activity.

Importantly, all strains tested in this study that are susceptible to PlyC lysis produce a cell wall

polysaccharide that contains the pRha backbone and a β-linked sugar substituent on the α1,2-

linked Rha (Fig 1A). We therefore suggest that the pRha backbone with and without a side

chain are both vital ligands to assist PlyC activity and the new SDSE_*gac* isolates will also be

susceptible to PlyC treatment due to production of the GAC.

Purified GAC precipitates PlyCB - but not PlyCB$_{R66E}$

The lysis assay of GAS cells, and in particular of the SDSE_*gac* variants, suggests that either

the ubiquitous pRha or GAC in GAS cells is the ligand for PlyCB. We propose that the PlyCB

octameric CBD binds GAC and/or the GAC pRha backbone. We tested this hypothesis by

investigating the binding of PlyC to partially purified GAC. We hypothesized if the GAC was able

to precipitate PlyCB, an interaction of the two systems must have occurred [45]. As a negative

control, we employed the previously published inactive mutant PlyCB$_{R66E}$, which lost the ability

to bind to GAS cells [13]. The purified proteins were incubated with the extracted GAC, and

precipitation was monitored at 340 nm, a standard wavelength for measuring protein

aggregation [46, 47] (Fig. 2A, B). Whilst keeping the PlyCB concentration constant, we varied

the concentration of GAC. Within five minutes at room temperature the solution became turbid,

suggesting aggregation (Fig. 2A). When the PlyCB concentration was kept constant and the

GAC concentration was varied, the turbidity correlated with PlyCB concentration in a dose

dependent manner, suggesting that PlyCB requires GAC to aggregate. Importantly, PlyCB did

not self-aggregate when no GAC was added in the assay. Furthermore, no aggregation was

detected when PlyCB$_{R66E}$ was incubated with purified GAC (Fig. 2B). To demonstrate the

presence of PlyCB in the precipitates, we analyzed the soluble and pellet fractions (Fig. 2C, D).

A higher yield of aggregated PlyCB was found in the pelleted samples when compared to the

soluble fraction (Fig. 2D). A similar precipitation effect was observed when we varied the PlyCB

concentration and kept the GAC concentration constant (Fig. 2E, F), demonstrating that both

species are necessary for an interaction.

When the L-Rha monosaccharide was mixed with PlyCB, no precipitation was observed,

suggesting that the GAC, or a moiety within the GAC, is a specific ligand required for PlyCB

precipitation.

PlyCB binds to recombinantly produced pRha backbone

The purified GAC from bacteria contains a mixture of carbohydrates, including the fully

decorated GAC with GroP[21] and a small proportion of the polysaccharide lacking the side

chains [48]. PlyC is able to lyse a number of GAS mutants including GAVS and d*gacI*_M1 [2,

11, 25, 49], that decorate the cell wall with the unmodified GAC lacking the side chains (Fig. 1A,

B), suggesting that the pRha backbone of the GAC is the minimal carbohydrate structure

required for PlyCB binding. To test this hypothesis, we recombinantly produced the pRha

backbone in *E. coli* cells. We and others have previously reported that the *S. mutans*

sccABCDEFG gene cluster, when transformed into *E. coli* cells, functionally produces the pRha

backbone attached to the lipid A [23, 28]. Additionally, to understand if PlyCB recognizes a

specific pRha backbone, we engineered *E. coli* cells expressing the GAC gene cluster

gacABCDEFG required for the GAC pRha production. *E. coli* cells carrying an empty plasmid

were used as a negative control.

Next, we investigated the binding of PlyCB conjugated with Alexa Fluor® 647 (PlyCB^{AF647}) to an

E. coli total cell lysate expressing the pRha backbone of the SCC or GAC, respectively (Fig.

3A). The blotted membranes were incubated with PlyCB^{AF647}, and a positive interaction between

PlyCB^{AF647} and the *E. coli* produced pRha is visualized as a prominent band around 40 kDa.

The size of the band agrees with the band detected by anti-GAC antibodies that were previously

reported to recognize pRha [23] (Fig. 3B). Importantly, PlyCB^{AF647} and GAC antibodies do not

interact with the cell lysate of *E. coli* expressing an empty plasmid (Fig. 3 A, B). We further

confirmed the ability of PlyCB to bind to *E. coli* cells decorated with the pRha by fluorescent

microscopy (S. F1). Only cells that produce the pRha are detected by the PlyCB^{AF647}, in

agreement with the results of the blot analysis.

To gather additional evidence that PlyCB interacts with the pRha backbone, we established a

flow cytometry assay to analyse the binding of PlyCB^{AF647} to pRha-producing *E. coli*. Unstained

E. coli cells that express pRha or carry an empty plasmid sort in the identical range (Fig. 4A).

The GAC antibodies label exclusively the cells producing pRha (Fig. 4A). A similar pattern of the

GAC antibodies binding is observed when the cells were incubated with PlyCB^{AF647} (Fig. 4B).

Contrary, the PlyCB$_{R66E}$^{AF645} mutant protein does not bind to *E. coli* cells, and PlyCB^{AF647} does

not interact with the cells expressing an empty vector (Fig. 4B). Taken together, these data

provide the first definitive evidence that the pRha backbone of GAC and SCC is a binding

receptor of the PlyCB octameric subunit.

PlyC lyses engineered *S. mutans* producing the GAC

Despite the fact that the SCC pRha backbone is identical to the GAC, *S. mutans* is resistant to

PlyC lysis (Fig. 1A). To get a better understanding why *S. mutans* is resistant to PlyC, we

compared PlyC-induced lysis of the sacculi purified from *S. mutans* WT and a number of mutant

strains producing different SCC variants (Fig. 5). First, we examined the Δ*sccH* mutant

producing the GroP-deficient SCC [21]. Similar to *S. mutans* WT, Δ*sccH* was resistant to PlyC-

mediated lysis (Fig. 5). Second, we tested the *sccN* deletion mutant, Δ*sccN*, that is deficient in

the enzyme required for generation of the Glc side chains [50]. A time dependent lysis is

observed for Δ*sccN*. Expression of the WT copy of *sccN* in Δ*sccN* (the Δ*sccN*:p*sccN* strain) fully

restored the resistance of the bacteria to PlyC (Fig. 5). These observations clearly suggest that

PlyC is able to bind to the *S. mutans* cells producing the unmodified pRha backbone, and the

Glc side chains in SCC hinders PlyC binding. We then investigated whether the addition of the

GAC GlcNAc side chains to the pRha backbone affects sensitivity of the engineered *S. mutans*

sacculi to PlyC-induced lysis. We expressed the GAS genes *gacHIJKL* required for the

formation and addition of the GlcNAc side chains and GroP to GAC [21], in the Δ*sccN*

background strain in two versions, creating the Δ*sccN:pgacHI*JKL* and Δ*sccN:pgacHIJKL*

strains [50]. The plasmid *pgacHI*JKL* contains an inserted stop codon in the *gacI* gene required

for generation of the GlcNAc side chain, and, therefore, the Δ*sccN:pgacHI*JKL* strain produces

the unmodified SCC lacking any side chains (Fig. 5). As expected, the sacculi isolated from this

strain remains susceptible to PlyC lysis. We previously showed that in Δ*sccN:pgacHIJKL*, the

Glc side chains are replaced with the GlcNAc side chains [50]. Interestingly, expression of

gacHIJKL in Δ*sccN* did not restore the resistance of the bacteria to PlyC (Fig. 5), indicating that

the GlcNAc side chains do not obstruct PlyC binding. Lastly, we analyzed PlyC-mediated lysis

of sacculi purified from the Δ*rgpG* mutant, which is deficient in SCC expression [50]. The RgpG

protein catalyzes the first step in SCC biosynthesis [51]. In comparison to Δ*sccN*, PlyC-induced

lysis of Δ*rgpG* was less pronounced (Fig. 5), indicating the importance of the pRha backbone of

SCC in PlyC activity and supporting the findings that the pRha backbone is a ligand contributing

to PlyC binding. These studies reveal that if the SCC is 'unmasked' *i.e.,* stripped of the Glc and

Glc-GroP side chains, it becomes a ligand for PlyCB and that *S. mutans* is PlyC susceptible if

SCC is replaced with GAC.

Molecular Dynamics simulations identify novel binding pocket in PlyC

Our biochemical investigations revealed that PlyCB interacts directly with the pRha backbone of the GAC as shown in the cellular context of the native streptococcal cells, engineered *E. coli* cells and *S. mutans* mutant strains. In contrast, traditional biochemical techniques such as size exclusion chromatography or isothermal titration calorimetry could not detect the binding of PlyCB when either the short or long pRha fragments were used (data not shown). These results corroborate the previous studies on related proteins, showing that the binding between endolysins to their respective isolated or purified cell wall ligands is challenging to characterize [52, 53]. However, the CBD is a carbohydrate binding module, which often contains the prominent binding sites to selectively coordinate the binding of linear polysaccharides (reviewed in [54]). To identify the prominent binding sites for pRha, we investigated the previously reported crystal structures of $PlyCB_{WT}$ and $PlyCB_{R66E}$ with the ligand site identification software SiteMap [55]. The software identifies cavities on the molecular surface of a protein structure and provides a 'druggability score', which indicates the likelihood that an identified site can bind a small molecule. Interestingly, for both $PlyCB_{WT}$ and $PlyCB_{R66E}$ protein structures, no druggable sites were identified (SF2). In the $PlyCB_{WT}$ structure, a small shallow cavity in proximity of R66 was identified, but with a very poor druggability score (0.4). Shen *et al.*, had previously reported a putative phosphate binding site in PlyCB [56], however, we concluded from our studies that no pocket or cavity is large enough to accommodate the pRha structure. Protein crystal structures represent a fixed conformation state that does not have to reflect the native fold in solution or in the context of the respective ligands. Riley *et al.* reported previously a PlyCB SAXS in solution structure [57] where a high degree of flexibility and movement of the PlyCB octameric ring was revealed. We therefore conducted Molecular Dynamics (MD) simulations on the $PlyCB_{WT}$ structure (PDB 4F87) [13] to investigate how the protein structure changes over time. The structure was simulated as a monomeric protein for 500 ns in the explicit solvent mode. Strikingly, the $PlyCB_{WT}$ structure changed into an 'open' conformation, where Y28 rotates and

remains stable in a new conformation, generating a novel pocket (SF3). In the WT crystal

structure, R29 establishes a salt bridge with E36 in the same β-hairpin and has a repulsive

interaction with R66 on the α-helix. This repulsive effect results in a higher mobility of the

terminal part of the β-strand which allows the stabilisation of an open conformation of Y28

(SF3). The stabilisation of the open conformation of Y28 opens a cavity that is clearly visible on

the molecular surface of the WT structure and is identified by SiteMap as a druggable binding

site (score 0.8) (SF2).

It is well reported in the literature that a PlyC holoenzyme formed by PlyCA and the PlyCB$_{R66E}$

mutant protein is not able to lyse GAS [56]. Our biochemical studies have revealed that the

PlyCB$_{R66E}$ mutant does not bind pRha. We therefore conducted MD simulations also on the

reported PlyCB$_{R66E}$ crystal structure. In agreement with the PlyCB$_{WT}$ protein simulations, the

structure was stable. Contrary to the PlyCB$_{WT}$ protein, the rotation of Y28 was not observed and

subsequently, no cavity/pocket is formed (SF4). We inspected the conformation of the mutated

residues in PlyCB$_{R66E}$. The side chain of R66E forms a stable salt bridge with R29, increasing

the structural rigidity and preventing the opening of Y28 (SF4). Interestingly, the site controlled

by Y28 is not at the protein-protein interface, but within the monomeric structure, suggesting

that a ligand binding pocket exists within the monomeric protein and is not at the interface of two

neighboring monomers (SF2 1B, Fig. 6).

PlyCB-rhamnose di- and tri-saccharide docking reveals potential interactions

The MD simulations revealed a potential ligand binding site in PlyCB$_{WT}$ that is generated after

the gatekeeper Y28 rotated towards the α-helix. We conducted ligand-protein docking using a

Rha di- and tri-saccharide, with both possible linkage configurations. The Rha disaccharide and

trisaccharide with a non-reducing end α1,2-Rha (SF5) produced the congruent docking results,

which we further analyzed (Fig. 6, SF6). The Rha disaccharide with the non-reducing end (1->3)

linkage appears to be stabilized by four hydrogen bond interactions (SF7 A, B). The terminal

reducing end Rha is within the hydrogen bond distance to S17 and T12, whilst the α1,3-linked

non-reducing end Rha is able to interact with the carbonyl backbone of F10. The identical

interactions are observed for the docked trisaccharide (SF7 C, D). In addition, the non-reducing

end Rha of the trisaccharide also interacts with the side chain hydroxyl groups of S54 and Q49

(SF7 C, D). When we carefully inspected the structural context of the Rha trisaccharide with the

terminal α1,2-linked Rha moieties, we detected the adjacent pockets that could potentially bind

the GlcNAc side chain linked to these Rha moieties (SF8 A, B). It is beyond the scope of this

study to confirm that these pockets have the ability to bind the GAC β-GlcNAc sidechain or in

fact the Group C Carbohydrate (GCC) β-GalNAc-GalNAc disaccharide. However, this model is

consistent with PlyCB binding the unmodified pRha backbone and the backbone decorated with

the GAC or GCC side chains. The model also explains why the *S. mutans* SCC — with the α-2-

Glc substitution on the α1,3-linked Rha — is not a ligand for the PlyCB protein. The identified

pocket does not allow the binding of the SCC with the side chain in this configuration (SF8 A,

C).

Polyrhamnose binding pocket suggests the PlyC catalytic mechanism

We further investigated the structural context of this binding pocket, which is perpendicular to

the PlyCB monomer-monomer interface. We superposed the reported PlyCA structure that also

contains the GlyH and CHAP amidase domains (Fig 7). This led to the intriguing discovery that

this model would not only explain the specificity of the enzyme, but also the substrate

recognition by the CHAP amidase and GlyH domains. The proposed pRha binding pocket is

perpendicular to the GlyH and CHAP amidase domains, that bind the PG backbone and

crosslinked peptides. Similar to wall teichoic acids[58], the GAC and GCC are likely directly

linked via their reducing end sugars to *N*-acetylmuramic acid (MurNAc) of PG. We observed that

the reducing end sugar of the docked complex points towards the GlyH and CHAP amidase

domains. When we compare this structural alignment with the architectural arrangement of the

crosslinked PG decorated with the GAC, the model greatly assists our understanding of

PlyCAactivity: the CHAP amidase domain cleaves the PG peptide bond. This allows the PG to

open up and slide into the GlyH domain, which removes the PG GlcNAc-MurNAc from the

reducing end. Concurrently, the GAC attached to the PG binds to the octameric PlyCB domains.

We docked a PG tetrasacharide into the reported PlyCA GlyH domain (Fig. 7). This positions

the PG in 5-7Å, the distance that the terminal GAC sugar, GlcNAc, could occupy [23].

Concluding remarks

Strikingly, our PlyC molecular dynamics simulations revealed a previously unknown

conformational change that exposes a potential carbohydrate binding site. Our simulations

agree with the reported flexibility of the PlyCB structure in solution, as determined by SAXS

studies [57]. The reported crystal structure represents a locked state that appears to be the

most stable configuration in the crystallization conditions, in absence of any ligand. The side

chain of Y28 acts as gatekeeper and conformational changes allow opening of a cavity that is

likely capable of recognizing the compounds in the size of several sugar residues. These

structural changes are possible due to an intrinsic flexibility introduced by the repulsion between

the side chains of R29 (adjacent to Y28) and R66 (SF3). Moreover, we propose a structural

feature of the PlyCB protein that explains why only certain streptococci are susceptible to PlyC's

lytic activity. The pRha decorated with an $\alpha-$linked side chain sugar is not compatible with the

PlyCB ligand binding site and therefore only those streptococci expressing pRha decorated with

β-linked substituents, such as GlcNAc and GlcNAc-GroP are susceptible. This could potentially

be exploited for diagnostic purposes or in the case of SCC and Group G Streptococci, opens up

the potential for novel therapeutic approaches. If SCC was treated by PlyC in combination with

an additional enzyme that removes the α–linked side chains in these streptococcal

carbohydrates, this would expose the pRha backbone and subsequently make these strains

susceptible. The identification of the pRha ligand site could also be further exploited by directed

evolution approaches to generate PlyCB protein variants that are capable to bind the

carbohydrates from, for example, SCC and Group G Streptococci. Since our proposed ligand

binding site is on the surface of each PlyCB monomer mutagenesis would most likely not impact

the multimerization interface of the PlyCB octameric ring.

Whilst much has been learned about the structure and function of PlyC in the past 20 years,

many questions remain, specifically with respect to its interaction with the PG. Considering the

average length of the cellular pRha is 7-10 kDa [25] and that our MD simulations shows an

individual PlyCB monomer can bind its target ligand, the α1,2-1,3-pRha with or without β-

configured GlcNAc/GalNAc-side chains, it is inviting to speculate that an element of avidity is

responsible for tight binding of the PlyCB octamer to the streptococcal surface. We have

envisioned a model in Fig. 7, although further proof is needed to substantiate this hypothesis.

Another question lies in the actions of the EADs relative to the PlyCB octamer. PlyC clearly has

a high turnover as demonstrated in multiple biochemical assays. However, it is unknown if

PlyCB "docks" to the surface and the flexibility of the holoenzyme allows the PlyC EADs to

cleave multiple bonds in a localized area weakening the overall superstructure of the PG.

Alternatively, the enzymatic turnover could be dictated by a balance of on and off rates of the

EADs and CBD monomers leading to widespread hydrolysis of the PG. Lastly, it is unknown

whether PlyC binds, cleaves, and releases the PG at random points on the streptococcal

surface or works its way down a single strand of PG in a processive manner. It is noteworthy

that cellulase enzymes, which cleave the β1,4 glycosidic linkages in cellulose, possess a

catalytic domain, a flexible linker, and a cellulose binding domain, analogous to the traditional

endolysins. It has been demonstrated that energy is stored in the flexible linker can adopt

compact and extended configurations that allows the cellulase to move in a "caterpillar-like"

motion down a chain of cellulose [59, 60]. Although PlyC does not contain an equivalent flexible

linker, the octameric nature of PlyCB invites the possibility that it may interact with the

successive pRha strands allowing PlyC to depolymerize the PG in a zipper-like fashion.

In conclusion, the α1,2-1,3-pRha is the definite, minimal carbohydrate substrate for the PlyCB

subunit. We validated this by comprehensive experiments, using pRha recombinantly produced

in *E. coli*, and the *S. mutans* variants of the Rha-based polysaccharides. Furthermore, the MD

simulations and subsequent docking studies revealed a potential binding site for Rha tri-

saccharides. Further studies are needed to detail the dynamics of the PlyC holoenzyme on the

streptococcal surface and the interdomain interactions. Nonetheless, the work described here

provides valuable insight into the molecular interactions that define a PlyC's host specificity,

which can inform the future studies as well as engineering approaches.

Data Availability Statement

No mandated datasets are associated with the paper.

 Figure Legend

Fig 1)

A) Symbolic drawings of the carbohydrate structures of GAS (GAC), polyrhamnose backbone

(pRha) in GAVS and M1_d*gacI*, GCS (GCC), GGS (GGC) and *S. mutans* (SCC). For simplicity,

the reported glycerol phosphate of occasionally present on the GAC and SCC side chains have

been omitted. Repeat units are marked with brackets. The pRha backbone with alternating ($\alpha1$-

>2) and ($\alpha1$->3) linkages, whilst the $\alpha1$->2 Rha being decorated with a $\beta1$->3 side chain is a

commonality among all PlyC susceptible strains. B) PlyC lysis of streptococcal pathogens.

Group A and Group C Streptococci serotypes are susceptible to PlyC-mediated lysis. Group G

Streptococci show limited susceptibility and *S. mutans* is resistant to PlyC lysis. Three SDSE

isolates that produce the GAC instead of GGC (SDSE_*gac*) are susceptible to PlyC treatment.

Fig 2)

Precipitation studies of purified PlyCB and GAC reveal direct interaction of PlyCB with GAC. A)

The PlyCB concentration is kept constant whilst the GAC concentration is varied. Visible

precipitate forms at the higher concentrations. B) The precipitate level is measured

spectrophotometrically at 340 nm and compared to the mutant $PlyCB_{R66E}$, which does not bind

the GAC. C) Coomassie stained and D) densitometry analysis of PlyCB protein from the

supernatant fraction and aggregates (pellets). E, F) The same dose dependency is observed

when the PlyCB concentration is varied. Arrowhead depicts PlyCB protein at 8 kDa.

Fig 3)

Representation of immunoblot analysis of the cell lysate of *E. coli* expressing the SCC and GAC

pRha and carrying an empty control plasmid (-ve). A) Blot was incubated with PlyCB-Alexa

Fluor® 647 ($PlyCB_{WT}^{AF647}$). B) Probing the same samples with the GAC antibodies confirms the

presence of GAC in the bands. Molecular mass markers are given in kDa.

Fig 4)

PlyCB binding to *E. coli* cells were investigated by flow cytometry after labelling with

PlyCB$_{WT}$^{AF647} and PlyCB$_{R66E}$^{AF645} mutant proteins. Blue: -ve control cells without pRha. Red:

pRha producing *E. coli* cells. Representative histograms are shown. A) Left panel: unstained

cells. Right panel: The anti-GAC antibodies (GAC-FITC) were used as a positive control to label

the *E. coli* cells producing pRha. The antibodies do not bind to the *E. coli* cells carrying an

empty plasmid (-ve). B) Left panel: PlyCB$_{WT}$^{AF647} binds to the *E. coli* cells producing pRha, but

not to the *E. coli* cells carrying an empty plasmid (-ve). Right panel: PlyCB$_{R66E}$^{AF645} does not

binds to the *E. coli* cells producing pRha.

Fig 5)

The PlyC-mediated lysis of sacculi purified from *S. mutans* strains. The lysis was monitored

after 10, 20, 30, 40, 50 and 60 min as a decrease in OD$_{600}$. Results are presented as a fold

change in OD$_{600}$ of the sacculi incubated with PlyC *vs.* the sacculi incubated without PlyC. Data

points and error bars represent mean values of four biological replicates and standard deviation,

respectively. P-values were determined by two-way ANOVA with Dunnett's multiple

comparisons test.

Fig 6)

Bottom view of the PlyCB protein structures in surface representation. The octameric PlyCB

structure with brown A, C, E, G and orange B, D, F, H monomers. A) WT (GH and CHAP

domain omitted, 4f87.pdb) B) WT after MD with docked tri-Rha. The MD simulations have

revealed a novel pocket that could accommodate the binding of short Rha-polysaccharides.

Fig 7)

A schematic model of the proposed PlyC binding to the streptococcal cell wall. A) PlyC in the

complex with docked PG tetra-saccharide in the GlyH domain (red) and pRha tri-saccharides in

PlyCB subunits. B) A schematic model showing PlyC binding to peptidoglycan and pRha.

Sugars are drawn to scale, with green triangles (Rha), blue square (GlcNAc) and beige square

(MurNAc). Proposed PlyC activity: The EAD cleaves the PG peptide to open up the PG

backbones that subsequently enter the GlyH domain (red) for PG cleavage.

FUNDING

The HCD laboratory is supported by Wellcome and Royal Society Grant 109357/Z/15/Z and the

University of Dundee Wellcome Trust Funds 105606/Z/14/Z and Tenovus Scotland Large

Research Grant [T17/17]. HK is supported by the National Institute of Standards and

Technology (NIST). VAF is supported by Rockefeller University laboratory funds. NK is

supported by NIH grants R01 DE028916 from the NIDCR and R01 AI143690 from the NIAID.

ACKNOWLEDGEMENTS

We thank the CDC Streptococcus Lab and the Active Bacterial Core surveillance program for

sharing isolates isolate 20170556 (stG6.0), 20171682 (stC74A.0), 20154376 (stG245.0),

20173686 (stG652.0), 20170560 and 20176966 (stG485.0). We also thank Ryan Heselpoth,

Yang Shen, and Sara Linden for reagents, technical advice, and helpful discussion.

References

1   Fernandes, S. and Sao-Jose, C. (2018) Enzymes and Mechanisms Employed by Tailed
Bacteriophages to Breach the Bacterial Cell Barriers. Viruses. **10**
2   Nelson, D., Loomis, L. and Fischetti, V. A. (2001) Prevention and elimination of upper
respiratory colonization of mice by group A streptococci by using a bacteriophage lytic enzyme.
Proc Natl Acad Sci U S A. **98**, 4107-4112

3   Dams, D. and Briers, Y. (2019) Enzybiotics: Enzyme-Based Antibacterials as
Therapeutics. Adv Exp Med Biol. **1148**, 233-253
4   Love, M. J., Bhandari, D., Dobson, R. C. J. and Billington, C. (2018) Potential for
Bacteriophage Endolysins to Supplement or Replace Antibiotics in Food Production and Clinical
Care. Antibiotics (Basel). **7**
5   Schmelcher, M. and Loessner, M. J. (2020) Bacteriophage endolysins - extending their
application to tissues and the bloodstream. Curr Opin Biotechnol. **68**, 51-59
6   Heselpoth, R. D., Swift, S. M., Linden, S. B., Mitchell, M. S. and Nelson, D. C. (2018)
Enzybiotics: endolysins and bacteriocins. In Bacteriophages - biology, technology, and therapy
(Harper, D., Abedon, S. T., Burrowes, B. and McConvill, M., eds.), Springer International
Publishing, Dordrecht, The Netherlands
7   Loessner, M. J., Kramer, K., Ebel, F. and Scherer, S. (2002) C-terminal domains of
Listeria monocytogenes bacteriophage murein hydrolases determine specific recognition and
high-affinity binding to bacterial cell wall carbohydrates. Molecular microbiology. **44**, 335-349
8   Mesnage, S., Dellarole, M., Baxter, N. J., Rouget, J. B., Dimitrov, J. D., Wang, N.,
Fujimoto, Y., Hounslow, A. M., Lacroix-Desmazes, S., Fukase, K., Foster, S. J. and Williamson,
M. P. (2014) Molecular basis for bacterial peptidoglycan recognition by LysM domains. Nat
Commun. **5**, 4269
9   Evans, A. C. (1934) *Streptococcus* bacteriphage: a study of four serological types.
Public Health Rep. **49**, 1386-1401
10   Krause, R. M. (1957) Studies on bacteriophages of hemolytic streptococci. I. Factors
influencing the interaction of phage and susceptible host cell. J Exp Med. **106**, 365-384
11   Maxted, W. R. (1957) The active agent in nascent phage lysis of streptococci. J Gen
Microbiol. **16**, 584-595
12   Nelson, D., Schuch, R., Chahales, P., Zhu, S. and Fischetti, V. A. (2006) PlyC: a
multimeric bacteriophage lysin. Proc Natl Acad Sci U S A. **103**, 10765-10770
13   McGowan, S., Buckle, A. M., Mitchell, M. S., Hoopes, J. T., Gallagher, D. T., Heselpoth,
R. D., Shen, Y., Reboul, C. F., Law, R. H., Fischetti, V. A., Whisstock, J. C. and Nelson, D. C.
(2012) X-ray crystal structure of the streptococcal specific phage lysin PlyC. Proc Natl Acad Sci
U S A. **109**, 12752-12757
14   Lancefield, R. C. (1933) A Serological Differentiation of Human and Other Groups of
Hemolytic Streptococci. J Exp Med. **57**, 571-595
15   Vandamme, P., Pot, B., Falsen, E., Kersters, K. and Devriese, L. A. (1996) Taxonomic
study of lancefield streptococcal groups C, G, and L (Streptococcus dysgalactiae) and proposal
of S. dysgalactiae subsp. equisimilis subsp. nov. Int J Syst Bacteriol. **46**, 774-781
16   Coligan, J. E., Kindt, T. J. and Krause, R. M. (1978) Structure of the streptococcal
groups A, A-variant and C carbohydrates. Immunochemistry. **15**, 755-760
17   Pritchard D G, G. R. L., Michlek S M, McGhee J R. . (1986) Biochemical aspects of
serotype carbohydrate antigens of Streptococcus mutans. In: Hamada S, Michalek S M, Kiyono
H, Menaker L, McGhee J R, editors. Molecular microbiology and immunology of Streptococcus
mutans. Amsterdam, The Netherlands: . Elsevier Science Publishers; , pp. 39–49.
18   Coligan, J. E., Schnute, W. C., Jr. and Kindt, T. J. (1975) Immunochemical and chemical
studies on streptococcal group-specific carbohydrates. J Immunol. **114**, 1654-1658
19   Huang, D. H., Rama Krishna, N. and Pritchard, D. G. (1986) Characterization of the
group A streptococcal polysaccharide by two-dimensional [1]H-nuclear-magnetic-resonance
spectroscopy. Carbohydrate research. **155**, 193-199
20   Neiwert, O., Holst, O. and Duda, K. A. (2014) Structural investigation of rhamnose-rich
polysaccharides from Streptococcus dysgalactiae bovine mastitis isolate. Carbohydr Res. **389**,
192-195
21   Edgar, R. J., van Hensbergen, V. P., Ruda, A., Turner, A. G., Deng, P., Le Breton, Y.,
El-Sayed, N. M., Belew, A. T., McIver, K. S., McEwan, A. G., Morris, A. J., Lambeau, G.,

Walker, M. J., Rush, J. S., Korotkov, K. V., Widmalm, G., van Sorge, N. M. and Korotkova, N. (2019) Discovery of glycerol phosphate modification on streptococcal rhamnose polysaccharides. Nature chemical biology. **15**, 463-471

van der Beek, S. L., Zorzoli, A., Canak, E., Chapman, R. N., Lucas, K., Meyer, B. H., Evangelopoulos, D., de Carvalho, L. P. S., Boons, G. J., Dorfmueller, H. C. and van Sorge, N. M. (2019) Streptococcal dTDP-L-rhamnose biosynthesis enzymes: functional characterization and lead compound identification. Molecular microbiology. **111**, 951-964

Zorzoli, A., Meyer, B. H., Adair, E., Torgov, V. I., Veselovsky, V. V., Danilov, L. L., Uhrin, D. and Dorfmueller, H. C. (2019) Group A, B, C, and G Streptococcus Lancefield antigen biosynthesis is initiated by a conserved alpha-d-GlcNAc-beta-1,4-l-rhamnosyltransferase. J Biol Chem. **294**, 15237-15256

van der Beek, S. L., Le Breton, Y., Ferenbach, A. T., Chapman, R. N., van Aalten, D. M., Navratilova, I., Boons, G. J., McIver, K. S., van Sorge, N. M. and Dorfmueller, H. C. (2015) GacA is essential for Group A Streptococcus and defines a new class of monomeric dTDP-4-dehydrorhamnose reductases (RmlD). Molecular microbiology. **98**, 946-962

Rush, J. S., Edgar, R. J., Deng, P., Chen, J., Zhu, H., van Sorge, N. M., Morris, A. J., Korotkov, K. V. and Korotkova, N. (2017) The molecular mechanism of N-acetylglucosamine side-chain attachment to the Lancefield group A carbohydrate in Streptococcus pyogenes. J Biol Chem. **292**, 19441-19457

Lood, R., Raz, A., Molina, H., Euler, C. W. and Fischetti, V. A. (2014) A highly active and negatively charged Streptococcus pyogenes lysin with a rare D-alanyl-L-alanine endopeptidase activity protects mice against streptococcal bacteremia. Antimicrob Agents Chemother. **58**, 3073-3084

Gera, K. and McIver, K. S. (2013) Laboratory growth and maintenance of Streptococcus pyogenes (the Group A Streptococcus, GAS). Curr Protoc Microbiol. **30**, 9D 2 1-9D 2 13

Shibata, Y., Yamashita, Y., Ozaki, K., Nakano, Y. and Koga, T. (2002) Expression and characterization of streptococcal rgp genes required for rhamnan synthesis in Escherichia coli. Infect Immun. **70**, 2891-2898

DuBois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A. and Smith, F. (1956) Colorimetric method for determination of sugars and related substances. Anal. Chem. **28**, 350-356

Heselpoth, R. D., Yin, Y., Moult, J. and Nelson, D. C. (2015) Increasing the stability of the bacteriophage endolysin PlyC using rationale-based FoldX computational modeling. Protein Eng Des Sel. **28**, 85-92

Case DA, e. a. (2016) AMBER2016. University of California, San Francisco

Maier, J. A., Martinez, C., Kasavajhala, K., Wickstrom, L., Hauser, K. E. and Simmerling, C. (2015) ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. J Chem Theory Comput. **11**, 3696-3713

Darden, T. A. and Pedersen, L. G. (1993) Molecular modeling: an experimental tool. Environ Health Perspect. **101**, 410-412

Jean-Paul Ryckaert, G. C., Herman J.C Berendsen. (1977) Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. Journal of Computational Physics. **23**, 327-341

DeLano, W. L. (2002) The PyMOL Molecular Graphics System. Delano Scientific, San Carlos.

William Humphrey, A. D., Klaus Schulten. (1996) VMD: Visual molecular dynamics. Journal of Molecular Graphics. **14**, 33-38

2021-1:, S. R. (2021) LigPrep Schrödinger, LLC, New York, NY

M., M. (1952) The lysis of group A hemolytic streptococci by extracellular enzymes of Streptomyces albus. II. Nature of the cellular substrate attacked by the lytic enzymes. . J Exp Med. . **96**, 569–580

39   Davies, M. R., McIntyre, L., Mutreja, A., Lacey, J. A., Lees, J. A., Towers, R. J.,
Duchene, S., Smeesters, P. R., Frost, H. R., Price, D. J., Holden, M. T. G., David, S., Giffard, P.
M., Worthing, K. A., Seale, A. C., Berkley, J. A., Harris, S. R., Rivera-Hernandez, T., Berking,
O., Cork, A. J., Torres, R., Lithgow, T., Strugnell, R. A., Bergmann, R., Nitsche-Schmitz, P.,
Chhatwal, G. S., Bentley, S. D., Fraser, J. D., Moreland, N. J., Carapetis, J. R., Steer, A. C.,
Parkhill, J., Saul, A., Williamson, D. A., Currie, B. J., Tong, S. Y. C., Dougan, G. and Walker, M.
J. (2019) Author Correction: Atlas of group A streptococcal vaccine candidates compiled using
large-scale comparative genomics. Nat Genet. **51**, 1295
40   Brandt, C. M., Haase, G., Schnitzler, N., Zbinden, R. and Lutticken, R. (1999)
Characterization of blood culture isolates of Streptococcus dysgalactiae subsp. equisimilis
possessing Lancefield's group A antigen. J Clin Microbiol. **37**, 4194-4197
41   Chochua, S., Metcalf, B. J., Li, Z., Rivers, J., Mathis, S., Jackson, D., Gertz, R. E., Jr.,
Srinivasan, V., Lynfield, R., Van Beneden, C., McGee, L. and Beall, B. (2017) Population and
Whole Genome Sequence Based Characterization of Invasive Group A Streptococci Recovered
in the United States during 2015. mBio. **8**
42   McMillan, D. J., Vu, T., Bramhachari, P. V., Kaul, S. Y., Bouvet, A., Shaila, M. S.,
Karmarkar, M. G. and Sriprakash, K. S. (2010) Molecular markers for discriminating
Streptococcus pyogenes and S. dysgalactiae subspecies equisimilis. Eur J Clin Microbiol Infect
Dis. **29**, 585-589
43   Lynskey, N. N., Jauneikaite, E., Li, H. K., Zhi, X., Turner, C. E., Mosavie, M., Pearson,
M., Asai, M., Lobkowicz, L., Chow, J. Y., Parkhill, J., Lamagni, T., Chalker, V. J. and
Sriskandan, S. (2019) Emergence of dominant toxigenic M1T1 Streptococcus pyogenes clone
during increased scarlet fever activity in England: a population-based molecular epidemiological
study. Lancet Infect Dis. **19**, 1209-1218
44   Pritchard, D. G., Rener, B. P., Furner, R. L., Huang, D. H. and Krishna, N. R. (1988)
Structure of the group G streptococcal polysaccharide. Carbohydr Res. **173**, 255-262
45   Mirica, K. A., Lockett, M. R., Snyder, P. W., Shapiro, N. D., Mack, E. T., Nam, S. and
Whitesides, G. M. (2012) Selective precipitation and purification of monovalent proteins using
oligovalent ligands and ammonium sulfate. Bioconjug Chem. **23**, 293-299
46   Sato, T., Mattison, K. W., Dubin, P. L., Kamachi, M. and Morishima, Y. (1998) Effect of
protein aggregation on the binding of lysozyme to pyrene-labeled polyanions. Langmuir. **14**,
5430-5437
47   Wilcken, R., Wang, G., Boeckler, F. M. and Fersht, A. R. (2012) Kinetic mechanism of
p53 oncogenic mutant aggregation and its inhibition. Proc Natl Acad Sci U S A. **109**, 13584-
13589
48   Michon, F., Moore, S. L., Kim, J., Blake, M. S., Auzanneau, F. I., Johnston, B. D.,
Johnson, M. A. and Pinto, B. M. (2005) Doubly branched hexasaccharide epitope on the cell
wall polysaccharide of group A streptococci recognized by human and rabbit antisera. Infect
Immun. **73**, 6383-6389
49   van Sorge, N. M., Cole, J. N., Kuipers, K., Henningham, A., Aziz, R. K., Kasirer-Friede,
A., Lin, L., Berends, E. T. M., Davies, M. R., Dougan, G., Zhang, F., Dahesh, S., Shaw, L., Gin,
J., Cunningham, M., Merriman, J. A., Hutter, J., Lepenies, B., Rooijakkers, S. H. M., Malley, R.,
Walker, M. J., Shattil, S. J., Schlievert, P. M., Choudhury, B. and Nizet, V. (2014) The classical
lancefield antigen of group a Streptococcus is a virulence determinant with implications for
vaccine design. Cell Host Microbe. **15**, 729-740
50   Svetlana Zamakhaeva, C. T. C., Jeffrey S. Rush, Sowmya Ajay Castro, Alexander E.
Yarawsky, Andrew B. Herr, Nina M. van Sorge, Helge C. Dorfmueller, Gregory I. Frolenkov,
Konstantin V. Korotkov, Natalia Korotkova. (2020) Modification of cell wall polysaccharide
spatially controls cell division in Streptococcus mutans. bioRxiv
51   De, A., Liao, S., Bitoun, J. P., Roth, R., Beatty, W. L., Wu, H. and Wen, Z. T. (2017)
Deficiency of RgpG causes major defects in cell division and biofilm formation, and deficiency of

LytR-CpsA-Psr family proteins leads to accumulation of cell wall antigens in culture medium by
Streptococcus mutans. Appl Environ Microbiol. **83**
52   Ganguly, J., Low, L. Y., Kamal, N., Saile, E., Forsberg, L. S., Gutierrez-Sanchez, G.,
Hoffmaster, A. R., Liddington, R., Quinn, C. P., Carlson, R. W. and Kannenberg, E. L. (2013)
The secondary cell wall polysaccharide of Bacillus anthracis provides the specific binding ligand
for the C-terminal cell wall-binding domain of two phage endolysins, PlyL and PlyG.
Glycobiology. **23**, 820-832
53   Shen, Y., Kalograiaki, I., Prunotto, A., Dunne, M., Boulos, S., Taylor, N. M. I., Sumrall, E.
T., Eugster, M. R., Martin, R., Julian-Rodero, A., Gerber, B., Leiman, P. G., Menendez, M., Dal
Peraro, M., Canada, F. J. and Loessner, M. J. (2021) Structural basis for recognition of bacterial
cell wall teichoic acid by pseudo-symmetric SH3b-like repeats of a viral peptidoglycan
hydrolase. Chemical Science. **12**, 576-589
54   Shoseyov, O., Shani, Z. and Levy, I. (2006) Carbohydrate binding modules: biochemical
properties and novel applications. Microbiol Mol Biol Rev. **70**, 283-295
55   Halgren, T. (2007) New method for fast and accurate binding-site identification and
analysis. Chem Biol Drug Des. **69**, 146-148
56   Shen, Y., Barros, M., Vennemann, T., Gallagher, D. T., Yin, Y., Linden, S. B., Heselpoth,
R. D., Spencer, D. J., Donovan, D. M., Moult, J., Fischetti, V. A., Heinrich, F., Losche, M. and
Nelson, D. C. (2016) A bacteriophage endolysin that eliminates intracellular streptococci. eLife.
5
57   Riley, B. T., Broendum, S. S., Reboul, C. F., Cowieson, N. P., Costa, M. G., Kass, I.,
Jackson, C., Perahia, D., Buckle, A. M. and McGowan, S. (2015) Dynamic Motion and
Communication in the Streptococcal C1 Phage Lysin, PlyC. PLoS One. **10**, e0140219
58   Brown, S., Santa Maria, J. P., Jr. and Walker, S. (2013) Wall teichoic acids of gram-
positive bacteria. Annu Rev Microbiol. **67**, 313-336
59   Rabinovich, M. L., Melnick, M. S. and Bolobova, A. V. (2002) The structure and
mechanism of action of cellulolytic enzymes. Biochemistry (Mosc). **67**, 850-871
60   McCabe, C., Zhao, X., Adney, W. S., Himmel, M. E. (2010) Energy Storage in Cellulase
Linker Peptides? Computational Modeling in Lignocellulosic Biofuel Production. **1052**, 119-134
834

Fig 1)

A) Symbolic drawings of the carbohydrate structures of GAS (GAC), polyrhamnose backbone (pRha) in GAVS and M1_d*gacI*, GCS (GCC), GGS (GGC) and S. mutans (SCC). For simplicity, the reported glycerol phosphate of occasionally present on the GAC and SCC side chains have been omitted. Repeat units are marked with brackets. The pRha backbone with alternating (α1->2) and (α1->3) linkages, whilst the α1->2 rhamnose being decorated with a β1->3 side chain is a commonality among all PlyC susceptible strains. B) PlyC lysis of streptococcal pathogens. Group A and Group C Streptococci serotypes are susceptible to PlyC-mediated lysis. Group G Streptococci show limited susceptibility and S. mutans is resistant to PlyC lysis. Three SDSE isolates that produce the GAC instead of GGC (SDSE_*gac*) are susceptible to PlyC treatment.

Fig 2)

Precipitation studies of purified PlyCB and GAC reveal direct interaction of PlyCB with GAC. A) The PlyCB concentration is kept constant whilst the GAC concentration is varied. Visible precipitate forms at the higher concentrations. B) The precipitate level is measured spectrophotometrically at 340 nm and compared to the mutant PlyCB_R66E, which does not bind the GAC. C) Coomassie stained and D) densitometry analysis of PlyCB protein from the supernatant fraction and aggregates (pellets). E, F) The same dose dependency is observed when the PlyCB concentration is varied. Arrowhead depicts PlyCB protein (8 kDa).

Fig 3)
Representation of immunoblot analysis of the cell lysate of E. coli expressing the SCC and GAC pRha and carrying an empty control plasmid (-ve). A) Blot was incubated with PlyCB-Alexa Fluor® 647 (PlyCB-WT-AF647). B) Probing the same samples with the GAC antibodies confirms the presence of GAC in the bands. Molecular mass markers are given in kDa.

Fig 4)

PlyCB binding to *E. coli* cells were investigated by flow cytometry after labelling with PlyCB$_{WT}^{AF647}$ and PlyCB$_{R66E}^{AF645}$ mutant proteins. Blue: -ve control cells without pRha. Red: pRha producing *E. coli* cells. Representative histograms are shown. A) Left panel: unstained cells. Right panel: The anti-GAC antibodies (GAC-FITC) were used as a positive control to label the *E. coli* cells producing pRha. The antibodies do not bind to the E. coli cells carrying an empty plasmid (-ve). B) Left panel: PlyCB$_{WT}^{AF647}$ binds to the *E. coli* cells producing pRha, but not to the *E. coli* cells carrying an empty plasmid (-ve). Right panel: PlyCB$_{R66E}^{AF645}$ does not binds to the *E. coli* cells producing pRha.

Fig 5)

The PlyC-mediated lysis of sacculi purified from *S. mutans* strains. The lysis was monitored after 10, 20, 30, 40, 50 and 60 min as a decrease in OD_{600}. Results are presented as a fold change in OD_{600} of the sacculi incubated with PlyC *vs.* the sacculi incubated without PlyC. Data points and error bars represent mean values of four biological replicates and standard deviation, respectively. P-values were determined by two-way ANOVA with Dunnett's multiple comparisons test.

Fig 6)

Bottom view of the PlyCB protein structures in surface representation. The octameric PlyCB structure with brown A, C, E, G and orange B, D, F, H monomers. A) WT (GH and CHAP domain omitted, 4f87.pdb) B) WT after MD with docked tri-rhamnose. The MD simulations have revealed a novel pocket that could accommodate the binding of short rhamnose-polysaccharides.

Fig 7)
A schematic model of the proposed PlyC binding to the streptococcal cell wall. A) PlyC in the complex with docked PG tetra-saccharide in the GH domain (red) and polyrhamnose tri-saccharides in PlyCB subunits. B) A schematic model showing PlyC binding to peptidoglycan and polyrhamnose. Sugars are drawn to scale, with green triangles (rhamnose), blue square (GlcNAc) and beige square (MurNAc). Proposed PlyC activity: The EAD cleaves the PG peptide to open up the PG backbones that subsequently enter the GH domain (red) for PG cleavage.