Multi-substrate DNA stable isotope probing reveals guild structure in bacterially mediated soil carbon cycling

Samuel E. Barnett[a*], Nicholas D. Youngblut[a,b*], Chantal N. Koechli[a,c], and Daniel H. Buckley[a]

    a. School of Integrative Plant Science, Cornell University, Ithaca, NY, USA

    b. Department of Microbiome Science, Max Planck Institute for Developmental Biology, Tübingen, Germany

    c. Department of Biological Sciences, University of the Sciences, Philadelphia, NY, USA

    * Co-first authors

Corresponding author: Daniel H. Buckley

dbuckley@cornell.edu

ORCIDs: 0000-0002-0096-0605, 0000-0002-7424-5276, NA, 0000-0003-1654-2907

Keywords: DNA-SIP, bacterial, soil, carbon cycling, ecology

Author contributions: NDY and DHB conceived of the study, NDY and CNK conducted the experiment and collected all the data, NDY and DHB analyzed the data, NDY, SEB, and DHB wrote the manuscript. All authors were involved in editing and reviewing manuscript.

Competing interests: The authors declare no competing interests.

Abstract

Soil microorganisms determine the fate of soil organic matter (SOM), and their activities

comprise a major component of the global carbon (C) cycle. We employed a multi-substrate

DNA-stable isotope probing experiment to track bacterial assimilation of C derived from distinct

sources that varied in bioavailability. This approach allowed us to measure microbial

contributions to SOM processing by measuring the C assimilation dynamics of diverse

microorganisms as they interact within soil. We identified and tracked 1,286 bacterial taxa that

assimilated ^{13}C in an agricultural soil over a period of 48 days. Overall ^{13}C-assimilation

dynamics of bacterial taxa, defined by the source and timing of the ^{13}C they assimilated,

exhibited low phylogenetic conservation. We identified bacterial guilds comprised of taxa that

had similar ^{13}C assimilation dynamics. We show that C source bioavailability explained

significant variation in both C mineralization dynamics and guild structure. In addition, guild

structure explained significant variation in bacterial growth dynamics. We demonstrate that the

observed guild structure is consistent with predictions made by bacterial life history theory. We

also demonstrate that the guild structure explains significant variation in the biogeographical

distribution of bacteria at continental and global scales. We interpret these findings in the context

of bacterial life history strategies and their relationship to terrestrial C-cycling.

Introduction

Soil organic matter (SOM) represents the largest terrestrial pool of organic carbon (C) on Earth (1), and SOM dynamics have major impacts on the global C-cycle. SOM is sensitive to land management (2–4) and climate change (5), and SOM pools influence soil health (6, 7), agricultural productivity (8), and climate stability (6, 9). However, the mechanisms that promote SOM persistence and loss remain uncertain and this introduces variability into global C-cycle models (10–13). Microorganisms are dominant drivers of SOM decomposition, stabilization, and mineralization (14), and C-cycle models increasingly seek to incorporate microbial traits to improve predictions of SOM dynamics (15, 16). For example, microbially explicit C-cycle models, such as MEND (17), MIMICS (18), and CORPSE (19), improve prediction of C-cycle dynamics when compared to models that ignore microbial processing (20).

Soil microbes play a critical role in governing global C flux, but the microbial mechanisms that determine SOM dynamics remain poorly characterized (13). Microbial contributions to soil C-cycling are typically determined in the aggregate (*e.g.,* soil respiration) without providing mechanistic insight into processes of individual microbes. Determining the characteristics of individual microbes typically requires laboratory cultivation, but most soil microbes remain uncultivated and poorly characterized (21–23). In the absence of cultivation, the characteristics of uncultivated microbes are often inferred by assuming phylogenetic conservation with cultivated representatives (24–26). However, phylogenetic conservation varies dramatically among microbial traits and functionalities (27), and many soil microbes lack closely related cultivated isolates (28). Microbial characteristics are also often inferred through metagenomic analyses that indicate the frequency of specific gene families and metabolic pathways in the environment (29, 30).

Inferences based on phylogenetic conservatism and metagenomic analyses, however,

predict 'potential metabolism', but this information cannot currently predict ecological

interactions that govern activity in the environment. Microbe-microbe and microbe-environment

interactions alter patterns of gene expression, activity, and growth, and so potential metabolism

tells us little of the actual microbial processes that drive soil C-cycling at any particular place and

time. The 'realized activity' of a microbe *in situ* depends on a complex pattern of biotic and

abiotic interactions. Therefore, realized activity in the C-cycle might correlate only weakly with

the metabolic potential of the community as inferred from phylogenetic conservation and

metagenomic analyses (31, 32).

DNA-stable isotope probing (DNA-SIP) can assess the *in situ* activity of microorganisms

within complex habitats based on patterns of C assimilation by identifying taxa that assimilate

^{13}C from labeled substrates (33). DNA-SIP can track the assimilation of C from multiple

substrates into thousands of microbial taxa in soil (34–36) with high specificity and sensitivity

(36). We used multiple-window high-resolution DNA-SIP (MW-HR-SIP) to track ^{13}C from nine

distinct carbon sources through the soil bacterial food web over a period of 48 days (Fig.1).

These nine sources of C were selected because they varied in bioavailability and are common in

soils. We initially defined bioavailability on the basis of solubility and hydrophobicity, which

predicts sorption to particulate organic matter (37) and ability to transport across the cell

membrane. The C sources we used were cellulose (a component of plant cell walls), glucose and

xylose (monomers of cellulose and hemicellulose respectively), vanillin (a phenolic acid derived

from lignin), glycerol and palmitic acid (components of plant lipids), an amino acid mixture

(monomers of protein), and lactate and oxalate (major fermentation products from soil bacteria

and fungi).

Our multi-substrate design, where only the identity of the ^{13}C-substrate was varied (Fig.

1), allowed us to systematically investigate C assimilation dynamics as a function of C source

while maintaining identical conditions across samples. In addition, by sampling over time we

focused on the ultimate fate of C within the bacterial community and not just the bacteria

responsible for initial substrate breakdown. That is, our goal was not to differentiate primary

assimilation from later stages, but rather to map bacterial guild structure by examining how C

from diverse sources infiltrates the community over time. Guilds are groups of organisms that

access resources in a similar manner (38). We used C assimilation dynamics to infer guild

structure based on the source and timing of the ^{13}C assimilated by bacterial taxa, such that

bacteria within a guild exhibit similar patterns of C assimilation over time. We predicted that

variations in C assimilation dynamics across bacterial taxa would be poorly explained by

phylogenetic relatedness. Rather, ecological and physiological characteristics would be shared

across taxa grouped into C assimilation guilds.

Results and discussion

Carbon source bioavailability explains carbon mineralization rates

We tracked C mineralization dynamics from each C source for up to 48 days by

measuring ^{13}CO$_2$ production. Peak ^{13}C mineralization rates occurred on day 1 for glucose and

amino acids; day 2 for xylose, glycerol, and vanillin; day 3 for lactate; day 7 for oxalate; day 10

for cellulose; and day 14 for palmitic acid (Fig. 2A, Fig. S1). Cumulative ^{13}C mineralization

varied substantially between sources, being lowest for vanillin (42% C mineralized) and highest

for oxalate (88% C mineralized, Fig. 2B, Fig. S2). As expected, total C mineralization (^{13}C +

^{12}C) was similar across all treatments (Fig. S1) since all microcosms received the same set of

substrates with only the identity of the [13]C-labeled substrate varied. These mineralization

dynamics were confirmed in an independent experiment with soils from the same site (Fig. S2).

Substrate bioavailability predicts degradation rate in soil (39), and so we selected C

sources that differ in their solubility and hydrophobicity (as defined by octanol-water partition

coefficients predicted using the XLogP3 model (40)) as described in Supplemental Methods and

indicated in Fig. 1. We found a significant and strong positive correlation between LogP and the

day of peak C mineralization across C sources (Pearson's r = 0.932, p-value = 0.001; Fig S3A)

and a significant negative correlation between LogP and the maximum C mineralization rate

(Pearson's r = -0.855, p-value = 0.007; Fig S3B). These findings confirm that C sources with low

bioavailability, such as cellulose (an insoluble polymer) and palmitic acid (a waxy fatty acid), are

degraded slowly, while those with high bioavailability, such as glucose, xylose, amino acids, and

lactate are degraded rapidly. Vanillin (an aromatic molecule) had intermediate bioavailability

allowing it to be mineralized rapidly, however, its hydrophobicity favors adsorption to organic

surfaces in soil (41, 42). Hence, while vanillin in the aqueous phase degrades rapidly, its sorption

to organic matter may limit microbial access over time. Highly oxidized compounds such as

oxalate favor allocation of C to catabolic processes, resulting in greater cumulative

mineralization (43, 44). Given that LogP cannot be determined for an insoluble material such as

cellulose, and that there was a strong relationship between LogP and day of peak mineralization

(Fig. S3A), in subsequent analyses we defined bioavailability operationally based on observed

day of peak mineralization.

Non-cultivated bacteria are major players in soil C-cycling

The temporal dynamics of bacterial C assimilation were assessed by performing MW-

HR-SIP for each of the nine ^{13}C-labeled C sources over 48 days. Sampling times were selected

based on the ^{13}C mineralization dynamics for each substrate. A total of 12,394 unique bacterial

OTUs were observed and 6,613 of these passed independent filtering on the basis of sparsity.

From this set, MW-HR-SIP identified 1,286 "incorporators" (*i.e.,* OTUs with significant

evidence of ^{13}C incorporation into DNA; Fig. 3; Supplemental Dataset). We also examined

unfractionated DNA (*i.e.,* DNA extracted directly from soil and not subjected to MW-HR-SIP) in

order to evaluate overall change in microbial community composition over time. The majority of

incorporators were also detected in unfractionated DNA (1,153, or about 90%), though 133

incorporators (~10%) were not observed in unfractionated DNA. This result indicates that rare

OTUs, not readily detected in 16S rRNA gene surveys, are active participants in soil C-cycling.

Density gradient centrifugation fractionates DNA by buoyant density with sequencing performed

across many gradient fractions, and this approach translates into far greater sequencing depth

than is typical in microbial community analyses, allowing for the detection of rare taxa not

readily observed when sequencing unfractionated soil DNA.

Only 13.6% of the incorporators matched cultivated isolates (*i.e., >* 97% 16S rRNA gene

sequence identity to known isolates), while 59.6% lacked a closely related isolate (*i.e., <* 97%

16S rRNA gene sequence identity), and 26.8% were only distantly related to any isolate (*i.e., <*

90% 16S rRNA gene sequence identity). C from low and intermediate bioavailability sources

(*i.e.,* cellulose, palmitic acid, and vanillin) was assimilated primarily by non-cultivated OTUs

(Fig. S4). In contrast, C from high bioavailability sources (*i.e.,* glucose, xylose, amino acids,

glycerol, oxalate, and lactate) was assimilated primarily by OTUs matching known isolates (Fig.

S4). Microbes that access C from low bioavailability sources are likely to have traits (*e.g.,* slow

growth, metabolic dependency, surface attachment) that diminish their tolerance for laboratory

growth media and may limit their representation in culture collections (45, 46).

Phylogenetic conservation of C assimilation dynamics

The ecophysiology of bacterial OTUs is often inferred from their phylogenetic

relatedness (47–50). We therefore sought to determine if C assimilation activity could be

predicted on the basis of phylogeny. We used two methods to test for the phylogenetic

conservation in C assimilation dynamics. First, we calculated dissimilarity in assimilation

dynamics between OTUs. Patterns of C assimilation were weakly conserved among closely

related taxa and dissimilarity in C assimilation dynamics increased rapidly with phylogenetic

distance (Fig. 4A-B). Even when analyzing C assimilation dynamics independently for each C

source, phylogeny was a poor predictor of realized activity (Fig. S5). Second, we used

consenTRAIT (49) to measure the phylogenetic depth (*i.e.,* 16S rRNA gene evolutionary

distance; τ_D) at which patterns of C assimilation were conserved. We found τ_D values ranging

from ~0.001 to 0.025 across the nine C sources (Fig. 4C), consistent with τ_D values of C

substrate utilization measured by Martiny et al. (49). There was a significant correlation between

τ_D and time of C assimilation (Pearson's $r = 0.791$, p-value < 0.001; Fig. 4C). More specifically,

phylogenetic depth was highest for C assimilation from vanillin, cellulose, and palmitic acid

when assimilated at later time points, though the phylogenetic clustering of bacteria that

assimilated C from these sources was not statistically significant across all time points

(consenTRAIT; adjusted p-value ≥ 0.05; Fig. 4C). These results suggest that the assimilation of

C from low bioavailability sources requires metabolic capabilities or traits that are more deeply

conserved than those required to assimilate C from high bioavailability sources. The results also

show that phylogenetic conservation of C-cycling process is highly variable, being higher for

some processes than for others. Even within a single substrate such as cellulose, the degree of

phylogenetic clustering depended on time of assimilation. High variation in phylogenetic

clustering for taxa that access C from one source at different times suggests that phylogenetic

conservation might be driven by shared ecology rather than shared catabolic pathways.

C assimilation dynamics define bacterial guild structure

Since phylogenetic relatedness was a weak predictor of C assimilation dynamics in soil,

we sought an alternative means to group bacteria into ecologically relevant clusters. We used

unsupervised clustering to group incorporators into operationally defined guilds based on their

^{13}C-labeling dynamics. This approach generated 28 guilds (Fig. 4D). Each guild contained

between 15 and 122 incorporators and represented 3 to 13 different phyla (Supplemental

Dataset). We then examined characteristic growth and ^{13}C assimilation dynamics across guilds

and related these characteristics with rRNA operon (*rrn*) copy number, a trait with known

implications for bacterial ecology and C-cycling activity (51, 52). In all cases, characteristics and

rrn copy number was averaged across all OTU within a guild (Supplemental Dataset).

After adjusting for multiple comparisons ($n = 4$), we found a positive correlation only

between *rrn* copy number and the maximum $\log_2$ fold change in normalized abundance (max

L_2FC; Pearson's r = 0.503, p-value = 0.025; Fig. 5D)). Max L_2FC was defined as the maximum

change in differential abundance before and after C sources were added to soil. High copy

number guilds exhibited large changes in normalized abundance in response to C input, while

low copy number guilds varied much less over time. In other words, guilds with high copy

number exhibited highly dynamic populations while populations of guilds with low copy number

were less dynamic. Since sequencing is compositional (53) and since several of the bacterial

groups with the largest growth response had high copy number (Supplemental Dataset), relative

abundance was normalized both by predicted *rrn* copy number and DNA yield for each sample.

The *rrn* copy number of bacteria grown in culture correlates with maximal growth rate as well as

C use efficiency, and has been proposed as a component of microbial life history strategies (51,

52). Our observation links the *rrn* copy number of guilds to their *in situ* population dynamics in

response to C addition. There were two guilds with high *rrn* copy numbers (guilds 1 and 19),

when removed from the analysis the relationship with maximum $\log_2$ fold change remained

marginally significant (Pearson's r = 0.458, p-value = 0.074; Fig. S6).

209   We did not find any correlation between *rrn* copy number and the number of sources

from which guilds derived their C (Pearson's r = 0.051, p-value = 0.797; Fig. 5A), the averaged

bioavailability of C source that led to ^{13}C-labeling of the guilds (Pearson's r = 0.428, p-value =

0.093; Fig 5B), or the 'latency' of ^{13}C-labeling (Pearson's r = -0.339, p-value = 0.077; Fig. 5C).

Here, we defined bioavailability operationally as a function of the time when peak ^{13}C-

mineralization was observed, such that higher values are more available (Fig. S3). We defined

latency as the relationship between the time of peak ^{13}C-mineralization and the time ^{13}C-labeling

was first detected. A lower value for latency indicates that C assimilation took place when C

mineralization was maximal, and hence the organism is likely using C directly from the source or

is coupled tightly to the other organisms participating in mineralization. A higher value for

latency indicates that C assimilation took place long after rates of C mineralization began to

decline, making it more likely that the organism was assimilating C that had previously been

transformed by microbial processing.

To further demonstrate the ecological relevance of this guild structure, we examined how

phylogenetic distance among guild members, measured with the nearest taxon index (NTI),

varied with respect to guild growth and ^{13}C-assimilation dynamics. We expect that variation in

selection pressure between guilds should alter phylogenetic clustering observed within guilds.

For example, we predict greater competition for C from low bioavailability sources than from

high bioavailable sources when C is added as a pulse. This expectation is based on the idea that

consumption of low bioavailability substrates is constrained by diffusion and defined by spatial

relationships. While high bioavailability pulsed resources (*e.g.,* glucose) are available in great

excess relative to the initial microbial biomass present in soil, causing competition to be

minimal. Competitive exclusion is expected to cause phylogenetic overdispersion (low NTI).

After adjusting for multiple comparisons ($n = 4$), we found positive correlations between NTI

and both C source bioavailability (Pearson's $r = 0.64$, p-value $= 0.001$; Fig. S7B) and population

dynamics (Pearson's $r = 0.51$, p-value $= 0.022$; Fig. S7D) of guilds. No correlation was observed

with the number of C sources (Pearson's $r = 0.453$, p-value $= 0.062$; Fig. S7A) nor C

assimilation latency (Pearson's $r = -0.226$, p-value $= 0.994$; Fig. S7C). These results indicate that

competitive interactions within guilds are linked to the manner of C acquisition (i.e., the

bioavailability of source C), and the growth dynamics of guild members.

While *rrn* copy number is one trait that can inform a guild's ecology and relationship to

SOC, traits often impose tradeoffs causing interactions that can confound simple correlations.

We therefore used a principal component analysis (PCA) to explore variation in guild responses

as a function of their characteristic growth and assimilation dynamics along with *rrn* copy

number (Fig. 5E). The first principal component (PC1) explained 36.8% of variation in guild

responses and this axis corresponds with variation in *rrn* copy number (43.1% loading), growth

dynamics (max L_2FC; 28.5% loading), latency (14.1% loading), C source bioavailability (11.9%

loading), and number of C sources (2.48% loading). The second principal component (PC2)

explains 22.7% of the variation in guild responses. This second axis largely corresponds with

variation in C source bioavailability (44.5% loading), latency (35.2% loading), number of C

sources (16.3% loading), growth dynamics (3.8% loading), and *rrn* copy number (0.3% loading).

The overall guild structure suggests three extremes in guild characteristics, with certain

characteristics maximized at each vertex (Fig. 5E). For example, guilds with the highest *rrn* copy

number and growth dynamics (max L_2FC) have low PC1 but intermediate PC2. Guilds

assimilating C from the least bioavailable and fewest C sources have high PC1 and low PC2.

Guilds with the highest latency in C assimilation, and greatest diversity of C assimilation

patterns (C sources) have high PC1 and high PC2. These vertices represent the characteristic

extremes of the bacterial guild structure, though many intermediate guilds exist.

We selected the most abundant OTUs (based on the summed normalized abundance

across time) that represented the three vertices of the guild structure to illustrate the maximal

differences between guilds. The most abundant OTU from guild 19 was OTU.7163, classified as

a *Pseudomonas* species. This pseudomonad was estimated to have ~5 *rrn* operons, its abundance

increased rapidly within a day following C input from a very low initial abundance, then dropped

dramatically in abundance over time, and it assimilated C from glucose and cellulose at early

time points (Fig. 6). Correspondingly, guild 19 has a high max L_2FC, intermediate C source

bioavailability, and low latency (Fig. 5). The most abundant OTU from guild 14 was OTU.22,

classified as an *Agromyces* species. This microbacterium was estimated to have ~1 *rrn* operon, it

increased in abundance after C input from a relatively high initial abundance, reaching maximal

abundance after 6 days followed by a decline over time, and it was consistently [13]C-labeled

throughout the experiment assimilating C from multiple sources (Fig. 6). Correspondingly, guild

14 has an intermediate max L_2FC, many C sources, and high latency (Fig. 5). The most abundant

OTU from guild 2 was OTU.197, classified as a species of *Verrucomicrobia*. This

verrucomicrobium was estimated to have ~2 *rrn* operons, it increased in abundance very slowly

reaching maximal abundance after 30 days, remaining at relatively low abundance throughout,

and assimilated C from glucose, vanillin, and cellulose at mostly later time points (Fig. 6).

Correspondingly, guild 2 has a low max L_2FC, low C source bioavailability, and low latency

(Fig. 5). Generalized growth responses for all guilds are visualized in Figure S8.

It is clear from Figures 5, 6, and S8 that guilds differ in their growth response (max

L_2FC), and access to carbon (latency, C sources, bioavailability) and these differences suggest

tradeoffs between growth dynamics and the ability to access C in the soil environment over time.

Thus, the guild structure we identified can be interpreted in the context of life history theory,

specifically Grime's C-S-R framework (54). While Grime's framework was conceived to

describe plant life history strategies, it has recently been applied to microbial systems (55–57).

Ruderal (R) strategists should have little investment in resource acquisition, being adapted for

rapid growth in response to a sudden influx of nutrients. From our guild structure, ruderal

bacteria are exemplified by the pseudomonads and bacilli of guild 19. These taxa have high *rrn*

copy number, highly dynamic growth response (max L_2FC), low latency, and [13]C-labelling tends

to occur immediately following nutrient addition, all together suggesting the ability to grow

rapidly as a strategy for exploiting episodic nutrient pulses. Competitive (C) strategists are

adapted for resource acquisition, likely allocating energy to features such as mycelia, biofilms,

extracellular enzymes, siderophores, and antimicrobials that require substantial investment of

energy but increase fitness under conditions defined by prolonged competition for resources.

Thus, competitors should grow more slowly (*i.e.,* lower maximum growth rate), with less

dynamic swings in population size. In our guild structure, competitors are exemplified by the

Agromyces from guild 14. These taxa have higher latency suggesting the ability to access ^{13}C

following metabolic processing by other microbes, and they access C from a greater diversity of

sources and over longer periods of time than ruderals. Both ruderals and competitors can be

considered copiotrophs because of their ability to quickly respond to nutrient inputs (55), but

these strategies would manifest differently in their population dynamics and sensitivity to non-

equilibrium states (*e.g.,* disturbance). Stress tolerators (S) were defined by Grime on the basis of

abiotic stress (54), but we propose that for soil bacteria these organisms are adapted for life at

low energy flux (*i.e.*, conditions that would be stressful and result in starvation or dormancy for

copiotrophs). In soil, we therefore propose that stress tolerators are synonymous with

oligotrophs, organisms adapted to access low bioavailability C sources that would be otherwise

unavailable to growth adapted copiotrophic organisms. Thus, in our guild structure, stress

tolerant (S) strategists are those exemplified by the less well characterized *Verrucomicrobia* of

guild 2. These taxa have low *rrn* copy number and very slow growth response, but also low

latency and a tendency to access C from low bioavailability sources. While we focus on guilds

19, 14, and 2 to illustrate the presence of tradeoffs, Figure 5 illustrates that there is a wide

diversity of strategies for accessing C from soil, including intermediate strategies not well binned

strictly as ruderals, competitors, or stress tolerators (Fig. 5, S8). Similarly, members from the

same taxonomic group may display dissimilar strategies. For example, while some *Pseudomonas*

species exemplify the ruderal strategy, others can be found in guilds representing competitors or

stress tolerators (Supplemental Dataset), demonstrating the physiological diversity and versatility

of this genus.

Finally, we hope that is abundantly clear that the ecological tradeoffs that define fitness

in soil are surely different from those that define fitness in other habitats (e.g. aquatic, gut, or

extreme environments, etc.), and the categories we define here might not reflect well the

determinants of microbial fitness in non-soil habitats. To illustrate this idea, consider the

copiotroph-oligotroph framework, which was first used to describe the life history strategies of

aquatic bacteria. Aquatic oligotrophs thrive at low substrate concentrations where substrate flux

across the membrane is low and this imposes selection for high metabolic efficiency (58–65).

Aquatic oligotrophic habitats are typically characterized by substrates that have low

concentration at steady state (66). Under such equilibrium conditions, oligotroph growth rates are

governed by substrate affinity as described by Michaelis-Menton kinetics (58). Soils, however,

represent a highly dynamic, non-equilibrium system in which substrate availability varies over

time and space in response to nutrient inputs, water availability, and substrate-mineral or

substrate-organic matter interactions (67). Further, soil drying limits diffusion dramatically,

resulting in periods of substrate limitation even if nutrients are otherwise abundant in bulk soil,

while soil wetting enhances diffusive transport of soluble nutrients. Hence, wetting and drying of

soil have major impacts on nutrient availability dynamics. Insoluble substrates, and those

substrates adsorbed to surfaces, have low bioavailability being unable to cross the cytoplasmic

membrane. Growth on these substrates is limited by the diffusion of enzymes and enzyme

products between cells and their substrates. As a result, we predict that access to C from low

bioavailability substrates favors oligotrophs in soil, because limitations on diffusion and

transport should select for metabolic efficiency. Furthermore, diffusive fluxes in soil

environments are favored by intimate contact between organism and substrate. Such close

contacts persist in water films that coat soil aggregates, and remain in place as soils dry and

diffusion is limited by increasing tortuosity. Hence, we predict that oligotrophic activity persists

over a wide range of moisture availability in soils and that such activity is mediated by diverse

collections of interacting organisms. Since aquatic oligotrophs are not constrained by diffusion

and tortuosity, but rather by substrate concentration in solution, we expect they will be

differentiated considerably from soil oligotrophs. In addition, we predict that access to C from

more soluble, bioavailable substrates in soil favors copiotrophs with ruderal or competitive

strategies. These substrates are ephemeral in nature outside of the rhizosphere, being transiently

available in high concentration as a pulsed resource following wetting events. When soils are

dry, water-soluble substrates are poorly available due to diffusional limitation. Wetting provides

a flush that delivers these substrates into pore water where they are more accessible and where

diffusion is less limited because tortuosity is low. The pulsed nature of soluble organic matter in

soil pore water will favor rapid utilization of the C when nutrients are transiently present in

excess, because the benefits of high affinity only matter when resources are present at low

concentration and the benefits of high efficiency only matter when resources are present at

steady state.

High turnover of bacterial biomass

In terms of understanding microbial contributions to soil C cycling, much focus has been

placed on primary microbial degradation of C derived from plants, and the degradation of plant

biomass and exudates is typically considered in relation to microbial catabolism. However,

recent evidence suggests that persistent SOM is largely composed from the products of microbial

anabolic reactions (68). Hence, microbial turnover and mortality is likely of considerable

importance in soil C-cycling. We observed rapid growth in response to C inputs, CO_2 flux

peaked at 2 days, and soil DNA increased over 6 days (a 52% increase), but there was also

evidence of substantial turnover and mortality. For example, soil DNA declined considerably

from day 6 through day 48 (a 20% decrease), and the normalized relative abundance of many

taxa decreased over time. Of the ^{13}C-labeled taxa, 387 were observed at high frequency over

time (detection in 6 of 7 time points) in unfractionated soil DNA. Of these high frequency OTUs

only 59 (15%) were observed to have a maximum in normalized relative abundance at the end of

the experiment (*i.e.*, no evidence of mortality), while 221 (57%) decreased by more than 50% in

normalized relative abundance from an early peak in abundance. Decreases in normalized

relative abundance were often dramatic and rapid (Fig. 6), particularly for guilds having high

max L$_2$FC (Fig. 5). In addition, among these frequently observed ^{13}C-labeled taxa, a total of 202

(52%) were ^{13}C-labeled at an early time point but were unlabeled by that same source at later

time points. Loss of ^{13}C-label can result from growth and isotope dilution over time, but when

coupled to evidence of decreases in mineralization rate, decreases in DNA yield, and decreases

in normalized relative abundance of most taxa over time, this result suggests mortality and

turnover of ^{13}C-DNA. Finally, the high latency for many ^{13}C-labeled taxa (Fig. 5) indicates that

many taxa became ^{13}C-labeled indirectly by consuming microbial products rather than by direct

assimilation of added C. Collectively these results argue that secondary processing, fueled by the

products of microbial anabolic metabolism, comprise a major component of the C that cycles

through soil communities. Based on our guild structure we would predict that much of this

microbial biomass C ultimately flows through taxa having more competitive life history

strategies (those with high latency in Fig. 5).

Guild structure explains bacterial biogeography

To test for a linkage between guild structure and biogeography, we mapped incorporators

to global bacterial surveys to determine whether guild membership explained variation in

biogeography of soil bacteria. We employed RLQ analysis, a multivariate approach that finds

correspondence between organism properties (R) and environmental parameters at various sites

(Q; *e.g.,* habitat or soil pH) based on organismal abundance within sites (L) (69, 70). We mapped

incorporator OTUs and their guild membership to two 16S rRNA gene surveys of soil bacterial

diversity spanning continental (QIITA study 619) and global scales (QIITA study 928) (71).

Guild membership explained significant variation in microbial relative abundance across

environmental parameters (Table S1; Fig. S9). For simplicity, guilds were divided into three

groups based on C assimilation profile. Group D assimilated C primarily from dissolved

substrates of high bioavailability (glucose, xylose, amino acids, glycerol, lactate, and oxalate).

Group P assimilated C primarily from particulate substrates of low bioavailability (cellulose and

palmitic acid). Group V assimilated C from vanillin; this group was treated separately as it was

likely present in both aqueous and noaqueous forms with transition between these two states

changing over time (owing to its insolubility and tendency to sorb to SOM). Guilds were further

grouped by time of labelling: early (E) and late (L). At continental scales, guilds that primarily

used dissolved substrates at late timepoints were associated with forest soils, and soils having

high organic C and N (DL; Fig. S9A). At global scales, these guilds were positively associated

with latitude (DL; Fig. S9B), likely as a result of the northern concentration of forest ecosystems

and SOM at global scales. The guilds most strongly associated with tropical and subtropical

broadleaf forests were those that assimilated C from particulate substrates early (PE; Fig. S9B).

The guilds that were most strongly associated with temperate grasslands, savannas, and

shrubland were those that assimilated C from diverse substrates (DL, PL, and VL; Fig. S9B).

Guilds that used C from particulate substrates and vanillin at later timepoints were most closely

associated with desert biomes (PL and VL; Fig. S9A). These results show that patterns of

microbial C assimilation can be linked to patterns of microbial biogeography and that guild

structure has relevance across large biogeographical scales.

Conclusions

We conducted a multi-substrate DNA-SIP experiment to measure the *in situ* activities of

bacteria in the soil C-cycle. We show that degradation dynamics and patterns of bacterial C

assimilation depended on the bioavailability of C inputs. In total, 1286 bacteria were found to

incorporate C derived from 9 different sources, and most of these bacteria were distantly related

to cultivated isolates. We found that C assimilation dynamics in soil had low phylogenetic

conservation among bacteria. Instead, C assimilation dynamics allowed us to group bacteria into

guilds whose properties were generally consistent with well-known life history strategies. We

also show that this guild structure can explain variation in bacterial biogeography. Our results

demonstrate that life history strategies are likely linked to C assimilation dynamics. SOM is

largely derived from the products of anabolic microbial metabolism (14), and therefore microbial

assimilation and turnover of C is a primary determinant of C fate in soil. Based on our results, we

hypothesize that C fate in soil is not determined solely, or even primarily, by the metabolic

pathways that enable its catalysis, but rather by the life history traits that govern microbial

growth and C acquisition *in situ*. As a result, life history theory could be useful in predicting the

C-cycle activity of microbial communities (18). While more research is necessary to better

understand the genetic and ecological underpinnings of life history strategies in soil bacteria, our

results add to the growing body of knowledge indicating the importance of life history theory to

bacterially mediated terrestrial C-cycling.

Materials and methods

Soil microcosm experiments

Soil microcosms were designed and incubated with ^{13}C-labeled and unlabeled substrates

as previously described (35) except that C amendment consisted of nine substrates: cellulose,

xylose, glucose, glycerol, vanillin, palmitic acid, amino acid mixture, lactate, and oxalate (SI;

Table S2). For most substrates, octanol-water partition coefficient (LogP) was determined using

XLogP3 through PubChem (40). Since amino acids were added as a mixture, this substrate LogP

was averaged across the 20 common amino acids. LogP is not available for cellulose as it is

insoluble. Each substrate was added at 0.4 mg C g^{-1} soil. Only one of the nine substrates was ^{13}C-

labeled in each treatment while the remainder were unlabeled (Fig. 1). Soil was from Penn Yan,

New York, USA , with site and collection methods previously described (35, 72) and outlined in

SI. ^{13}C and ^{12}C mineralization dynamics were determined based on concentrations of $^{12}CO_2$ and

$^{13}CO_2$ in microcosm headspace measured over time as detailed in SI. Soil from microcosms were

destructively sampled depending on substrate mineralization dynamics over days 1, 3, 6, 14, 30,

and 48 after substrate addition.

DNA extraction and isopycnic centrifugation

DNA extraction and isopycnic centrifugation was conducted as previously described (35)

and detailed in SI. In short, for each treatment, size selected DNA (≥ 4 kb) (73) was centrifuged

in a CsCl density gradient and 100 µl fractions were collected. An aliquot of DNA taken before

isopycnic centrifugation was used for whole microcosm bacterial community sequencing (*i.e.,*

unfractionated DNA).

16S rRNA gene amplification and sequencing

We amplified and sequenced the V4 region of the 16S rRNA gene as previously

described (74) and detailed in SI using dual indexed primers (515f and 806r) developed by

Kozich *et al.* (75). 16S rRNA amplicon libraries were processed as described previously (35) and

detailed in SI. OTUs were clustered at 97% sequence similarity. Raw sequencing reads can be

accessed at the NCBI Short Read Archive (accession PRJNA668741).

Identifying incorporators

OTUs that incorporated ^{13}C into their DNA (*i.e.,* incorporators) were identified with

MW-HR-SIP (36) using development code for the HTSSIP R package (76). In short, we used

DESeq2 (77) to identify OTUs significantly enriched in the "heavy" gradient fractions of the

^{13}C-labeled treatments compared to the "heavy" gradient fractions in corresponding ^{12}C-control.

Incorporators were defined as OTUs with a $\log_2$ fold enrichment greater than 0.25 and a

Benjamini-Hochberg adjusted p-value less than 0.1. We used the overlapping BD windows 1.70-

1.73, 1.72-1.75, and 1.74-1.77 g/ml, which were shown via simulations to have the highest

sensitivity across all tested scenarios while not losing specificity (36). Independent sparsity

filtering was conducted for each BD window to minimize the number of comparisons performed.

The taxonomic novelty of incorporators was determined by aligning OTU representative

sequences to the Silva all-species living tree project small subunit database version 123,

representing all sequenced type strains curated at that time (78). Alignment was performed using

nucleotide BLAST (79) with an expected value cutoff of 1E-20. Incorporators with no hits $\geq$

97% sequence identity were considered to have no closely related isolates, while those with no

hits ≥ 90% were considered to have no related isolates. Statistical analyses were conducted in R

version 3.1.2 (80).

Analyses of phylogenetic conservation

Fasttree v2.1.9 (81) was used to infer a maximum likelihood phylogeny from the 16S

rRNA sequence alignment of representative OTUs. The phylogeny was rooted with *Sulfolobus*

solfataricus DSM1616 (Genbank accession X90478) as the outgroup. Phylogenetic conservation

of realized function was examined both by comparing functional distance with phylogenetic

distance and testing for a phylogenetic signal with a modified version of consenTRAIT (49)

(https://github.com/nick-youngblut/consenTRAIT).

Analysis of guild structure

Incorporators were grouped into guilds based on their ^{13}C-labelling patterns as

determined by $\log_2$ fold change in ^{13}C-labeled treatment fractions versus corresponding

unlabeled control fractions. Specifically, clustering was performed by: *i*) calculating pairwise

Gower distances for each OTU pair, *ii*) hierarchically clustering (UPGMA agglomeration

method) based on the resulting distance matrix, *iii*) generating a dendrogram of the hierarchical

associations, *iv*) detecting clusters in the dendrogram using the *cutreeHybrid* function in the

DynamicTreeCut R package (82).

The *rrn* copy number of each OTU was predicted as described previously (83). Briefly,

pplacer v1.1 (84) was used to insert OTU representative sequences into a 16S rRNA gene

reference phylogeny of bacteria with known *rrn* copy number. The resulting phylogeny was used

by guppy v1.1 to predict *rrn* copy numbers of each OTU based on its relatedness to the reference

taxa. The natural log of the copy number was used for all comparisons (51).

Four guild C assimilation and growth characteristics were calculated. All characteristics

were first calculated for individual OTUs then averaged within each guild. The number of C

sources from which C was assimilated was simply the number of C sources from which an OTU

was ^{13}C-labeled at any timepoint. Since logP is not available for cellulose, C source

bioavailability was defined operationally based on the day of maximal ^{13}C mineralization rate for

each substrate (as detailed in SI). The C source bioavailability was determined as the average of

all sources from which ^{13}C was assimilated. Latency of C assimilation was determined by the

natural log of the ratio between the first day of ^{13}C-labelling and the day of peak C

mineralization for each C source. This value was then averaged across all sources from which C

was assimilated. The maximum $\log_2$ fold change in OTU abundance in the unfractionated DNA

was calculated following abundance normalization to minimize bias due to compositional data.

Normalization included two steps: normalizing for predicted *rrn* copy number and normalizing

by sample DNA yield. OTU relative abundances were first divided by their predicted rRNA

operon copy number, then by the estimate copy number for the entire community. Within each

timepoint, normalized relative abundances were then multiplied by the average DNA yield

across the replicate microcosms. The extracted DNA yield (ng DNA g^{-1} dry weight soil) was

quantified with the Quant-iT PicoGreen dsDNA Assay Kit (Thermo Fisher Scientific, Waltham,

MA, USA). We recognize that this calculation does not take into account inflation in DNA yield

due to the growth of fungi and other organisms not accounted for in our study, however we

believe that these effects will be small compared to the effects of bacterial growth, particularly as

we are simply comparing between bacterial OTUs within this study. Untreated bulk soils were

used as our baseline abundance, with OTUs undetected in these soils assigned the lowest

abundance measured. For each OTU, the $\log_2$ fold change in abundance between this baseline

and the timepoint when abundance was highest was then calculated. If abundance at all

timepoints was less than the baseline or undetected, the maximum abundance was assigned the

baseline abundance, making the $\log_2$ fold change = 0 for such OTUs. If an OTU is ^{13}C-labeled, it

is growing on the substrates provided. A decrease in abundance at a time when taxa are labeled

indicates either that the rate of mortality for the population exceeds the division rate (deaths >

births), that labelling occurred prior to growth decline, or that normalization was not entirely

successful at eliminating all variance due to compositional sequencing.

After averaging across all member OTUs within each guild, we measured the relationship

between the natural log of the rRNA operon copy number and each of the four C assimilation

characteristics using the Pearson's correlation. We also measured the relationship between all

four C assimilation characteristics and the nearest taxon index (NTI) within guilds. NTI was

calculated using R package picante (85). *rrn* copy number and all four of characteristics were

used in a principal component analysis (PCA) to more fully assess the variation in these

characteristics across guilds.

Mapping incorporators to biogeography datasets

Detailed descriptions of the independent datasets and their analysis for this study are

found in SI. In short, OTU count tables, representative 16S rRNA gene sequences, and metadata

was downloaded from https://qiita.ucsd.edu/ for QIITA studies 619 (continental dataset) and 928

(global dataset). Incorporators were mapped to external dataset OTUs separately for each study

with the mothur alignment tool (86) using the Silva reference alignment as a template. To

examine the biogeography of guilds at continental (QIITA 619) and global (QIITA 928) scales,

guilds were grouped by assimilation dynamics (as described in results). Then for each QIITA

dataset, guild designations (R) of mapped OTUs were used in combination with their OTU count

tables (L) and sample metadata (Q) for RLQ and forth corner analyses (70, 87) as detailed in SI.

Acknowledgements

We thank Chuck Pepe-Ranney, Ashley Campbell, Marquessa Henry, and Braulio Castillo, Anay

Hindupur, and Jillian Waters for their help with experimental design and implementation.

Funding

This material is based upon work supported by the Department of Energy Office of Science,

Office of Biological & Environmental Research Genomic Science Program under Award

Numbers DE-SC0004486 and DE-SC0010558.

References

1. E. G. Jobbágy, R. B. Jackson, The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecol. Appl.* **10**, 423–436 (2000).

2. V. Brovkin, *et al.*, Effect of anthropogenic land-use and land-cover changes on climate and land carbon storage in CMIP5 projections for the twenty-first century. *J. Climate* **26**, 6859–6881 (2013).

3. J. P. W. Scharlemann, E. V. J. Tanner, R. Hiederer, V. Kapos, Global soil carbon: understanding and managing the largest terrestrial carbon pool. *Carbon Manag.* **5**, 81–91 (2014).

4. Z. Luo, G. Wang, E. Wang, Global subsoil organic carbon turnover times dominantly controlled by soil properties rather than climate. *Nat. Commun.* **10**, 3688 (2019).

5. M. A. Bradford, *et al.*, Managing uncertainty in soil carbon feedbacks to climate change. *Nat. Clim. Change* **6**, 751–758 (2016).

6. R. Lal, Soil carbon sequestration impacts on global climate change and food security. *Science* **304**, 1623–1627 (2004).

7. R. Lal, Soil health and carbon management. *Food Enrgy. Security* **5**, 212–222 (2016).

8. D. W. Reeves, The role of soil organic matter in maintaining soil quality in continuous cropping systems. *Soil Till. Res.* **43**, 131–167 (1997).

9. P. Smith, Soils as carbon sinks: the global context. *Soil Use Manage.* **20**, 212–218 (2004).

10. M. Heimann, M. Reichstein, Terrestrial ecosystem carbon dynamics and climate feedbacks. *Nature* **451**, 289–292 (2008).

11. K. E. O. Todd-Brown, *et al.*, Changes in soil organic carbon storage predicted by Earth system models during the 21st century. *Biogeosciences* **11**, 2341–2356 (2014).

574   12.   Y. Luo, T. F. Keenan, M. Smith, Predictability of the terrestrial carbon cycle. *Glob.*

575   *Change Biol.* **21**, 1737–1751 (2015).

576   13.   W. R. Wieder, *et al.*, Carbon cycle confidence and uncertainty: Exploring variation among

577   soil biogeochemical models. *Glob. Change Biol.* **24**, 1563–1579 (2018).

578   14.   J. Schimel, S. Schaeffer, Microbial control over carbon cycling in soil. *Front. Microbiol.*

579   **3**, 348 (2012).

580   15.   K. E. O. Todd-Brown, *et al.*, Causes of variation in soil carbon simulations from CMIP5

581   Earth system models and comparison with observations. *Biogeosciences* **10**, 1717–1736

582   (2013).

583   16.   W. R. Wieder, *et al.*, Explicitly representing soil microbial processes in Earth system

584   models. *Global Biogeochem. Cy.* **29**, 1782–1800 (2015).

585   17.   G. Wang, W. M. Post, M. A. Mayes, Development of microbial-enzyme-mediated

586   decomposition model parameters through steady-state and dynamic analyses. *Ecol. Appl.*

587   **23**, 255–272 (2013).

588   18.   W. R. Wieder, A. S. Grandy, C. M. Kallenbach, G. B. Bonan, Integrating microbial

589   physiology and physio-chemical principles in soils with the MIcrobial-MIneral Carbon

590   Stabilization (MIMICS) model. *Biogeosciences* **11**, 3899–3917 (2014).

591   19.   B. N. Sulman, R. P. Phillips, A. C. Oishi, E. Shevliakova, S. W. Pacala, Microbe-driven

592   turnover offsets mineral-mediated storage of soil carbon under elevated CO_2. *Nat. Clim.*

593   *Change* **4**, 1099–1102 (2014).

594   20.   W. R. Wieder, G. B. Bonan, S. D. Allison, Global soil carbon projections are improved by

595   modelling microbial processes. *Nat. Clim. Change* **3**, 909–912 (2013).

596   21.   L. A. Hug, *et al.*, A new view of the tree of life. *Nat. Microbiol.* **1**, 16048 (2016).

597   22.   K. G. Lloyd, A. D. Steen, J. Ladau, J. Yin, L. Crosby, Phylogenetically novel uncultured

598   microbial cells dominate earth microbiomes. *mSystems* **3**, e00055-18 (2018).

599   23.   A. D. Steen, *et al.*, High proportions of bacteria and archaea across most biomes remain

600   uncultured. *ISME J.* **13**, 3126–3130 (2019).

601   24.   M. G. I. Langille, *et al.*, Predictive functional profiling of microbial communities using

602   16S rRNA marker gene sequences. *Nat. Biotechnol.* **31**, 814–821 (2013).

603   25.   K. P. Aßhauer, B. Wemheuer, R. Daniel, P. Meinicke, Tax4Fun: predicting functional

604   profiles from metagenomic 16S rRNA data. *Bioinformatics* **31**, 2882–2884 (2015).

605   26.   M. Brbić, *et al.*, The landscape of microbial phenotypic traits and associated genes.

606   *Nucleic Acids Res.* **44**, 10074–10090 (2016).

607   27.   J. B. H. Martiny, S. E. Jones, J. T. Lennon, A. C. Martiny, Microbiomes in light of traits:

608   A phylogenetic perspective. *Science* **350**, aac9323 (2015).

609   28.   J. Choi, *et al.*, Strategies to improve reference databases for soil microbiomes. *ISME J.* **11**,

610   829–834 (2017).

611   29.   N. Fierer, *et al.*, Cross-biome metagenomic analyses of soil microbial communities and

612   their functional attributes. *P. Natl. Acad. Sci. U.S.A.* **109**, 21390–21395 (2012).

613   30.   J. Alneberg, *et al.*, Genomes from uncultivated prokaryotes: a comparison of

614   metagenome-assembled and single-amplified genomes. *Microbiome* **6**, 173 (2018).

615   31.   D. L. Wohl, S. Arora, J. R. Gladstone, Functional redundancy supports biodiversity and

616   ecosystem function in a closed and constant environment. *Ecology* **85**, 1534–1540 (2004).

617   32.   R. Evans, *et al.*, Defining the functional traits that drive bacterial decomposer community

618   productivity. *ISME J.* **11**, 1680–1687 (2017).

33.   M. G. Dumont, J. C. Murrell, Stable isotope probing — linking microbial identity to

function. *Nat. Rev. Microbiol.* **3**, 499–504 (2005).

34.   B. A. Hungate, *et al.*, Quantitative microbial ecology through stable isotope probing. *Appl.*

Environ. Microb. **81**, 7570–7581 (2015).

35.   C. Pepe-Ranney, A. N. Campbell, C. N. Koechli, S. Berthrong, D. H. Buckley, Unearthing

the ecology of soil microorganisms using a high resolution DNA-SIP approach to explore

cellulose and xylose metabolism in soil. *Front. Microbiol.* **7**, 703 (2016).

36.   N. D. Youngblut, S. E. Barnett, D. H. Buckley, SIPSim: A modeling toolkit to predict

accuracy and aid design of DNA-SIP experiments. *Front. Microbiol.* **9**, 570 (2018).

37.   D. Sijm, R. Kraaij, A. Belfroid, Bioavailability in soil or sediment: exposure of different

organisms and approaches to study it. *Environ. Pollut.* **108**, 113–119 (2000).

38.   D. Simberloff, T. Dayan, The guild concept and the structure of ecological communities.

Annu. Rev. Ecol. Syst. **22**, 115–143 (1991).

39.   B. Berg, K. Hannus, T. Popoff, O. Theander, Changes in organic chemical components of

needle litter during decomposition. Long-term decomposition in a Scots pine forest. *Can.*

J. Botany **60**, 1310–1319 (1982).

40.   T. Cheng, *et al.*, Computation of octanol–water partition coefficients by guiding an

additive model with knowledge. *J. Chem. Inf. Model.* **47**, 2140–2148 (2007).

41.   T. Polubesova, Y. Chen, R. Navon, B. Chefetz, Interactions of hydrophobic fractions of

dissolved organic matter with Fe^{3+}- and Cu^{2+}-montmorillonite. *Environ. Sci. Technol.* **42**,

4797–4803 (2008).

42.   M. Keiluweit, M. Kleber, Molecular-level interactions in soils and sediments: the role of

aromatic π-systems. *Environ. Sci. Technol.* **43**, 3421–3429 (2009).

43.   J. B. Brant, E. W. Sulzman, D. D. Myrold, Microbial community utilization of added

carbon substrates in response to long-term carbon input manipulation. *Soil Biol. Biochem.*

38, 2219–2232 (2006).

44. S. Manzoni, P. Taylor, A. Richter, A. Porporato, G. I. Ågren, Environmental and

stoichiometric controls on microbial carbon-use efficiency in soils. *New Phytol.* **196**, 79–

91 (2012).

45. K. Alain, J. Querellou, Cultivating the uncultured: limits, advances and future challenges.

Extremophiles **13**, 583–594 (2009).

46. V. H. T. Pham, J. Kim, Cultivation of unculturable soil bacteria. *Trends Biotechnol.* **30**,

475–484 (2012).

47. J. T. Lennon, Z. T. Aanderud, B. K. Lehmkuhl, D. R. Schoolmaster Jr., Mapping the niche

space of soil microorganisms using taxonomy and traits. *Ecology* **93**, 1867–1879 (2012).

48. S. A. Placella, E. L. Brodie, M. K. Firestone, Rainfall-induced carbon dioxide pulses

result from sequential resuscitation of phylogenetically clustered microbial groups. *P.*

Natl. Acad. Sci. U.S.A. **109**, 10931–10936 (2012).

49. A. C. Martiny, K. Treseder, G. Pusch, Phylogenetic conservatism of functional traits in

microorganisms. *ISME J.* **7**, 830–838 (2013).

50. K. L. Dolan, J. Peña, S. D. Allison, J. B. H. Martiny, Phylogenetic conservation of

substrate use specialization in leaf litter bacteria. *PLOS ONE* **12**, e0174472 (2017).

51. B. R. Roller, T. M. Schmidt, The physiology and ecological implications of efficient

growth. *ISME J.* **9**, 1481–1487 (2015).

52. N. Fierer, M. A. Bradford, R. B. Jackson, Toward an ecological classification of soil

bacteria. *Ecology* **88**, 1354–1364 (2007).

53. G. B. Gloor, J. M. Macklaim, V. Pawlowsky-Glahn, J. J. Egozcue, Microbiome datasets

are compositional: and this is not optional. *Front. Microbiol.* **8**, 2224 (2017).

54.   J. P. Grime, Evidence for the existence of three primary strategies in plants and its

relevance to ecological and evolutionary theory. *Am. Nat.* **111**, 1169–1194 (1977).

55.   J. I. Prosser, *et al.*, The role of ecological theory in microbial ecology. *Nat. Rev.*

Microbiol. **5**, 384–392 (2007).

56.   A. Ho, *et al.*, Conceptualizing functional traits and ecological characteristics of methane-

oxidizing bacteria as life strategies. *Env. Microbiol. Rep.* **5**, 335–345 (2013).

57.   N. Fierer, Embracing the unknown: disentangling the complexities of the soil microbiome.

Nat. Rev. Microbiol. **15**, 579–590 (2017).

58.   D. K. Button, Biochemical basis for whole-cell uptake kinetics: specific affinity,

oligotrophic capacity, and the meaning of the Michaelis constant. *Appl. Environ. Microb.*

57, 2033–2038 (1991).

59.   D. K. Button, Nutrient-limited microbial growth kinetics: overview and recent advances.

A. Van Leeuw. J. Microb. **63**, 225–235 (1993).

60.   C. E. Zobell, C. W. Grant, Bacterial utilization of low concentrations of organic matter. *J.*

Bacteriol. **45**, 555–564 (1943).

61.   J. R. Postgate, J. R. Hunter, The survival of starved bacteria. *Microbiology* **29**, 233–263

(1962).

62.   H. W. Jannasch, Growth of marine bacteria at limiting concentrations of organic carbon in

seawater. *Limnol. Oceanogr.* **12**, 264–271 (1967).

63.   S. I. Kuznetsov, G. A. Dubinina, N. A. Lapteva, Biology of oligotrophic bacteria. *Annu.*

Rev. Microbiol. **33**, 377–387 (1979).

64.   Y. Akagi, U. Simidu, N. Taga, Growth responses of oligotrophic and heterotrophic marine

bacteria in various substrate concentrations, and taxonomic studies on them. *Can. J.*

Microbiol. **26**, 800–806 (1980).

65. D. K. Button, Nutrient uptake by microorganisms according to kinetic parameters from

theory as related to cytoarchitecture. *Microbiol. Mol. Biol. R.* **62**, 636–645 (1998).

66. C. Arnosti, Microbial extracellular enzymes and the marine carbon cycle. *Annu. Rev. Mar.*

Sci. **3**, 401–425 (2011).

67. P. Sollins, P. Homann, B. A. Caldwell, Stabilization and destabilization of soil organic

matter: mechanisms and controls. *Geoderma* **74**, 65–105 (1996).

68. C. Liang, W. Amelung, J. Lehmann, M. Kästner, Quantitative assessment of microbial

necromass contribution to soil organic matter. *Glob. Change Biol.* **25**, 3578–3590 (2019).

69. S. Dolédec, D. Chessel, C. J. F. ter Braak, S. Champely, Matching species traits to

environmental variables: a new three-table ordination method. *Environ. Ecol. Stat.* **3**, 143–

166 (1996).

70. S. Dray, *et al.*, Combining the fourth-corner and the RLQ methods for assessing trait

responses to environmental variation. *Ecology* **95**, 14–21 (2014).

71. S. T. Bates, *et al.*, Examining the global distribution of dominant archaeal populations in

soil. *ISME J.* **5**, 908–917 (2011).

72. S. T. Berthrong, D. H. Buckley, L. E. Drinkwater, Agricultural management and labile

carbon additions affect soil microbial community structure and interact with carbon and

nitrogen cycling. *Microb. Ecol.* **66**, 158–170 (2013).

73. N. D. Youngblut, D. H. Buckley, Intra-genomic variation in G + C content and its

implications for DNA stable isotope probing. *Env. Microbiol. Rep.* **6**, 767–775 (2014).

74. S. E. Barnett, N. D. Youngblut, D. H. Buckley, Soil characteristics and land-use drive

bacterial community assembly patterns. *FEMS Microbiol. Ecol.* **96**, fiz194 (2020).

75. J. J. Kozich, S. L. Westcott, N. T. Baxter, S. K. Highlander, P. D. Schloss, Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl. Environ. Microb.* **79**, 5112–5120 (2013).

76. N. D. Youngblut, S. E. Barnett, D. H. Buckley, HTSSIP: An R package for analysis of high throughput sequencing data from nucleic acid stable isotope probing (SIP) experiments. *PLOS ONE* **13**, e0189616 (2018).

77. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

78. P. Yarza, *et al.*, The All-Species Living Tree project: A 16S rRNA-based phylogenetic tree of all sequenced type strains. *Syst. Appl. Microbiol.* **31**, 241–250 (2008).

79. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

80. R Core Team, R: A language and environment for statistical computing (2018).

81. M. N. Price, P. S. Dehal, A. P. Arkin, FastTree: Computing large minimum evolution trees with profiles instead of a distance matrix. *Mol. Biol. Evol.* **26**, 1641–1650 (2009).

82. P. Langfelder, B. Zhang, S. Horvath, Defining clusters from a hierarchical cluster tree: the Dynamic Tree Cut package for R. *Bioinformatics* **24**, 719–720 (2008).

83. S. W. Kembel, M. Wu, J. A. Eisen, J. L. Green, Incorporating 16S gene copy number information improves estimates of microbial diversity and abundance. *PLOS Comput. Biol.* **8**, e1002743 (2012).

84. F. A. Matsen, R. B. Kodner, E. V. Armbrust, pplacer: linear time maximum-likelihood

and Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*

Bioinformatics **11**, 538 (2010).

85.   S. W. Kembel, *et al.*, Picante: R tools for integrating phylogenies and ecology.

Bioinformatics **26**, 1463–1464 (2010).

86.   P. D. Schloss, *et al.*, Introducing mothur: open-source, platform-independent, community-

supported software for describing and comparing microbial communities. *Appl. Environ.*

Microb. **75**, 7537–7541 (2009).

87.   S. Dray, D. Chessel, J. Thioulouse, Co-inertia analysis and the linking of ecological data

tables. *Ecology* **84**, 3078–3089 (2003).

Figure 1: Diagram of the design of the DNA-SIP experiment. This experiment employed soil
microcosms, each amended with all nine C sources (added at 0.4 mg C per gram of soil), only
one of which was > 99% ^{13}C-labeled in treatment microcosms. Control microcosms had all nine
C sources added but none were isotopically labeled. Microcosms were destructively sampled at
multiple timepoints (black bars) based on mineralization rates from preliminary experiment.
Headspace samples were taken every 1-7 days. DNA extracted from microcosm soil was used
both for DNA-SIP and whole bacterial community sequencing (unfractionated). Bioavailability
estimated based on mineralization dynamics (Note that vanillin is highly hydrophobic and it
occurred in both aqueous and nonaqueous (sorbed or solid) phases, but only the aqueous phase
bioavailability is indicated here. Cellulose and palmitic acid were insoluble and were added as
powders.).
758
759

760
761
Figure 2: C source mineralization dynamics varied based on bioavailability. (A) [13]C
mineralization rates and (B) cumulative amounts of [13]C mineralized per gram dry weight of soil.
Inset for (A) displays a finer scale representation of the [13]C mineralization rates over days 0-10.
Error bars represent ± standard deviation among microcosm replicates ($n = 3$).
766
767

768
769
Figure 3: The dynamics of C assimilation varied across the 1,286 OTUs that exhibited
significant ^{13}C-labelling. OTUs are ordered by their phylogeny with Phylum, or Class for
Proteobacteria, indicated by different colors in the phylogenetic tree. Only Phyla/Classes with
10 or more ^{13}C-labeled OTUs are colored in this way with others colored grey. Each vertical
block indicates ^{13}C-labelling results for a single day as specified at the top. Each column within a
block indicates results for a given substrate as specified on the bottom. Rows display the ^{13}C-
labelling dynamics for each OTU as follows: red bars indicate ^{13}C-labelling, light grey bars
indicate OTU detection in the gradient but no evidence of labeling, white indicates the OTU was
not detected in the gradient. Phylum/Class abbreviations are Bact. = *Bacteroidetes*, Acido. =
Acidobacteria, Gemm. = *Gemmatimonadetes*, Chl-flx. = *Chloroflexi*, Firmi. = *Firmicutes*,
Actino. = *Actinobacteria*, Verruc. = *Verrucomicrobia*, Plancto. = *Planctomycetes*, Beta. =
Betaproteobacteria, Gamma. = *Gammaproteobacteria*, Delta. = *Deltaproteobacteria*, Alpha. =
Alphaproteobacteria.

Figure 4: (A) The Gower's distance in C assimilation dynamics (*i.e.,* realized functional distance) between OTUs increases rapidly in relation to their phylogenetic distance. The vertical dashed lines indicate distance values that correspond to estimates of taxonomic ranks for genus, family, order, class, and phylum (from left to right). OTU pairs are grouped into phylogenetic distance bins with error bars indicating ± standard error. (B) Mantel test comparing realized functional distance and phylogenetic distance between OTUs with points colored by significance of correlation. Note very low Mantel r. (C) Phylogenetic depth (Tau D) of C assimilation from each substrate at each timepoint derived from consenTRAIT tests. All p-values are globally adjusted with the Benjamini-Hochberg correction. There is a positive correlation between Tau D and time (Pearson's r = 0.79, p-value < 0.001). This linear relationship is represented by the red line with the ribbon representing the standard error. D) As phylogenetic relatedness poorly explained C assimilation dynamics, we instead grouped OTUs by their [13]C-labeling pattern into guilds. Here C assimilation dynamics of the 28 guilds (ordered based on similarity in C assimilation dynamics) is displayed, with red color intensity indicating the percentage of member OTUs who have detectable [13]C-labelling from each substrate on each day.

801
802
Figure 5: We propose that guild structure corresponds with life history strategies of bacteria. We
tested whether the *rrn* copy number of guilds was correlated with other guild characteristics. A)
The average number of C sources from which C was assimilated by each guild. B) The averaged
bioavailability of the sources from which C was assimilated by each guild. Bioavailability is
defined here as a unitless value determined operationally based on mineralization dynamics for
each substrate, with higher values for more rapidly mineralized substrates. C) The averaged
latency of C assimilation for each guild. Latency is defined operationally and it is proportional to
the time delay between peak mineralization and the time of ^{13}C assimilation into DNA for each C
source. D) The dynamic growth response of each guild as measured by increase in abundance in
response to substrate addition. Red and grey lines indicate the statistically significant and non-
significant linear relationship between factors respectively with shading representing standard
error. Pearson's r and p-values for these relationships are listed above each plot. All p-values
were corrected for multiple comparisons with Benjamini-Hochberg procedure ($n = 4$). Error bars
for points indicate ± standard error across OTUs within each guild. E) The PCA differentiating
guild structure on the basis of *rrn* copy number (ln(rrn)), C source bioavailability, latency of C
assimilation, dynamic growth response (Max l2FC), and number of C sources (C sources).
Circles represent guilds with numbers indicating identity and arrows representing loadings of
each variable onto the principal component axes (PC1 and PC2).
821
822

823
824
Figure 6: Growth and [13]C-labeling dynamics of the most abundant OTUs from guilds at the
extremes indicated on the PCA (Fig. 5E). A) The [13]C-labeling pattern of the exemplar OTUs.
Times when [13]C-labeling was detected for each substrate are indicated by circles with colors
indicating the different OTUs defined in panel B. Black rectangular borders indicate the
timepoints in in which microcosms were harvested for DNA-SIP (*i.e.,* where [13]C-labeling can be
detected). Red rectangles indicate the peak C mineralization timepoints for each C source (Fig.
2). B) The growth dynamics of the exemplar OTUs. Abundances are normalized by *rrn* copy
number and DNA yield from the microcosms. OTU.7163 is a member of guild 19 and proposed
to exemplify a ruderal strategist. OTU. 22 is a member of guild 14 and proposed to exemplify a
competitor. OTU.197 is a member of guild 2 and proposed to exemplify a stress tolerator.