A bispecific monomeric nanobody induces SARS-COV-2 spike trimer dimers

Leo Hanke[1,†], Hrishikesh Das[2,†], Daniel J Sheward[1], Laura Perez Vidakovics[1], Egon Urgard[1],

Ainhoa Moliner-Morro[1], Vivien Karl[1], Alec Pankow[1], Changil Kim[1], Natalie L Smith[1,] Gabriel

K Pedersen[3], Jonathan M Coquet[1], B Martin Hällberg[2,4,*], Ben Murrell[1,‡,*], Gerald M

McInerney[1, ‡,*]

[1] Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, Stockholm,

Sweden.

[2] Department of Cell and Molecular Biology, Karolinska Institutet, Stockholm, Sweden.

[3] Center for Vaccine Research, Statens Serum Institut, Copenhagen, Denmark

[4] Karolinska Institutet VR-RÅC, Centre for Structural Systems Biology, Notkestraße 85, 22607,

Hamburg, Germany

[†] Contributed equally
[‡] Contributed equally

* Correspondence to: martin.hallberg@ki.se; benjamin.murrell@ki.se; gerald.mcinerney@ki.se

Abstract

Antibodies binding to the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) spike have therapeutic promise, but emerging variants show the potential for virus escape. Thus, there is a need for therapeutic molecules with distinct and novel neutralization mechanisms. Here we isolated a nanobody that potently neutralizes SARS-CoV-2, including the B.1.351 variant, and cross-neutralizes SARS-CoV. We demonstrate the therapeutic potential of the nanobody in a human ACE2 transgenic mouse model. Using biochemistry and electron cryomicroscopy we show that this nanobody simultaneously interacts with two RBDs from different spike trimers, rapidly inducing the formation of spike trimer-dimers. This naturally elicited bispecific monomeric nanobody establishes a novel strategy for potent immobilization of viral antigens.

Introduction

The SARS-CoV-2 pandemic and the vast societal and economic consequences of the connected lockdowns have had devastating consequences for the world's populations. Despite rapid progress in vaccine development, some emerging virus variants appear to be less effectively neutralized by antibodies elicited by the first-generation vaccines [1] and the emergence of further variants is expected. To prepare for different scenarios and to have emergency alternatives at hand, novel ways to inhibit and neutralize the virus and variants of concern are needed. As therapeutic agents in the early phases of SARS-CoV-2 infection, camelid-derived single-domain antibody fragments are promising candidates. These so-called 'nanobodies' not only bind their antigen with very high specificity and affinity, but they can also be easily produced to large quantities in *E.coli* [2] and show favourable biochemical characteristics, including good thermal

stability. Importantly, they can easily be combined to form homo- and heterodimers or other

multimers to increase potency and target multiple epitopes simultaneously [3–5]. This strategy

allows the neutralization of multiple known or unknown virus variants and reduces the risk for

viral escape [5].

The attachment of SARS-CoV-2 to the host cell receptor ACE2 and subsequent membrane

fusion is mediated by the viral spike (S) protein [6]. In the prefusion conformation, the spike is a

large homotrimer. Each S monomer consists of two subunits: S1 and S2. The S1 subunit contains

the receptor binding domain (RBD) and S2 contains the fusion peptide, as well as the

transmembrane domain. The RBD is responsible for the initial attachment to ACE2 and is found

in two distinct conformational states: The receptor-accessible 'up' conformation and the receptor

inaccessible 'down' conformation [6,7]. Receptor binding induces the rather unstable 3-up state,

which leads to shedding of S1 and refolding of S2 thereby presenting the fusion peptide to the

opposing membrane [6,8,9]. SARS-CoV-2 is easily neutralized by high-affinity binding antibodies

and other spike-targeting molecules, including nanobodies [10–13]. Many of them bind to, or close

to, the receptor binding motif of the RBD and neutralize by directly hindering virus attachment

[14,15]. Other mechanisms include locking the RBDs in the down conformation or preventing

productive fusion [5,16]. Still, the emergence of new virus variants that evade antibody-mediated

neutralization mandates the identification of molecules with a more diverse set of binding

epitopes and neutralization mechanisms allowing for neutralization of a variety of virus variants.

Here, we generated a nanobody that has two non-overlapping interaction sites on the SARS-

CoV-2 RBD, each targeting a different epitope, which induces the formation of dimers of spike

trimers and suppresses SARS-CoV-2 *in vivo*.

Results

Isolation and characterization of a potent RBD-specific nanobody.

To identify nanobodies with unique neutralization potential, we immunized one alpaca (*Funny*) with SARS-CoV-2 spike protein and recombinant RBD. We isolated RNA from peripheral blood mononuclear cells and created phagemid libraries. Phage display screens with immobilized RBD yielded one promising nanobody, called Fu2. Since multimerization of nanobodies can substantially improve virus neutralization potency [3,17], we created three different dimers (illustrated in Fig. 1A): One Fu2-Fc fusion, expressed in mammalian cells, the second a chemically linked Fu2 homodimer, and thirdly a heterodimer of Fu2 with a previously described RBD-specific neutralizing nanobody Ty1 [14]. Chemically linked constructs were generated using a combination of sortase A labelling and Cu-free click-chemistry as described in detail previously [3].

Using pseudotyped lentivirus (PSV) neutralization assay (Fig. 1B), we found that monomeric Fu2 was approximately 10-times more potent than Ty1, with an IC_{50} in the range of 106 ng/ml (7 nM). The Fu2-Fc fusion only slightly increased the neutralization potency of this nanobody to 61 ng/ml (0.75 nM), and a chemical homo-dimerization to an IC_{50} of 26 ng/ml (0.8 nM). Thus, the observed enhancement in potency is attributable to dimerization, rather than the Fc-domain itself. Interestingly, the Fu2-Ty1 heterodimer was extremely potent, with an IC_{50} approximately 6-fold lower than that of the Fu2 homodimer, in the range of 4 ng/ml (140 pM). Fu2 directly prevented spike binding to ACE2 as confirmed by flow cytometry of HEK293T-hACE2 cells stained with recombinant, fluorescently labelled spike (+/- Fu2) (Fig. 1C). Analysis of binding kinetics by surface plasmon resonance (SPR) revealed that Fu2 bound the RBD with sub-nanomolar affinity (Fig. 1D). The 1:1 binding model fit the sensorgrams well at the used concentration range, indicating a single dominant interaction. Analysis of next-generation sequencing data for other

members of the Fu2 lineage yielded several Fu2-variants with similar properties (Supplementary Fig. 1A-E).

To test the integrity of the recombinant spike protein after the addition of Fu2, we analysed the size-exclusion chromatography elution profiles in combination with different nanobody constructs (Fig. 1E). The SARS-CoV-2 spike trimer eluted in a single peak and the addition of Ty1 [14] did not result in a shift in the position of this peak, which is expected given the small relative size addition of the 14 kDa nanobody to the large (>400 kDa) spike trimer. In contrast, the addition of Fu2 resulted in a substantial shift of the peak, suggesting the formation of a larger complex, likely a dimer of the spike trimer. Next, we combined the spike with the Fu2-Ty1 heterodimer. This also resulted in earlier elution, and besides a possible spike dimer, this construct gave rise to an even earlier elution peak suggesting the induction of higher-order spike multimers.

Structural basis for Fu2 induction of spike trimer dimers.

To better understand neutralization and the aggregation of spike by Fu2, we used electron cryomicroscopy (cryo-EM) to determine the structure of the Fu2-spike complex. The structure revealed a remarkable head-to-head dimerization of the trimeric spike, bound by six molecules of Fu2 (Fig. 2). This distinct structural state appeared as prominent 2D classes in our analyses (Fig. 2A). Binding of Fu2 to the RBD is unique as it can simultaneously interact with two different RBDs from different spike trimers in the up conformation (Figs. 2B and 2C). To our knowledge, Fu2 is the only nanobody or spike-binding molecule developed against the SARS-CoV-2 spike to have this property. Our high resolution (2.9Å; 0.143 FSC) localized reconstruction of the RBD-Fu2-RBD-Fu2 interface helped us elucidate the molecular details of this interaction. Each RBD in the complex provides two binding interfaces for two Fu2 molecules, which we call

'interface-major' and 'interface-minor' according to surface area covered (Fig. 2D and

Supplementary Fig 2A and Supplementary 2B). The interface-major and interface-minor have a

buried surface area of ~740 Å^2 and ~495 Å^2, respectively. Parts of the Fu2 framework regions

drive the dimerization by shape-complementarity and coupled solvent exclusion in the two

interfaces. In addition, the interface-major consists of 15 hydrogen bonds and one salt bridge

between Fu2 D117 and RBD K378, while the interface-minor comprises five hydrogen bonds.

Furthermore, a region around the Fu2 β-turn A40-K43 plays a key role in dimerization through

interactions with two proximal RBDs (Supplementary Figs. 2C and 2D). Specifically, one of the

two proximal RBDs' T500-Y505 region packs against the Fu2 A40-K43 β-turn and the RBD

Y505 is in addition bound by Fu2 Q39. Meanwhile, the other proximal RBD is bound through a

bidentate hydrogen bond between Fu2 E44 and the backbone amides of RBD V503-G504 (Fig.

2E and Supplementary Figs. 2C and 2D). Interestingly, the overall structure of Fu2-bound spike

had no significant deviations from previous structures (Supplementary Fig. 3) and we did not find

any contact between any two Fu2 molecules or between spike trimers suggesting that the

observed spike trimer dimerization is solely driven RBD-Fu2 interactions.

Fu2 blocks the RBD from binding ACE2.

Structural alignment of an ACE2-RBD structure to the Fu2-spike structure (single RBD) shows

that binding of ACE2 would be hindered by Fu2 bound to any of the two interfaces (Fig. 2F).

Fu2 bound to the interface-major would clash with ACE2 residues ~303-331 and the N-glycan at

position 322, while Fu2 bound to interface-minor would mask the interaction surface of ACE2,

making it sterically impossible for ACE2 to access the RBD. A comparison of the Fu2 and Ty1

binding site revealed that interface-minor is partially shared between Ty1 and Fu2 (Supplementary

Figs. 2E and 2F), suggesting that in context of the Fu2-Ty1 heterodimer, Ty1 is displaced upon

dimerization of the spike trimer. Ty1 can then potentially bind additional spike molecules,

explaining the elution profile showing both dimers and higher order spike complexes (Fig. 1E).

In the complex, Fu2's long CDR3 (G99 to Y120) plays a significant role in RBD binding. Here,

the Fu2 S107-R111 region adopts a β-strand conformation that binds the RBD T376-Y380 β-

strand to give rise to an energetically favourable inter-chain extension of the RBD anti-parallel

β-sheet (Figs. 2G and 2H). Interestingly, the Fu2 CDR1 (G26-Y32) and CDR2 (T52-S57) do not

participate in RBD binding, and none of the CDRs is directly involved in the formation of the

spike dimer.

Fu2 stabilizes the RBD in the 'up' conformation.

Cryo-EM analyses, both of the purified spike or of intact virions has revealed that unliganded

spike is mostly present in all-down (~60%) or 1-up (~40%) conformation [6]. Our Fu2-spike

structure shows the presence of two trimeric spike molecules, both in the 3-up conformation.

This arrangement allows Fu2 to access all six possible binding sites on RBDs without steric

conflict. Alignment of RBD-Fu2 to RBD-down in a 2-up spike structure confirms that binding of

Fu2 to RBD in the down conformation is not possible (Supplementary Fig. 2G). Fu2 bound to

either interaction interface on an RBD in the down conformation would clash with neighbouring

RBDs. Likewise, an alignment of the 2-up spike structure to the Fu2-trimer-dimer shows that

RBD-down would clash with the neighbouring Fu2 molecule (Supplementary Fig. 2H). In

conclusion, for Fu2 binding and for trimer-dimer assembly, all RBDs must be in the up

conformation.

Fu2-Ty1 provides therapeutic relief in a mouse model of SARS-CoV-2 infection

An important requirement for the development of antivirals is neutralization potential *in vivo*. To address this, we used K18-hACE2 transgenic mice that express human ACE2 as a model for SARS-CoV-2 infection. To determine if neutralizing nanobodies could prevent infection, or reduce disease severity when administered in early infection, we performed a prophylactic and a therapeutic experiment *in vivo* (Fig. 3). Mice in the prophylactic treatment group (N=4) were injected intraperitoneally (i.p.) with 370 µg of the Fu2-Ty1 heterodimer (11-20 mg/kg), 6 h before intranasal challenge with 10^3 plaque forming units (PFU) of SARS-CoV-2. The therapeutic group (N=4) was injected i.p. with 280 µg (9-14 mg/kg) of Fu2-Ty1 daily, for four days, starting 24 h post infection. Viability and weight change were monitored daily for 12 days post-challenge. Weight loss for infected mice was first observed on day 4. Prophylactic treatment did not greatly alter weight loss or disease severity. In contrast, weight loss was delayed in mice receiving Fu2-Ty1 heterodimers on a therapeutic regimen (10% weight loss endpoint, $p<0.05$, Cox Proportional Hazard model). One of four mice treated post infection fully recovered from infection with an otherwise-lethal challenge dose, with weight rebounding to pre-infection levels. From day 6 oropharyngeal samples, qPCR of SARS-CoV-2 E-gene showed significantly reduced genomic and subgenomic RNA in the therapeutic group compared to the untreated (Fig. 3C). The data shows that the Fu2-Ty1 dimer suppresses SARS-CoV-2 *in vivo*.

Fu2 cross-neutralizes SARS-CoV, and clinical isolates of SARS-CoV-2 including the B.1.351 variant of concern

To test the breadth of neutralization, we evaluated cross-neutralization of SARS-CoV PSV (Fig. 4A). While Ty1 did not neutralize SARS-CoV, Fu2 and dimeric Fu2 constructs were neutralizing, but less so than against SARS-CoV-2 PSV. Interestingly, the neutralization curves displayed a much flatter slope as compared to those against SARS-CoV-2 PSV. It required 6.1

μg /ml of monomeric Fu2 to reach 50% neutralization in this assay, while 570 ng /ml of Fu2-Fc

and 57 ng/ml of the Fu2-dimer was sufficient for 50% neutralization.

To understand the molecular mechanism of cross-neutralization of SARS-CoV we compared the

RBD sequences and performed structural alignments (Figs. 4B, 4C and Supplementary Fig. 4A).

Interestingly, the interface-major residues are conserved, while interface-minor has significant

differences, suggesting that the divergent interface-minor of SARS-CoV RBD would not permit

Fu2 binding. From these analyses we conclude that Fu2 binding would be unlikely to form

SARS-CoV trimer dimers.

To demonstrate neutralization of replication competent SARS-CoV-2 we performed a plaque

reduction neutralization test (PRNT) using infectious SARS-CoV-2, isolated from the first

Swedish COVID-19 patient [18] (Fig. 4D). Similar to neutralization of PSV, monomeric Fu2

neutralized SARS-CoV-2 much better than Ty1, a dimeric construct of Fu2 provided a 2.8-fold

improvement over the monomeric version of Fu2, while the heterodimer of Ty1 and Fu2 showed

a 50-fold improvement compared to monomeric Fu2 in this assay.

A PRNT using the B.1.351 (501Y.V2) [1] variant showed that Fu2 very potently neutralized this

novel variant (Fig. 4E) and that the Fu2 dimer and Fu2-Fc again improved neutralization

efficiency. The IC_{50} values were even slightly better than when using the Swedish isolate, but the

difference is within the intraassay variability. Ty1 failed to neutralize this virus, and the Fu2-Ty1

heterodimer did not neutralize better than Fu2 alone. The variant B.1.351 harbours four amino

acid substitutions in the RBD at position K417N, N439K, E484K and N501Y. Only N501 and

E484 are located in proximity of the Fu2 interaction interface. *In silico* analyses with the PISA

server (25) confirmed that the two interfaces do not change in these mutants. Both buried surface

areas of the interfaces as well as the number of possible hydrogen bonds remain unchanged (Fig.

4F). These results and analysis combined suggests that Fu2 likely also neutralizes the recently

emerged variants P1 (K417T, E484K, N501Y) and B.1.1.7 (N501Y).

Discussion

Nanobodies represent a valuable alternative to monoclonal antibodies for virus neutralization

[19,20]. Their ease of production, thermal stability, and robust biochemical behaviour, combined

with straight-forward functionalization and multimerization possibilities, make them suitable

candidates for therapeutic applications.

The newly emerged SARS-CoV-2 B.1.351 variant shows nine amino acid changes in the spike

protein. While biological properties of this variant remain to be determined, two amino acid

substitutions in the receptor binding motif are of potential concern for vaccine efficacy,

neutralization efficiency of existing monoclonal antibody therapies, and changed dynamics of

virus spread [21–23]. The novel nanobody described here, Fu2, can effectively neutralize the

B.1.351 variant and has therefore a strong and unique potential to form the basis for development

of therapeutic interventions against this emerging variant of concern.

Fu2 is also capable of neutralizing SARS-CoV, highlighting its specificity for a more conserved

epitope. However, the potency was significantly reduced for the monomeric nanobody, likely

because of decreased affinity to the RBD of this virus. That the dimeric constructs significantly

increased potency further supports this hypothesis. A similar observation was made for VHH-72

originally developed against SARS-CoV, but with some neutralization potential in dimeric form

against SARS-CoV-2 [24].

Nanobodies typically have short serum half-lives, ranging from 1-3 hours [25], compared to 6-8

days of IgG1 [26]. We show that a high dose of nanobody heterodimer is not suitable as a molecule

for passive immunization as it did not diminish the severity of disease of hACE2 expressing

mice. In contrast, if administered daily for four days post challenge, the heterodimer significantly

delayed disease onset and severity, suggesting that nanobodies and dimeric nanobody constructs

have the potential for SARS-CoV-2 antiviral therapy. Half-life extending strategies such as

fusion to serum albumin moieties could be used to extend half-life [27] and reduce the number of

infusions.

The RBD-specific nanobody Fu2 displays a particularly unique mode of virus neutralization by

rapidly inducing stable spike trimer dimers that are incapable of binding to ACE2. In contrast to

spike trimer dimers reported for the full-length spike that are based on S1-S1 interactions placing

the trimers 'side-by-side' [28], Fu2 induced a dimer in a 'head-to-head' orientation with the C-

termini at the distal ends. Targeting two epitopes increases the binding surface area, but, more

importantly, when locked in such a conformation the function of each glycoprotein is sterically

blocked both by the nanobody and by the other glycoprotein in such a way that the whole

complex must dissociate for the glycoprotein to regain its function.

The interaction between Fu2 and the RBD is highly unusual, and for a monomer to induce

antigen dimerization in this manner, two copies of an antigen must be bound at different epitopes

with precisely the right spatial arrangement. This novel mode of antigen binding may be

uncommon in naturally elicited antibody repertoires but could be achievable by structure-based

antibody design and might present a general strategy for potent immobilization of viral

glycoproteins and other antigens. It remains to be seen whether 'designer' bispecific monomer

nanobodies can push the potency envelope beyond what is currently attainable.

Fig. 1. An RBD-specific nanobody neutralizes SARS-CoV-2. (**A**) Overview of different nanobody constructs used in this study: (1) nanobody monomer, (2) nanobody Fc-fusion, (3) chemically linked nanobody homodimer, (4) chemically linked nanobody heterodimer. (**B**) A SARS-CoV-2 spike pseudotyped lentivirus (PSV) was incubated with a dilution series of the indicated nanobodies at 37°C for one hour before infecting HEK293T-hACE2 cells. Neutralization (in %) compared to untreated PSV is shown. Data from at least two replicates is shown and the error bars represent the standard deviation. (**C**) Recombinantly expressed, prefusion stabilized, and fluorescently labeled SARS-CoV-2 spike protein was incubated with a control nanobody specific for IAV NP [29], or Fu2 and used to stain ACE2 expressing HEK293T cells. Cells were analyzed by flow cytometry and a representative histogram is shown. (**D**) Binding kinetics of Fu2 to the RBD were measured by surface plasmon resonance (SPR). Sensorgram is color-coded based on concentration. The fit is based on the 1:1 Langmuir model and is shown in dark grey solid lines. (**E**) Recombinantly expressed, prefusion stabilized SARS-CoV-2 spike protein was run alone or preincubated with the indicated nanobody constructs on a Superose 6 size-exclusion column. Elution profiles (A280) are shown.

Fig. 2. Cryo-EM provides structural evidence of Fu2-mediated dimerization of trimeric spike in '3-up' conformation (**A**) Representative 2D class averages. (**B**) Cryo-EM map (two different side views) and (**C**) Atomic model of 'dimeric' spike in complex with Fu2; and two opposite protomers with two Fu2 molecules. Both spike trimers in the complex are present in '3-up' conformation (six Fu2 molecules are numbered). (**D**) Close-up view of the RBD-Fu2 interaction interfaces showing the interface-major and interface-minor. (**E**) A β-hairpin (gray sticks) from the Fu2 framework region (residues 39-45) contacts two RBDs simultaneously in the Fu2-mediated spike trimer-dimer. (**F**) Modelling of simultaneous spike-Fu2-ACE2 binding shows that Fu2 blocks the binding of ACE2 to the RBD from two directions. Clashes marked with dashed circles. (**G**) Fu2 binds the RBD through an extended inter-chain β-sheet interaction mediated by the CDR3 region (green). CDR1 (gray) and CDR2 (blue) does not take part in RBD binding (**H**) Closeup of a part of the CDR3-RBD interface displaying the anti-parallel β-strand hydrogen bond pairing between the Fu2 CDR3 region and the RBD (light orange).

Fig. 3. Therapeutic nanobody reduces disease severity in a SARS-CoV-2 challenge model. (**A**) Timeline of the challenge experiment. K18-hACE2 transgenic mice were challenged with 1000 plaque forming units (PFU) of SARS-CoV-2 and received prophylactic (blue) or therapeutic (red) Fu2-Ty1 at the indicated time points. (**B**) Weight of mice during the challenge experiment. The mean weight of each mouse of day 0 to day 2 served as baseline and the weight loss relative this baseline is shown. Uninfected mice are shown in grey, untreated infected mice in black, prophylactic treatment group in blue and therapeutic group in red. (**C**) Analysis of oropharyngeal samples from mice at day 6 in infected groups. Ratios of E-gene to ABL1 is shown for both genomic and subgenomic RNA (** p < 0.01, Mann–Whitney U test, one-tailed).

Fig. 4. Fu2 cross-neutralizes SARS-CoV and a B.1.351 variant of concern clinical isolate
(**A**) SARS-CoV S pseudotyped lentivirus (PSV) was incubated with the indicated constructs before infecting HEK293T-ACE2 cells, and neutralization is shown compared to untreated PSV. (**B** and **C**) Structural basis of SARS-CoV cross-neutralization by Fu2. Structural alignment of SARS-CoV-2 Spike (RBD bound Fu2) with SARS-CoV RBD (PDB:6ACD). Residues of interface-major and interface-minor shown as sticks, divergent residues are labelled. (**D and E**) PRNTs with clinical isolates of SARS-CoV-2. 100 plaque forming units (PFU) of SARS-CoV-2 [18] or B.1.351 (501Y.V2) [1] were incubated with dilution series of the indicated nanobody constructs for 1 h at 37°C before infecting monolayers of Vero E6 cells. Plaques were quantified 72 h post infection. Neutralization representing the reduction of plaques relative to the control wells is shown. Error bars represent standard deviation (SD) across replicate experiments. (**F**) Assessment of Fu2-RBD interaction in RBD variants. *In silico* mutational analyses of RBD residues and probable impact on Fu2 binding. Fu2 is shown in red and variant residues are indicated.

Materials and Methods

Cells and viruses.

HEK293T cell (ATCC-CRL-3216) and Vero E6 cells (ATCC-CRL-1586) were maintained in DMEM (Gibco) supplemented with 10% fetal calf serum and 1% penicillin-streptomycin in a humidified incubator with 5% CO_2 at 37°C. The HEK293T cell line was further engineered to overexpress human ACE2 by lentiviral transduction [14]. All cell lines used for the experiments were negative for mycoplasma as determined by PCR. Infectious SARS-CoV-2 [18] and the B.1.351 [1] isolate were propagated in Vero E6 cells and titrated by plaque assay.

Proteins and probes.

The plasmid for the expression of the SARS-CoV-2 prefusion-stabilized spike (2P) ectodomain was a kind gift from the McLellan lab [6]. The plasmid was used for transient transfection of FreeStyle 293F cells using the FreeStyle MAX reagent (Thermo Fisher Scientific). The spike trimer was purified from filtered supernatant on Streptactin XT resin (IBA Lifesciences) or Ni-NTA resin and purified by size-exclusion chromatography on a Superdex 200 (Cytiva). For cryo-EM a similar construct with 6 proline substitutions (S6-P) was used [30], and the protein was purified on His-Pur Ni-NTA resin (Thermo Fisher Scientific) followed by size-exclusion chromatography.

The RBD domain was cloned upstream of a sortase A motif (LPETG) and a 6xHIS tag. The plasmid was used for transient transfection of FreeStyle 293F cells as described above. The protein was purified from filtered supernatant on His-Pur Ni-NTA resin followed by size-exclusion chromatography on a Superdex 200.

Nanobodies were cloned in the pHEN plasmid with a C-terminal sortase motif (LPETG) and a

6xHIS tag. BL21 *E. coli* were transformed with this plasmid, and expression was induced with 1

mM IPTG at OD600 = 0.6 and cells were grown overnight at 30 °C. Nanobodies were retrieved

from the periplasm by osmotic shock and purified on Ni-NTA resin and size-exclusion

chromatography.

Sortase A 5M was produced as described before in BL21 *E. coli* and purified by Ni-NTA and

size exclusion chromatography [3].

Fluorescent spike ectodomain was generated by first attaching dibenzocyclooctyine-*N*-

hydroxysuccinimidyl ester (DBCO-NHS) to the spike trimer in a 3:1 molar ratio, before

attaching AbberiorStar-635P-azide by click-chemistry. The final product was purified from

unreacted DBCO and fluorophore on a PD-10 desalting column.

The biotinylated RBD was generated using sortase A and amine-PEG3-biotin as a nucleophile.

The reaction was performed with 50 µM RBD, 5 µM sortase A 5M, and 8 mM amine-PEG3-

biotin in 50 mM Tris pH 7.5, 150 mM NaCl, for 6 hours at 4 °C. Sortase A and unreacted RBD

was removed on Ni-NTA resin and excess nucleophile was removed by two consecutive

purifications on PD-10 desalting columns.

Dimer generation using click chemistry.

To generate homo- and hetero-dimers, the different nanobodies were first site-specifically

functionalized on the C-terminus using sortase A with either an azide or a dibenzocyclooctine

(DBCO) as described in detail here [3]. In brief, for functionalization with DBCO, 70 µM of

nanobody was incubated with 5 µM sortase A, 8 mM DBCO-amine (Sigma-Aldrich) in 150 mM

NaCl, 50 mM Tris pH 7.5 and 10 mM $CaCl_2$ for 3 h at 25 °C. To functionalize the nanobody

with an azide, 70 µM of nanobody, was incubated with 5 µM sortase A 5M, 10 mM 3-azido-1-

propanamine (Sigma-Aldrich #762016) in 150 mM NaCl, 50 mM Tris pH 7.5 and 10 mM $CaCl_2$

for 3 h at 25 °C. In both reactions, unreacted nanobody, sortase A and excess nucleophile were

removed using Ni-NTA resin and PD-10 columns or size-exclusion chromatography. The click

reaction was initiated by mixing azide and DBCO labeled nanobody in a 1:1 molar ratio for 48 h

at 4 °C. Dimers were purified from unreacted monomeric nanobody by size-exclusion

chromatography on a Superdex S200 16/600 column (Cytiva).

Alpaca immunization, library generation and nanobody isolation.

One adult female alpaca (*Funny*) at PreClinics, Germany, was immunized four times in a 60-day

immunization schedule. Each immunization consisted of two injections. For the first

immunization, 200 µg of prefusion stabilized spike and 200 µg of S1+S2 domain (Sino

Biologicals) was used. Remaining immunizations each consisted of one injection with 200 µg

RBD and one injection with 200 µg prefusion stabilized spike, both produced in Freestyle 293F

cells as described above. Serum from the final bleed at day 60 neutralized SARS-CoV-2

pseudoviruses with an ID50 >20 000. The animal study protocol was approved by the PreClinics

animal welfare officer commissioner and registered under the registration No. 33.19-42502-05-

17A210 at the Lower Saxony State Office for Consumer Protection and Food Safety—LAVES

and is compliant with the Directive 2010/63/EU on animal welfare.

The nanobody phage library was generated as described [14]. Phage display was performed on

RBD immobilized on magnetic beads.

Neutralization assays.

Pseudotyped virus neutralization assay: Pseudotyped viruses were generated by co-transfection

of HEK293T cells with plasmids encoding firefly luciferase, a lentiviral packaging plasmid

(Addgene cat#8455), and a plasmid encoding either the SARS-CoV or SARS-CoV-2 spike

protein, both harboring a C-terminal truncation of 18 amino acids. Media was changed 12-16 h

post-transfection, and pseudotyped viruses were harvested at 48 and 72 hours, filtered through a

0.45 μm filter and stored at -80 °C until use. Pseudotyped viruses sufficient to generate 100,000

relative light units (RLU) were incubated with serial dilutions of nanobody for 60 min at 37 °C.

15,000 HEK293T-ACE2 cells were then added to each well, and the plates were incubated for 48

h at 37 °C. Luminescence was measured using Bright-Glo (Promega) on a GM-2000

luminometer (Promega) with an integration time of 0.3 s.

Plaque reduction neutralization test: One day before the experiment, 100,000 Vero E6 cells were

seeded per well of a 24-well plate. A 3-fold serial dilution of the nanobody constructs starting at

10 μg/ml were incubated with 100 plaque-forming units (PFU) of the first Swedish clinical

isolate of SARS-CoV-2 or variant B.1.351 for 1 h at 37°C. The nanobody-virus mixture was then

added to the cells and incubated at 37 °C for 1 h. Cells were washed with PBS, and an overlay of

1% CMC/DMEM supplemented with 2% FBS was pipetted into the wells. Three days post-

infection, cells were fixed in 10% formaldehyde, followed by staining with crystal violet.

Plaques were counted and inhibition determined by comparing to control wells on the same

plate.

Flow cytometry

HEK293T-hACE2 cells were trypsinized and fixed in 4% formaldehyde in PBS for 20 min. Cells

were stained with spike- AbberiorStar-635P not premixed or premixed with Fu2 or control

nanobody. Fluorescence was quantified using a BD FACSCelesta and the FlowJo software

package.

Gel shift assays

For gel shift assay, 80 µg of prefusion stabilized spike protein was incubated with or without nanobody for 15 minutes and was run on a Superose 6 increase 10/300 GL (Cytiva), and absorbance at 280 nm was detected. Sample and injection volumes were kept identical between runs to obtain comparable elution profiles.

Next generation sequencing

Next generation sequencing on pre- and post-enrichment nanobody libraries was performed as described in Hanke et al. 2020 [14] on an Illumina MiSeq.

Surface plasmon resonance

Binding kinetics were determined by surface plasmon resonance using a Biacore 2000. All experiments were performed at 25 °C in a running buffer of 10 mM HEPES, 150 mM NaCl, pH 7.4, and 0.005% Tween-20 (v/v). Site-specifically biotinylated RBD was immobilized on streptavidin sensor chips (Series S sensor Chip SA, GE Healthcare) to a level of ~200 resonance units (RU). A 2-fold dilution series of the nanobodies was injected at a flow rate of 30 µl/min (association 180 s, dissociation 900 s), and the immobilized RBD was regenerated using 0.1 M glycine-HCL buffer pH 2 for 2x 10 seconds. Data were analyzed using BIAevaluation Software and fitted using the 1:1 Langmuir model with mass transfer.

SARS-CoV-2 challenge experiments

K18-hACE2 transgenic mice were purchased from Jackson laboratories and maintained as a hemizygous line. Experiments were conducted in BSL3 facilities at the Comparative Medicine

department (KM-F) at Karolinska Institutet. Ethics for studies of virus infection and therapeutic

intervention were obtained from Stockholms Jordbruksverket (10513-2020). Mice were

prophylactically or therapeutically administered nanobodies as described in the main text and

challenged intranasally with 1000 PFU SARS-CoV-2 (Swedish isolate) in 40 µl PBS following

isoflurane sedation. Oropharyngeal sampling was performed at the indicated timepoints under

light anesthesia with isoflurane. Weight and general body condition was monitored daily until

weight drop started, whereupon mice were monitored twice daily. During the experiment, weight

loss, changes in general health, breathing, body movement and posture, piloerection and eye

health were monitored. Mice were typically sacrificed when they achieved 20% weight loss.

However, some mice were sacrificed before losing 20% in body weight, when movement was

greatly impaired and/or they experienced difficulty breathing that was considered to reach a

severity level of 0.5 on Karolinska Institutet's veterinary plan for monitoring animal health.

RNA Extraction and RT-qPCR for SARS-CoV-2 detection.

Viral RNA was isolated from buccal swabs collected on dpi 6 and stored in 500 µl of TRIzol™

Reagent (Invitrogen). Total RNA extractions from buccal swab samples were performed using

an adapted TRIzol™ manufacturers protocol with a 45 min precipitation step at -20° C. RNA

pellets were resuspended in 20 µl of RNase-free water.

RT-PCR reactions were performed using 4 µl of resuspended RNA in a 20 µl reaction volume

using the Superscript III one step RT-qPCR system with Platinum Taq Polymerase (Invitrogen)

with 400 nM concentrations of each primer and 200 nM of probe. Primers and probes for the

CoV-E gene target were as previously described [31]. Primers and probes for the ABL1 target were

adapted from Ishige et al. [32] to enable detection of the murine homolog: (ABL1_ENF1003_deg:

5'-TGGAGATAACACTCTCAGCATKACTAAAGGT-3', ABL1_ENR1063: 5'-

GATGTAGTTGCTTGGGACCCA-3', ABL1_ENPr1043_deg: 5'-HEX-

CCATTTTTSGTTTGGGCTTCACACCATT-BHQ1-3'). The CoV-E and ABL1 TaqMan assays

were run in multiplex. Detection of the subgenomic CoV-E target was adapted from Wölfel et al.

[33], using a leader/E gene junction specific forward primer: (sgEjunc_SARSCoV2_F: 5'-

CGATCTCTTGTAGATCTGTTCTCTAAACG-3'). All oligonucleotides were synthesized by

Eurofins Genomics.

Thermal cycling conditions for all assays consisted of RT at 55 °C for 10m, denaturation at 95

°C for 3m, and 45 cycles of 95C, 15s and 58C, 30s. Reactions were carried out using a CFX96

Connect Real-Time PCR Detection System (Bio-Rad) following manufacturer instructions. To

generate standard curves, a synthetic DNA template gBlock (Integrated DNA Technologies) was

transcribed using the mMessage mMachine™ T7 Transcription Kit (Invitrogen) and serially

diluted. To reduce sampling-related variability, SARS-CoV-2 RNA copies were normalized by

ABL1 copies, and this ratio was used for comparisons. ABL1 copies were not significantly

different between groups.

Cryo-EM sample preparation and imaging

Spike trimer (1.8mg/ml) S6-P and Fu2 (2.1mg/ml) were mixed in a 1:6 molar ratio followed by

incubation on ice for 10 minutes. Prior to cryo-EM grid preparation, grids were glow-discharged

with 25 mA for 2 minutes using an EMS 100X (Electron Microscopy Sciences) glow-discharge

unit. CryoMatrix® holey grids with amorphous alloy film (R 2/1 geometry; Zhenjiang Lehua

Technology Co., Ltd) were used. 3 µl aliquots of sample solutions were applied to the grids and

the grids with sample were then vitrified in a Vitrobot Mk IV (Thermo Fisher Scientific) at 4°C

and 100% humidity (blot 10 s, blot force 3, 595 filter paper (Ted Pella Inc.)). Cryo-EM data

collection was performed with EPU (Thermo Fisher Scientific) using a Krios G3i transmission-

electron microscope (Thermo Fisher Scientific) operated at 300 kV in the Karolinska Institutet's

3D-EM facility (https://ki.se/cmb/3d-em). Images were acquired in 165 kx nanoprobe EFTEM

SA mode with a slit width of 10 eV using a K3 Bioquantum during 1.5 s during which 60 movie

frames were collected with a flux of 0.81 e-/Å2 per frame (Supplementary Table 1). Motion

correction, CTF-estimation, Fourier binning (to 1.01 Å/px), picking and extraction in 600-pixel

boxes (size threshold 400 Å, distance threshold 100 Å, using the pretrained

BoxNet2Mask_20180918 model) were performed on the fly using Warp [34]. Due to preferential

orientation, we collected a part of the dataset with 25-degree stage tilt.

A total of 14,081 micrographs were selected based on an estimated resolution cut-off of 4 Å and

defocus below 2 microns as estimated by Warp. Furthermore, Warp picked 1,035,962 particles

(dimer of trimeric spike with Fu2) based on above-mentioned criteria. Extracted particles were

imported into cryoSPARC v3.0.1 [35] for 2D classification, 3D classification and non-uniform 3D

refinement. After 2D classification, clean classes with high-resolution features (and with

characteristic trimeric spike views) were retained and used to generate ab-initio 3D

reconstructions. In total 277,372 particles were retained for refinement steps and these particles

were processed with C1 symmetry (image processing scheme Supplementary Fig. 5).

These particles were further processed using heterogeneous refinement that resulted in a

reconstruction with high-resolution structural features in the core of the spike. One round of

homogeneous refinement was followed by non-uniform refinement. All final reconstructions

were analysed using 3D-FSC [36] and there were moderate anisotropy in the full map

reconstructions while the localized reconstruction displayed no significant anisotropy

(Supplementary Fig. 6A and 6B). All CTF refinements were per-particle CTF refinements

interspersed with global aberration correction (beamtilt, trefoil, tetrafoil and spherical

aberration). Please see Supplementary Table 1 for data collection and processing statistics and

the respective cryo-EM data processing schemes. For the Spike-Fu2 dimer interface we used a

particle set with partial-signal subtraction of all parts except for Fu2-RBD dimer interface

(containing two RBDs and two Fu2). From this we performed local reconstruction (non-

uniform). The local reconstruction resulted in a map with 2.89Å overall resolution as compared

to 3.2Å overall resolution for the full map.

Cryo-EM model building and structure refinement

The structure of the spike protein trimer PDB: 7KSG [5] was used as a starting model for model

building. A homology model for Fu2 was generated by swiss model [37] with PDB: 5LHN as

template (chain B) [38]. Structure refinement and manual model building were performed using

COOT [39] and PHENIX [40] in interspersed cycles with secondary structure, Ramachandran,

rotamers and bond geometry restrains. Structure figures and EM density-map figures were

generated with UCSF ChimeraX [41] and COOT, respectively. Please see Supplementary Table 2

for refinement and validation statistics.

References

1.   Cele, S. *et al.* Escape of SARS-CoV-2 501Y.V2 from neutralization by convalescent plasma.

medRxiv 2021.01.26.21250224 (2021) doi:10.1101/2021.01.26.21250224.

2.   Muyldermans, S. Nanobodies: Natural Single-Domain Antibodies. *Annu. Rev. Biochem.* **82**, 775–

797 (2013).

3.   Moliner-morro, A. *et al.* Picomolar SARS-Cov-2 neutralization using multi-arm PEG nanobody

constructs. *Biomolecules* **10**, 1–11 (2020).

4.   Hultberg, A. *et al.* Llama-derived single domain antibodies to build multivalent, superpotent and

broadened neutralizing anti-viral molecules. *PLoS One* **6**, 1–12 (2011).

5.   Koenig, P.-A. *et al.* Structure-guided multivalent nanobodies block SARS-CoV-2 infection and

suppress mutational escape. *Science (80-.).* **371**, eabe6230 (2021).

6.   Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.

Science (80-.). **367**, (2020).

7.   Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell*

181, 281-292.e6 (2020).

8.   Walls, A. C. *et al.* Unexpected Receptor Functional Mimicry Elucidates Activation of Coronavirus

Fusion. *Cell* **176**, 1026-1039.e15 (2019).

9.   Pallesen, J. *et al.* Immunogenicity and structures of a rationally designed prefusion MERS-CoV

spike antigen. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E7348–E7357 (2017).

10. Wang, C. *et al.* A human monoclonal antibody blocking SARS-CoV-2 infection. *Nat. Commun.* **11**,

1–6 (2020).

11. Liu, L. *et al.* Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2 spike. *Nature*

584, 450–456 (2020).

12. Lei, C. *et al.* Neutralization of SARS-CoV-2 spike pseudotyped virus by recombinant ACE2-Ig. *Nat.*

Commun. **11**, (2020).

13. Custódio, T. F. *et al.* Selection, biophysical and structural analysis of synthetic nanobodies that

effectively neutralize SARS-CoV-2. *Nat. Commun.* **11**, (2020).

14. Hanke, L. *et al.* An alpaca nanobody neutralizes SARS-CoV-2 by blocking receptor interaction. *Nat.*

Commun. **11**, 4420 (2020).

15. Xiang, Y. *et al.* Versatile and multivalent nanobodies efficiently neutralize SARS-CoV-2. *Science*

(80-.). **370**, 1479–1484 (2021).

16. Schoof, M. *et al.* An ultrapotent synthetic nanobody neutralizes SARS-CoV-2 by stabilizing

inactive Spike. *Science (80-.).* **370**, 1473–1479 (2021).

17. Ibañez, L. I. *et al.* Nanobodies with in vitro neutralizing activity protect mice against H5N1

influenza virus infection. *J. Infect. Dis.* **203**, 1063–1072 (2011).

18.   Monteil, V. *et al.* Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using Clinical-

Grade Soluble Human ACE2. *Cell* **181**, 905-913.e7 (2020).

19.   Cardoso, F. M. *et al.* Single-Domain Antibodies Targeting Neuraminidase Protect against an H5N1

Influenza Virus Challenge. *J. Virol.* **88**, 8278–8296 (2014).

20.   Power, U. F., Stortelers, C., Allosery, K. & Melero, J. A. Generation and Characterization of ALX-

0171 , a Potent Novel Therapeutic Nanobody for the Treatment of Respiratory Syncytial Virus

Infection. *Antimicrob. Agents Chemother.* **60**, 6–13 (2016).

21.   Planas, D. *et al.* Sensitivity of infectious SARS-CoV-2 B.1.1.7 and B.1.351 variants to neutralizing

antibodies. doi:10.1101/2021.02.12.430472.

22.   Tada, T. *et al.* Neutralization of viruses with European, South African, and United States SARS-

CoV-2 variant spike proteins by convalescent sera and BNT162b2 mRNA vaccine-elicited

antibodies. doi:10.1101/2021.02.05.430003.

23.   Hu, J. *et al.* Emerging SARS-CoV-2 variants reduce neutralization sensitivity to convalescent sera

and monoclonal antibodies. *Cell. Mol. Immunol.* 1–3 (2021) doi:10.1038/s41423-021-00648-1.

24.   Wrapp, D. *et al.* Structural Basis for Potent Neutralization of Betacoronaviruses by Single-Domain

Camelid Antibodies. *Cell* **181**, 1004-1015.e15 (2020).

25.   Cortez-Retamozo, V. *et al.* Efficient tumor targeting by single-domain antibody fragments of

camels. *Int. J. Cancer* **98**, 456–462 (2002).

26.   Vieira, P. & Rajewsky, K. The half-lives of serum immunoglobulins in adult mice. *Eur. J. Immunol.*

18, 313–316 (1988).

27.   Tijink, B. M. *et al.* Improved tumor targeting of anti-epidermal growth factor receptor

Nanobodies through albumin binding: Taking advantage of modular Nanobody technology. *Mol.*

Cancer Ther. **7**, 2288–2297 (2008).

28.   Bangaru, S. *et al.* Structural analysis of full-length SARS-CoV-2 spike protein from an advanced

vaccine candidate. *Science (80-.).* **370**, 1089–1094 (2020).

29. Hanke, L. *et al.* The Antiviral Mechanism of an Influenza A Virus Nucleoprotein-Specific Single-

Domain Antibody Fragment. *MBio* **7**, e01569--16 (2016).

30. Hsieh, C. L. *et al.* Structure-based design of prefusion-stabilized SARS-CoV-2 spikes. *Science (80-.*

). **369**, 1501–1505 (2020).

31. Corman, V. M. *et al.* Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR.

Eurosurveillance **25**, 2000045 (2020).

32. Ishige, T. *et al.* Highly sensitive detection of SARS-CoV-2 RNA by multiplex rRT-PCR for molecular

diagnosis of COVID-19 by clinical laboratories. *Clin. Chim. Acta* **507**, 139–142 (2020).

33. Wölfel, R. *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* **581**,

465–469 (2020).

34. Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data preprocessing with Warp. *Nat.*

Methods **16**, 1146–1152 (2019).

35. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. CryoSPARC: Algorithms for rapid

unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

36. Zi Tan, Y. *et al.* Addressing preferred specimen orientation in single-particle cryo-EM through

tilting. *Nat. Methods* **14**, 793–796 (2017).

37. Waterhouse, A. *et al.* SWISS-MODEL: Homology modelling of protein structures and complexes.

Nucleic Acids Res. (2018) doi:10.1093/nar/gky427.

38. Kromann-Hansen, T. *et al.* Discovery of a novel conformational equilibrium in urokinase-type

plasminogen activator. *Sci. Rep.* **7**, 1–11 (2017).

39. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta*

Crystallogr. Sect. D Biol. Crystallogr. **66**, 486–501 (2010).

40. Adams, P. D. *et al.* PHENIX: A comprehensive Python-based system for macromolecular structure

solution. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 213–221 (2010).

41.   Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and analysis.

631   *Protein Sci.* **27**, 14–25 (2018).

632   42.   Chen, V. B. *et al.* MolProbity: All-atom structure validation for macromolecular crystallography.

633   *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 12–21 (2010).

634   43.   Barad, B. A. *et al.* EMRinger: Side chain-directed model and map validation for 3D cryo-electron

635   microscopy. *Nat. Methods* **12**, 943–946 (2015).

639
640

Acknowledgments: Experiments with replication-competent SARS-CoV-2 were performed in the Biomedicum BSL3 core facility, Karolinska Institutet. All cryo-EM data were collected in the Karolinska Institutet 3D-EM facility. We thank Agustin Ure for assistance with figure generation and Tomas Nyman (Protein Science Facility at KI) for providing access to SPR instruments.

Funding: LH was supported by the David och Astrid Hageléns stiftelse, the Clas Groschinskys Minnesfond and a Jonas Söderquist's scholarship. This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No. 101003653 (CoroNAb), to BM and GMM. BMH is supported by the Knut and Alice Wallenberg Foundation (KAW 2017.0080 and KAW 2018.0080). The work was supported by project grants from the Swedish Research Council to BMH (2017-6702 and 2018-3808), BM (2018-02381) and to GMM (2018-03914 and 2018-03843).

Author contributions: Conceptualization: LH, BM, GMM. Methodology: LH, DJS, JMC, BMH, BM, GMM. Investigation: LH, HD, DJS, LPV, EU, AMM, VK, NS, AP, CK, GKP, BMH. Data Curation: LH, BM. Visualization: LH, HD, BHM, BM. Funding acquisition: LH, BMH, BM, GMM. Supervision: LH, DJS, JMC, BM, GMM. Writing – original draft: LH, HD. Writing – review & editing: LH, HD, DJS, LPV, EU, AMM, VK, GKP, JMC, BMH, BM, GMM.

Competing interests: LH, DJS, LPV, BM, and GMM are listed as inventors on a patent application describing SARS-CoV-2 nanobodies.

Data and materials availability: The cryo-EM density maps have been deposited in the Electron Microscopy Data Bank under accession codes EMD-12561 (dimer of spike trimer + 6 Fu2) and EMD-12465 (localized reconstruction of 2 RBDs + 2 Fu2). The atomic coordinates have been deposited in the Protein Data Bank under IDs 7NS6 (dimers of spike trimers + 6 Fu2) and 7NLL (localized reconstruction of 2 RBDs + 2 Fu2). All nanobody sequences will be made available on public databases upon publication.

Supplementary Figures

Supplementary Fig. 1. Fu2 variants identified by next-generation sequencing (NGS) display similar neutralization potential and binding kinetics. (**A**) Average difference of Fu2 variants is shown as a tree diagram. (**B**) Fu2 and seven Fu2-variants were expressed in *E.coli*, purified, and analyzed by SDS-PAGE and Coomassie staining. (**C**) A dilution series of Fu2 and its variants were incubated with SARS-CoV-2 pseudotyped lentivirus (PSV) for 1 hour at 37 °C before infecting HEK293T-hACE2 cells. Neutralization (in %) compared to the untreated PSV is shown. (**D**) Binding kinetics of Fu2 and variants to the RBD were measured by surface plasmon resonance (SPR). Site-specifically biotinylated RBD was immobilized on streptavidin sensor chips, and kinetics for a dilution series of the indicated monomeric nanobodies were measured. Sensorgrams are color-coded based on concentration. The fit is based on the 1:1 Langmuir model and is shown in dark grey solid lines. (**E**) Kinetic parameters of Fu2 and Fu2 variants binding to the RBD.

Supplementary Fig. 2. (**A**) Surface view of the RBD with interface-major and interface-minor color coded. (**B**) Relative positions of two Fu2 molecules binding the RBD. (**C**) The interaction between two RBDs and two Fu2s showing in gray the β-hairpins in the Fu2 framework region (39-45) that help to mediate the dimer-of-trimers Fu2-spike complex. (**D**) Close-up view of (**C**) (**E** and **F**) Assessment of Fu2-Ty1 heterodimer binding. Alignment of the Spike-Fu2 structure to the Spike-Ty1 structure (RBD in 'up' conformation) (PDB:6ZXN). Arrows indicate partial overlap of binding surface on RBD. (**G**) Assessment of Fu2-RBD interaction in RBD-down conformation. RBD-Fu2 structure superimposed to RBD-down of spike in 2-up conformation (PDB:7A29) (**H**) Assessment of Fu2-RBD interaction in RBD-down conformation in dimeric spike. 2-up spike (PDB:7A29) is shown superimposed with the spike dimer-of-trimers.

699
700

Align whole Spike protomer

Align only RBD

C

Fu2-RBD

PDB:7KSG
3-up Spike + VhhE

PDB:7K4N
3-up Spike +
S2E12 Fab fragment

PDB:7A98
3-up Spike + ACE2

PDB:7A93
2-up Spike

~6Å

~6Å

~8Å

PDB:7A29
2-up Spike + Sb23

PDB:6XM4
1-up Spike

PDB:6ZXN
1-up Spike + Ty1

PDB:6XM3
1-up Spike

701
702

Supplementary Fig. 3. Assessment of Fu2 mediated structural changes in spike structure.
(**A**) Alignment of dimeric spike trimer with spikes in 1-up, 2-up and 3-up conformation (one
protomer in 'up' state shown). (**B**) Structural alignment of dimeric spike trimer with RBDs of 1-
up, 2-up and 3-up spikes (protomer in 'up' conformation). (**C**) Structural comparison of RBD
(Fu2-bound dimeric spike) with eight different RBDs structures.

Supplementary Fig. 4. (**A**) Sequence alignment of RBDs from SARS-CoV and SARS-CoV-2. Interface-major and interface-minor residues are highlighted. Arrows indicate the different amino acids of the interface. (**B**) Fu2 variants and analysis of interface residues. Distinct residues of different variants shown as sticks (color coded).

Supplementary Fig. 5 Cryo-EM image processing scheme. Particles picked by Warp were processed in cryoSPARC v3.1. Representative 2D class averages are shown. Particles contributing to the clean classes were used to generate ab-initio reconstructions (five classes) followed by heterogeneous refinement. One class showing high-resolution features was refined further (C1 symmetry). For localized reconstruction, particle subtraction followed by local refinement (Non-uniform) was performed.

Supplementary Fig. 6 Cryo-EM validation (**A**) and (**B**) 3D FSC and sphericity of 'dimer' of spike trimer and of localized map, (**C**) and (**D**) Local resolution estimation of 'dimer' of spike trimer and local resolution estimation of localized map (**E**) and (**F**) Angular distribution of 'dimer' of spike trimer reconstruction and localized reconstruction map (**G**) and (**H**) FSC curves of 'dimer' of spike trimer and of localized map (**I**) model-map fitting atomic resolution.

Supplementary Table 1. Cryo-EM data collection and processing

	Dimer of spike trimer + 6 Fu2	Localized reconstruction of 2 RBDs + 2 Fu2
EMDB	EMD-12561	EMD-12465
PDB	7NS6	7NLL
Data collection		
Microscope	TFS Krios G3i	
Voltage (kV)	300	
Detector	Bioquantum K3	
Recording mode	Counting	
EFTEM SA Magnification	165kx	
Calibrated pixel size (Å)	0.51	
Movie micrograph pixel size (Å)	0.51	
Flux (e$^-$/[(camera pixel)*s])	8.4	
Number of frames per movie micrographs	60	
Total movie micrograph exposure time (s)	1.5	
Fluency (e$^-$/Å^2)	48.6	
Underfocus range (μm)	0.2 – 1.3	
Energy filter slit width (eV)	10	
Data collection automation software	EPU	
EM Data processing		
Number of micrographs	14,081	14,081
Number of total projection images	1,035,962	1,035,962
Number of projection images used for reconstruction	277,372	277,372
Symmetry	C1	C1
Map resolution (FSC 0.143; Å)	3.18	2.9
Map resolution no mask (FSC 0.143; Å)	4.2	8.9
B-factor applied to map (Å^2)	68.1	37.3
3D-FSC [36]		
3D-FSC resolution (3D-FSC 0.143; Å)	4.0	3.2
Sphericity (3D-FSC)	0.831	0.924

Supplementary Table 2. Cryo-EM model refinement and validation statistics

	Dimer of spike trimers + 6 Fu2	Localized reconstruction of 2 RBDs + 2 Fu2
EMDB	EMD-12561	EMD-12465
PDB	7NS6	7NLL
Refinement		
Refinement Package	PHENIX 1.19	PHENIX 1.19
Refinement space	Real	Real
Highest resolution used	3.2	2.9
Model composition		
Atoms refined	55904	5058
Residues SARS-CoV-2 Spike (or RBD) and nanobody Fu2	7109	656
Glycans	50	2
Full Map-model correspondence		
Map correlation coefficient	0.87	0.90
Map to model FSC (Å) (FSC 0.143)	3.2	2.6
Mean B-factor		
Residues SARS-CoV-2 Spike (or RBD) and nanobody Fu2	140	85.89
Glycans	129	96.22
Model geometry		
RMSD ideal bond length (Å)	0.004	0.003
RMSD ideal bond angles (˚)	0.826	0.529
Side chain rotamer outliers (%)	0	0
C-ß outliers (%)	0.38	0
Ramachandran plot		
Favoured (%)	97.76	99.38
Outliers (%)	0	0
General macromolecular model validation		
MolProbity score [42]	1.66	1.38
Clashscore [42]	12.38	6.92
EM specific model validation		
CaBLAM (%) [43]	2.55	0.94
EM-ringer score [43]	1.73	3.16