TITLE

CSF1R inhibitor levels determine sex-specific phenotype of resilient microglia and

neurofunctional rescue leading to extended survival in tauopathy mice

AUTHORS

Noah R. Johnson[1,2†], Peng Yuan[1,3†], Erika Castillo[1], T. Peter Lopez[1], Weizhou Yue[1], Annalise

Bond[1], Brianna M. Rivera[1], Miranda C. Sullivan[1], Masakazu Hirouchi[1,4], Kurt Giles[1,5], Atsushi

Aoyagi[1,4], and Carlo Condello[1,5†*]

[1] Institute for Neurodegenerative Diseases, UCSF Weill Institute for Neurosciences, University

of California, San Francisco, CA 94158, USA.

[2] University of Colorado Alzheimer's and Cognition Center, Linda Crnic Institute for Down

Syndrome, Department of Neurology, University of Colorado Anschutz Medical Campus,

Aurora, CO 80045, USA.

[3] Department of Rehabilitation Medicine, Huashan Hospital, State Key Laboratory of Medical

Neurobiology, Institute for Translational Brain Research, MOE Frontiers Center for Brain

Science, Fudan University, Shanghai 200032, China.

[4] Daiichi Sankyo Co., Ltd., Tokyo, 140-8710, Japan.

[5] Department of Neurology, UCSF Weill Institute for Neurosciences, University of California, San

Francisco, CA 94158, USA.

[†]N.J., P.Y. and C.C. contributed equally to this work. *Correspondence to:

carlo.condello@ucsf.edu

ABSTRACT

Microglia are central to pathogenesis in many neurological conditions. Drugs targeting colony-stimulating factor-1 receptor (CSF1R) to block microglial proliferation in preclinical disease models have shown mixed outcomes, thus the therapeutic potential of this approach remains unclear. Here, CSF1R inhibitors were evaluated in tauopathy mice using multiple dosing schemes, drug analogs, and longitudinal measurements in the brain and plasma. A sex-independent reduction in pathogenic tau was seen in several models and non-microglial gene expression patterns reverted toward a normal wild type signature. Surprisingly, despite greater drug exposure in male mice, functional rescue and extended survival was only observed in female mice. A dose-dependent upregulation of immediate early genes and neurotransmitter dysregulation were observed in the brains of male mice only, indicating that excitotoxicity may have precluded functional benefits. Drug-resilient microglia in male mice exhibited morphological and gene expression patterns consistent with increased neuroinflammatory signaling, suggesting a mechanistic basis for sex-specific excitotoxicity. These data argue that complete microglial ablation is neither required nor desirable for neuroprotection and that therapeutics targeting microglia must consider sex-dependent effects on functional outcomes.

INTRODUCTION

Microglia, the resident innate immune cells of the central nervous system (CNS), are important for neurodevelopment and homeostasis, and are a fundamental component to pathogenesis in many neurological conditions. We now appreciate that microglia are heterogeneous cells, are influenced by the periphery, have sex-dependent biology, and can be helpful or harmful depending on the disease stage or specific pathology[1-4]. Gene mutations affecting the expression and sequence of microglial genes (e.g. *TREM2*, *CD33*, and *MS4A*) increase risk for Alzheimer's disease (AD), and implicate microglia in several disease pathways including toxic protein aggregation (Aβ and tau) and neuroinflammation[5,6]. Thus, for the first time, there is unequivocal evidence in humans that certain microglial functions are robustly involved in the pathogenesis of neurodegenerative disease. However, the precise mechanisms governing microglia function in disease are still not well understood.

In tauopathy (a family of neurodegenerative disorders characterized by tau inclusions in neural cells), there is growing evidence that microglia play an early and constant role in tau

aggregation and neuronal loss. Disease-activated microglia can secrete pro-inflammatory

cytokines that regulate neuronal kinases and phosphotases causing tau hyperphosphorylation,

aggregation and consequent neurodegeneration[7-9]. Genome-wide transcriptomic studies have

identified innate immune pathways that implicate early and robust involvement of microglia in

human tauopathy[10,11] and related mouse models[12,13]. Deletion of microglia-specific genes or

genetic ablation of microglial cells in rodents have been useful approaches to dissect microglia-

mediated mechanisms in disease models, but pharmacologic tools to more dynamically

manipulate microglial function have been limited. Recently developed small-molecule drugs

targeting colony-stimulating factor-1 receptor (CSF1R), a receptor kinase critical for survival and

proliferation of CNS microglia, peripheral tissue macrophages and blood myeloid cells[14], are

approved for clinical use in various oncology indications[15], and have now been adopted by the

neuroscience community to study microglia biology. In the past few years, there have been

numerous studies using CSF1R inhibitors in models of neurological disease, but only a few

studies in models of primary tauopathy[16-19]. While important first steps, these studies only

explored single, static time points of treatment, or used only one sex. Given the dynamic nature

and complexity of microglial activation, the timing of CSF1R inhibition in tauopathy and its

translational relevance is still an open question.

Thus, the goal of our study was to define a therapeutic window that not only reduced

pathological markers, but also led to functional improvement. Moreover, we questioned whether

complete or continuous microglial ablation using CSF1R inhibitors was necessary given the

important and diverse roles these cells play in brain health and disease. Here, we systematically

test CSF1R inhibition using multiple drug analogs at several time points in transgenic mice

developing spontaneous tauopathy, and in an inoculation model of induced tauopathy. We

demonstrate a reduction of tau pathology in multiple dosing schemes without complete

microglial ablation; drug exposure levels correlated with the extent of tau-prion[20] and microglial

reduction. Surprisingly, we observed suppressed plasma biomarkers of neurodegeneration,

rescue of aberrant behavior, and extended survival in female mice only. These data reveal a

previously unrecognized sex-dependent therapeutic benefit of pharmacological CSF1R

inhibition. Transcriptome analyses showed that treated tauopathy mice exhibited a restored

gene expression profile similar to wild type mice; however, we observed a specific module of

sex- and drug concentration-dependent gene expression that might explain the lack of

functional rescue in male mice. Interestingly, residual microglia had a morphology similar to wild

type microglia and their gene expression pattern indicated a unique, sex-specific signature in

response to CSF1R inhibition. These data highlight yet another context for microglial

heterogeneity with implications for novel microglial biology, and argue that tempering microglial

activation with drugs, rather than cellular ablation, is a better therapeutic strategy with clinical

relevance.

RESULTS

CSF1R inhibition reduces pathogenic tau in the brains of Tg2541 mice

Building on previous findings[16-19], we first evaluated the effect of CSF1R inhibition on the levels

of pathogenic tau in the brains of transgenic mice expressing human tau, using a cell-based tau-

prion bioassay, enzyme-linked immunosorbent assay (ELISA), and immunohistochemical (IHC)

analysis. To deplete microglia, Tg2541 mice were dosed with one of two potent, orally

bioavailable, and brain-penetrant CSF1R inhibitors: PLX3397 (pexidartinib), which binds

receptor tyrosine kinases CSF1R, and to lesser extent, KIT and FLT3[21], and PLX5622, which

selectively binds CSF1R[22]. Three different treatment paradigms were evaluated: acute (2−4

months old), chronic (2−7 months old), and terminal (2 months old until death) (Fig. 1a−c).

Transgenic B6-Tg(Thy1-MAPT*P301S)2541 mice[23], referred to here as Tg2541 mice, express

the 0N4R isoform of human tau with the familial frontotemporal lobar degeneration (FTLD)-

linked P301S mutation[24], which increases its aggregation propensity and prion-like

characteristics[25,26]. We previously demonstrated that the levels of pathogenic tau in hindbrain

regions of Tg2541 mice were greater than in forebrain regions[27]. This observation is consistent

with the neuropathological staging of human FTLD and specifically of progressive supranuclear

palsy (PSP) where tau deposition begins and predominates in subcortical and brainstem

nuclei[28]. Therefore, the forebrain and hindbrain regions were examined separately in this study

(Fig. 1d).

We confirmed that CSF1R inhibition effectively reduced microglial markers Iba1 and P2yr12 by

an average of ~60% in both the forebrains and hindbrains of Tg2541 mice compared to vehicle

treatment, and that they had similar effects in the brains of C57BL/6J wild type mice

(Supplementary Fig. 1a−p). Principal component analysis of all Iba1 and P2yr12 data combined

showed that sex did not have a significant effect on the extent of microglial depletion by CSF1R

inhibitors (Supplementary Fig. 1q and Supplementary Data File 1), and therefore male and

female mice were grouped together for analysis unless otherwise noted.

We next employed a reproducible and rapid cell-based bioassay[27,29,30] to measure the activity of

replication-competent tau-prions in brain homogenates from Tg2541 mice. To ensure an

appropriate dynamic range in this bioassay, we optimized for dilution factor and assay duration

using aged Tg2541 mouse brain samples, which showed greater than 100-fold higher signal

than wild type mouse brain samples (Supplementary Fig. 2). Following acute, chronic, or

terminal treatment with PLX3397 or PLX5622, tau-prion activity in the forebrains of Tg2541 mice

was significantly decreased compared to vehicle-treated mice (Fig. 1e−g).

Hyperphosphorylation and aggregation of tau occurs first in hindbrain regions of Tg2541 mice,

especially in the brainstem and spinal cord, leading to motor deficits caused by severe

paraparesis[23]. This is consistent with our previous report of early and aggressive tau-prion

activity in hindbrain regions of Tg2541 mice[27]; as such, we found that acute CSF1R inhibition

was insufficient to reduce tau-prion activity in the hindbrain (Fig. 1e). However, chronic or

terminal treatment with PLX3397 did significantly reduce tau-prion activity in hindbrain regions

(Fig. 1f,g) and the spinal cords of Tg2541 mice (Supplementary Fig. 3). To examine other

markers of pathogenic tau, we measured the levels of tau phosphorylated at Ser396 (pS396) by

ELISA, and tau phosphorylated at Ser202/Thr205 (pS202/T205) by IHC. Acute, chronic, or

terminal PLX3397 treatment robustly reduced pS396 tau in both forebrain and hindbrain regions

of Tg2541 mice (Fig. 1h−j), and also reduced pS202/T205 tau in forebrain regions (Fig. 1k−m)

and in the spinal cord (Supplementary Fig. 3d). Since the various measures of tau pathology

represent different steps of tau pathogenesis (hyperphosphorylation vs. oligomerization vs.

filament formation), they may be differentially impacted by CSF1R inhibition with different

treatment regimens. Thus, to consider all tau measurements and both brain regions together,

we performed principal component analysis which revealed that pathogenic tau was reduced by

both CSF1R inhibitors, and that there was no significant effect of sex on drug efficacy (Fig. 1n

and Supplementary Data File 1).

Having verified the benefits of microglial depletion at an early disease stage, we next wondered

whether initiating CSF1R inhibition at a more advanced stage of disease would have similar

effects, simulating an interventional drug treatment. Thus, we dosed Tg2541 mice with PLX3397

in a delayed treatment paradigm (4−7 months old). Similar to terminal treatment, interventional

treatment also significantly reduced tau-prion activity in both the forebrain and hindbrain

(Supplementary Fig. 4a,b). Although pS396 tau levels were unchanged after interventional

treatment, levels of tau phosphorylated at Thr231 (pT231) were reduced in the forebrain

(Supplementary Fig. 4c,d). Considering the potential off-target effects of continuous, long-term

microglial depletion on brain function, we also wondered whether periodic CSF1R inhibition

might provide a safer, yet similarly efficacious therapy. Thus, we also tested PLX3397 dosed

intermittently by repeating dosing cycles of three weeks on followed by three weeks off. The

intermittent dosing interval was selected based on prior studies showing that there is rapid

microglial repopulation of the brain, and that morphological and transcriptional changes in the

microglia return to baseline levels within 21 days of removing PLX[31]. Intermittent treatment

produced similar reductions in the levels of microglial markers in both brain regions as for

continuous treatment (Supplementary Fig. 1r–t), but tau-prion activity and pT231 levels were

reduced only in the forebrain (Supplementary Fig. 4e–h). Taken together, these data suggest

that intermittent/interventional dosing is sufficient to reduce pathogenic tau in the forebrain of

Tg2541 mice, likely due to slower disease kinetics; however, continuous CSF1R inhibition is

necessary for the extended reduction of pathogenic tau in the hindbrain.

Tau has been shown to propagate throughout the brain in a prion-like fashion along

interconnected neural networks[32,33]. To test the hypothesis that microglial depletion may reduce

the propagation of tau-prions[20] in the brains of Tg2541 mice, we inoculated fibrils of the

microtubule-binding repeat domain of tau, referred to as K18 fibrils[34], into the hippocampus and

overlying cortex (forebrain regions) of Tg2541 mice and then treated them with PLX3397.

Compared to un-inoculated mice, K18-inoculated mice had significantly increased tau-prion

levels in the ipsilateral (inoculated) forebrain, as well as in the contralateral forebrain and in the

hindbrain (Supplementary Fig. 5), which suggests that tau-prions had propagated from the

inoculation site to those brain regions. However, acute PLX treatment was sufficient to

significantly reduce tau-prion levels in the ipsilateral forebrain and hindbrain, as well as in the

contralateral forebrain. Furthermore, tau-prion levels in the contralateral forebrain of PLX-

treated mice were not significantly different from the forebrain of un-inoculated, vehicle-treated

mice (Supplementary Fig. 5), which indicates that CSF1R inhibition prevented the spreading of

tau-prions from the inoculation site to this brain region.

CSF1R inhibition can affect peripheral immune cells such as blood myeloid cells and tissue

macrophages, in addition to microglia[17,35]. To determine if the effects of PLX3397 and PLX5622

on pathogenic tau in the brain are due, at least in part, to depletion of peripheral CSF1R-

expressing cells we dosed Tg2541 mice with PLX73086[36], a non-brain penetrant CSF1R

inhibitor analog of PLX3397 and PLX5622. Chronic treatment with PLX73086 had no significant

effect on microglial markers Iba1 and P2yr12, or on levels of tau-prions or pTau[S396] or

pTau[T231] in the forebrain or hindbrain of Tg2541 mice (Supplementary Fig. 6). Therefore, the

effects of CSF1R inhibitors in peripheral compartments do not significantly contribute to their

reduction of pathogenic tau in the CNS. We also evaluated the numbers of Iba1[+]/CD206[+]

perivascular macrophages (PVMs) and found that PLX3397 treatment did not significantly

deplete this cellular population, although there was a trend ($P = 0.0947$) towards reduced PVMs

in female Tg2541 mice (Supplementary Fig. 7). Lastly, because there is limited data for CSF1R

expression in neurons after injury[37], we considered whether PLX3397 or PLX5622 might affect

neurons or their expression of tau protein in Tg2541 mice. Acute, chronic, and terminal CSF1R

inhibition did not significantly reduce levels of neuronal nuclei (NeuN), detected by IHC, or total

tau, detected by ELISA (Supplementary Fig. 8). Therefore, CSF1R inhibitors do not directly

affect measures of neuronal viability or tau expression, consistent with a prior report using

PLX3397 in cultured primary neurons[17]. Together, these data confirm that drug effects on

biological and functional end points are due to inhibition of CSF1R in CNS microglia.

199   CSF1R inhibition extends survival in female Tg2541 mice

We next focused on the terminal treatment paradigm with PLX3397 to evaluate the long-term

effects of CSF1R inhibition on lifespan and behavior. Tg2541 mice develop paraparesis from

5−6 months of age which makes feeding difficult, resulting in a loss of body weight and thus, a

greatly reduced lifespan compared to wild type mice. We found that terminal PLX treatment

significantly extended the median survival of female Tg2541 mice [16.5 days longer median

survival; $P = 0.0004$], but not male Tg2541 mice, compared to vehicle treatment (Fig. 2a,b). The

extended survival in PLX-treated female mice was preceded by significantly reduced weight

loss, which was not observed in male mice (Fig. 2c). Body weight at 180 days of age,

irrespective of treatment, was predictive of lifespan in female mice but not in male mice, with

less weight loss being correlated with longer survival (Fig. 2d). Lower forebrain tau-prion levels

were also correlated with longer survival in female mice but not in male mice (Fig. 2e),

suggesting that Tg2541 mice have a sex-specific physiological response to tauopathy. To

confirm the effect of PLX treatment on survival in a different experimental paradigm, we used a

midbrain inoculation model. Since Tg2541 mice spontaneously develop substantial tau

pathology in the midbrain[27], we predicted that K18 inoculation in the midbrain would accelerate

and synchronize the disease course, which would be ideal for studying mouse survival. Indeed,

female Tg2541 mice inoculated with K18 tau fibrils died significantly earlier than mice inoculated

with diluent, though no difference was observed in male mice (Fig. 2f,g). Consistent with our

prior result, PLX treatment significantly extended the median survival of female mice inoculated

with K18 tau fibrils [29.5 days longer median survival; $P = 0.0095$], but not male mice, compared

to vehicle treatment (Fig. 2h). Taken together, these data indicate that CSF1R inhibition robustly

extends the lifespan of female Tg2541 mice, even during an accelerated disease course.

CSF1R inhibition reduces aberrant behavioral phenotypes in Tg2541 mice

To examine the relationship between drug exposure and markers of disease progression more
closely, we collected blood plasma at monthly intervals from mice receiving terminal treatment
with PLX3397 or vehicle (Fig. 3a). Consistent with previous reports[17], male Tg2541 mice had
higher plasma (25.3%; $P < 0.0001$) and brain (44.9%; P = 0.0250) concentrations of PLX than
did female mice (Fig. 3b and Supplementary Fig. 9a–c); we also observed this difference in wild
type mice (Supplementary Fig. 9d,e). Male and female mice had *ad libitum* access to food, and
had similar rates of food consumption relative to body weight, independent of whether it
contained PLX or vehicle (Supplementary Fig. 10a,b). However, female mice were consistently
more active than male mice (Supplementary Fig. 10c,d). Thus, the reduced PLX exposure in
female mice is likely due to a higher metabolic and drug clearance rate compared with male
mice. In line with the hypothesis that drug exposure was excessive in male mice, we observed a
trend towards reduced body weight in male wild type mice receiving terminal PLX treatment, but
not in female wild type mice (Supplementary Fig. 9f). In spite of this sex-specific difference in
PLX exposure, we found that higher plasma concentrations of PLX were correlated with greater
microglial depletion in both forebrain and hindbrain regions, independent of sex (Fig. 3c).
Furthermore, higher PLX exposure was correlated with reduced tau-prion levels in the forebrain
regions of both male and female mice (Fig. 3d). Together, these data indicate that PLX has
dose-dependent on-target effects in both male and female mice.

Previous studies have demonstrated a common hyperactive phenotype in the early stages of
tauopathy in transgenic rodent models[38,39]. While the precise mechanism that leads to this
deficit is unclear, this phenotype is causally linked with tau aggregate burden[40]. Based on the
reduction of tau deposition we observed with PLX treatment, we sought to also examine its
effect on this hyperactive phenotype. Using an automated home-cage monitoring system, we
longitudinally tracked the activity levels in Tg2541 mice at different ages, measuring their
amounts of rearing, locomotion, and wheel running. We confirmed the previous reports, finding
that at early ages the Tg2541 mice displayed a hyperactive phenotype relative to wild type mice
(90−150 days old in females, 90−120 days old in males), while at later ages their activity was
significantly reduced (Fig. 3e), likely due to the accumulation of pathogenic tau in brain regions
associated with motor function. PLX treatment led to a consistent reduction in Tg2541 mouse
hyperactivity, but did not change their hypoactivity at later ages (Fig. 3e), indicating the activity

reduction is not due to a general weakening effect. Detailed examination of the individual activity

measurements revealed that PLX treatment normalized the amounts of wheel running (Fig. 3f)

and active time (Fig. 3g). These data indicate that PLX treatment corrects the aberrant behavior

of Tg2541 mice towards that of wild type mice.

257   Sex-dependent effects of CSF1R inhibition on a biomarker of CNS injury

To further interrogate CNS damage caused by tauopathy, or potentially caused by the observed

sex-dependent PLX exposure, we also evaluated the plasma levels of neurofilament light chain

(NfL). NfL is a validated blood-based biomarker of neuronal injury[41] which correlates with

disease progression and tau burden in human tauopathy[42,43]. Female PLX-treated mice had

reduced levels of NfL compared to vehicle-treated mice (Fig. 4a,b), consistent with reduced

CNS injury due to tauopathy. Conversely, NfL levels were increased in male mice following PLX

treatment (Fig. 4c), suggestive of PLX-induced toxicity. Consistent with these findings, PLX

treatment resulted in significantly increased NfL levels in male mice that received midbrain

inoculation of K18 tau fibrils, but not female mice (Fig. 4d–f). We found no correlation between

NfL level and survival in female mice, but in male mice, higher NfL levels were strongly

correlated with reduced survival in both PLX and vehicle treatment groups (Fig. 4g,h).

Furthermore, the concentrations of NfL in both the forebrain and hindbrain at death were

positively correlated with PLX3397 concentration in the brains of male, but not female, Tg2541

mice (Fig. 4i,j). Interestingly, intermittent treatment, resulting in a 50% lower total dosage of

PLX, produced a significant decrease in NfL levels in male mice, but had no effect in female

mice (Fig. 4k–m). Taken together, these data suggest that in female Tg2541 mice, tauopathy

drives CNS injury and its reduction by PLX effectively masks any effect of PLX toxicity, whereas

in male mice, excessive exposure causes CNS injury that supersedes any benefit of PLX.

Consistent with this premise, we found that PLX treatment increased plasma NfL levels in male

wild type mice, as expected, but also in female wild type mice (Fig. 4n–p), albeit at substantially

lower levels than tau-induced NfL in Tg2541 mice.

279   Peripheral CSF1R inhibition does not cause therapeutic or adverse effects in the CNS

CSF1R inhibitors have known adverse effects in the periphery including anemia, leukopenia,

and hepatotoxicity, which have been observed in human clinical trials[44]. To determine if

peripheral PLX toxicity was causing NfL release or reducing drug efficacy, we evaluated mice

receiving chronic treatment with PLX73086, the non-brain penetrant CSF1R inhibitor analog of

PLX3397 and PLX5622. We observed no effect of PLX73086 treatment on body weight or

plasma NfL levels in either male or female Tg2541 mice (Fig. 5a–d), indicating that peripheral

CSF1R inhibition does not impact tauopathy-driven phenotypes. Furthermore, we observed no

effect of PLX73086 treatment on body weight or plasma NfL levels in wild type mice (Fig. 5e–h),

demonstrating that the drug-induced toxicity indicated by NfL rise is dependent on the drug

entering the brain. Histopathological evaluation of liver sections of PLX3397-dosed Tg2541

mice stained with hematoxylin and eosin (H&E), Masson's trichrome, or Picosirius red did not

reveal any overt signs of liver injury or fibrosis (Fig. 5j), nor did quantification of the Picosirius

red-stained liver sections (Fig. 5k). Alkaline phosphatase (ALP) in the plasma, a different

indicator of liver damage, was elevated in Tg2541 mice following PLX3397 treatment; however,

the ALP levels were elevated in both male and female mice (Fig. 5l), indicating that the sex-

specific toxicity of PLX3397 in male mice likely occurs in the brain rather than in the periphery.

Taken together, these data further confirm that the drug exposure of brain-penetrant CSF1R

inhibitors was appropriate for female Tg2541 mice to reduce tauopathy and significantly extend

survival; however, the drug exposure was too high for male Tg2541 mice, and the resulting

neurotoxicity outweighed its therapeutic benefit.

CSF1R inhibition shifts gene expression patterns in Tg2541 mice towards wild type

To better characterize global molecular changes in the CNS due to CSF1R inhibition, we used

the Nanostring platform to analyze a curated panel of gene transcripts related to

neuroinflammation, myeloid cell function and neuropathology in bulk brain tissue following

chronic treatment of Tg2541 mice with PLX5622. The chronic treatment group was selected for

this analysis because the collection time point was synchronized (unlike in the terminal

treatment group) and seven months of age is close to the average lifespan of Tg2541 mice. We

measured mRNA transcripts of 1,841 genes, many of them shown to be regulated by tau or Aβ

pathology in previous genome-wide gene expression studies[12,13]. To validate the Nanostring

approach, we identified 53 genes with a broad range of expression level changes and measured

mRNA transcripts with quantitative reverse-transcription PCR (RT-qPCR) in the same sample

used for sequencing. The RT-qPCR results generally matched the trends shown in the

Nanostring data (Supplementary Fig. 11), supporting the validity of our transcriptomic data.

Since microglial cells are directly impacted by PLX treatment, we first excluded the microglial-

specific genes (see Methods) and examined the general trend of expression patterns among

different treatment groups. Pearson's correlation matrix showed high similarity among wild type

brains with or without PLX treatment (Fig. 6a), indicating that the gene expression pattern we

measured is not affected by the treatment itself. In contrast, PLX treatment in Tg2541 mice

caused a distinct shift in the gene expression pattern away from the vehicle-treated group.

Interestingly, the gene expression patterns of PLX-treated Tg2541 mice showed a stronger

correlation with wild type mice than with vehicle-treated Tg2541 mice (Fig. 6a dashed boxes

and Fig. 6b). To further quantify this shift, we performed partial-least squares (PLS) regression

analysis using the gene expression data from vehicle-treated Tg2541 and wild type mice (Fig.

6c filled circles), and projected the data from PLX-treated mice onto the PLS dimensions (Fig.

6c empty circles). This allowed us to represent the transgene-specific gene expression pattern

in a relatively low-dimensional space, and to quantify the changes associated with treatment by

calculating the population vector distances and angles in this space. We found that PLX

treatment significantly normalized gene expression patterns in Tg2541 mice towards those of

wild type mice (Fig. 6d,e, only two out of five dimensions are shown, covering >95% of the total

variance). The normalization in gene expression was further confirmed by similar trends in

neuron-specific genes (Supplementary Fig. 12). Importantly, PLX treatment in wild type mice

showed negligible changes in the gene expression patterns. These results indicate that PLX

treatment specifically suppresses the abnormal transcriptome associated with transgene

overexpression, consistent with its effects ameliorating pathogenic tau deposition.

Evidence for excitotoxicity with increased drug exposure

As described above, although we observed consistent reduction in the levels of pathogenic tau

in the brains of both male and female Tg2541 mice with PLX treatment, only female mice

benefited from extended survival, functional rescue, and reduced NfL levels (Figs. 1–4). We

hypothesized that excessive PLX dosing may underlie this sex-specific effect, as male mice

consistently had higher drug exposure in the plasma and CNS (Fig. 3 and Supplementary Fig.

9), and also benefitted from a lower, intermittent PLX dosing paradigm (Supplementary Fig. 4).

Indeed, in our transcriptomic analysis we identified individual genes whose expression was

associated with brain PLX5622 concentration or with sex (Fig. 6f). First, we ruled out the

possibility that sex-dependent functional benefits were caused by differential expression of the

drug target, CSF1R, or its ligand, CSF1 (Supplementary Fig. 13). Using a PLS regression of all

non-microglia genes to brain PLX concentration and sex for each sample, we calculated the

variable importance score along each of these dimensions. Interestingly, many immediate early

genes (IEGs) showed high importance scores (Fig. 6g), suggesting that IEGs might be a

module that is altered by CNS drug exposure. To further test this possibility, we examined all

IEGs (56 genes overlapped in our dataset) and found that their expression patterns fit closely

with the brain PLX concentration (Fig. 6h). Importantly, when we excluded the IEGs and

examined the PLX-induced transcriptome shift along the wild type-to-Tg2541 dimension, the

correlation of gene expression changes and brain PLX concentration was no longer significant

(Fig. 6i), indicating that the IEGs contribute substantially to the PLX treatment effects. Notably,

relative to wild type vehicle-treated mice, only male PLX5622-treated Tg2541 mice had

significantly upregulated expression of IEGs (Fig. 6j), which we also validated using qPCR of

the five most highly expressed IEGs (Supplementary Fig. 14a-c), as well as by *in situ* labeling of

Fos mRNA (Fig. 6k,l). Furthermore, we observed the same drug-dependent IEG up-regulation in

male mice in a completely independent study using PLX3397 (Supplementary Fig. 14d-f),

thereby confirming that this is a robust sex-specific effect of CSF1R inhibition. As increased IEG

expression can be indicative of neuronal hyperactivity, these data provide a plausible

mechanism by which excessive PLX dosing may have led to excitotoxicity, thereby masking its

therapeutic effect in male mice.

To interrogate this mechanism directly, we performed hydrophilic interaction liquid

chromatography tandem mass spectrometry (HILIC-MS/MS)-based metabolomics to measure

the levels of relevant neurotransmitters in forebrain lysates of the mice. PLX3397-treated male

Tg2541 mice, but not PLX-treated wild type or female Tg2541 mice, had increased brain levels

of the excitatory neurotransmitter glutamate (Fig. 7a), the most common cause of neuronal

excitotoxicity[45]. Unexpectedly, PLX-treated male Tg2541 mice also had increased levels of the

inhibitory neurotransmitter γ-aminobutyric acid (GABA) (Fig. 7b). However, these PLX-induced

alterations were not due to general neurotransmitter dysregulation because we did not observe

altered levels of acetylcholine, glycine, or histamine in the brains of male Tg2541 mice

(Supplementary Fig. 15). These data provide evidence of PLX-induced glutamate-driven

excitotoxicity in the brains of male Tg2541 mice which may, in part, underlie the sex-specific

benefit of CSF1R inhibition on mouse survival.

CSF1R inhibition ameliorates pathological activation of astrocytes

As suggested by the reduction of tau deposition, we hypothesized that astrocyte-driven

neuroinflammation would also be reduced by PLX treatment. Therefore, we examined

transcriptome shifts in astrocyte-specific genes upon PLX treatment. Similar to the neuronal-

specific genes, we observed a normalization of astrocyte-specific gene expression patterns

towards wild type in both forebrain and hindbrain regions (Fig. 8a,b). Using previously described

astrocytic gene signatures of disease[46], we found that PLX treatment led to a dose-dependent

reduction in the expression of the A1 astrocytic gene cluster associated with neurotoxic effects

(Fig. 8c,d). In addition, we measured astrogliosis over time using longitudinal bioluminescence

imaging (BLI) methods based on a previously established transgenic reporter system of glial

fibrillary acidic protein (GFAP)-driven luciferase[47], which we validated by IHC and mRNA

analyses (Supplementary Fig. 16). To perform reliable BLI in Tg2541 mice, we intercrossed

each transgenic line to an albino background and refined the method, using a synthetic luciferin

substrate to increase signal from deep hindbrain regions (see Methods and Supplementary Fig.

17). This technique allowed us to non-invasively measure astrogliosis in live mice longitudinally

over the course of PLX treatment. In vehicle-treated Tg2541 mice, the BLI signal gradually

increased with age (Supplementary Fig. 18), in accordance with the accumulation of tau

pathology and gliosis reported in Tg2541 mice[23,48]. Consistent with our hypothesis and with

measurement of GFAP using other methods (Supplementary Fig. 16), CSF1R inhibition

suppressed the BLI signal in both the forebrain and the hindbrain (Fig. 8e−g). Together, these

data suggest that astrocytic inflammation, in particular the neurotoxic astrocytes, driven by

microgliosis was attenuated by CSF1R inhibition, thus leading to a general neuroprotective

effect.

CSF1R inhibition preferentially eliminates a highly activated microglia subpopulation

Given reported roles of microglia in driving astrocytic inflammation and neurotoxicity in disease,

we next interrogated whether resilient microglial phenotypes could be responsible for the sex-

specific neurofunctional effects. Thus, we examined the morphological and transcriptional

changes in microglia following CSF1R inhibition. In tauopathy, microglia acquire an activated

morphology in brain regions where neurons contain tau aggregates, a phenomenon seen in

many focal neuropathologies[49]. Interestingly, the elimination of microglia in the Tg2541 mouse

brain following PLX3397 treatment was not uniform nor complete, but was the most effective in

the vicinity of tau aggregates (Fig. 9a,b). The microglial density near tau-laden neurons was

reduced by more than 60%, but in distal regions (>200 microns) the microglial density was not

significantly changed (Fig. 9c), indicating that microglia in the vicinity of the tau aggregates may

have increased sensitivity towards CSF1R inhibition. We then compared the morphologies of

PLX-resistant (residual microglia in PLX-treated mice) and PLX-sensitive microglia (in vehicle-

treated mice). Notably, we found that PLX resistance was associated with more abundant and

intricate microglial cell processes, close to the levels seen in wild type mice (Fig. 9d−g). The

number of microglial process branches ($P = 0.0075$), process lengths ($P = 0.028$), and territory

sizes ($P = 0.011$) in the forebrain were also significantly different between male and female

PLX-treated mice, with female mice showing more abundant and intricate microglial processes
by all three metrics (Fig. 9d−g). These data suggest that microglia associated with tau pathology
may be in an "activated" state, with a reduced number of processes. This view is consistent with
previous reports that activated microglia adopt a "disease-associated microglial" (DAM)
phenotype, with some of the signature genes associated with inflammatory responses that are
detrimental to adjacent neural cells[50,51]. Interestingly, our data suggests that DAM in tauopathy
are more vulnerable to PLX treatment, and that surviving microglia might be neuroprotective,
particularly in female mice.

In support of this view, transcriptome analyses showed that many microglial-specific genes were
upregulated in Tg2541 mice (Fig. 10a), among which the most notable were signature DAM
genes such as *Tyrobp*, *Clec7a*, *Trem2* and *CD68*. By correlation analysis among different
samples using a generalized Louvain algorithm[52], we found that the microglia-specific genes in
our dataset were clustered into three groups (Fig. 10b and Supplementary Fig. 19). The red-
cluster genes showed the highest degree of modulation by transgene overexpression, while the
blue-cluster genes showed a moderate degree of modulation, and the green-cluster genes
showed almost no difference between Tg2541 and wild type mice (Fig. 10b). Transgene-
modulated genes clustered into red and blue groups, consistent with a recent finding that tau
pathology activates both immune-activation and immune-suppression gene expression
modules[10].

We next compared the gene expression patterns in vehicle- and PLX5622-treated brains. We
examined all previously reported DAM signature genes[51] and found a partial match with the
activation markers in each of our identified gene clusters (Fig. 10c). Regardless of the
designation of homeostatic or activation genes reported in previous studies, the red-cluster
genes showed a stereotypical pattern of transgene activation and sensitivity to PLX treatment,
while green- and blue-cluster genes did not appear to be modulated by these factors. Given that
PLX predominantly eliminated microglia in the vicinity of the tau deposits (Fig. 9c), these data
suggest that red-cluster genes are preferentially expressed by tau-associated microglia.
Interestingly, when we examined PLX-induced expression changes in male and female mice
separately, we found that while the expression of red cluster genes were substantially
diminished by PLX independent of sex, green cluster genes and some blue cluster genes were
markedly increased in treated male mice (Fig. 10d,e). We validated some of the sex-specific
microglial genes affected by PLX treatment using real-time PCR as well as *in situ* labeling of
mRNA (Fig. 10f,g). To better describe the microglial gene expression in the surviving population

and account for the cell number reduction, we calculated the gene expression of resilient

microglia by normalization to six microglia-specific housekeeping genes. The normalized data

showed a trend towards increased gene expression in male mice, yet most of the male-specific

PLX-induced genes belonged to the green cluster (Fig. 10e), many of which are known to be

pro-inflammatory. Indeed, subsequent Ingenuity Pathway Analysis revealed a higher activation

of inflammation-related pathways in PLX-treated male mice compared to female mice (Fig. 10h).

These inflammation-related pathways included tau-activated NF-κB[53] and excitotoxicity[54]

pathways, but not amyloid-induced inflammasome genes[55] (Supplementary Fig. 20). Pathways

related to phagocytosis or microglia growth, on the other hand, did not show such consistent

sex-specific change (Fig. 10h). Taken together, our data show morphological and transcriptional

changes in microglia associated with tau deposition, consistent with a pattern of pathological

activation. CSF1R inhibition appears to preferentially eliminate these microglia in female mice,

leaving the brain with a more quiescent and less inflammatory microglial population. Male mice,

on the other hand, showed a drug-induced inflammatory microglia phenotype, which might

contribute to neuronal excitotoxicity and diminished therapeutic effect.

DISCUSSION

Our study reveals several major findings from a comprehensive evaluation of CSF1R inhibitors

in preclinical models of tauopathy. Importantly, we present the first line of evidence that CSF1R

inhibition reduces pathology that leads to functional improvements associated with longer

lifespan and reduced behavioral deficits in tauopathy mice (Figs. 2 and 3). Overall, our data

showing a reduction of pathogenic tau is consistent with prior studies using a different drug

scaffold targeting CSF1R (JNJ-527; edicotinib) in Tg2541 mice[18], or using PLX3397 in a

different mouse model of tauopathy (TgPS19)[17]. However, in our study, neuroprotection

occurred despite incomplete microglia depletion (~60%); upon deeper analysis, we identified

distinct microglial-specific gene clusters suggesting subsets of microglia responsive to

tauopathy or resilient to CSF1R inhibition (Fig. 10). This finding is in line with the wealth of data

demonstrating that microglia exist as unique subsets in different brain regions, sexes, ages or

disease states[2-4]. From this perspective, our data suggest that it may be possible to target

specific subsets of activated microglia in tauopathy, without affecting other beneficial microglia

populations. Taken together, we argue that CSF1R inhibition causing complete microglia

ablation is unnecessary for therapeutic benefits, and may possibly be detrimental in humans

given that microglia are important for brain homeostasis and defending against other insults.

Several prior studies have suggested that therapeutic outcomes may only be achieved with

complete ablation of microglia, although these studies used different mouse models or dosing

regimens[17,19]. Nevertheless, these disparate findings in prior literature are now more

interpretable alongside our study, which sheds light on the intricate relationships between

CSF1R inhibitor dosing, microglial depletion and therapeutic outcomes.

The precise mechanism of CSF1R inhibitors causing reduced tau pathology is still unclear, but

our data indicates that activated microglia are the primary target resulting in reduced numbers of

cells producing pro-inflammatory cytokines[7-9] and other disease-associated microglial factors

(Fig. 10) that stoke tau pathogenesis in neurons, such as apolipoprotein E[17,56] and complement

proteins[57]. In addition, it seems plausible that CSF1R inhibitors may also block microglia-

mediated activation of A1 astrocytes[46], which in turn secrete factors that also drive tau

pathogenesis; blocking this cellular feed-forward pathway using a different drug targeting

microglia led to neuroprotection in a synucleinopathy model of Parkinson's disease[58].

Consistent with this view, PLX-treated mice exhibited a restored astrocyte phenotype,

suggesting that therapeutic benefits in our study may also be due, in part, to quelling neurotoxic

astrocytes (Fig. 8). Pharmacological CSF1R inhibition has been reported to also deplete

PVMs[59]. Although we did not detect a significant reduction in the numbers of cortical blood

vessel-associated PVMs in our studies (Supplementary Fig. 7), their roles in tauopathy remain

to be determined. While there is cross-talk between peripheral immune cells and microglia[1], we

show that a non-brain penetrant analog, PLX73086, did not affect CNS microglia, tau pathology

or NfL levels (Fig. 5 and Supplementary Fig. 6), and it is thus unlikely that CSF1R inhibition in

the periphery contributes to the phenotypic rescue observed in our study. Lastly, it is possible

that chronic PLX treatment caused depletion of some oligodendrocyte progenitor cells (OPCs)

in our study, but we expect that PLX did not affect mature oligodendrocytes or myelination[60].

The relationship between OPC biology and tau pathology in neurons is largely unknown, and

thus it remains unclear how CSF1R inhibition in OPCs contributes, if at all, to the mechanism of

action. Nevertheless, this topic warrants further investigation.

Tau pathology in Tg2541 mice is associated with moderate microgliosis and an up-regulation of

transcriptomic signatures of microglial activation[10,50]. Our transcriptome analysis showed

activation of two major clusters of microglial-specific genes in Tg2541 mice. These two clusters

had a high degree of overlap with the immune activation and suppression modules recently

described in tauopathy mice and FTLD patients[10], indicating a specific reactive transcriptional

program of microglia towards tau pathology. Consistent with this view, genes in the activated
clusters also matched transcriptome modules described in activated microglia in
neurodegeneration models (such as *Itgax* and *Clec7a*), but not in tumor or acute inflammation
models[50]. We found an additional cluster of microglial genes that had similar expression in
Tg2541 and wild type mice. Intriguingly, this cluster was not affected by PLX treatment in female
mice, while in male mice this cluster exhibited marked activation (Fig. 10). Considering that PLX
eliminates more than half of the total microglia population, a parsimonious explanation for this
sustained gene cluster is that they are preferentially expressed by an inert microglial
subpopulation that does not respond to tauopathy or lower dose CSF1R inhibition[61]. Consistent
with this notion, we found that PLX treatment preferentially eliminates activated microglia in the
vicinity of tau deposits, and thus most surviving microglia are not in direct contact with tau-laden
neurons (Fig. 9). This is in contrast to Aβ mice, in which the surviving microglia are usually
associated with Aβ plaques following PLX treatment[22,62]. We found that surviving microglia in
female mice were non-inflammatory, and had longer and more elaborate processes compared
to vehicle-treated microglia, showing functional and morphological features more similar to
those of wild type microglia (Fig. 9). In sum, our data describe a microglial genetic signature that
remains stable in Tg2541 mice with or without PLX treatment, likely representing a "dormant"
microglial subpopulation that are less dependent on CSF1R for survival, or are less sensitive to
CSF1R inhibition at the doses administered in our study.

Sex-specific differences exist in mouse microglial morphology, function, gene expression, and
response to tauopathy, and the differences increase with age[63-66]. Our data identify a sex-
dependent effect on therapeutic exposure and efficacy of CSF1R inhibition in Tg2541 mice. A
difference in the plasma levels of PLX3397 during *ad libitum* oral dosing in male and female
mice has been noted previously[17]; however, only male mice were evaluated further. We
examined both male and female Tg2541 mice and found that, despite similar food intake,
plasma and brain levels of PLX3397 were higher in male mice compared to female mice (Fig. 3
and Supplementary Fig. 9). However, at this level of drug exposure, only female mice received
a functional benefit from CSF1R inhibition, an unexpected and clinically relevant outcome that
would have been overlooked had our analysis been focused on a single sex. In male Tg2541
mice, despite a robust reduction of microglia and pathogenic tau, PLX treatment did not slow
weight loss or extend survival, and plasma NfL levels were significantly increased, indicative of
neuronal damage[41]. Neurodegeneration in Tg2541 mice has been shown to be largely limited to
spinal cord motor neurons[23,67], and we found that NeuN and total tau immunoreactivity in the

brain were largely unchanged by CSF1R inhibition (Supplementary Fig. 8). Thus, the functional

deficits we measured are likely caused by neuronal dysfunction rather than neuronal death, but

can be rescued by CSF1R inhibition. It has been suggested before that microglia from male

animals may exhibit an increased responsiveness to CSF1R depletion compared to microglia

from female animals[68]. Our results indicate that despite robust on-target effects for microglial

depletion, male mice developed a PLX-induced inflammatory microglial phenotype (Fig. 10).

Furthermore, we observed a concentration-dependent activation of IEGs in PLX treated Tg2541

mice, suggesting that excessive PLX dosing in male mice may lead to excitotoxicity (Fig. 6),

thus masking the beneficial effect of tau reduction. Microglia in male mice adopted a

transcriptome and morphological phenotype that have been previously linked to

excitotoxicity[54,69]. Curiously, IEGs were not significantly upregulated in male wild type mice,

indicating that their activation may not be due to high concentration of PLX alone, but may also

be dependent on tau deposits. Consistent with premise, we identified increased glutamate and

GABA neurotransmitter levels in the brains of male Tg2541 mice, but not male wild type mice or

any female mice (Fig. 7), indicative of excitotoxicity following PLX dosing. Previous studies have

linked tau accumulation and aberrant neural activity *in vivo*[70]. On the other hand, microglia are

known to mediate neuroprotection against excitotoxicity[69,71] and elimination of microglia can

exacerbate seizures and related neuronal degeneration[72]. Therefore, the concurrent tau

removal and drug-induced inflammation driven by surviving microglia may increase the risk for

hyperactivity, resulting in excitotoxicity in male mice with high PLX concentrations. Other sex-

specific differences may also contribute to microglial sensitivity to CSF1R inhibition by a

currently unknown mechanism. Future translational studies of pharmacological CSF1R

inhibitors will need to carefully evaluate the role of sex in both safety and therapeutic outcomes.

CSF1R inhibitors were shown to be protective in mouse models of other neurodegenerative

diseases, such as AD and Down syndrome[22,73,74]. However, under different treatment

conditions, CSF1R inhibition did not affect Aβ plaque burden, but did rescue some functional

deficits[62,75]. Therefore, microglia play a dynamic role in the brain's response to Aβ pathogenesis,

and their attenuation may impart distinct benefits at different stages of disease. Our results

suggest that, in primary human tauopathies, a subset of microglia play a net negative role

before, during, and after disease onset and that their removal may be a viable therapeutic

strategy. It remains to be determined if similar benefits should be expected for tauopathy in AD

given the preceding comorbid Aβ pathology, but this may be elucidated in rodent models co-

expressing human tau and Aβ. Nevertheless, because CSF1R inhibition has not been reported

to be detrimental in Aβ mice, CSF1R inhibitors could ameliorate AD-related tauopathy even if

caused by different disease mechanisms. Ongoing human clinical trials of CSF1R inhibitors in

AD (e.g. NCT04121208) may provide additional mechanistic insights.

Primary human tauopathies constitute a class of neurodegenerative diseases caused by tau

misfolding and aggregation and include progressive supranuclear palsy (PSP), corticobasal

degeneration (CBD) and Pick's disease, among others. When combined with AD, in which tau

aggregation follows Aβ deposition, tauopathies afflict a significant proportion of the human

population, and thus novel approaches to directly or indirectly block tau pathogenesis or its

downstream effects are urgently needed. Our study highlights several aspects of

pharmacological CSF1R inhibition that bolster its therapeutic potential for human tauopathies.

First, we observed greater efficacy of early (acute) CSF1R inhibition to restrict tau-prion levels in

forebrain regions (Fig. 1e), likely due to reduced disease severity there relative to hindbrain

regions. At later stages (chronic and terminal), CSF1R inhibition with PLX3397 did reduce tau-

prion levels in the hindbrain, albeit a modest reduction relative to the effects in the forebrain

(Fig. 1f,g). These findings suggest that early and long-term CSF1R inhibition (though not

necessarily continuous) would most effectively mitigate human tauopathy. Second, we

demonstrated that interventional dosing of Tg2541 mice, initiated at a stage when robust tau

deposition had already occurred[23,27], led to a significant reduction in pathogenic tau

(Supplementary Fig. 4). Therefore, our data support some potential clinical benefit of CSF1R

inhibitors for treatment, in addition to prevention, of tauopathy. This is important because

prophylactic prevention of non-autosomal dominant neurodegenerative diseases may be difficult

due to a lack of definitive prognostic biomarkers paired with the fact that aggregation of the

causative proteins can occur years or decades prior to symptom onset[30]. Third, we found that

intermittent dosing of Tg2541 mice at three-week intervals produced a significant reduction in

pathogenic tau (Supplementary Fig. 4). Despite relatively minimal off-target effects from

continuous, long-term dosing of CSF1R inhibitors in mice[22], non-human primates[76], and

humans[21], intermittent dosing would be clinically preferable if a similar therapeutic outcome was

achieved, given the important functions for microglia and related peripheral cells in innate

immunity. Because neurodegenerative tauopathies are slow, protracted diseases and microglia

are long-lived[77], it is conceivable that breaks in dosing may occur on the order of months or

years and be informed by medical imaging probes for microglia activation[78]. Imaging will be

highly valuable for guiding intermittent dosing because microglial repopulation in a pathological

setting (e.g., disease or normal aging) may not necessarily result in a return to baseline

transcriptional profiles. Fourth, we found CSF1R inhibition to extend the survival of female

Tg2541 mice (Fig. 2), indicating that the reduction in pathogenic tau in this model system

translates to an improved clinically relevant outcome. We postulate that if CSF1R inhibitor

dosing was optimized for male Tg2541 mice, any adverse effects in the CNS or periphery would

likely be diminished and their survival extended. Fifth, we showed that complete microglial

depletion is not necessary, or even desirable, for a therapeutic benefit. As discussed above, the

microglia that survive CSF1R inhibition represent a unique microglial sub-population that likely

serves important functions in brain homeostasis. Future preclinical studies may pinpoint the

precise level, timing, and frequency of CSF1R inhibition such that the detrimental effects of

microglial activation are minimized while an appropriate number of homeostatic microglia

remain for brain surveillance. Lastly, CSF1R inhibitors applied in conjunction with tau

immunotherapy may prove to be a successful combination therapy; because microglia are not

needed for antibody effector function[79], removing tauopathy-activated microglia would slow tau

pathogenesis and may also increase the efficacy of tau immunotherapy. Taken together, our

data strongly support the therapeutic modulation of microglial activation by CSF1R inhibitors as

a promising approach to treating human tauopathies.

METHODS

Animals

The Tg2541 transgenic mouse line expresses the human 0N4R tau isoform under the Thy1.2

genetic promoter. Tg2541 mice were originally generated on a mixed C57BL/6J × CBA/Ca

background[23] and were then bred onto a congenic C57BL/6J background using marker-assisted

backcrossing for eight generations before intercrossing to generate homozygous mice. Albino

Tg2541 mice were generated by intercrossing Tg2541 with C57BL/6J mice expressing a

spontaneous mutation in the tyrosinase gene (homozygous for Tyr^{c-2J}) causing albinism

(Jackson Laboratory; 000058). To generate mice for *in vivo* bioluminescence imaging, we

employed Tg(*Gfap*-luc) mice, which express firefly luciferase under the control of the murine

Gfap promoter (gift from Caliper Life Sciences). These reporter mice were originally on the FVB

background, but we backcrossed them to a congenic B6 background, and then crossed them to

the B6-albino background. To create bigenic mice, albino Tg2541 mice were crossed with albino

Tg(*Gfap*-luc) animals to produce double hemizygotes; next, double hemizygotes were crossed

and the progeny were screened for the presence of both transgenes expressed at

homozygosity. Animals were maintained in a facility accredited by the Association for

Assessment and Accreditation of Laboratory Animal Care International in accordance with the

Guide for the Care and Use of Laboratory Animals. All procedures for animal use were

approved by the University of California, San Francisco's Institutional Animal Care and Use

Committee.

PLX compound formulation in mouse chow

PLX3397 was provided by Plexxikon Inc. and was formulated in AIN-76A standard chow by

Research Diets Inc. at 275 mg/kg as previously described[80]. PLX5622 was provided by

Plexxikon Inc. and was formulated in AIN-76A standard chow by Research Diets Inc. at 1200

mg/kg as previously described[62]. PLX73086 was provided by Plexxikon Inc. and was formulated

in AIN-76A standard chow by Research Diets Inc. at 200 mg/kg as recommended by Plexxikon

Inc.

Immunohistochemistry and slide scanning

Formalin-fixed samples were embedded into paraffin (FFPE) using standard procedures and

microtome-cut into 8 μm sagittal brain sections or coronal spinal cord sections and mounted

onto slides. To reduce tissue autofluorescence, paraffin slides were photobleached for 48 hours.

Slides were deparaffinized in a 61°C oven for 15 minutes and rehydrated through alcohols.

Antigen retrieval was performed by autoclaving for 10 minutes at 121°C in 0.01 M citrate buffer.

Sections were blocked in 10% normal goat serum (NGS) (Vector Labs) for 1 hour at room

temperature. All primary antibodies were used at 1:250 dilution and included rabbit monoclonal

anti-Iba1 (Abcam, ab178847), rabbit polyclonal anti-P2yr12 (Atlas, HPA014518), mouse

monoclonal anti-NeuN (Millipore, MAB377), chicken anti-GFAP (Abcam, ab4674), and mouse

monoclonal anti-pS202/T205 tau (AT8; Thermo Fisher, MN1020). Primary antibodies were

diluted in 10% NGS in PBS and allowed to incubate on the slides overnight at room

temperature. Primary antibody detection was performed using goat secondary antibodies with

conjugated AlexaFluor 488, AlexaFluor 555, or AlexaFluor 647 (Life Technologies) at 1:500

dilution. Slides were cover-slipped using PermaFluor mounting medium (Thermo). Whole-

section tiled images were acquired with an Axioscan.Z1 slide scanner (Zeiss) at 20✕

magnification, and quantification was performed with Zen 2.3 software (Zeiss).

Cellular bioassay to measure tau-prion activity

A HEK293T cell line expressing the repeat domain of 4R human tau (aa 243–375) containing

the P301L and V337M mutations and C-terminally fused to YFP was previously generated as

described[29]. A stable monoclonal line was maintained in DMEM, supplemented with 10% FBS

and 1% penicillin/streptomycin. To perform the bioassay, 3,000 cells (containing 0.1 µg/ml

Hoechst 33342) were plated in 70 µl per well into 384-well plates (Greiner) and incubated for 2

hours before treatment with samples. Clarified brain lysate at a final concentration of 1.25 µg/mL

total protein was first incubated with Lipofectamine 2000 (0.2% final concentration) and

OptiMEM (9.8% final concentration) for 90 minutes, and then added to the plated cells in

quadruplicate. Plates were incubated at 37°C for 1–3 days, and then the live cells were imaged

using an INCell Analyzer 6000 Cell Imaging System (GE Healthcare) and custom algorithms

were used to detect fluorescent YFP-positive puncta (aggregates).

Mechanical tissue homogenization

Postmortem brains and spinal cords were thawed and weighed to determine the mass in grams.

Brains were bisected into forebrain and hindbrain pieces using a single cut with a scalpel blade

between the striatum and hypothalamus. Tissue was homogenized in nine volumes of cold

DPBS containing Halt Protease Inhibitor Cocktail (1x, Thermo Fisher Scientific) using a

Precellys 24-bead beater (Bertin Instruments) with metal bead lysing matrix (MP Biomedical).

Where necessary, brain lysates were clarified by centrifugation at 10,000 × g for 10 min at 4°C.

All tissue and samples were stored at −80°C until further use.

Formic acid extraction of insoluble proteins in brain tissue for ELISA

Fifty microliters of formic acid were added to 25 µL of 10% brain homogenate and placed in an

ultracentrifuge tube. The samples were vortexed, sonicated for 20 minutes at 37°C in a water-

bath sonicator, and then centrifuged at 100,000 x g for 1 hour. Fifty microliters of supernatant

were recovered to a low-binding tube and neutralized with 950 µL of neutralization buffer (1 M

Tris base and 500 mM dibasic sodium phosphate). Samples were aliquoted into low-binding

tubes and flash frozen in liquid nitrogen. The following ELISA kits from Thermo Fisher Scientific

were used according to the manufacturer's protocols: total tau (KHB0041), p-tau S396

(KHB7031), and p-tau T231 (KHB8051). Each sample was analyzed in duplicate. Raw ELISA

values were adjusted to total brain protein (grams) in the clarified 10% brain homogenate as

determined by bicinchoninic acid (BCA) assay (Pierce/Thermo Fisher Scientific).

Quantification of total protein in brain homogenate

Total protein content in the PBS-soluble (clarified 10% brain homogenate) and detergent-

soluble fractions was quantified using the Pierce BCA Protein Assay Kit (Thermo Fisher

Scientific) following the manufacturer's protocol.

Generation of tau K18*P301L fibrils

Production, purification and fibrillization of recombinant tau K18*P301L fibrils were performed as

previously described[81].

Stereotaxic injections in Tg2541 mice

Forebrain inoculation: Ten-week-old Tg2541 mice received unilateral inoculations of 10 µl of 1.5

mg/ml tau K18 P301L fibrils using stereotaxic methods. Injections followed a two-step process:

the needle was first advanced to the hippocampus (Bregma -2.5mm, Lateral 2.0mm; Depth -2.3

mm from the skull surface) to deliver 5 µl over three minutes, then the Hamilton syringe pump

was paused for five minutes to allow for diffusion prior to retracting the needle to the overlying

cortex (Depth -1.3 mm) where the remaining 5 µl was injected. After fibril injection, the needle

remained in place for five minutes to allow for diffusion of fibrils before retraction, patching the

skull and suturing the scalp. Midbrain inoculation: Ten-week old Tg2541 mice received bi-lateral

inoculations of 10 µl of 1.5 mg/ml tau K18 P301L fibrils using stereotaxic methods. Five

microliters was injected at each site in the midbrain (Bregma, -4.3 mm; Lateral, 1.0 mm, Depth, -

2.5 mm) and (Bregma, -4.3 mm; Lateral, -1.0 mm, Depth, -2.5 mm).

Automated home cage monitoring of behavior

Total activity measurements of freely moving mice were made every 30 days after PLX dosing

in Promethion cages (Sable Systems International). At each time point, mice were first

randomized and placed individually in Promethion cages for 4 to 6 days. Real-time cage activity

recording was continuous during the entire session using a combination of a running wheel with

sensors to measure speed and distance traveled, three balances to measure body weight, food

and water consumption, and a matrix of infrared light beams to measure XYZ movements with

0.25 cm resolution. Analysis of these metrics was used to detect behaviors such as sleep,

rearing and general locomotion. For each mouse, data used for analyses were average

readings per light or dark cycle. Data from the first circadian cycle were excluded due to variable

behavior during habituation. To calculate the activity scores, wheel use, locomotion and rearing

were first normalized to a 0−1 scale by the maximum value in the whole dataset, and then the

geometric mean of the normalized values for each session was calculated.

Quantification of PLX compound levels in brain tissue and plasma

Brain homogenates (20% w/v) were prepared in PBS by one 30-second cycle of bead beating at 5500 rpm with a Precellys 24-bead beater (Bertin Instruments) or plasma samples were prepared by dilution to 25% with PBS. Compounds were recovered by mixing equal parts of brain homogenate with a 50/50 (v/v) solution of acetonitrile (ACN) and methanol containing 1 mM niflumic acid. Precipitated proteins were removed by vacuum filtration (Captiva ND, Agilent). Analysis was performed using a liquid chromatography-tandem mass spectrometry system consisting of an API4500 triple quadra-pole instrument (AB Sciex, Foster City, CA) interfaced with a CBM-20A controller, LC20AD 230 pumps, and a SIL-5000 auto-sampler (Shimadzu Scientific, Columbia, MD). Samples were injected onto a BDS Hypersil C8 column maintained at room temperature. The amount of ACN in the gradient was increased from 75–95% ACN over two minutes, held for one minute, and then re-equilibrated to 75% ACN over 1.4 minutes. Data acquisition used multiple reaction monitoring in the positive ion mode. Specific methods were developed for each compound (PLX3397 and PLX5622), enabling the determination of absolute concentrations.

Blood plasma neurofilament light (NfL) protein measurement using SIMOA

At monthly time points, 150 µl blood was collected in EDTA-coated tubes. The plasma was centrifuged at 1,000 x g for ten minutes to clarify the samples, and was then diluted with sample diluent buffer included in the kit by 25-fold and 100-fold, respectively, prior to the measurement. Plasma NfL concentration was measured and analyzed using the NfL kit (Quanterix) with the SIMOA HD-1 analyzer (Quanterix). Briefly, samples, magnetic beads coated with capture antibody, and biotinylated detector antibodies were combined. Thereafter, the capture beads were resuspended with streptavidin-β-galactosidase (SBG) and resorufin β-D-galactopyranoside (RGP) and transferred to the SIMOA disk. Each bead fit into a microwell in the disk and if NfL was captured then the SBG hydrolysed the RGP substrate which generated a fluorescent signal, and then the concentration was measured against a standard curve derived from known concentrations of recombinant NfL included in the kit. The lower limit of quantification of the assay for plasma was 17.15 pg/mL.

Masson's trichrome and Picosirius red staining

FFPE liver sections (eight µm) were deparaffinized through xylenes and graded alcohols and then rehydrated in distilled water. The Masson's trichrome staining kit (Abcam #ab150686) was

used according to the manufacturer's protocol. The Picro Sirius Red Stain Kit (Abcam

#ab150681) was used to stain tissue sections for 60 min at room temperature. The slides were

rinsed in two changes of acetic acid, three changes of ethanol, and then mounted using

PermaFluor mounting medium (Thermo). Whole-section tiled images were acquired with an

Axioscan.Z1 slide scanner (Zeiss) at 20× magnification, and quantification was performed with

Zen 2.3 software (Zeiss).

Alkaline phosphatase (ALP) ELISA of plasma samples

Mouse plasma samples were diluted 1:100 in the provided dilution buffer and measured using

the ALP ELISA kit (Biovision #E4572-100) according to the manufacturer's protocol.

RNA extraction and Nanostring RNA expression measurements

RNAlater-preserved samples were homogenized in PBS and total RNA was extracted from

samples using the Quick-RNA Miniprep Kit (Zymo Research). RNA extracts were evaluated for

concentration and purity using a Nanodrop 8000 instrument (Thermo Fisher Scientific) and

diluted to a concentration of 20 ng/μl. Hybridizations were performed for the mouse

Neuroinflammation, Myeloid cell, and Neuropathology panels according to the nCounter XT

Assay user manual (Nanostring). The hybridizations were incubated at 65°C for 16 hours, and

then were added to the nCounter SPRINT Cartridge for data collection using the nCounter

SPRINT Profiler. Counts were analyzed using the nSolver Analysis Software.

RNA expression analysis

In total, there were 10 mice in the Tg2541 vehicle group, 10 mice in the Tg2541 PLX5622

group, 6 mice in the wild type vehicle group, and 6 mice in the wild type PLX5622 group. Each

mouse had separate forebrain and hindbrain samples and three panels of Nanostring

sequencing were performed on each sample. Data from the three panels were pooled together

to form the final dataset. When pooling data, if a gene appeared in more than one panel then

the average read value was used in subsequent analysis, unless one panel failed to detect the

gene.

To assign cell-type specificity of each gene, we used the transcriptome dataset reported in a

previous study[82], inspired by previously reported approaches in bulk tissue samples[50]. We set a

specificity threshold in which a gene qualifies to be cell-type specific if its expression in a cell

type is greater than five times the sum in all other cell types. Using this standard, our dataset

had 242 microglia-specific genes, 47 astrocyte-specific genes and 70 neuron-specific genes. All

cell-type specific gene analyses were repeated with a three-time threshold and all results were

consistent (data not shown).

We used partial least-square (PLS) regression (MATLAB) to extract the gene expression pattern

aligned with Tg2541-wild type axis, using individual gene reads from each mouse as predictors

and genotype as responses. Only vehicle groups were used in constructing the PLS regression.

Forebrain and hindbrain were calculated separately. Five output dimensions were chosen for all

PLS analyses, as they covered 99.99% of the total variance in all cases. The scores in the first

two dimensions were plotted. To project PLX3397-treated groups to the PLS dimensions, we

used the following formula:

$Score_{projection} = (Loading_{predictor} \backslash (raw - mean_{predictor})')'$

To calculate population vector distance, we use the "mhal" command in MATLAB. All five

dimensions were used for each mouse. The wild type vehicle group was used as a target.

To calculate the vector angle, each mouse's gene expression pattern was regarded as a five-

dimension vector in the PLS space, and the angle between each mouse and the average vector

of the wild type vehicle group was calculated with the following formula:

$$A = cos^{-1}((u \cdot v)/(|u||v|))$$

To calculate the PLS regression along the PLX concentration and sex-correlated dimensions,

we constructed regressions using all non-microglial genes or only immediate early genes[83] to

measured brain PLX concentrations and sex of each sample. We then calculated variable

importance in projection to isolate the genes important for the regression. To calculate the

projected PLX concentrations, we used the products of gene expression levels and coefficients

estimated from PLS regression.

To calculate clusters in the microglial-specific genes, we calculated pairwise Pearson's

correlation coefficients across 32 samples among each gene. The resulting similarity matrix was

then processed with a generalized Louvain community detection algorithm[52].

To estimate the gene expression levels of the resilient microglia, we used 6 genes (Tmem119,

P2ry12, Fcrls, Olfml3, Itgam v1, and Itgam v2) as microglial-specific house-keeping genes[84] and

used their levels relative to the vehicle-treated group to scale the expression levels of other

microglial genes. We used these estimated expression levels as input for the Ingenuity Pathway

Analysis (QIAGEN).

Gene expression analysis by RT-qPCR

Mouse brains were collected at endpoints and flash frozen in DNA/RNA shield reagent. Tissue
was homogenized as described above and total RNA was purified using a commercial isolation
kit (Zymo Research). RNA concentration and the RNA integrity number (RIN) were determined
using a Bioanalyzer 2100 instrument and an Agilent RNA 6000 Pico Kit (Agilent 5067-1513).
Only samples with a RIN score ≥7.0 were used for gene expression analysis. To confirm
transcriptome profiling results, 2.5 ng of sample mRNA was applied to triplicate RT-qPCR
reactions consisting of 1x TaqPath 1-Step Multiplex Master Mix (ThermoFisher Scientific
A28526), Taqman primer/probe sets and a normalizing human MAPT Taqman assay. Reactions
were run on a QuantStudio 6 and 7 Pro instrument and amplification yielding cycle threshold
(C_T) values were corrected with Mustang Purple passive reference dye for each target gene.
Gene expression of PLX-treated mice relative to vehicle-treated mice was determined by the
comparative C_T method and values were expressed as fold-change.

Comparative C_T equation:

$2-\Delta\Delta C_T$=[(C_T gene of interest – C_T hMAPT internal control)]PLX-treated mice − [(C_T gene of
interest – C_T hMAPT internal control)]vehicle-treated mice

In situ **mRNA hybridization (RNAscope) analysis**

All RNAscope experiments were performed using FFPE tissue sections of age-matched male
and female mice from transgenic tau and wild type control lines. The RNAscope Multiplex
Fluorescent V2 Reagent Kit (#323100) was used according to the manufacturer's instructions,
other than the target retrieval incubation step, which was carried out at 95°C for 20 minutes. The
following probes, all from ACD Bio (Newark, CA), were used: *MAPT* (#522491), *Fos* (#506931-
C2), and *Csf1r* (#428191-C3).

HILIC-MS/MS metabolomics analysis

HILIC-MS/MS analysis was carried out by the West Coast Metabolomics Center at the
University of California, Davis. Briefly, four milligrams of mouse cortex were homogenized with
3.2mm stainless steel beads using a bead-beater in 750 µl of methyl-tertiary butyl ether and 225
µl of methanol. Samples were then vortexed for 10 s, shaken on an orbital shaker at 4°C for 5

min, and then 188 μl of liquid chromatography mass spectrometry (LCMS)-grade water was added. Samples were vortexed again for 20 s and centrifuged for 2 min at 14,000xg. 110 μl of the lower phase (all polar metabolites) were then moved to a new 1.5 ml tube and dried using a centrivap. Dried samples were resuspended in 100 μL buffer containing internal standards and then analyzed using an Agilent 1290 ultra-high pressure liquid chromatography (UHPLC)/Sciex Triple-time-of-flight (TOF) 6600 mass spectrometer. Briefly, a Waters Acquity Premier UHPLC bridged ethylene hybrid (BEH) Amide Column (1.7 μm, 2.1 mm x 50 mm) was used with a mobile phase A of ultrapure water with 10mM ammonium formate (AF) and 0.125% formic acid (FA) (pH 3.0), and a mobile phase B of 95:5 (v/v) acetonitrile:ultrapure water with 10 mM AF + 0.125% FA (pH 3.0). A column temperature of 45°C and a gradient of 0 min, 100% B; 0.5 min, 100% B; 1.95 min, 70% B; 2.55 min, 30% B; 3.15 min, 100% B; 3.8 min, 100% B, were used with a flow rate of 0.8 ml/min.

Following data collection, internal standard chromatograms were examined for consistency of peak height and retention time as quality control measures. Raw data files were then processed using an updated version of the MS-DIAL software[85] using an in-house mzRT library and MS/MS spectral matching with NIST/MoNA libraries. All MS/MS annotations were then manually curated to ensure that only high-quality metabolite identifications were included.

In vivo bioluminescence imaging

Bioluminescence imaging was performed on the brains and spinal cords of albino bigenic Tg(2541:*Gfap*-luc) homozygous mice after receiving an intraperitoneal injection of 25 mg/kg cyclic luciferin-1 (CycLuc1) sodium salt solution (Aobious; AOB6377) prepared in PBS, pH 7.4. After CycLuc1 injection, mice were placed in an anesthetization chamber and exposed to an isoflurane/oxygen gas mix for ten minutes. During this time, the heads of the mice were shaved to enhance the bioluminescence signal. After anesthetization, mice were placed in an IVIS Lumina III small animal imaging system (PerkinElmer) and were kept under constant anesthesia. Mice were imaged for 60 s duration at three time points (14, 16 and 18 minutes) following CycLuc1 injection as determined in one-hour time-lapse calibration studies. After image acquisition, the mice were allowed to recover in their home cages. Brain and spinal cord bioluminescence values were calculated from images displaying surface radiance using standardized regions of interest and were then converted to total photon flux (photons per second) using Living Image software version 4.4 (PerkinElmer).

Confocal imaging of thick tissue sections

Vibratome-sectioned brain slices (40 µm thick) were immunolabeled with Iba1, AT8, and/or

CD206 (Biorad, MCA2235) antibodies using standard protocols for free-floating sections in

multi-well plates. Sections were mounted using PermaFluor and #1.5 coverglass. Using a Leica

SP8 confocal microscope equipped with HyD detectors and an AOBS, samples were first

visualized using Navigator function to acquire an overview image of each slice using a 20×

water-immersion lens (0.95 NA). From the mosaic image, smaller tiled-ROIs were marked in the

forebrain and hindbrain to acquire high-resolution, sequential-scanned image stacks using a

63× water-immersion lens (1.2 NA). Eight-bit image z-stacks (1 µm steps) were collected at

512×512-pixel resolution. Images were processed using custom MATLAB code.

Microglial morphology analysis

Microglia morphology was analyzed using a custom script in MATLAB. Briefly, raw confocal

image stacks were smoothed and then maximally projected. Isolated microglia cells were

manually selected for analysis. The selected microglia region was binarized with an intensity

threshold, and then the cell body was detected by fitting a largest circle in the binary mask. After

excluding the cell body region, the remaining microglia processes were skeletonized and branch

number, branch length and bounding box were measured using "regionprops" and "bwmorph"

commands.

Statistical analysis

Statistical analyses were performed using GraphPad Prism 8. Comparisons between two

groups were performed by two-tailed unpaired t test or by Mann-Whitney nonparametric test.

For comparisons of more than two groups, one-, two-, or three-way ANOVA was performed with

Holm-Šidák post hoc analysis. Following ANOVA, residuals were evaluated for normal

distribution using the Anderson-Darling test and the data were evaluated for equal variance

using the Brown-Forsythe test. If both assumptions were violated ($P < 0.05$), the data was

reanalyzed using Welch's ANOVA with Dunnett T3 post hoc analysis. For repeated-measures

ANOVA, sphericity was not assumed and the Geisser-Greenhouse correction was applied. If

any data points were missing, a mixed-effects model (Restricted maximum likelihood; REML)

was used instead. Pearson's correlation tests were performed as one-tailed tests as, in each

case, we had a directional hypothesis of either positive or negative correlation. Sample sizes

are shown in graphs with each data point representing an individual mouse, or are reported in

the figure legends. Experimental replication and exact statistical tests used are detailed in the

figure legends.

ACKNOWLEDGEMENTS

We thank Plexxikon, Inc for providing the PLX3977, PLX5622 and PLX73086 compounds, and
Andrey Reymar and Brian West for consulting on drug dosing and chow formulation. We thank
Julian Castaneda, Karina Walker and Lyn Batia (and their staff) at the UCSF Hunter's Point
animal facility for coordinating transgenic mouse development, breeding, and drug efficacy
studies. We thank Masahiro Inoue (Daiichi Sankyo, Inc.) for insightful discussions on
pharmacokinetics and pharmacodynamics in our study. We thank Stanley Prusiner and David
Ramsay at the UCSF Institute for Neurodegenerative Diseases (IND) for access to equipment
and technical resources critical for the completion of this study. We thank the following UCSF
IND staff for technical assistance: Abby Oehler, Rigoberto Roman-Albarran, Julia Becker, Marta
Gavidia and Manuel Elepano. The study was funded by grants from the National Institutes of
Health: (# RF1 AG061874), the Rainwater Charitable Foundation, the Sherman Fairchild
Foundation, the Henry M. Jackson Foundation, the Edward N. & Della L. Thome Memorial
Foundation, and Daiichi Sankyo, Inc. P.Y. was funded by Shanghai Municipal Science and
Technology Major Project and Shanghai Natural Science Foundation (22ZR1415000).
Competing interests: The Institute for Neurodegenerative Diseases (UCSF) had a research
collaboration with Daiichi Sankyo, Inc. (Tokyo, Japan).

AUTHOR CONTRIBUTIONS

941   C.C. conceived the study. N.R.J and C.C. designed the experiments. N.R.J., E.C., T.P.L., W.Y.,
942   A.B., B.M.R., M.C.S., H.M., K.G., A.A., and C.C. performed experiments and prepared data.
943   N.R.J., P.Y., and C.C. analyzed and interpreted data. N.R.J., P.Y., and C.C. wrote the paper.
944   C.C. supervised the study.

DATA AVAILABILITY

947   The authors declare that all other data supporting the findings of this study are available within
948   the paper and its supplementary files.

CODE AVAILABILITY

The transcriptome data (Nanostring) that support the findings of this study, specifically in
Figures 6-9, are available from Github with the following link: https://gitfront.io/r/user-
8849465/665dd65fd9d9e78650ed02b9f30236d99240de39/UCSF-PLX-nanostring/

REFERENCES

1. Dionisio-Santos, D.A., Olschowka, J.A. & O'Banion, M.K. Exploiting microglial and peripheral immune cell crosstalk to treat Alzheimer's disease. *J Neuroinflammation* **16**, 74 (2019).
2. Masuda, T.*, et al.* Spatial and temporal heterogeneity of mouse and human microglia at single-cell resolution. *Nature* **566**, 388-392 (2019).
3. Kodama, L. & Gan, L. Do Microglial Sex Differences Contribute to Sex Differences in Neurodegenerative Diseases? *Trends Mol Med* **25**, 741-749 (2019).
4. Stratoulias, V., Venero, J.L., Tremblay, M.E. & Joseph, B. Microglial subtypes: diversity within the microglial community. *EMBO J* **38**, e101997 (2019).
5. Schwabe, T., Srinivasan, K. & Rhinn, H. Shifting paradigms: The central role of microglia in Alzheimer's disease. *Neurobiol Dis* **143**, 104962 (2020).
6. Shi, Y. & Holtzman, D.M. Interplay between innate immunity and Alzheimer disease: APOE and TREM2 in the spotlight. *Nat Rev Immunol* **18**, 759-772 (2018).
7. Maphis, N.*, et al.* Reactive microglia drive tau pathology and contribute to the spreading of pathological tau in the brain. *Brain* **138**, 1738-1755 (2015).
8. Stancu, I.C.*, et al.* Aggregated Tau activates NLRP3-ASC inflammasome exacerbating exogenously seeded and non-exogenously seeded Tau pathology in vivo. *Acta Neuropathol* **137**, 599-617 (2019).
9. Ising, C.*, et al.* NLRP3 inflammasome activation drives tau pathology. *Nature* **575**, 669-673 (2019).
10. Rexach, J.E.*, et al.* Tau Pathology Drives Dementia Risk-Associated Gene Networks toward Chronic Inflammatory States and Immunosuppression. *Cell Rep* **33**, 108398 (2020).
11. Allen, M.*, et al.* Divergent brain gene expression patterns associate with distinct cell-specific tau neuropathology traits in progressive supranuclear palsy. *Acta Neuropathol* **136**, 709-727 (2018).
12. Wang, H.*, et al.* Genome-wide RNAseq study of the molecular mechanisms underlying microglia activation in response to pathological tau perturbation in the rTg4510 tau transgenic animal model. *Mol Neurodegener* **13**, 65 (2018).
13. Matarin, M.*, et al.* A genome-wide gene-expression analysis and database in transgenic mice during development of amyloid or tau pathology. *Cell Rep* **10**, 633-644 (2015).
14. Stanley, E.R. & Chitu, V. CSF-1 receptor signaling in myeloid cells. *Cold Spring Harb Perspect Biol* **6**(2014).

15.   Cannarile, M.A.*, et al.* Colony-stimulating factor 1 receptor (CSF1R) inhibitors in cancer
therapy. *J Immunother Cancer* **5**, 53 (2017).
16.   Asai, H.*, et al.* Depletion of microglia and inhibition of exosome synthesis halt tau
propagation. *Nat Neurosci* **18**, 1584-1593 (2015).
17.   Shi, Y.*, et al.* Microglia drive APOE-dependent neurodegeneration in a tauopathy mouse
model. *J Exp Med* **216**, 2546-2561 (2019).
18.   Mancuso, R.*, et al.* CSF1R inhibitor JNJ-40346527 attenuates microglial proliferation and
neurodegeneration in P301S mice. *Brain* **142**, 3243-3264 (2019).
19.   Bennett, R.E.*, et al.* Partial reduction of microglia does not affect tau pathology in aged
mice. *J Neuroinflammation* **15**, 311 (2018).
20.   Condello, C., DeGrado, W.F. & Prusiner, S.B. Prion biology: implications for Alzheimer's
disease therapeutics. *Lancet Neurol* **19**, 802-803 (2020).
21.   Tap, W.D.*, et al.* Structure-Guided Blockade of CSF1R Kinase in Tenosynovial Giant-Cell
Tumor. *N Engl J Med* **373**, 428-437 (2015).
22.   Spangenberg, E.*, et al.* Sustained microglial depletion with CSF1R inhibitor impairs
parenchymal plaque development in an Alzheimer's disease model. *Nat Commun* **10**,
3758 (2019).
23.   Allen, B.*, et al.* Abundant tau filaments and nonapoptotic neurodegeneration in transgenic
mice expressing human P301S tau protein. *J Neurosci* **22**, 9340-9351 (2002).
24.   Bugiani, O.*, et al.* Frontotemporal dementia and corticobasal degeneration in a family with
a P301S mutation in tau. *J Neuropathol Exp Neurol* **58**, 667-677 (1999).
25.   Goedert, M., Jakes, R. & Crowther, R.A. Effects of frontotemporal dementia FTDP-17
mutations on heparin-induced assembly of tau filaments. *FEBS Lett* **450**, 306-311 (1999).
26.   Sperfeld, A.D.*, et al.* FTDP-17: an early-onset phenotype with parkinsonism and epileptic
seizures caused by a novel mutation. *Ann Neurol* **46**, 708-715 (1999).
27.   Johnson, N.R.*, et al.* Evidence for sortilin modulating regional accumulation of human tau
prions in transgenic mice. *Proc Natl Acad Sci U S A* **114**, E11029-E11036 (2017).
28.   Kovacs, G.G.*, et al.* Distribution patterns of tau pathology in progressive supranuclear
palsy. *Acta Neuropathol* **140**, 99-119 (2020).
29.   Sanders, D.W.*, et al.* Distinct tau prion strains propagate in cells and mice and define
different tauopathies. *Neuron* **82**, 1271-1288 (2014).
30.   Aoyagi, A.*, et al.* Abeta and tau prion-like activities decline with longevity in the Alzheimer's
disease human brain. *Sci Transl Med* **11**(2019).
31.   Elmore, M.R., Lee, R.J., West, B.L. & Green, K.N. Characterizing newly repopulated
microglia in the adult mouse: impacts on animal behavior, cell morphology, and
neuroinflammation. *PLoS One* **10**, e0122912 (2015).
32.   Iba, M.*, et al.* Synthetic tau fibrils mediate transmission of neurofibrillary tangles in a
transgenic mouse model of Alzheimer's-like tauopathy. *J Neurosci* **33**, 1024-1037 (2013).
33.   Seeley, W.W., Crawford, R.K., Zhou, J., Miller, B.L. & Greicius, M.D. Neurodegenerative
diseases target large-scale human brain networks. *Neuron* **62**, 42-52 (2009).
34.   Gustke, N., Trinczek, B., Biernat, J., Mandelkow, E.M. & Mandelkow, E. Domains of tau
protein and interactions with microtubules. *Biochemistry* **33**, 9511-9522 (1994).
35.   Lei, F.*, et al.* CSF1R inhibition by a small-molecule inhibitor is not microglia specific;
affecting hematopoiesis and the function of macrophages. *Proc Natl Acad Sci U S A* **117**,
23336-23338 (2020).
36.   Bellver-Landete, V.*, et al.* Microglia are an essential component of the neuroprotective
scar that forms after spinal cord injury. *Nat Commun* **10**, 518 (2019).
37.   Luo, J.*, et al.* Colony-stimulating factor 1 receptor (CSF1R) signaling in injured neurons
facilitates protection and survival. *J Exp Med* **210**, 157-172 (2013).
38.   Jul, P.*, et al.* Hyperactivity with Agitative-Like Behavior in a Mouse Tauopathy Model. *J
Alzheimers Dis* **49**, 783-795 (2016).

39.   Dumont, M*., et al.* Behavioral deficit, oxidative stress, and mitochondrial dysfunction
1043   precede tau pathology in P301S transgenic mice. *FASEB J* **25**, 4063-4072 (2011).
1044   40.   Wang, X*., et al.* Early intervention of tau pathology prevents behavioral changes in the
1045   rTg4510 mouse model of tauopathy. *PLoS One* **13**, e0195486 (2018).
1046   41.   Bacioglu, M*., et al.* Neurofilament Light Chain in Blood and CSF as Marker of Disease
1047   Progression in Mouse Models and in Neurodegenerative Diseases. *Neuron* **91**, 56-66
1048   (2016).
1049   42.   Ashton, N.J*., et al.* Increased plasma neurofilament light chain concentration correlates
1050   with severity of post-mortem neurofibrillary tangle pathology and neurodegeneration. *Acta*
1051   *Neuropathol Commun* **7**, 5 (2019).
1052   43.   Rojas, J.C*., et al.* CSF neurofilament light chain and phosphorylated tau 181 predict
1053   disease progression in PSP. *Neurology* **90**, e273-e281 (2018).
1054   44.   Benner, B*., et al.* Pexidartinib, a Novel Small Molecule CSF-1R Inhibitor in Use for
1055   Tenosynovial Giant Cell Tumor: A Systematic Review of Pre-Clinical and Clinical
1056   Development. *Drug Des Devel Ther* **14**, 1693-1704 (2020).
1057   45.   Dong, X.X., Wang, Y. & Qin, Z.H. Molecular mechanisms of excitotoxicity and their
1058   relevance to pathogenesis of neurodegenerative diseases. *Acta Pharmacol Sin* **30**, 379-
1059   387 (2009).
1060   46.   Liddelow, S.A*., et al.* Neurotoxic reactive astrocytes are induced by activated microglia.
1061   *Nature* **541**, 481-487 (2017).
1062   47.   Watts, J.C*., et al.* Bioluminescence imaging of Abeta deposition in bigenic mouse models
1063   of Alzheimer's disease. *Proc Natl Acad Sci U S A* **108**, 2528-2533 (2011).
1064   48.   Bellucci, A*., et al.* Induction of inflammatory mediators and microglial activation in mice
1065   transgenic for mutant human P301S tau protein. *Am J Pathol* **165**, 1643-1652 (2004).
1066   49.   Ishizawa, K. & Dickson, D.W. Microglial activation parallels system degeneration in
1067   progressive supranuclear palsy and corticobasal degeneration. *J Neuropathol Exp Neurol*
1068   **60**, 647-657 (2001).
1069   50.   Friedman, B.A*., et al.* Diverse Brain Myeloid Expression Profiles Reveal Distinct Microglial
1070   Activation States and Aspects of Alzheimer's Disease Not Evident in Mouse Models. *Cell*
1071   *Rep* **22**, 832-847 (2018).
1072   51.   Keren-Shaul, H*., et al.* A Unique Microglia Type Associated with Restricting Development
1073   of Alzheimer's Disease. *Cell* **169**, 1276-1290 e1217 (2017).
1074   52.   Mucha, P.J., Richardson, T., Macon, K., Porter, M.A. & Onnela, J.P. Community structure
1075   in time-dependent, multiscale, and multiplex networks. *Science* **328**, 876-878 (2010).
1076   53.   Wang, C*., et al.* Microglial NF-kappaB drives tau spreading and toxicity in a mouse model
1077   of tauopathy. *Nat Commun* **13**, 1969 (2022).
1078   54.   Hong, J*., et al.* Microglial Toll-like receptor 2 contributes to kainic acid-induced glial
1079   activation and hippocampal neuronal cell death. *J Biol Chem* **285**, 39447-39457 (2010).
1080   55.   Heneka, M.T*., et al.* NLRP3 is activated in Alzheimer's disease and contributes to
1081   pathology in APP/PS1 mice. *Nature* **493**, 674-678 (2013).
1082   56.   Krasemann, S*., et al.* The TREM2-APOE Pathway Drives the Transcriptional Phenotype
1083   of Dysfunctional Microglia in Neurodegenerative Diseases. *Immunity* **47**, 566-581 e569
1084   (2017).
1085   57.   Dejanovic, B*., et al.* Changes in the Synaptic Proteome in Tauopathy and Rescue of Tau-
1086   Induced Synapse Loss by C1q Antibodies. *Neuron* **100**, 1322-1336 e1327 (2018).
1087   58.   Yun, S.P*., et al.* Block of A1 astrocyte conversion by microglia is neuroprotective in models
1088   of Parkinson's disease. *Nat Med* **24**, 931-938 (2018).
1089   59.   Kerkhofs, D*., et al.* Pharmacological depletion of microglia and perivascular macrophages
1090   prevents Vascular Cognitive Impairment in Ang II-induced hypertension. *Theranostics* **10**,
1091   9512-9527 (2020).

60.   Liu, Y.*, et al.* Concentration-dependent effects of CSF1R inhibitors on oligodendrocyte
progenitor cells ex vivo and in vivo. *Exp Neurol* **318**, 32-41 (2019).
61.   Zhan, L.*, et al.* A MAC2-positive progenitor-like microglial population is resistant to CSF1R
inhibition in adult mouse brain. *Elife* **9**(2020).
62.   Spangenberg, E.E.*, et al.* Eliminating microglia in Alzheimer's mice prevents neuronal loss
without modulating amyloid-beta pathology. *Brain* **139**, 1265-1281 (2016).
63.   Kodama, L.*, et al.* Microglial microRNAs mediate sex-specific responses to tau pathology.
Nat Neurosci **23**, 167-171 (2020).
64.   Guneykaya, D.*, et al.* Transcriptional and Translational Differences of Microglia from Male
and Female Brains. *Cell Rep* **24**, 2773-2783 e2776 (2018).
65.   Kang, S.S.*, et al.* Microglial translational profiling reveals a convergent APOE pathway
from aging, amyloid, and tau. *J Exp Med* **215**, 2235-2245 (2018).
66.   Colombo, G.*, et al.* A tool for mapping microglial morphology, morphOMICs, reveals brain-
region and sex-dependent phenotypes. *Nat Neurosci* **25**, 1379-1393 (2022).
67.   Hampton, D.W.*, et al.* Cell-mediated neuroprotection in a mouse model of human
tauopathy. *J Neurosci* **30**, 9973-9983 (2010).
68.   Berve, K., West, B.L., Martini, R. & Groh, J. Sex- and region-biased depletion of
microglia/macrophages attenuates CLN1 disease in mice. *J Neuroinflammation* **17**, 323
(2020).
69.   Vinet, J.*, et al.* Neuroprotective function for ramified microglia in hippocampal
excitotoxicity. *J Neuroinflammation* **9**, 27 (2012).
70.   Busche, M.A.*, et al.* Tau impairs neural circuits, dominating amyloid-beta effects, in
Alzheimer models in vivo. *Nat Neurosci* **22**, 57-64 (2019).
71.   Badimon, A.*, et al.* Negative feedback control of neuronal activity by microglia. *Nature* **586**,
417-423 (2020).
72.   Liu, M.*, et al.* Microglia depletion exacerbates acute seizures and hippocampal neuronal
degeneration in mouse models of epilepsy. *Am J Physiol Cell Physiol* **319**, C605-C610
(2020).
73.   Sosna, J.*, et al.* Early long-term administration of the CSF1R inhibitor PLX3397 ablates
microglia and reduces accumulation of intraneuronal amyloid, neuritic plaque deposition
and pre-fibrillar oligomers in 5XFAD mouse model of Alzheimer's disease. *Mol
Neurodegener* **13**, 11 (2018).
74.   Pinto, B.*, et al.* Rescuing Over-activated Microglia Restores Cognitive Performance in
Juvenile Animals of the Dp(16) Mouse Model of Down Syndrome. *Neuron* **108**, 887-904
e812 (2020).
75.   Dagher, N.N.*, et al.* Colony-stimulating factor 1 receptor inhibition prevents microglial
plaque association and improves cognition in 3xTg-AD mice. *J Neuroinflammation* **12**, 139
(2015).
76.   Hillmer, A.T.*, et al.* Microglial depletion and activation: A [(11)C]PBR28 PET study in
nonhuman primates. *EJNMMI Res* **7**, 59 (2017).
77.   Reu, P.*, et al.* The Lifespan and Turnover of Microglia in the Human Brain. *Cell Rep* **20**,
779-784 (2017).
78.   Horti, A.G.*, et al.* PET imaging of microglia by targeting macrophage colony-stimulating
factor 1 receptor (CSF1R). *Proc Natl Acad Sci U S A* **116**, 1686-1691 (2019).
79.   Lee, S.H.*, et al.* Antibody-Mediated Targeting of Tau In Vivo Does Not Require Effector
Function and Microglial Engagement. *Cell Rep* **16**, 1690-1700 (2016).
80.   Elmore, M.R.*, et al.* Colony-stimulating factor 1 receptor signaling is necessary for
microglia viability, unmasking a microglia progenitor cell in the adult brain. *Neuron* **82**,
380-397 (2014).

81. Barghorn, S., Biernat, J. & Mandelkow, E. Purification of recombinant tau protein and
preparation of Alzheimer-paired helical filaments in vitro. *Methods Mol Biol* **299**, 35-51
(2005).
82. Zhang, Y.*, et al.* Purification and Characterization of Progenitor and Mature Human
Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron* **89**,
37-53 (2016).
83. Wu, Y.E., Pan, L., Zuo, Y., Li, X. & Hong, W. Detecting Activated Cell Populations Using
Single-Cell RNA-Seq. *Neuron* **96**, 313-329 e316 (2017).
84. Chiu, I.M.*, et al.* A neurodegeneration-specific gene-expression signature of acutely
isolated microglia from an amyotrophic lateral sclerosis mouse model. *Cell Rep* **4**, 385-
401 (2013).
85. Tsugawa, H.*, et al.* MS-DIAL: data-independent MS/MS deconvolution for comprehensive
metabolome analysis. *Nat Methods* **12**, 523-526 (2015).
1154

FIGURES

 Fig. 1| CSF1R inhibition by three treatment paradigms reduces pathogenic tau levels in

 the brains of Tg2541 mice. a−c, Schematics of acute (**a**), chronic (**b**), or terminal (**c**) PLX

 treatment of Tg2541 mice from 2−4 mo of age, 2−7 mo of age, or 2 mo of age until death,

 respectively. **d**, Sagittal view of the mouse brain divided into two regions: the forebrain,

 containing the cortex, hippocampus, striatum, and olfactory bulb; and the hindbrain, containing

 the thalamus, hypothalamus, midbrain, cerebellum, and brain stem. **e−g**, Tau-prion levels in

 forebrain and hindbrain tissue homogenates of Tg2541 mice receiving acute (**e**), chronic (**f**), or

 terminal (**g**) treatment with vehicle, PLX3397 (275 mg/kg oral), or PLX5622 (1200 mg/kg oral),

 measured using the HEK293T cell tau-prion bioassay and normalized to the vehicle-treated

 group. **h−j**, Levels of pTau [S396] measured by ELISA in formic acid extracts of forebrain and

 hindbrain tissue homogenates of Tg2541 mice receiving acute (**h**), chronic (**i**), or terminal

 treatment (**j**) with vehicle, PLX3397, or PLX5622, normalized to total protein concentration.

 k−m, Quantification of pTau [S202/T205]-positive area by IHC analysis of forebrain and

 hindbrain areas of Tg2541 mice receiving acute (**k**), chronic (**l**), or terminal (**m**) treatment with

vehicle, PLX3397, or PLX5622. Welch ANOVA with Dunnett T3 post hoc testing was used in **e**,

f, **h**, **i**, and **l**. Two-way ANOVA with Holm-Šidák post hoc testing was used in **g**, **j**, **k**, and **m**. *P*

values for all statistically significant differences ($P < 0.05$) are shown. **n**, Principal component

analysis was performed, using all tau-prion and pTau[S396] data presented in Fig. 1e−j to

calculate a 'tau score' that represents the amount of pathogenic tau in both the forebrains and

hindbrains of Tg2541 mice. All data was first standardized to the respective vehicle-treated

group of the same sex and same dosing paradigm. Then, two principal components (PC1 and

PC2) were identified which accounted for 70.7% of the total variance in the data. Multiple linear

regression was performed on PC1 of the drug-treated groups to evaluate the main effects sex

and dosing paradigm, and the dosing*sex interaction effect. Multiple linear regression was

performed on all groups to determine the main effect of drug and the drug*sex interaction effect.

P values for all statistically significant differences ($P < 0.05$) are shown. n.s. indicates not

statistically significant. PC2 was also evaluated, but only the drug main effect was statistically

significant. In **e−n**, each symbol represents the forebrain or hindbrain of an individual mouse,

with female mice shown as closed or open circles and male mice shown as closed or open

triangles. Error bars represent the s.d. of the mean.

1188

Fig. 2| CSF1R inhibition extends survival of female Tg2541 mice. a, Schematic of terminal
PLX3397 treatment (275 mg/kg oral) of Tg2541 mice from 2 mo of age until death. **b**, Kaplan-
Meier plot showing percent survival of female or male Tg2541 mice treated with vehicle or
PLX3397. n=7 mice for Female Vehicle; n=6 mice for Female PLX3397; n=9 mice for Male
Vehicle; n=9 mice for Male PLX3397. **c**, Body weights of female or male Tg2541 mice treated
with vehicle or PLX3397. Differences in weight between vehicle and PLX3397 treatment in
female or male mice were evaluated by mixed-effects analysis (Restricted maximum likelihood).
Each symbol represents an individual mouse and lines indicate group means. **d**, **e**, Correlation
plots for body weight at 180 d of age (**d**) or forebrain tau-prion activity at death (**e**) and survival
for female or male Tg2541 mice treated with vehicle or PLX3397. Each symbol represents an
individual mouse and linear regression lines are shown for female or male mice, with vehicle-
and PLX3397-treated mice combined. Pearson's correlation analysis was performed and the

results are shown. **f**, Schematic of terminal PLX3397 treatment of Tg2541 mice from 2.5 mo of

age until death, following inoculation of K18 tau fibrils into the midbrain (hindbrain region) at 2.5

mo of age. **g**, Kaplan-Meier plot showing percent survival of female or male Tg2541 mice

inoculated with K18 tau fibrils or diluent. n=8 mice for Female Vehicle Diluent; n=7 mice for

Female Vehicle K18; n=8 mice for Male Vehicle Diluent; n=7 mice for Male Vehicle K18.

Differences in survival between diluent and K18 inoculation in male or female mice treated with

vehicle were evaluated by Log-rank (Mantel-Cox) test. **h**, Kaplan-Meier plot showing percent

survival of female or male Tg2541 mice inoculated with K18 tau fibrils and then receiving

terminal treatment of vehicle or PLX3397 (275 mg/kg oral). n=7 mice for Female Vehicle K18;

n=6 mice for Female PLX3397 K18; n=7 mice for Male Vehicle K18; n=8 mice for Male

PLX3397 K18. In **b**, **g**, and **h**, differences in survival between treatment groups were evaluated

by Log-rank (Mantel-Cox) test.

1213

Fig. 3| PLX3397 has sex-dependent pharmacokinetics and reduces hyperactivity in Tg2541 mice. a, Schematic of terminal PLX3397 treatment (275 mg/kg oral) of Tg2541 mice from 2 mo of age until death and blood plasma collected at 3, 4, 5, and 6 mo of age. **b**, Plasma concentration of PLX3397 in female or male Tg2541 mice. Each symbol represents an individual mouse and the lines indicate group means. The difference between female and male mice was assessed by two-way repeated measures ANOVA. **c**, **d**, Correlation plots for plasma concentration of PLX3397 at 90 d age and Iba1 area fraction by IHC (**c**) or tau-prion activity (**d**) in the forebrains or hindbrains of Tg2541 mice at death. Female mice are shown as open circles and male mice shown as open triangles. Linear regression was performed with female and male mice combined and best-fit lines are shown. Pearson's correlation analysis was performed and

the results are shown. **e**, Longitudinal energy expenditure indices (see Methods) of female or

male Tg2541 mice treated with vehicle or PLX3397. Symbols represent the group means and

shaded regions indicate the s.d. of the mean. Group sizes are the same as shown in **f** and **g**.

Differences between vehicle and PLX3397 treatment in female or male mice were assessed by

three-way repeated measures ANOVA. **f**, **g**, Average distance traveled on the running wheel (**f**)

or still time during the dark cycle (**g**) in female or male Tg2541 mice treated with vehicle or

PLX3397, and measured between 90-d-old and 150-d-old. Each symbol represents an

individual mouse and the dashed lines indicate the same measurements in 90-d-old wild type

mice. Differences between vehicle and PLX3397 treatment in female or male mice were

assessed by Mann-Whitney test.

1234

Fig. 4| Plasma NfL levels are reduced by PLX3397 in female Tg2541 mice and increased in male mice. a, Schematic of terminal PLX3397 treatment (275 mg/kg oral) of Tg2541 mice from 2 mo of age until death and blood plasma collected at 3, 4, 5, and 6 mo of age. **b**, **c**, Plasma concentration of NfL in female (**b**) or male (**c**) Tg2541 mice treated with vehicle or PLX3397. **d**, Schematic of terminal PLX3397 treatment of Tg2541 mice from 2.5 mo of age until death, following inoculation of K18 tau fibrils into the midbrain (hindbrain region) at 2.5 mo of age, and blood plasma collected monthly thereafter until death. **e**, **f**, Plasma concentration of NfL in female (**e**) or male (**f**) Tg2541 mice inoculated with K18 followed by terminal treatment

with vehicle or PLX3397, plotted over days post-inoculation (dpi). **g**, **h**, Correlation plots for plasma NfL concentration and survival in female (**g**) or male (**h**) Tg2541 mice. Each symbol represents an individual mouse. **i**, **j**, Correlation plots for brain concentration of PLX3397 and NfL concentration in the (**i**) forebrains or (**j**) hindbrains of Tg2541 mice receiving midbrain inoculation with diluent. Each symbol represents the forebrain or hindbrain of an individual mouse. In **g**−**j**, linear regression and Pearson's correlation analysis were performed and the results are shown. **k**, Schematic of intermittent PLX treatment of Tg2541 mice from 2−7 mo of age, with three weeks on treatment followed by three weeks off of treatment, and blood plasma collected at 3, 4, 5, and 6 mo of age. **l**, **m**, Plasma concentration of NfL in female (**l**) or male (**m**) Tg2541 mice receiving intermittent treatment with vehicle or PLX3397. **n**, Schematic of terminal PLX treatment of C57BL/6J wild type mice (Wt) from 2 mo of age until death and blood plasma collected at 3, 4, 5, 6, and 7 mo of age. **o**, **p**, Plasma concentration of NfL in female (**o**) or male (**p**) Wt mice treated with vehicle or PLX3397. In **b**, **c**, **e**, **f**, **l**, **m**, **o**, and **p**, the differences between vehicle or PLX3397 treatment were evaluated by non-linear regression using exponential growth models (**b**, **c**, **l**, **m**, **o**, and **p**) or third-order polynomial models (**e** and **f**) and the best-fit lines and statistical results are shown. Each symbol represents an individual mouse.

Fig. 5| Sex-dependent effects of CSF1R are not caused by peripheral toxicity. a,
Schematic of chronic treatment of Tg2541 mice from 2−7 mo of age with PLX73086 (200 mg/kg
oral), a non-brain penetrant analog of PLX3397 and PLX5622. **b**, **c**, Plasma concentration of
NfL in female (**b**) or male (**c**) Tg2541 mice treated with vehicle or PLX73086. **d**, Body weights
of female or male T2541 mice treated with PLX73086. **e**, Schematic of chronic treatment of wild
type (Wt) mice from 2−7 mo of age with PLX73086 (200 mg/kg oral). **f**, **g**, Plasma concentration
of NfL in female (**f**) or male (**g**) Wt mice treated with vehicle or PLX73086. **h**, Body weights of
female or male wild type mice treated with PLX73086. In **b**, **c**, **f**, and **g**, the differences between
vehicle or PLX3397 treatment were evaluated by non-linear regression using quadratic models.

Each symbol represents an individual mouse and the best-fit lines and statistical results are

shown. In **d** and **h**, differences in weight between vehicle and PLX73086 treatment in male or

female mice were evaluated by two-way repeated measures ANOVA and *P* values are shown.

Each symbol represents an individual mouse and lines indicate group means. **i**, Schematic of

terminal treatment of Tg2541 mice from 2 mo of age until death with PLX3397. **j**, Representative

histopathology images of liver sections of Tg2541 mice receiving terminal treatment with vehicle

or PLX3397, stained with hematoxylin and eosin (H&E), Masson's trichrome, or Picosirius red.

High magnification insets are shown of the regions outlined with a white box. Scale bars, 200

μm. **k**, Sirius red-stained liver sections of Tg2541 mice that received acute or terminal treatment

with vehicle or PLX3397 were quantified for percent positive area. The acute and terminal

treatment groups were combined for the analysis due to a limited sample size of terminal

treatment groups and because the group means were similar for the two treatment paradigms.

Mice receiving terminal treatment are shown as black symbols and mice receiving acute

treatment are shown as purple or green symbols. **l**, Alkaline phosphatase (ALP) levels were

measured at eight months of age in the plasma of Tg2541 mice receiving midbrain inoculation

of diluent at 2.5 mo of age and then treated with vehicle or PLX3397 until death. The data are

presented as a percent change from ALP levels at two months of age in the same mice. In **k**

and **l**, the differences between vehicle and PLX3397 treatment were evaluated by ANOVA with

Holm-Šidák post hoc analysis and *P* values are shown. Each symbol represents an individual

mouse and error bars indicate the s.d. of the mean.

1289

Fig. 6| PLX treatment normalizes gene expression profile, but dose-dependent

excitotoxicity occurs with higher drug exposure. a, Pearson's correlations among individual

mice for non-microglia gene expression patterns (1,599 genes for each mouse). Mice from the

same treatment group (black boxes) show a high degree of correlations. Dashed boxes show

correlations of Tg2541 mice and wild type vehicle group mice. **b**, Distributions of correlation

coefficients in the two groups in dashed boxes in **a**. **c**, Partial least square (PLS) regression

scores and projections of non-microglia gene expression patterns in different groups (1,599

genes for each mouse). The first two dimensions are shown covering >95% of the total

variance. Vehicle-treated mice (filled dots) were used for regression (five dimensions covering

99.99% of the total variance) and PLX5622-treated (1200 mg/kg oral) mice (empty circles) were

projected onto the regression dimensions. **d**, Mahalanobis population vector distances of gene

expression patterns in PLS dimensions relative to the wild type vehicle group. Mann-Whitney

test was used for statistical comparison. **e**, Population vector angles of gene expression

patterns in PLS dimensions from different treatment groups. **f**, Example plots of measured brain

PLX concentration against mRNA counts for a PLX-modulated gene (*Fos*) and a sex-modulated

gene (*Uty*). Linear regression and Pearson's correlation analysis were performed and the

results are shown. N/A indicates that there was no detectable expression of the *Uty* gene in

female mice. See Supplementary Data File 2 for statistical comparisons between male and

female PLX-treated mice of all genes. **g**, Variable importance in projection (VIP) scores in a PLS

regression using non-microglial gene expression patterns to brain PLX concentration and sex.

Genes with VIP scores above 2 are labeled. Red fonts indicate a subset that belongs to

immediate early genes (IEGs). **h**, Scatter plot of measured vs. projected brain PLX

concentration, which was based on a PLS regression with IEG expression. **i**, Correlation of

brain PLX concentration and the population vector distances between PLX-treated individual

mouse transcriptome pattern to WT, with and without the IEGs (1,599 and 1,543 genes for each

mouse, respectively). Linear regressions were calculated for each group. **j**, Quantification of

normalized expression levels from all IEGs (56 genes for each mouse) in the forebrain. Mann-

Whitney tests were used for comparing between groups. **k**, Representative confocal images of

in situ labeling of *MAPT* and *Fos* mRNA (RNAscope) in the forebrains of male and female

Tg2541 mice treated with PLX3397 from 2 mo of age until death. Zoomed regions of interest are

shown to the right. Arrows indicate labeled *Fos* puncta. **l**, Quantification of *Fos* mRNA levels

measured in RNAscope images shown in **k**. Mann-Whitney tests were used for comparing

between groups. In **c**, **f**, **h−j**, and **l**, each symbol represents an individual mouse.

1323

Fig. 7| PLX dysregulates neurotransmitter levels in the brains of male Tg2541 mice. **a**,
Glutamate levels in forebrain lysates of wild type (Wt) or Tg2541 (Tg) mice, dosed for 6-12
weeks with PLX3397 (275 mg/kg oral) or vehicle beginning at 4 months of age, were measured
by hydrophilic interaction liquid chromatography tandem mass spectrometry (HILIC-MS/MS). **b**,
γ-Aminobutyric acid (GABA) levels in forebrain lysates of Wt or Tg mice, dosed for 6-12 weeks
with PLX3397 (275 mg/kg oral) or vehicle beginning at 4 months of age, were measured HILIC-
MS/MS. In **a** and **b,** Mann-Whitney tests were used for comparing vehicle and PLX3397
treatment groups within mouse genotype and sex, and the *P* values of statistically significant
differences (*P* < 0.05) are shown. Each symbol represents an individual mouse and error bars
indicate the s.d. of the mean.

Fig. 8| CSF1R inhibition ameliorates tau-induced pathological astrocyte activation. **a**, **b**,

Analyses similar to Fig. 6c−e using astrocyte-specific genes (47 genes in each mouse) in (**a**)

forebrain and (**b**) hindbrain regions. For other cell types and brain regions, see Supplementary

Fig. 12. **c**, Quantifications of featured astrocyte genes in different conditions. Data are

represented as mean ± S.D. T-tests were used to compare vehicle and PLX-treated groups for

each gene, with 5% false discovery rate correction for multiple comparisons. **d**, Correlation of

brain PLX concentration and different groups of astrocyte gene expression. Based on a prior

study[46], A1 genes are associated with neurotoxic astrocytes following lipopolysaccharide

exposure and A2 genes are associated with neuroprotective function in an artery occlusion

model. Linear regressions were calculated for each group. **e**, Representative images of *in vivo*

bioluminescence imaging of GFAP activity in Tg2541 mice with vehicle or PLX3397 treatment.

Arrows indicate hindbrain regions. **f**, **g**, Quantifications of longitudinal measurements of

astrogliosis-driven bioluminescence in the (**f**) forebrains or (**g**) hindbrains of Tg2541 mice with

vehicle or PLX3397 treatment. Differences in BLI signal between vehicle and PLX treatment

were evaluated by mixed-effects analysis (Restricted maximum likelihood). In **a**, **b**, **d**, **f**, and **g**

each symbol represents an individual mouse.

1351

Fig. 9| PLX3397 treatment preferentially eliminates reactive microglia around tau

deposits. a, Representative confocal images of immunostaining of tau protein (magenta) and

microglia (Iba1, green) in the brain of a vehicle-treated (left) or PLX3397-treated (right) 210-d-

old Tg2541 mouse. **b**, Representative confocal images, similar to panel **a**, of a vehicle-treated

210-d-old Tg2541 mouse. The right panel shows the zoomed image from the dashed box in the

left panel. The white lines in the right panel show distances from the tau deposit at the center. **c**,

Quantification of microglial densities at different distances from the nearest tau deposit in

Tg2541 mice treated with vehicle or PLX3397 (275 mg/kg oral). Two-way ANOVA was used to

compare statistical differences between treatment groups and distance bins. **d**, Representative

confocal images of microglial processes labeled by Iba1 immunohistochemistry. **e**−**g**,

Quantification of microglial processes branch numbers (**e**), total lengths (**f**) and territory sizes (**g**)

in Tg2541 mice treated with vehicle or PLX3397. Each data point shows the average value of

the microglia measured in an individual mouse. Error bars represent the s.d. of the mean.

Circles indicate female mice and triangles indicate male mice. Dotted lines show the average

measurements from microglia in wild type mice. Mann-Whitney tests were used to compare

between groups. The forebrain regions of PLX3397-treated male and female mice were also

compared directly and the *P* values are shown on each plot.

1369

Fig. 10| Selective ablation of tau-activated microglia gene expression by PLX5622. a,

Volcano plot of gene expression changes between Tg2541 and wild type mice. Many microglial-

specific genes (magenta dots) show trends of up-regulation in Tg2541 mice. **b**, Schemaball

graphs of microglial-specific genes comparing Tg2541 to wild type. Individual circle sizes

indicate fold changes. Microglial-specific genes (242 genes in our dataset) are clustered into

three groups: red, green and blue. Connecting arcs between genes represent the degree of

correlation. The pie charts on the right show the percentage of genes in each group that are

differentially expressed in Tg2541 and wild type mice, and the Venn diagrams indicate the brain

regions of differential expression. **c**, Heatmap showing expression level changes in DAM

genes[51]. Gene names are color coded to show their group assignments. **d**, Schemaball graphs

of microglial-specific genes comparing Tg2541 vehicle to PLX5622 treatment (1200 mg/kg oral)

in male and female mice. **e**, Estimated gene expression levels in resilient microglia comparing

between genders and treatment. Each dot represents a different gene, and is color-coded

according to the cluster to which it belongs. **f**, Real-time PCR quantification of example

microglial genes *Gadd45b* and *Csf1r* comparing vehicle to PLX5622 treatment in Tg2541 mice.

Bars indicate medians and student's t-tests were used to compare groups. **g**, Representative

images and quantification of *Csf1r* gene expression by *in situ* mRNA hybridization (RNAscope)

analyses of Tg2541 mice treated with PLX3397 from 2 mo of age until death. Bars indicate

medians and two-way ANOVA was used to compare groups. **h**, Ingenuity Pathway Analysis of

the gene expression patterns in the resilient microglia in male and female mice. Paired t-tests

were used in each category to compare sexes.

SUPPLEMENTARY INFORMATION

1392

Supplementary Fig. 1| Microglial depletion by CSF1R inhibitors in Tg2541 and wild type

mice. a, Sagittal and superior diagrams of the mouse brain divided into two regions: the
forebrain, containing the cortex, hippocampus, striatum, and olfactory bulb; and the hindbrain,
containing the thalamus, hypothalamus, midbrain, cerebellum, and brain stem. **b−d**, Schematics
of acute (**b**), terminal (**c**), or chronic (**d**) PLX treatment of Tg2541 mice from 2−4 mo of age, 2
mo of age until death, or 2−7 mo of age, respectively. **e−g**, Quantification of the Iba1-positive
area fraction by IHC in the forebrains or hindbrains of Tg2541 mice receiving acute (**e**), terminal
(**f**), or chronic (**g**) treatment with vehicle, PLX3397 (275 mg/kg oral), or PLX5622 (1200 mg/kg
oral). **h−j**, Quantification of the P2yr12-positive area fraction by IHC in the forebrains or
hindbrains of Tg2541 mice receiving acute (**h**), terminal (**i**), or chronic (**j**) treatment with vehicle,
PLX3397, or PLX5622. **k**, **l**, Schematics of acute (**k**) or terminal (**l**) PLX treatment of C57BL/6J
wild type mice (Wt) from 2−4 mo of age, or 2 mo of age until death, respectively. **m**, **n**,
Quantification of the Iba1-positive area fraction by IHC in the forebrain and hindbrain of Wt mice
receiving acute (**m**) or terminal (**n**) treatment with vehicle, PLX3397, or PLX5622. **o**, **p**,
Quantification of the P2yr12-positive area fraction by IHC in the forebrains or hindbrains of Wt
mice receiving acute (**o**) or terminal (**p**) treatment with vehicle, PLX3397, or PLX5622. *P* values
for all statistically significant differences ($P < 0.05$) are shown. **q**, Principal component analysis
was performed, using all data presented in **e−j** and **m−p** to calculate a 'microglial score' that
represents the amount of Iba1 and P2yr12 staining in both the forebrains and hindbrains of
Tg2541 and Wt mice. All data was first standardized to the respective vehicle-treated group of
the same sex and same dosing paradigm. Then, two principal components (PC1 and PC2) were
identified which accounted for 79.0% of the total variance in the data. Multiple linear regression
was performed on PC1 of the drug-treated groups to evaluate the main effects sex and dosing
paradigm, and the dosing*sex interaction effect. Multiple linear regression was performed on all
groups to determine the main effect of drug and the drug*sex interaction effect. *P* values for all
statistically significant differences ($P < 0.05$) are shown. n.s. indicates not statistically significant.
PC2 was also evaluated, but only the drug main effect was statistically significant. **r**, Schematic
of intermittent PLX treatment of Tg2541 mice from 2−7 mo of age, with three weeks on
treatment followed by three weeks off of treatment. **s**, **t**, Quantification of the Iba1-positive (**s**) or
P2yr12-positive (**t**) area fractions by IHC in the forebrains or hindbrains of Tg2541 mice
receiving intermittent treatment with vehicle or PLX3397. In **e−j**, **m−p**, **s**, and **t**, each symbol
represents the forebrain or hindbrain of an individual mouse, with female mice shown as closed
or open circles and male mice shown as closed or open triangles. Error bars represent s.d. of
the mean. Two-way ANOVA with Holm-Šidák post hoc testing was used in **e**, **f**, **h**, **i**, **m−p**, **s**, and
t. Welch ANOVA with Dunnett T3 post hoc testing was used in **g** and **j**.

1428

Supplementary Fig. 2| Optimization of HEK293T cell bioassay for measuring tau-prions in

Tg2541 mouse brain homogenates. a−c, Percent of HEK293T cells expressing YFP-tau-

RD*P301L/V337M with tau aggregates at one (**a**), two (**b**), or three (**c**) days post-infection with

Tg2541 mouse forebrain or hindbrain homogenates at concentrations ranging from 0.0075−7.5

µg/ml. Error bars represent s.e.m. of three terminal Tg2541 mice. **d**, Percent of cells infected

with tau aggregates over time following infection with 0.075, 0.25, or 0.75 µg/ml Tg2541 mouse

forebrain or hindbrain homogenates. Lines represent the means of three terminal Tg2541 mice.

e, Tau-prion levels in 0.25 µg/ml forebrain or hindbrain homogenates of Tg2541 (Tg) or wild

type (Wt) mice, measured using the optimized HEK293T cell tau-prion bioassay. Each symbol

represents the forebrain or hindbrain of an individual mouse and female mice are shown as

closed circles while male mice are shown as closed triangles. Two-way ANOVA with Holm-

Šidák post hoc testing was used and *P* values for all statistically significant differences (*P* <

0.05) are shown.

1442

Supplementary Fig. 3| CSF1R inhibition reduces microglia and pathogenic tau levels in the spinal cords of Tg2541 mice. a, Schematic of chronic PLX treatment of Tg2541 mice from 2−7 mo of age. **b**, Quantification of the Iba1-positive area fraction by IHC in the spinal cords of Tg2541 mice receiving chronic treatment with vehicle or PLX3397 (275 mg/kg oral). **c**, Tau-prion levels in spinal cord tissue homogenates of Tg2541 mice receiving chronic treatment with vehicle or PLX3397, measured using the HEK293T cell tau-prion bioassay and normalized to the vehicle group. **d**, Quantification of pTau [S202/T205]-positive area by IHC in the spinal cords of Tg2541 mice receiving chronic treatment with vehicle or PLX3397. In **b−d**, each symbol represents an individual mouse, with female mice shown as closed or open circles and male mice shown as closed or open triangles. Error bars represent s.d. of the mean. Unpaired t tests were used and *P* values for all statistically significant differences (*P* < 0.05) are shown.

1454

Supplementary Fig. 4| Interventional or intermittent CSF1R inhibition reduces pathogenic

tau levels in the brains of Tg2541 mice. a, Schematic of interventional PLX treatment of

Tg2541 mice from 4−7 mo of age. **b**, Tau-prion levels in forebrain and hindbrain tissue

homogenates of Tg2541 mice receiving interventional treatment with vehicle or PLX3397 (275

mg/kg oral), measured using the HEK293T cell tau-prion bioassay and normalized to the vehicle

group. **c**, Levels of pTau [S396] measured by ELISA in formic acid extracts of forebrain and

hindbrain tissue homogenates of Tg2541 mice receiving interventional treatment with vehicle or

PLX3397, normalized to total protein concentration. **d**, Levels of pTau [T231] measured by

ELISA in formic acid extracts of forebrain and hindbrain tissue homogenates of Tg2541 mice

receiving interventional treatment with vehicle or PLX3397, normalized to total protein

concentration. **e**, Schematic of intermittent PLX treatment of Tg2541 mice from 2−7 mo of age,

with three weeks on treatment followed by three weeks off of treatment. **f**, Tau-prion levels in

forebrain and hindbrain tissue homogenates of Tg2541 mice receiving intermittent treatment

with vehicle or PLX3397, measured using the HEK293T cell tau-prion bioassay and normalized

to the vehicle group. **g**, Levels of pTau [S396] measured by ELISA in formic acid extracts of

forebrain and hindbrain tissue homogenates of Tg2541 mice receiving intermittent treatment

with vehicle or PLX3397, normalized to total protein concentration. **h**, Levels of pTau [T231]

measured by ELISA in formic acid extracts of forebrain and hindbrain tissue homogenates of

Tg2541 mice receiving intermittent treatment with vehicle or PLX3397, normalized to total

protein concentration. In **b−d** and **f−h**, each symbol represents the forebrain or hindbrain of an

individual mouse, with female mice shown as closed or open circles and male mice shown as

closed or open triangles. Error bars represent s.d. of the mean. Two-way ANOVA with Holm-

Šidák post hoc testing was used and *P* values for all statistically significant differences (*P* <

0.05) are shown.

1479

Supplementary Fig. 5| CSF1R inhibition reduces pathogenic tau spreading following K18

forebrain inoculation. a, Schematic of acute PLX treatment of Tg2541 mice from 2.5−4 mo of

age following inoculation of K18 tau fibrils into the hippocampus and overlying cortex (forebrain

regions) at 2.5 mo of age. **b**, Tau-prion levels in the ipsilateral (inoculated side) or contralateral

(un-inoculated side) forebrain or hindbrain tissue homogenates of Tg2541 mice receiving acute

treatment with vehicle or PLX3397 (275 mg/kg oral) following K18 forebrain inoculation,

measured using the HEK293T cell tau-prion bioassay and presented as percent of cells with tau

aggregates. Horizontal lines across bars indicate the mean tau-prion level in forebrain (12.54%

cells with aggregates) or hindbrain (26.06% cells with aggregates) tissue homogenates of

Tg2541 mice that did not undergo K18 inoculation and received acute treatment with vehicle

(Un-inoc. Veh). Each symbol represents the ipsilateral or contralateral forebrain or hindbrain of

an individual mouse, with female mice shown as closed or open circles and male mice shown

as closed or open triangles. Error bars represent s.d. of the mean. Three-way ANOVA with

Holm-Šidák post hoc testing was used and P values for all statistically significant differences (P

< 0.05) between Veh and 3397 are shown above the bars. P values for all differences between

each group and the respective brain region of Un-inoc. Veh mice are shown on the bars.

1496

Supplementary Fig. 6| Non-brain penetrant analog of CSF1R inhibitors does not reduce microglia or pathogenic tau levels in the brains of Tg2541 mice. a, Schematic of chronic treatment of Tg2541 mice from 2−7 mo of age with PLX73086, a non-brain penetrant analog of PLX3397 and PLX5622. **b**, **c**, Quantification of the Iba1-positive (**b**) or P2yr12-positive (**c**) area fractions by IHC in the forebrains or hindbrains of Tg2541 mice receiving chronic treatment with vehicle or PLX73086 (200 mg/kg oral). **d**, Tau-prion levels in forebrain and hindbrain tissue homogenates of Tg2541 mice receiving chronic treatment with vehicle or PLX73086, measured

using the HEK293T cell tau-prion bioassay and normalized to the vehicle group. **e**, **f**, Levels of

pTau [S396] (**e**) or pTau [T231] (**f**) measured by ELISA in formic acid extracts of forebrain and

hindbrain tissue homogenates of Tg2541 mice receiving chronic treatment with vehicle or

PLX73086, normalized to total protein concentration. In **b−f**, each symbol represents the

forebrain or hindbrain of an individual mouse, with female mice shown as closed or open circles

and male mice shown as closed or open triangles. Error bars represent s.d. of the mean. Two-

way ANOVA with Holm-Šidák post hoc testing was used and P values for all statistically

significant differences ($P < 0.05$) are shown.

1512

Supplementary Fig. 7| Pharmacological CSF1R inhibition does not deplete perivascular

macrophages. a, Representative immunofluorescence images of Tg2541 mouse cortical blood

vessels labeled for perivascular macrophages (Iba1$^+$/CD206$^+$) and microglia (Iba1$^+$/CD206$^-$).

Iba1 labels both PVMs and microglia. Mice were treated with PLX3397 from 2.5 mo of age until

death, following inoculation of K18 tau fibrils into the midbrain (hindbrain region) at 2.5 mo of

age. Scale bar, 100 μm. **b**, Quantification of Iba1$^+$/CD206$^+$ cell density in images of Tg2541

mouse cortical blood vessels. Each symbol represents the average of six blood vessel images

from each individual mouse. Error bars represent s.d. of the mean. Two-way ANOVA was used

and the *P* values for Holm-Šidák post hoc testing between treatment groups are shown.

1522

Supplementary Fig. 8| CSF1R inhibition by three treatment paradigms does not affect

neurons or total tau levels. a−c, Schematics of acute (**a**), chronic (**b**), or terminal (**c**) PLX

treatment of Tg2541 mice from 2−4 mo of age, 2−7 mo of age, or 2 mo of age until death,

respectively. **d−f**, Quantification of neuronal nuclei (NeuN)-positive area fraction by IHC analysis

of forebrain and hindbrain areas of Tg2541 mice receiving acute (**d**), chronic (**e**), or terminal (**f**)

treatment with vehicle, PLX3397 (275 mg/kg oral), or PLX5622 (1200 mg/kg oral). **g−i**, Levels of

total tau measured by ELISA in forebrain and hindbrain tissue homogenates of Tg2541 mice

receiving acute (**g**), chronic (**h**), or terminal treatment (**i**) with vehicle, PLX3397, or PLX5622,

normalized to total protein concentration. In **d−i**, each symbol represents the forebrain or

hindbrain of an individual mouse, with female mice shown as closed or open circles and male

mice shown as closed or open triangles. Error bars represent s.d. of the mean. Two-way

ANOVA with Holm-Šidák post hoc testing was used in **d**, and **f−i**. One-way ANOVA with Holm-

Šidák post hoc testing was used in **e**. P values for all statistically significant differences ($P <$

0.05) are shown.

1537

Supplementary Fig. 9| PLX3397 levels are increased in male Tg2541 and wild type mice.

a, Schematic of terminal PLX3397 treatment of Tg2541 mice from 2.5 mo of age until death,

following inoculation of K18 tau fibrils into the midbrain (hindbrain region) at 2.5 mo of age.

Blood plasma was collected at monthly intervals from 3−9 mo of age. **b**, Plasma concentration

of PLX3397 in female or male Tg2541 mice inoculated with K18 followed by terminal treatment

with PLX3397, plotted over days post inoculation (dpi). Mixed-effects analysis (Restricted

maximum likelihood) was used to compare female and male mice and the *P* value is shown. **c**,

Brain concentration of PLX3397 at death in female or male Tg2541 mice inoculated with K18

followed by terminal treatment with PLX3397. Unpaired t test was used to compare female and

male mice and the *P* value is shown. Each symbol represents an individual mouse and the error

bars represent the s.d. of the mean. **d**, Schematic of terminal PLX treatment of C57BL/6J wild

type mice (Wt) from 2 mo of age until death and blood plasma collected at 3, 4, 5, 6, and 7 mo

of age. **e**, Plasma concentration of PLX3397 in female or male Wt mice. The difference between

female and male mice was assessed by two-way repeated measures ANOVA and the *P* value is

1552 shown. **f**, Body weights of female or male Wt mice treated with vehicle or PLX3397. Differences

1553 in weight between vehicle and PLX3397 treatment in male or female mice were evaluated by

1554 two-way repeated measures ANOVA and *P* values are shown. In **b**, **e**, and **f**, each symbol

1555 represents an individual mouse and the lines represent the group means.

1556

Supplementary Fig. 10| Food consumption and activity of Tg2541 mice treated with

CSF1R inhibitors. a, **b**, Food consumption measured in Tg2541 mice receiving terminal

treatment with vehicle or PLX3397 (275 mg/kg oral) (**a**), or vehicle or PLX5622 (1200 mg/kg

oral) (**b**), reported as grams of food per gram of mouse body weight per day (g/g/d). **c**, **d**,

Running wheel activity measured in Tg2541 mice receiving terminal treatment with vehicle or

PLX3397 (**c**), or vehicle or PLX5622 (**d**), reported as total distance traveled in meters per day

(m/d). In **a**–**d**, each symbol represents an individual mouse, with female mice shown as closed

or open circles and male mice shown as closed or open triangles. Error bars represent s.d. of

the mean. Two-way ANOVA was used to compare female and male mice and *P* values are

shown.

Supplementary Fig. 11| Validation of Nanostring sequencing results with real-time PCR.

a, Delta Ct values of 24 selected genes in hindbrains of Tg2541 mice treated with PLX5622

(1200 mg/kg) and vehicle. A lower Delta Ct reading indicates higher expression level. **b**, Delta

Ct values of 29 selected genes in hindbrains of Tg2541 and wild type mice treated with vehicle.

A lower Delta Ct reading indicates higher expression level. **c**, Expression ratios of the 53 genes

in (**a**) and (**b**) as measured by qPCR and Nanostring sequencing. **d**, Scatter plot of the data in

(**c**). Red dotted line shows a linear trend line.

Supplementary Fig. 12| Gene expression pattern analyses of non-microglial and neuron-
specific genes. **a−c,** Analyses similar to Fig. 6c−e, using (**a**) all non-microglial genes in
hindbrain, or neuron-specific genes in (**b**) forebrain and (**c**) hindbrain. **d−g**, Analyses similar to
Fig. 6g−j, using data from hindbrains.

1580

Supplementary Fig. 13| PLX treatment led to increased CSF1 expression and decreased CSF1R expression. Quantification of CSF1 expression showed slight increase after PLX treatment, while CSF1R expression showed marked decrease. Two-way ANOVA was used for

1584   statistical tests. Bottom panels showed scatter plots of brain PLX concentration against CSF1 or

1585   CSF1R expression. Linear regressions were calculated for each group.

Supplementary Fig. 14| Validation of IEG up-regulation following PLX treatment. a, **b**,

Real-time PCR quantifications of the mRNA levels of five IEGs in forebrains (**a**) and hindbrains

(**b**) from PLX5622- or vehicle-treated Tg2541 or wildtype mice. **c,** Correlation of brain PLX

concentration and expression levels from the five immediate early genes in the forebrains from

PLX5622-treated Tg2541 mice. **c**, **d**, Correlation of brain PLX concentration and normalized

expression levels from all immediate early genes (IEGs, 56 genes for each mouse) in the (**d**)

forebrains and (**e**) hindbrains of vehicle- and PLX3397-treated Tg2541 mice. Pearson's

correlation analysis was performed and the *P* values are shown. **f**, Quantification of normalized

expression levels from all IEGs (56 genes for each mouse) in the forebrain and hindbrain of

vehicle- and PLX3397-treated Tg2541 mice. Mann-Whitney tests were used for comparisons

between groups. Each symbol represents an individual mouse.

1598

Supplementary Fig. 15| Brain neurotransmitter levels unaffected by PLX3397 in male

Tg2541 mice. a, Forebrain lysates of wild type (Wt) or Tg2541 (Tg) mice, dosed for 6-12 weeks

with PLX3397 (275 mg/kg oral) or vehicle beginning at 4 months of age, were measured by

hydrophilic interaction liquid chromatography tandem mass spectrometry (HILIC-MS/MS). **b**,

HILIC-MS/MS analysis of the same samples for glycine. **c**, HILIC-MS/MS analysis of the same
samples for histamine. **a–c,** Mann-Whitney tests were used for comparing vehicle and PLX3397
treatment groups within mouse genotype and sex, and the P values of statistically significant
differences ($P < 0.05$) are shown. Each symbol represents an individual mouse and error bars
indicate the s.d. of the mean.

1608

Supplementary Fig. 16| GFAP expression is reduced by PLX3397 treatment in Tg2541

mice. a, **b**, GFAP levels measured by IHC in five different brain regions, three forebrain regions

and two hindbrain regions, of (**a**) wild type (Wt) and (**b**) Tg2541 mice treated with PLX3397 (275

mg/kg oral) or vehicle. GFAP levels in Tg2541 mice were corrected for the average levels in

each brain region of PLX3397- and vehicle-treated Wt mice. **c**, **d**, *GFAP* mRNA levels

measured by Nanostring in the forebrain or hindbrain of (**c**) wild type and (**d**) Tg2541 mice

treated with PLX3397 (275 mg/kg oral) or vehicle. *GFAP* mRNA levels in Tg2541 mice were

corrected for the average levels in each brain region of PLX3397- and vehicle-treated Wt mice.

In **a**−**d**, Two-way ANOVA with Holm-Šidák post hoc testing was used to evaluate the differences

between vehicle- and PLX3397-treated groups within each brain region and *P* values of

statistically significant differences ($P < 0.05$) are shown. Each symbol represents an individual

mouse with female mice shown as closed or open circles and male mice shown as closed or

open triangles. **e**−**h**, Representative GFAP IHC images of a brain section from (**e**) a Wt vehicle-

treated mouse, (**f**) a Wt PLX3397-treated mouse, (**g**) a Tg2541 vehicle-treated mouse, and (**h**) a

Tg2541 PLX3397-treated mouse. High-magnification images of the indicated hippocampus

(Hip) and midbrain (Mid) regions are shown below each image. Scale bars, 500 μm.

1625

Supplementary Fig. 17| Generation, optimization, and validation of bigenic Tg2541/GFAP-luciferase mice for in vivo bioluminescent imaging (BLI) of astrocytosis. a, Kaplan-Meier plot shows that the survival curve (kinetics of disease) is unchanged for Tg2541 homozygous mice bred to B6-albino background and homozygous for the reporter GFAP-luciferase transgene. **b**, **c**, Kaplan-Meier plots for (**b**) female and (**c**) male mice showing that survival curves of homozygous and hemizygous B6-albino bigenic Tg2541 mice is not sex-dependent; homozygous Tg2541 mice have a median survival of 212 days, and hemizygous Tg2541 mice have a median survival of 378 days. **d**, **e**, As a crude surrogate of general health, longitudinal measurements of mouse body weight (grams) shows that in contrast to Wt mice, (**d**) female and (**e**) male Tg2541 mice lose weight as a result of decreased food intake from increasing

paraparesis with disease progression. Because earlier studies gave a standard volume of d-

luciferin substrate regardless of changes in individual mouse weight, we optimized the protocol

to give a 25mg/kg of CycLuc1 to increase consistency in BLI measurements. **f**, To validate the

GFAP-luciferase reporter gene and the synthetic luciferin substrate (CycLuc1) *in vivo*, we

performed BLI in Tg2541 homozygous mice (~200 days old) with advanced disease pathology

and showed the BLI signal is significantly increased in the forebrain, hindbrain and spinal cord

as compared to similar aged Tg2541 hemizygous and Wt mice. Welch ANOVA with Dunnett T3

post hoc testing was used to compare groups and *P* values for all statistically significant

differences ($P < 0.05$) are shown. Error bars represent the s.d. of the mean. **g**, Time-lapse

imaging (two-minute intervals) of BLI signal from brain and spinal cord in Tg and Wt mice

showed that peak BLI signal occurred between 14 and 20 minutes after CycLuc1 injection to the

peritoneum. Thus, to decrease variability in our study, we collect images at 14, 16, and 18

minutes post-injection and average all three time points to account for subtle differences in time

of injection and individual mouse pharmacokinetic of CycLuc1. **h**, **i**, Longitudinal BLI plots in (**h**)

female and (**i**) male mice show kinetics of gliosis in the Tg2541 brain and spinal cord (3 mice

per field of view). **j**, Example image of the field of view (magnification) used to capture both the

brain and spinal cord in three mice per time point. In **d**, **e**, **h**, and **i**, differences between groups

were evaluated by repeated measures ANOVA and *P* values are shown. Each symbol

represents an individual mouse and lines indicate group means.

1655

Supplementary Fig. 18| Brain and spinal cord BLI of PLX3397-treated wild type and

Tg2541 mice. a, **b**, Longitudinal brain BLI plots of (**a**) female and (**b**) male wild type (Wt) mice

treated with PLX3397 (275 mg/kg oral) or vehicle. **c**, **d**, Longitudinal brain BLI plots of (**c**) female

and (**d**) male Tg2541 mice (Tg) treated with PLX3397 (275 mg/kg oral) or vehicle. The data

were corrected for brain BLI levels of PLX3397- and vehicle-treated Wt mice, shown in **a** and **b**.

Due to resolution and/or sensitivity of this magnification, we did not observe any differences

between PLX3397 and vehicle groups. Higher magnification imaging on single mice from the

same cohort was performed as shown in Fig. 8e−g. **e**, **f**, Longitudinal spinal cord BLI plots of (**e**)

female and (**f**) male Wt mice treated with PLX3397 (275 mg/kg oral) or vehicle. **g**, **h**,

Longitudinal spinal cord BLI plots of (**g**) female and (**h**) male Tg2541 mice (Tg) treated with

PLX3397 (275 mg/kg oral) or vehicle. The data were corrected for spinal cord BLI levels of

PLX3397- and vehicle-treated Wt mice, shown in **e** and **f**. **i**, **j**, High magnification BLI plots of the

(**i**) forebrain and (**j**) hindbrain of female Wt mice treated with PLX3397 (275 mg/kg oral) or

vehicle, and used to correct the BLI data of female Tg2541 mice presented in Fig. 8f,g. In **a**−**j**,

differences between vehicle and PLX3397 treatment were evaluated by repeated measures

ANOVA and *P* values are shown. Each symbol represents an individual mouse and lines

1672   indicate group means.

1673

1674

1675 **Supplementary Fig. 19| Clustering of microglia specific genes.** The schemaball graph of
1676 microglial-specific genes as shown in Fig. 10. The name of each gene is shown next to the dot.

Female   Male

Supplementary Fig. 20| Tau pathology or PLX treatment did not lead to marked alteration

of inflammasome-related genes. Quantification of 14 inflammasome-related genes in different

experimental groups.

Supplementary Data File 1| Statistical tests of sex-specific drug effects on microglia, tau,

and neurons. **a**, Statistical test results for 3-way ANOVA analyses for each figure panel where

female and male mice were combined. Statistical test results include the type III sum-of-squares

(SS), the F statistic, and the *P* value for the main effect of sex and for the sex*drug interaction

effect. **b**, **c**, Statistical test results for 2-way ANOVA analyses for (**b**) female or (**c**) male mice

separately, for each figure panel where female and male mice were combined. Statistical test

results include the type III sum-of-squares (SS), the F statistic, and the *P* value for the main

effect of drug, and the *P* value for the multiple comparisons post-hoc testing of each drug

compared to the vehicle in the forebrain, hindbrain, or spinal cord. 'na' indicates that the data is

not available because the drug was not tested in that particular experiment.

1691

Supplementary Data File 2| Differentially expressed genes in male and female PLX-

treated Tg2541 mice. Forebrain samples from male and female Tg2541 mice were analyzed

for gene expression by Nanostring following chronic treatment with PLX5622. The data

represent the average expression levels of five male and five female mice, which were then

compared by unpaired t-test and the *P* value is shown. Please note that the entire transcriptome

dataset which also includes hindbrain data, wild type mice, and vehicle-treated mice are

available from Github with the following link: https://gitfront.io/r/user-

8849465/665dd65fd9d9e78650ed02b9f30236d99240de39/UCSF-PLX-nanostring/

1700