An epigenome atlas of neural progenitors within the embryonic mouse forebrain

Christopher T. Rhodes[1], Joyce J. Thompson[2], Apratim Mitra[3], Dhanya Asokumar[1,2], Dongjin R. Lee[1], Daniel J. Lee[1,2], Yajun Zhang[1], Eva Jason[3], Ryan K. Dale[3], Pedro P. Rocha[2,4], Timothy J. Petros[1]

[1]Unit on Cellular and Molecular Neurodevelopment

[2]Unit on Genome Structure and Regulation

[3]Bioinformatics and Scientific Programming Core

Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD),

National Institutes of Health (NIH), Bethesda, MD 20892, USA.

[4]National Cancer Institute (NCI), NIH, Bethesda, MD 20982, USA.

*Correspondence: tim.petros@nih.gov

Keywords: Interneuron, Neurodevelopment, Epigenetic, Chromatin Accessibility, Single cell sequencing

ABSTRACT

A comprehensive characterization of epigemonic organization in the embryonic mouse

forebrain will enhance our understanding of neurodevelopment and provide insight into

mechanisms of neurological disease. We collected single-cell chromatin accessibility profiles from

four distinct neurogenic regions of the embryonic mouse forebrain using single nuclei ATAC-Seq

(snATAC-Seq). We identified thousands of differentially accessible peaks, many restricted to

distinct progenitor cell types or brain regions. We integrated snATAC-Seq and single cell

transcriptome data to characterize changes of chromatin accessibility at enhancers and

promoters with associated transcript abundance. Multi-modal integration of histone modifications

(CUT&Tag and CUT&RUN), promoter-enhancer interactions (Capture-C) and high-order

chromatin structure (Hi-C) extended these initial observations. This dataset reveals a diverse

chromatin landscape with region-specific regulatory mechanisms and genomic interactions in

distinct neurogenic regions of the embryonic mouse brain and represents an extensive public

resource of a 'ground truth' epigenomic landscape at this critical stage of neurogenesis.

INTRODUCTION

Inhibitory GABAergic interneurons are a heterogeneous cell population that can be classified based on electrophysiological properties, morphologies, synaptic connectivity, neurochemical markers and transcriptomes[1-3]. In the forebrain, GABAergic neurons are born from transient embryonic structures in the ventral telencephalon known as the medial, caudal and lateral ganglionic eminences (MGE, CGE, and LGE, respectively), whereas glutamatergic projection neurons arise from the dorsal telencephalon. The MGE and CGE (and adjacent preoptic area) generate nearly all cortical and hippocampal interneurons, with each region generating almost entirely distinct, non-overlapping interneuron subtypes[4-6].

The embryonic brain contains two primary classes of neural progenitors: multipotent self-renewing apical progenitors (APs, also known as radial glia cells) located in the ventricular zone (VZ) and basal progenitors (BPs) that undergo neurogenic divisions within the subventricular zone (SVZ)[7]. Both APs and BPs give rise to postmitotic immature neurons (Ns) within the GEs that migrate tangentially to populate the telencephalon. Recent evidence demonstrates that initial interneuron subtype fate is specified within the GEs as interneuron progenitors exit the cell cycle[8-11]. It is well established that changes in a cell's epigenetic landscape alters cell fate decisions throughout normal development[12,13] and can be associated with neurodevelopmental disorders[14-16]. In fact, many neurological and psychiatric disease-associated genes are expressed during embryonic development[17,18] and are enriched specifically in APs and immature interneurons[19,20]. Furthermore, many neurological disorders have been linked directly to polymorphisms in enhancer regions[21,22], and GWAS indicate that >90% of disease-associated single nucleotide polymorphisms (SNPs) are located outside of coding regions[23]. Thus, a thorough characterization of the epigenomic landscape during neurogenesis is necessary to understand normal development and potential disease etiologies.

Using a single nuclei assay for transposase accessible chromatin followed by sequencing (snATAC-Seq)[24], we characterized the chromatin accessibility of cells during the transition from

progenitors to lineage restricted neurons within the GEs and dorsal telencephalon of the

embryonic mouse brain. We identified differentially accessible peaks (DA peaks) enriched in

specific brain regions and/or distinct progenitor cell types. Among chromatin accessibility profiles,

individual loci smoothly transition from open to closed chromatin (or vice versa) during the initial

stages of neurogenesis. We validated our snATAC-Seq and single cell RNA sequencing (scRNA-

Seq) observations with orthogonal epigenomic methods. Genome-wide histone modification

profiles associated with promoters (H3K4me3), active enhancers (H3K27ac) and gene repression

(H3K27me3) were highly concordant with our snATAC-Seq profiles showing spatially restricted

enrichment patterns. Our single-cell derived gene-enhancer models largely agreed with direct

observations of promoter-enhancer interactions by Capture-C and higher-order chromatin

domains by Hi-C. These data are available as a UCSC Genome Browser track hub, providing an

important new resource for the field to explore spatial differences in the chromatin landscape of

distinct neuronal progenitors within the embryonic mouse forebrain.

RESULTS

Identifying chromatin accessibility profiles in the embryonic forebrain

To ascertain the chromatin accessibility landscape of differentiating neurons, we dissected

the MGE, CGE, LGE and cortex from wild-type mice at embryonic day 12.5 (E12.5) when cells in

the GEs are undergoing neurogenesis[4] and processed single nuclei on the 10X Genomics

platform (Fig. 1a). Since cortical neurogenesis occurs later, we also harvested E14.5 dorsal cortex

to compare both temporally (E12.5 GEs vs. E12.5 cortex) and neurogenically (E12.5 GEs vs.

E14.5 cortex) matched dorsal and ventral forebrain. Sequencing libraries contained 39,253 single

nuclei, with 10,310 from MGE, 8,543 from CGE, 11,346 from LGE and 9,054 from cortex. Libraries

were aggregated, downsampled to equal numbers of median fragments per nuclei and normalized

by latent semantic analysis (LSA) before peak calling, construction of cell-by-peak count matrices

and integration of different samples (Extended Data Fig. 1a-g).

Using uniform manifold approximation and projection (UMAP), nuclei segregated largely

by tissue region (Fig. 1b). The smart local moving (SLM) algorithm[25] detected 27 clusters, of which

three non-neuronal clusters were removed to retain 96.8% of nuclei in 24 clusters (Fig. 1c). Cell

types were assigned by inspecting promoter accessibility (PA) (defined as the sum of reads

mapping within -2000 bp of a TSS) of canonical cell type markers and were further refined by

transferring cell type assignments from droplet-based scRNA-Seq data of E12.5 embryos (Fig.

1d and Extended Data Fig. 1i-n) to the snATAC-seq dataset. PA for markers of interneuron and

excitatory glutamatergic pyramidal cell maturation segregated clusters into mitotic APs and BPs,

and postmitotic Ns (Fig. 1D and Extended Data Fig. 1o-q). MGE and most CGE and LGE nuclei

displayed accessible chromatin at GABAergic neuron markers, while virtually all cortical nuclei

have accessible chromatin at markers of glutamatergic neurons (Extended Data Fig. 1o and

Extended Data Fig. 2). A group of LGE and CGE nuclei displayed accessibility profiles more

similar to glutamatergic neuron markers and were labeled as a 'mixed' neuron population

(Extended Data Fig. 1o and Extended Data Fig. 2). This was likely contamination from the pallial-

subpallial boundary (PSB), a region that gives rise to cells located in the piriform cortex, claustrum

and amygdala[26,27].

To quantify temporal differentiation programs, a minimum spanning tree was constructed

in Monocle3[28,29] (Fig. 1e). The progression along pseudotime largely recapitulated known

maturation markers, from *Nes*+ and *Ccnd2*+ cycling progenitors to *Dcx*+ and *Rbfox3*+ postmitotic

neurons (Fig. 1f-i). Additionally, region restricted genes such as *Nkx2-1* and *Lhx6* in the MGE,

and *Neurod6* in the cortex, displayed open accessibility profiles restricted to these regions (Fig.

1j-m). We examined pseudobulk ATAC read pileups within each cluster for regionally restricted

genes for the MGE (*Nkx2-1*) and cortex (*Neurod6*) (Fig. 1n-o) and for two pan-neuronal

maturation markers for APs (*Nes*) and BPs/Ns (*Dcx*) (Fig. 1p-q). High signal strength for *Nkx2-1*

and *Neurod6* reads were restricted to the MGE and cortex/mixed clusters, respectively. As

expected, *Nes* reads were enriched in AP clusters with diminished signals in BP and N clusters,

whereas *Dcx* displayed the inverse low-AP to high-N accessibility profile. Notably, strong

accessibility was detected in the second intron of *Nes* which contains a known enhancer[30] (Fig.

1p). These observations were in agreement with ENCODE H3K4me3 ChIP-Seq data from E12.5

mouse forebrain (Fig. 1n-q).

Differentially accessible peak profiles during neurogenesis

When comparing differentially accessible (DA) peaks among all possible peaks (intergenic

peaks and those in promoters/gene bodies), we sought to detect cluster and cell type specific

markers, detecting a total of 30,046 DA peaks (FDR <= 0.05, average log(fold change) > 0) across

all clusters (Fig. 2a, Extended Data Fig. 3a, and Supplementary Table 1). These DA peaks

represent accessible genomic loci that are potentially unique to specific cell types. To characterize

DA peak profiles across clusters, we asked whether the genomic coordinates bounding DA peaks

of one cluster had reads in any other cluster. If there were reads in a DA peak from one cluster in

another cluster, this peak was considered overlapping between the clusters. No minimum

threshold for overlapping peak counts was used before calculating the percentage of overlapping

DA peaks from one cluster compared to all peaks from the same cluster (Fig. 2a, "% Overlap").

We also counted the number of DA peaks from each cluster to assess if there were differences

in the number of DA peaks per cluster (Fig. 2a, "Peak counts").

Unsupervised hierarchical clustering (HC) of DA peak profiles created a dendrogram that

segregated clusters initially by maturation state (AP, BP and N) and secondarily by tissue origin.

Since the overall profiles of LGE and CGE nuclei were very similar, these two regions were

labeled 'LGE/CGE' for this analysis. The 'mixed' neuron population (Extended Data Fig. 1o) was

also left as an individual group for this analysis. The dendrogram generated by HC is very similar

to cluster relationships in LSA/UMAP space, which is encouraging since different features were

used in each analysis (HC: DA peaks in cluster pairs, LSA: all peaks across libraries). As

expected, dot plot positions containing both high DA peaks counts and high percent overlap were

almost exclusively grouped along the diagonal, while positions with limited numbers of DA peaks

or low between-cluster peak overlap populated the off-diagonals, indicating high specificity of DA

peaks to specific clusters (Fig. 2a). The cell type- and brain region-specific DA peak profiles are

consistent with previous models of chromatin reorganization during cellular differentiation[31,32].

We also visualized binarized peak signals (i.e., "open" or "closed" regions) per cluster

using a heatmap and again observed high peak signals primarily along the diagonal (Extended

Data Fig. 3a). Intriguingly, the mean number of DA peaks decreased as maturation progressed,

with a significant decrease in DA peaks in BP and N nuclei compared to APs in each tissue

(Extended Data Fig. 3c-d). Despite this decrease, DA peak profiles became more distinct as

maturation progressed, as indicated by the low between-cluster peak overlap along the dot plot

off-diagonals (Fig. 2a and Extended Data Fig. 3a). The decrease in global accessibility over time

is consistent with previous observations during cellular differentiation[33]. We observed a greater

number of DA peaks in all maturation stages in the cortex compared to GEs (Extended Data Fig.

3e-f).

Regions of accessible chromatin are enriched for transcription factor (TF) binding motifs

that often play essential roles in driving cell specification. To characterize region-enriched TF

motifs, we performed motif analysis on DNA sequences within DA peaks using the JASPAR

CORE[34] vertebrates collection. Motif analysis detected TF motifs that have lineage- and tissue-

specific roles during interneuron neurogenesis (Fig. 2a, Extended Data Fig. 3b and

Supplementary Table 2). Further, among motifs enriched in specific clusters that also had a

corresponding differentially accessible promoter, the top 5 matches contain one or more TF motifs

with known spatial or temporal expression profiles correlating with the cell cluster (Fig. 2b and

Supplementary Table 3). For example, UMAP visualization of motif enrichment by chromVAR[35]

for NR2F1 (MGE and CGE-specific)[36], ASCL1 (GABAergic BP-enriched), EMX1 (cortical

progenitor AP/BP-specific) and NEUROD2 (cortical postmitotic N-specific) motifs demonstrated

the expected neuronal lineage progression and/or region-restricted patterns (Fig. 2b-e). We

further quantified genome-wide DA peak distributions within annotated gene regions and found

the majority of DA peaks were constrained to promoter and intronic regions of gene bodies, and

distal intergenic loci (Extended Data Fig. 3g). These findings indicate that DA peaks are specific

to brain region and cell-state, and importantly, that DA peaks contain lineage-specific TF motifs

that may regulate cell fate decisions differentiation.

Candidate cis-regulatory elements are dynamic and cell state dependent

The global decrease of DA peak numbers from cycling neural progenitors to postmitotic

immature neurons prompted us to examine changes to DA peaks during maturation and lineage

commitment. For this and all future analysis, we removed the 'mixed' population to characterize

only LGE and CGE-derived GABAergic cells. The Monocle3 extension Cicero[37] detects

interactions or 'connections' between any 2 genomic loci and then assigns a co-accessibility score

between the 2 sites, thereby calculating the proportion of nuclei containing a given co-accessibility

interaction within a population. We quantified such interactions within all nuclei and detected

92,414 connections that had a co-accessibility score equal to or greater than 0.25, meaning a

given locus-locus interaction is detectable in $\geq$ 25% of all nuclei in the defined population

(henceforth "Cicero connections").

To explore co-accessibility changes during neurogenesis, we used a heatmap to visualize

Cicero connections in which at least one of the 2 interacting loci was a DA peak, representing

potential interactions between DA peaks and putative cis-regulatory elements (cREs). Nuclei were

divided into 10 bins of equal pseudotime intervals along the Y-axis, with individual DA peaks

grouped via hierarchical clustering along the X-axis. The heatmap color represents the proportion

of nuclei with a given DA peak (interacting via a Cicero connection) with a co-accessibility score

$\geq$ 0.25 (Fig. 3a). Nearly half of these Cicero connections were enriched in AP nuclei, which is

consistent with an overall decrease in accessibility as development progresses (Extended Data

Fig. 3c-d). As chromatin regions with dynamic accessibility are associated with gene regulation

during neural stem cell activation[32], we hypothesized that Cicero connections enriched in

immature neurons encode lineage-specific cREs that may play a role in neuronal lineage

commitment.

To identify candidate cREs at specific genes, we examined Cicero connections within 0.5

Mb windows of gene TSSs after filtering for co-accessibility scores $\geq$ 0.25. We identified Cicero

connections in which only one of the 2 interacting loci overlapped a TSS, representing potential

interactions between a TSS and putative cREs (henceforth "TSS-cRE connections"). All Cicero

connections within this 0.5 Mb window are visualized as orange arcs whereas TSS-cRE

connections for a selected gene are highlighted in purple (Fig. 3b-c). To characterize Cicero

connections that were spatially restricted, we downsampled tissues to equal nuclei numbers and

detected 91,904, 76,858, 89,366, and 148,942 Cicero connections with co-accessibility scores $\geq$

0.25 in MGE, LGE, CGE and cortex, respectively. *Lhx6* and *Neurod6* Cicero connections are

restricted to the MGE and cortex, respectively, and TSS-cRE connections for *Lhx6* and *Neurod6*

are virtually exclusive to these regions (Fig. 3b). *Nr2f2* (*CoupTF-II*) is a marker for APs in the VZ

of the CGE and caudal MGE[38] whereas *Sp8* is a marker for LGE progenitors that is excluded from

the MGE[39]; both genes displayed highest co-accessibility scores and TSS-cRE connections

counts in the expected regions (Fig. 3b). Overall, tissue specific TSS-cRE connectivity patterns

were similar to regionally restricted gene expression patterns that are critical to neuronal

development.

To characterize Cicero connections that varied between neurogenic cell types, we

downsampled to equal nuclei numbers and detected 89,312, 88,980, and 110,362 Cicero

connections with co-accessibility scores $\geq$ 0.25 in AP, BP and N nuclei, respectively. Among these

Cicero connections, postmitotic genes *Lhx6* and *Neurod6* had their highest co-accessibility scores

and TSS-cRE connections in BP and N nuclei (Fig. 3c). Progenitor-enriched genes *Nr2f2* and *Sp8*

had their highest co-accessibility scores and TSS-cRE connections in APs, with decreased

connections and co-accessibility scores in Ns. The pan-AP marker nestin (*Nes*) had its highest

co-accessibility scores and TSS-cRE connections in APs throughout all regions compared to BPs

and Ns. Conversely, the postmitotic GABAergic marker *Gad1* had its highest co-accessibility

scores and connection counts in BPs and Ns. As with regional specificity of TSS-cRE

connections, the temporal connection patterns largely recapitulate known gene expression

patterns as neurons mature. In sum, cREs likely interact with DA peaks and TSSs to regulate

genes in regionally and temporally restricted patterns, and the co-accessibility patterns of TSS-

cREs connections closely resemble known spatial and temporal expression patterns in the

embryonic forebrain.

Integrative analysis of chromatin accessibility and gene expression profiles

To enhance our understanding of the relationship between chromatin accessibility and

gene expression during neurogenesis, we combined chromatin accessibility profiles from

snATAC-Seq data with age and region-matched scRNA-Seq data. Integrating ATAC and RNA

data involves quantifying ATAC reads in or near gene bodies by a Gene Activity Score (GAS) as

a proxy for transcript abundance. After testing multiple GAS metrics for snATAC-Seq/scRNA-Seq

integration, we defined GAS as the sum of all ATAC reads mapping to the promoter, first exon

and presumptive enhancers of a given gene because this GAS metric produced the highest

concordance between ATAC and RNA assays (Extended Data Fig. 4). Following integration, the

clustering of snATAC-Seq and scRNA-Seq cells were highly concordant, with the MGE and cortex

integrated cells formed distinct clusters whereas the LGE and CGE cells were largely overlapping

(Fig. 4a-b). The Louvain algorithm detected 26 clusters (Fig. 4c) and Monocle3 assigned

pseudotime (Fig. 4d) which largely recapitulated temporal and spatially restricted expression

patterns expected in embryonic forebrain neurogenic regions (Fig. 4e-j and Extended Data Fig.

5).

Prior to integration, we refined our cRE predictions to detect presumptive enhancers by

combining our Cicero TSS-cRE analysis with TSS-cRE predictions from the SnapATAC

algorithm[40] (Supplementary Table 4). SnapATAC predictions link distal regulatory elements to

target genes based on the transcript count of a gene and chromatin accessibility at peaks flanking

the gene using gene expression as input variable to predict the binarized chromatin state at peaks.

Our rationale for using multiple algorithms was TSS-cREs connections common to both methods

likely improves cRE predictions. After combining TSS-cRE connections from both methods, we

retained common cREs to generate a list of all detectable presumptive enhancers (Extended Data

Fig. 6). We took the intersection of these loci and ENCODE H3K27ac ChIP-Seq peaks[41] from

E12.5 and E14.5 forebrain, resulting in a set of 'high-confidence' enhancer candidates (henceforth

"presumptive enhancers") (Supplementary Table 5). We detected previously validated VISTA

enhancers[42] interacting with genes in the MGE (hs704 and hs1538 regulating *Nkx2-1*), cortex

(hs627 regulating *Neurod2*), GABAergic progenitors (hs967, hs998, hs1114, hs1354, and hs1540

regulating *Ascl1*) and glutamatergic progenitors (hs1025 regulating *Emx1*) (Fig. 4k-n and

Supplementary Table 6).

To characterize the temporal expression and chromatin accessibility profiles during

neurogenesis, we utilized a hierarchical clustering based approach from the DEGReport package[43]

to group RNA, GAS and enhancer counts from our integrated scRNA-Seq/snATAC-Seq data (Fig. 5a-

r). As not all genes had detectable snATAC based GAS and/or enhancer counts, we selected ~300

of the most differentially expressed genes (DEGs) (from scRNA) among APs, BPs and Ns that

had corresponding GAS and enhancer counts. DEGReports hierarchical clustering uncovered 5

groups containing at least 6 genes (Extended Data Fig. 7a-b), of which over 85% of the genes fell

into two categories: one cluster is consistent with high expression and accessibility profiles within

progenitors (APs and BPs) that are downregulated in postmitotic neurons (156 DEGs, Fig. 5a),

and another cluster with the complimentary profile (90 DEGs, Fig. 5b). Genes within these clusters

displayed similar patterns of expression and chromatin accessibility over pseudotime.

Visualization of representative early-expressed genes *Hes1* (Fig. 5c-f) and *Lmo1* (Fig. 5g-j) and

later-expressed genes *Myt1l* (Fig. 5k-n) and *Lhx6* (Fig. 5o-r) demonstrate consistent trends for

transcript, GAS and enhancer counts over pseudotime. We quantified the number of high-

confidence enhancers associated with upregulated DEGs as maturation progressed and

identified 200 enhancers associated with DEGs that had a positive fold change from APs-to-BPs,

175 enhancers for BPs-to-Ns, and 269 enhancers from APs-to-Ns (Extended Data Fig. 7c).

Likewise, we found 188 enhancers associated with DEGs that had a negative fold change from

APs-to-BPs, 179 enhancers for BPs-to-Ns, and 320 enhancers from APs-to-Ns (Extended Data

Fig. 7c). There is a gradual decrease in the ratio of the number of enhancers being activated

versus the number being decommissioned as maturation progressed (Extended Data Fig. 7d),

suggesting that a greater number of genes and associated enhancers are repressed as

progenitors exit the cell cycle. Taken together, DEGs and associated enhancers exhibit

reorganization during the transition from progenitors to lineage committed postmitotic immature

neurons.

We characterized the differentiation processes by visualizing matched heatmaps for RNA,

GAS and presumptive enhancer counts of highly variable genes (Fig. 5s-u). We selected

transcript counts and corresponding GAS and enhancer counts for the top 500 most variable

genes from the E12.5 integrated dataset, of which 210 had corresponding GAS and enhancer

counts (Extended Data Fig. S7e-g). RNA, GAS and enhancer count matrices for these genes

were co-clustered using hierarchical clustering with a correlation distance metric and average

linkage and visualized in matched heatmaps (Fig. 5s-u). Partitioning early, transitional and late

expressing gene profiles with respect to pseudotime largely followed a continuous progression as

cells matured from APs through Ns. Overall, there was high similarity between 1) early and late

gene expression patterns detected by degPatterns (Fig. 5a-b and Extended Data Fig. 7a-b) and

2) early and late expressing genes visualized in heatmaps (Fig. 5s-u), indicating distinct, dynamic

expression and chromatin accessibility in APs, BPs and Ns. By integrating multiple single cell

modalities, we characterized the chromatin accessibility and gene expression profile of distinct

forebrain regions during neurogenesis.

Histone modifications and higher-order chromatin organization reveal region-specifc chromatin states

Predicting enhancers from snATAC data has enormous potential for mapping regulatory elements in heterogeneous cell populations. To validate some of these predictions, we carried out two additional sets of experiments. First, we preformed CUT&RUN[44] and CUT&Tag[45] on E12.5 MGE, LGE, CGE and cortex to detect histone modifications associated with active/poised promoters (H3K4me3), active enhancers (H3K27ac) and repressed genes (H3K27me3)[46-48]. Most genes with spatially-restricted mRNA and promoter accessibility profiles contained corresponding H3K4me3 peaks whereas spatially-repressed genes were enriched with H3K27me3 (Fig. 6a). More globally, we observed ~70% of ATAC peaks at promoters overlapped with H3K4me3 marks in all brain regions (Fig. 6b).

To further identify candidate enhancers, we combined our Cicero analysis with H3K27ac enrichment. There was less overlap between ATAC peaks and H3K27ac marks (29.2%-45.1%), as expected due to the weaker correlation between accessibility and H3K27ac marks throughout the genome. However, if we restricted analysis to ATAC peaks with a Cicero connection to a promoter (indicative of possible enhancers), then the percent overlap of ATAC peaks with H3K27ac marks increased considerably (54.9%-68.1%) (Fig. 6b).

We observed region-specific colocalization between ATAC peaks, Cicero connections and H3K27ac marks at many genes, some of which represent novel candidate enhancers. For example, there are VISTA enhancers downstream of the GE-enriched *Ascl1*, with one site (hs1540) showing co-accessibilty in nearly all nuclei (Fig. 6c). However, none of these VISTA reporters displayed the expected GE-restricted *Ascl1* expression pattern (https://enhancer.lbl.gov) nor contained GE-enriched H3K27ac marks (Fig. 6c). Instead, we identified two other regions with Cicero interactions with GE-enriched H3K27ac marks compared to cortex (Fig. 6c, grey bars). We identified similar loci near *Lhx6* and *Neurog2* with enriched

H3K27ac marks specifically in the MGE and cortex, respectively, representing potentially novel

cREs (Fig. 6d-e).

We performed Hi-C to characterize chromatin structure genome wide (Fig. 7a) and

Capture-C to directly quantify promoter interactions at ~50 genes with tissue-specific expression

patterns (Fig. 7b and Supplementary Table 7). At the *Nkx2-1* locus, Hi-C data revealed formation

of an MGE-specific chromatin domain compared to other brain regions (Fig. 7a). Capture-C

confirmed these distinct interactions, with the *Nkx2-1* promoter interacting directly with a region

near the *Mbip* gene specifically in the MGE (Fig. 7b). Notably, *Mbip* expression is also restricted

to the MGE during development[49]. Conversely, interactions between the *Nkx2-1* promoter and the

Nkx2-9 and *Pax9* loci (genes not expressed in the embryonic forebrain) were specifically detected

in the LGE, CGE and cortex (Fig. 7b). While the exact nature of these interactions is unclear

(promoter-enhancer, promoter-promoter, etc.), the formation of region-specific chromatin

domains is not observable from the other assays, as the snATAC and histone modifications at

the *Mbip*, *Nkx2-9* and *Pax9* locus are quite similar between the different brain regions (Fig. 7c).

Both Hi-C and Capture-C data identified a direct interaction between the *Nr2f1* promoter

and an intron within *2210408I21Rik* specific to the CGE and MGE, where *Nr2f1* is expressed[36]

(Fig. 7d-e). This locus also contains stronger K27ac signal in the CGE and MGE (Fig. 7f),

providing additional evidence for formation of region-specific promoter-enhancer interactions.

Similarly, we observe cortex-enriched interactions of the *Lhx2* promoter with two putative

enhancers within *Dennd1a* introns, with both loci displaying stronger H3K27ac signals in the

cortex compared to other regions (Fig. 7g-i).

Thus, the combination of single cell accessibility and transcriptomes, histone modifications

and higher-order chromatin interactions represents a comprehensive epigenomic 'ground truth'

of distinct neurogenic regions of the embryonic mouse brain that give rise to specific neuronal

subtypes. These data are publicly available and searchable as a UCSC Genome Browser track

hub (See Data Availability section).

DISCUSSION

We characterized the single cell chromatin accessibility and transcriptomic profiles, histone modifications and higher order chromatin organization in four distinct neurogenic regions of the mouse embryonic forebrain. While recent studies performed single cell sequencing experiments to characterize chromatin accessibility in the mouse and human forebrain[31,50-52], our dataset represents the most comprehensive analysis of the chromatin landscape in the developing brain to date. With this combinatorial approach, we characterized the variation and dynamic reconfiguration of mRNA, gene accessibility and active enhancers during neurogenesis and across different neurogenic cell types. We identified numerous candidate enhancers for genes involved in well-characterized neuronal subtypes, many with region-specific direct genomic interactions verified by Hi-C and Capture-C. These data are publicly available in an easily searchable platform on the UCSC genome browser (See Data Availability section; https://genome.ucsc.edu/cgi-bin/hgTracks?hubUrl=https://hpc.nih.gov/~BSPC-Petros/track_hub/chromatin_quantification.hub.txt&hgS_loadUrlName=https://hpc.nih.gov/~BSPC-Petros/track_hub/session.txt&hgS_doLoadUrl=submit). This dataset will be an important resource for the field leading to a greater understanding of the genetic and epigenetic mechanisms regulating initial neuronal fate decision in the embryonic forebrain.

Gene expression and DA peaks were strongly correlated with H3K4me3 and H3K27me3 peaks at active and repressed promoters, respectively, in specific brain regions. There was also a high correlation between ATAC peaks with Cicero connections to gene promoters and H3K27ac marks at these ATAC peaks (Fig. 6b), indicative of likely active enhancers. In most instances, there was a lag for chromatin at genes and enhancers to become inaccessible compared to RNA downregulation, both at the individual gene level (Fig. 5c-r) and global level (Fig. 5s-u). This observation implies that some repressive mechanisms (i.e. repressor TFs, DNA methylation, etc.) likely precede repressive histone modifications and decreased chromatin accessibility at cREs.

We observed numerous instances where accessibility of promoters, gene body and/or enhancers

precede transcript upregulation (e.g., *Nkx2-1* in Fig. 4g-h and *Myt1l* and *Lhx6* in Fig. 5k-r), which

is in agreement with several recent reports[51,53]. Future multiomics studies that can simultaneously

capture the epigenome and transcriptome within single cells during development should provide

significant insight into this relationship.

By performing Hi-C and Capture-C on dissected MGE, LGE, CGE and cortex, we

characterized region-specific chromatin domains and enhancer-promoter interactions *in vivo* that

were not previously identifiable in ENCODE or other studies that do not distinguish between

different forebrain regions. For example, the *Nkx2-1* chromatin domain is markedly different

between the MGE (where *Nkx2-1* directly contacts the *Mbip* locus) and non-MGE (where *Nkx2-1*

directly contacts the *Nkx2-9* and *Pax9* locus) (Fig. 7). Perturbing these types of interactions could

reveal important insights into how chromatin organization affects promoter-enhancer interactions

and gene function in a region-specific manner. As we only examined ~50 genes with Capture-C,

the realm of region-specific interactions between genes and cREs in the developing forebrain is

only beginning to be explored. Previous comparative analyses of chromatin structure have

described organ and cell type-specific spatial configurations, but have focused mostly on adult

tissues[54,55]. Analysis of cell populations representing earlier stages of differentiation trajectories

have been mostly restricted to the immune system[56,57], limb differentiation[58] and other

organisms[59,60] and have revealed lower variation of chromatin structure between different cell

types. In contrast, our data show that spatially adjacent cells representing early neuronal

specification processes can present vastly heterogenous 3D chromatin strcutures.

We note several additional intriguing observations from our data. First, the chromatin

accessibility profiles reveal significant diversity in AP clusters from the GEs, much more so than

cortical APs which have greater similarity between each other compared to other clusters (Fig.

2). Much of the reported transcriptional diversity within the GEs has been restricted to postmitotic

cells[8,9], so our data suggests that there may be greater transcriptional and chromatin state

diversity in GE APs than previously appreciated[61].

Second, the number and score of global Cicero connections near a particular gene

(orange arcs from Fig. 3b-c) are only loosely correlated to TSS-cRE connections. In some

instances, the number of global Cicero connections at certain genomic loci appear to be tissue

specific (greater co-accessibility in regions flanking *Neurod6* restricted to cortex and *Lhx6*

restricted to MGE), while global Cicero connections near other genes appear more similar

between tissues (similar co-accessibility for all brain regions flanking *Sp8* and *Nr2f2*). This may

suggest a role of Cis-Co-accessibility Networks (CCANs)[37], modules of connection sites that are

highly co-accessible with one another in specific brain regions during neuronal fate determination.

Third, the population of 'mixed' cells that were collected with the LGE and CGE tissue

expressed markers for both GABAergic and glutamatergic cells yet formed a distinct cluster from

the cortex and GE populations (Fig. 2 and Extended Data Fig. 2). These 'mixed' cells likely reside

at the PSB as they were not detected in the MGE population. The diversity of cells arising from

the lateral/ventral pallium remains poorly characterized, but this region appears to give rise to

glutamatergic cells of the insular cortex, piriform cortex, claustrum and pallial amygdala[26]. A recent

scRNAseq study may shed light on the heterogeneity of this PSB region[27], but further work is

needed to better characterize this cell population.

In sum, the single cell chromatin accessibility and transcript profiles, histone modifications,

and higher-order chromatin structure define the epigenetic "ground truth" of distinct forebrain

regions during initial neuronal fate decisions. This resource will aid our understanding of normal

development and neurological disease as many disease-associated genes are enriched in neural

progenitors and immature neurons[19,20], and many disease-associated SNPs are located in non-

coding enhancer regions[21-23].

ACKNOWLEDGEMENTS

414 We thank S. Henikoff for pA-Tn5 transposome; S. Coon, J. Iben, T. Li and other members of the

415 NICHD Molecular Genomic Core; C. J. McBain, K. Pelkey, J. A. Kassis and members of the Petros

416 Lab for helpful discussions. Further information and requests for resources and reagents should

417 be directed to and will be fulfilled by the Lead Contact, Timothy J. Petros (tim.petros@nih.gov).

419 AUTHOR CONTRIBUTIONS

420 C.T.R and T.J.P. designed the study and wrote the paper. C.T.R., Y.Z. and T.J.P. extracted and

421 purified nuclei. D.R.L. and T.J.P. extracted and purified embryonic cells. Y.Z. prepared single cell

422 sequencing libraries. C.T.R., A.M., D.R.L. and R.K.D. analyzed single cell data. D.J.L., D.A.,

423 J.J.T., C.T.R. and P.P.R. performed and analyzed CUT&Tag and CUT&RUN experiments. J.J.T.

424 and P.P.R. performed and analyzed Hi-C and Capture-C experiments. E.J. prepared the UCSC

425 browser page. P.P.R., R.K.D. and T.J.P. supervised the project.

427 COMPETING INTERESTS STATEMENT

428 The authors declare no competing interests.

METHODS

Animals. The following mouse lines were used in this study: C57BL/6J (JAX# 000664). For timed matings, noon on the day a vaginal plug was observed was denoted E0.5. All experimental procedures were conducted in accordance with the National Institutes of Health guidelines and were approved by the NICHD Animal Care and Use Committee (ACUC). Both male and female embryonic and adult mice were used without bias for the single cell sequencing reactions.

Tissue dissection. To recover embryonic tissue, dams were anesthetized with isoflurane and then euthanized by cervical dislocation. Embryos were removed from the uterus and kept in chilled artificial cerebrospinal fluid (ACSF, in mM: 87 NaCl, 26 NaHCO$_3$, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 0.5 CaCl$_2$, 7 MgCl$_2$, 10 glucose, 75 sucrose, saturated with 95% O$_2$, 5% CO$_2$, pH 7.4). Brains were extracted from E12.5 and E14.5 embryos, hemisected, microdissected to obtain the MGE, LGE, CGE and somatosensory cortex, and kept in ACSF.

Nuclei extraction for single nuclei ATAC-seq, CUT&Tag and CUT&RUN. Nuclei isolation followed the 10X Genomics ATAC nuclei isolation protocol with several modifications. All steps were performed on ice. For each brain region, tissue was transferred to a Dounce homogenizer containing 1mL ATAC lysis buffer (10 mM Tris-HCL, pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% Tween-20, 0.1% IGEPAL, 2% BSA). Samples were dounced with 10 strokes pestle A and 10 strokes pestle B. Lysate was strained through a 40 µm filter pre-wetted with ATAC wash buffer (10 mM Tris-HCL, pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% Tween-20, 2% BSA) and neutralized with 2 mL wash buffer.

For snATAC-Seq, lysates were centrifuged 500 g for 5 min at 4^{o}C, supernatant removed, nuclei pellet washed once with 1 mL wash buffer, and centrifuged 500 g for 5 min. A diluted aliquot of nuclei solution was mixed with Trypan Blue (1:1) and counted on a hemocytometer. For all

samples, we prepared 3,000 nuclei/μL samples for snATACseq reactions, with 5 μL (~15,000

nuclei) used for each snATAC reaction.

For CUT&Tag/CUT&RUN, tissue was homogenized as described above. Nuclei

suspensions were centrifuged 500 g for 5 min at 4°C, supernatant removed, and washed once

with 1 mL wash buffer, centrifuged 500 g for 5 min and washed a final time with 1 mL 1X CUT&Tag

wash buffer (from CUT&Tag IT Assay Kit, Active Motif, #53610) or 1 mL CUT&RUN 1X wash

buffer (1 mL HEPES pH 7.5, 1.5 m 5M NaCl, 12.5 uL 2M spermidine and 47.5 mL dH$_2$O with 1

Roche Complete Protease Inhibitor EDTA-Free tablet). A diluted aliquot of nuclei solution was

mixed with Trypan Blue (1:1) and counted on a hemocytometer. Equal numbers of MGE, CGE,

LGE and cortex nuclei were pooled and diluted to a final concentration of 1000 nuclei/μL in wash

buffer, with 100 μL (~100,000 nuclei) used for each CUT&Tag/CUT&RUN reaction.

Cell dissociation for single cell RNA-Seq, Hi-C and Capture-C. Embryonic tissue was

dissected as described above. To collect whole cells, embryonic MGE, LGE, CGE and cortex

tissue was incubated in 1 mg/mL Pronase (Roche #10165921001) in ACSF for 20 minutes at RT.

Pronase solution was removed and 2 ml of reconstitution solution (1% fetal bovine serum +

DNAse (1:10,000, Roche #47167280001)) in oxygenated ACSF was added. For Hi-C and

Capture-C preparations, DNAse was not included in the reconsititution solution. Cells were

triturated sequentially with fire-polished large, medium and small bore Pasteur pipettes to

mechanically dissociate tissue.

For scRNA-Seq, DRAQ5 (20 μM) and DAPI (1:10,000) was added to the single cell

suspension, which was passed through a pre-wet 30 μm filter and then processed on an SH800

cell sorter to purify the sample. DRAQ5+/DAPI- live cells were collected in low-bind 1.5 mL tubes

containing 100 μL ACSF. Cell solutions were centrifuged at 300 g for 5 min at 4 °C in a swinging

bucket centrifuge and then counted on a hemocytometer. 15,000 cells (or highest amount

recovered after sorting) was used for 10X Genomics scRNA-Seq experiments.

snATAC library preparation and sequencing. snATAC reaction was carried out following 10X

Genomics ATAC User Guide (revision C), libraries were prepared following 10X Genomics and

Illumina guidelines, and sequenced on an Illumina HiSeq2500. Sequencing metrics were as

follows: **CGE**: Replicate 1: Read pairs: 469,280,227; Estimated number of cells: 4,013; Median

fragments per cell: 13,003; Fraction of fragments in peaks: 73.3%, Fraction of transposition events

in peaks: 58.4%. Replicate 2: Read pairs: 115,917,541; Estimated number of cells: 4,530; Median

fragments per cell: 11,082; Fraction of fragments in peaks: 80.3%, Fraction of transposition events

in peaks: 72.1%. **MGE**: Replicate 1: Read pairs: 429,523,963; Estimated number of cells: 6,845;

Median fragments per cell: 12,807; Fraction of fragments in peaks: 72.0%, Fraction of

transposition events in peaks: 54.2%. Replicate 2: Read pairs: 107,989,853; Estimated number

of cells: 3,465; Median fragments per cell: 12,278; Fraction of fragments in peaks: 77.4%, Fraction

of transposition events in peaks: 67.3%. **LGE**: Replicate 1: Read pairs: 491,904,518; Estimated

number of cells: 6,577; Median fragments per cell: 13,426; Fraction of fragments in peaks: 67.7%,

Fraction of transposition events in peaks: 47.4%. Replicate 2: Read pairs: 115,545,951; Estimated

number of cells: 4,769; Median fragments per cell: 10,660; Fraction of fragments in peaks: 77.7%,

Fraction of transposition events in peaks: 65.8%. **Cortex (E12.5)**: Read pairs: 112,120,408;

Estimated number of cells: 4,946; Median fragments per cell: 11,257; Fraction of fragments in

peaks: 78.0%, Fraction of transposition events in peaks: 68.1%. **Cortex (E14.5)**: Read pairs:

433,510,039; Estimated number of cells: 4,108; Median fragments per cell: 16,810; Fraction of

fragments in peaks: 74.2%, Fraction of transposition events in peaks: 63.6%.

scRNA library preparation and sequencing. cDNA libraries were prepared using 10X

Genomics 3' RNA v3 chemistry. Library preparation was carried out following 10X Genomics RNA

User Guide (rev C) and sequenced following 10X Genomics and Illumina guidelines. Samples

were sequenced to the following depths: **CGE**: Reads: 154,348,231; Estimated number of cells:

4,522; Median reads per cell: 34,133; Median genes per cell: 2,805. **MGE**: Reads: 192,833,398;

Estimated number of cells: 6,331; Median reads per cell: 30,459; Median genes per cell: 2,877.

LGE: Reads: 180,580,907; Estimated number of cells: 6,843; Median reads per cell: 26,389;

Median genes per cell: 2,726. **Cortex (E12.5)**: Reads: 176,542,467; Estimated number of cells:

7,453; Median reads per cell: 23,687; Median genes per cell: 2,318.

CUT&Tag library preparation and sequencing. Embryonic tissue was homogenized as

described above before proceeding with CUT&Tag. For each CUT&Tag replicate, 100,000 nuclei

were resuspended in 1.5 mL 1X Wash Buffer and then processed with the Active Motif CUT&Tag

IT Assay Kit (#53610) following manufacturer's instructions. Primary antibodies used: H3K4me3

(Active Motif, 39159), H3K27ac (Abcam, ab4729), H3K27me3 (Cell Signaling, 9733T). Following

library amplification, DNA quantity was determined with a Thermo Qubit and library quality

characterized with an Agilent Tapestation. Libraries were balanced for DNA content and pooled

before performing a final SPRIselect bead 1x left size selection and paired-end sequenced (50 x

50 bp) on an Illumina NovaSeq. Samples were sequenced to a following depths per library: **CGE**:

H3K27me3 Replicate 1: 13,604,105; H3K27me3 Replicate 2: 47,136,470; H3K4me3 Replicate 1:

50,541,513; H3K4me3 Replicate 2: 25,035,394; **CTX**: H3K27me3 Replicate 1: 13,640,572;

H3K27me3 Replicate 2: 36,097,462; H3K4me3 Replicate 1: 18,35,546; H3K4me3 Replicate 2:

30,114,008; **LGE**: H3K27me3 Replicate 1: 13,539,124; H3K27me3 Replicate 2: 30,498,855;

H3K4me3 Replicate 1: 14,760,336; H3K4me3 Replicate 2: 17,211,681; **MGE**: H3K27me3

Replicate 1: 17,214,723; H3K27me3 Replicate 2: 16,336,430; H3K4me3 Replicate 1: 5,518,769;

H3K4me3 Replicate 2: 14,720,921.

CUT&RUN library preparation and sequencing. Embryonic tissue was homogenized as

described above. Single nuclei suspensions were centrifuged 500g for 5 min at 4°C, supernatant

removed, and washed with 1 mL 1X Wash Buffer (1 mL HEPES pH 7.5, 1.5 m 5M NaCl, 12.5 uL

2M spermidine and 47.5 mL dH_2O with 1 Roche Complete Protease Inhibitor EDTA-Free tablet).

For each CUT&RUN replicate, 100,000 nuclei were resuspended in 1.5 mL 1X Wash buffer.

BioMag® Plus Concanavalin A beads (Bangs Laboratories) were washed in in Binding Buffer

(20mM HEPES pH 7.5, 10mM KCl, 1mM CaCl2, 1mM MnCl2). Nuclei were resuspended in Wash

Buffer, mixed with a slurry of the Concavalin A coated magnetic beads, and rotated for 10 minutes

at room temperature. 10 µl of Concavalin A bead slurry was used per 100,000 cells. The beads

were resuspended in Wash Buffer containing 2 mM EDTA, 0.1% bovine serum albumin, 0.05%

Digitonin, and 1:50 dilution of primary antibody, which was then incubated on a nutating platform

for 2 hours at room temperature. The beads were then washed twice in Digitonin Buffer (20 mM

HEPES pH 7.5, 150 mM NaCl, 0.5 mM Spermidine, 1x Roche Complete Protease Inhibitor no

EDTA, 0.05% Digitonin and 0.1% bovine serum albumin). Then they were incubated with pA-MN

(600 µg/ml, 1:200, either home-made or a gift from S. Henikoff) in Digitonin Buffer for 1 hour at

4ºC. Following this incubation, beads were washed twice with Digitonin Buffer and finally

resuspended in 150 µl of Digitonin Buffer, and equilibrated to 0ºC before adding CaCl2 (2mM).

The beads were then incubated for 1 hour at 0ºC. After this hou,150 µl of 2X Stop Buffer (200 mM

NaCl, 20 mM EDTA, 4 mM EGTA, 50 µg/ml RNase A, 40 µg/ml glycogen), was added. Beads

were incubated for 30 minutes at 37ºC and then pelleted at 16,000g for 5 minutes at 4ºC.

Supernatant was transferred, mixed with 3 µL 10% SDS and 1.8U Proteinase K (NEB), and

incubated for 1 hour at 50ºC, shaking at 900 rpm. 300 µl of 25:24:1 Phenol/Chloroform/Isoamyl

Alcohol was added, solutions were vortexed, and transferred to Maxtrack phase-lock tubes

(Qiagen). The samples in the phase-lock tubes were centrifuged at 16,000g for 3 minutes at room

temperature. 300 µl of Chloroform was added, solutions were mixed by inversion, and centrifuged

at 16,000g for 3 minutes at room temperature. Aqueous layers were transferred to new tubes and

DNA isolated through Ethanol precipitation. These samples were resuspended in 10mM Tris-HCl

pH 8.0 (ThermoFisher). CUT&RUN libraries were prepared following the SMARTer ThruPlex

TAKARA Library Prep kit with small modifications. Double stranded DNA (10 µl), Template

Preparation D Buffer (2 µl), and Template Preparation D Enzyme (1 µl) were combined and added

to each sample. End Repair and A-tailing was performed in a Thermocycler with a heated lid

(22ºC, 25 min; 55ºC, 20 min). To each sample, library Synthesis D Buffer (1 µl) and Library

Synthesis D Enzyme (1 µl) and library synthesis was performed (22ºC, 40 min). Library

Amplification D Buffer (25 µl), Library Amplification D Enzyme (1 µl), Nuclease-free water (4 µl),

and a unique Illumina-compatible indexed primer (5 µl) were added. Library amplification was

performed using the following conditions: 72ºC for 3 min; 85ºC for 2 min; 98ºC for 2 min

(denaturation); 4 cycles of 98ºC for 20 s, 67ºC for 20 s, 72ºC for 10 s (addition of indexes); 14

cycles of 98ºC for 20 s, 72ºC for 10 s (library amplification). Post-PCR clean-up involved

SPRIselect bead 0.6X left/1x right double size selection then washed twice gently in 80% ethanol,

and eluted in 10-12 µl 10 mM Tris pH 8.0. 1:50 Dilutions of primary antibodies against the following

active and repressive histone modifications were used: H3K4me3 (Active Motif, 39159), H3K27ac

(Abcam, ab4729) and H3K27me3 (Cell Signaling, 9733T). Following library amplification, DNA

quantity was determined with a Thermo Qubit and library quality characterized with an Agilent

Tapestation. Libraries were balanced for DNA content and pooled before performing a final

SPRIselect bead 1x left size selection and paired-end sequenced (50 x 50 bp) on an Illumina

NovaSeq. **CGE**: H3k27ac Replicate 1: 39,950,806; H3k27ac Replicate 2: 68,704,235; **CTX**:

H3k27ac Replicate 1: 45,279,956; H3k27ac Replicate 2: 73,920,447; **LGE**: H3k27ac Replicate 1:

46,933,225; H3k27ac Replicate 2: 55,438,848; **MGE**: H3k27ac Replicate 1: 55,871,849; H3k27ac

Replicate 2: 77,034,587.

Hi-C and Capture-C library preparation and sequencing. HiC and Capture-C were performed

and analyzed as described previously[62]. Embryonic tissue dissected and cells were dissociated

as described above. After dissociation, 1 million cells per region were fixed with 1% formaldehyde

(Thermo: 28908) made in 1 ml HBSS media. Fixation was carried out at room temperature on a

nutator, for 10 mins, protected from light. To stop fixation, glycine was added at a final

concentration of 0.13 M and samples were incubated for 5 mins at room temperature followed by

15 mins on ice. Fixed cells were then washed once with ice-cold PBS. After spinning cells at

2500g 4°C for 5 mins, the pellet was flash frozen in liquid nitrogen and stored at -80°C. To perform

Hi-C and Capture-C, pellets were first thawed on ice and then incubated with 1 ml lysis buffer (10

mM Tris-HCL pH8, 10 mM NaCl, 0.2% Igepal CA-630, Roche Complete EDTA-free Sigma

#11836170001). After lysis, cells were dounced and washed with cold PBS. Nuclei extracted with

this method were then collected by centrifugation and subjected to DpnII digest in 50 µl 0.5% SDS

and incubated at 62°C for 10 minutes after which 150 µl of 1.5% Triton-X was added and cells

incubated for 15 minutes at 37°C while shaking at 900 rpm. 25 µl of 10X DpnII restriction buffer

(NEB) was then added, and cells were incubated for 15 minutes while shaking. After that, 200U

of DpnII (NEB R0543M) were added and incubated for 2 hours, then 200U more and incubated

overnight. Next morning 200U more were added and incubated for 3h (total 600U of DpnII). DpnII

was inactivated at 62°C for 20 minutes. For Hi-C, biotin fill-in was done by incubating cells with a

mixture of 4.5 µl dCTP dTTP and dGTP at 3.3 mM, 8 µl klenow polymerase (NEB M0210L) and

37.5 µl Biotin-14-dATP (Thermo 19524016) for 4h at RT while shaking at 900 rpm for 10 seconds

every 5 minutes. Ligation was done overnight at 16°C also rotating at 900 rpm for 10 seconds

every 5 minutes by adding 120 µl of 10X ligation buffer (NEB), 664 µl water, 100 µl 10% Triton-

X, 6 µl BSA 20mg/ml, and 2 µl T4 ligase (NEB cat #M0202M). For Capture-C, biotin fill-in step

was skipped and 50 µl more of water was added to the ligation mix. Crosslink removal was done

overnight with 50 µl of proteinase K in 300 µl of following buffer (10 mM Tris-HCl pH8.0, 0.5 M

NaCl, 1%SDS) while shaking at 1400 rpm at 65°C. Following Sodium Acetate and 100% Ethanol

-80°C precipitation, DNA was resuspended in 50 µl 10 mM Tris HCL for Hi-C or 130 µl for Capture-

C. Sonication for Hi-C was done using Covaris onetube-10 AFA strips using the following

parameters for a 300 bp fragment size (Duration: 10 secs, repeat for 12 times, total time 120 secs,

peak power-20W, duty factor 40%, CPB-50). Sonication for Capture-C was done using Covaris

AFA microtubes 130 with following settings for a fragment size of 200 bp fragments (Duration:

225 secs, peak power-75W, duty factor 25%, Cycles per Burst-1000). Sonications were

performed in a Covaris ME220 sonicator. Sonicated material was then size selected using SPRI

beads with the following ratios: 0.55X and 1X for Capture-C and 0.55X and 0.75X for Hi-C. Hi-C

material was then bound, washed and recovered to 150 µl Streptavidin C1 beads (Thermo 65002)

per sample following manufacturers recommendations. Bead-bound DNA was resuspended in 50

µl 10 mM Tris HCl. Library preparation was done using Kapa Hyper Prep KK8502 kit. 10 µl of

End-repair buffer and enzyme mix were added to resuspended beads and incubated for 30

minutes at RT and then 30 minutes at 65°C. 1 µl of 15 mM annealed-Illumina adaptors, containing

an universal p5 and an indexed p7 oligo, were then incubated with a mixture containing 40 µl of

ligase and ligation buffer at RT for 60 minutes. Libraries were then amplified using 4 reactions per

sample for a total of 200 µl and 10 cycles, as recommended by manufacturer. For Capture-C,

following sonication and size selection, 1 µg of template material was resuspended in 50 µl of 10

mM Tris and used for library prep with 10 µl of End-Repair reaction. 5 µl of 15 mM annealed -

Illumina adaptors were ligated to the Capture-C material. Using a total volume of 100 µl, library

was amplified by PCR using 6 cycles. For capture, 1 µg of Capture-C library per sample was

mixed with mouse COT1 DNA and universal as well as index-specific blocking oligos from

SeqCap EZ HE-oligo (Roche). 4.5 µl pool of biotinylated probes (xGen Lockdown Probe Pools

from IDT), with each probe at 0.4 fmol/µl targeting the promoters of our loci of interested were

added to this mixture and incubated for 3 days at 47°C. Following binding to Streptavidin C1

beads, material was washed as recommended by the SeqCap EZ Hybridization and Wash Kits.

Following washes material was amplified by PCR using Kapa polymerase and 14 cycles. Material

from different samples was then combined and 1 µg of pooled libraries was recaptured in a single

reaction and amplified with 8 cycles. Probes for Capture-C were designed using Capsequm

(capsequm.molbiol.ox.ac.uk/cgi-bin/CapSequm.cgi) by selecting regions adjacent to the gene

promoters of interest. The list of probes used can be found in Supplementary Table 7.

scRNA-Seq Analysis. *Cellranger*: The Cellranger (v3.0.0) pipeline was used to process single

cell/nuclei RNA-Seq libraries, all steps used default parameters unless otherwise noted.

Cellranger mkfastq converted BCL files generated from sequencing to demultiplexed FASTQ files.

Reads were aligned to 10X Genomics's pre-built mouse (GRCm38/mm10) reference genome

with Cellranger count. For single cell RNA-Seq libraries, 10X Genomics's pre-built mRNA model

of GRCm38/mm10 annotation (refdata-cellranger-mm10-3.0.0) was used to map reads to genes,

while single nuclei RNA-Seq libraries used a corresponding pre-mRNA model constructed per

10X Genomics guidelines. Aligned reads were de-duplicated, filtered for valid cell barcodes and

used to construct gene-by-barcode matrices.

Seurat: Filtered gene-by-barcode matrices were used as input to Seurat (v3.0.0,

https://satijalab.org/seurat/) in R (v.4.0.0, https://cran.r-project.org/). For each cell barcode,

summary statistics were calculated for the metadata columns n_Features, n_genes, and percent

reads mapping to mitochondrial genes (if present). Outliers within the metadata columns were

detected using Tukey's fence method for far-out outliers ([Q1 - k(Q3 - Q1), Q3 + k(Q3 - Q1)],

where k = 3) which is resistant to extreme values (https://datatest.readthedocs.io/en/stable/how-

to/outliers.html) and barcodes with any metadata column that contained outliers were removed.

The remaining barcodes were processed using the SCTransform workflow in Seurat.

snATAC-Seq Analysis. *Cellranger-atac*: The Cellranger-atac (v1.2.0) pipeline was used to

process single nuclei ATAC-Seq data, all steps used default parameters unless otherwise noted.

Cellranger-atac mkfastq converted BCL files generated from sequencing to demultiplexed FASTQ

files. Reads were aligned to 10X Genomics's pre-built mouse (GRCm38/mm10) reference

genome and genomic annotation (refdata-cellranger-atac-mm10-1.2.0) by Cellranger-atac count.

Libraries were aggregated and downsampled to equal numbers of median fragments per nuclei

by Cellranger-atac aggr. Aligned reads were de-duplicated, filtered for valid cell barcodes, and

constructed peak-by-barcode and TF-by-barcode matrices and fragments file.

Signac: Filtered peak-by-barcode matrix and fragments file were analyzed using Signac

(v1.0.0) (https://satijalab.org/signac/index.html), all steps used default parameters unless

otherwise noted. For each cell, summary statistics were calculated for the metadata columns

n_Features, n_genes, and percent reads mapping to mitochondrial genes (if present). Outliers

within the metadata columns were detected using Tukey's fence method for far-out outliers, as

previously described, and barcodes with any metadata column that contained outliers were

removed.

Following pre-processing, peak-by-barcode matrices were processed as instructed by

Signac documentation: highly variable peaks were detected by Signac::FindTopFeatures() and

peak-by-barcode matrices normalized by Term Frequency - Inverse Document Frequency (TF-

IDF) method by Signac:: RunTFIDF().

To address batch specific variation, the integration framework from the Seurat package

was used on normalized peak-by-barcode matrices with the functions

Seurat::FindIntegrationAnchors(), which detects features common to all batches, and

Seurat::IntegrateEmbeddings(), which computes a weights matrix using the low dimensional cell

embeddings (LSI coordinates), prior to merging counts matrices across batches. All downstream

Seurat/Signac functions included "latent.vars" arguments to further regress out confounding

variables.

Following integration steps, dimensional reduction with Singular Value Decomposition

(SVD) and Uniform Manifold Approximation and Projection (UMAP), and cluster detection using

smart local moving (SLM) algorithm were performed. RNA cell type predictions from scRNA-Seq

were transferred to snATAC-Seq nuclei using the label transfer method outlined by Signac.

Following the standard Signac workflow, promoter accesbility (PA) for each gene was calculated

by summing all reads mapping to a gene body plus promoter (2 Kb upstream from gene TSSs).

Differential testing we performed for peak counts and PA by Seurat::FindAllMarkers(min.pct=0.2,

test.use='LR', latent.vars='nCount_peaks') and Seurat::FindAllMarkers(), respectively. Motif

analysis was performed using Signac by Seurat::FindMarkers(only.pos=TRUE, min.pct=0.2,

test.use='LR', latent.vars='nCount_peaks') followed by Signac::FindMotifs() and using the

JASPAR (http://jaspar.genereg.net/) CORE vertebrates collection as a reference database. Per

cell motif deviation scores were computed using chromVAR (v1.12.0) and the UCSC mm10

genome sequences (BSgenome.Mmusculus.UCSC.mm10, v1.4.0), both from Bioconductor

(v3.12), with the Signac::RunChromVAR() wrapper.

Cicero: Following outlier detection and dimensional reduction, the filtered peak-by-

barcode counts matrix was transferred from the Seurat object to a Monocle3 (https://cole-trapnell-

lab.github.io/monocle3/) cell_data_set (CDS) object. The counts object was processed with

Cicero (v1.9.1) (https://cole-trapnell-lab.github.io/cicero-release/) following user documentation

with defaults unless otherwise specified. Prior to calculating Cicero co-accessibility scores, a

Cicero CDS object was created using the newly created Monocle3 cds object and UMAP

coordinates   from   the   Seurat   object   (i.e.   cicero::make_cicero_cds(cds.object,

reduced_coordinates=seurat.object@reductions$umap@cell.embeddings)).

SnapATAC: snATAC-Seq reads were processed with SnapATAC (v1.0.0) independently

of   Seurat/Signac/Cicero   workflows   following   user   documentation

(https://github.com/r3fang/SnapATAC) with default parameters unless otherwise specified.

FASTQ reads were converted into snap files using Snaptools (v1.4.1) and Python 2.7 and ATAC

accessible regions were binned into 5000 kB windows and counted to create cell-by-bin matrices.

Snap files were analyzed by SnapATAC. Valid cell barcodes with log10(UMI) between 3 to 6, and

promoter ratio between 0.05 to 0.6 were retained for further analysis. All other barcodes were

discarded. cell-by-bin matrices were merged, binarized, filtered against ENCODE blacklist

regions (http://mitra.stanford.edu/kundaje/akundaje/release/blacklists/), and barcodes with bin

counts less than 1000 and greater than the 95th percentile were removed. Dimensional reduction

was performed using Nyström landmark diffusion maps algorithm, batch reduction was performed

using   Harmony   (https://github.com/immunogenomics/harmony)   by

SnapATAC::runHarmony(eigs.dim=1:40), a K Nearest Neighbor (KNN) graph constructed by

SnapATAC::runKNN(eigs.dims=1:dims_use, k=15) and the Louvain algorithm used to detect

clusters. Gene activity scores were calculated by counting snATAC-Seq reads mapping to known

gene bodies. scRNA-Seq/snRNA-Seq based cell type predictions were transferred to

corresponding snATAC-Seq barcodes by

SnapATAC::snapToSeurat(eigs.dims=1:40,norm=TRUE,scale=TRUE),

Seurat::FindTransferAnchors() and Seurat::TransferData() as described in SnapATAC

documentation. Cluster specific peaks were detected using MACS2 by

SnapATAC::runMACS(macs.options="--nomodel --shift 100 --ext 200 --qval 5e-2 -B --SPMR").

Gene-enhancer pairs were predicted by SnapATAC::predictGenePeakPair() for every TSS that

had a non-zero peak count across all barcodes. The gene-enhancer pairs list was filtered by

removing pairs where FDR > 0.05.

Enhancer Prediction: The greatest SnapATAC -log10(adjusted p-value) value was

multiplied by 2 as an upper limit, and all -log10(adjusted p-values) were percentile ranked per

gene. Similarly, Cicero co-accessibility scores were percentile ranked for each gene. Peaks

detected by SnapATAC or Cicero that did not overlap gene TSSs were extended 500 bp upstream

and downstream before merging overlapping peak coordinates using the GenomicRanges

(v1.42.0) Bioconductor package with GenomicRanges::reduce(). To generate a list of all

presumptive enhancers, merged SnapATAC peaks and merged Cicero peaks were

concatenated. To generate a more stringent list of enhancers for snATAC-Seq/scRNA-Seq

integration, the intersection SnapATAC and Cicero peaks was taken by

GenomicRanges::intersect(). To detect H3K27ac+ presumptive enhancers, concatenated peaks

were further filtered by retaining only peaks that intersected forebrain H3K27ac ChIP-Seq peaks

for age E12.5 (UCSC Database: mm10, Primary Table: encode3Ren_forebrain_H3K27ac_E12,

Big Bed File: /gbdb/mm10/encode3/histones/ENCFF957YEE.bigBed) or E14.5 (UCSC Database:

mm10, Primary Table: encode3Ren_forebrain_H3K27ac_E14, Big Bed File:

/gbdb/mm10/encode3/histones/ENCFF088LWR.bigBed)   by

GenomicRanges::SubsetByOverlaps().

scRNA-Seq/snATAC-Seq Analysis. *scRNA-Seq/snATAC-Seq Integration*: For each data type,

barcodes were filtered to remove far-out outliers, as previously described by Tukey's fence

method. Prior to integrating data, for snATAC-Seq barcodes, gene activity score (GAS) was

calculated by counting reads that mapped to the promoter regions (2kb upstream of TSSs), the

first exon of each transcript, and (if detected) presumptive enhancer loci associated to each gene.

Integration of scRNA-Seq and snATAC-Seq datasets was performed using the Seurat integration

workflow. GAS (snATAC-Seq) and RNA (scRNA-Seq) count matrices were normalized with

Seurat::NormalizeData(), highly variable genes detected with Seurat::FindVariableFeatures() and

common features across all samples selected with Seurat::SelectIntegrationFeatures().

Integration of assays was performed with Seurat::FindIntegrationAnchors() and IntegrateData().

degPatterns: To obtain a list of differentially expressed genes (DEGs),

Seurat::FindMarkers() was used to detect DEGs from embryonic and P30 scRNA-Seq data.

Barcodes were ordered by increasing pseudotime, as assigned by Monocle3. The Seurat object

containing integrated snATAC-Seq/scRNA-Seq datasets was subset into 2 Seurat objects, one

containing scRNA-Seq and one containing snATAC-Seq. degPatterns were quantified using the

previously detected DEGs list and 3 separate matrices: gene-by-barcode counts (from scRNA-

Seq Seurat object), gene-by-barcode counts (GAS slot from snATAC-Seq Seurat object) and

gene-by-barcode counts (enhancer slot from snATAC-Seq Seurat object) by DEGReport::

degPatterns() using the DEGReport (https://github.com/lpantano/DEGreport) package.

RNA/GAS/Enhancers Heatmaps: The top 500 highly variable genes from embryonic

scRNA-Seq data were detected by Seurat::FindVariableFeatures(nfeatures = 500). Barcodes in

the Seurat object containing integrated snATAC-Seq/scRNA-Seq were ordered by increasing

pseudotime, as assigned by Monocle3. This Seurat object was subset into 2 Seurat objects, one

containing scRNA-Seq and one containing snATAC-Seq, which were further split into 3 matrices:

gene-by-barcode counts (from scRNA-Seq Seurat object), gene-by-barcode counts (GAS slot

from snATAC-Seq Seurat object) and gene-by-barcode counts (enhancer slot from snATAC-Seq

Seurat object). RNA, GAS and enhancer gene-by-barcode matrices were concatenated together,

and hierarchical clustering of genes was performed using correlation distance and average

linkage metrics. Following clustering, the concatenated matrix was re-split into RNA, GAS and

enhancer gene-by-barcode matrices. Matrices were centered and scaled for each gene prior to

rendering heatmaps and the gene dendrograms were constructed using the distances calculated

with the concatenated matrix. Manually annotated color bars were based on gene cohorts

detected by degPatterns().

CUT&Tag Analysis. Reads were aligned to mouse genome (GRCm38/mm10 build) with the

Bowtie2 aligner with the following parameters (-p 40 -N 1 --local --very-sensitive-local --no-unal -

-no-mixed --no-discordant --phred33 -I 10 -X 700), aligned reads mapping to blacklisted regions

were removed, and PCR duplicates were removed with PICARD. Quality control for replicates

was assessed using deepTools plotHeatmap and plotProfile, and reproducibility was assessed

by peak calling per replicate with MACS and calculating pair-wise consensus peak counts with

bedtools intersect using the parameter (-f 0.50). Replicates were merged with samtools merge,

deduplicated with PICARD, and peaks called with MACS using the parameter (--broad). To

generate signal tracks for visualization, the merged bam files were converted to normalized

bigWig files using deeptools bamCoverage with the following parameters (--normalizeUsing

RPKM -p 10 --binSize 5 --minFragmentLength 150).

Hi-C and Capture-C analysis. Hi-C and Capture-C libraries were sequenced with paired-end

reads of 51 nucleotides. Data was processed using the HiC-Pro pipeline[63] to produce a list of valid

interactions pairs. This list was converted into cool and mcool files for visualization with higlass[64].

For Capture-C data, the make_viewpoints Hicpro script was used to obtain individual Capture-C

bigwig files for each replicate of each viewpoint with 2kb-sized bins and excluding 500bp

surrounding the DpnII-fragment where probes hibridize. For visualization, averages from

replicates were used.

Data Availability. Gene Expression Omnibus accession numbers for the sequencing data

reported in the paper are GSE167047 (snATAC-Seq), GSE167013 (scRNA-Seq), GSEXXXXXX

(H3K4me3 CUT&Tag), GSEXXXXXX (H3K27me3 CUT&Tag), GSEXXXXXX (H3K27ac

CUT&RUN), GSEXXXXXX (Hi-C) and GSEXXXXXX (Capture-C). A searchable platform with all

single cell accessibility and transcriptomic, CUT&Tag, CUT&Run, Hi-C and Capture-C data can

be found on the UCSC Genome Browser:

https://genome.ucsc.edu/cgi-bin/hgTracks?hubUrl=https://hpc.nih.gov/~BSPC-

Petros/track_hub/chromatin_quantification.hub.txt&hgS_loadUrlName=https://hpc.nih.gov/~BSP

C-Petros/track_hub/session.txt&hgS_doLoadUrl=submit.

REFERENCES

DeFelipe, J. *et al.* New insights into the classification and nomenclature of cortical GABAergic interneurons. *Nat Rev Neurosci* **14**, 202-216, doi:10.1038/nrn3444 (2013).

Gouwens, N. W. *et al.* Integrated Morphoelectric and Transcriptomic Classification of Cortical GABAergic Cells. *Cell* **183**, 935-953 e919, doi:10.1016/j.cell.2020.09.057 (2020).

Tasic, B. *et al.* Shared and distinct transcriptomic cell types across neocortical areas. *Nature* **563**, 72-78, doi:10.1038/s41586-018-0654-5 (2018).

Bandler, R. C., Mayer, C. & Fishell, G. Cortical interneuron specification: the juncture of genes, time and geometry. *Curr Opin Neurobiol* **42**, 17-24, doi:10.1016/j.conb.2016.10.003 (2017).

Pelkey, K. A. *et al.* Hippocampal GABAergic Inhibitory Interneurons. *Physiol Rev* **97**, 1619-1747, doi:10.1152/physrev.00007.2017 (2017).

Miyoshi, G. Elucidating the developmental trajectories of GABAergic cortical interneuron subtypes. *Neurosci Res* **138**, 26-32, doi:10.1016/j.neures.2018.09.012 (2019).

Lui, J. H., Hansen, D. V. & Kriegstein, A. R. Development and evolution of the human neocortex. *Cell* **146**, 18-36, doi:10.1016/j.cell.2011.06.030 (2011).

Mayer, C. *et al.* Developmental diversification of cortical inhibitory interneurons. *Nature* **555**, 457-462, doi:10.1038/nature25999 (2018).

Mi, D. *et al.* Early emergence of cortical interneuron diversity in the mouse embryo. *Science* **360**, 81-85, doi:10.1126/science.aar6821 (2018).

Fishell, G. & Kepecs, A. Interneuron Types as Attractors and Controllers. *Annu Rev Neurosci* **43**, 1-30, doi:10.1146/annurev-neuro-070918-050421 (2020).

Bandler, R. C. *et al.* Single-cell delineation of lineage and genetic identity in the mouse brain. *Nature*, doi:10.1038/s41586-021-04237-0 (2021).

Podobinska, M. *et al.* Epigenetic Modulation of Stem Cells in Neurodevelopment: The Role of Methylation and Acetylation. *Front Cell Neurosci* **11**, 23, doi:10.3389/fncel.2017.00023 (2017).

Sokpor, G., Castro-Hernandez, R., Rosenbusch, J., Staiger, J. F. & Tuoc, T. ATP-Dependent Chromatin Remodeling During Cortical Neurogenesis. *Front Neurosci* **12**, 226, doi:10.3389/fnins.2018.00226 (2018).

Dall'Aglio, L. *et al.* The role of epigenetic modifications in neurodevelopmental disorders: A systematic review. *Neurosci Biobehav Rev* **94**, 17-30, doi:10.1016/j.neubiorev.2018.07.011 (2018).

15   Sokpor, G., Xie, Y., Rosenbusch, J. & Tuoc, T. Chromatin Remodeling BAF (SWI/SNF)
Complexes in Neural Development and Disorders. *Front Mol Neurosci* **10**, 243,
doi:10.3389/fnmol.2017.00243 (2017).

16   Wang, D. *et al.* Comprehensive functional genomic resource and integrative model for
the human brain. *Science* **362**, doi:10.1126/science.aat8464 (2018).

17   Cross-Disorder Group of the Psychiatric Genomics Consortium, C. Genomic
Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric
Disorders. *Cell* **179**, 1469-1482 e1411, doi:10.1016/j.cell.2019.11.020 (2019).

18   Grove, J. *et al.* Identification of common genetic risk variants for autism spectrum
disorder. *Nat Genet* **51**, 431-444, doi:10.1038/s41588-019-0344-8 (2019).

19   Schork, A. J. *et al.* A genome-wide association study of shared risk across psychiatric
disorders implicates gene regulation during fetal neurodevelopment. *Nat Neurosci* **22**,
353-361, doi:10.1038/s41593-018-0320-0 (2019).

20   Trevino, A. E. *et al.* Chromatin accessibility dynamics in a model of human forebrain
development. *Science* **367**, doi:10.1126/science.aay1645 (2020).

21   Carullo, N. V. N. & Day, J. J. Genomic Enhancers in Brain Health and Disease. *Genes*
(Basel) **10**, doi:10.3390/genes10010043 (2019).

22   Kikuchi, M. *et al.* Enhancer variants associated with Alzheimer's disease affect gene
expression via chromatin looping. *BMC Med Genomics* **12**, 128, doi:10.1186/s12920-
019-0574-8 (2019).

23   Perenthaler, E., Yousefi, S., Niggl, E. & Barakat, T. S. Beyond the Exome: The Non-
coding Genome and Enhancers in Neurodevelopmental Disorders and Malformations of
Cortical Development. *Front Cell Neurosci* **13**, 352, doi:10.3389/fncel.2019.00352
(2019).

24   Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A Method for
Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**, 21 29 21-21
29 29, doi:10.1002/0471142727.mb2129s109 (2015).

25   Waltman, L. & van Eck, N. J. A smart local moving algorithm for large-scale modularity-
based community detection. *Eur Phys J B* **86**, 471, doi:10.1140/epjb/e2013-40829-0
(2013).

26   Wullimann, M. F. Should we redefine the classic lateral pallium? *J Comp Neurol* **525**,
1509-1513, doi:10.1002/cne.24127 (2017).

27   Moreau, M. X., Saillour, Y., Cwetsch, A. W., Pierani, A. & Causeret, F. Single-cell
transcriptomics of the early developing mouse cerebral cortex disentangle the spatial
and temporal components of neuronal fate acquisition. *Development* **148**,
doi:10.1242/dev.197962 (2021).

28   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.
Nature **566**, 496-502, doi:10.1038/s41586-019-0969-x (2019).

29 Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386,
doi:10.1038/nbt.2859 (2014).

30 Lothian, C. & Lendahl, U. An evolutionarily conserved region in the second intron of the
human nestin gene directs gene expression to CNS progenitor cells and to early neural
crest cells. *Eur J Neurosci* **9**, 452-462, doi:10.1111/j.1460-9568.1997.tb01622.x (1997).

31 Preissl, S. *et al.* Single-nucleus analysis of accessible chromatin in developing mouse
forebrain reveals cell-type-specific transcriptional regulation. *Nat Neurosci* **21**, 432-439,
doi:10.1038/s41593-018-0079-3 (2018).

32 Maybury-Lewis, S. Y. *et al.* Changing and stable chromatin accessibility supports
transcriptional overhaul during neural stem cell activation. *bioRxiv* (2020).

33 Stergachis, A. B. *et al.* Developmental fate and cellular maturity encoded in human
regulatory DNA landscapes. *Cell* **154**, 888-903, doi:10.1016/j.cell.2013.07.020 (2013).

34 Khan, A. *et al.* JASPAR 2018: update of the open-access database of transcription
factor binding profiles and its web framework. *Nucleic Acids Res* **46**, D260-D266,
doi:10.1093/nar/gkx1126 (2018).

35 Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat
Methods* **14**, 975-978, doi:10.1038/nmeth.4401 (2017).

36 Hu, J. S. *et al.* Coup-TF1 and Coup-TF2 control subtype and laminar identity of MGE-
derived neocortical interneurons. *Development* **144**, 2837-2851, doi:10.1242/dev.150664
(2017).

37 Pliner, H. A. *et al.* Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell
Chromatin Accessibility Data. *Mol Cell* **71**, 858-871 e858,
doi:10.1016/j.molcel.2018.06.044 (2018).

38 Kanatani, S., Yozu, M., Tabata, H. & Nakajima, K. COUP-TFII is preferentially expressed
in the caudal ganglionic eminence and is involved in the caudal migratory stream. *J
Neurosci* **28**, 13582-13591, doi:10.1523/JNEUROSCI.2132-08.2008 (2008).

39 Ma, T. *et al.* A subpopulation of dorsal lateral/caudal ganglionic eminence-derived
neocortical interneurons expresses the transcription factor Sp8. *Cereb Cortex* **22**, 2120-
2130, doi:10.1093/cercor/bhr296 (2012).

40 Fang, R. *et al.* Fast and Accurate Clustering of Single Cell Epigenomes Reveals Cis-
Regulatory Elements in Rare Cell Types. *bioRxiv*, doi:10.1101/615179 (2019).

41 Gorkin, D. U. *et al.* An atlas of dynamic chromatin landscapes in mouse fetal
development. *Nature* **583**, 744-751, doi:10.1038/s41586-020-2093-3 (2020).

42 Visel, A., Minovitsky, S., Dubchak, I. & Pennacchio, L. A. VISTA Enhancer Browser--a
database of tissue-specific human enhancers. *Nucleic Acids Res* **35**, D88-92,
doi:10.1093/nar/gkl822 (2007).

43   Pantano, L. DEGreport: Report of DEG analysis. R package version 1.13.8.,
doi:10.18129/B9.bioc.DEGreport (2017).

44   Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution
mapping of DNA binding sites. *Elife* **6**, doi:10.7554/eLife.21856 (2017).

45   Kaya-Okur, H. S. *et al.* CUT&Tag for efficient epigenomic profiling of small samples and
single cells. *Nat Commun* **10**, 1930, doi:10.1038/s41467-019-09982-5 (2019).

46   Hughes, A. L., Kelley, J. R. & Klose, R. J. Understanding the interplay between CpG
island-associated gene promoters and H3K4 methylation. *Biochim Biophys Acta Gene*
Regul Mech **1863**, 194567, doi:10.1016/j.bbagrm.2020.194567 (2020).

47   Calo, E. & Wysocka, J. Modification of enhancer chromatin: what, how, and why? *Mol*
Cell **49**, 825-837, doi:10.1016/j.molcel.2013.01.038 (2013).

48   Schuettengruber, B., Bourbon, H. M., Di Croce, L. & Cavalli, G. Genome Regulation by
Polycomb and Trithorax: 70 Years and Counting. *Cell* **171**, 34-57,
doi:10.1016/j.cell.2017.08.002 (2017).

49   Tucker, E. S. *et al.* Molecular specification and patterning of progenitor cells in the lateral
and medial ganglionic eminences. *J Neurosci* **28**, 9504-9518,
doi:10.1523/JNEUROSCI.2341-08.2008 (2008).

50   Markenscoff-Papadimitriou, E. *et al.* A Chromatin Accessibility Atlas of the Developing
Human Telencephalon. *Cell* **182**, 754-769 e718, doi:10.1016/j.cell.2020.06.002 (2020).

51   Ziffra, R. S. *et al.* Single cell epigenomic atlas of the developing human brain and
organoids. *bioRxiv* (2019).

52   Allaway, K. C. *et al.* Genetic and epigenetic coordination of cortical interneuron
development. *Nature* **597**, 693-697, doi:10.1038/s41586-021-03933-1 (2021).

53   Ma, S. *et al.* Chromatin Potential Identified by Shared Single-Cell Profiling of RNA and
Chromatin. *Cell* **183**, 1103-1116 e1120, doi:10.1016/j.cell.2020.09.056 (2020).

54   Schmitt, A. D. *et al.* A Compendium of Chromatin Contact Maps Reveals Spatially Active
Regions in the Human Genome. *Cell Rep* **17**, 2042-2059,
doi:10.1016/j.celrep.2016.10.061 (2016).

55   Winick-Ng, W. *et al.* Cell-type specialization is encoded by specific chromatin topologies.
Nature **599**, 684-691, doi:10.1038/s41586-021-04081-2 (2021).

56   Javierre, B. M. *et al.* Lineage-Specific Genome Architecture Links Enhancers and Non-
coding Disease Variants to Target Gene Promoters. *Cell* **167**, 1369-1384 e1319,
doi:10.1016/j.cell.2016.09.037 (2016).

57   Webb, M. R., Hibberd, M. G., Goldman, Y. E. & Trentham, D. R. Oxygen exchange
between Pi in the medium and water during ATP hydrolysis mediated by skinned fibers
from rabbit skeletal muscle. Evidence for Pi binding to a force-generating state. *J Biol*
Chem **261**, 15557-15564 (1986).

58   Bonev, B. *et al.* Multiscale 3D Genome Rewiring during Mouse Neural Development.
Cell **171**, 557-572 e524, doi:10.1016/j.cell.2017.09.043 (2017).

59   Ghavi-Helm, Y. *et al.* Enhancer loops appear stable during development and are
associated with paused polymerase. *Nature* **512**, 96-100, doi:10.1038/nature13417
(2014).

60   Ing-Simmons, E. *et al.* Independence of chromatin conformation and gene regulation
during Drosophila dorsoventral patterning. *Nat Genet* **53**, 487-499, doi:10.1038/s41588-
021-00799-x (2021).

61   Lee, D. R. *et al.* Transcriptional heterogeneity of ventricular zone cells throughout the
embryonic mouseforebrain. *bioRxiv*, doi:doi.org/10.1101/2021.07.05.451224 (2021).
993
62   Thompson, J. J. *et al.* Rapid redistribution and extensive binding of NANOG and GATA6
at shared regulatory elements underlie specification of divergent cell fates. *bioRxiv*,
2021.2007.2028.454132, doi:10.1101/2021.07.28.454132 (2021).

63   Servant, N. *et al.* HiC-Pro: an optimized and flexible pipeline for Hi-C data processing.
Genome Biol **16**, 259, doi:10.1186/s13059-015-0831-x (2015).

64   Kerpedjiev, P. *et al.* HiGlass: web-based visual exploration and analysis of genome
interaction maps. *Genome Biol* **19**, 125, doi:10.1186/s13059-018-1486-1 (2018).

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

Fig 1. Chromatin accessibility in the embryonic forebrain is cell type and state specific. a,
Schematic of snATAC-Seq workflow and neurogenic cell types: apical progenitors (APs), basal
progenitors (BPs) and neurons (Ns). **b-e,** UMAP visualization of single nuclei clustered by brain
region (**b**), SLM (**c**), neurogenic cell type (**d**) and pseudotime (**e**). In **d**, PA = promoter accessibility,
representing reads mapping within 2 kb upstream of TSSs. **f-m**, PA scores for genes enriched in
specific neurogenic cell types (**f-i**) or distinct brain regions (**j-m**). **n-q**, Aggregated reads per SLM
cluster. *Nkx2-1* (MGE), *Neurod6* (cortex), *Nes* (APs) and *Dcx* (BPs/Ns), arranged by either brain
region (**n-o**) or neurogenic cell type (**p-q**). The y-axis range for chromatin accessibility tracks are
normalized to the maximum reads per gene. Peaks: differentially accessible peak coordinates,
H3K4me3: H3K4me3 signal from E12.5 forebrain ENCODE ChIP-Seq data.

1026
Fig 2. Differentially accessible peaks are cluster and lineage specific. a, Embryonic snATAC-
Seq dot plot of differentially accessible peaks (DA peaks) for each cluster. Dot diameter indicates
the percent of DA peaks from one cluster (column cluster labels) which are detectable in any other
cluster (row cluster labels). Color intensity represents the total DA peak count per cluster.
Hierarchical clustering was performed using correlation distance and average linkage. Names of
representative cluster-enriched transcription factor binding motifs in DA peaks listed to right of dot
plot. **b-e**, UMAP plots of nuclei colored by ChromVAR global motif deviations (Z-scores) for
NR2F1, ASCL1, EMX1 and NEUROD2.

1035
Fig. 3. Detection of cis-regulatory elements within the developing mouse forebrain. a,
Heatmap depicting DA peaks binned along pseudotime from embryonic snATAC-Seq nuclei. DA
peaks were filtered to retain only peaks that had at least one Cicero peak-peak connection.
Stacked bar plots to left of heatmap depicts proportion of AP, BP and N nuclei per bin. **b-c**, Cicero
connections within 0.5 Mb window centered around TSS of *Lhx6* (postmitotic MGE marker),
Neurod6 (postmitotic cortex marker), *Nr2f2* (CGE progenitors), *Sp8* (LGE progenitors), *Nes* (pan-
AP) and *Gad1* (pan-interneuron) broken down based on tissue (**b**) or neurogenic cell type (**c**).
Cicero connections overlapping the TSS of selected genes shown as purple arcs, all Cicero
connections in genomic region shown as orange arcs. Y-axis unit is co-accessibility score. Only
connections with co-accessibility scores greater than 0.25 are depicted. Peaks: snATAC peaks
used by Cicero to quantify peak-peak connections. Gene models are visualized at the bottom of
each column with gene of interest highlighted.

1048

Fig. 4. Integrative analysis of embryonic snATAC-Seq and scRNA-Seq. a, Workflow depicting integration of embryonic snATAC-Seq (top left) and scRNA-Seq (top right) data. Bottom, UMAP plot showing integrated snATAC-Seq nuclei and scRNA-Seq cells colored by tissue. **b-d**, UMAP visualization of integrated snATAC-Seq/scRNA-Seq data colored by assay (**b**), Louvain cluster (**c**), and pseudotime (**d**). **e-j**, UMAP visualization of integrated scRNA-Seq cells and snATAC-sec nuclei colored by transcript counts or GAS for *Nes* (**e-f**), *Nkx2-1* (**g-h**) and *Neurod2* (**i-j**). Grey dots in background represent cells/nuclei from other assay. **k-m**, Genome browser tracks displaying enhancer predictions regulating *Nkx2-1* (**k**), *Neurod2* (**l**), *Ascl1* (**m**) and *Emx1* (**n**). Gene-Peak interactions are visualized in blue arcs, arc heights indicate the relative interaction scores between gene TSS (red line) and peaks. snATAC-Seq Peaks: displays co-accessible coordinates, H3K27ac ENCODE: Forebrain E12.5 H3K27ac peaks from ENCODE project, VISTA: enhancers from the VISTA Genome Browser project.

Fig. 5. Transcription, gene accessibility and active enhancer utilization are highly correlated and dynamic during neuron lineage commitment. a-b, Line charts of 'early expressed' (**a**) and 'late expressed' (**b**) DEG clusters detected by degPatterns using embryonic integrated snATAC-Seq/scRNA-Seq data. Y-axis is Z-score for RNA, GAS, or enhancers counts per gene. X-axis is binned pseudotime periods. RNA, GAS, or enhancers for individual genes in these clusters are plotted (156 'early' and 90 'late' genes). **c-r**, Line chart and UMAP visualizations of RNA, GAS and enhancer read counts for representative 'early' genes *Hes1* and *Lmo1* (**c-j**) and 'late' genes *Myt1l* and *Lhx6* (**k-r**). For line charts, Y-axis is Z-score for RNA, GAS, or enhancers counts per gene. X-axis is binned pseudotime periods. Grey dots in background of UMAP plots represent cells/nuclei from other assay. **s-u**, Heatmaps of RNA (**s**), GAS (**t**) and enhancers (**u**) counts for 210 variable genes. Heatmap columns were ordered by hierarchical clustering of 210 variable genes with correlation distance and average linkage. Rows were ordered by pseudotime (assigned by Moncole3). Color bars in **s** indicate genes grouped together based on 'early', 'transition' and 'late' profiles (as assigned by degPatterns and manually refined).

1076

Fig. 6. Histone modifications within embryonic neural progenitors. a, Tracks of scRNA-Seq,
snATAC-Seq and histone modifications correlated with active promoters (H3K4me3), repressed
genes (H3K27me3) and active enhancers (H3K27ac) in MGE, LGE, CGE and cortex (CTX); pan-
GE *Dlx2*, MGE-restricted *Nkx2-1*, CGE-enriched *Nr2f2* and cortex-restricted *Emx1*. **b**, Venn
diagrams comparing ATAC peaks at promoter regions vs. H3K4me3 peaks at promoters (left);
ATAC peaks outside promoters vs. H3K27ac peaks outside promoters (middle); and ATAC peaks
with Cicero connections to a gene promoter vs. H3K27ac peaks outside promoters. Percentages
represent % of ATAC peaks overlapping with histone marks per brain region. **c-e**, Top, Integration
of gene-enhancer predictions using Cicero/SnapATAC interactions as in Extended Figure 6. Arc
height of Cicero/SnapATAC interactions track indicates relative interaction scores between gene
TSS and predicted cis-regulatory elements. Middle, tissue-specific Cicero connections with
snATAC peaks. Cicero connections were filtered to retain scores > 0.25, and connections where
one anchor intersects a gene TSS while the second anchor does not intersect promoter regions
of any genes. Bottom, H3K27ac tracks with VISTA hits for *Ascl1* (**c**), *Lhx6* (**d**) and *Neurog2* (**e**).
Vertical blue line denotes TSS for each gene, grey shaded rectangles indicate loci of interest
related to TSS. VISTA hits near genes are depicted with dark blue bars. Black scale bars above
Cicero/SnapATAC tracks = 50 kb.

1094
Fig. 7. Higher-order chromatin structure within embryonic neural progenitors. a, Hi-C 1D
interaction frequency heatmaps for MGE, LGE, CGE and CTX at MGE-restricted *Nkx2-1*. Black
triangle denotes an MGE-specific interaction, blue arrows denote non-MGE specific interactions.
b, Capture-C pileup at *Nkx2-1* locus extending to vertical dashed lines in **a**. Y-axis represents
reads from loci interacting with *Nkx2-1* promoter bait. Blue line indicates gene TSS, grey bars
indicate potential regulatory elements directly interacting with *Nkx2-1* promoter. **c**, Signal tracks
for single cell assays (scRNA-Seq and snATAC-Seq) and histone modifications correlated with
active promoters (H3K4me3), active enhancers (H3K27ac) and repressed genes (H3K27me3) in
each tissue. VISTA track includes tested enhancers. **d-f**, Hi-C heatmaps (**d**), Capture-C
interactions (**e**) and chromatin landscape (**f**) of CGE-enriched *Nr2f1*. Black triangles in **d** denote
MGE and CGE-enriched interactions. **g-i**, Hi-C heatmaps (**g**), Capture-C interactions (**h**) and
chromatin landscape (**i**) of cortex-restricted *Lhx2*. Black triangles in **g** denote CTX-enriched
interactions. Black scale bars above Hi-C plots in **a**, **d**, **g** = 500 kb, black scale bars below MGE
Capture-C tracks in **b**, **e**, **h** = 100 kb.

1109

Extended Data Fig 1. snATAC-Seq and scRNA-Seq quality control and batch reduction,
and scRNA-Seq based snATAC-Seq annotation. a, Violin plots of snATAC-Seq QC metrics
before outlier removal. **b**, Violin plots of snATAC-Seq QC metrics after outlier removal. **c-e**, UMAP
visualization of snATAC-Seq nuclei prior to batch removal colored by replicate (**c**), tissue origin
(**d**) and SLM cluster (**e**). **f-h**, UMAP visualization of batch corrected snATAC-Seq nuclei colored
by replicate (**f**), tissue origin (**g**) and SLM cluster (**h**). **i**, Violin plots of scRNA-Seq QC metrics
before outlier removal. **j**, Violin plots of scRNA -Seq QC metrics after outlier removal. **k-n**, UMAP
visualization of scRNA-Seq cells colored by tissue (**k**), neuronal class (**l**), Louvain cluster (**m**) and
cell type (**n**). **o-q**, UMAP visualization of snATAC-Seq nuclei colored by neuronal class (**o**),
scRNA-Seq cell type predictions (**p**) and assigned neurogenic cell type (**q**).

1120
Extended Data Fig. 2. Visualization of GAS for critical interneuron markers. UMAP
visualization of GAS from snATAC-Seq nuclei for specified genes. Legend depicts normalized
read counts.

1124
1125
1126

1127
Extended Data Fig. 3. DA Peaks per cluster and read distribution across gene features. a,
Embryonic snATAC-Seq signal plots of DA peaks for each cluster (same cluster order as Fig 2).
Signal was binarized to indicate DA peaks from one cluster (column cluster labels) which are
detectable in any other cluster (row cluster labels). **b**, For each cluster (rows of the dot plot), motifs
representative of lineage and maturation stage, corresponding motif name and P-value are
displayed to the right of the dot plot. All motifs are in the top five hits for each cluster, with top hits
indicated with an asterisk. **c**, Bar chart of DA peak counts where -log(Fold Change) > 0 per SLM
cluster colored by neurogenic cell type. **d**, Bar graph of mean DA peak counts where -log(Fold
Change) > 0 per SLM cluster colored by neurogenic cell type and grouped by tissue. **e**, Bar chart
of DA peak counts where -log(Fold Change) > 0 per SLM cluster colored by tissue. **f**, Bar chart of
mean DA peak counts where -log(Fold Change) > 0 per SLM cluster colored by tissue and
grouped by neurogenic cell type. **G**, Pie chart of relative enrichment of DA peaks within annotated
gene features.

1141
1142

Extended Data Fig. 4. Gene activity score comparisons. UMAP visualizations of integrated snATAC-Seq and scRNA-Seq data colored by assay (left) and tissue samples (right). For integration, snATAC-Seq Gene Activity Score (GAS) was quantified using four different strategies: (**a**) quantified peaks within gene promoters (-2000 bp to 0 bp of 5' transcript coordinate) and gene bodies, (**b**) quantified peaks within gene promoters and the first exon of all transcripts, (**c**) quantified peaks within gene promoters only, and (**d**) quantified peaks within promoters, the first exon of all transcripts, and any associated putative enhancers (H3K27ac+ chromatin).

1150
1151

Extended Data Fig 5. Visualization of RNA counts and GAS following integration. UMAP
visualization of transcript abundance from scRNA-Seq cells (left) and GAS from snATAC-seq
nuclei (right) for genes enriched in specific brain regions (**a-j**) or in distinct neurogenic cell types
(**k-p**).

1152
1153
1154
1155

1162

Extended Data Fig. 6. Characterization of enhancers in embryonic neurogenic tissue. a,
Gene-enhancer interactions originating at the *Sox2* locus as defined by Cicero (cyan arcs),
SnapATAC (magenta arcs), the combination of Cicero and SnapATAC (black arcs) and the
intersection of Cicero and SnapATAC (dark blue arcs, representing high confidence enhancers).
High confidence enhancers ("Intersection") were used for further analyses. Window is 0.5 Mb
centered on TSS. Arc height represents interaction score. **b-e**, Visualization of intersection of
Cicero and SnapATAC gene-peak interactions for *Nes* (**b**), *Neurog2* (**c**), *Gsx2* (**d**) and *Dlx1* (**e**).
Window is 100 Kb, centered on the gene TSS. Peaks: displays co-accessible coordinates,
Encode: displays H3K27ac signal from E12.5 forebrain from ENCODE ChIP-Seq data, VISTA:
enhancers with positive activity in the forebrain and midbrain from the VISTA Enhancer Browser.
f, Bar chart depicting the total number of gene-peak interactions detected by Cicero, SnapATAC
and high-confidence interactions via the intersection of Cicero and SnapATAC ("merged" label).
G, Bar chart of the number of gene-peak interactions binned by gene-peak distance for high-
confidence gene-peak interactions.

1177
Extended Data Fig. 7. degPatterns analysis and heatmaps of RNA, GAS and enhancer
counts for top 500 highly variable genes. a-b, Line graphs of genes within each cluster
detected by degPatterns that contained > 5 genes per cluster using embryonic integrated
snATAC-Seq/scRNA-Seq data. Y-axis is Z-score for RNA, GAS, or enhancers counts per gene.
X-axis is binned pseudotime periods with origin corresponding to early development and higher
pseudotime values corresponding to more differentiated cells. RNA, GAS, or enhancers for each
gene in this cluster are plotted individually. **c**, Bar graph of change of enhancer usage for all
detected H3K27ac+ presumptive enhancers associated with DEGs per state transition: AP to BP,
BP to N and AP to N. All detectable H3K27ac+ presumptive enhancers defined as genomic
intersection of 1) the genomic union of SnapATAC cREs and Cicero cREs with co-accessibility
score >0.25) and 2) E12.5 ENCODE H3K27ac ChIP-Seq peaks. **d**, Bar graph of ratio of enhancer
activation versus enhancer deactivation per state transition: AP to BP, BP to N and AP to N. **e-g**,
Heatmaps of RNA (**e**), GAS (**f**) and enhancer (**g**) counts for top 500 highly variable genes from
scRNA-Seq data and corresponding genes from snATAC-Seq GAS and enhancer matrices.
Heatmap columns were ordered by hierarchical clustering with correlation distance and average
linkage. Rows were ordered by pseudotime (assigned by Moncole3).

 SUPPLEMENTARY TABLES