Somewhere I belong: phylogeny and morphological evolution in a species-rich lineage of

ectoparasitic flatworms infecting cichlid fishes.

Armando J. Cruz-Laufer[1], Antoine Pariselle[2,3], Michiel W. P. Jorissen[1,4], Fidel Muterezi

Bukinga[5], Anwar Al Assadi[6], Maarten Van Steenberge[7,8], Stephan Koblmüller[9], Christian

Sturmbauer[9], Karen Smeets[1], Tine Huyse[4,7], Tom Artois[1], Maarten P. M. Vanhove[1,7,10]

1 UHasselt – Hasselt University, Faculty of Sciences, Centre for Environmental Sciences,

Research Group Zoology: Biodiversity and Toxicology, Agoralaan Gebouw D, 3590

Diepenbeek, Belgium.

2 ISEM, Université de Montpellier, CNRS, IRD, Montpellier, France.

3 Faculty of Sciences, Laboratory "Biodiversity, Ecology and Genome", Research Centre

"Plant and Microbial Biotechnology, Biodiversity and Environment", Mohammed V

University, Rabat, Morocco.

4 Department of Biology, Royal Museum for Central Africa, Tervuren, Belgium.

5 Section de Parasitologie, Département de Biologie, Centre de Recherche en Hydrobiologie,

Uvira, Democratic Republic of the Congo.

6 Fraunhofer Institute for Manufacturing Engineering and Automation IPA, Nobelstraße 12,

70569 Stuttgart, Germany.

7 Laboratory of Biodiversity and Evolutionary Genomics, KU Leuven, Charles

Deberiotstraat 32, B-3000, Leuven, Belgium.

8 Operational Directorate Taxonomy and Phylogeny, Royal Belgian Institute of Natural

Sciences, Vautierstraat 29, B-1000 Brussels, Belgium.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

9 Institute of Biology, University of Graz, Universitätsplatz 2, 8010, Graz, Austria.

10 Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2,

CZ-611 37, Brno, Czech Republic.

Corresponding author: Armando J. Cruz-Laufer, armando.cruzlaufer@uhasselt.be

ABSTRACT

A substantial portion of biodiversity evolved through adaptive radiation. However, the effects

of explosive speciation on species interactions remain poorly understood. Metazoan parasites

infecting radiating host lineages could improve our knowledge because of their intimate host

relationships. Yet limited molecular, phenotypic, and ecological data discourage multivariate

analyses of evolutionary patterns and encourage the use of discrete characters. Here, we

assemble new molecular, morphological, and host range data widely inferred from a species-

rich lineage of parasites (*Cichlidogyrus*, Platyhelminthes: Monogenea) infecting cichlid

fishes to address data scarcity. We infer a multi-marker (28S/18S rDNA, ITS1, COI mtDNA)

phylogeny of 58/137 species and characterise major lineages through synapomorphies

inferred from mapping morphological characters. We predict the phylogenetic position of

species without DNA data through shared character states, a combined molecular-

morphological phylogenetic analysis, and a classification analysis with support vector

machines. Based on these predictions and a cluster analysis, we assess the systematic

informativeness of continuous characters, search for continuous equivalents for discrete

characters, and suggest new characters for morphological traits not analysed to date. We also

model the attachment/reproductive organ and host range evolution using the data of 136/137

described species and multivariate phylogenetic comparative methods (PCMs). We show that

discrete characters can mask phylogenetic signals but can be key for characterising species

groups. Regarding the attachment organ morphology, a divergent evolutionary regime for at

least one lineage was detected and a limited morphological variation indicates host and

environmental parameters affecting its evolution. However, moderate success in predicting

phylogenetic positions, and a low systematic informativeness and high multicollinearity of

morphological characters call for a revaluation of characters included in species

characterisations.

KEY WORDS

Adaptive radiation; *Cichlidogyrus*; Dactylogyridae; Monogenea; systematic informativeness.

ACKNOWLEDGEMENTS

We thank J.-F. Agnèse (Institut de Recherche pour le Développement & Université de

Montpellier) for taking responsibility for part of the molecular work, and F. A. M. Volckaert

(KU Leuven) and J. Snoeks (Royal Museum for Central Africa & KU Leuven) for their

guidance since the early stages of this research. J. Bamps, A. F. Grégoir, L. Makasa, J. K.

Zimba (Department of Fisheries), C. Katongo (University of Zambia), T. Veall, O. R.

Mangwangwa (Rift Valley Tropicals), V. Nshombo Muderhwa, T. Mulimbwa N'sibula, D.

Muzumani Risasi, J. Mbirize Ndalozibwa, V. Lumami Kapepula (Centre de Recherche en

Hydrobiologie-Uvira), the Schreyen-Brichard family (Fishes of Burundi), F. Willems

(Kasanka Trust), S. Dessein (Botanic Garden Meise), A. Chocha Manda, G. Kapepula

Kasembele, E. Abwe, B. Katemo Manda, C. Mukweze Mulelenu, M. Kasongo Ilunga Kayaba

and C. Kalombo Kabalika (Université de Lubumbashi), M. Collet and P. N'Lemvo (Institut

Congolais pour la Conservation de la Nature), D. Kufulu-ne-Kongo (École Muilu Kiawanga),

L. Matondo Mbela (Université Kongo), S. Wamuini Lunkayilakio, P. Nguizani Bimbundi, B.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Boki Fukiakanda, P. Ntiama Nsiku (Institut Supérieur Pédagogique de Mbanza-Ngungu), P.

Nzialu Mahinga (Institut National pour l'Etude et la Recherche Agronomiques—

Mvuazi/Institut Supérieur d'études agronomiques—Mvuazi) and M. Katumbi Chapwe are

thanked for administrative, field and lab support, making this study possible. We thank A.

Avenant-Oldewage, M. Geraerts, T. Gobbin, P. C. Igeh, N. Kmentová, M. Mendlová, and C.

Rahmouni for providing raw species measurements used for their respective publications.

Furthermore, we thank J. M. Mirande and T. Moons for technical advice on the use of *TNT*.

Field work was supported by travel grants V.4.096.10.N.01, K.2.032.08.N.01 and K220314N

from the Research Foundation—Flanders (FWO-Vlaanderen) (to MPMV), two travel grants

from the King Leopold III Fund for Nature Conservation and Exploration (to MPMV and

MVS), FWO-Vlaanderen Research Programme G.0553.10, the University Development

Cooperation of the Flemish Interuniversity Council (VLIR-UOS) South Initiative

ZRDC2014MP084, the OCA type II project S1_RDC_TILAPIA and the Mbisa Congo

project (2013–2018), the latter two being framework agreement projects of the RMCA with

the Belgian Development Cooperation. When data collection for this study commenced,

MVS and MPMV were PhD fellows, and TH a post-doctoral fellow, of FWO-Vlaanderen.

Part of the research leading to results presented in this publication was carried out with

infrastructure funded by the European Marine Biological Research Centre (EMBRC)

Belgium, Research Foundation – Flanders (FWO) project GOH3817N. MWPJ was supported

by the Belgian Federal Science Policy Office (BRAIN-be Pioneer Project

(BR/132/PI/TILAPIA), and a BOF Reserve Fellowship from Hasselt University. AJCL is

funded by Hasselt University (BOF19OWB02), and MPMV receives support from the

Special Research Fund of Hasselt University (BOF20TT06). We thank the reviewers

including J. M. Mirande and an anonymous reviewer for suggesting improvements to the

manuscript.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

DATA AVAILABILITY STATEMENT

The morphological data that support the findings of this study are openly available in

MorphoBank at www.morphobank.org, at https://dx.doi.org/XXXXXXXX. The DNA

sequence data are openly available in the GenBank Nucleotide Database at

https://www.ncbi.nlm.nih.gov/genbank, accession numbers XXXXXX–XXXXXX.

Phylogenetic trees and data matrices are openly available in TreeBase at https://treebase.org,

accession number XXXXXX.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

INTRODUCTION

Adaptive radiations and host-parasite interactions

Adaptive radiation is one of the most important processes of species formation. These

explosive speciation events might explain a substantial part of the biodiversity on the planet

(Glor, 2010). Adaptive radiations are characterised by a rapid diversification resulting from

adaptation to newly available ecological niches (Losos, 2010). Famous examples include

Darwin's finches (Grant, 1999), Caribbean lizards of the genus *Anolis* Daudin, 1802 (Mahler

et al., 2013), cichlid fishes (Salzburger, 2018), and Hawaiian silverworths (Landis et al.,

2018). Despite the attention these species have receive, few studies have investigated the

evolution of ecological interactions involving these groups and other organisms of other

groups beyond the feeding ecology of the former (e.g. Guerrero and Tye, 2009; Takahashi

and Koblmüller, 2011; but see Karvonen and Seehausen, 2012; Blažek et al., 2018). How do

organisms evolve that accompany the rapid diversification process of an adaptive radiation?

Metazoan parasites could provide answers to this question as they form often intimate

relationships with their hosts. Thus, their evolutionary regime can be strongly impacted by

the host's evolutionary history (Huyse et al., 2005). Multiple evolutionary regimes might

apply to parasites infecting radiating host lineages. First, parasites that are strongly dependent

on their hosts, might experience a selection pressure to remain competitive in the arms race

with the host's defences (Kaltz and Shykoff, 1998). Structures relevant to this arms race such

as attachment organs might experience a strong stabilising selection pressure as they are the

parasite's main physical connection to the host. In this scenario, the attachment organ

evolution would be heavily impacted by host and environmental parameters (Kaltz and

Shykoff, 1998). Second, parasites could follow the example of their hosts in terms of an

(adaptive) radiation. Such co-radiations have previously been reported for insects parasitic on

plants or other insects (Weiblen and Bush, 2002; Forbes et al., 2009) but quantitative support

for these hypotheses in the form of evolutionary models has not been provided to date. Third,

parasites could have structures that are not under strong selection pressure and, therefore,

their morphology might randomly diverge over time following a pattern associated with a

'random walk', i.e. genetic drift or randomly fluctuating selection (Losos, 2008).

Reproductive organ structures, e.g. in flatworms infecting the gills of cyprinid fish (Šimková

et al., 2002), have been suggested to follow this pattern causing reproductive isolation

between species.

Species-rich but missing data: a case study for host-parasite systems

Extensive evolutionary analyses often require large molecular and morphological

datasets. However, such datasets remain scarce for parasitic organisms. Metazoan parasites

are often small, which makes identifying these organisms to species level notoriously

difficult (de Meeûs et al., 2007). Furthermore, only a fraction of the known species is

genetically characterised (Poulin et al., 2019) and most species are yet to be discovered and

described (Poulin et al., 2020). Among the best-known models in adaptive radiation research,

African cichlids (Cichliformes, Cichlidae) are possibly the best-studied system (Salzburger,

2018) with approximately 2000 (described and undescribed) species reported from Eastern

Africa alone (Turner et al., 2001; Salzburger et al., 2014). This knowledge is rooted in a long

and productive tradition of international cichlid research (see Ronco et al., 2020; Van

Steenberge et al., 2011). Resulting from this scientific interest, one group of gill parasites

(Fig. 1a) infecting these fishes, the monogenean flatworms belonging to *Cichlidogyrus* sensu

Paperna, 1960 [incl. the nested genus *Scutogyrus* Pariselle & Euzet, 1995 (Wu et al., 2007),

together referred to as *Cichlidogyrus* in the following] (Fig. 1b, c), has been studied in more

depth than other species-rich parasite genera, in particular from the African continent (Cruz-

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Laufer et al., 2021). Species of *Cichlidogyrus* rival (Cruz-Laufer et al., 2021) and possibly

exceed (see Poulin, 2014) their hosts in terms of species numbers: 137 parasite species have

been described on 126 fish species (Cruz-Laufer et al., 2021). Monogenean flatworms have

provided insight into parasite speciation (Meinilä et al., 2004; Šimková et al., 2013; Vanhove

et al., 2015), population dynamics (Kmentová et al., 2021b), anthropogenic introductions

(Šimková et al., 2019; Jorissen et al., 2020) as well as host biogeography (Barson et al., 2010;

Pariselle et al., 2011; Vanhove et al., 2013, 2016). This variety of evolutionary research

paired with the model system status of the host species of species of *Cichlidogyrus*, has led to

the suggestion of the cichlid-*Cichlidogyrus* species network as model system for parasite

speciation research and the evolution of host-parasite interactions (Pariselle et al., 2003;

Vanhove et al., 2016). Recent advances in immunological (Zhi et al., 2018), pathological

(Igeh and Avenant⬜Oldewage, 2020), genomic (Vanhove et al., 2018; Caña-Bozada et al.,

2021), and microscopy research (Fannes et al., 2015) have brought the vision of a cichlid-

Cichlidogyrus model system closer to reality [see Cruz-Laufer et al. (2021) for a detailed

review of model system qualities].

The species-richness of *Cichlidogyrus* has previously been attributed to co-divergence

with their hosts (Pariselle et al., 2003; Vanhove et al., 2015), host switching (Pariselle et al.,

2003; Messu Mandeng et al., 2015), and within host speciation (Mendlová et al., 2012;

Vanhove et al., 2015). Similar to other parasites, morphological characters of monogenean

attachment and reproductive organ structures have formed an essential part of species

descriptions. Evolutionary studies have paid considerable attention to the evolutionary

patterns of these (mostly) sclerotised structures (e.g. Šimková et al., 2002; Mendlová et al.,

2012). For species of *Cichlidogyrus*, the attachment organ includes a haptor with several

anchors, hooks, and bars and reproductive organs with a male copulatory organ (MCO) and,

at times, a sclerotised vagina (Fig. 1b, c). Based on the similar phenotypes observed in related

species, some studies have suggested phylogenetic constraints, i.e. ancestry of the parasite

species, as the sole determinants of the morphology of the attachment organ in *Cichlidogyrus*

(Vignon et al., 2011). This claim has been questioned more recently. At least one species

appears to have changed its attachment organ morphology compared to its close relatives as a

response to a host switch towards non-cichlid fishes (Messu Mandeng et al., 2015). These

host switches are considered rare as monogenean representatives have been reported for 27

(Carvalho Schaeffner, 2018) of 48 (Lévêque et al., 2008) fish families in Africa and most

harbour some other dactylogyridean lineage of gill parasites (see Carvalho Schaeffner, 2018).

Despite this discovery, no study has investigated the macroevolution of the morphological

characters of the attachment and reproductive organs for almost a decade (see Mendlová et

al., 2012).

The present study aims to investigate the evolutionary processes that have shaped the

morphology of species of *Cichlidogyrus*. However, out of the 137 species that are currently

described, DNA sequences of only 18 species were included in the most recent phylogenetic

study (Messu Mandeng et al., 2015). Furthermore, morphological and ecological data have

not been collected in centralised databases inhibiting the progress of large-scale meta-

analytical studies (Cruz-Laufer et al., 2021). Previous studies have evaded this scarcity of

information by loosely defining discrete states for the attachment organ morphology and the

host repertoire base on the taxonomic literature. For instance, Vignon et al. (2011) and

Mendlová et al. (2012) grouped species by the relative size of the sclerotised structures of the

attachment organ (for an overview of the terminology, see Fig. 1c) using generic terms like

'large' and 'small' instead of coding these characters based on continuous measurements.

Similarly, Mendlová and Šimková (2014) proposed an index of specificity (IS) to group

species as species-, genus-, and tribe-specific, or generalist based on the most recent host

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

classification at the time. Yet discretisation has been known to cause information loss as the

variability of the otherwise continuous parameters is largely ignored (Altman and Royston,

2006; Goloboff et al., 2006; Parins-Fukuchi, 2018). These obstacles emphasise the need for

new approaches and more extensive and accessible datasets to gain insight into the evolution

of *Cichlidogyrus* and the phylogenetic information of morphological and ecological

characters.

To address data scarcity, we provide new morphological, molecular, and host range data,

which are made available in public databases. We perform multiple analyses to investigate

the evolution of phenotypic characters to answer the following questions:

(i) Which main lineages can we infer from the phylogeny? To which lineages do species

without molecular data belong? Which predictions can be made through character

mapping, a combined (molecular and morphological) parsimony analysis, and a machine

learning algorithm?

(ii) How systematically informative are the morphometric measurement most widely applied

to this genus? Does the use of discrete characters lead to information loss in the

morphometric and host range data?

(iii)As species of *Cichlidogyrus* infect a host lineage known for its rapid speciation, which

evolutionary processes have shaped the attachment and reproductive organ morphology?

Are these structures under stabilising selection pressure (H_2), do the parasites mirror the

radiations of their hosts either as an early (H_3) or a late burst (H_4) in the character

evolution, or do these structures follow a pattern associated with a random walk (H_1)?

Furthermore, do species infecting cichlids from the Eastern African radiations follow a

different evolutionary regime than the species infecting other hosts (H_5)?

MATERIALS AND METHODS

Sampling

Fish specimens belonging to Cichlidae Bonaparte, 1835 (Cichliformes), the

nothobranchiid genus *Aphyosemion* Myers, 1924 (Cyprinodontiformes: Nothobranchiidae),

and *Polycentropsis abbreviata* Boulenger, 1901 (Ovalentaria, *incertae sedis*: Polycentridae)

were collected during various expeditions across Africa between 2008 and 2019 and from

few specimens from aquaculture facilities in France (Appendix 1). As samples collected

during these expeditions have been included in previous studies, sampling details can be

found in the respective publications (Vanhove et al., 2011, 2013; Muterezi Bukinga et al.,

2012; Pariselle et al., 2015a, 2015b; Jorissen et al., 2018a, 2018b, 2020). We dissected the

gills of the fish and stored the samples in 96% ethanol to preserve the DNA of the parasites

attached to the gills. Then, we screened the gills for parasitic infections of flatworm species

belonging to *Cichlidogyrus*, removed the parasites from the gills using dissection needles

under a stereomicroscope, mounted them temporarily in water for species-level identification

at a magnification of 1000x (100x magnification with oil immersion and 10x ocular) and

stored the parasite specimens individually in 96% ethanol for DNA extraction. As mostly

entire specimens were used for DNA extraction, we refer to the above-mentioned articles for

type and voucher specimens from the same host individuals deposited in curated collections

(see Appendix 1).

DNA extraction, amplification, and sequencing

For the DNA extraction of recent samples, we used the Nucleospin Kit (Macherey-

Nagel, USA) following manufacturer guidelines but with 60 µl instead of 100 µl of elution

buffer added in the final step. For some samples processed at an earlier stage, we suspended

DNA samples through homogenisation with no added extraction steps following the

procedures described by Marchiori et al. (2015). We amplified and sequenced three partial

nuclear ribosomal genes including a fragment of the large subunit ribosomal DNA (28S

rDNA) with the primers C1 (5'-ACCCGCTGAATTTAAGCAT-3') and D2 (5'-

TGGTCCGTGTTTCAAGAC-3') (Hassouna et al., 1984), a fragment of the small subunit

ribosomal DNA (18S rDNA) and the internal transcribed spacer 1 (ITS1) with the primers S1

(5'-ATTCCGATAACGAACGAGACT-3') (Matejusová et al., 2001) and IR8 (5'-

GCAGCTGCGTTCTTCATCGA-3') (Šimková et al., 2003), and a fragment of the

mitochondrial gene coding for the cytochrome oxidase *c* subunit 1 protein (COI mtDNA)

with the primers ASmit1 (5'-TTTTTTGGGCATCCTGAGGTTTAT-3') (Littlewood et al.,

1997), Cox1_Schisto_3 (5'-TCTTTRGATCATAAGCG-3') (Lockyer et al., 2003), and

ASmit2 (5'-TAAAGAAAGAACATA ATGAAAATG-3') (Littlewood et al., 1997). Reaction

protocols followed the procedures of Messu Mandeng et al. (2015) for 28S rDNA, Mendlová

et al. (2012) for 18S rDNA, and Vanhove et al. (2015) for COI mtDNA. We purified PCR

products with a NucleoFast 96 PCR kit (Macherey-Nagel, USA) or with a GFX PCR DNA

kit and Gel Band Purification kit (GE Healthcare, USA) following manufacturer guidelines.

Bidirectional Sanger sequencing was conducted according to the Big Dye Terminator v3.1

sequencing protocol (Applied Biosystems, USA) at a 1:8 dilution with an ABI PRISM 3130

Avant Genetic Analyser automated sequencer (Applied Biosystems, USA) and the primers

included in the PCR protocols.

Morphological and host range data collection

We assembled morphometric (Fig. 1c) and host range data for 136 species belonging

to *Cichlidogyrus*. Raw measurements taken through light microscopy were assembled from

AP's personal morphometric database and from the raw data of previous publications kindly

provided by the authors of the cited studies (Pariselle and Euzet, 2003; Vanhove et al., 2011;

Gillardin et al., 2012; Muterezi Bukinga et al., 2012; Pariselle et al., 2013; Řehulková et al.,

2013; Van Steenberge et al., 2015; Messu Mandeng et al., 2015; Kmentová et al., 2016a,

2016b, 2016c; Rahmouni et al., 2017, 2018; Igeh et al., 2017; Geraerts et al., 2020; Gobbin et

al., 2021). These raw data were deposited at MorphoBank (www.morphobank.org, project

XXXXX). We calculated the standard errors and standard deviations for each measurement

from the raw data. However, for species where no or only partial raw data were available, we

used the mean values provided in the literature (Douëllou, 1993; Pariselle and Euzet, 2003;

Řehulková et al., 2013; Rahmouni et al., 2017; Jorissen et al., 2018b, 2018a) without errors.

If no mean was reported, we inferred the measures from drawings in the literature (Dossou,

1982; Dossou and Birgi, 1984; Birgi and Lambert, 1986; Muterezi Bukinga et al., 2012) and

calibrating measurements through the included scale bars. Measurements of *C. dionchus*

Paperna. 1968 were not included as no mean values were provided in respective publications

(Paperna, 1960, 1968; Paperna and Thurston, 1969) and drawings were incomplete with some

structures, e.g. the second marginal hook, entirely missing. For the auxiliary plate (AuP) of

the male copulatory organ, we used the surface area and its relative error resulting of the sum

of relative errors of the original measurements inferred from the plate length and width

assuming an ellipsoid shape.

Beyond these continuous characters, we assigned all species to previously suggested

discrete characters for the haptor morphology and host range, which include the configuration

of the hooks (Vignon et al., 2011), the similarity of the anchors, the shape of the ventral bar

(Mendlová et al., 2012), and the index of specificity (IS) (Mendlová and Šimková, 2014).

Hook configurations were coded according to the secondary growth in the first and third hook

pair ('small-small', 'small-large', 'large-small', 'large-large') relative to the second pair,

which retains it embryonal size in adult specimens (Llewellyn, 1963). The anchor pairs were

categorised as 'similar' or 'dissimilar' in shape and size based on the original species

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

descriptions. The ventral bar shape was defined according to the presence of membranous

extensions (see Mendlová et al., 2012) and size ('with membranous extensions', 'without

membranous extensions', 'massive with membranous extension', 'bar supports large

extended plate (*Scutogyrus*)'). We inferred IS data from the host-parasite interactions listed

by Cruz-Laufer et al. (2021) with species exclusively reported from hosts belonging to a

single species, genus, or tribe classified as species-, genus-, and tribe-specific, or generalist if

hosts belonging to two different tribes (or even families) were infected by the same parasite

species. For the classification of cichlid species into tribes, we followed Dunz and Schliewen

(2013). All discrete and continuous morphometric and host specificity data are made

available at TreeBase alongside the DNA sequence alignments and tree topologies

(www.treebase.org, accession number: XXXXX). Raw morphometric data are made

available at MorphoBank (www.morphobank.org, project XXXXX).

We also surveyed the taxonomic literature to detect additional features used to

recognise groups of related species. First, we attempted to find equivalents for characters and

character states frequently mentioned in species descriptions among the morphometrics. For

instance, the root lengths of the anchors in the attachment organ are frequently mentioned as

'large' or 'small', which can be reflected through the ratio of the two roots in each anchor

pair. If no measurement reflected the described features, we developed new discrete

variables. Detailed information on the continuous and proposed new discrete characters and

their character states can be found in Appendix 2.

Phylogenetic analyses

We assembled a four-locus concatenated multiple alignment from the sequences

generated during this study and sequences available at GenBank (see Appendix 3). The

alignment includes partial ITS1, 18S rDNA, 28S rDNA, and COI mtDNA sequence data.

Partial DNA sequence data (i.e. specimens with less than those four loci sequenced) were

included with the lacking fragments coded as missing data. We aligned the sequences of each

locus using the algorithm *L-INS-i* in *MAFFT* v.7.409 (Katoh and Standley, 2013) as

recommended for ribosomal DNA by the *MAFFT* manual, and removed poorly aligned

positions and divergent regions using the options for less stringent parameters in *Gblocks*

v0.91b (Talavera and Castresana, 2007).

We estimated tree topologies under maximum parsimony (MP) through *TNT* v1.5

(Goloboff et al., 2008b; Goloboff and Catalano, 2016) using extended implied weighting

(Goloboff, 2014) in a range of values for the concavity constant K (20, 21, 23, 26, 30, 35, 41,

48, 56). Extended implied weighting reduces the impact of characters with missing data that

were weighted artificially high in the original implied weighting method (Goloboff, 1993).

For DNA sequence data, collectively weighting all sites in a partition (gene or codon) was

suggested as a more appropriate method (Goloboff et al., 2008a). Therefore, we explored

three weighting schemes proposed by Mirande (2019): all characters weighted separately

(SEP), all characters weighted according to the average homoplasy of the marker (BLK), and

characters in the protein-coding marker, i.e. COI, weighted according to the average

homoplasy of their position with other characters were weighted as in BLK (POS). Similar to

Mirande (2009), we selected the parameter combinations (k value and weighting scheme) that

produced the most stable topology to compute a strict consensus tree. The distortion

coefficient and subtree pruning and regrafting (SPR) distance were used as selection criteria

by computing the similarity of each consensus tree obtained under the different parameters to

the rest. MP tree searches involved rounds of tree fusing, sectorial searches, tree drifting, and

tree ratchet (Goloboff, 1999; Nixon, 1999) under default settings and each round was stopped

following three hits of the same optimum. Gaps were treated as missing data. As

bootstrapping and jackknifing are reported to be distorted by differently weighted characters

(Goloboff, 2003), branch support was estimated through symmetric resampling with a

probability of change of 0.33 and values expressed as differences in frequencies (GC:

'Groups present/Contradicted'). *Cichlidogyrus berrebii* Pariselle & Euzet, 1994, *C. kothiasi*

Pariselle & Euzet, 1994, and *C. pouyaudi* Pariselle & Euzet, 1994, parasites of tylochromine

cichlids, were used to root the phylogenetic trees due to the well-documented early diverging

phylogenetic position of these species (Mendlová et al., 2012; Messu Mandeng et al., 2015).

To infer the phylogenetic position of species of *Cichlidogyrus* without DNA sequence

data, we performed a second phylogenetic analysis including the morphometric

measurements of the attachment and reproductive organs as a separate block in *TNT* and

extended implied weighting using the same set of k values and the weighting scheme that

produced the most stable tree topologies for the DNA data. The homoplasy of each

morphometric character was estimated independently as recommended by Goloboff et al.

(2006) for continuous characters. All other settings in *TNT* remained the same as above. The

full *TNT* data matrix is provided in Supporting Information (Table S1).

For downstream analyses of character evolution, we also estimated phylogenies

through Bayesian inference (BI) and maximum likelihood (ML) methods using *MrBayes*

v3.2.6 (Ronquist and Huelsenbeck, 2003) on the CIPRES Science Gateway online server

(Miller et al., 2010) and *IQ-Tree* v1.6.12 (Nguyen et al., 2015) respectively. We used these

model-based approaches (BI and ML) to provide a consistent approach to the downstream

multivariate phylogenetic comparative analyses, which are themselves model-based. DNA

sequence data were partitioned by gene and, for the COI mtDNA, by codon position. We

selected the substitution models for each partition according to the Bayesian information

criterion (BIC) as rendered by *ModelFinder* in *IQ-TREE* (Kalyaanamoorthy et al., 2017)

using partition merging (Chernomor et al., 2016) (Appendix 4). For BI analyses, we selected

only models implemented in MrBayes (Appendix 4). We used two parallel runs and four

chains of Metropolis-coupled Markov chain Monte Carlo iterations. We ran 20 million

generations with a burn-in fraction of 0.25 and sampled the trees every 1000^{th} generation. We

checked convergence criteria by assessing the average standard deviation of split frequencies

(< 0.01 in all datasets) and the effective sample size (> 200) using *Tracer* v1.7 (Rambaut et

al., 2018). For ML analyses, we estimated branch support values using both ultrafast

bootstrap approximation (Hoang et al., 2018) and Shimodaira-Hasegawa-like approximate

likelihood ratio tests (SH-aLRT) (Guindon et al., 2010) with 1000 replicates as recommended

by the *IQ-Tree* manual. We considered a BI posterior probability (PP) ≥ 0.95, an ultrafast

bootstrap values ≥ 95, and SH-aLRT statistic ≥ 80 as well-supported (Hoang et al., 2018). We

plotted the graphs and phylogenetic trees using the *R* packages *ggplot2* v3.3.5 (Wickham,

2016) and *ggtree* v3.13 (Yu et al., 2017, 2018).

To verify congruence of the final hypothesis of the parsimony analysis (molecular

data) with the BI and ML consensus trees, we analysed the congruence of the MP, BI, and

ML tree topologies using the Congruence Among Distance Matrices (CADM) tests

(Legendre and Lapointe, 2004; Campbell et al., 2011). We used the package *ape* v5.3

(Paradis and Schliep 2019) in *R* v4.1.0 (R Core Team, 2021) to calculate phylogenetic pair-

wise distance matrices and to conduct the CADM test.

Clade affiliation and discriminative power of morphometrics: statistical classification,

literature review, and cluster analysis

Beyond the clades supported in the combined MP analysis, we also used two

additional approaches to assess the phylogenetic position of species of *Cichlidogyrus* that

have not been sequenced to date. First, we characterised the morphology and host repertoires

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

of the clades inferred from the molecular phylogeny (Fig. 2) based on character maps of the

continuous and discrete morphological characters and the host repertoires surveyed in the

literature. To map continuous characters on the consensus MP tree, we estimated ancestral

character states through the function *anc.ML* in the *R* package *phytools* v0.7-80 (Revell,

2012). To map the discrete characters, we estimated ancestral states under maximum

parsimony with all rates set to equal as we could not make any assumption on the transition

costs between character states. This analysis was implemented in the function

asr_max_parsimony in the *R* package *castor* v1.6.9 (Louca and Doebeli, 2018). All character

maps were plotted using the *ggplot2* and *ggtree*. Species of *Cichlidogyrus* not included in

molecular phylogenies were assigned to the clades based upon the measurements and their

character states while taking host repertoires (by tribe of cichlids or family of non-cichlids)

into consideration. These last criteria (natural and invasive host repertoires) were taken into

account as we expected the host environment to be a relevant factor in speciation processes of

these parasites (see McCoy, 2003; Huyse et al., 2005). We used the host-parasite list and

literature databases provided by Cruz-Laufer et al. (2021) to summarise the host repertoires.

Second, we predicted clade affiliation through a supervised machine learning

algorithm using support vector machines (SVMs). Machine learning can be used to assess of

the predictive power of morphometric measurements in taxonomic studies, in particular with

support vector machines (SVMs) (e.g. Zischke et al., 2016; Fang et al., 2018). In these

supervised machine learning approaches, a dataset is provided to train the learning algorithm

to classify individuals in distinct groups. The predictions generated by the algorithm are then

validated against a set of samples with a known affiliation. The performance can then provide

an insight into the predictive power of the input data. Subsequently, the optimised algorithm

can be applied to a test dataset of samples with unknown group affiliation. Here, we carried

out the SVM analysis with a radial basis kernel function as implemented in the method

svmRadial in the *R* package *caret* (Kuhn, 2008; Meyer et al., 2020). Missing data in the

measurements were imputed through k-nearest neighbour imputation, scaled, and centred as

implemented in the function *preProcess*. The *C* and σ parameters of the kernel function were

optimised through a grid-search with exponentially growing sequences as suggested by Hsu

et al. (2003) and through tenfold cross-validation with ten repetitions. As specimens

belonging to different clades were observed in unequal numbers (class imbalance), we

optimised the parameters based on Cohen's κ, a multiclass performance metric accounting for

class imbalance (Landis and Koch, 1977). We considered κ < 0.2 a *slight*, κ between 0.2 and

0.4 as *fair*, κ between 0.4 and 0.6 as *moderate*, κ between 0.6 and 0.8 as *substantial*, and κ

above 0.8 as *almost perfect* agreement (Landis and Koch, 1977). Prediction provided by the

optimised algorithm were compared to the clade affiliations provided the combined

molecular-morphological parsimony analyses and the literature survey. Finally, we inferred

the variable importance for SVM predictions through a pairwise receiver operating

characteristic (ROC) curve analysis as implemented in the function *filtervarImp*. We

considered an area under the curve (AUC) between below 0.7 *poor*, between 0.7 and 0.8

acceptable, between 0.8 and 0.9 *excellent*, and above 0.9 *outstanding*.

To further assess the systematic informativeness of the morphometrics, we

investigated the amount of possible redundancies by identifying clusters of multicollinear

morphometric measurements. We used a Pearson pairwise correlation matrix (Dormann et al.,

2013) and the *ward.d2* clustering algorithm (Murtagh and Legendre, 2014) to detect clustered

variables in the *R* package *ComplexHeatmap* v2.8.0 (Gu et al., 2016). If the absolute values

of the Pearson correlation coefficients (|r|) exceeded 0.7 (Dormann et al., 2013), we

considered measurements multicollinear. Hence, we split the variables into clusters along the

dendrograms provided by *ward.d2* until all coefficients within the same cluster exceeded this

threshold. Heatmaps were plotted using the *R* package *ComplexHeatmap* v2.8.0 (Gu et al.,

2016).

Character evolution of the attachment and reproductive organs and phylogenetic signal

Using the morphometric measurements, we tested for possible patterns of a random walk

(H_1), stabilising selection (H_2), and adaptive radiation with a decelerating (H_3) and

accelerating divergence of characters (H_4) on the evolution of the attachment and

reproductive organs. We also tested if species from the East African lakes followed a

different evolutionary regime (H_5) as a result of the multiple host radiations in this region. To

detect these patterns, we employed multivariate phylogenetic comparative methods as

implemented in the *R* package *mvmorph* v1.1.4 (Clavel et al., 2015) to account for potential

interactions between characters. This software package addresses the sensitivity of previous

multivariate approaches (e.g. Khabbazian et al., 2016; Goolsby et al., 2017) to trait

covariation, data orientation, and trait dimensions (Adams and Collyer, 2018).

We fitted a range of multivariate models of continuous character evolution on a single

sample of 100 randomly selected tree topologies drawn from the post–burn-in phase of the BI

analysis of the molecular markers. Morphometric measurements were averaged by species.

Furthermore, we excluded all but one specimen per species from the phylogenetic trees to

avoid that specimens of the same species are assigned identical values, which might

otherwise artificially decrease the estimated variability of the measurements per taxon

affecting model performance. Specimens included in the subset trees, were chosen at random

and are highlighted in Appendix 3. The tested models include:

• The Brownian motion model (BM) (Felsenstein, 1973) simulates a 'random walk', i.e.

genetic drift or randomly fluctuating selection (see Hansen and Martins, 1996) (H_1). In

this case, the parasite morphology would not be affected by strong selection pressures and

would randomly diverge over time.

• The Ornstein-Uhlenbeck model (OU) (Butler and King, 2004) approximates a character

evolution under stabilising selection (H_2). Here, the parasite morphology experiences a

selection pressure towards a selective optimum due to a high host-dependence.

• The Early-Burst (EB) (Harmon et al., 2010) or accelerate-decelerate model (ACDC)

(Blomberg et al. 2003) models an early rapid evolution of characters followed by a slow-

down such as might occur during adaptive radiation events (H_3). Here, the parasite

morphology has rapidly diverged possibly mirroring the explosive speciation events

reported for some of the host lineages.

• The Late-Burst model (LB), a variation of the ACDC model with an accelerating

divergence of characters (Blomberg et al., 2003) (H_4), simulates evolution as expected

under a recent, ongoing radiation event. The speciation of the parasites would mirror the

host radiations but would still be ongoing.

• The multi-rate Brownian motion model (BMM) represents two different multi-selective

and multi-rate Brownian motion regimes for East African parasite lineages and other

lineages (O'Meara et al., 2006). Here, the parasite morphology would have undergone an

evolutionary rate shift because species infecting the host radiations in Eastern African

have developed differently than their congeners in the rest of Africa. The regimes were

defined using the function *paintSubTree* in the *R* package *phytools* v0.7-80 (Revell,

2012).

We fitted all models using the *mvgls* function (Clavel et al., 2019) in *mvmorph* as

recommended for independently evolving measurements with a natural orientation.

Therefore, we used a restricted maximum likelihood (REML) estimation with a leave-one-out

cross-validation of the penalized log-likelihood (PL-LOOCV), an estimated within-species

variation, an archetypal ridge estimator, and a diagonal unequal variance target matrix

(Clavel et al., 2019). To run the LB model with *mvgls*, we set the upper bound of the EB

search algorithm to 10. For the BMM model, we defined a separate regime for the *EAR* clade

(Fig. 2), which encompasses species infecting cichlids of the East African radiation (Fig. 3b).

As traditional information criteria such as the Akaike information criterion (AIC) are not

applicable to penalised likelihoods (Clavel et al., 2019), we assessed the model performance

with the extended information criterion (EIC) (Ishiguro et al., 1997; Kitagawa and Konishi,

2010) using the function *EIC* in *mvmorph* with 100 bootstrap replicates. Smaller EIC values

were considered indicative of a better model performance. To test for phylogenetic signal in

the datasets, we also inferred Pagel's λ (Pagel, 1999) through the *lambda* option in *mvgls*.

Finally, we modelled the character evolution of discrete characters proposed in previous

studies (see *Morphological and host range data collection*) under a univariate Pagel's λ

model (Pagel, 1999) using the *fitDiscrete* function in the *R* package *geiger* v2.0 (Pennell et

al., 2014) to assess the effects of data discretisation. We applied this function to the same set

of 100 randomly selected BI subset trees fitted to the morphometric data to account for

phylogenetic uncertainty. Model performance was assessed through the sample size–

corrected Akaike information criterion in relation to a white noise model (ΔAICc) that

assumes absolute phylogenetic independence (Pennell et al., 2014). We also produced

character maps for the discrete morphological characters (HC, VBS, AS) and the equivalent

sets of continuous characters using *asr_max_parsimony*, *anc.ML*, *ggplot2*, and *ggtree* as

mentioned above.

RESULTS

Phylogenetic analyses: Molecular data

We generated 60 sequences for 33 species and 64 specimens of *Cichlidogyrus* and

Scutogyrus including 47 28S, 14 combined 18S and ITS1 rDNA, and five COI mtDNA

sequences. For 21 of these species, these sequences are the first to be published representing

a 57% increase in taxon coverage to 42% of all currently described species, 45% of described

species of *Cichlidogyrus* and 71% of *Scutogyrus*. The alignments include 93 sequences/820

base pairs for 28S rDNA, 54 sequences/481 base pairs for 18S rDNA, 57 sequences/383 base

pairs for ITS rDNA, and 19 sequences/489 base pairs for COI mtDNA. The combined data

set includes sequences of 103 specimens belonging to 58 species (see Appendix 3), and has a

length of 2173 base pairs following the removal of poorly aligned positions and divergent

regions (missing sequence data were treated as gaps). We deposited sequences generated for

this study in GenBank (XXXXXX – XXXXXX). All sequences are listed in Appendix 3

including the respective GenBank accession numbers, the host species, and the sampling

locations. The concatenated alignments and MP, BI, and ML tree topologies can be accessed

at Treebase (www.treebase.org, accession number: XXXXX). Substitution models used for

the different partitions are provided in Appendix 4.

The BLK weightings scheme produced the most stable tree topologies for the

molecular markers (distortion coefficients and SPR distances approximately 1.000) for all k

values. For the combined parsimony analysis, k = 48 produced the most stable tree topology

(distortion coefficient: 0.901; SPR distance: 0.743 with 46 SPR moves). Thus, the final

hypotheses (molecular and combined trees) were inferred from the strict consensus the trees

produced under these parameters (BLK; k = 48) (Fig. 2a; Fig. 4). Model selection for BI and

ML analyses resulted in the merging of all COI codon positions. The BI and ML

concatenated tree topologies were congruent with the final MP consensus tree (Kendall's W

= 0.76, χ^2 = 12066, p < 0.01). Thus, we display support values of the ML analysis (UFBoot)

alongside the posterior probabilities in the BI phylogram (Fig. 2b).

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

We found 11 well-supported clades within *Cichlidogyrus* based on the MP analysis of

the molecular dataset (Fig. 2a). These clades were also supported the BI and ML analyses

suggesting a high stability of the phylogenies produced by the molecular dataset provided

here. The clades and the their respective node support values (GC value/PP/UFBoot/SH-

aLRT) are (Fig. 2a): the *'EAR'* clade (27/1/99/99) (#1), the *'CPO'* clade (91/1/100/100) (#2),

the *'Hemi'* clade (94/1/100/100) (#3), the *'Tilapiae'* clade (99/1/100/100) (#4), the *'Oreo1'*

clade (76/1/99/98) (#5), the *'Halli'* clade (100/1/100/100) (#6), *Scutogyrus* (86/0.99/*/90)

(#7), the *'Cop'* clade (48/1/100/100/*) (#8), the *'Bulb'* clade (92/1/100/100) (#9), the *'Oreo2'*

clade (81/1/97/94) (#10), and the basal *'Tylo'* clade (100/1/100/100) (#11) (see

Characterisation of species groups for details of morphology and host range). Beyond the

well-supported monophyly of all species excluding the *Tylo* group (100/1/100/100), the MP

tree shows the *Bulb* clade as sister group to all the other clades (51/*/*/*). Furthermore, *C.*

tilapiae and *C. halli* are positioned as sister clades of *Hemi* and *Oreo1* respectively albeit

with moderate support (GC values of 11 and 9). The BI and MP analyses also provides partial

support for two main lineages containing *Oreo1*, *Cop*, *Scutogyrus*, and *Hemi* (0.99/*/*/*) or

Cop, *Scutogyrus*, and *Hemi* (*/*/87/*). Both main lineages would also include *C. tilapiae*

Paperna, 1960.

Literature survey and character mapping

For a more extensive classification of species of *Cichlidogyrus* into the proposed

species groups, we developed four new discrete characters for the reproductive organs: the

shape of the penis, its diameter, the shape of the accessory piece, and the shape of the

sclerotised vagina (for the respective character states, see Appendix 2). We mapped all

morphometric and the newly proposed discrete characters on the molecular MP phylogeny

(Fig. 5). Out of the 11 species groups (Fig. 2a), we found shared morphological

characteristics in eight. Species of the *Oreo2* and the *EAR* groups shared no apparent features

in their attachment and reproductive organs with species of the same group apart from the

characteristics mentioned in the genus diagnosis (Pariselle and Euzet, 2009). We were able to

affiliate all but 12 species without available DNA sequences to these clades (Appendix 5)

based on character states shared with species included in the molecular phylogeny. Some

morphological structures were particularly relevant as synapomorphies of the species groups

(Appendix 6; Appendix 7; also Fig. 5a and Fig. 6). Based on the number of species groups

defined by a structure, the male copulatory organ (MCO) was the most distinctive structure as

phenotypes were specific to eight species groups (and several subgroups) (Fig. 5b; Fig. 6;

Appendix 6). The first hook pair (U1) was enlarged in four species groups (Fig. 6) and one

subgroup of the morphologically diverse *EAR* group. The length of the auricles of the dorsal

bar (DB h) was distinct in two species groups (Fig. 6) and one subgroup of the *EAR* group.

Hook pairs 3–7 were enlarge in two groups (Fig. 6). For the detailed morphological

characterisations of all mentioned groups and a discussion on the host and geographical

ranges, see *Characterisation of species groups* as well as several character maps (Fig. 5) and

an overview of the morphological features (Fig. 6) (full species names are cited in Appendix

5 and reported alongside host names; for abbreviations of morphometric measures, see Fig.

1c).

Characterisation of species groups

In the following section we summarise the results of the literature survey. All continuous

and discrete characters and their states are explained in Appendix 2. Ancestral character

states for the different species groups (and subgroups) are listed in Appendix 6 and Appendix

7. The following characterisations also discuss morphological differences to species groups

with similar characteristics. We also refer to publications that have previously mentioned or

discussed shared characteristics of these species groups. For 12 species, we found no

morphological similarities that could place them near any of the species groups characterised

above.

1. *EAR: Species infecting cichlids from the East African Radiation.* The *EAR* species

group comprises a large heterogeneous group of species that infect cichlids from East African

lineages, i.e. the East African Radiation (Schedel et al., 2019). The *EAR* clade consists of

multiple potential subgroups based on morphological features (Fig. 6) and node support

values from the phylogenetic analyses. The subgroups of the *EAR* group include species with

a long heel infecting non-haplochromine cichlids such as bathybatine, ectodine,

boulengerochromine, and possibly cyphotilapiine cichlids (*Heel*), species infecting

cyprichromine and trematocarine and potentially eretmodine cichlids (*CT*), and species

infecting tropheine cichlids (*Troph*) (see Fig. 6). The morphology of the attachment and

reproductive organs vary substantially between and within the subgroups (Fig. 5b). Hence,

we observed no characteristic features for the whole group with rather general and simple

ancestral features (e.g. small marginal hooks; straight, simple penis; no sclerotised vagina,

see Appendix 6 and Appendix 7). However, some subgroups display the following shared

morphological features.

• *Troph* (90/*/99/80). These species share few similarities and were placed in a subgroup

due to the host species, which all belong to the tropheine radiation. Species without

available DNA sequences, infecting tropheines, and sharing morphological features with

species in this group, were also added to this subgroup. Dorsal anchor unlike ventral with

elongated inner root (except for *C. antoineparisellei*) and deeper indentation between the

roots. However, indentation generally shallow, sometimes with fused roots (see parasites

of *Interochromis loocki* including *C. antoineparisellei*, *C. buescheri*, *C.*

schreyenbrichardorum, *C. vealli* on the ventral and *C. antoineparisellei* and *C. buescheri*

on the dorsal anchor) (Fig. 6). Dorsal bar with well-developed but not elongated auricles.

Hooks generally small but can be slightly more developed (e.g. *C. irenae, C. gistelincki,*

C. masilyai, C. salzburgeri). The MCO consists of penis and accessory piece, both

variably shaped. Penis thin and arched (*C. antoineparisellei, C. buescheri*) or looped (*C.*

schreyenbrichardorum) or slightly broadened (*C. vealli*) (Fig. 5b; Fig. 6) for parasite of

Interochromis loocki or thin and slightly sinuous for *C. gistelincki*. Penis broadened to

wide and short (reminiscent of *C. halli*) for other species. Accessory piece simple,

elongated with few remarkable structures, sometimes with cap-like (*C. gistelincki*),

forked (*C. antoineparisellei, C. buescheri, C. masilyai, C. salzburgeri*), or hook-shaped

(*C. raeymaekersi*) distal end.

• *Heel* (97/1/99/97). Dorsal anchor unlike ventral with elongated inner root. Dorsal bar with

short auricles. Well-developed first hook pair. Male copulatory organ with short penis

sometimes with spirally screw thread–like thickened wall (*C. casuarinus, C. centesimus,*

C. nshomboi), characteristic elongated heel (Fig. 5a) of one third to twice the length of the

penis, and reduced filiform or sometimes missing (only *C. centesimus*) accessory piece

(Fig. 5b; Fig. 6). The species of the *Heel* subgroup have previously been grouped together

(Muterezi Bukinga et al., 2012), most recently by Rahmouni et al. (2018b).

• *CT* (97/1/100/90). Dorsal and ventral anchors similar in shape but can vary in size (see *C.*

brunnensis) with deep to shallow (*C. brunnensis*) indentation. Dorsal bar with short

auricles. Hooks generally short. Male copulatory organ consists of penis and accessory

piece and sometimes a marked heel (*C. brunnensis*). Penis medium-length, broadened,

and mostly straight (*C. brunnensis, C. evikae, C. jeanloujustinei, C. milangelnari, C.*

sturmbaueri) with thickened wall. Accessory piece with bifurcate distal end (*C.*

brunnensis), or in two-parts (*C. evikae, C. jeanloujustinei, C. milangelnari*). Similarities

of some species of this subgroup with species infecting cyphotilapiine and ectodine

cichlids with elongated auricles (*C. adkoningsi*, *C. discophonum*, *C. glacicremoratus*, *C.*

koblmuelleri, *C. makasai*, *C. vandekerkhovei*) or elongated marginal hooks (*C.*

rectangulus, *C. sturmbaueri*) have previously been discussed (Rahmouni et al., 2017,

2018) but the phylogenetic analysis provided no support for a monophyletic group

including species of the *CT* subgroup and *C. vandekerkhovei* (Fig. 2a).

• *Other species*. The remaining species of the *EAR* species group shared no evident features

with any of the proposed subgroups. *Cichlidogyrus attenboroughi* appears to be the sister

taxon to the *Heel* group with moderate support (56/0.99/*/*) but shares no notable

characteristics with these species. Several species infecting cyphotilapiine and ectodine

cichlids display a broadened (*C. glacicremoratus*, *C. koblmuelleri*) or straight (*C.*

adkoningsi, *C. rectangulus, C. sturmbaueri*) penis with an accessory piece in two-parts

(*C. glacicremoratus*, *C. koblmuelleri*) or with a bifurcated distal end (*C. makasai*, *C.*

rectangulus, C. sturmbaueri, *C. vandekerkhovei*) similar to species of the *CT* group. Yet

many of these species also display elongated auricles at the dorsal bar (*C. adkoningsi*, *C.*

discophonum, *C. glacicremoratus*, *C. koblmuelleri*, *C. makasai*) similar to *C.*

vandekerkhovei, which appears to be unrelated to species of the *CT* group (Fig. 2a) but

also displays a bifurcate distal end. According to the phylogenetic analyses (Fig. 2a), *C.*

consobrini and *C. gillardinae* form a monophyletic group. However, we found few shared

characteristics except for a simple tubular penis with a small heel and a simple accessory

piece. These characteristics might place these two species close to *C. haplochromii* and *C.*

longipenis, both of which have been reported from haplochromine cichlids in Lake

Victoria (Pariselle and Euzet, 2009). Yet these simple features might also reflect ancestral

character states of the *EAR* group (Appendix 6; Appendix 7).

2. *CPO: Species infecting coptodonine, pelmatolapiine, oreochromine and other cichlids*.

Dorsal anchor with inner root longer than outer root and V-shaped indentation between the

roots but inner root of *C. arthracanthus* considerably longer than of other species. Roots of *C.*

cubitus and *C. louipaysani* fused. Ventral anchor similar shape to dorsal anchor but slightly

larger. However, roots of ventral anchor of *C. tiberianus* fused. Dorsal bar slightly arched

with large medium-sized to large auricles. Ventral bar always with membranous extensions.

First hook pair (U1) small and hook pairs 3–7 (U3–7) very long, i.e. more than triple the size

of first pair (Fig. 5a; Fig. 6) but U1 of *C. arthracanthus* large and U3–7 of *C. cubitus* and *C.*

louipaysani short (Fig. 6). The MCO consists of long, arched, or sometimes spiralled (*C.*

arthracanthus), tubular penis with a well-marked irregularly shaped heel, and a massive,

roughly S-shaped accessory piece (Fig. 5b; Fig. 6) that is frequently connected to the heel.

The accessory piece has an extension or thickening at the first turn in the proximal half and

frequently displays a folded back (*C. paganoi, C. vexus*), straight and pointy (*C. guirali*), or

hook-like (*C. bilongi, C. douellouae, C. ergensi, C. gallus, C. legendrei, C. microscutus*)

distal end, or sometimes additional terminations resulting in a furcate ending with two (*C.*

aegypticus, C. agnesi, C. anthemocolpos, C. bouvii, C. cubitus, C. flexicolpos, C. lemoallei,

C. louipaysani, C. ouedraogoi, C. testificatus, C. thurstonae, C. tiberianus) or three (*C.*

bonhommei, C. hemi, C. kouassii) digitations. However, the first turn is never V-shaped or

knee-like as in species of the *Hemi* clade and the hook-shaped termination is never sickle-like

such as in species of the *Cop* clade. Several species display an auxiliary plate (AuP) close to

the distal end of the MCO (Fig. 5a) including *C. aegypticus, C. agnesi, C. bilongi, C. gallus,*

C. guirali (two pieces), *C. microscutus, C. paganoi,* and *C. thurstonae*. A sclerotised vagina

is mostly present but has not been reported for *C. arthracanthus* (Paperna, 1960; Ergens,

1981). Other species with more developed hook pairs 3–7 include *Cichlidogyrus*

bulbophallus, C. flagellum, C. lobus, and *C. ranula,* some species of the *Hemi* clade, and all

species of the *Heel* subgroup of the *EAR* clade. *Cichlidogyrus bulbophallus* and species of the

Hemi clade and the *Heel* subgroup have been likened to *C. arthracanthus* because of the

additional well-developed first hook pair (U1) (Geraerts et al., 2020). However, all of these

species differ from the confirmed clade members as follows.

• The dorsal and ventral anchors differ in shape and size but are similar for species of the

CPO clade.

• The penis is never long, arched, and tubular but short (*Heel*), bulbous (*C. bulbophallus,*

C, flagellum, C. ranula), or draws a loop (Fig. 6) or long spiral (*Hemi*).

• The accessory piece is not complex and S-shaped with an extension in the proximal half

but mostly reduced (*Heel*), simple (*C. bulbophallus, C, flagellum, C. ranula*), or has a

sharp knee-like bend (*Hemi*).

Previous phylogenetic studies (Mendlová et al., 2012; Messu Mandeng et al., 2015) have

placed *C. cubitus* alongside species of the *CPO* clade with long marginal hooks ("*tiberianus*"

group, see Pouyaud et al., 2006) but morphological similarities were not further discussed.

3. *Hemi: Species infecting hemichromine cichlids and non-cichlid fishes.* Dorsal

anchor with elongated narrow inner root and rounded outer root. Ventral anchor slightly

larger with short and robust inner root. Dorsal bar slightly arched with short auricles. Ventral

bar arched with small membranous extensions. First hook pair (U1) large and well-developed

(Fig. 5a, Fig. 6) except for *C. amieti*. Hook pairs 3–7 medium-sized to large with bases more

developed than for second hook pair. The MCO consists of thin and long penis that

frequently forms loops (*C. amieti, C.* cf. *bychowskii, C. dracolemma, C. inconsultans, C.*

kmentovae, C. nandidae) (Fig. 6) or spirals with one (*C. calycinus, C. teugelsi*) to multiple

turns (*C. euzeti, C. longicirrus, C. polyenso, C sanseoi*) and re-joins the accessory piece in its

distal portion, and an accessory piece of two distinct portions. These portions can be shaped

like a V or a simple spiral with an expanded knee-like bend (*C. amieti, C. calycinus, C.* cf.

bychowskii, C. dageti, C. dionchus, C. dracolemma, C. falcifer, C. kmentovae, C. nandidae,

C.teugelsi) (see Dossou and Birgi, 1984; Pariselle and Euzet, 2004), or a large spiral followed

by a non-spiralled distal portion (*C. euzeti, C longicirrus, C. polyenso, C. sanseoi*). In the

presence of the knee-like bend a heel is visible. The sclerotised vagina is long, sinuous

(except for *C. dageti, C. falcifer, C. kmentovae*) or spiralled (*C. longicirrus, C. sanseoi*).

Species of the *Cop* and *Tylo* groups share some characteristics with the species of the *Hemi*

group but have either a substantially shorter penis or do not display the characteristic knee-

like bend in the accessory piece. Previous studies have grouped species of the *Hemi* species

group together based on phylogenetic (Mendlová et al., 2012; Řehulková et al., 2013) and

morphological (Dossou and Birgi, 1984; Pariselle and Euzet, 2004; Jorissen et al., 2018a)

analyses.

4. *Tilapiae: The species complex of Cichlidogyrus tilapiae*. For a detailed

characterisation, see species descriptions and characterisations by Paperna (1960) and

Douëllou (1993). *C. tilapiae* have both been hypothesised to form a species complex

(Pouyaud et al., 2006).

5. *Oreo1: Species infecting oreochromine and coptodonine cichlids*. Anchor pairs

similar in shape and size but dorsal anchors somewhat smaller. Anchors with short and

broadened outer root and long inner root. Dorsal bar arched with two broad and large

auricles. Ventral bar with thin mid-portion, and thickened towards the ends includes

triangular membranous extensions. Hooks small with small shaft. The MCO consists of long

tubular (*C. cirratus, C. giostrai, C. mbirizei, C. mvogoi, C. ornatus*) (Fig. 6) or short and

thickened (*C. acerbus, C. lagoonaris, C. nageus, C. njinei, C. slembroucki*) penis (Fig. 5b)

sometimes with bulbous extension at the base of the tube (*C. njinei, C. ornatus, C. giostrai*)

and complicated roughly C-shaped (Fig. 6) accessory piece often with finger or hook-shaped

outgrowths and marked heel. Proximal end of accessory piece attached to penis bulb and

distal end holds the distal portion or a mid-portion of the penis. Vagina simple (*C. acerbus,*

C. lagoonaris, C. slembroucki), sinuous (*C. giostrai, C. njinei, C. ornatus*), or coiled (*C.*

cirratus, C. mbirizei, C. mvogoi) tube and only slightly sclerotised.

6. *Halli*: *The species complex of Cichlidogyrus halli*. For detailed characterisation see

original species description by Price and Kirk (1967). *Cichlidogyrus halli* has both been

hypothesised to be a species complex (Douëllou, 1993; Jorissen et al., 2018b).

7. *Scutogyrus: Species infecting oreochromine and pelmatolapiine cichlids*. This

group has been diagnosed as the separate genus *Scutogyrus* including a detailed

morphological characterisation (Pariselle and Euzet, 1995). Concerning the haptor

morphology, species of *Scutogyrus* display well-developed hook pairs 3–7, elongated auricles

at the dorsal bar (Fig. 5a; Fig. 6) and a ventral bar that supports a large plate (Fig. 6). The

accessory piece presents a distal flap (Fig. 5b; Fig. 6).

8. *Cop: Species infecting coptodonine cichlids*. Anchor pairs dissimilar in shape.

Dorsal anchor with V-shaped indentation between the roots and an elongated inner root at

least four times longer (Fig. 5a) than the outer root. Ventral anchor with shallower

indentation, and relatively short roots including a considerably shorter inner root. Dorsal bar

thick with large and broad auricles. Ventral bar V-shaped and truncated at the ends and

thinner towards the mid-portion, displays membranous extensions. First hook pair (U1) well-

developed (Fig. 5a, Fig. 6). Hook pairs 3–7 (U3–7) small. The MCO consists of short tubular

penis with a marked heel and an accessory piece with a sickle-like terminal hook connected

to the basal bulb of the penis. No sclerotised vagina has been observed for species belonging

to this group except for *C. digitatus* in the original description (Dossou 1982). However, none

of the more recent reports of species of the *Cop* clade (Pariselle and Euzet, 1996, 2003)

remark on this report. Previous studies have grouped species of the *Hemi* clade, e.g. *C.*

dionchus (Pariselle and Euzet, 2003), and the *Tylo* clade, e.g. *C. kothiasi* (Pariselle et al.,

2013), with the species of the *Cop* clade because of the similar morphology of the hook pairs,

i.e. large U1 and small U3–7. However, none of these species displays an accessory piece

with a sickle-like terminal hook. Species of the *Cop* clade have previously been grouped

together (Pariselle and Euzet, 2003; Jorissen et al., 2018b) but without *C. berminensis*.

9. *Oreo2: Two species infecting oreochromine and coptodonine cichlids*. Both species

display simple features such as similar anchor pairs and small hook pairs. Most other features

differ considerably between the species. Dorsal and ventral anchors are similar in size and

shape, the roots are variable in length, i.e. a long inner root in *C. amphoratus* (see drawing

Pariselle and Euzet, 1996) or barely to no distinct roots in *C. sclerosus* (Douëllou, 1993).

Dorsal bar X-shaped with two auricles of variable shape and length. Ventral bar variable in

shape and size with membranous extensions (Mendlová et al., 2012). Hooks small with

developed but small shaft except for the second pair (Douëllou, 1993; Pariselle and Euzet,

1996). The MCO consists of arched, tubular penis (Fig. 5b) with large heel ('subspherical' in

C. amphoratus or shaped as a 'serrated plate' in *C. sclerosus*), and associated relatively

simple accessory piece (Douëllou, 1993; Pariselle and Euzet, 1996) (Fig 5b). Sclerotised

vagina present. *Cichlidogyrus amphoratus* presents a swollen middle portion of the penis

reminiscent of species of the *Bulb* clade (Pariselle and Euzet, 1996). However, in the latter

group, the dorsal anchors have long inner roots and the first hook pair is well developed (Fig.

6). The close relationship of these species has previously been reported but was not further

discussed (see Messu Mandeng et al., 2015).

10. *Bulb: Species from Southern Africa with a bulbous penis*. Dorsal and ventral anchors

dissimilar. Dorsal anchors with deep indentation between short outer root and a long inner

root (Fig. 6). Dorsal bar short and stout with auricles medium-sized to large (average > 20

µm for *C. zambezensis*). First hook pair (U1) well-developed (Fig. 5a; Fig. 6) albeit

somewhat less in *C. zambezensis*. Hook pairs 3–7 small to slightly developed. Male

copulatory organ consists of a broad penis with a swollen, bulbous middle portion, a small

heel (Fig. 6), and a sharp tube-like (*C. philander*) or filiform (*C. maeander, C.*

papernastrema, *C. pseudozambezensis, C. zambezensis*) termination, and a simple accessory

piece of similar length, which is articulated to the base of the penis bulb (Price et al., 1969;

Douëllou, 1993; Jorissen et al., 2018b). *Cichlidogyrus amphoratus*, *C. bulbophallus*, *C.*

flagellum, *C. giostrai*, *C. karibae*, *C. lagoonaris*, *C. njinei*, *C. ornatus*, and *C. sanjeani* also

present a swollen portion in the penis but differ from the species of the group as follows.

• *C. amphoratus* has similar dorsal and ventral anchors with no well-developed inner root

of the dorsal anchor (Fig. 6).

• The first hooks (U1) of *C. amphoratus* (Fig. 6), *C. flagellum, C. giostrai, C. karibae, C.*

lagoonaris, C. njinei, *C. ornatus*, *C. ranula*, and *C. sanjeani* are considerably shorter and

less developed.

• The hooks U3–U7 of *C. bulbophallus* and *C. ranula* are considerably longer and resemble

the marginal hooks of the species of the *CPO* species group.

• The tubular distal portion of the penis of *C. amphoratus* (Fig. 6)*, C. bulbophallus, C.*

flagellum, C. giostrai, C. njinei, and *C. ornatus* exceed the length of the swollen portion

of the penis considerably.

• *C. amphoratus, C. giostrai, C. lagoonaris, C. njinei, C. ornatus,* and *C. sanjeani* have

only been reported from Western Africa not Southern Africa (see Pariselle and Euzet

2009).

• *C. amphoratus* and *C. njinei* belong to the *Oreo2* and *Oreo1* species groups respectively.

11. *Tylo*: *Species infecting tylochromine cichlids.* Dorsal and ventral anchors dissimilar.

Roots of anchors frequently fused forming fenestrae (windows), e.g. for *Cichlidogyrus*

berrebii, C. chrysopiformis, C. dijetoi, C. kothiasi, C. pouyaudi (Fig. 6), and *C. sergemorandi*

(Pariselle and Euzet, 1994; Pariselle et al., 2014; Rahmouni et al., 2018). Dorsal anchors with

short outer root and a long inner root, median portion of blade hollow in *C. berrebii, C.*

pouyaudi (Pariselle and Euzet, 1994), and *C. sigmocirrus* (Pariselle et al., 2014). Dorsal bar

with two generally short auricles fused to the bar (Fig. 6; Fig. 7a) is one of the most

distinctive features of species of this group (Pariselle et al., 2014; Rahmouni et al., 2018).

The hooks are generally small but the first pair is developed in *C. bixlerzavalai, C.*

chrysopiformis, C. dijetoi, C. kothiasi, and *C. muzumanii* (Fig. 5a). The MCO consists of a

coiled tubular penis, a heel, and a flat ribbon-like (*C. berrebii, C. kothiasi, C. pouyaudi, C.*

sigmocirrus) (Pariselle and Euzet, 1994; Pariselle et al., 2014), drape-like (*C. chrysopiformis,*

C. dijetoi, C. mulimbwai, C. omari, C. sergemorandi) (Muterezi Bukinga et al., 2012;

Pariselle et al., 2014; Jorissen et al., 2018a; Rahmouni et al., 2018), or reduced (*C.*

bixlerzavalai, C. muzumanii) (Muterezi Bukinga et al., 2012; Jorissen et al., 2018a) accessory

piece. Unlike other species of *Cichlidogyrus*, the penis and accessory piece of the MCO are

separated at the base but might be connected through a filament in *C. chrysopiformis* and *C.*

sigmocirrus (Pariselle et al., 2014). Species of the *Tylo* group have regularly been grouped

together since the descriptions by Pariselle and Euzet (1994) with most recent additions

provided by Jorissen et al. (2018a) and Rahmouni et al. (2018b).

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Phylogenetic position of 'unsequenced' species and systematic informativeness: literature

survey, combined phylogeny, machine learning algorithm, and cluster analysis

The addition of the morphometric data to the parsimony analysis resulted in an overall

less resolved and supported tree (Fig. 4). Overall, 30% (24/79) of the 'unsequenced' species

formed well-supported clades with sequenced species allowing us to affiliate these with the

respective clades from the molecular phylogeny. Phylogenetic positions inferred from the

combined MP consensus tree were in moderate accordance ($\kappa = 0.48$) with the classifications

in species groups inferred from the character maps. Meanwhile, the optimised SVMs ($C = 2^{-7}$,

$\sigma = 0.5$) predicted species group affiliation of the sequenced species with moderate success (κ

$= 0.68$) with 1035 observations included. For the unsequenced species (833 observations),

accordance with the parsimony analysis ($\kappa = 0.36$) and literature survey ($\kappa = 0.31$) was

comparatively low.

Variable importances of the optimised SVMs indicated an acceptable to excellent

discriminatory power on average (Fig. 7a) with exception of the surface of the auxiliary plate

(AuP) in the MCO that performed poorly (mean AUC < 0.6). Collapsing the AUC values by

species groups (Fig. 7b), indicates an uneven distribution with some groups such as *CPO*

being easier to discriminate against their congeners than others such as *Tylo*. Lastly, based on

the Pearson pairwise correlation matrix (Fig. 7c), we detected 15 clusters of multicollinear

measurements indicating that almost half of the measurements (14) are multicollinear.

Character evolution of the attachment and reproductive organs: morphometrics and

discretisation

For the attachment organ measurements, we detected a strong phylogenetic signal (λ

$= 0.91$; CI: 0.77–1.00). Brownian Motion (BM), Ornstein-Uhlenbeck (OU), and Early-Burst

(EB) models all performed equally (Fig. 4b). The Late Burst (LB) model performed worse.

Only the multi-rate BM model (BMM) performed better albeit with a slight overlap in EIC

estimates. For the reproductive organ measurements, the phylogenetic signal was

considerably weaker ($\lambda = 0.73$; CI: 0.43–1.00) and no model outperformed the BM model.

Out of the discrete characters, only the hook configuration (HC) and ventral bar shape

(VBS) produced strong phylogenetic signals (Fig. 4c). In contrast, we detected a relatively

weak phylogenetic signal for the anchor similarity (AS) and no signal for the index of

specificity (IS). Character maps (Fig. 9a) illustrate these phylogenetic pattern. Based on the

literature survey, we suggested a series of measurements that are equivalents of HC, VBS,

and AS. The HC summarises the lengths of the marginal hooks (U1–U7), VBS is described

the width of the ventral bar in relation to its size ('massive bar' in Mendlová et al., 2012) but

also includes information on the presence of membranous extensions, and AS likely refers to

a difference in lengths of the inner roots of the dorsal vs. the ventral anchor pair and possibly

a difference in size between the anchors pairs based on a drawing in Mendlová et al., (2012)

but the publication only mentions a 'difference in shape'. Character maps of these continuous

measurements are plotted next to the equivalent discrete characters (Fig. 9b).

DISCUSSION

We investigated the evolution of the attachment and reproductive organs of parasites

infecting a host lineage that has undergone multiple adaptive radiation events (Seehausen,

2006). Species of *Cichlidogyrus* (incl. the nested *Scutogyrus*) form a morphologically diverse

lineage of parasitic flatworms infecting the gills of African cichlid fishes. This diversity

alongside the model system status of the hosts led to the suggestion to use these parasites as a

model system for parasite macroevolution (Pariselle et al., 2003; Vanhove et al., 2016). In

this context, the present study is the most extensive reconstruction of intra-generic

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

evolutionary relationships of African metazoan parasites to date providing new

morphological, molecular, and host range data, which are made available in public databases.

We were able to identify and characterise 11 different species groups (Fig. 2a; Fig. 6)

amongst the 137 species that are currently known (Cruz-Laufer et al., 2021). For some of

these groups, member species have previously been hypothesised to be closely related, e.g.

species within the *Tylo, Hemi*, *CPO*, and '*Scutogyrus*' groups. However, this study is the first

to summarise all these groups and to analyse the evolution of the attachment and reproductive

organ morphology using sequence data of 58 species and morphometric data of 136 species.

We here hypothesise the phylogenetic position of described species of *Cichlidogyrus* and

Scutogyrus without available molecular data through a synthesis of morphological,

molecular, and host range data. However, due to missing data, e.g. a limited coverage of the

four gene regions and a limited taxon coverage with the majority of species yet undiscovered

(Cruz-Laufer et al., 2021), the relationship between the proposed species groups remains

mostly unresolved with only moderate support for a few larger clades (Fig. 2a, b). In this

light, we opted against taking any nomenclatural act such as splitting the genus into several

new genera or synonymising *Scutogyrus* with *Cichlidogyrus* [for a more detailed discussion

of this matter and similar cases within Dactylogyridae, see Kmentová et al. (2021a)].

Systematic informativeness of morphometric measurements and host range data

To identify and classify species of *Cichlidogyrus,* previous publications have applied

a specific set of measurements of the sclerotised attachment and reproductive organs (Cruz-

Laufer et al., 2021). We investigated the systematic informativeness of the 29 measurements

most frequently provided in the literature (Fig. 1c). Generally, the measurements appear to

have a low systematic informativeness as a substantial portion of unsequenced species (70%)

could not be placed in any of the clades inferred from the molecular phylogeny through a

parsimony analysis combining molecular and morphometric data. Phylogenetic positions

inferred from the combined MP consensus tree were in moderate accordance with the

classifications based on character maps that considered additional discrete characters for the

reproductive organs proposed here (Fig. 5b). The limited systematic informativeness of the

measurements is also reflected by the performance of the classification analysis using support

vector machines (SVMs). The optimised machine learning algorithm predicted species group

affiliation within the sequenced species with moderate success ($\kappa = 0.48$). Yet for the

unsequenced species, accordance of the SVM-based classification with the parsimony

analysis ($\kappa = 0.36$) and literature survey ($\kappa = 0.31$) was comparatively low. One cause for this

performance might be the low number of systematically informative measurements in the

dataset as ten of these showed a moderate or low discriminatory power on average (AUC $<$

0.8, see Fig. 7a). Furthermore, the discriminative power of the measurements varied

considerably between the different clades (Fig. 7b).

Notably, the phylogenetic signal was low for any of the evolutionary models applied

to the reproductive organ in the context of the phylogenetic comparative method. This lack of

information presents a stark contrast to the importance of the reproductive organ morphology

to the character map–based classification. The phenotype of the male copulatory organ was

crucial for characterising groups of related species in the form of several additional discrete

characters proposed here (Appendix 2). These characters allowed us to consider information

on the shape and diameter of the reproductive organs not included in the parsimony analysis

and SVM-based approach. Therefore, we could provide hypotheses for all but 12 species

regarding their phylogenetic position. This relative success highlights the value of

phylogenetic information conveyed by reproductive organ structures; e.g. Rahmouni et al.

(2017) noted that the heel of the male copulatory organ was shown to be similarly shaped in

possibly related Eastern African species. Similar conclusions have been drawn for other

dactylogyrid monogeneans such as species of *Characidotrema* Paperna & Thurston, 1968

(Řehulková et al., 2019). Regarding the lack of a phylogenetic signal of reproductive organ

measurements, Pouyaud et al. (2006) proposed that this issue might be explained by a fast

evolutionary rate but we suggest that this observation might be rather linked to a limited

taxon coverage as more species have been described since. Instead, the low systematic

informativeness indicated here more likely reflects a lack of systematically informative

metrics as, for example, the number of reproductive organ measurements is low compared to

the attachment organ (5 vs. 24). The measurements that are currently employed, fail to

incorporate the systematic information offered by the attachment and reproductive organ

morphology.

The low predictive power of the morphometrics has encouraged the use of loosely

defined discrete characters in the past (Fig. 9). Several studies have categorised species

according to their attachment organ morphology (Vignon et al., 2011; Mendlová et al., 2012)

and host repertoires (Mendlová and Šimková, 2014). Here, we compared the systematic

informativeness of these discrete characters against their continuous counterparts (Fig. 9).

Two of the discrete characters lacked a phylogenetic signal (Fig. 8b) despite the moderate

discriminatory power of the continuous characters suggested by the machine learning

algorithm (Fig. 7a), a problem possibly caused by merging the information of multiple

continuous variables into a single discrete character (Fig. 9). Discretisation can also introduce

a researcher bias and produce misleading results. For instance, Mendlová et al. (2012)

proposed that similar anchor shapes were an ancestral state of the binary character 'shape of

anchors' ('anchor similarity' in the present study) based on a character map but the

phylogenetic informativeness of the 'shape of anchors' character was never investigated

further. In fact, the cluster analysis of the present study shows that dorsal anchor

measurements and the respective ventral anchor measurements are highly multicollinear (Fig.

7c) with the exception of the inner roots (DAd and VAd). The ventral bar shape was also

introduced as a variable for character mapping but no ancestral state could be inferred

(Mendlová et al., 2012). Yet species of *Scutogyrus* have a unique phenotype with an extended

plate associated to the bar structure (Pariselle and Euzet, 1995). These inconsistent results

between continuous characters and their discretised counterparts underline that interpretations

based on these discretised characters should be treated with caution. For instance, the hook

configuration character states introduced by Vignon et al. (2011) are founded on gaps

detected between the hook measurements of different species (see Pariselle and Euzet, 2009).

New species discoveries in the last decade demonstrate that these character states might not

be as clear-cut (see Rahmouni et al., 2017; Geraerts et al., 2020). Nonetheless, hook

measurements were highly relevant in a machine learning context (Fig. 7a) indicating that,

while an update to the coding of this character might be required, some character states might

still prove useful for characterising species groups of *Cichlidogyrus*, e.g. the monophyletic

CPO group species with well-developed hook pairs 3–7. If researchers wish to use discretised

morphological characters in the future, character states should be redefined through

quantitative character coding techniques such as proposed by Garcia-Cruz and Sosa (2006).

The resulting character states would be more independent from researcher bias, which could

increase the repeatability of such studies.

A discrete variable proposed to represent host specificity has also led to doubtful

conclusions. Mendlová and Šimková, (2014) used the index of specificity (IS) to summarise

the phylogenetic specificity of species of *Cichlidogyrus* (and *Scutogyrus*) into four categories

including species-, genus-, and tribe-specific species, and generalists (Fig. 3a). Their study

concluded that host specificity is related to parental care of the cichlid hosts. However, this

observation was made due to an inherent sampling bias towards hosts raising their offspring

through mouthbrooding (Cruz-Laufer et al., 2021), a specific type of parental care. Mendlová

and Šimková (2014) also observed a correlation of the IS with the host phylogeny. However,

this observation might again reflect a study bias in cichlid parasitology, in this case towards

large economically relevant host species (Cruz-Laufer et al., 2021). Because of their

importance to fisheries, these fishes unlike others have been introduced to various new

habitats. Anthropogenic introductions might also increase the chances of co-introduction for

their parasites and create ecological opportunities to expand host repertoires (e.g. Jorissen et

al., 2020). To avoid these biases, future studies should treat host specificity as a complex

parameter encompassing structural, phylogenetic, and geographical aspects (Poulin et al.,

2011) as well as temporal fluctuations (Brooks et al., 2019). We suggest that these parameters

could be included, e.g., in a genus-wide comparative study in the future.

In general, the size of morphological structures should first be treated as continuous,

unmodified characters before further processing to avoid losing systematic information.

Discretisation should only be applied with sufficient statistical evidence to back up character

coding. We suggest increasing sampling and sequencing efforts to address data deficiency.

Attachment organ evolution: rate heterogeneity, co-divergence, and host switching

To infer evolutionary patterns from the attachment organ morphology, we applied a

penalised likelihood framework for multivariate phylogenetic comparative methods (PCMs)

to the continuous character data. The evolutionary models representing different evolutionary

hypotheses performed almost equally well (Fig. 4b). However, we found no significantly

increased performance of the Ornstein-Uhlenbeck (OU), Early Burst (EB), and Late Burst

(LB) models compared to the Brownian Motion (BM) model. Only the multi-rate BM

(BMM) model performed slightly but significantly better than the other models (Fig. 4b). The

OU, EB, and LB models present extensions of the Brownian Motion (BM) model (Butler and

King 2004; Harmon et al. 2010). Therefore, the character evolution of the attachment organs

measurements might simply follow a pattern associated with genetic drift or randomly

fluctuating natural selection (Losos, 2008). This pattern might also be reflected in

morphological features that are frequently shared by related species (Fig. 6). The relatedness

of the parasite species could determine the evolution of attachment organ as suggested by a

previous study (Vignon et al. 2011). Studies on a related lineage of monogeneans

(*Ligophorus* Euzet & Suriano, 1977) also indicate a correlation of morphometrics and

phylogenetic relationships (Rodríguez-González et al. 2017). However, the increased support

for the BMM model suggests that the evolution of these parasites might have been shaped by

different evolutionary regimes. This rate heterogeneity appears logical as host lineages

outside the Great African lakes have not experienced the same explosive speciation as their

East African relatives (see Brawand et al., 2015). In fact, purely single-rate models appear

increasingly unlikely to capture the real evolutionary history of *Cichlidogyrus* because of its

pan-African distribution (Cruz-Laufer et al., 2021) and ecological diversity of hosts (Burress,

2014). In the present study, we only modelled a multi-rate BM model because a lack of

software packages suited for high-dimensional dataset with pre-set hypothesis. Yet other

multi-rate models including OU, EB, and LB processes could provide better fits. In

particular, species of the *EAR* group (Fig. 2a) that infect cichlid radiations in East Africa (Fig.

3) might mirror the explosive speciation of their hosts (see Vanhove et al., 2015). In recent

years, multiple tools have been developed to investigate these more complex models but

often these methods represent maximum likelihood (ML) approaches (e.g. Beaulieu et al.,

2012; Ingram and Mahler, 2013; Puttick, 2018) that are sensitive to trait covariation, data

orientation, and trait dimensions, particularly for high-dimensional datasets (Adams and

Collyer, 2018). Other methods were developed to detect evolutionary rate shifts (e.g. Ingram

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

and Mahler, 2013; Uyeda and Harmon, 2014; Khabbazian et al., 2016) rather than to test

hypotheses based on previous knowledge (such as for the *EAR* clade in the present case).

Therefore, penalised log-likelihood approaches that address these issues remain currently

limited to multi-rate BM models but future updates will most likely close this technological

gap.

Highly similar morphological characteristics can be found in multiple species groups

(Fig. 5a; Fig. 6) such as a well-developed outer root of the dorsal anchor (e.g. species of the

Bulb and *Cop* groups), large first hooks (e.g. species of *Tylo*, *EAR – Heel*, and *Hemi* groups),

and long third to seventh hooks (e.g. species of the *EAR – Heel*, *CPO* groups). Character

shifts observed within species groups appear to be adaptations to divergent host repertoires or

environments (see Messu Mandeng et al., 2015). Host- and environmentally induced shifts in

the attachment organ morphology have also been observed amongst other dactylogyrids, e.g.

species of *Kapentagyrus* (Kmentová et al., 2018) and *Thaparocleidus* (Šimková et al., 2013),

and other parasite lineages, e.g. cymothoid isopods (Baillie et al., 2019). For *Cichlidogyrus*,

host species infected by parasites from the same species group frequently belong to the same

tribe or related tribes of cichlids (Fig. 3; Appendix 5). Consequently, the phylogenetic

congruence amongst some clades of *Cichlidogyrus* and their hosts is strong (Vanhove et al.,

2015). Examples for host-induced character shifts amongst species of *Cichlidogyrus* can be

found across the genus:

• *Cichlidogyrus amieti* has undergone a recent character shift as it displays small first

hooks unlike other species of the *Hemi* clade, where they are well-developed. This change

might result from a host switch (Messu Mandeng et al., 2015).

• *Cichlidogyrus sclerosus,* a generalist infecting fishes belonging to various cichlid tribes

(Fig. 3), shares almost no attachment organ features with its sister species *C. amphoratus*,

a specialist infecting only coptodonine cichlids (Fig. 3), beyond those shared by all

species of the genus, e.g. auricles associated with the dorsal bar of the attachment organ

(see Pariselle and Euzet, 2009).

• The diverse morphological features of species of the *EAR* group might relate to the

ecological and morphological diversity of the hosts, e.g. related to feeding behaviour

(Burress, 2014). The parasites, which we grouped into subgroups of the *EAR* clade, might

have adapted to these ecologically diverse hosts resulting in divergent attachment and

reproductive organ morphologies.

• Co-infections of different species of *Cichlidogyrus* can result in niche segregation on the

gills of a host such as in species infecting Lake Victoria cichlids (Gobbin et al., 2020). At

least some co-infecting species can be told apart based on their attachment organ

morphology (see Gobbin et al., 2021). Thus, phenotypic differences might be linked to

specific infection sites. Morphological adaptations to gill microhabitats have been

reported for other monogeneans (Rohde, 1976; Ramasamy et al., 1985) but no study has

so far investigated these mechanisms for species of *Cichlidogyrus*.

The host repertoires of the parasites offer clues for evolutionary processes frequently

associated with host-induced effects in the evolutionary history of *Cichlidogyrus* including

co-divergence and host switching. For instance, co-divergence might be reflected in the host

(Fig. 3; Appendix 5) and geographical repertoires of species of *Cichlidogyrus* (Vanhove et

al., 2013) (e.g. *EAR* group species are exclusive to Eastern Africa and *Bulb* group species to

Southern Africa). Furthermore, a small yet increasing number of natural and invasion-

induced host switches are reported in the literature from cichlids to non-cichlids, e.g. in the

Hemi clade (Birgi and Euzet, 1983; Birgi and Lambert, 1986), and from African cichlids to

cichlids and non-cichlids in the Americas, the Levant, and Madagascar (*C. arthracanthus, C.*

sclerosus, C. tiberianus, C. tilapiae, C. thurstonae, and *Scutogyrus longicornis*) (Paperna,

1960; Jiménez-Garcia et al., 2001; Šimková et al., 2019). While the literature suggests that

most African fish families have their own monogenean lineages (Carvalho Schaeffner, 2018)

and the number of similar host switches might, hence, be limited, more host switches might

be discovered in the future if more (also non-cichlid) host species are studied.

Despite these host-induced character shifts, the evolutionary models employed for

multivariate PCMs detected no changes in the evolutionary rate of the attachment organ

measurements. Apart from the limitation regarding multi-rate models, mentioned above, we

suggest that this observation might be a result of incomplete sampling, which has affected

similar studies in the past. For instance, previous studies were limited to single structures, i.e.

the anchors (Rodríguez-González et al., 2017), or to the much lower number of species of

Cichlidogyrus described at the time (77 vs. 130 today), which comprised less phenotypical

diversity than is currently known to exist (Vignon et al., 2011). In fact, the phylogenetic

position of some species with features diverging from their respective species groups (e.g. *C.*

amieti from other *Hemi* group species) were not known. Observed similarities of the

attachment organ morphology in related species reflected the specificity of the parasites to

the respective host lineage rather than only the relatedness of the monogeneans. While the

present analysis includes more species, DNA sequence data of less than half of all described

species are available at GenBank and many species are likely to be discovered in the future

(Cruz-Laufer et al., 2021). This data deficiency could affect the performance of the models

implemented through PCMs as, e.g., the lack of lineages might lower the resolution of

phylogenetic inference and increase phylogenetic uncertainty. A more complete taxon

coverage of morphological, molecular, and ecological data will provide more insight into the

evolutionary patterns of the genus. Furthermore, a closer look into the evolution of single

species groups might reveal different patterns than for a genus-wide study. For instance, the

high variability in the attachment and reproductive organ morphology of the *EAR* group

might indicate a late burst in the divergence of characters induced by phenotypic adaptation

to the rapidly speciating host lineages. A slightly better performance of the multi-rate model

provides reason to further investigate the evolutionary history of this group, as it might fall

under a divergent evolutionary regime. Lastly, convergent evolution of morphological

characters might also have affected PCM model performance. Convergence can mask

evolutionary processes that have occurred in the past as phenotypes become more similar

(Losos, 2008). Monogenean parasites infecting the haplochromine radiations in the Eastern

African lakes might be particularly impacted by this effect because of the convergent

evolution of their hosts. These fishes have evolved via replicated adaptive radiation events

resulting in similar yet unrelated species occupying the same niches in different lakes

(McGee et al., 2016). This setting makes their parasites also potential targets to study

convergent evolution in parasites. Conversely, the Eastern African diversity of *Cichlidogyrus*

has mostly been investigated in Lake Tanganyika, the oldest lake, limiting the current

possibilities to study parasite evolution in the context of replicate host radiations in this

region [but see Gobbin et al. (2021) for a study focusing on Lake Victoria].

Reproductive organ evolution

Unlike the attachment organ, the reproductive organs show no character changes

correlated to host relatedness. Reproductive organ characters are evolutionarily stable within

the different clades as evidenced by the role of the male copulatory organ in the

morphological characterisation of the species groups. The evolution of the reproductive

organs, unlike the attachment organ, has no apparent connection to the host range and seems

to be determined mostly by phylogenetic constraints. Differences within species groups seem

to reflect phylogenetic positions, e.g. the short penis of *C. falcifer* (*Hemi*) and simple

accessory piece of *C. falcifer* and *C. cubitus* (*CPO*) are ancestral states (Fig. 5) of the

respective clade. We detected a marginally better support for the BMM model (Fig. 8a)

suggesting that the *EAR* group, with its host lineages that have resulted from adaptive

radiation events, might have experienced an evolutionary regime slightly divergent from its

congeners. However, the low phylogenetic signal in these data provides far less confidence in

this result. Previous studies on the reproductive organs of monogenean flatworms have

suggested that the complex surface of the reproductive organ might correlate with

environmental factors, e.g. to avoid separation during copulation through the water flow

(Kearn and Whittington, 2015). This hypothesis would suggest that the sclerotised structures

evolve under selective pressure. However, while the screw thread–like thickening of the penis

wall for some species (*C. casuarinus, C. centesimus*, *C. nshomboi*) (Fig. 6, *Heel* subgroup)

might present such an example, we see no general trend to confirm this hypothesis. Neither

generalist species such as *C. halli, C. tilapiae*, and *C. sclerosus* nor species infecting

potentially more motile hosts living in fast-flowing waters, e.g. *C. consobrini, C. maeander,*

and *C. ranula* infecting species of *Orthochromis*, display more complex MCOs than their

congeners. Consequently, the diversification of the reproductive organs might be explained

by reproductive isolation such as previously suggested for other dactylogyrid monogeneans

(Šimková et al., 2002). A quantitative approach to measure reproductive organ shape could

provide a more detailed answer to this question by capturing the shape variation currently

only recorded as qualitative information (Fig. 5b) in taxonomic literature (see below for

proposals).

Classification, prediction, and research prospects

We were able to demonstrate the existing yet limited predictive power of the

measurements currently employed by most taxonomists through a phylogenetic analysis

under maximum parsimony accompanied by a machine learning approach. To improve the

accuracy of these predictions overall, datasets with a more complete taxon coverage and

higher number of observations are needed. The results of the character map–based

classification vs. the parsimony analysis and the statistical classification approaches have

highlighted a discrepancy between the systematic informativeness of the attachment and

reproductive organ morphology in taxonomic literature and the low number of

phylogenetically informative measurements. Future studies should address this discrepancy

by increasing the number of systematically informative metrics.

Several possible solutions have been proposed to increase the number of metrics.

First, some studies included novel measures such as the lengths of the shaft of the hooks, the

curved length of the accessory piece, or the width of the penis tube and the bulbous portion of

the penis (e.g. Geraerts et al., 2020) to describe species of *Cichlidogyrus* in more detail.

Second, geomorphometric analyses have been employed for the anchors of some species of

Cichlidogyrus (Rahmouni et al., 2021) as well as other monogeneans (Vignon and Sasal,

2010; Rodríguez-González et al., 2017; Kmentová et al., 2020) to infer the most

systematically informative structures. Third, 3D imaging techniques as previously developed

for monogenean sclerotised structures (e.g. García-Vásquez et al., 2012) have been proposed

to improve characterisations of species of *Cichlidogyrus* (Cruz-Laufer et al., 2021) using

methods employed for shape analyses (Klingenberg, 2010; Dryden and Mardia, 2016). Key

characters that could be inferred using this approach are for example elongation (long vs

round), torsion (straight vs coiled), and complexity (few vs many extensions). However, all

of these solutions remain currently unavailable for comparative analyses. Data resulting from

the proposed additional measures, geomorphometric measurements, and 3D imaging

techniques are available for only few morphological structures, species, and studies. Future

research should increase the taxon coverage for proposed methods and improve

quantification methods to address these limitations. Future datasets should encompass an

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

increasing number of DNA sequences and morphometric measurements from all known

species as this study highlights the limitations of using morphological characters to infer the

evolution of monogenean parasites. Nonetheless, raw data should regularly be published in

supplements or data repositories (e.g. Kmentová et al., 2016b) to improve the accessibility of

morphometric data. This approach could enhance information flow and academic capacity for

research into species of monogenean parasites in the future (Cruz-Laufer et al., 2021).

CONCLUSION

Due to the species-richness and the model system status of their cichlid hosts, the

parasite lineage including species of *Cichlidogyrus* and *Scutogyrus* could be an ideal

macroevolutionary model system for speciation in parasitic organisms in the future. Notably,

these species belong to one of the more extensively investigated lineages in close interaction

with an adaptive radiation of other species, namely cichlid fishes. As adaptive radiations

might explain a substantial part of the biodiversity on the planet, many species potentially

interact with radiating lineages. Therefore, understanding evolutionary history in this context

could be key for understanding the diversity of species interactions in general.

Similar to many parasite systems, the limited availability of phenotypic and genotypic

data for the 137 species of *Cichlidogyrus* currently described has hampered investigations

into the evolutionary history of the lineage. This study provides an example of how an

increased availability of morphological and DNA sequence data can promote computational

approaches such as parsimony analyses, phylogenetic comparative methods, and machine

learning classification to answer basic questions of evolutionary research. The

phylogenetically informative measurements (continuous and discrete, established and newly

proposed) and characterisation of multiple clades within *Cichlidogyrus* have highlighted key

information required to describe new species as well as the shortcomings of the

morphological characters that are currently used. Features of the attachment organ used to

characterise the proposed species groups remain key descriptors alongside the features of the

reproductive organs. However, they only serve as predictors in a multivariate context. This

conclusion presents a stark contrast to previous classification systems for this taxon, which

used to predict species groups based on single discrete morphological characters.

Using multivariate phylogenetic comparative methods, we were able to model

evolutionary mechanisms of the attachment and reproductive organs suggested by previous

studies. No support for alternative models suggests that the variation of the morphology of

these organs across the genus has been shaped by different evolutionary regimes.

Morphological patterns in different species groups further indicate that the attachment organ

morphology can be shaped by host parameters. In contrast, the reproductive organs appear to

follow a more random evolutionary regime. Other mechanisms might play a role in the

evolution of this parasite lineage, e.g. within-host speciation as suggested by other studies.

Therefore, more morphometric and DNA sequence data will be needed to provide a more

detailed picture of the evolution of the genus. We encourage researchers to publish

morphometric raw data regularly to improve the data accessibility for the scientific

community, and to sequence systematically informative DNA regions of the remaining

described or undiscovered species.

1207

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

REFERENCES

Adams, D.C., Collyer, M.L., 2018. Multivariate phylogenetic comparative methods:

evaluations, comparisons, and recommendations. Syst. Biol. 67, 14–31.

Altman, D.G., Royston, P., 2006. The cost of dichotomising continuous variables. Br. Med. J.

332, 1080.

Baillie, C., Welicky, R.L., Hadfield, K.A., Smit, N.J., Mariani, S., Beck, R.M.D., 2019.

Hooked on you: shape of attachment structures in cymothoid isopods reflects parasitic

strategy. BMC Evol. Biol. 19, 207.

Barson, M., Přikrylová, I., Vanhove, M.P.M., Huyse, T., 2010. Parasite hybridization in

African *Macrogyrodactylus* spp. (Monogenea, Platyhelminthes) signals historical host

distribution. Parasitology 137, 1585–1595.

Beaulieu, J.M., Jhwueng, D.C., Boettiger, C., O'Meara, B.C., 2012. Modeling stabilizing

selection: expanding the Ornstein-Uhlenbeck model of adaptive evolution. Evolution 66,

2369–2383.

Birgi, E., Euzet, L., 1983. Monogènes parasites des poissons des eaux douces du Cameroun.

Présence des genres Cichlidogyrus et Dactylogyrus chez Aphyosemion

(Cyprinodontidae). Bull. la Soc. Zool. Fr. 108, 101–106.

Birgi, E., Lambert, A., 1986. Présence chez un Nandidae (Téléostéen), Polycentropsis

abbreviata Boulenger, 1901, du genre Cichlidogyrus (Monogenea, Monopisthocotylea,

Ancyrocephalidae). Ann. Parasitol. Hum. Comp. 61, 521–528.

Bitja-Nyom, A.R., Agnèse, J.F., Pariselle, A., Bilong-Bilong, C.F., Gilles, A., Snoeks, J.,

2021. A systematic revision of the five-spotted *Hemichromis* complex (Cichliformes:

Cichlidae) from West Africa and Lower Guinea, with the description of a new species

from Cameroon. Hydrobiologia 5, 3779–3803.

Blažek, R., Polačik, M., Smith, C., Honza, M., Meyer, A., Reichard, M., 2018. Success of

cuckoo catfish brood parasitism reflects coevolutionary history and individual

experience of their cichlid hosts. Sci. Adv. 4, eaar4380.

Blomberg, S.P., Garland, T., Ives, A.R., 2003. Testing for phylogenetic signal in comparative

data: behavioral traits are more labile. Evolution 57, 717–745.

Brawand, D., Wagner, C.E., Li, Y.I., Malinsky, M., Keller, I., Fan, S., Simakov, O., Ng,

A.Y., Lim, Z.W., Bezault, E., Turner-Maier, J., Johnson, J., Alcazar, R., Noh, H.J.,

Russell, P., Aken, B., Alföldi, J., Amemiya, C., Azzouzi, N., Baroiller, J.F., Barloy-

Hubler, F., Berlin, A., Bloomquist, R., Carleton, K.L., Conte, M.A., D'Cotta, H., Eshel,

O., Gaffney, L., Galibert, F., Gante, H.F., Gnerre, S., Greuter, L., Guyon, R., Haddad,

N.S., Haerty, W., Harris, R.M., Hofmann, H.A., Hourlier, T., Hulata, G., Jaffe, D.B.,

Lara, M., Lee, A.P., MacCallum, I., Mwaiko, S., Nikaido, M., Nishihara, H., Ozouf-

Costaz, C., Penman, D.J., Przybylski, D., Rakotomanga, M., Renn, S.C.P., Ribeiro, F.J.,

Ron, M., Salzburger, W., Sanchez-Pulido, L., Santos, M.E., Searle, S., Sharpe, T.,

Swofford, R., Tan, F.J., Williams, L., Young, S., Yin, S., Okada, N., Kocher, T.D.,

Miska, E.A., Lander, E.S., Venkatesh, B., Fernald, R.D., Meyer, A., Ponting, C.P.,

Streelman, J.T., Lindblad-Toh, K., Seehausen, O., Di Palma, F., 2015. The genomic

substrate for adaptive radiation in African cichlid fish. Nature 513, 375–381.

Brooks, D.R., Hoberg, E.P., Boeger, W.A., 2019. The Stockholm Paradigm: climate change

and emerging disease. University of Chicago Press, Chicago, USA.

Burress, E.D., 2014. Cichlid fishes as models of ecological diversification: patterns,

mechanisms, and consequences. Hydrobiologia 748, 7–27.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Butler, M.A., King, A.A., 2004. Phylogenetic comparative analysis: A modeling approach for

adaptive evolution. Am. Nat. 164, 683–695.

Campbell, V., Legendre, P., Lapointe, F.J., 2011. The performance of the Congruence

Among Distance Matrices (CADM) test in phylogenetic analysis. BMC Evol. Biol. 11.

Caña-Bozada, V., Llera-Herrera, R., Fajer-Ávila, E.J., Morales-Serna, F.N., 2021.

Mitochondrial genome of *Scutogyrus longicornis* (Monogenea: Dactylogyridea), a

parasite of Nile tilapia *Oreochromis niloticus*. Parasitol. Int. 81, 102281.

Carvalho Schaeffner, V.C., 2018. Part 5: Host-parasite list. In: Scholz, T., Vanhove, M.P.M.,

Smit, N., Jayasundera, Z., Gelnar, M. (Eds.), A Guide to the parasites of African

freshwater fishes. CEBioS, Royal Belgian Institute of Natural Sciences, Brussels,

Belgium, pp. 361–402.

Chernomor, O., von Haeseler, A., Minh, B.Q., 2016. Terrace aware data structure for

phylogenomic inference from supermatrices. Syst. Biol. 65, 997–1008.

Clavel, J., Aristide, L., Morlon, H., 2019. A Penalized Likelihood Framework for High-

Dimensional Phylogenetic Comparative Methods and an Application to New-World

Monkeys Brain Evolution. Syst. Biol. 68, 93–116.

Clavel, J., Escarguel, G., Merceron, G., 2015. mvMORPH: an R package for fitting

multivariate evolutionary models to morphometric data. Methods Ecol. Evol. 6, 1311–

1319.

Cruz-Laufer, A.J., Artois, T., Smeets, K., Pariselle, A., Vanhove, M.P.M., 2021. The cichlid–

Cichlidogyrus network: a blueprint for a model system of parasite evolution.

Hydrobiologia 848, 3847–3863.

de Meeûs, T., McCoy, K.D., Prugnolle, F., Chevillon, C., Durand, P., Hurtrez-Boussès, S.,

Renaud, F., 2007. Population genetics and molecular epidemiology or how to *débusquer*

la bête. Infect. Genet. Evol. 7, 308–332.

Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G.,

Gruber, B., Lafourcade, B., Leitão, P.J., Münkemüller, T., Mcclean, C., Osborne, P.E.,

Reineking, B., Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S., 2013.

Collinearity: A review of methods to deal with it and a simulation study evaluating their

performance. Ecography 36, 27–46.

Dossou, C., 1982. Parasites de poissons d'eau douce du Bénin. III. Espèces nouvelles du

genre *Cichlidogyrus* (Monogenea) parasites de Cichlidae. Bull. Inst. Fondam. Afr. Noire

Ser. A Sci. Nat. 44, 295–322.

Dossou, C., Birgi, E., 1984. Monogènes parasites d'Hemichromis fasciatus Peters, 1857

(Teleostei, Cichlidae). Ann. Sci. Nat. Zool. 6, 101–109.

Douëllou, L., 1993. Monogenean of the genus Cichlidogyrus Paperna, 1960 (Dactylogyridae:

Ancyrocephalinae) from cichlid fishes of Lake Kariba (Zimbabwe) with description of

five new species. Syst. Parasitol. 25, 159–186.

Dryden, I.L., Mardia, K. V, 2016. Statistical shape analysis: with applications in R, 2nd ed. ,

Wiley series in probability and statisticsJohn Wiley & Sons, Chichester, UK.

Dunz, A.R., Schliewen, U.K., 2013. Molecular phylogeny and revised classification of the

haplotilapiine cichlid fishes formerly referred to as *"Tilapia."* Mol. Phylogenet. Evol.

68, 64–80.

Ergens, R., 1981. Nine species of the genus *Cichlidogyrus* Paperna, 1960 (Monogenea:

Ancyrocephalinae) from Egyptian fishes. Folia Parasitol. 28, 205–214.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Fang, Z., Fan, J., Chen, X., Chen, Y., 2018. Beak identification of four dominant octopus

species in the East China Sea based on traditional measurements and geometric

morphometrics. Fish. Sci. 84, 975–985.

Fannes, W., Vanhove, M.P.M., Huyse, T., Paladini, G., 2015. A scanning electron

microscope technique for studying the sclerites of *Cichlidogyrus*. Parasitol. Res. 114,

2031–2034.

Felsenstein, J., 1973. Maximum likelihood estimation of evolutionary trees from continuous

characters. Am. J. Hum. Genet. 25, 471–492.

Firmat, C., Alibert, P., Mutin, G., Losseau, M., Pariselle, A., Sasal, P., 2016. A case of

complete loss of gill parasites in the invasive cichlid Oreochromis mossambicus.

Parasitol. Res. 115, 3657–3661.

Forbes, A.A., Powell, T.H.Q., Stelinski, L.L., Smith, J.J., Feder, J.L., 2009. Sequential

sympatric speciation across trophic levels. Science 323, 776–779.

Garcia-Cruz, J., Sosa, V., 2006. Coding quantitative character data for phylogenetic analysis:

a comparison of five methods. Syst. Bot. 31, 302–309.

García-Vásquez, A., Shinn, A.P., Bron, J.E., 2012. Development of a light microscopy stain

for the sclerites of *Gyrodactylus* von Nordmann, 1832 (Monogenea) and related genera.

Parasitol. Res. 110, 1639–1648.

Geraerts, M., Muterezi Bukinga, F., Vanhove, M.P.M., Pariselle, A., Chocha Manda, A.,

Vreven, E.J., Huyse, T., Artois, T., 2020. Six new species of *Cichlidogyrus* Paperna,

1960 (Platyhelminthes: Monogenea) from the gills of cichlids (Teleostei: Cichliformes)

from the Lomami River Basin (DRC: Middle Congo). Parasit. Vectors 13, 187.

Gillardin, C., Vanhove, M.P.M., Pariselle, A., Huyse, T., Volckaert, F.A.M., 2012.

Ancyrocephalidae (Monogenea) of Lake Tanganyika: II: description of the first

Cichlidogyrus spp. parasites from Tropheini fish hosts (Teleostei, Cichlidae). Parasitol.

Res. 110, 305–313.

Glor, R.E., 2010. Phylogenetic insights on adaptive radiation. Annu. Rev. Ecol. Evol. Syst.

41, 251–270.

Gobbin, T.P., Vanhove, M.P.M., Seehausen, O., Maan, M.E., 2020. Microhabitat

distributions and species interactions of ectoparasites on the gills of cichlid fish in lake

Victoria, Tanzania. Int. J. Parasitol.

Gobbin, T.P., Vanhove, M.P.M., Seehausen, O., Maan, M.E., Pariselle, A., 2021. Four new

species of *Cichlidogyrus* (Platyhelminthes, Monogenea, Dactylogyridae) from Lake

Victoria haplochromine cichlid fishes, with the redescription of *C. bifurcatus* and *C.*

longipenis [preprint]. bioRxiv 2021.01.29.428376.

Goloboff, P.A., 1993. Estimating character weights during tree search. Cladistics 9, 83–91.

Goloboff, P.A., 1999. Analyzing large data sets in reasonable times: solutions for composite

optima. Cladistics 15, 415–428.

Goloboff, P.A., 2003. Improvements to resampling measures of group support. Cladistics 19,

324–332.

Goloboff, P.A., 2014. Extended implied weighting. Cladistics 30, 260–272.

Goloboff, P.A., Carpenter, J.M., Arias, J.S., Esquivel, D.R.M., 2008a. Weighting against

homoplasy improves phylogenetic analysis of morphological data sets. Cladistics 24,

758–773.

Goloboff, P.A., Catalano, S.A., 2016. TNT version 1.5, including a full implementation of

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

phylogenetic morphometrics. Cladistics 32, 221–238.

Goloboff, P.A., Farris, J.S., Nixon, K.C., 2008b. TNT, a free program for phylogenetic

analysis. Cladistics 24, 774–786.

Goloboff, P.A., Mattoni, C.I., Quinteros, A.S., 2006. Continuous characters analyzed as such.

Cladistics 22, 589–601.

Goolsby, E.W., Bruggeman, J., Ané, C., 2017. Rphylopars: fast multivariate phylogenetic

comparative methods for missing data and within-species variation. Methods Ecol. Evol.

8, 22–27.

Goren, M., 2014. *Tristramella sacra*. IUCN Red List Threat. Species 2014

e.T61372A19010617.

Grant, P.R., 1999. Ecology and evolution of Darwin's finches, 2nd ed. Princeton Science

Library, Princeton University Press, Princeton, USA.

Gu, Z., Eils, R., Schlesner, M., 2016. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. Bioinformatics 32, 2847–2849.

Guerrero, A.M., Tye, A., 2009. Darwin's finches as seed predators and dispersers. Wilson J.

Ornithol. 121, 752–764.

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010.

New algorithms and methods to estimate maximum-likelihood phylogenies: assessing

the performance of PhyML 3.0. Syst. Biol. 59, 307–321.

Hansen, T.F., Martins, E.P., 1996. Translating between microevolutionary process and

macroevolutionary patterns: the correlation structure of interspecific data. Evolution 50,

1404–1417.

Harmon, L.J., Losos, J.B., Jonathan Davies, T., Gillespie, R.G., Gittleman, J.L., Bryan

Jennings, W., Kozak, K.H., McPeek, M.A., Moreno-Roark, F., Near, T.J., Purvis, A.,

Ricklefs, R.E., Schluter, D., Schulte Ii, J.A., Seehausen, O., Sidlauskas, B.L., Torres-

Carvajal, O., Weir, J.T., Mooers, A.Ø., 2010. Early bursts of body size and shape

evolution are rare in comparative data. Evolution 64, 2385–2396.

Hassouna, N., Michot, B., Bachellerie, J.-P., 1984. The compelte nucleotide sequences of

mouse 28S rRNA gene. Implications for the process of size increase of the large subunit

rRNA in higher eukaryotes. Nucleic Acids Res. 12, 3563–3583.

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., Le Vinh, S., 2018. UFBoot2:

Improving the Ultrafast Bootstrap Approximation. Mol. Biol. Evol. 35, 518–522.

Hsu, C., Chang, C., Lin, C., 2003. A practical guide to support vector machines. Technical

report. Department of Computer Science and Information Engineering, National Taiwan

University, Taipei, Taiwan. Available at:

https://www.csie.ntu.edu.tw/~cjlin/papers/guide/guide.pdf (accessed Mar 6, 2021).

Huyse, T., Poulin, R., Théron, A., 2005. Speciation in parasites: a population genetics

approach. Trends Parasitol. 21, 469–475.

Igeh, P.C., Avenant☐Oldewage, A., 2020. Pathological effects of *Cichlidogyrus philander*

Douëllou, 1993 (Monogenea, Ancyrocephalidae) on the gills of *Pseudocrenilabrus*

philander (Weber, 1897) (Cichlidae). J. Fish Dis. 43, 177–184.

Igeh, P.C., Dos Santos, Q.M., Avenant-Oldewage, A., 2017. Redescription of *Cichlidogyrus*

philander (Monogenea, Ancyrocephalidae) using scanning electron microscopy (SEM)

and molecular analysis. Parasite 24, 49.

Ingram, T., Mahler, D.L., 2013. SURFACE: Detecting convergent evolution from

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

comparative data by fitting Ornstein-Uhlenbeck models with stepwise Akaike

Information Criterion. Methods Ecol. Evol. 4, 416–425.

Ishiguro, M., Sakamoto, Y., Kitagawa, G., 1997. Bootstrapping log likelihood and EIC, an

extension of AIC. Ann. Inst. Stat. Math. 49, 411–434.

Jiménez-Garcia, M.I., Vidal-Martínez, V.M., Lopez-Jiménez, S., Jiménez-García, M.I.,

Vidal-Martínez, V.M., López-Jiménez, S., 2001. Monogeneans in introduced and native

cichlids in México: Evidence for transfer. J. Parasitol. 87, 907.

Jorissen, M.W.P., Huyse, T., Pariselle, A., Wamuini Lunkayilakio, S., Muterezi Bukinga, F.,

Chocha Manda, A., Kapepula Kasembele, G., Vreven, E.J., Snoeks, J., Decru, E., Artois,

T., Vanhove, M.P.M., 2020. Historical museum collections help detect parasite species

jumps after tilapia introductions in the Congo Basin. Biol. Invasions 11, 1123.

Jorissen, M.W.P., Pariselle, A., Huyse, T., Vreven, E.J., Snoeks, J., Decru, E., Kusters, T.,

Wamuini Lunkayilakio, S., Muterezi Bukinga, F., Artois, T., Vanhove, M.P.M., 2018a.

Six new dactylogyrid species (Platyhelminthes, Monogenea) from the gills of cichlids

(Teleostei, Cichliformes) from the Lower Congo Basin. Parasite 25, 64.

Jorissen, M.W.P., Pariselle, A., Huyse, T., Vreven, E.J., Snoeks, J., Volckaert, F.A.M.,

Chocha Manda, A., Kapepula Kasembele, G., Artois, T., Vanhove, M.P.M., 2018b.

Diversity and host specificity of monogenean gill parasites (Platyhelminthes) of cichlid

fishes in the Bangweulu-Mweru ecoregion. J. Helminthol. 92, 417–437.

Jorissen, M.W.P., Vanhove, M.P.M., Pariselle, A., Snoeks, J., Vreven, E.J., Šimková, A.,

Wamuini Lunkayilakio, S., Chocha Manda, A., Kapepula Kasembele, G., Muterezi

Bukinga, F., Artois, T., Huyse, T., 2021. Molecular footprint of parasite co-introduction

with Nile tilapia in the Congo Basin [preprint]. Research Square rs.3.rs-995291/v1.

Kaltz, O., Shykoff, J.A., 1998. Local adaptation in host–parasite systems. Heredity 81, 361–
370.

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermiin, L.S., 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods 14,
587–589.

Karvonen, A., Seehausen, O., 2012. The role of parasitism in adaptive radiations—When
might parasites promote and when might they constrain ecological speciation? Int. J.
Ecol. 2012.

Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol. Biol. Evol. 30, 772–780.

Kearn, G., Whittington, I., 2015. Sperm transfer in monogenean (platyhelminth) parasites.
Acta Parasitol. 60, 567–600.

Khabbazian, M., Kriebel, R., Rohe, K., Ané, C., 2016. Fast and accurate detection of
evolutionary shifts in Ornstein–Uhlenbeck models. Methods Ecol. Evol. 7, 811–824.

Kitagawa, G., Konishi, S., 2010. Bias and variance reduction techniques for bootstrap
information criteria. Ann. Inst. Stat. Math. 62, 209–234.

Klingenberg, C.P., 2010. Evolution and development of shape: integrating quantitative
approaches. Nat. Rev. Genet. 11, 623–635.

Kmentová, N., Cruz-Laufer, A.J., Pariselle, A., Smeets, K., Artois, T., Vanhove, M.P.M.,
2021a. Dactylogyridae 2021: seeing the forest through the (phylogenetic) trees.
EcoEvoRxiv 10.32942/osf.io/5mh3d.

Kmentová, N., Gelnar, M., Koblmüller, S., Vanhove, M.P.M., 2016a. First insights into the
diversity of gill monogeneans of "*Gnathochromis*" and *Limnochromis* (Teleostei,

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van
Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Cichlidae) in Burundi: Do the parasites mirror host ecology and phylogenetic history?

PeerJ 2016, e1629.

Kmentová, N., Gelnar, M., Koblmüller, S., Vanhove, M.P.M., 2016b. Deep-water parasite

diversity in Lake Tanganyika: Description of two new monogenean species from

benthopelagic cichlid fishes. Parasit. Vectors 9, 1–9.

Kmentová, N., Gelnar, M., Mendlová, M., Van Steenberge, M., Koblmüller, S., Vanhove,

M.P.M., 2016c. Reduced host-specificity in a parasite infecting non-littoral Lake

Tanganyika cichlids evidenced by intraspecific morphological and genetic diversity. Sci.

Rep. 6, 39605.

Kmentová, N., Hahn, C., Koblmüller, S., Zimmermann, H., Vorel, J., Artois, T., Gelnar, M.,

Vanhove, M.P.M., 2021b. Contrasting host-parasite population structure: Morphology

and mitogenomics of a parasitic flatworm on pelagic deepwater cichlid fishes from Lake

Tanganyika. Biology (Basel). 10, 797.

Kmentová, N., Koblmüller, S., Van Steenberge, M., Artois, T., Muterezi Bukinga, F.,

Mulimbwa N'sibula, T., Muzumani Risasi, D., Masilya Mulungula, P., Gelnar, M.,

Vanhove, M.P.M., 2020. Failure to diverge in African Great Lakes: the case of

Dolicirroplectanum lacustre gen. nov. comb. nov. (Monogenea, Diplectanidae) infecting

latid hosts. J. Great Lakes Res. 46, 1113–1130.

Kmentová, N., Van Steenberge, M., Raeymaekers, J.A.M., Koblmüller, S., Hablützel, P.I.,

Bukinga, F.M., N'sibula, T.M., Mulungula, P.M., Nzigidahera, B., Ntakimazi, G.,

Gelnar, M., Vanhove, M.P.M., 2018. Monogenean parasites of sardines in Lake

Tanganyika: diversity, origin and intraspecific variability. Contrib. Zool. 87, 105–132.

Kuhn, M., 2008. Building predictive models in R using the caret package. J. Stat. Softw. 28,

1–26.

Landis, J.R., Koch, G.G., 1977. The measurement of observer agreement for categorical data.

Biometrics 33, 159.

Landis, M.J., Freyman, W.A., Baldwin, B.G., 2018. Retracing the Hawaiian silversword

radiation despite phylogenetic, biogeographic, and paleogeographic uncertainty.

Evolution 72, 2343–2359.

Legendre, P., Lapointe, F.-J., 2004. Assessing congruence among distance matrices: single-

malt scotch whiskies revisited. Aust. N. Z. J. Stat. 46, 615–629.

Lévêque, C., Oberdorff, T., Paugy, D., Stiassny, M.L.J., Tedesco, P.A., 2008. Global

diversity of fish (Pisces) in freshwater. Hydrobiologia 595, 545–567.

Littlewood, D.T.J., Rohde, K., Clough, K.A., 1997. Parasite speciation within or between

host species? - Phylogenetic evidence from site-specific polystome monogeneans. Int. J.

Parasitol. 27, 1289–1297.

Llewellyn, J., 1963. Larvae and larval development of monogeneans. In: Dawes, B. (Ed.),

Advances in parasitology. Academic Press, London and New York, pp. 287–326.

Lockyer, A.E., Olson, P.D., Østergaard, P., Rollinson, D., Johnston, D.A., Attwood, S.W.,

Southgate, V.R., Horak, P., Snyder, S.D., Le, T.H., Agatsuma, T., McManus, D.P.,

Carmichael, A.C., Naem, S., Littlewood, D.T.J., 2003. The phylogeny of the

Schistosomatidae based on three genes with emphasis on the interrelationships of

Schistosoma Weinland, 1858. Parasitology 126, 203–224.

Losos, J.B., 2008. Phylogenetic niche conservatism, phylogenetic signal and the relationship

between phylogenetic relatedness and ecological similarity among species. Ecol. Lett.

11, 995–1003.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van
Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Losos, J.B., 2010. Adaptive radiation, ecological opportunity, and evolutionary determinism:
American society of naturalists E. O. Wilson Award address. Am. Nat. 175, 623–639.

Louca, S., Doebeli, M., 2018. Efficient comparative phylogenetics on large trees.
Bioinformatics 34, 1053–1055.

Mahler, D.L., Ingram, T., Revell, L.J., Losos, J.B., 2013. Exceptional convergence on the
macroevolutionary landscape in island lizard radiations. Science 341, 292–295.

Marchiori, N.C., Pariselle, A., Pereira, J., Agnèse, J.F., Durand, J.D., Vanhove, M.P.M.,
2015. A comparative study of *Ligophorus uruguayense* and *L. saladensis* (Monogenea:
Ancyrocephalidae) from *Mugil liza* (Teleostei: Mugilidae) in southern Brazil. Folia
Parasitol. 62, 024.

Matejusová, I., Gelnar, M., McBeath, A.J.A., Collins, C.M., Cunningham, C.O., 2001.
Molecular markers for gyrodactylids (Gyrodactylidae: Monogenea) from five fish
families (Teleostei). Int. J. Parasitol. 31, 738–745.

McCoy, K.D., 2003. Sympatric speciation in parasites – what is sympatry? Trends Parasitol.
19, 400–404.

McGee, M.D., Neches, R.Y., Seehausen, O., 2016. Evaluating genomic divergence and
parallelism in replicate ecomorphs from young and old cichlid adaptive radiations. Mol.
Ecol. 25, 260–268.

Meinilä, M., Kuusela, J., Ziⵜtara, M.S., Lumme, J., 2004. Initial steps of speciation by
geographic isolation and host switch in salmonid pathogen *Gyrodactylus salaris*
(Monogenea: Gyrodactylidae). Int. J. Parasitol. 34, 515–526.

Mendlová, M., Desdevises, Y., Civáňová, K., Pariselle, A., Šimková, A., 2012. Monogeneans

of West African cichlid fish: evolution and cophylogenetic interactions. PLoS One 7.

Mendlová, M., Šimková, A., 2014. Evolution of host specificity in monogeneans parasitizing

African cichlid fish. Parasit. Vectors 7, 69.

Messu Mandeng, F.D., Bilong Bilong, C.F., Pariselle, A., Vanhove, M.P.M., Bitja Nyom,

A.R., Agnèse, J.F., 2015. A phylogeny of *Cichlidogyrus* spp. (Monogenea,

Dactylogyridea) clarifies a host-switch between fish families and reveals an adaptive

component to attachment organ morphology of this parasite genus. Parasit. Vectors 8,

582.

Meyer, D., Dimitriadou, E., Hornik, K., Weingessel, A., Leisch, F., 2020. e1071: Misc

Functions of the Department of Statistics, Probability Theory Group (Formerly: E1071),

TU Wien: R package version 1.7-4. Vienna, Austria. Available at: https://cran.r-

project.org/web/packages/e1071/e1071.pdf (accessed Oct 27, 2020).

Miller, M.A., Pfeiffer, W., Schwartz, T., 2010. Creating the CIPRES Science Gateway for

inference of large phylogenetic trees. 2010 Gatew. Comput. Environ. Work. GCE 2010.

Mirande, J.M., 2009. Weighted parsimony phylogeny of the family Characidae (Teleostei:

Characiformes). Cladistics 25, 574–613.

Mirande, J.M., 2019. Morphology, molecules and the phylogeny of Characidae (Teleostei,

Characiformes). Cladistics 35, 282–300.

Murtagh, F., Legendre, P., 2014. Ward's hierarchical agglomerative clustering method:

Which algorithms implement Ward's criterion? J. Classif. 31, 274–295.

Muterezi Bukinga, F., Vanhove, M.P.M., Van Steenberge, M., Pariselle, A., 2012.

Ancyrocephalidae (Monogenea) of Lake Tanganyika: III: *Cichlidogyrus* infecting the

world's biggest cichlid and the non-endemic tribes Haplochromini, Oreochromini and

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Tylochromini (Teleostei, Cichlidae). Parasitol. Res. 111, 2049–2061.

Nguyen, L.-T., Schmidt, H.A., von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: a fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol.

Biol. Evol. 32, 268–274.

Nixon, K.C., 1999. The parsimony ratchet, a new method for rapid parsimony analysis.

Cladistics 15, 407–414.

O'Meara, B.C., Ané, C., Sanderson, M.J., Wainwright, P.C., 2006. Testing for different rates

of continuous trait evolution using likelihood. Evolution 60, 922.

Pagel, M., 1999. Inferring the historical patterns of biological evolution. Nature 401, 877–

884.

Paperna, I., 1960. Studies on monogenetic trematodes in Israel. 2. Monogenetic trematodes of

cichlids. Bamidgeh 12, 20–33.

Paperna, I., 1968. Monogenetic trematodes collected from fresh water fish in Ghana. Second

report. Bamidgeh 17, 107–111.

Paperna, I., 1979. Monogenea of inland water fish in Africa. Ann. Mus. R. Afr. Cent.

Tervuren Belg. Ser. IN-8 Sci. Zool. 226, 1–131.

Paperna, I., Thurston, J.P., 1969. Monogenetic trematodes collected from cichlid fish in

Uganda; including the description of five new species of *Cichlidogyrus*. Rev. Zool. Bot.

Afr. 79, 15–33.

Parins-Fukuchi, C., 2018. Use of continuous traits can improve morphological phylogenetics.

Syst. Biol. 67, 328–339.

Pariselle, A., Bitja Nyom, A.R., Bilong Bilong, C.F., 2014. Four new species of

Cichlidogyrus (Monogenea, Ancyrocephalidae) from *Sarotherodon mvogoi* and

Tylochromis sudanensis (Teleostei, Cichlidae) in Cameroon. Zootaxa 3881, 258–266.

Pariselle, A., Bitja Nyom, A.R., Bilong Bilong, C.F., Bilong, C.F., 2013. Checklist of the

ancyrocephalids (Monogenea) parasitizing *Tilapia* species in Cameroon, with the

description of three new species. Zootaxa 3599, 78–86.

Pariselle, A., Boeger, W.A., Snoeks, J., Bilong Bilong, C.F., Morand, S., Vanhove, M.P.M.,

2011. The monogenean parasite fauna of cichlids: a potential tool for host biogeography.

Int. J. Evol. Biol. 2011, 471480.

Pariselle, A., Euzet, L., 1994. Three new species of *Cichlidogyrus* Paperna, 1960

(Monogenea, Ancyrocephalidae) parasitic on *Tylochromis jentinki* (Steindachner, 1895)

(Pisces, Cichlidae) in West Africa. Syst. Parasitol. 29, 229–234.

Pariselle, A., Euzet, L., 1995. *Scutogyrus* gen.n. (Monogenea: Ancyrocephalidae) for

Cichlidogyrus longicornis minus Dossou, 1982, *C. l. longicornis*, and *C. l. gravivaginus*

Paperna and Thurston, 1969, with description of three new species parasitic on African .

J. Helminthol. Soc. Washingt. 62, 157–173.

Pariselle, A., Euzet, L., 1996. *Cichlidogyrus* Paperna, 1960 (Monogenea, Ancyrocephalidae):

gill parasites from West African Cichlidae of the subgenus *Coptodon* Regan, 1920

(pisces), with descriptions of six new species. Syst. Parasitol. 34, 109–124.

Pariselle, A., Euzet, L., 2003. Four new species of *Cichlidogyrus* (Monogenea:

Ancyrocephalidae), gill parasites of *Tilapia cabrae* (Teleostei: Cichlidae), with

discussion on relative length of haptoral sclerites. Folia Parasitol. 50, 195–201.

Pariselle, A., Euzet, L., 2004. Two new species of *Cichlidogyrus* Paperna, 1960 (Monogenea,

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Ancyrocephalidae) gill parasites on *Hemichromis fasciatus* (Pisces, Cichlidae) in Africa,

with remarks on parasite geographical distribution. Parasite 11, 359–364.

Pariselle, A., Euzet, L., 2009. Systematic revision of dactylogyridean parasites (Monogenea)

from cichlid fishes in Africa, the Levant and Madagascar. Zoosystema 31, 849–898.

Pariselle, A., Morand, S., Deveney, M.R., Pouyaud, L., 2003. Parasite species richness of

closely related hosts: historical scenario and "genetic" hypothesis. In: Combes, C.,

Jourdane, J., Ducreux-Modat, A., Pages, J.-R. (Eds.), Taxonomie, écologie et évolution

des métazoaires parasites: Livre-hommage à Louis Euzet. Presses Universitaires de

Perpignan, Perpignan, France, pp. 147–166.

Pariselle, A., Muterezi Bukinga, F., Van Steenberge, M., Vanhove, M.P.M., 2015a.

Ancyrocephalidae (Monogenea) of Lake Tanganyika: IV: *Cichlidogyrus* parasitizing

species of Bathybatini (Teleostei, Cichlidae): reduced host-specificity in the deepwater

realm? Hydrobiologia 748, 99–119.

Pariselle, A., Van Steenberge, M., Snoeks, J., Volckaert, F.A.M., Huyse, T., Vanhove,

M.P.M., 2015b. Ancyrocephalidae (Monogenea) of Lake Tanganyika: Does the

Cichlidogyrus parasite fauna of *Interochromis loocki* (Teleostei, Cichlidae) reflect its

host's phylogenetic affinities? Contrib. Zool. 84, 25–38.

Pennell, M.W., Eastman, J.M., Slater, G.J., Brown, J.W., Uyeda, J.C., Fitzjohn, R.G., Alfaro,

M.E., Harmon, L.J., 2014. Geiger v2.0: an expanded suite of methods for fitting

macroevolutionary models to phylogenetic trees. Bioinformatics 30, 2216–2218.

Poulin, R., 2014. Parasite biodiversity revisited: frontiers and constraints. Int. J. Parasitol. 44,

581–589.

Poulin, R., Hay, E., Jorge, F., 2019. Taxonomic and geographic bias in the genetic study of

helminth parasites. Int. J. Parasitol. 49, 429–435.

Poulin, R., Krasnov, B.R., Mouillot, D., 2011. Host specificity in phylogenetic and

geographic space. Trends Parasitol. 27, 355–361.

Poulin, R., Presswell, B., Jorge, F., 2020. The state of fish parasite discovery and taxonomy:

a critical assessment and a look forward. Int. J. Parasitol. 50, 733–742.

Pouyaud, L., Desmarais, E., Deveney, M., Pariselle, A., 2006. Phylogenetic relationships

among monogenean gill parasites (Dactylogyridea, Ancyrocephalidae) infesting tilapiine

hosts (Cichlidae): systematic and evolutionary implications. Mol. Phylogenet. Evol. 38,

241–249.

Price, C.E., Kirk, R.G., 1967. First description of a monogenetic trematode from Malawi.

Rev. Zool. Bot. Afr. 76, 137–144.

Price, C.E., Peebles, H.E., Bamford, T., 1969. The monogenean parasites of African fishes.

IV. Two new species from South African hosts. Rev. Zool. Bot. Afr. 79, 117–124.

Puttick, M.N., 2018. Mixed evidence for early bursts of morphological evolution in extant

clades. J. Evol. Biol. 31, 502–515.

R Core Team, 2021. R: a language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. Available at: https://www.r-project.org/

(accessed Jul 5, 2021).

Rahmouni, C., Van Steenberge, M., Vanhove, M.P.M., Šimková, A., 2021. Intraspecific

morphological variation in *Cichlidogyrus* (Monogenea) parasitizing two cichlid hosts

from Lake Tanganyika exhibiting different dispersal capacities. Hydrobiologia 848,

3833–3845.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Rahmouni, C., Vanhove, M.P.M., Šimková, A., 2017. Underexplored diversity of gill

monogeneans in cichlids from Lake Tanganyika: eight new species of Cichlidogyrus

Paperna, 1960 (Monogenea: Dactylogyridae) from the northern basin of the lake, with

remarks on the vagina and the heel of the male copulatory . Parasit. Vectors 10, 591.

Rahmouni, C., Vanhove, M.P.M., Šimková, A., 2018. Seven new species of *Cichlidogyrus*

Paperna, 1960 (Monogenea: Dactylogyridae) parasitizing the gills of Congolese cichlids

from northern Lake Tanganyika. PeerJ 6, e5604.

Ramasamy, P., Ramalingam, K., Hanna, R.E.B., Halton, D.W., 1985. Microhabitats of gill

parasites (Monogenea and copepoda) of teleosts (*Scomberoides* spp.). Int. J. Parasitol.

15, 385–397.

Rambaut, A., Drummond, A.J., Xie, D., Baele, G., Suchard, M.A., 2018. Posterior

summarization in Bayesian phylogenetics using Tracer 1.7. Syst. Biol. 67, 901–904.

Řehulková, E., Kičinjaová, M.L., Mahmoud, Z.N., Gelnar, M., Seifertová, M., 2019. Species

of *Characidotrema* Paperna & Thurston, 1968 (Monogenea: Dactylogyridae) from

fishes of the Alestidae (Characiformes) in Africa: new species, host-parasite associations

and first insights into the phylogeny of the genus. Parasit. Vectors 12, 366.

Řehulková, E., Mendlová, M., Šimková, A., 2013. Two new species of *Cichlidogyrus*

(Monogenea: Dactylogyridae) parasitizing the gills of African cichlid fishes

(Perciformes) from Senegal: morphometric and molecular characterization. Parasitol.

Res. 112, 1399–1410.

Revell, L.J., 2012. phytools: an R package for phylogenetic comparative biology (and other

things). Methods Ecol. Evol. 3, 217–223.

Rodríguez-González, A., Sarabeev, V., Balbuena, J.A., 2017. Evolutionary morphology in

shape and size of haptoral anchors in 14 *Ligophorus* spp. (Monogenea: Dactylogyridae).

PLoS One 12, e0178367.

Rohde, K., 1976. Monogenean gill parasites of *Scomberomorus commersoni* Lacépède and

other mackerel on the Australian East Coast. Zeitschrift für Parasitenkd. 51, 49–69.

Ronco, F., Büscher, H.H., Indermaur, A., Salzburger, W., 2020. The taxonomic diversity of

the cichlid fish fauna of ancient Lake Tanganyika, East Africa. J. Great Lakes Res. 46,

1067–1078.

Ronquist, F., Huelsenbeck, J.P., 2003. MrBayes 3: Bayesian phylogenetic inference under

mixed models. Bioinformatics 19, 1572–1574.

Salzburger, W., 2018. Understanding explosive diversification through cichlid fish genomics.

Nat. Rev. Genet. 19, 705–717.

Salzburger, W., van Bocxlaer, B., Cohen, A.S., 2014. Ecology and evolution of the African

Great Lakes and their faunas. Annu. Rev. Ecol. Evol. Syst. 45, 519–545.

Schedel, F.D.B., Musilova, Z., Schliewen, U.K., 2019. East African cichlid lineages

(Teleostei: Cichlidae) might be older than their ancient host lakes: new divergence

estimates for the East African cichlid radiation. BMC Evol. Biol. 19, 1–25.

Seehausen, O., 2006. African cichlid fish: a model system in adaptive radiation research.

Proc. R. Soc. B Biol. Sci. 273, 1987–1998.

Šimková, A., Ondračková, M., Gelnar, M., Morand, S., 2002. Morphology and coexistence of

congeneric ectoparasite species: reinforcement of reproductive isolation? Biol. J. Linn.

Soc. 76, 125–135.

Šimková, A., Plaisance, L., Matějusová, I., Morand, S., Verneau, O., 2003. Phylogenetic

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

relationships of the Dactylogyridae Bychowsky, 1933 (Monogenea: Dactylogyridea):

The need for the systematic revision of the Ancyrocephalinae Bychowsky, 1937. Syst.

Parasitol. 54, 1–11.

Šimková, A., Řehulková, E., Rasoloariniaina, J.R., Jorissen, M.W.P., Scholz, T., Faltýnková,

A., Mašová, Š., Vanhove, M.P.M., 2019. Transmission of parasites from introduced

tilapias: a new threat to endemic Malagasy ichthyofauna. Biol. Invasions 21, 803–819.

Šimková, A., Serbielle, C., Pariselle, A., Vanhove, M.P.M., Morand, S., 2013. Speciation in

Thaparocleidus (Monogenea: Dactylogyridae) parasitizing Asian pangasiid catfishes.

Biomed Res. Int. 2013, 353956.

Takahashi, T., Koblmüller, S., 2011. The adaptive radiation of cichlid fish in Lake

Tanganyika: a morphological perspective. Int. J. Evol. Biol. 2011, 1–14.

Talavera, G., Castresana, J., 2007. Improvement of phylogenies after removing divergent and

ambiguously aligned blocks from protein sequence alignments. Syst. Biol. 56, 564–577.

Thurston, J.P., 1970. The incidence of Monogenea and parasitic Crustacea on the gills of fish

in Uganda. Rev. Zool. Bot. Afr. 82, 111–130.

Turner, G.F., Seehausen, O., Knight, M.E., Allender, C.J., Robinson, R.L., 2001. How many

species of cichlid fishes are there in African lakes? Mol. Ecol. 10, 793–806.

Uyeda, J.C., Harmon, L.J., 2014. A novel Bayesian method for inferring and interpreting the

dynamics of adaptive landscapes from phylogenetic comparative data. Syst. Biol. 63,

902–918.

Van Steenberge, M., Pariselle, A., Huyse, T., Volckaert, F.A.M., Snoeks, J., Vanhove,

M.P.M., 2015. Morphology, molecules, and monogenean parasites: an example of an

integrative approach to cichlid biodiversity. PLoS One 10, e0124474.

Van Steenberge, M., Vanhove, M.P.M., Muzumani Risasi, D., Mulimbwa N'Sibula, T.,

Muterezi Bukinga, F., Pariselle, A., Gillardin, C., Vreven, E.J., Raeymaekers, J.A.M.,

Huyse, T., Volckaert, F.A.M., Nshombo Muderhwa, V., Snoeks, J., 2011. A recent

inventory of the fishes of the north-western and central western coast of Lake

Tanganyika (Democratic Republic Congo). Acta Ichthyol. Piscat. 41, 201–214.

Vanhove, M.P.M., Briscoe, A.G., Jorissen, M.W.P., Littlewood, D.T.J., Huyse, T., 2018. The

first next-generation sequencing approach to the mitochondrial phylogeny of African

monogenean parasites (Platyhelminthes: Gyrodactylidae and Dactylogyridae). BMC

Genomics 19, 520.

Vanhove, M.P.M., Hablützel, P.I., Pariselle, A., Šimková, A., Huyse, T., Raeymaekers,

J.A.M., 2016. Cichlids: a host of opportunities for evolutionary parasitology. Trends

Parasitol. 32, 820–832.

Vanhove, M.P.M., Pariselle, A., Van Steenberge, M., Raeymaekers, J.A.M., Hablützel, P.I.,

Gillardin, C., Hellemans, B., Breman, F.C., Koblmüller, S., Sturmbauer, C., Snoeks, J.,

Volckaert, F.A.M., Huyse, T., 2015. Hidden biodiversity in an ancient lake:

phylogenetic congruence between Lake Tanganyika tropheine cichlids and their

monogenean flatworm parasites. Sci. Rep. 5, 13669.

Vanhove, M.P.M., Van Steenberge, M., Dessein, S., Volckaert, F.A.M., Snoeks, J., Huyse,

T., Pariselle, A., 2013. Biogeographical implications of Zambezian *Cichlidogyrus*

species (Platyhelminthes: Monogenea: Ancyrocephalidae) parasitizing Congolian

cichlids. Zootaxa 3608, 398–400.

Vanhove, M.P.M., Volckaert, F.A.M., Pariselle, A., 2011. Ancyrocephalidae (Monogenea) of

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

1705 Lake Tanganyika: I: four new species of *Cichlidogyrus* from *Ophthalmotilapia ventralis*

1706 (Teleostei: Cichlidae), the first record of this parasite family in the basin. Zoologia 28,

1707 253–263.

1708 Vignon, M., Pariselle, A., Vanhove, M.P.M., 2011. Modularity in attachment organs of

1709 African *Cichlidogyrus* (Platyhelminthes: Monogenea: Ancyrocephalidae) reflects

1710 phylogeny rather than host specificity or geographic distribution. Biol. J. Linn. Soc. 102,

1711 694–706.

1712 Vignon, M., Sasal, P., 2010. The use of geometric morphometrics in understanding shape

1713 variability of sclerotized haptoral structures of monogeneans (Platyhelminthes) with

1714 insights into biogeographic variability. Parasitol. Int. 59, 183–191.

1715 Weiblen, G.D., Bush, G.L., 2002. Speciation in fig pollinators and parasites. Mol. Ecol. 11,

1716 1573–1578.

1717 Wickham, H., 2016. ggplot2: Elegant graphics for data analysis, 2nd ed. , Use R!Springer,

1718 New York, USA.

1719 Wu, X.Y., Zhu, X.Q., Xie, M.Q., Li, A.X., 2006. The radiation of *Haliotrema* (Monogenea:

1720 Dactylogyridae: Ancyrocephalinae): molecular evidence and explanation inferred from

1721 LSU rDNA sequences. Parasitology 132, 659–668.

1722 Wu, X.Y., Zhu, X.Q., Xie, M.Q., Li, A.X., 2007. The evaluation for generic-level monophyly

1723 of Ancyrocephalinae (Monogenea, Dactylogyridae) using ribosomal DNA sequence

1724 data. Mol. Phylogenet. Evol. 44, 530–544.

1725 Yu, G., Lam, T.T.-Y., Zhu, H., Guan, Y., 2018. Two methods for mapping and visualizing

1726 associated data on phylogeny using ggtree. Mol. Biol. Evol. 35, 3041–3043.

Yu, G., Smith, D.K., Zhu, H., Guan, Y., Lam, T.T.-Y., 2017. ggtree□: An R package for

visualization and annotation of phylogenetic trees with their covariates and other

associated data. Methods Ecol. Evol. 8, 28–36.

Zhi, T., Xu, X., Chen, J., Zheng, Y., Zhang, S., Peng, J., Brown, C.L., Yang, T., 2018.

Expression of immune-related genes of Nile tilapia *Oreochromis niloticus* after

Gyrodactylus cichlidarum and *Cichlidogyrus sclerosus* infections demonstrating

immunosupression in coinfection. Fish Shellfish Immunol. 80, 397–404.

Zischke, M.T., Litherland, L., Tilyard, B.R., Stratford, N.J., Jones, E.L., Wang, Y.-G., 2016.

Otolith morphology of four mackerel species (*Scomberomorus* spp.) in Australia:

species differentiation and prediction for fisheries monitoring and assessment. Fish. Res.

176, 39–47.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

LIST OF FIGURES

Figure 1. Overview of morphology of species of *Cichlidogyrus* Paperna, 1960

(Platyhelminthes: Monogenea, Dactylogyridae). (a) Multiple specimens attached to the gills

of *Sarotherodon melanotheron* Rüppel, 1852 (Cichliformes: Cichlidae). (b) Microscopic

image of *Cichlidogyrus agnesi* Pariselle & Euzet, 1995 with sclerotised structures of

reproductive (male copulatory organ and vagina) and attachment organs indicated by arrows.

(c) Overview of hard part morphology and most widely applied measurements of these

structures in taxonomic literature. Abbreviations: a, anchor total length; b, anchor blade

length; c, anchor shaft length; d, anchor guard length; e, anchor point length; 1–7, hook

lengths; w, bar width; x, bar length; h, auricle length; y, distance between auricles; AuP,

surface of auxiliary plate of male copulatory organ; AP, accessory piece length; Pe, penis

length; He, heel length; l, length; w, width. Terminology and methodology of measurements

according to Fannes et al. (2017).

Figure 2. Phylograms of three monogenean flatworms belonging to *Cichlidogyrus* and

Scutogyrus (Platyhelminthes, Monogenea, Dactylogyridae) based on three nuclear (18S, 28S,

and ITS rDNA) and one mitochondrial (CO1 mtDNA) DNA sequence markers. Sequences of

specimens in bold have been generated for this study. (a) Final hypothesis of analysis under

maximum parsimony and extended implied weighting (k = 48, weighting scheme BLK) with

node support estimated through symmetric resampling (p = 0.33). (b) Bayesian phylogram

with Bayesian posterior probabilities (PP) followed by ultrafast bootstrap values (UFBoot)

and Shimodaira-Hasegawa-like approximate likelihood ratios (SH-aLRT) inferred from

maximum likelihood estimation indicated at nodes; asterisk (*) indicates low or moderate

support below the threshold (PP < 0.95, UFBoot < 95, SH-aLRT < 80); black dots, internal

nodes with strong support across all support values. Node labels (1-11), monophyletic clades

considered strongly supported: species infecting (mostly) hemichromine cichlids (*Hemi*),

species belonging to *Scutogyrus* (*Scutogyrus*), species infecting (mostly) coptodonine cichlids

among others (*Cop*), the first species group infecting oreochromine cichlids among others

(*Oreo1*), species infecting cichlids belonging to the East African Radiation (*EAR*), species

infecting coptodonine, pelmatolapiine oreochromine, tilapiine, heterotilapiine, and

gobiocichline cichlids (*CPO*), species from Southern Africa with a bulbous penis (*Bulb*), the

second species group infecting mainly oreochromine cichlids among others (*Oreo2*), *C. halli*–

clade (*Halli*), and species infecting tylochromine cichlids (*Tylo*). Abbreviation: *C.*

schreyenbrichard., *C. schreyenbrichardorum.*

Figure 3. Host repertoire of species of *Cichlidogyrus.* (a) Character map of index of

specificity according to Mendlová and Šimková (2014). (b) Host range matrix of the species

by tribe or subfamily of cichlid hosts and family of non-cichlid hosts (recent anthropogenic

host range expansions italicised). APLO, Aplocheilidae; NOTH, Nothobranchiidae; POLY,

Polycentridae; Pare, Paretroplinae; Ptyc, Ptychochrominae; Cich, Cichlasomatini; Chro,

Chromidotilapiini; Tylo, Tylochromini; Hemi, Hemichromini; Gobi, Gobiocichlini; Copt,

Coptodonini; Hete, Heterotilapiini, PelL, Pelmatolapiini; Oreo, Oreochromini; Tila, Tilapiini;

Boul, Boulengerochromini; Bath, Bathybatini; Trem, Trematocarini; Bent, Benthochromini;

Cypr, Cyprichromini; Ecto, Ectodini; Hapl, Haplochromini.

Figure 4. Consensus tree of phylogenetic analysis under maximum parsimony and extended

implied weighting (k = 48, BLK weighting scheme) combining molecular and morphometric

data; red tip points indicate species with and blue without DNA sequence data; node support

values constitute GC values inferred from symmetric resampling. Compared to the molecular

tree (Fig. 2a), this tree is less resolved and less supported.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Figure 5. Character maps of morphological characters. (a) Scaled means of morphological

continuous characters with abbreviations referring to measurements of sclerotised structures

in the attachment and reproductive organ used to characterise species belonging to

Cichlidogyrus (see Fig. 1). (b) Character maps of newly proposed discrete characters for the

reproductive organs; indices in legend refer to character states suggested in Appendix 2. For

details on all characters, see numbers in Appendix 2. Numbers on nodes refer to clades in Fig.

2a.

Figure 6. Morphology of attachment and reproductive organs of representative species

belonging to proposed species groups and subgroups (Fig. 2) of *Cichlidogyrus* (incl.

Scutogyrus) including cladogram and illustrations of sclerotised structures of attachment

organ (anchors, bars, hooks) and the MCO of selected species (vagina morphology not shown

as little unifying or contrasting morphological patterns were detected between groups, see

"Characterisation of species groups"). Arrows indicate key features of each species group

(blue) and subgroup of the *EAR* group (yellow), dashed lines indicate typical shapes. Species

of the *EAR* group that are not displayed are labelled in grey. Scale = 30 μm.

Figure 7. Systematic informativeness of morphometric measurements (a) Variable

importance (AUC) according the optimised support vector machines (SVM) classifying

specimens belonging to *Cichlidogyrus* into proposed species groups; colours indicate

contribution by species group; AUC values > 0.8 are considered excellent discriminators;

AUC values between 0.7 and 0.8 are considered acceptable discriminators; AUC values < 0.7

are considered poor discriminators. (b) AUC values clustered by clade showing that the

SVMs could distinguish species groups with different accuracy. (c). Cluster analysis of

morphometric measurements of attachment and reproductive organs; clusters were detected

using the *ward.d2* clustering algorithm. For abbreviation of measurements, see Fig. 1. Species

groups were proposed based on phylogenetic analysis (Fig. 2).

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

Figure 8. Results of phylogenetic comparative analysis applied to one set of 100 randomly

sampled tree topologies from the Bayesian post–burn in phase. Fitted continuous models

include Brownian Motion (BM), Ornstein-Uhlenbeck (OU), Early-Burst (EB), Late-Burst

(LB), and multi-rate Brownian Motion (BMM) models simulating a random walk (H_1),

stabilising selection (H_2), adaptive radiation with a decelerating (H_3) and accelerating (H_4)

divergence of characters, and divergent BM regimes in parts of the phylogeny (H_5). (a) Model

fits were assessed through the Extended Information Criterion (EIC) for multivariate analysis.

All models assuming a single evolutionary regime across the phylogeny performed similarly

(LB performed worse). However, the BMM model suggesting a different regime for the EAR

clade outperformed the latter. (b) Univariate PCMs for discrete characters including the hook

configuration (HC) (Vignon et al. 2011), anchor similarity (AS), ventral bar shape (VBS)

(Mendlová et al. 2012), and index of (host) specificity (IS) (Mendlová and Šimková 2014).

HC and VBS show a strong phylogenetic signal, AS and IS shows no detectable phylogenetic

signal. Model performance is assessed through difference in the sample size–corrected Akaike

Information Criterion (ΔAICc) compared to a white noise model under the assumption of a

star-like phylogeny (absolute phylogenetic independence).

Figure 9. Character maps of discrete characters for the attachment organ morphology as

defined by Vignon et al. (2011) and Mendlová et al. (2012) vs. the respective set of

continuous measurement suggested in this study. The character maps indicate that

summarising multiple variables into a single discrete character might cause the information

loss observed as a lack of a phylogenetic signal in Fig. 8b. Numbers on nodes refer to clades

in Fig. 2a. Hook configuration A, B, and C assigned according to criteria by Vignon et al.

(2011) with relative size of hook pair 1 vs pairs 3–7; haptor configuration D equates to the

configuration of *C. arthracanthus* with large hooks 1 and 3–7. Anchor similarity either similar

(1) or dissimilar (2). Ventral bar shape with (1) or without (2) membranous extensions,

massive with extensions (3) or support large plate (4).LIST OF APPENDICES (IN PRINT)

Appendix 1. Sampling localities and dates of specimens belonging to *Cichlidogyrus* and

Scutogyrus collected for this study including a reference for the samples previously included

in taxonomic studies. Freshwater ecoregions are assigned according to Thieme et al. (2005).

Appendix 2. List of continuous and discrete characters and character states inferred from the

taxonomic literature and used for character mapping in Fig. 1. All characters represent

characteristics that are commonly used to described species of *Cichlidogyrus* and *Scutogyrus*.

Appendix 3. Specimen data of cichlid parasites of the genera *Cichlidogyrus* and *Scutogyrus*

used for phylogenetic analyses including host species, GenBank accession numbers, locality

by country, and reference. Voucher/isolate ID and accession numbers in italics indicate

specimens not included in subset trees used for phylogenetic comparative methods.

Appendix 4. Substitution models of molecular evolution and partitions for Bayesian inference

(BI) and maximum likelihood estimation (ML) of phylogeny of species of *Cichlidogyrus* and

Scutogyrus. Models include the general time reversible model (GTR), the Kimura 1980 model

(K80), the transitional model 3 with unequal base frequencies (TIM3e), the Tamura-Nei

model (TN), and the three-parameter model 2 (TPM2) plus empirical base frequencies (+ F), a

proportion of invariable sites (+ I), a discrete Γ model with four rate categories (Γ4), or a

FreeRate model with three categories (+ R3). For model specification see the IQ-TREE

ModelFinder manual (Kalyaanamoorthy et al., 2017).

Appendix 5. Species groups of *Cichlidogyrus* with species included, species potentially

included, and the respective host ranges reported in the taxonomic literature including host

species of candidate species.

Cruz-Laufer AJ, Pariselle A, Jorissen MWP, Muterezi Bukinga F, Al Assadi A, Van

Steeberge M, Koblmüller S, Sturmbauer C, Huyse T, Smeets K, Artois T, Vanhove MPM

1857   **Appendix 6.** Ancestral states of continuous characters inferred from a literature survey and

1858   used for character mapping. Character IDs refer to numbers in Appendix 2.

1859   **Appendix 7.** Ancestral states of discrete characters inferred from a literature survey and used

1860   for character mapping. Values represent probabilities for different character states. Character

1861   IDs refer to numbers in Appendix 2.

1862

1863   SUPPORTING INFORMATION

1864   **File S1.** Input data matrix for use in *TNT* including morphometric and DNA sequence data.

1865   **File S2.** Raw morphometric measurements for species of *Cichlidogyrus* and *Scutogyrus*.

c Attachment organ   Reproductive organs

Male copulatory organ

AP

Ventral bar

Vagina

Vagina

b

Male copulatory organ

Attachment organ

(a)

(b)

0.03

(a)

(b)

1: species-specific
2: genus-specific
3: tribe-specific
4: generalist

0.9

(a)

1: DAa/VAb 2: DAa/VAa 3: DAd/DAc 4: VAd/VAc 5: DBh/DBx 6: VBw/VBx 7: U1/U2 8: U7/U2 9: Pe 12: AP 14: He 15: AuP 16: VgL 17: VgW

Relative size
0   4   8

(b)

10: Shape of penis
- 10a
- 10b
- 10c
- 10d
- 10e
- 10f
- 10g

11: Diameter of penis
- 11a
- 11b
- 11c

13: Shape of accessory piece
- 13a
- 13b
- 13c
- 13d
- 13e
- 13f
- 13g
- 13h
- 13i
- 13j
- 13k
- 13l
- 13m
- 13n

18: Shape of sclerotised vagina
- 18a
- 18b
- 18c
- 18d

(a)

(b)

(a)

Hook configuration

- A
- B
- C
- D

Anchor similarity

- 1
- 2

Ventral bar shape

- 1
- 2
- 3
- 4

(b)

U1–U7

DAd, VAd

VBw, VBx