A protease-activatable luminescent biosensor and

reporter cell line for authentic SARS-CoV-2 infection

Pehuén Pereyra Gerber[1,2], Lidia M Duncan[1,2], Edward JD Greenwood[1,2], Sara Marelli[1,2], Adi

Naamati[1,2], Ana Teixeira-Silva[1,2], Thomas WM Crozier[1,2], Ildar Gabaev[1,2], Jun R Zhan[1,2],

Thomas E Mulroney[3], Emily C Horner[3], Rainer Doffinger[4], Anne E Willis[3], James ED

Thaventhiran[1,3], Anna V Protasio[5], Nicholas J Matheson[1,2,6,*]

1. Department of Medicine, University of Cambridge, UK

2. Cambridge Institute for Therapeutic Immunology and Infectious Disease (CITIID), Jeffrey

Cheah Biomedical Centre, University of Cambridge, UK

3. MRC Toxicology Unit, University of Cambridge, UK

4. Department of Immunology, Cambridge University Hospitals NHS Foundation Trust, UK

5. Department of Pathology, University of Cambridge, UK

6. NHS Blood and Transplant, Cambridge, UK

*Correspondence: njm25@cam.ac.uk

Short title: *A luminescent reporter cell line for SARS-CoV-2 replication*

Abstract

Efforts to define serological correlates of protection against COVID-19 have been hampered by the lack of a simple, scalable, standardised assay for SARS-CoV-2 replication and antibody neutralisation. Plaque assays remain the gold standard, but are impractical for high-throughput screening. In this study, we show that expression of viral proteases may be used to quantitate infected cells. Our assays exploit the cleavage of specific oligopeptide linkers, leading to the activation of cell-based optical biosensors. First, we characterise these biosensors using recombinant SARS-CoV-2 proteases. Next, we confirm their ability to detect viral protease expression during infection with authentic virus. Finally, we generate reporter cells stably expressing an optimised luciferase-based biosensor, enabling viral replication to be measured within 24 h in a 96- or 384-well plate format, including variants of concern. We have therefore developed a luminescent SARS-CoV-2 reporter cell line, and demonstrated its utility for quantitation of infectious virus and neutralising antibodies.

Introduction

Severe respiratory syndrome coronavirus 2 (SARS-CoV-2) infection or vaccination with spike protein or mRNA are able to induce neutralising antibodies and prevent coronavirus disease 2019 (COVID-19). Recently, SARS-CoV-2 variants have emerged with the ability to escape antibody neutralisation *in vitro*, and protective immunity *in vivo* [1]. Defining serological correlates of protection against these variants has been complicated by variability in the assays used to measure neutralising antibody titres [2]. Plaque reduction neutralisation tests (PRNT) remain the gold standard, but their utility for large-scale screening of sera is limited by low throughput and long turnaround times limit. Fluorescent [3] and luminescent [4] SARS-CoV-2 reporter viruses have been described, but the reverse genetic modification of SARS-CoV-2 is laborious, and these approaches are restricted to re-engineered reference strains. Lentiviral particles pseudotyped with SARS-CoV-2 spike proteins are therefore commonly

used as surrogates, but these assays are not standardised between laboratories, and

correlate imperfectly with results for wildtype virus [5].

There is therefore an urgent unmet need for a simple, high-throughput method to quantitate

replication and neutralisation of authentic SARS-CoV-2 isolates, including new variants of

concern. Cell-based optical reporter systems have been instrumental in the response to a

previous global pandemic, caused by the human immunodeficiency virus-1 (HIV-1). In

particular, luciferase-based neutralisation assays using TZM-bl reporter cells are employed

worldwide for the assessment of vaccine-elicited neutralising antibodies, monoclonal

antibodies and the neutralising antibody response to HIV-1 infection [6, 7]. Cited by >1000

publications, they have also been instrumental in the development of antiviral therapeutics,

and facilitated a wealth of fundamental research on retrovirology. In this study, we aim to

develop an equivalent luminescent reporter cell line for SARS-CoV-2, by exploiting the

expression of viral protease activity for the detection and quantitation of infected cells.

Results

Detection of recombinant SARS-CoV-2 protease activity using FlipGFP-based reporters

During the SARS-CoV-2 replication cycle, polyproteins pp1a and pp1ab are processed into

16 non-structural proteins (nsp1 to nsp16) by the action of two virally-encoded proteases on

sequence-specific cleavage sites: Papain-like Protease (PLPro, or nsp3) and Main or 3C-like

Protease (MPro, or nsp5) [8]. The "flip" GFP (FlipGFP) reporter comprises an inactive, split

eGFP molecule, in which cleavage of an oligopeptide linker is able to restore GFP

fluorescence [9]. To test the feasibility of protease-activatable biosensors for SARS-CoV-2,

we therefore first replaced a tobacco etch virus (TEV) protease cleavage site present in the

FlipGFP oligopeptide linker with candidate MPro (WT3c and Opt3c) or PLPro (PLP1-3)

cleavage sites (**Fig 1A**), focussing on sequences conserved across SARS-CoV-2 viral isolates

(**S1A Fig**).

Candidate biosensors were co-transfected with blue fluorescent protein (BFP) plus/minus

cognate SARS-CoV-2 protease into HEK293T cells, and analysed by flow cytometry after 24

h (**Fig 1B-D**). To control for background FlipGFP fluorescence, mCherry was co-expressed

with FlipGFP using a T2A peptide linker, and the ratio of FlipGFP/mCherry fluorescence in

BFP+ (transfected) cells used to quantitate reporter activation (**S2 Fig**). All reporters showed

an increase in FlipGFP fluorescence, but the magnitude of effect was variable (**Fig 1B**). We

therefore selected the Opt3c-FlipGFP (**Fig 1C**, MPro) and PLP2-FlipGFP (**Fig 1D**, PLPro)

biosensors for further evaluation. Similar results were obtained using epifluorescence

microscopy (**Fig 1E**).

To confirm the specificity of these biosensors for their cognate protease, the same co-

transfection assay was used to measure Opt3c-FlipGFP, PLP2-FlipGFP and TEV-FlipGFP

activation in pair-wise combination with MPro, PLPro or TEV protease. In each case, an

increase in FlipGFP fluorescence was observed only in the presence of cognate protease (**Fig**

1F). As a further check, we generated non-cleavable mutants of Opt3c-FlipGFP and PLP2-

FlipGFP, by changing key residues in their respective cleavage sites. Neither non-cleavable

control showed an increase in FlipGFP fluorescence in the presence of cognate protease (**Fig**

1G). Taken together, these results confirm that the selected SARS-CoV-2 biosensors are

activated in a strictly protease and sequence-dependent manner.

Activation of FlipGFP-based reporters by SARS-CoV-2 replication

To test whether Opt3c-FlipGFP and PLP2-FlipGFP could be activated by viral protease

expression during SARS-CoV-2 infection, we made use of a permissive HEK293T cell line

over-expressing ACE2 and furin [10]. These HEK293T-ACE2 cells are readily transfectable

with reporter constructs, and form large syncytia within 24 h of SARS-CoV-2 infection (**S3A**

Fig). Since syncytia are lost during flow cytometric analysis (**S3B-C Fig**), we instead used

fluorescent microscopy to analyse changes in FlipGFP fluorescence. HEK293T-ACE2 cells

were transfected with reporter constructs, then infected after 12 h with SARS-CoV-2. After a

further 24 h incubation, reporter activation was quantitated as the ratio of FlipGFP/mCherry

fluorescence in spike-positive syncytia, compared with uninfected cells. As expected, a

consistent increase in FlipGFP fluorescence was observed in spike-positive syncytia

transfected with either Opt3c-FlipGFP or PLP2-FlipGFP, but not their corresponding non-

cleavable controls (**Fig 2A-B**).

These fluorescent biosensors therefore provide proof-of-concept data confirming that viral

protease activity may be used to signal SARS-CoV-2 infection. Nonetheless, the magnitude

of effect is markedly reduced compared with over-expression of recombinant viral proteases

(compare **Fig 2B** with **Fig 1G**). This likely reflects lower levels of protease expression during

viral infection (**Fig 2C**). In addition, it is possible that the localisation of proteases and/or

presence of other viral proteins and endogenous polyprotein substrates during authentic viral

infection reduces their likelihood of encountering reporter molecules.

Luciferase-based biosensors of authentic SARS-CoV-2 infection

To enzymatically amplify the signal from viral protease activity, we next sought to generate

equivalent luciferase-based biosensors. The 30F-GloSensor reporter comprises an inactive,

circularly permuted firefly luciferase (FFluc) molecule, in which cleavage of a specific

oligopeptide linker is able to restore luminescence [11]. Co-expression with Renilla luciferase

(Rluc) from the same vector allows reporter levels and cell viability to be normalised between

conditions. We therefore generated 30F-GloSensor constructs with the same sequence-

specific oligopeptide linkers as our FlipGFP-based reporters, together with the equivalent non-

cleavable controls. These constructs were co-transfected with/without cognate SARS-CoV-2

protease into HEK293T cells, and luminescence analysed after 24 h (**Fig 3A**). Normalising for

Rluc luminescence, a 139-fold increase in luminescence was observed for the 30F-Opt3c

biosensor (MPro), and a 74-fold increase for the 30F-PLP2 biosensor (PLPro) – much greater

than the changes in fluorescence with FlipGFP-based reporters (compare **Fig 3A** with **Fig**

1G).

To test whether 30F-Opt3c and 30F-PLP2 could also be efficiently activated by viral protease

expression during SARS-CoV-2 infection, HEK293T-ACE2 cells were again transfected with

reporter constructs, then infected after 12 h with SARS-CoV-2. After a further 24 h incubation,

reporter activation was quantitated as the ratio of FFluc/Rluc luminescence. Compared with

uninfected cells, a marked increase in luminescence was observed in infected cells

transfected with either 30F-Opt3 or 30F-PLP2, but not non-cleavable controls (**Fig 3B**). In the

context of viral infection, 30F-PLP2 showed a better dynamic range (29-fold increase in

luminescence) than 30F-Opt3c (12-fold increase in luminescence). We therefore selected

30F-PLP2 for further validation in assays of SARS-CoV-2 replication.

First, we titrated a SARS-CoV-2 viral stock in HEK293T-ACE2 cells transfected with 30F-

PLP2, and compared the luciferase signal with automated microscopy for spike. As expected,

the ratio of FFluc/Rluc luminescence increased in accordance with the amount of virus, and

correlated closely with the proportion of spike-positive (infected) cells (**Fig 3C-E**). Next,

HEK293T-ACE2 cells transfected with 30F-PLP2 were infected with SARS-CoV-2 in the

presence of remdesivir, GC376, favipiravir, lopinavir or DMSO (**Fig 3F-G**). Similar to published

data [12-14], these assays readily distinguished antiviral compounds abrogating SARS-COV-

2 replication *in vitro* (remdesivir and GC376) from those that did not (favipiravir and lopinavir).

IC50s for remdesivir and GC376 were in the low nM range, corresponding with previous

results from PRNTs [12, 13, 15]. Importantly, since the luminescent signal was inhibited by

remdesivir and GC376 – neither of which targets PLPro directly – biosensor activation must

genuinely reflect viral replication.

Generation of a luminescent SARS-CoV-2 reporter cell line for quantitation of infectious

virus and neutralising antibodies

To simplify our assay and adapt it for large-scale serological screening, we optimised the 30F-

PLP2/Rluc reporter cassette for lentiviral transduction (**S4 Fig**), and used it to generate a

stable cell line expressing the 30F-PLP2 biosensor (**Fig 4A**). When these HEK293T-ACE2-

30F-PLP2 cells were infected with wild-type SARS-CoV-2, a 20-fold increase in luminescence

was seen at 24 h (**Fig 4B**). To further increase the dynamic range of the assay, we screened

single-cell clones obtained from the bulk population of reporter cells. Clones B7 and G7

showed the maximum response to SARS-CoV-2 infection (>100-fold increase in FFluc/Rluc

ratio at 24 h), and clone B7 was selected for the remaining experiments shown in this paper

(**Fig 4C**). Assays using these cells are readily conducted in a 96-well or 384-well plate format

(**Fig 4D**), and – as expected based on PLP2 sequence conservation (**S1 Fig A**) – they may

be activated by different isolates of wildtype SARS-CoV-2, including the B.1.1.7 variant of

concern (**Fig 4E**).

Finally, to demonstrate the utility of our reporter cell line for measuring SARS-CoV-2

neutralising activity, we tested serum samples from 5 healthy control donors 21 days after

their first or second doses of Pfizer-BioNTech BNT162b2 mRNA vaccine (**Fig 5A-B**).

Consistent with previous results [16], whilst anti-spike antibodies were readily detectable by

immunoassay in all individuals (**Fig 5C**), sigmoidal neutralisation curves were observed in only

3/5 donors after their first dose (**Fig 5A**, upper panels). The highest neutralising titre at 50%

inhibition (NT50) was observed for donor 2 (NT50=5037), who also had detectable anti-

nucleocapsid antibodies, suggesting previous SARS-CoV-2 infection (**Fig 5C**, left panel).

Conversely, after the second dose, strong neutralising activity was seen for all donors tested

(NT50 range 667-2592, **Fig 5A**, lower panels, and **Fig 5B**).

Discussion

Taken together, our data establish the feasibility of protease-activatable biosensors for the

quantitation of authentic SARS-CoV-2 replication, and demonstrate the practical utility of a

luciferase-based reporter cell line for the measurement of neutralising antibody activity in

clinical samples. As well as serological surveys, they are ideally suited for both general

virological assays, and screens of candidate antiviral compounds. Luminescent assays may

be readily adapted to high-throughput platforms and, whilst the mechanism of reporter

activation is different, the range of potential applications is similar to TZM-bl reporter cells for
HIV infection.

Prior to the COVID-19 pandemic, protease-activatable reporters had been used successfully
for the detection of recombinant proteases from a range of viruses including HIV-1, HCV,
DENV-1, and MERS-CoV [17-20], but not endogenous viral proteases expressed during
authentic infection. Whilst other groups have also described protease-activatable reporters for
SARS-CoV-2 MPro [21-24], the primary aim of this study was to develop a simple, optical
biosensor for authentic SARS-CoV-2 viral infection. Consistent with this, our luminescent
reporter cell line allows assays to be conducted in 24 h, in a 96-well or 384-well plate format,
without the need for sophisticated high-content microscopy.

Compared with FlipGFP biosensors, the window for luciferase-based biosensor activation was
greatly increased. In addition, whilst the 30F-Opt3/MPro biosensor was readily activated by
recombinant protease expression, the 30F-PLP2/PLPro biosensor performed markedly better
during SARS-CoV-2 infection. This may reflect inversion of the relative expression levels of
MPro and PLPro, or differential accessibility of the reporters (substrates) to viral proteases in
the context of infection. Since PLPro of SARS-CoV has deubiquitylation (DUB) activity, and is
able to regulate whole cell levels of ubiquitylation during viral replication [25], we suspect that
PLPro of SARS-CoV-2 may also be able to readily access a wide variety of cellular proteins,
including cytosolic reporters.

Unlike reverse-engineered fluorescent or luminescent reporter viruses, our reporter cells are
able to detect different clinical SARS-CoV-2 isolates without modification of the viral genome,
including emerging variants of concern. Beyond SARS-CoV-2, the PLP2 cleavage sequence
is highly conserved in SARS-CoV, and partially conserved in MERS-CoV (**S1A Fig**). This
suggests that the Papain-like proteases of these viruses will also be able to activate 30F-
PLP2, and we confirmed this in the case of recombinant SARS-CoV PLPro (**S1B Fig**). Our
protease-activatable luminescent biosensor and reporter cell line therefore provide an "off the

shelf" solution for the quantitation of authentic betacoronavirus replication and neutralising

antibody activity during this and (potentially) future coronavirus pandemics.

Materials and methods

Reagents

The following chemical compounds were used in this study: lopinavir (APExBIO, A8204);

GC376 (a kind gift from Wayne Vuong, John C. Vederas and M. Joanne Lemieux); remdesivir

(APExBIO, B8398); favipiravir (BioVision, 2778-5, a kind gift from Aartjan te Velthuis).

The following commercial assays and kits were used in this study: Dual-Glo Luciferase Assay

System (Promega, E2920); NEBuilder HiFi DNA Assembly Cloning Kit (NEB, E5520S); 5-

alpha Competent E. coli (NEB, C2987); Micro BCA Protein Assay Kit (Thermo Fisher, 23235).

The following antibodies were used in this study: anti-NSP3 (743 - 1072) SARS-CoV-2 (MRC

PPU reagents, sheep DA126, 1st bleed, for immunoblot, 1:500); anti-NSP5 SARS-CoV-2

(MRC PPU reagents, sheep DA118, 2nd bleed, for immunoblot, 1:500); anti-SARS-CoV-2

spike [1A9] (Genetex, GTX632604, for microscopy and flow cytometry, 1:500); anti-SARS-

CoV-2 nucleocapsid (Novus Biologicals, NB100-56683, for immunoblot, 1:1,000); anti-β-actin-

HRP (Sigma, A5316, for immunoblot, 1:20,000); anti-mouse Alexa Fluor 647 (AF647)

secondary (Jackson ImmunoResearch, #715-605-150, for microscopy and flow cytometry,

1:1,000); anti-mouse Alexa Fluor 594 (AF594) secondary (Jackson Immunoresearch, #715-

585-150, for microscopy, 1:1,000).

Cell culture

HEK293T cells (a kind gift from Paul Lehner, authenticated by STR profiling [26, 27]) and

VeroE6 cells (a kind gift from Rupert Beale, authenticated by species-specific PCR (IDEXX

BioAnalytics)) were cultured in DMEM supplemented with 10% fetal calf serum (FCS), 100

units/ml penicillin, and 0.1 mg/ml streptomycin at 37℃ in 5% CO2. All cells were regularly

screened and confirmed to be mycoplasma negative (Lonza MycoAlert and IDEXX

BioAnalytics).

Vectors for transgene expression

<u>FlipGFP-based reporters</u>

To generate FlipGFP-based reporters, the TEV FlipGFP plasmid PCDNA3-FlipGFP(TEV

cleavage seq) T2A mCherry (Addgene, #124429, a gift from Xiaokun Shu [9]) was used as a

template for pairs of PCR reactions including primers designed to generate overlapping

products replacing the TEV cleavage site with the indicated cleavage sequence (**S5 Table**).

For example, to replace the TEV cleavage site with the PLP2 cleavage sequence, products

generated by PCR reactions including 'Near AfeI'/'PLP2 Rv' and 'PLP2 Fw'/'Near AflIII' primer

pairs were used. The same plasmid was then digested with AfeI and AflIII and assembled with

the gel purified PCR products using the HiFi Assembly Master Mix (NEB, E5520).

For MPro, we selected a wildtype self-cleavage sequence (SAVLQ/SGF, herein termed WT3c)

present between nsp4 and nsp5, and an "optimal" cleavage sequence (TVRLQ/SGF, herein

termed Opt3c) found by substrate profiling of SARS-CoV MPro [28]. For PLPro, we selected

all three cognate cleavage sequences present in the pp1a polyprotein (ELNGG/AYT, herein

termed PLP1; TLKGG/APT, herein termed PLP2; and ALKGG/KIV, herein termed PLP3).

To generate a non-cleavable Opt3c-FlipGFP (MPro) reporter, the critical glutamine residue in

the cleavage site was changed to isoleucine (TVRLI/SGF). For the PLP2-FlipGFP (PLPro)

reporter, the critical LKGG sequence was scrambled to GLGK (TGLGK/APT).

<u>Luciferase-based reporters</u>

To generate circularly permuted firefly luciferase (FFluc)-based reporters, the pGloSensor-

30F plasmid (Promega; discontinued, but available on request) was digested with BamHI and

HindIII, then assembled (as above) with oligonucleotides encoding the indicated cleavage

sequences (**S6 Table**).

To generate a lentiviral expression vector for the 30F-PLP2 reporter (**S4 Fig**), the pCMV-

intron-30F-PLP2-pA-pSV40-hRluc reporter cassette was first amplified from the pGloSensor-

30F PLP2, then assembled (as above) with pHRSIN-pCMV-EGFP-PGK-Puro [29] digested

with EcoRI and NotI. Next, the SV40 poly(A) signal and promoter were replaced with an

internal ribosome entry site (IRES) to generate the final pHRSIN-30F-PLP2-IRES-hRluc-

WPRE-PGK-Puro lentiviral expression vector. In brief, the pA-pSV40-hRluc fragment was

excised by digestion with SalI, and replaced by three-way HiFi assembly (as above) with

Renilla luciferase amplified by PCR from pGloSensor-30F and a 586 bp encephalomyocarditis

virus (EMCV) IRES. The final sequence of the 30F-PLP2-IRES-hRluc-WPRE-PGK-Puro

reporter cassette is shown (**S7 Appendix**).

<u>Viral proteases</u>

To generate an expression vector for SARS-CoV-2 Main protease (MPro), the open reading

frame was amplified by PCR from the pDONR223 SARS-CoV-2 NSP5 plasmid (Addgene,

#141259, a gift from Fritz Roth [30]), including an ATG start codon in the forward primer and

a TAA stop codon in the reverse primer (MPro_Fw: GCTTGGTACCGAGCTCGCACATGT

CCGGATTCCGCAAGATG; MPro_Rv: GTGATGGATATCTGCAGTTACTGGAAGGTCACAC

CAGAGC), then assembled (as above) with EcoRI-linearised pcDNA3.1.

To generate an expression vector for the SARS-CoV-2 Papain-like protease catalytic domain

(PLPro c.d.), we used the SARS-CoV nsp3 catalytic domain as a guide [31]. The coding

sequence from amino acid 747E to 1061K was amplified by PCR from the pDONR207 SARS-

CoV-2 NSP3 plasmid (Addgene, #141257, a gift from Fritz Roth[30]), including a Kozak

sequence and ATG start codon in the forward primer and a TAA stop codon in the reverse

primer   (PLPcd_Fw: GCTTGGTACCGAGCTCGGCCACCATGGAGGTGAGGACCATCAA

GGTGTT; PLPcd_Rv: GTGTGATGGATATCTGCAGTTACTTGATGGTGGTGGTGTAGCT),

then assembled (as above) with EcoRI-linearised pcDNA3.1. Where indicated, empty
pcDNA3.1 was used as a control.

For expression of TEV protease, pcDNA3.1 TEV (full-length) (Addgene, #64276, a gift from
Xiaokun Shu [32]) was used.

To generate an expression vector for the SARS-CoV Papain-like protease catalytic domain
(PLPro c.d.) [31], a gene block was synthesized by GenScript (**S8 Appendix**), then assembled
(as above) with pcDNA3.1 digested with BamHI and EcoRI.

<u>Other transgenes</u>

For expression of BFP, pTAG-BFP-N (Evrogen, FP172) was used.

To generate a lentiviral expression vector for human ACE2, the open reading frame was
amplified from a HepG2 cDNA library by PCR using the indicated primers (ACE2_Fw:
CGCCCGGGGGGGATCCACTAGGTACCATGTCAAGCTCTTCCTGGCTCC;   ACE2_Rv:
CTAGAGTCGCGGCCGCTCTACTCGAGCTAAAAGGAGGTCTGAACATCATCAGTGTTTT
G), then assembled (as above) with pHRSIN-pSFFV-GFP-PGK-Hygro [33] digested with KpnI
and XhoI to generate pHRSIN-ACE2-Hygro.

For expression of human furin, pHRSIN-Furin-Puro (a kind gift from Paul Lehner) was used.

For lentiviral packaging, psPAX2 (Addgene, #12260, a gift from Didier Trono) and pCMV-VSV-
G (Addgene, #8454, a gift from Bob Weinberg [34]) were used.

All constructs generated for this study were verified by Sanger sequencing (Source
BioScience).

Generation of HEK293T-ACE2 and HEK293T-ACE2-30F-PLP2 cells

For transduction of HEK293T cells with ACE2, a pHRSIN-ACE2-Hygro lentiviral stock was
generated by co-transfection of HEK293T cells with psPAX2 and pCMV-VSV-G using
standard methods. After selection with hygromycin for 2 weeks, cells were sorted for high cell

surface ACE2 expression and single-cell cloned. Following expansion, a clone with stable,

homogeneously high expression of ACE2 (herein termed clone 22) was selected for further

experiments. For transduction with furin, a pHRSIN-Furin-Puro lentiviral stock was generated

as above, and clone 22 cells (herein termed HEK293T-ACE2 cells) were selected with

puromycin for a further 2 weeks. Finally, to generate HEK293T-ACE2-30F-PLP2 reporter

cells, HEK293T-ACE2 cells were further transduced with a pHRSIN-30F-PLP2-IRES-hRluc-

WPRE-PGK-Puro lentiviral stock at amultiplicity of infection (MOI) of 3, generated as above.

100 single-cell clones were obtained by limiting dilution, then screened for sensitivity to SARS-

CoV-2 infection (maximum increase in ratio of FFluc/Rluc luminescence).

Production and titration of SARS-CoV-2 viral stocks

Unless otherwise stated, the virus used in this study was the clinical isolate SARS-CoV-

2/human/Liverpool/REMRQ0001/2020, a kind gift from Ian Goodfellow (University of

Cambridge), isolated by Lance Turtle (University of Liverpool) and David Matthews and

Andrew Davidson (University of Bristol) [35, 36]. Where indicated, the B.1.1.7 viral isolate

SARS CoV-2 England/ATACCC 174/2020 was used, a kind gift from Greg Towers [37].

Viral stocks were prepared by passaging once in VeroE6 cells. In brief, cells were infected at

a low multiplicity of infection (MOI) with the original viral stock, and incubated for 72 h (by

which time cytopathic effect was evident). Virus-containing culture supernatants were then

clarified by centrifugation at 600 g for 5 min, and immediately frozen in aliquots at -80°C. Viral

stocks were then titrated in VeroE6 cells by 50% tissue culture infectious dose (TCID50)

assays using standard methods.

Flow cytometric analysis of FlipGFP-based reporters in the presence of recombinant

proteases

HEK293T cells were seeded at least 12 h in advance and transfected at approximately 50%

confluency. Plasmids and TransIT-293 transfection reagent (Mirus) were mixed at a ratio of 1

µg DNA : 3 µLTransit-T293 in Opti-MEM (Gibco). For an experiment conducted in triplicate at

a 48-well scale, we typically used (per condition): 150 ng BFP expression vector, 300 ng

FlipGFP-based reporter construct, and 450 ng of either empty pcDNA3.1 (control) or

pcDNA3.1 encoding the indicated protease; 2.7 µL TransIT-293; and 150 µL Opti-MEM (for

50 µL of transfection/well).

For titration of SARS-CoV-2 proteases, the total amount of DNA transfected was kept constant

by adding additional empty pcDNA3.1. Where indicated, DMSO or candidate inhibitors of

recombinant MPro or PLPro were added immediately after transfection.

Unless otherwise indicated, HEK293T cells were dissociated with trypsin 24 h post-

transfection, resuspended in PBS + 0.5% FBS and analysed immediately by flow cytometry

using a BD LSRFortessa equipped with 405 nm, 488 nm, 561 nm and 640 nm lasers. The

ratio of FlipGFP/mCherry mean fluorescence intensity (MFI) in BFP+ (transfected) cells was

used to quantitate reporter activation (FlowJo 10.7). An indicative gating strategy is shown in

S2 Fig.

Flow cytometric analysis of spike protein expression in SARS-CoV-2-infected cells

HEK293T-ACE2 cells were seeded at a density of 9×10^4 cells/48-well in 250 µL complete

media. The following morning, cells were infected with SARS-CoV-2 at MOI=1 and incubated

for 24 h.

To measure the fraction of infected cells using flow cytometry, cells were first dissociated with

trypsin and fixed for 15 min by incubation in 4% paraformaldehyde (PFA). Cells were then

permeabilised with Perm/Wash buffer (BD), stained for SARS-CoV-2 spike protein using a

mouse monoclonal antibody (GeneTex, GTX632604) for 30 min at room temperature, washed

twice, stained with an anti-mouse Alexa Fluor 647 (AF647) secondary antibody (Jackson

ImmunoResearch, #715-605-150) for 30 min at room temperature, washed twice,

resuspended in PBS + 0.5% FBS and analysed by flow cytometry as above.

Automated microscopic analysis of spike protein expression in SARS-CoV-2-infected

cells

HEK293T-ACE2 cells were seeded at a density 9 x 10^4 cells/well of an 8-well μ-Slide (Ibidi,

80826) in 250 μL complete media. The following morning, cells were infected with SARS-CoV-

2 at MOI=1 and incubated for 24 h.

To measure the fraction of infected cells using automated microscopy, cells were first fixed for

15 min by incubation in 4% PFA. Cells were then permeabilised with Perm/Wash buffer (BD),

stained for SARS-CoV-2 spike protein using a mouse monoclonal antibody (GeneTex,

GTX632604) for 30 min at room temperature, washed twice, stained with an anti-mouse Alexa

Fluor 594 (AF594) secondary antibody (Jackson ImmunoResearch, #715-585-150) for 30 min

at room temperature, washed extensively, mounted with 200 μL/well of Fluoroshield Mounting

Media (Sigma, F6057), and analysed by automated microscopy.

In brief, images were acquired using a Cellomics ArrayScan XTI high-throughput imaging

platform (Thermo Fisher) using a 386 nm excitation/emission filter to detect DAPI-stained

nuclei and a 560 nm excitation/emission filter to detect AF594. Images were then analysed

with built-in high content HCS Studio software (by Thermo Fisher) using the Target Activation

application. For this, cellular objects were identified by applying overlays (masks) based on

DAPI intensity. Necessary steps to exclude non-cellular artefacts (large or small objects) were

activated based on average nuclei size. Additionally, background correction was performed

on both channels. The generated nuclei masks were then applied to the AF594 channel and

the threshold for AF594 staining was determined using stained mock-infected cells. Finally,

cells were considered infected if their AF594 signal was above this threshold. 42 fields were

scanned for each sample/condition to ensure the analysis of a sufficient number of cells.

Confocal microscopic analysis of FlipGFP-based reporters and spike protein

expression in SARS-CoV-2-infected cells

HEK293T-ACE2 cells were seeded at a density 9 x 10^4 cells/well of an 8-well μ-Slide (Ibidi,

80826) in 250 μL complete media. After 1 h, cells were transfected in duplicate with the

indicated FlipGFP-based reporter constructs as above. The following morning, cells were

infected with SARS-CoV-2 at MOI=1 and incubated for 24 h.

To measure reporter activation in infected cells using confocal microscopy, cells were first

fixed for 15 min by incubation in 4% PFA. Cells were then permeabilised with Perm/Wash

buffer (BD), stained for SARS-CoV-2 spike protein using a mouse monoclonal antibody

(GeneTex, GTX632604) for 30 min at room temperature, washed twice, stained with an anti-

mouse AF647 secondary antibody (Jackson ImmunoResearch, #715-605-150) for 30 min at

room temperature, mounted with 200 µL/well of Fluoroshield Mounting Media (Sigma, F6057),

and analysed by confocal microscopy using a Zeiss LSM 710 Inverted confocal microscope

equipped with 405, 458, 543 and 633 nm lasers and a Plan Apochromat 63X/1.40 Oil DIC

M27 objective. For each reporter construct, the ratio of FlipGFP/mCherry MFI was calculated

manually using Fiji (ImageJ), by creating a mask around syncytiated cells that were both

spike+ (infected) and mCherry+ (transfected).

Immunoblotting

Washed cell pellets were lysed in PBS + 1% Triton supplemented with Halt Protease and

Phosphatase Inhibitor Cocktail (Thermo Scientific) for 30 min on wet ice. Post-nuclear

supernatants were heated in Laemelli Loading Buffer for 5 min at 95℃, separated by SDS-

PAGE and transferred to Immobilon-P membrane (Millipore). Membranes were blocked in

PBS/5% non-fat dried milk (Marvel)/0.2% Tween and probed with the indicated primary

antibody overnight at 4℃. Reactive bands were visualised using HRP-conjugated secondary

antibodies and SuperSignal West Pico or Dura chemiluminescent substrates (Thermo

Scientific). Typically, 10–20 µg total protein was loaded per lane.

Luminescent analysis of luciferase-based reporters in the presence of recombinant

proteases

HEK293T cells were transfected in triplicate at a 48-well scale essentially as for transfection

of FlipGFP-based reporter constructs, typically using 150 ng of the indicated pGloSensor-30F

reporter construct and 150 ng of empty pcDNA3.1 (control) or pcDNA3.1 encoding the

indicated protease.

After 24 h incubation, media was aspirated from each well, and cells lysed with 50 µL/well

Dual-Glo Luciferase Buffer (Promega) diluted 1:1 with PBS + 1% NP-40, for 10 min at room

temperature. Lysates were then transferred to opaque half-area 96-well plates, and reporter

activation quantitated as the ratio of firefly luciferase (FFluc)/Renilla luciferase (Rluc) activity

measured using the Dual-Glo kit (Promega) according to the manufacturer's instructions. In

brief, FFluc activity was first measured using a ClarioStar microplate reader. 25 µL Stop and

Glo Buffer and Substrate (Promega) was then added to each well. After incubation for 10 min

at room temperature, Rluc activity was measured using the same ClarioStar microplate reader

and the ratio of FFluc/Rluc activity calculated for each condition.

Luminescent analysis of luciferase-based reporters in SARS-CoV-2-infected cells

HEK293T-ACE2 cells were reverse-transfected with plasmids and TransIT-293 at a ratio of 1

µg DNA : 3 µLTransit-T293 in Opti-MEM (Gibco). For an experiment conducted in triplicate at

a 48-well scale, we typically used (per condition): 900 ng pGloSensor-30F reporter construct;

2.7 µL TransIT-293; and 150 µL Opti-MEM (for 50 µL of transfection mix/well). 2.7×10^5 cells

were dissociated with Accutase, combined with the transfection mix, and seeded at 9×10^4

cells/well.

The following morning, cells were infected with SARS-CoV-2 at MOI=0.01 (or for the titration

experiments, with the indicated volume of viral stock) and incubated for 24 h. Media was

aspirated from each well, and cells lysed with 50 µL/well of Dual-Glo Luciferase Buffer

(Promega) diluted 1:1 with PBS + 1% NP-40 for 10 min at room temperature. Lysates were

then transferred to opaque half-area 96-well plates, and reporter activation quantitated as the

ratio of FFluc/Rluc activity as above. Where indicated, candidate antivirals were added to the

cells 1 h before infection with SARS-CoV-2.

Luminescent analysis of SARS-CoV-2-infected HEK293T-ACE2-30F-PLP2 reporter cells

Unless otherwise stated, HEK293T-ACE2-30F-PLP2 reporter cells (either the bulk population,

or indicated clone) were seeded at a density of 4×10^4 cells/96-well in 100 µL complete media.

The following morning, cells were infected with SARS-CoV-2 at MOI=0.01 and incubated for

24 h. Media was aspirated from each well, and cells lysed with 25 µL/well of Dual-Glo

Luciferase Buffer (Promega) diluted 1:1 with PBS + 1% NP-40 for 10 min at room temperature.

Lysates were then transferred to opaque half-area 96-well plates, and reporter activation

quantitated as the ratio of FFluc/Rluc activity as above. For evaluation of candidate antivirals,

compounds were added to the cells 1 h before infection with SARS-CoV-2. To measure SARS-

CoV-2 neutralising antibody titres, SARS-CoV-2 viral stock (MOI=0.01) was pre-incubated

with a 3-fold dilution series of heat-inactivated serum for 2 h at 37°C, prior to addition to

reporter cells. Where indicated, assays were performed in 384-well plates, with cell numbers

and reagent volumes scaled according to surface area, and reporter activation quantitated

directly as FFluc activity.

Serological assessment

Serum samples from healthy control donors were obtained 21 days after their 1st or 2nd dose

of BNT162b2 mRNA vaccine (Pfizer-BioNTech), heat-inactivated at 56°C for 30 min, then

frozen in aliquots at -80°C.

Antibodies to SARS-CoV-2 nucleocapsid, trimeric spike and spike receptor binding domain

were quantitated by Luminex bead-based immunoassay as previously described [38]. In brief,

recombinant SARS-CoV-2 proteins were covalently coupled to carboxylated bead sets

(Luminex; Netherlands) to form a multiplex assay. Coupled bead sets were incubated with

serum samples at a dilution of 1/100 for 1 h at room temperature, washed three times with

10mM PBS/0.05% Tween-20, incubated for 30 min with a PE-labeled anti–human IgG-Fc

antibody (Leinco/Biotrend), washed again, then resuspended in 100 µl PBS/Tween-20. They

were then analyzed on a Luminex analyzer (Luminex / R&D Systems) using Exponent

Software V31. Specific binding was reported as mean fluorescence intensity (MFI). Serum

samples from patients with PCR-confirmed COVID-19 were used as positive controls, and

stored serum samples collected prior to November 2019 were used as negative controls.

Diagnostic cutoffs were determined by receiver operating characteristic (ROC) curve analysis.

Ethics statement

Ethical approval for this study was granted by the East of England – Cambridge Central

Research Ethics Committee (08/H0308/176). Written informed consent was obtained from all

volunteers prior to providing serum samples.

SARS-CoV-2, SARS-CoV, and MERS-CoV sequence logos

A SARS-CoV-2 genomic alignment file was retrieved from the GISAID database

(https://www.epicov.org/). At the time of access (30/06/2020), this alignment (msa_0630)

contained 50,387 sequences. Genome sequences and alignments for MERS-CoV and SARS-

CoV were accessed via the NCBI Virus portal

(https://www.ncbi.nlm.nih.gov/labs/virus/vssi/#/virus?SeqType_s=Nucleotide). For the MERS-

CoV sequences, we searched the database for Virus with taxonid=1335626 and Host with

taxonid='human'. We set the nucleotide completeness option to 'complete' and alignments

were generated using the NCBI alignment function. Consensus sequences automatically

generated and included in the alignment file where removed. For the SARS-CoV sequences,

we searched the database for Virus with taxonid=694009 and Host taxonid= 'human'. Because

of the large number of sequences (~15,000), it was not possible to perform the alignment

using the NCBI alignment function. Instead, FASTA sequences were downloaded and aligned

using MAFFT[39] with default parameters.

Having manually inspected the genomic alignments to identify the regions of interest, we used

the 'extractalign' function in the European Molecular Biology Open Software Suite (EMBOSS)

[40]. In the case of the GISAID alignment, we removed entries that were incomplete or that

presented nucleotide ambiguities (non-CTGA bases). The resulting sets (one per region of

interest and four per alignment inspected) were conceptually translated using the 'transeq'

function in EMBOSS. The resulting amino acid sequences were used as input for the WebLogo application (http://weblogo.berkeley.edu) [41, 42].

Statistical analysis

General data manipulation was conducted using Microsoft Excel, and statistical analysis using Prism 8.0 (GraphPad). Unless otherwise stated, sample means were compared by one-way ANOVA followed by Tukey's multiple comparison test. To compare the results of luminometry and automated microscopy, linear regression and Pearson's correlation coefficient were used. To calculate half-maximal inhibitory concentrations (IC50s), FFluc/Rluc ratios were analysed using the log(inhibitor) vs. response -- Variable slope (four parameters) function. To calculate neutralising antibody titres at 50% inhibition (NT50s), FFluc/Rluc ratios were analysed using the Sigmoidal, 4PL, X is log(concentration) function.

Availability of materials

Biological materials from this study are available from specified commercial sources, or from the corresponding author on execution of an appropriate Material Transfer Agreement (MTA).

Acknowledgements

This work was supported by the MRC (CSF ref. MR/P008801/1 to NJM), NHSBT (grant ref. WPA15-02 to NJM), Addenbrooke's Charitable Trust (grant ref. to 900239 NJM), the Rosetree Trust (grant ref. G103718 to JRZ) and the NIHR Cambridge BRC. The authors thank Mark Wills for overseeing the Containment Level 3 facility, Nika Romashova for assistance with automated microscopy, Paul Lehner, James Nathan, Wayne Vuong, John C. Vederas, M. Joanne Lemieux, Aartjan te Velthuis, Rupert Beale, Ian Goodfellow, Greg Towers and Ravi Gupta for supplying key reagents, and members of the Matheson and Lehner laboratories for critical discussion.

Author contributions

PPG and NJM conceived and designed the study with input from IG, LMD, SM and AN. PPG, LMD, EJDG, SM, AN, TEM, ECH and RD performed the experiments and analysed the data. AVP conducted the bioinformatic analysis. AT-S, TWMC, JRZ, AEW and JEDT provided key resources. NJM supervised the work. PPG and NJM drafted the manuscript. NJM edited the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

Data availability

Source data from this study are available from the corresponding author on reasonable request.

References

1.   Harvey WT, Carabelli AM, Jackson B, Gupta RK, Thomson EC, Harrison EM, et al. SARS-CoV-2 variants, spike mutations and immune escape. Nat Rev Microbiol. 2021;19(7):409-24. Epub 2021/06/03. doi: 10.1038/s41579-021-00573-0. PubMed PMID: 34075212; PubMed Central PMCID: PMCPMC8167834.

2.   Altmann DM, Boyton RJ, Beale R. Immunity to SARS-CoV-2 variants of concern.

Science. 2021;371(6534):1103-4. Epub 2021/03/13. doi: 10.1126/science.abg7404. PubMed

PMID: 33707254.

3.   Xie X, Muruato A, Lokugamage KG, Narayanan K, Zhang X, Zou J, et al. An Infectious

cDNA Clone of SARS-CoV-2. Cell Host Microbe. 2020;27(5):841-8 e3. Epub 2020/04/15. doi:

10.1016/j.chom.2020.04.004.   PubMed   PMID: 32289263;   PubMed   Central   PMCID:

PMCPMC7153529.

4.   Xie X, Muruato AE, Zhang X, Lokugamage KG, Fontes-Garfias CR, Zou J, et al. A

nanoluciferase SARS-CoV-2 for rapid neutralization testing and screening of anti-infective

drugs for COVID-19. Nat Commun. 2020;11(1):5214. Epub 2020/10/17. doi: 10.1038/s41467-

020-19055-7. PubMed PMID: 33060595; PubMed Central PMCID: PMCPMC7567097.

5.   Donal TS, Adam CH, Javier G-J, Michael LK, Stephanie L, Anthony B, et al. Nature

Portfolio. 2021. doi: 10.21203/rs.3.rs-224655/v1.

6.   Sarzotti-Kelsoe M, Bailer RT, Turk E, Lin CL, Bilska M, Greene KM, et al. Optimization

and validation of the TZM-bl assay for standardized assessments of neutralizing antibodies

against   HIV-1.   J   Immunol   Methods.   2014;409:131-46.   Epub   2013/12/03.   doi:

10.1016/j.jim.2013.11.022.   PubMed   PMID: 24291345;   PubMed   Central   PMCID:

PMCPMC4040342.

7.   Wei X, Decker JM, Liu H, Zhang Z, Arani RB, Kilby JM, et al. Emergence of resistant

human   immunodeficiency   virus   type   1   in   patients   receiving   fusion   inhibitor   (T-20)

monotherapy. Antimicrob Agents Chemother. 2002;46(6):1896-905. Epub 2002/05/23. doi:

10.1128/aac.46.6.1896-1905.2002.   PubMed   PMID: 12019106;   PubMed   Central   PMCID:

PMCPMC127242.

8.   Chen Y, Liu Q, Guo D. Emerging coronaviruses: Genome structure, replication, and

pathogenesis. J Med Virol. 2020;92(4):418-23. Epub 2020/01/23. doi: 10.1002/jmv.25681.

PubMed PMID: 31967327; PubMed Central PMCID: PMCPMC7167049.

9.   Zhang Q, Schepis A, Huang H, Yang J, Ma W, Torra J, et al. Designing a Green

Fluorogenic Protease Reporter by Flipping a Beta Strand of GFP for Imaging Apoptosis in

Animals. J Am Chem Soc. 2019;141(11):4526-30. Epub 2019/03/02. doi:

10.1021/jacs.8b13042. PubMed PMID: 30821975; PubMed Central PMCID:

PMCPMC6486793.

10.   Condor Capcha JM, Lambert G, Dykxhoorn DM, Salerno AG, Hare JM, Whitt MA, et

al. Generation of SARS-CoV-2 Spike Pseudotyped Virus for Viral Entry and Neutralization

Assays: A 1-Week Protocol. Front Cardiovasc Med. 2020;7:618651. Epub 2021/02/02. doi:

10.3389/fcvm.2020.618651. PubMed PMID: 33521067; PubMed Central PMCID:

PMCPMC7843445.

11.   Wigdal SS, Anderson JL, Vidugiris GJ, Shultz J, Wood KV, Fan F. A novel

bioluminescent protease assay using engineered firefly luciferase. Curr Chem Genomics.

2008;2:16-28. Epub 2008/01/01. doi: 10.2174/1875397300802010016. PubMed PMID:

20161840; PubMed Central PMCID: PMCPMC2803436.

12.   Ma C, Sacco MD, Hurst B, Townsend JA, Hu Y, Szeto T, et al. Boceprevir, GC-376,

and calpain inhibitors II, XII inhibit SARS-CoV-2 viral replication by targeting the viral main

protease. Cell Res. 2020;30(8):678-92. Epub 2020/06/17. doi: 10.1038/s41422-020-0356-z.

PubMed PMID: 32541865; PubMed Central PMCID: PMCPMC7294525.

13.   Vuong W, Khan MB, Fischer C, Arutyunova E, Lamer T, Shields J, et al. Feline

coronavirus drug inhibits the main protease of SARS-CoV-2 and blocks virus replication. Nat

Commun. 2020;11(1):4282. Epub 2020/08/29. doi: 10.1038/s41467-020-18096-2. PubMed

PMID: 32855413.

14.   Simonis A, Theobald SJ, Fatkenheuer G, Rybniker J, Malin JJ. A comparative analysis

of remdesivir and other repurposed antivirals against SARS-CoV-2. EMBO Mol Med.

2021;13(1):e13105. Epub 2020/10/06. doi: 10.15252/emmm.202013105. PubMed PMID:

33015938; PubMed Central PMCID: PMCPMC7646058.

15.   Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, et al. Remdesivir and chloroquine

effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. Cell Res.

2020;30(3):269-71. Epub 2020/02/06. doi: 10.1038/s41422-020-0282-0. PubMed PMID:

32020029; PubMed Central PMCID: PMCPMC7054408.

16.   Sahin U, Muik A, Vogler I, Derhovanessian E, Kranz LM, Vormehr M, et al. BNT162b2

vaccine induces neutralizing antibodies and poly-specific T cells in humans. Nature. 2021.

Epub 2021/05/28. doi: 10.1038/s41586-021-03653-6. PubMed PMID: 34044428.

17.   Arias-Arias JL, MacPherson DJ, Hill ME, Hardy JA, Mora-Rodriguez R. A fluorescence-

activatable reporter of flavivirus NS2B-NS3 protease activity enables live imaging of infection

in single cells and viral plaques. J Biol Chem. 2020;295(8):2212-26. Epub 2020/01/11. doi:

10.1074/jbc.RA119.011319.   PubMed   PMID: 31919100;   PubMed   Central   PMCID:

PMCPMC7039541.

18.   Lindsten K, Uhlikova T, Konvalinka J, Masucci MG, Dantuma NP. Cell-based

fluorescence assay for human immunodeficiency virus type 1 protease activity. Antimicrob

Agents Chemother. 2001;45(9):2616-22. Epub 2001/08/15. doi: 10.1128/aac.45.9.2616-

2622.2001. PubMed PMID: 11502538; PubMed Central PMCID: PMCPMC90701.

19.   Sabariegos R, Picazo F, Domingo B, Franco S, Martinez MA, Llopis J. Fluorescence

resonance energy transfer-based assay for characterization of hepatitis C virus NS3-4A

protease activity in live cells. Antimicrob Agents Chemother. 2009;53(2):728-34. Epub

2008/12/10. doi: 10.1128/AAC.01029-08. PubMed PMID: 19064890; PubMed Central PMCID:

PMCPMC2630637.

20.   Kilianski A, Mielech AM, Deng X, Baker SC. Assessing activity and inhibition of Middle

East respiratory syndrome coronavirus papain-like and 3C-like proteases using luciferase-

based biosensors. J Virol. 2013;87(21):11955-62. Epub 2013/08/30. doi: 10.1128/JVI.02105-

13. PubMed PMID: 23986593; PubMed Central PMCID: PMCPMC3807373.

21.   Froggatt HM, Heaton BE, Heaton NS. Development of a Fluorescence-Based, High-

Throughput SARS-CoV-2 3CL(pro) Reporter Assay. J Virol. 2020;94(22). Epub 2020/08/28.

doi: 10.1128/JVI.01265-20.   PubMed   PMID: 32843534;   PubMed   Central   PMCID:

PMCPMC7592234.

22.   Rawson JMO, Duchon A, Nikolaitchik OA, Pathak VK, Hu WS. Development of a Cell-

Based Luciferase Complementation Assay for Identification of SARS-CoV-2 3CL(pro)

Inhibitors. Viruses. 2021;13(2). Epub 2021/01/28. doi: 10.3390/v13020173. PubMed PMID:

33498923; PubMed Central PMCID: PMCPMC7911889.

23.   O'Brien A, Chen DY, Hackbart M, Close BJ, O'Brien TE, Saeed M, et al. Detecting

SARS-CoV-2 3CLpro expression and activity using a polyclonal antiserum and a luciferase-

based biosensor. Virology. 2021;556:73-8. Epub 2021/02/07. doi: 10.1016/j.virol.2021.01.010.

PubMed PMID: 33548599; PubMed Central PMCID: PMCPMC7837110.

24.   Pahmeier F, Neufeldt CJ, Cerikan B, Prasad V, Pape C, Laketa V, et al. A Versatile

Reporter System To Monitor Virus-Infected Cells and Its Application to Dengue Virus and

SARS-CoV-2. J Virol. 2021;95(4). Epub 2020/12/02. doi: 10.1128/JVI.01715-20. PubMed

PMID: 33257477; PubMed Central PMCID: PMCPMC7851548.

25.   Yan S, Wu G. Spatial and temporal roles of SARS-CoV PL(pro) -A snapshot. FASEB

J. 2021;35(1):e21197. Epub 2020/12/29. doi: 10.1096/fj.202002271. PubMed PMID:

33368679; PubMed Central PMCID: PMCPMC7883198.

26.   Menzies SA, Volkmar N, van den Boomen DJ, Timms RT, Dickson AS, Nathan JA, et

al. The sterol-responsive RNF145 E3 ubiquitin ligase mediates the degradation of HMG-CoA

reductase together with gp78 and Hrd1. Elife. 2018;7. doi: 10.7554/eLife.40009. PubMed

PMID: 30543180; PubMed Central PMCID: PMCPMC6292692.

27.   Miles AL, Burr SP, Grice GL, Nathan JA. The vacuolar-ATPase complex and assembly

factors, TMEM199 and CCDC115, control HIF1alpha prolyl hydroxylation by regulating cellular

iron levels. Elife. 2017;6. doi: 10.7554/eLife.22693. PubMed PMID: 28296633; PubMed

Central PMCID: PMCPMC5391204.

28.   Chuck CP, Chong LT, Chen C, Chow HF, Wan DC, Wong KB. Profiling of substrate

specificity of SARS-CoV 3CL. PLoS One. 2010;5(10):e13197. Epub 2010/10/16. doi:

10.1371/journal.pone.0013197. PubMed PMID: 20949131; PubMed Central PMCID:

PMCPMC2950840.

29.   van den Boomen DJH, Sienkiewicz A, Berlin I, Jongsma MLM, van Elsland DM, Luzio

JP, et al. A trimeric Rab7 GEF controls NPC1-dependent lysosomal cholesterol export. Nat

Commun. 2020;11(1):5559. Epub 2020/11/05. doi: 10.1038/s41467-020-19032-0. PubMed

PMID: 33144569; PubMed Central PMCID: PMCPMC7642327.

30. Kim DK, Knapp JJ, Kuang D, Chawla A, Cassonnet P, Lee H, et al. A Comprehensive,

Flexible Collection of SARS-CoV-2 Coding Regions. G3 (Bethesda). 2020. Epub 2020/08/09.

doi: 10.1534/g3.120.401554. PubMed PMID: 32763951.

31. Bekes M, van der Heden van Noort GJ, Ekkebus R, Ovaa H, Huang TT, Lima CD.

Recognition of Lys48-Linked Di-ubiquitin and Deubiquitinating Activities of the SARS

Coronavirus Papain-like Protease. Mol Cell. 2016;62(4):572-85. Epub 2016/05/21. doi:

10.1016/j.molcel.2016.04.016. PubMed PMID: 27203180; PubMed Central PMCID:

PMCPMC4875570.

32. To TL, Piggott BJ, Makhijani K, Yu D, Jan YN, Shu X. Rationally designed fluorogenic

protease reporter visualizes spatiotemporal dynamics of apoptosis in vivo. Proc Natl Acad Sci

U S A. 2015;112(11):3338-43. Epub 2015/03/04. doi: 10.1073/pnas.1502857112. PubMed

PMID: 25733847; PubMed Central PMCID: PMCPMC4371907.

33. Tchasovnikarova IA, Timms RT, Matheson NJ, Wals K, Antrobus R, Gottgens B, et al.

GENE SILENCING. Epigenetic silencing by the HUSH complex mediates position-effect

variegation in human cells. Science. 2015;348(6242):1481-5. Epub 2015/05/30. doi:

10.1126/science.aaa7227. PubMed PMID: 26022416; PubMed Central PMCID:

PMCPMC4487827.

34. Stewart SA, Dykxhoorn DM, Palliser D, Mizuno H, Yu EY, An DS, et al. Lentivirus-

delivered stable gene silencing by RNAi in primary cells. RNA. 2003;9(4):493-501. Epub

2003/03/22. doi: 10.1261/rna.2192803. PubMed PMID: 12649500; PubMed Central PMCID:

PMCPMC1370415.

35. Daly JL, Simonetti B, Klein K, Chen KE, Williamson MK, Anton-Plagaro C, et al.

Neuropilin-1 is a host factor for SARS-CoV-2 infection. Science. 2020;370(6518):861-5. Epub

2020/10/22. doi: 10.1126/science.abd3072. PubMed PMID: 33082294.

36. Patterson EI, Prince T, Anderson ER, Casas-Sanchez A, Smith SL, Cansado-Utrilla C,

et al. Methods of Inactivation of SARS-CoV-2 for Downstream Biological Assays. J Infect Dis.

2020;222(9):1462-7. Epub 2020/08/17. doi: 10.1093/infdis/jiaa507. PubMed PMID: 32798217;

PubMed Central PMCID: PMCPMC7529010.

37.   Reuschl AK, Thorne L, Zuliani Alvarez L, Bouhaddou M, Obernier K, Soucheray M, et

al. Host-directed therapies against early-lineage SARS-CoV-2 retain efficacy against B.1.1.7

variant. bioRxiv. 2021. Epub 2021/01/28. doi: 10.1101/2021.01.24.427991. PubMed PMID:

33501437; PubMed Central PMCID: PMCPMC7836107.

38.   Buckland MS, Galloway JB, Fhogartaigh CN, Meredith L, Provine NM, Bloor S, et al.

Treatment of COVID-19 with remdesivir in the absence of humoral immunity: a case report.

Nature Communications. 2020;11(1):6385. doi: 10.1038/s41467-020-19761-2.

39.   Katoh K, Misawa K, Kuma Ki, Miyata T. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Research.

2002;30(14):3059-66. doi: 10.1093/nar/gkf436.

40.   Rice P, Longden I, Bleasby A. EMBOSS: the European Molecular Biology Open

Software Suite. Trends Genet. 2000;16(6):276-7. Epub 2000/05/29. doi: 10.1016/s0168-

9525(00)02024-2. PubMed PMID: 10827456.

41.   Crooks GE, Hon G, Chandonia JM, Brenner SE. WebLogo: a sequence logo

generator. Genome Res. 2004;14(6):1188-90. Epub 2004/06/03. doi: 10.1101/gr.849004.

PubMed PMID: 15173120; PubMed Central PMCID: PMCPMC419797.

42.   Schneider TD, Stephens RM. Sequence logos: a new way to display consensus

sequences. Nucleic Acids Res. 1990;18(20):6097-100. Epub 1990/10/25. doi:

10.1093/nar/18.20.6097. PubMed PMID: 2172928; PubMed Central PMCID:

PMCPMC332411.

Figure captions

Fig 1. Cell-based biosensors of SARS-CoV-2 protease activity.

(**A**) Diagram of pp1a polyprotein showing candidate SARS-CoV-2 cleavage sequences for Papain-like Protease (PLPro; PLP1-3) and Main Protease (MPro; WT3c and Opt3c). Further details on sequence selection are available in the **Materials and methods**. Sites of cleavage are highlighted in red (C-terminal side of indicated amino acid). (**B-D**) Activation of FlipGFP-based reporters by recombinant SARS-CoV-2 protease expression. HEK293T cells were co-transfected with BFP and indicated FlipGFP-based reporter constructs encoding candidate MPro or PLPro cleavage sequences ± MPro, PLPro c.d., or empty pcDNA3.1. Illustrative flow cytometry data for Opt3c-FlipGFP (**C**) and PLP2-FlipGFP (**D**) are shown. (**E**) Detection of FlipGFP-based reporter activation by epifluorescence microscopy. HEK293T cells were co-transfected with Opt3c-FlipGFP biosensor plus/minus MPro (left panel) or PLP2-FlipGFP biosensor plus/minus PLPro (right panel). FlipGFP and mCherry fluorescence were analysed by epifluorescence microscopy 24 h post-transfection. mCherry, red. FlipGFP, green. Representative of 3 independent experiments. (**F**) Protease-specificity of FlipGFP-based reporters. HEK293T cells were co-transfected with BFP and indicated FlipGFP biosensors ± MPro, PLPro c.d., TEV protease or empty pcDNA3.1. (**G**) Sequence specificity of FlipGFP-based reporters. HEK293T cells were co-transfected with BFP and indicated FlipGFP biosensors ± MPro, PLPro or empty pcDNA3.1. Further details on non-cleavable mutants are available in the **Materials and methods**. For all flow cytometry experiments (**B-D** and **F-G**), FlipGFP and mCherry fluorescence in BFP+ cells were analysed 24 h post-transfection. An indicative gating strategy is shown in **S2 Fig**. Mean values ± SEM are shown for experiments performed in triplicate, representative of at least 3 independent experiments. **** $p<0.0001$. MPro, recombinant SARS-CoV-2 Main Protease. PLPro c.d., catalytic domain of recombinant SARS-CoV-2 Papain-Like Protease. TEV, recombinant TEV protease.

Fig 2. Activation of FlipGFP-based reporters by SARS-CoV-2 replication.

(**A-B**) HEK293T-ACE2 cells were transfected with indicated FlipGFP biosensors, incubated for 12 h, then infected with SARS-CoV-2 at MOI=1. Cells were fixed, permeabilised and stained for SARS-CoV-2 spike protein 24 h post-infection. FlipGFP and mCherry fluorescence in spike+ syncytia versus spike- cells were analysed by confocal microscopy. Illustrative microscopy data from spike+ syncytia (**A**) and mean values ± SEM for at least 15 syncytia/cells from each condition (**B**) are shown, representative of 2 independent experiments. **** p<0.0001. mCherry, red. FlipGFP, green. Spike, yellow. DAPI, blue. DIC, differential interference contrast. (**C**) Expression levels of SARS-CoV-2 proteases. HEK293T-ACE2 cells were mock-infected, infected with SARS-CoV-2 at MOI=1 or transfected with MPro (left panel) or PLPro (right panel) under standard conditions. Cells were lysed after 24 h and analysed by immunoblot using antibodies specific for nsp5 (MPro), the PLPro c.d. of nsp3 or nucleocapsid. β-actin was included as a loading control. Arrows indicate presumed full-length nsp3 (215 kDa) and intermediate cleavage products present in SARS-CoV-2-infected cells. Asterisk indicates recombinant PLPro c.d.. Representative of 2 independent experiments. O.E., over-expression. MPro, recombinant SARS-CoV-2 Main Protease. PLPro c.d., catalytic domain of recombinant SARS-CoV-2 Papain-Like Protease.

Fig 3. Luminescent biosensors of authentic SARS-CoV-2 infection.

(**A**) Activation of luciferase-based reporters by recombinant SARS-CoV-2 protease expression. HEK293T cells were co-transfected with indicated 30F biosensors ± MPro, PLPro c.d., or empty pcDNA3.1. (**B**) Activation of luciferase-based reporters by SARS-CoV-2 infection. HEK293T-ACE2 cells were transfected with indicated 30F biosensors, incubated for 12 h, then mock-infected or infected with SARS-CoV-2 at MOI=0.01. (**C-E**) Quantitation of infected cells. HEK293T-ACE2 cells were transfected with the 30F-PLP2 biosensor, incubated for 12 h, then mock-infected or infected with increasing doses of SARS-CoV-2. Cells were analysed in parallel 24 h post-infection by either epifluorescence microscopy for SARS-CoV-2 spike protein (**c**), or luminometry for Firefly and Renilla luciferase activities (**D**). Spike+ cells

were enumerated by automated microscopy (Cellomics). Illustrative microscopy data (**C**) and mean values ± SEM (**D**) are shown for an experiment performed in duplicate (microscopy) or triplicate (luminometry). The correlation between the ratio of Firefly/Renilla luminescence and the proportions of spike+ cells is shown (**E**). Spike, red. DAPI, blue. DIC, differential interference contrast. R^2, Pearson's correlation coefficient. Representative of 2 independent experiments. (**F**) Inhibition of SARS-CoV-2 replication by candidate antivirals. HEK293T-ACE2 cells were transfected with the 30F-PLP2 biosensor, incubated for 12 h, then infected with SARS-CoV-2 at MOI=0.01 in the presence of DMSO or decreasing doses of candidate antivirals. Titration curves and IC50s are shown for remdesivir and GC376. Representative of 2 independent experiments. For all experiments, Firefly and Renilla luciferase activities were measured by luminometry 24 h post-transfection (**A**) or infection (**B-G**). Unless otherwise stated, mean values ± SEM are shown for experiments performed in triplicate, representative of at least 3 independent experiments. **** $p < 0.0001$. MPro, recombinant SARS-CoV-2 Main Protease. PLPro c.d., catalytic domain of recombinant SARS-CoV-2 Papain-Like Protease.

Fig 4. An optimised luminescent SARS-CoV-2 reporter cell line.

(**A**) Schematic showing activation of luminescent reporter cell line by authentic SARS-CoV-2 infection. (**B**) Activation of reporter cells by SARS-CoV-2 infection. HEK293T-ACE2-30F-PLP2 cells (bulk population) were mock-infected or infected with SARS-CoV-2 at MOI=0.01. (**C**) Selection of optimised cell lines. HEK2932-ACE2-30F-PLP2 cells (bulk population) or single-cell clones (B7 and G7) were mock-infected or infected with increasing doses of SARS-CoV-2. (**D**) Miniaturisation to 384-well plate format. Clone B7 reporter cells were infected with SARS-CoV-2 at MOI=0.01 in the presence of decreasing doses of remdesivir in either a 96-well or 384-well plate format. Titration curves and IC50s for remdesivir are shown. Representative of 2 independent experiments. (**E**) Detection of SARS-CoV-2 B.1.1.7 variant. Clone B7 reporter cells were mock-infected or infected with increasing doses of SARS-CoV-2 variant B.1.1.7. Representative of 2 independent experiments. For all experiments, Firefly (FFluc) and Renilla (Rluc) luciferase activities were measured by luminometry 24 h post-

infection. Unless otherwise stated, mean values ± SEM are shown for experiments performed

in triplicate, representative of at least 3 independent experiments. **** $p < 0.0001$. MPro,

recombinant SARS-CoV-2 Main Protease. PLPro c.d., catalytic domain of recombinant SARS-

CoV-2 Papain-Like Protease.

Fig 5. Quantitation of neutralising activity in human serum samples.

(**A-B**) Measurement of neutralising antibody titres after inoculation with BNT162b vaccine. (**A**)

Neutralisation curves and NT50s for serum samples from 5 healthy control donors 21 days

after their 1st dose (upper panels) or 2nd dose (lower panels). SARS-CoV-2 viral stock

(MOI=0.01) was pre-incubated with a 3-fold dilution series of heat-inactivated serum before

addition to clone B7 reporter cells. Firefly and Renilla luciferase activities were measured by

luminometry 24 h post-infection. Mean values ± SEM are shown for experiments performed in

triplicate, representative of 2 independent experiments. (**B**) Comparison of NT50s after 1st and

2nd doses. Samples with no detectable neutralising activity at the lowest dilution (dotted line)

are plotted at an arbitrary NT50 of 2. (**C**) Quantification of SARS-CoV-2 antibodies in serum

samples after inoculation with BNT162b vaccine. Antibodies to SARS-CoV-2 nucleocapsid,

trimeric spike and spike receptor binding domain were quantitated by Luminex bead-based

immunoassay. Negative controls, stored serum samples collected prior to 2019. Vaccinated,

serum samples from 5 healthy control donors 21 days after their 1st dose of BNT162b vaccine

(same serum samples as **A-B**). Donor 2 is highlighted in red. Dotted lines indicate diagnostic

cutoffs determined by receiver operating characteristic (ROC) curve analysis. MFI, mean

fluorescence intensity.

Supporting information

S1 Fig. Cleavage sequences of highly pathogenic human betacoronaviruses.

(**A**) Conservation of cleavage sequences. Amino acid sequence logos for cleavage sequences between nsp1/nsp2 (PLP1), nsp2/nsp3 (PLP2), nsp3/nsp4 (PLP3), and nsp4/nsp5 (WT3c) of SARS-CoV-2, SARS-CoV and MERS-CoV viruses. Relative letter heights indicate conservation across the sequences analysed. (**B**) Luminescent biosensor activation by PLPro of SARS-CoV. HEK293T cells were co-transfected with the 30F-PLP2 biosensor and the catalytic domain of recombinant Papain-Like Protease (PLPro c.d.) of either SARS-CoV-2 or SARS-CoV. Firefly and Renilla luciferase activities were measured by luminometry 24 h post-transfection. Mean fold-changes in Firefly/Renilla luminescence ± SEM in the presence or absence of protease are shown for an experiment performed in triplicate. Representative of 2 independent experiments.

S2 Fig. Indicative gating strategy for flow cytometry experiments using FlipGFP-based reporters.

Cells were typically gated on FSC-H and SSC-H, and doublets excluded using FSC-W. FlipGFP and mCherry fluorescence were analysed in BFP+ (transfected) cells. Data illustrate change in fluorescence of Opt3c-FlipGFP biosensor in the presence (lower panels) or absence (upper panels) of Main Protease (MPro).

S3 Fig. Infection of HEK293T-ACE2 cells by SARS-CoV-2.

(**A**) Cytopathic effect in HEK293T-ACE2 cells. HEK293T or HEK293T-ACE2 cells were infected with SARS-CoV-2 at MOI=1 then examined for cytopathic effect using brightfield microscopy after 24 h. Representative of >10 independent experiments. (**B-C**) Loss of spike+ HEK293T-ACE2 cells during flow cytometric analysis. HEK293T-ACE2 cells were infected with SARS-CoV-2 at MOI=1. Cells were analysed in parallel 24 h post-infection for SARS-CoV-2 spike protein by either automated microscopy (Cellomics) or flow cytometry. In each

case, the proportion of spike+ cells was measured. Illustrative data (**B**) and mean values ±

SEM (**C**) are shown for an experiment performed in triplicate. **** p<0.0001. Spike, yellow.

DAPI, blue. Representative of 2 independent experiments.

S4 Fig. 30F-PLP2 constructs.

Diagrams of 30F-PLP2 luciferase-based reporter in pGloSensor-30F and pHRSIN-30F-PLP2-

IRES-hRluc-WPRE-PGK-Puro expression vectors. 30-FF-358-544/30-FF-4-354, circularly

permuted firefly luciferase-based reporter, orange (numbers indicate positions of amino acids

in wildtype FFluc). hRluc, codon-optimised (humanised) Renilla luciferase, cyan. LTR, HIV-1

long terminal repeat, yellow. IRES, internal ribosome entry site, brown. WPRE, woodchuck

hepatitis virus post-transcriptional regulatory element (WPRE), purple. PuroR, puromycin

resistance, green.

S5 Table. PCR primers for generation of FlipGFP-based reporters.

S6. Table. Oligonucleotides for generation of luciferase-based reporters.

S7 Appendix. Complete sequence of 30F-PLP2-IRES-hRluc-WPRE-PGK-Puro reporter

cassette.

S8 Appendix. Gene block for cloning of SARS-CoV Papain-like protease catalytic

domain.

Fig 1

Fig 2

Fig 3

Fig 4

A

Vaccinee number

B

C

Fig 5

A

	PLP1	PLP2	PLP3	WT3C

SARS-CoV-2
(n= 50387)

SARS-CoV
(n= 15930)

MERS-CoV
(n = 258)

B

Fold change with/without protease

■ SARS-CoV-2 PLPro c.d.
■ SARS-CoV PLPro c.d.

S1 Fig

S2 Fig

A

HEK293T   HEK293T-ACE2

Uninfected   SARS-CoV-2   Uninfected   SARS-CoV-2

B

Cellomics

Spike   DAPI   Merge

Flow cytometry

Mock-infected   SARS-CoV-2

SSC-H

0.21%   9.63%

Spike

C

■ Cellomics
□ Flow cytometry

Apparent % Spike+ cells

****

S3 Fig

pGlo-30F-PLP2

pHRSIN-30F-PLP2-IRES-hRluc-WPRE-PGK-Puro

S4 Fig