Microglial refinement of A-fibre projections in the postnatal spinal cord dorsal horn is required for normal maturation of dynamic touch.

Yajing Xu, Stephanie C. Koch, Alexander Chamessian, Qianru He, Mayya Sundukova, Paul Heppenstall, RuRong Ji, Maria Fitzgerald, Simon Beggs*

Affiliations

Neuroscience, Physiology and Pharmacology, UCL, London, UK

Yajing Xu, Stephanie C. Koch, Maria Fitzgerald and Simon Beggs

UCL Great Ormond Street Institute of Child Health, London, UK

Simon Beggs

Duke University School of Medicine, Duke University, Durham, NC, USA

Alexander Chamessian, Qianru He, RuRong Ji

SISSA (International School for Advanced Studies), Trieste, Italy

Paul Heppenstall, Mayya Sundukova

Impact statement

Microglia phagocytose superfluous A-fibres in the superficial spinal dorsal horn during normal development, the disruption of which leads to long term aberrant dynamic touch processing and behaviour.

Abstract

Sensory systems are shaped in postnatal life by the refinement of synaptic connections. In the dorsal horn of the spinal cord, sensory circuits undergo postnatal activity dependent reorganisation, including the retraction of primary afferent A-fibres from superficial to deeper laminae which is accompanied by decreases in cutaneous sensitivity. Here we show that microglia, the resident immune cells in the CNS, phagocytose A-fibre terminals in superficial laminae in the first weeks of life. Genetic perturbation of microglial engulfment at that time prevents the normal process of A-fibre retraction, resulting in increased sensitivity of dorsal horn cells to dynamic tactile cutaneous stimulation, and behavioural hypersensitivity to dynamic touch. Thus, functional microglia are necessary for normal postnatal development of dorsal horn sensory circuits. In the absence of microglial engulfment, superfluous A-fibre projections remain in the dorsal horn and the balance of sensory connectivity is disrupted, leading to lifelong hypersensitivity to dynamic touch.

Introduction

The neonatal spinal dorsal horn differs substantially from that in adults and undergoes extensive structural and functional reorganisation over the postnatal period. One notable change is the termination zone of primary afferent A-fibres, the large myelinated afferents that encompass many low threshold cutaneous tactile afferents [1]. These afferent terminals occupy both superficial and deep laminae of the dorsal horn in neonatal rodents and gradually retract to terminate in deeper laminae III-IV by the end of the 3rd postnatal week and in adulthood [2-4]. This retraction of A-fibre terminals is accompanied by a reduction in dorsal horn cell receptive field sizes on the skin, as well as a decline in tactile sensitivity and an increase in reflex behaviour precision, with similar changes in somatosensory behaviour observed in human infants (Fitzgerald 1985, 2015; Fitzgerald et al. 1988). This suggests that structural refinement in the dorsal horn likely underlies behavioural maturation.

The process of A-fibre terminal retraction in the first weeks of life is activity dependent: Blocking neuronal input through spinal NMDAR inhibitors or increasing the noise of neuronal input through random vibration to the skin over extended period both prevented the normal retraction of A-fibres [8,9]. However, the exact mechanism underlying the retraction of A-fibre terminals is not known. Microglia cells, the major phagocytes in the CNS, have been shown to remove superfluous neurons both by driving apoptosis, removing apoptotic cells, and phagocytosing synapses and neurites during postnatal refinement [10]. To date, such studies have been largely restricted to the brain [11-15], with two studies reporting a role of microglia in the postnatal development of spinal cord ventral horn motor circuits [16,17]. Whether microglia are also involved in the maturation of somatosensory circuits in the dorsal horn is not known. Here we hypothesise that the retraction of A-fibres from

superficial laminae in the postnatal period is driven by microglia which prune A-fibre

terminals in the dorsal horn as part of normal postnatal development of spinal

sensory circuits.

Microglia undergo postnatal maturation during which they not only change in density

and morphology but also alter their transcriptional and functional identity [18,19]. Brain

microglia show particularly high expression of lysosome associated genes at P4/5

suggesting a specialised role of microglial phagocytosis during development [20]. In

addition, microglia exhibit spatial heterogeneity, as they populate different brain

regions at different rates postnatally, and express distinct local genetic profiles and

phenotype in adulthood [21–23]. So far, most research has focused on the brain and

relatively little is known about spinal cord microglia.

To test whether A-fibre terminals in the developing spinal dorsal horn are pruned by

microglial phagocytosis, we used a transgenic mouse line that expresses tdTomato

(tdT) under the *Vglut1* promoter which labels a subset of A-fibres [24] and mapped

morphological changes in dorsal horn microglia across the first two postnatal weeks

using immunofluorescence and confocal microscopy. We next tested whether normal

microglial phagocytosis was required for the A-fibre pruning. Constitutive knock-out

of the gene *Tmem16f* in microglia was shown to reduce microglial phagocytosis and

motility in the adult spinal cord in a neuropathic pain model, resulting in reduced pain

behaviour [25]. Therefore, we blocked microglial phagocytosis during the first postnatal

week using a tamoxifen inducible Cre-mediated deletion of the *Tmem16f* gene in

microglia and measured the effects on A-fibre pruning, the subsequent maturation of

dorsal horn synaptic connections, and behavioural reflex sensitivity to mechanical

skin stimulation. The results suggest that microglia prune A-fibre terminals in the

developing spinal dorsal horn and that this postnatal microglial refinement of A-fibre

terminals is required for normal somatosensory maturation.

Results

Dorsal horn microglia phagocytose A-fibre terminals during postnatal

development

In the developing spinal cord, tactile-encoding A-fibres initially project throughout the

dorsoventral extent of the dorsal horn and refine over the first few postnatal weeks to

terminate in deeper laminae, segregated from the more superficial terminals of

noxious-encoding C-fibres [8]. We confirmed this using a transgenic reporter mouse in

which tdTomato is expressed in vesicular glutamate transporter 1 (*Vglut1*)

expressing neurons (VGLUT1-tdT), a subpopulation of large myelinated sensory

neurons with features consistent with Aβ-low threshold mechanoreceptors (LTMRs)

[24]. This tdT-expression was sparse at P0 and increased with age, likely due to the

increasing developmental expression profile of *Vglut1* [26,27] (Figure 1—figure

supplement 1a). Despite this, the mice clearly revealed A-fibre terminals with flame-

shaped arbors extending into laminae I-II up to P7 (Figure 1a), which are no longer

present in these superficial laminae by P28, consistent with previous findings [8].

We hypothesised that microglia might be involved in this refinement process by

removing A-fibres in the superficial laminae via phagocytosis and determined the

phagocytic activity of dorsal horn microglia over this period using the lysosome

associated molecular marker CD68 (Figure 1b-f, Figure 1—figure supplement 1i-k).

CD68 lysosome volume increased in both laminae I-II and III-IV over the postnatal

period, reaching a peak at P10 and declining thereafter at P17 and P28 (LI-II P10 vs

P0 unpaired mean difference 117.16 [95.00% CI 92.26, 137.61], LIII-LIV unpaired

mean difference 136.02 [95.00% CI 82.89, 184.41]. It is known that the dorsal horn

undergoes developmental apoptosis around birth, extending into the first postnatal

week. Therefore, to determine whether the rise in CD68 could be contributed to

engulfment of apoptotic cells, we also quantified the number of apoptotic cells and

phagocytic cups per microglia. In contrast to the rise in CD68, the number of

phagocytic cups per microglia decreased over the first postnatal week concomitant

with a decrease in apoptotic cell counts (Figure 1—figure supplement 2), suggesting

that the increase in CD68 is not due to phagocytosis of apoptotic cells but other

material (Phagocytic cups: $F(3, 12)=10.97$, $P=0.0009$, apoptotic cells: $F(3,$

$12)=14.18$, $P=0.0003$). In parallel, microglial density and their ratio to neurons

increased over this period (Figure 1—figure supplement 1b-h). Together this

suggests that the first postnatal week represents a distinct period of microglial

activity, characterised by high levels of microglial CD68.

128
Figure 1. Spinal dorsal horn microglia engulf A-fibres during normal postnatal development
a. VGLUT1-tdTomato (red) expression in spinal laminae I-II decreases across age. Dashed white line
indicates border between lamina II and lamina III. Scale bar = 50µm.
b. Representative z-projected super-resolution image of A-fibre engulfment by microglia within the cell
body. White inset box show location of higher magnification panels in **c** and **d**. Scale bar = 5 µm.
c, d. High magnification images of microglial A-fibre engulfment in **b** stained for microglia (Iba1, cyan),
microglial lysosomes (CD68, white) and endogenously fluorescent A-fibres (VGLUT1-tdT, red). Cross-
hairs show position of the xz and yz side-view panels. Bottom panels show surface rendering of the
super-resolution image revealing pieces of tdT labelled fibres engulfed inside lysosome. Scale bar = 1
µm.

e. Microglial lysosome volume peaks at P10 and decreases thereafter for LI-II. P3 vs. P0: mean
difference 0.48 [95% CI 23.52, 24.35]; P7 vs. P0: mean difference 52.19 [95% CI 23.71, 79.47]; P10
vs. P0: mean difference 117.16 [95% CI 92.26, 137.61]; P17 vs. P0: mean difference 2.52 [95% CI -
15.07, 18.38]; P28 vs. P0: mean difference 31.84 [95% CI 11.42, 49.16]. Field of view (FOV) = 245μm
x 65μm.
f. Microglial lysosome volume peaks at P10 and decreases thereafter for LIII-IV. P3 vs. P0: mean
difference -66.67 [95% CI -121.07, -18.14]; P7 vs. P0: mean difference -19.48 [95% CI -67.54, 21.66];
P10 vs. P0: mean difference 136.02 [95% CI 82.89, 184.41]; P17 vs. P0: mean difference -100.85
[95% CI -145.40, -62.42]; P28 vs. P0: mean difference -74.95 [95% CI -119.97, -36.61]. Field of view
(FOV) = 245μm x 65μm.
g. Engulfed fibre volume peaks at P7 and decreases thereafter for LI-II. P3 vs. P0: mean difference
1.46 [95% CI -6.86, 6.29]; P7 vs. P0: mean difference 4.05 [95% CI -4.11, 8.18]; P10 vs. P0: mean
difference -3.28 [95% CI -10.76, 1.11]; P17 vs. P0: mean difference -3.69 [95% CI -11.57, -0.10]; P28
vs. P0: mean difference -4.11 [95% CI -11.79, -0.38]. Field of view (FOV) = 245μm x 65μm.
h. Engulfed fibre volume peaks at P7 and decreases thereafter for LIII-IV. P3 vs. P0: mean difference
-13.47 [95% CI -31.97, 2.15]; P7 vs. P0: mean difference 11.03 [95% CI -7.37, 26.33]; P10 vs. P0:
mean difference -7.05 [95% CI -24.52, 6.62]; P17 vs. P0: mean difference -24.55 [95% CI -40.82, -
11.27]; P28 vs. P0: mean difference -21.49 [95% CI -37.93, -7.92]. Field of view (FOV) = 245μm x
65μm.
N-numbers as indicated. Black and colourful data points indicate females and males respectively.
159

We next asked whether the high level of CD68 expression in microglia during the

first postnatal week is associated with the refinement of neural connectivity in the

dorsal horn through the removal of superfluous A-fibre terminals. Engulfment of A-

fibres was measured by quantifying VGLUT1-tdT fluorescence volume within CD68-

positive lysosomes inside Iba1-labelled microglia cells (Figure. 1 b-d, g-h).

Consistent with the postnatal increase in microglial lysosome volume, the volume of

engulfed A-fibres was high during the first postnatal week, peaking at P7 and

decreasing thereafter in both laminae I-II, (unpaired mean difference P28 vs. P0 -

4.11 [95% CI -11.79, -0.38]) and laminae III-IV (unpaired mean difference P28 vs. P0

-21.49 [95% CI -37.93, -7.92]). This effect was more pronounced in the deeper

laminae LIII-IV, which has higher volumes of fibre engulfment at all ages compared

to laminae I-II likely due to the higher density of A-fibres in the deeper laminae.

Microglial activity is required for normal A-fibre pruning in laminae I-III

Having established that microglia participate in postnatal dorsal horn remodeling of A-fibre afferent terminals, we sought to determine whether microglial function is necessary for this process to occur normally. The phospholipid scramblase TMEM16F is required for normal microglial phagocytic activity[10]. To address this question we disrupted microglial function postnatally using a Cre-inducible conditional knocking out of *Tmem16f* in microglia cells [25], and observed the structural, functional and behavioural consequences in adult animals. A *R26$^{LSL-Ai9}$* reporter line was used to label *Cxc3cr1^{Cre}*-expressing microglial cells [28,29] and *Thy1^{eGFP}* allele was used to identify A-fibres [30,31].

To control for off-target effects of Cre expression and tamoxifen administration, both cKO mice and control mice (*Cx3cr1$^{CreER/+}$; Tmem16f$^{fl/fl}$; R26$^{LSL-Ai9}$; Thy1^{eGFP}* and *Cx3cr1$^{CreER/+}$; Tmem16f$^{+/+}$; R26$^{LSL-Ai9}$; Thy1^{eGFP}* respectively) received 4-hydroxytamoxifen (4-HT) daily from P1 to P3 and were assessed at 3-4 months old. Specificity of Cre expression was confirmed with *R26$^{LSL-Ai9}$* tdT expression.

Adult microglial Tmem16f cKO mice have increased VGLUT1+ terminals in superficial but not deep laminae

Postnatal deletion of microglial *Tmem16f* resulted in adults with increased A-fibre terminal occupancy in the superficial dorsal horn, as revealed by both Thy1-GFP expression and VGLUT1 immunohistochemistry (Figure 2a-b, d). As Thy1-GFP is expressed in only a small number of sensory neurons and varied across animals, all subsequent quantification was performed using VGLUT1 immunolabelling. Synaptic density of this increased input was quantified using VGLUT1 immunohistochemistry in laminae I-III (Figure 2a upper panels). *Tmem16f* cKO mice had greater primary

afferent VGLUT1 synaptic density throughout the dorsal horn than control mice as

measured by both total synapse volume and synapse number (Figure 2b) (unpaired

mean difference VGLUT1 volume: 799.28 [95.00% CI 450.09, 1136.35], VGLUT1

number: 533.67 [95.00% CI 356.12, 695.50]). In contrast, local inhibitory VGAT

synapse density was unaltered (Figure. 2a, c) (unpaired mean difference VGAT

volume: -168.31 [95.00% CI -562.61, 253.44], VGAT number: -159.19 [95.00% CI -

416.42, 148.88]).

As synaptic engulfment was also observed in the deeper laminae we re-analysed the

data by cropping the original images to only contain LI or LIII to assess potential

differences across laminae (Figure 2e). Surprisingly, VGLUT1 density in LIII was

unaltered in contrast to LI (Figure 2g-h), suggesting that the changes observed are

mainly driven by an increase in superficial VGLUT1 terminal projections (unpaired

mean difference superficial/LI VGLUT1 count: 201.33 [95.00% CI 110.90, 292.00],

deep/LIII VGLUT1 count: 56.61 [95.00% CI -13.81, 118.42]).

To investigate whether the superfluous VGLUT1 terminals indeed form synaptic

contacts, we also co-stained spinal cord sections from *Tmem16f* cKO and control

animals with HOMER, which marks postsynaptic densities. Indeed, we found

colocalization of VGLUT1-positive Thy1-GFP labelled A-fibre terminals with HOMER

were increased in *Tmem16f* cKO animals, suggesting that a surplus of functional

VGLUT1 positive synaptic contacts are formed in the superficial dorsal horn of

Tmem16f cKO animals (Figure 2—figure supplement 1).

Thus, targeted deletion of microglial *Tmem16f*, and consequently impaired

phagocytosis specifically disrupted the normal developmental pruning of excitatory

afferent A-fibre projections in the superficial dorsal horn.

221
Figure 2. Neonatal *Tmem16f* cKO in microglia increases dorsal horn A-fibre terminals.
a. Representative images of VGLUT1 and VGAT puncta from the spinal dorsal horn of adult *Tmem16f*
control and cKO animals. Field of view (FOV) = 94µm x 94µm (VGLUT1), 96µm x 96µm (VGAT).
Scale bars = 10µm.
b. VGLUT1 puncta volume and count were both increased in adult *Tmem16f* cKO animals compared
to controls. Control vs cKO volume: mean difference 799.28 [95.00% CI 450.09, 1136.35]; control vs
cKO count: mean difference 533.67 [95.00% CI 356.12, 695.50].
c. VGAT puncta volume and count were not significantly different between adult *Tmem16f* cKO
animals and controls. Control vs cKO volume: mean difference -168.3105 [95.00% CI -562.61,
253.44]; control vs cKO count: mean difference -159.19 [95.00% CI -416.42, 148.88].
d. Thy1-GFP labelled A-fibres (yellow) are present in the superficial dorsal horn (delimited by IB4,
blue) of adult cKO mice, but not in adult controls. tdTomato-labelled microglia (cyan) are present in
both cKO and control animals. Lower panels show high power images of the boxed areas in the
upper panels. Scale bar = 100µm in the top panel and 20µm in the lower panel.
e. Thy1-GFP labelled A-fibres (yellow) in the superficial dorsal horn laminae of adult *Tmem16f* cKO
mice express VGLUT1 (magenta). (A) Low magnification image showing superficially-projecting Thy1-
GFP labelled fibres. White box indicates where images were taken for analysis in **a-c**. (B) High
magnification of boxed region in A. Cyan boxes indicate the cropped areas used for analysis in **g** and

h. (C) Thy1-GFP labelling. (D) VGLUT1 immunoreactivity. Scale bar in A = 100µm. Scale bar in D =
10µm.
f. A-D. High-magnification examples for VGLUT1-expression (magenta) in Thy1-GFP labelled A-fibres
(yellow) in adult *Tmem16f* cKO mice. Scale bar in A = 20µm, scale bar in D = 5µm.
g, h. VGLUT1 count in adult *Tmem16f* cKO and control animals in superficial lamina I (**g**) and deep
lamina III (**h**). LI VGLUT1 count mean difference 201.33 [95.00% CI 110.90, 292.00]; LIII VGLUT1
count mean difference 56.61 [95.00% CI -13.81, 118.42]. FOV = 31µm x 94µm.
N-numbers as indicated. Black and red data points indicate females and males respectively.
248

Dorsal horn sensory neurons in adult microglial Tmem16f cKO mice are less

responsive to dynamic tactile stimulation of the skin, but have larger receptive

fields

We reasoned that the reduced A-fibre pruning due to impaired neonatal microglial

phagocytosis in *Tmem16f* cKO mice would alter behavioural sensitivity to hindpaw

tactile stimulation. The hindlimb withdrawal reflex was measured in response to

brushing of the plantar surface of the paw. *Tmem16f* cKO mice showed a greater

number of withdrawals in response to dynamic brushing than controls (unpaired

mean difference 0.83 [95% CI 0.072, 1.57]), but not to static vF stimulation (Figure 3,

Figure 3—figure supplement 1). *Tmem16f* cKO animals therefore displayed

cutaneous hypersensitivity specifically towards dynamic low-threshold tactile

stimulation.

We hypothesised that excessive A-fibre terminals in dorsal horn might lead to

increased dorsal horn activity which could underlie the behavioural hypersensitivity.

To test this, we used *in vivo* single unit extracellular recordings in the dorsal horn of

anaesthetised *Tmem16f* cKO and control mice. We recorded from adapting and non-

adapting wide dynamic range neurons (WDR) in the deep dorsal horn, which

according to the criteria by Lee et al. 2019 [32] are presumed to be excitatory and

inhibitory neurons respectively. Of 30 cells recorded from control mice, 8 were non-

adapting and 22 were adapting. Of 28 cells recorded from *Tmem16f* cKO mice, 5

were non-adapting and 23 were adapting. This is consistent with the expected ratio

of 1:2 of inhibitory to excitatory cells as reported in previous studies [32–34] (P=0.56 for

control and P=0.11 for cKO in Binomial test against expected ratio). We quantified

spinal neuronal response to 1) dynamic innocuous tactile stimuli (brush), and 2)

static von Frey hair (vF) stimuli applied to the plantar hindpaw as well as mapped

dynamic touch-sensitive receptive field areas on the plantar hindpaw.

276
Figure 3. Neonatal *Tmem16f* deletion in microglia increases dynamic brush sensitivity
a. Brush withdrawal response for *Tmem16f* cKO animals are higher than control animals, mean
difference 0.83 [95% CI 0.072, 1.57].
b. vF withdrawal response did not differ between *Tmem16f* cKO and control animals, mean difference
0.12 [95% CI -0.39, 0.65].
N-numbers as indicated. Black and red data points indicate females and males respectively.
283

We reasoned that excessive A-fibre presence in the dorsal horn would increase their

spatial connectivity with postsynaptic neurons, leading to larger dorsal horn receptive

field areas. Consistent with increased A-fibre synaptic contacts, mean brush

receptive field sizes for adapting neurons were increased by almost 50% (unpaired

mean difference 13.7 [95.00% CI 3.3, 27.2]), with a notable subpopulation of cKO

neurons expanding their receptive fields to cover the entire, or majority of the plantar

surface (Figure 4a, b). The expansion of receptive field area was accompanied by a

reduced number of spikes in response to both brush and vF stimulation in *Tmem16f*

cKO adapting neurons (unpaired mean difference brush: -11.4 [95% CI -19.2, -4.5],

vF: -6.4 [95% CI -8.5, -4.5]) as well as a reduction in spontaneous activity (Figure 4c-

f, Figure 4—figure supplement 1a).

In contrast, brush receptive field sizes and brush-evoked responses did not differ

between control and *Tmem16f* cKO mice in non-adapting (inhibitory) neurons (Figure

4g-j), though an increase in vF-evoked responses in *Tmem16f* cKO animals was

observed (mean difference 2.74 [95.00% CI -0.074, 6.25], $F_{(1; 110)}$ = 7.014, P =

0.0093) (Figure 4k, l). Spontaneous activity was also unaltered (Figure 4—figure

supplement figure 1b).

301
Figure 4. Neonatal *Tmem16f* deletion in microglia decreases evoked activity, but increases
brush receptive field size
a. Brush receptive field sizes was increased for adapting neurons in *Tmem16f* cKO, mean difference
13.7 [95.00% CI 3.3, 27.2].
b. Overlay of receptive fields in **a**.
c. Brush evoked response was decreased for adapting neurons in *Tmem16f* cKO, mean difference -
11.4 [95% CI -19.2, -4.5].
d. Example firing trace of adapting cell from control and cKO animals to brush stimulation with raster
plots underneath.
e. vF evoked activity was decreased for adapting neurons in *Tmem16f* cKO, mean difference -6.4
[95% CI -8.5, -4.5].

f. Example firing trace of an adapting cell from control and cKO animals to vF stimulation with raster
plots underneath.
g. Brush receptive field sizes was unchanged for non-adapting neurons in *Tmem16f* cKO, mean
difference 5.53 [95.00% CI -15.91, 40.43].
h. Overlay of receptive fields in **g**.
i. Brush evoked response was unchanged for non-adapting neurons in *Tmem16f* cKO, mean
difference 8.62 [95.00% CI -5.78, 22.72].
j. Example firing trace of non-adapting cell from control and cKO animals to brush stimulation with
raster plots underneath.
k. vF evoked activity was decreased for non-adapting neurons in *Tmem16f* cKO, mean difference
2.74 [95.00% CI -0.074, 6.25].
l. Example firing trace of a non-adapting cell from control and cKO animals to vF stimulation with
raster plots underneath.
N-numbers as indicated. Black and colourful data points indicate females and males respectively.
327

Discussion

In summary, we report that dorsal horn microglia have a distinct phenotype during

the first postnatal week characterized by high phagocytic activity, which coincides

with postnatal engulfment of A-fibre terminals in the dorsal horn, supporting a role of

microglia in developmental dorsal horn remodelling. We further show that disruption

of microglial function by targeted deletion of *Tmem16f* during early postnatal life

impairs microglia mediated A-fibre refinement in the dorsal horn leading to long-term

changes in dorsal horn function and behaviour that persist into adulthood. Together,

our data suggests that microglia mediated refinement of A-fibres during the early

postnatal period is critical to both normal dorsal horn development and appropriate

spatial encoding of dynamic touch.

In our study we looked at a subset of VGLUT1 positive A-fibres, which are

predominantly low-threshold, myelinated Aβ mechanoreceptors [24,35], and primarily

transduce innocuous mechanical stimulation of the skin, such as stroking and

brushing. In adults, the superficial laminae is occupied by both A- and C-fibre

projections, but the during the neonatal period A-fibre inputs dominate the superficial

dorsal horn, as nociceptive C-fibre inputs are weak and only start to strengthen

between P5-P10 [36,37]. Functionally, this results in lower cutaneous mechanical

thresholds for activity in the superficial laminae, which together with immature local

and descending inhibition leads to exaggerated reflex behaviour in neonates [5–7,38].

This early sensitivity to low threshold stimuli might be particularly important during

the neonatal period for maternal bonding, as skin to skin contact after birth was

shown to reduce neonatal stress and pain response and improve mother-infant

interaction [39]. Therefore, appropriate A-fibre maturation might play an important role

in overall infant development.

Microglia cells show the strongest increase in density over the first postnatal week

which coincides with a decrease in phagocytic activity. Hammond et al. 2019

revealed a specialised phagocytic population of microglia at P4/5 in the brain, while

our data suggests that the peak of microglial phagocytic activity might be later in the

spinal cord around P10. Moreover, dorsal horn microglial density observed in our

study appears to be higher than in various brain regions studied at comparable ages

[21,40] (Table 1). Together, this suggests that dorsal horn microglia follow a different

developmental trajectory than brain microglia.

Paper	Brain area	Age	Density per μm^3
Schwarz et al. 2012 [21]	Parietal cortex	P4	2.23×10^{-7}
	Amygdala	P4	4.77×10^{-6}
	CA1 of hippocampus	P4	4.88×10^{-7}
	Paraventricular nucleus	P4	1.37×10^{-6}
Perez-Pouchoulen et al. 2015 [40]	Cerebellum	P5	3×10^{-6}
Present study (Figure 1—figure supplement 1d)	Superficial dorsal horn	P3	5×10^{-6}

Table 1. Microglial density across different CNS regions (in rats). Estimated numbers
from the graphs in respective paper, with male and female microglia counts pooled.

While microglial CD68 volume increased postnatally, the amount of microglial

phagocytic cup numbers decreased. Phagocytic cups are relatively large structures

involved in phagocytosing cell bodies [41], and their reduction is consistent with the

decline in apoptotic cell numbers postnatally. This shows that the postnatal increase

in CD68 cannot be due to removal of apoptotic cell bodies but rather due to the

refinement of neuronal connections.

Dorsal horn microglia phagocytose A-fibre projections both in the superficial (laminae

I-II) and deeper (laminae III-IV) laminae during normal postnatal development. The

data here shows that most of the A-fibre engulfment occurs before P10 and

decreases thereafter. The overall amount of A-fibre engulfment is likely

underestimated, as reporter mice crossed with the *Vglut1*-Cre line used here only

express tdT in a subset of A-fibres [24]. Further, the developmental upregulation of

Vglut1 expression [26,27] means that engulfment will be especially underestimated in

younger animals, as their A-fibres might be present but not yet expressing tdT. In the

brain, it has been shown that VGLUT1 protein expression at P0 is only 6.6% of the

adult level, reaching 47% by P10 and 92% by P20 [27]. Therefore, the fall in A-fibre

engulfment over the first postnatal week is likely much sharper in reality than Figure

1 g, h suggests.

Although microglial CD68 volume peaks at P10, the engulfed VGLUT1-tdT volume

seems to peak earlier at P7. While the percentage of lysosome volume occupied by

VGLUT1-tdT is about 15% in the superficial laminae and 40% in the deep laminae at

P7, by P10 it has become less than a quarter of that. It is possible that a slowing rate

in engulfment of VGLUT1-tdT labelled fibres between P10 and P7 allows VGLUT1-

tdT to be degraded in the lysosomes at a faster rate than new VGLUT1-tdT material

is engulfed, thus causing a reduction of VGLUT1-tdT at P7 before the reduction in

lysosome volume follows suit by P10. It is also possible that developmental events at

P10 require additional microglial phagocytosis of materials other than A-fibres.

Neonatal *Tmem16f* cKO increased the number of Thy1-GFP positive VGLUT1

synapses in the superficial dorsal horn in adults, suggesting that *Tmem16f* function

is necessary for microglia mediated refinement of A-fibres.

However, increase in presynaptic terminals seems to predominantly affect lamina I,

but not lamina III, even though developmental engulfment of A-fibres was observed

both in LI-II and LIII-IV, suggesting that either only the engulfment of superficial

synapses is dependent on *Tmem16f* or alternatively that superfluous synapses in the

deeper laminae were more efficiently compensated and removed at later stages

before animals reached adulthood.

The density of VGAT presynaptic terminals was also unaltered in *Tmem16f* cKO

animals. This suggests that *Tmem16f* mediated engulfment is synapse specific. A

recent study in the hippocampus suggested that complement mediated synapse

elimination by microglia was targeted to VGLUT2, but not VGLUT1 or VGAT

synapses instead [42]. Together with our data this suggest that developmental

synapse elimination is both specific to synapse identity and location.

Consistent with the increase in dorsal horn VGLUT1, *Tmem16f* cKO increased brush

receptive field size in adapting/excitatory neurons, which mirrors the large receptive

field sizes in immature neonatal animals [5]. During the neonatal period A-fibre inputs

dominate the superficial dorsal horn [36,37] and hindpaw receptive fields are initially

large, reducing in size over the first two postnatal weeks [5]. Functionally, this results

in lower cutaneous mechanical thresholds, which together with immature local and

descending inhibition leads to exaggerated reflex behaviour in neonates [5–7,38]. Both

enlarged receptive field sizes and hypersensitivity were observed in Tmem16f cKO

animals. This is unlikely due to local disinhibition [43,44], as spontaneous and evoked

activity of adapting WDR neurons were not increased, suggesting that the larger

receptive field sizes reflect a lack of primary afferent refinement instead.

Consistent with the increased A-fibre input and brush receptive field size, which likely

affects spatial summation of dynamic inputs, *Tmem16f* cKO increased behavioural

sensitivity to brush. However, counterintuitively, the increase in A-fibre projections

and behavioural sensitivity was accompanied by a reduction in evoked activity for

brush and vF as well as spontaneous activity in adapting neurons.

How does decreased evoked activity result in higher reflex sensitivity? One

possibility is that the activation pattern of WDR neurons may be more important than

firing rate per se for the coding of sensory information. Although the activity of

individual neurons is decreased, potentially more neurons are activated in total due

to the lack of A-fiber refinement. Therefore, the activation of WDR neurons on a

population level might be a better predictor of the spinal reflex response. A potential

mechanism underlying this effect is homeostatic synaptic downscaling, which

maintains homeostasis by reducing neuronal firing rate when the overall network

activity is increased [45]. Our data shows that activity recorded from single neurons do

not always positively correlate with behavioural changes and need to be interpreted

with care.

In contrast to the adapting neurons, little change was observed for non-

adapting/inhibitory neurons, apart from an increase in vF-evoked activity. This is

likely due to the low sample size, as fewer non-adapting/inhibitory neurons were

recorded. The increased firing rate of non-adapting/inhibitory neurons during vF-

evoked activity would likely decrease the activity of adapting/excitatory neurons and

thus contribute to the decreased overall activity observed in cKO animals.

We propose therefore that disrupting microglial function with *Tmem16f* cKO causes

an increase in VGLUT1 positive A-fibre terminals in adult animals, potentially due to

failed pruning of A-fibres in the neonatal period. This excess of peripheral input might

preferentially target excitatory neurons which leads to larger receptive field sizes

thereof, and potentially activates larger numbers of excitatory neurons upon

peripheral stimulation. In compensation, through homeostatic scaling or increased A-

fibre input onto non-adapting/inhibitory neurons, inhibitory activity onto excitatory

neurons is increased, decreasing the firing rate of individual neurons, but not

necessarily the total output as a population. This is also consistent with increased

behavioural sensitivity in *Tmem16f* cKO animals.

Limitations

Several caveats need to be considered when interpreting the data. In our

experiments we used a CNS-wide KO of microglial *Tmem16f*. Therefore, we cannot

rule out cortical contributions to the observed changes in the spinal cord – for

example, top-down facilitation of inhibitory inputs or inhibition of excitatory inputs

could also explain the decreased firing rate in adapting and non-adapting neurons in

Tmem16f cKO animals. However, a purely top-down mediated overall inhibition of

dorsal horn activity would not explain the increase in receptive field size following

Tmem6f 1 cKO, though it could additionally contribute to the changes observed.

Therefore, at least part of the observed changes must be mediated by changes in

peripheral afferent input to the dorsal horn.

Furthermore, it is possible that *Tmem16f* cKO will also affect the engulfment and

removal of apoptotic cells by microglia, which contribute to the behavioural and

functional changes seen in adulthood. However, the mechanisms involved in

engulfing whole cell bodies vs synaptic material are likely distinct (in our study CD68

volume was not correlated with apoptotic clearance), and therefore *Tmem16f* cKO

might affect microglial pruning without affecting the phagocytosis of apoptotic cells.

We used VGLUT1 as a marker for presynaptic boutons of A-fibres as well as the

driver of tdT expression. However, it has been shown that cortico-spinal projections

also contribute to VGLUT1 synapses throughout the dorsal horn [33]. Therefore, we

cannot exclude that the behavioural and functional effects seen in the *Tmem16f* cKO

animals are due to disruption of cortico-spinal refinement. However, during early

postnatal development, corticospinal projections do not enter in the grey matter

lumbar segment until P7, while peripheral A-fibre afferents reach the lumbar dorsal

horn by embryonic day (E)14 [46,47]. Therefore, we can conclude that at least prior to

P7, where the majority of VGLUT1-tdT engulfment was observed, the engulfment of

VGLUT1-tdT labelled fibres are indeed afferent A-fibres.

An important open question is what determines the engulfment or survival of

synapses and fibres in the dorsal horn. It is known that microglia can sense and

respond to neuronal activity [48–51]. For example, microglia increase process motility

both to neuronal hyper- and hypoactivity [49], and in the brain, microglial engulfment

seems to preferentially target weak synapses [12,13]. Given that perturbation of

neuronal activity can indeed impair A-fibre refinement [8,9], neuronal activity seems a

likely signal for microglial engulfment in the dorsal horn. Several molecules

expressed by neurons and microglia have been identified as "eat me" and "keep me"

signals [12,13,52,53], among which members of the complement cascade, e.g. C1q, have

been shown to tag synapses for microglial engulfment both in the brain and the

spinal cord [12,17]. Whether C1q also plays a role in dorsal horn remodeling remains to

be investigated.

Conclusions

In summary, we have shown that dorsal horn microglia phagocytose A-fibres during

normal postnatal development and that disruption of microglial function can lead to

long-term structural and functional changes in the dorsal horn and behavioural

changes towards dynamic touch. This has important implications for perinatal care of

infants.

Low-threshold stimuli such as skin-to-skin contact have been shown to reduce infant

stress and improve infant-mother interaction and maternal bonding [39]. Therefore,

appropriate maturation of A-fibres might play an important role in overall infant

development. In addition, previous research have shown that peripheral injury during

the first postnatal week (but not after P10) cause hyperalgesia upon re-injury in

adulthood, and that this could be prevented with intrathecal minocycline injection,

which is a non-specific microglial inhibitor [54,55]. It is tempting to speculate that injury

alters microglial pruning of A-fibres as described here which could underlie part of

the long-term changes seen following injury. Thus, the role of microglia in normal

and aberrant neonatal development could be investigated for therapeutic potential in

perinatal care.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Genetic reagent (*M. musculus*)	Ai9; *R26$^{LSL-Ai9/+}$*	Jackson Laboratory	Stock # 007909 RRID:IMSR_JAX:007909	PMID: 20023653
Genetic reagent (*M. musculus*)	Ai9; *R26$^{LSL-Ai9/+}$*	Jackson Laboratory stock	Stock # 007905 RRID:IMSR_JAX:007905	PMID: 20023653
Genetic reagent (*M. musculus*)	*Slc17a7*-IRES2-Cre; *Vglut1$^{Cre/+}$*	Jackson Laboratory stock	Stock # 023527 RRID:IMSR_JAX:023527	PMID: 25071457
Genetic reagent (*M. musculus*)	*Cx3cr1*-CreERT2-YFP; *Cx3cr1$^{CreER/+}$*	Jackson Laboratory stock	Stock # 021160 RRID:IMSR_JAX:021160	PMID: 35045285
Genetic reagent (*M. musculus*)	*Thy1*-EGFP-M; *Thy1^{eGFP}*	Jackson Laboratory stock	Stock # 007788 RRID:IMSR_JAX:007788	PMID: 11086982
Genetic reagent (*M. musculus*)	*Tmem16f*-floxed; *Tmem16f$^{fl/fl}$*	PMID: 27332874		Dr Paul Heppenstall (SISSA, Trieste, Italy)
Antibody	anti-Caspase-3 (Rabbit monoclonal)	Cell Signalling	Cat# Asp175, 5A1E	IF (1:100)
Antibody	anti-CD68 (Mouse monoclonal)	BIO-RAD	Cat# MCA341R	IF(1:500)
Antibody	anti-CD68 (Rabbit polyclonal)	Abcam	Cat# ab125212	IF(1:500)

Antibody	anti-GFAP (Mouse, monoclonal)	Sigma	Cat# G3893	IF(1:2000)
Antibody	anti-Iba1 (Rabit, polyclonal)	Wako	Cat# 019-19741	IF(1:1000)
Antibody	anti-Iba1 (Goat, polyclonal)	Abcam	Cat# ab5076	IF(1:1000)
Antibody	anti-NeuN (Mouse, monoclonal)	Millipore	Cat# MAB377	IF(1:2000)
Antibody	anti-VGAT (Mouse, monoclonal)	Synaptic Systems	Cat# 131011	IF(1:2000)
Antibody	anti-VGAT (Rabbit, polyclonal)	Synaptic Systems	Cat# 131002	IF(1:2000)
Antibody	anti-VGLUT2 (Guinea pig, polyclonal)	Millipore	Cat# AB2251-I	IF(1:2000)
Antibody	anti-Synaptophysin (Mouse, monoclonal)	Abcam	Cat# Ab8049	IF(1:2000)
Antibody	anti-Synaptophysin (Guinea pig, polyclonal)	Synaptic Systems	Cat# 101004	IF(1:2000)
Antibody	anti-goat Cy2 (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 705-225-147	IF(1:500)
Antibody	anti-goat Cy3 (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 705-165-147	IF(1:500)
Antibody	anti-goat Cy5 (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 705-175-147 1	IF(1:500)

Antibody	anti-goat Alexa 488 (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 705-545-003	IF(1:500)
Antibody	anti-guinea pig Cy3 (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 706-165-148	IF(1:500)
Antibody	anti-guinea pig Alexa 647 (Donkey polyclonal)	Millipore	Cat# AP193SA6	IF(1:500)
Antibody	anti-mouse Cy2 Jackson (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 715-225-150	IF(1:500)
Antibody	anti-mouse Cy3 Jackson (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 715-165-150	IF(1:500)
Antibody	anti-mouse Cy5 (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 715-175-150	IF(1:500)
Antibody	anti-rabbit Cy2 (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 711-225-152	IF(1:500)
Antibody	anti-rabbit Alexa 594 (Donkey polyclonal)	Jackson ImmunoResearch	Cat# 711-585-152	IF(1:500)
Antibody	anti-rabbit Alexa 647 (Donkey polyclonal)	Millipore	Cat# AP182SA6	IF(1:500)
Antibody	anti-rabbit-biotinylated (Horse polyclonal)	Vector	Cat# BA11-00	IF(1:500)
peptide, recombinant protein	Streptavidin-594	Life Technologies	Cat# S32356	IF(1:500)

Software, algorithm	ImageJ; FIJI	PMID:22930834, PMID:29187165	RRID:SCR_003070	
Software, algorithm	Volocity	PerkinElmer	RRID:SCR_002668	
Software, algorithm	LabChart 7	ADInstruments		
Software, algorithm	Graphpad prism	GraphPad Software, Inc.	RRID:SCR_002798	
Software, algorithm	Estmationstats. com (web application)	PMID: 31217592	RRID:SCR_018321	Free web application
other	Carbostar-1 (micro-electrode)	Kation Scientific		

Animals

Sprague Dawley rats of both sexes were used for experiments in Figure 1—figure

supplement 1d-h and Figure 1—figure supplement 2. Transgenic mice on C57BL/6J

background of both sexes were used in all other experiments.

Experiments used the following transgenic mouse lines:

1. Ai9 / Rosa26-CAG::loxP-STOP-loxP-tdTomato-WPRE (Jackson Laboratory

stock 007909 & 007905)

2. *Slc17a7*-IRES2-Cre (Jackson Laboratory stock 023527)

3. *Cx3cr1*-CreERT2-YFP (Jackson Laboratory stock 021160)

4. *Thy1*-EGFP-M (Jackson Laboratory stock 007788)

5. *Tmem16f*-floxed (flx) animals (Batti et al., 2016)

For visualisation of A-fibres, *Slc17a7*-IRES2-Cre (*Vglut1*-Cre) males (JAX stock no.

023527) were crossed with Ai9 females (JAX stock no. 007909) to obtain animals

that expressed the tdTomato fluorophore under the *Vglut1* promoter (*Vglut1$^{Cre/+}$;*

R26$^{LSL-Ai9/+}$).

To generate tamoxifen inducible microglia-specific *Tmem16f* knock-out mice

(*Tmem16f* cKO), Cx3cr1-CreER-YFP (JAX stock no. 021160) mice were crossed to

Tmem16f-flx animals (generated by P. Heppenstall, see Batti et al. 2016), as well as

Ai9 (JAX stock no. 007905) and *Thy1*-EGFP-M (JAX stock no. 007788).

Experimental animals were heterozygous for *Cx3cr1*-CreER-YFP ($^{CreER/+}$),

homozygous for mutant conditional allele *Tmem16f*-flx ($^{fl/fl}$), and carrying Ai9 (*R26^{LSL-}*

^{Ai9}) and *Thy1*-eGFP (^{eGFP}) alleles (zygocity was not determined for *R26$^{LSL-Ai9}$* and

Thy1^{eGFP}). This produced the following genotype: *Cx3cr1$^{CreER/+}$; Tmem16f$^{fl/fl}$; R26^{LSL-}*

^{Ai9}; *Thy1^{eGFP}*. Control animals were homozygous for the wild type *Tmem16f* allele:

Cx3cr1$^{CreER/+}$; Tmem16f$^{+/+}$; R26$^{LSL-Ai9}$; Thy1^{eGFP}.

Both females and males were used. No sex differences were expected and animals

of both sexes were pooled together for analysis, but data points are presented as

black (female) or red/magenta/blue (male) to indicate the sexes. Numbers of animals

used for each experiment are indicated in the figures. For a table with detailed

species, ages, sexes, and numbers of animals used in each experiment, please see

supplementary Table 1. All procedures were carried out in accordance with the

guidelines of the UK Animals (Scientific Procedures) Act 1986 and subsequent

amendments.

Drugs

4-hydroxytamoxifen (4-HT) was dissolved at 1mg/ml in corn oil, and 50µl was

injected intragastrically per pup daily on three consecutive days from P1-3, following

a previously described protocol [56]. Both control and experimental animals

(*Tmem16*[+/+] and *Tmem16f*[fl/fl] respectively, see animals section above) received 4-HT

injections to control for any effects of 4-HT itself. The dam was given a protein

enriched diet a few days before and following delivery to aid milk production and pup

survival.

Immunohistochemistry

Animals were overdosed with pentobarbital and transcardially perfused with saline

followed by ice-cold 10% formalin. The sciatic nerve was exposed and traced to

locate L4 & L5 dorsal root ganglia (DRG) and the corresponding region of the lumbar

spinal cord was dissected and post-fixed in 10% formalin overnight, followed by

immersion in 30% sucrose until they sank. 50µm free-floating spinal cord sections

were cut on the microtome with every 2nd section collected.

Tissue sections were washed 3 × 10 min in PBS and then incubated in blocking

solution (10% donkey serum, 0.2% Triton X-100 in PBS) for 2.5h at room

temperature. The sections were then incubated with primary antibodies at 4°C

overnight followed by secondary antibodies at room temperature for 2h, both diluted

in 3% blocking solution (3% donkey serum, 0.2% Triton X-100 in PBS) (for list of

antibodies and their respective concentrations used, see Key Resources Table).

Samples were mounted in Fluoromount Aqueous Mounting Medium (Sigma) or

ProLong[TM] Diamond Antifade Mountant (Thermo Fischer), if the tissue contained

endogenous fluorophores.

Image acquisition and analysis

Confocal z-stacks were taken with a Zeiss LSM880 confocal microscope or

Yokogawa CSU22 spinning disk microscope using a 20× water immersion objective

(NA 1.0) for imaging of A-fibres and 63× oil immersion objective (NA 1.4) followed by

analysis in Fiji software. Details on microscope settings can be found in

supplementary Table 1 and in the metadata of example images online at

https://github.com/Yajing826/A-fibre-engulfment. Only intact sections with an even

stain were analysed, and at least 6 sections were imaged and analysed per animal

to reduce variability for all figures (apart from S3, where 1-3 sections were analysed

per animal).

Cell counts and phagocytic cup counts were performed manually on confocal images

in Fiji or Volocity, with the experimenter blinded to the age groups. Phagocytic cups

were defined as any rounded structure at the end of a microglial process.

A-fibre engulfment by microglia and synapse density were analysed with automated

batch processing in Fiji using the 3D-ROI manager plugin and custom written macros

[57–60]. For A-fibre engulfment, each of the channels containing staining for microglia,

lysosomes, or A-fibres were binarized and the volume of their overlap measured. For

synapse density measures, the channel containing synaptic stain was binarized and

segmented, following which volume and object numbers were recorded. Macro

scripts for the automated analysis are available online at

https://github.com/Yajing826/A-fibre-engulfment.

Behaviour

Behavioural testing was carried out on adult mice of both sexes between 3-4 months old, with the experimenter blinded to animal genotype/treatment. Animals were placed on a mesh platform (Ugo Basile) within individual transparent plastic chambers (6cm x 6cm x 12cm) for sensory testing of the plantar surface of the hind paw. Habituation and testing happened over five consecutive days. Animals were habituated to the testing environment for 1h per day on the first two days within individual plexicon chambers on the mesh platform. On the remaining days, animals were habituated for 30 min before being tested on each day. Brush response and von-Frey (vF) threshold were determined on the 3rd day, while repeated vF response testing was spread over the remaining 2 days to avoid sensitisation. Number of withdrawal reflexes were scored in each case, where only a rapid paw lifting was scored as a reflex. Animals were allowed to rest at least 20 seconds between each stimulus. For brush response, a fine brush (Pro Arte, series 007, size 2) was moved over the plantar surface of the hind paw from heel to toe over a 2 second period [61]. This was repeated five times and the number of withdrawal reflexes out of five was recorded. vF threshold was assessed using the simplified up and down method [62]. Filaments were aimed at the region between the foot pads. Force was applied until the filament bent, and held in place for 2 seconds.

To generate a response curve to vF stimulation, repeated vF response was recorded by applying each of filaments no. 3 - 7 (0.04g - 0.6g) five times on the plantar surface, directed at the region between the foot pads. The sequence of vF filaments was randomised. Number of withdrawal reflexes out of five times was recorded.

In vivo extracellular recording

Animals subjected to behavioural testing were reused in electrophysiological

recordings. Experimenter was blinded to animal genotype/treatment. All recordings

were performed on adult mice (3-4 months old) of both sexes in the deep dorsal

horn. Cells were not recorded beyond 550 µm depth from the surface of the spinal

cord (see Figure 4—figure supplement 1c for depth of all recorded neurons). 3-5

mice were used per sex and treatment group. All experiments were carried out by

the same experimenter to ensure consistency.

<u>Animal preparation</u>

Mice were anaesthetised with intraperitoneal urethane injection (10% in saline,

1.5g/kg). 100µl of 0.6 g/ml atropine and 200µl saline were injected subcutaneously to

respectively counteract the mucus-driving side effect of urethane and to prevent

dehydration. The animal was constantly monitored for depth of anaesthesia

throughout the experiment and supplemented with 50µl (5µg) urethane as needed.

200µl of saline was supplemented every 2 hours. Body temperature of the animal

was kept close to 37°C with a heating pad throughout.

After cessation of reflexes, a tracheotomy was performed and a short plastic tube of

about 1cm inserted to aid free breathing of the animal. The animal was then

transferred onto a stereotactic frame and fixed with ear and hip bars. A laminectomy

was carried out at vertebral level T13-L1 which corresponds to the spinal segments

L4-L5 underneath. The spinal column was clamped for stability, the dura was

removed, and the exposed spinal cord was covered with mineral oil to prevent

drying.

<u>Single unit extracellular recordings</u>

A carbon micro-electrode (Carbostar-1, Kation Scientific) was lowered with a

motorised manipulator (Scientifica) into the exposed spinal cord at a constant speed.

A reference electrode was inserted into the back muscle close to the laminectomy for

differential recording. Recorded neural activity was amplified 2000 times, and filtered

for signals between 1kHz-10kHz (NL104 amplifier and NL125/6 band-pass filter

modules from NeuroLog Digitimer). The signal was sampled at 20kHz and digitised

using Powerlab 4/30 (ADInstruments). The trace was recorded and analysed in the

software LabChart 7 (ADInstruments).

To isolate single neurons, the plantar surface of the animal's hind paw was gently

continuously stroked as a searching signal, while the electrode is being lowered

through the dorsal horn of the spinal cord. Once a cell has been identified by equal

amplitude of the spikes recorded, the brush receptive field of the cell was mapped

out by a fine brush (Pro Arte, series 202, size 1, brush tip cut short to 7mm length x

1mm width).

Spontaneous activity was recorded for 10 min before and 5 min after stimulation. To

record brush and von-Frey (vF) fibre evoked activity from single neurons, each

stimulus was manually applied for 2 seconds over the receptive field of the cell and

repeated 3 times, with a minimum of 10 second interval in between. (vF filament

strength were as follows: 1 = 0.008g, 2 = 0.02g, 3 = 0.04, 4 = 0.07g, 5 = 0.16g, 6 =

0.40g, 7 = 0.60g, 8 = 1g, 9 = 1.7, 10 = 2g).

Cells were not recorded if they exhibited very high spontaneous firing rates that did

not allow evoked activity to be clearly distinguished from spontaneous activity. Only

wide dynamic range neurons responding both to brush and pinch stimulation were

recorded. Animals were euthanised at the end of the experiment and the spinal cord

was collected in neutral buffered 10% formalin (overnight) followed by 30% sucrose

solution for re-use in immunohistochemistry.

<u>Analysis & cell type categorisation</u>

Analysis was carried out in the LabChart 7 software (ADInstruments). For

spontaneous activity, firing rate was analysed over a 10min window prior to applying

any stimuli. Firing rate for evoked responses were analysed over the first second of

the stimulus duration and averaged over three trials.

Cells were divided into adapting and non-adapting groups based on their firing

properties towards a threshold vF-stimulus [32]. This threshold vF was defined as the

first vF filament that evokes a firing rate of 10Hz or more. The response within the

first second of vF application was

analysed to calculate an adaptive ratio R, which was defined as

$$R = \frac{Number\ of\ spikes\ fired\ between\ 0.5 - 1\ sec}{Number\ of\ spikes\ fired\ between\ 0 - 0.5\ sec}$$

If a cell adapts rapidly to stimulation, one would expect R to be close to zero, as

barely any spikes should be fired between 0.5 - 1 sec, however, if a cell is non-

adapting and firing continuously, one would expect R to be close to 1. To decide the

boundary between adapting and non-adapting cells we used k-means cluster

analysis, which sorted the values into two groups that is equivalent to a boundary at

R = 0.33. For the k-means clustering we included cells from a previous experiment.

Statistical Analysis

Estimation statistics for the 95% confidence intervals (95% CI) of the mean

difference were calculated on estimationstats.com [63] using 5000 samples of bias-

corrected and accelerated bootstrapping. As bootstrapping is less accurate for small

samples sizes [64], confidence intervals were only calculated for samples with N≥5.

For Figure 1—figure supplement 1d, h and Figure 1—figure supplement 2 where

N=4, one-way ANOVA was used with post-hoc comparisons carried out using

Dunnett's method.

Additionally, conventional null-hypothesis significance testing was carried out on

estimationstats.com and GraphPad Prism 6 for all comparisons (significance level

was set at α=0.05), which are listed in the Appendix 1—table 1.

Data are presented as mean ± SD in all figures. Where applicable, the effect size is

presented as 95% CI of the mean difference on a separate but aligned axis. The

mean difference is plotted as a dot on the background of its probability distribution,

and the 95% confidence interval is indicated by the ends of the error bar. All values

in text and figures are given with two decimals or rounded to two significant figures.

N-numbers are as indicated in figures. For a comprehensive list with exact statistical

values and analyses, see Appendix 1—table 1.

Data availability

All data generated or analysed during this study are included in the manuscript and

supporting   files.   Source   files   for   all   data   are   available   at:

https://www.ebi.ac.uk/biostudies/ under the accession number S-BSST609. Macro

scripts   for   automated   analysis   in   Fiji   are   available   online   at

https://github.com/Yajing826/A-fibre-engulfment.

Funding Information

This work was supported by the NIAA W1071H (SB) and Wellcome Trust 109006/Z/15/A (YX).

Author Contributions

YX and SB performed the experiments, YX, SK, MF and SB designed the study, analysed and interpreted data and co-wrote the manuscript. AC, QH and RRJ developed the mouse model and prepared tissue for analysis. MS and PH developed the mouse model.

Competing Interest Statement

The authors declare no competing interests.

References

References

1. Abraira, V. E. & Ginty, D. D. The Sensory Neurons of Touch. *Neuron* **79**, 618–639
(2013).
2. Fitzgerald, M., Butcher, T. & Shortland, P. Developmental changes in the laminar
termination of A fibre cutaneous sensory afferents in the rat spinal cord dorsal horn. *J.*
Comp. Neurol. **348**, 225–233 (1994).
3. Pignatelli, D., Ribeiro-da-Silva, A. & Coimbra, A. Postnatal maturation of primary
afferent terminations in the substantia gelatinosa of the rat spinal cord. An electron
microscopic study. *Brain Res.* **491**, 33–44 (1989).
4. Jennings, E. & Fitzgerald, M. C-fos can be induced in the neonatal rat spinal cord by
both noxious and innocuous peripheral stimulation. *Pain* **68**, 301–306 (1996).
5. Fitzgerald, M. The post-natal development of cutaneous afferent fibre input and
receptive field organization in the rat dorsal horn. *J. Physiol.* **364**, 1–18 (1985).
6. Fitzgerald, M., Shaw, A. & MacIntosh, N. Postnatal development of the cutaneous
flexor reflex: comparative study of preterm infants and newborn rat pups. *Dev. Med.*
Child Neurol. **30**, 520–526 (1988).
7. Fitzgerald, M. What do we really know about newborn infant pain? *Exp. Physiol.* **100**,
1451–1457 (2015).
8. Beggs, S., Torsney, C., Drew, L. J. & Fitzgerald, M. The postnatal reorganization of
primary afferent input and dorsal horn cell receptive fields in the rat spinal cord is an
activity-dependent process. *Eur. J. Neurosci.* **16**, 1249–1258 (2002).
9. Granmo, M., Petersson, P. & Schouenborg, J. Action-Based Body Maps in the Spinal
Cord Emerge from a Transitory Floating Organization. *J. Neurosci.* **28**, 5494–5503
(2008).
10. Salter, M. W. & Stevens, B. Microglia emerge as central players in brain disease. *Nat.*
Med. **23**, 1018–1027 (2017).
11. Paolicelli, R. C. *et al.* Synaptic pruning by microglia is necessary for normal brain
development. *Science (80-.).* **333**, 1456–1458 (2011).
12. Schafer, D. P. *et al.* Microglia sculpt postnatal neural circuits in an activity and
complement-dependent manner. *Neuron* **74**, 691–705 (2012).
13. Gunner, G. *et al.* Sensory lesioning induces microglial synapse elimination via
ADAM10 and fractalkine signaling. *Nat. Neurosci.* **22**, 1075–1088 (2019).
14. Milinkeviciute, G. *et al.* Microglia Regulate Pruning of Specialized Synapses in the
Auditory Brainstem. *Front. Neural Circuits* **13**, 1–19 (2019).
15. Lui, H. *et al.* Progranulin Deficiency Promotes Circuit-Specific Synaptic Pruning by
Microglia via Complement Activation. *Cell* **165**, 921–935 (2016).
16. Vainchtein, I. D. *et al.* Astrocyte-derived interleukin-33 promotes microglial synapse
engulfment and neural circuit development. *Science (80-.).* **359**, 1269–1273 (2018).
17. Vukojicic, A. *et al.* The classical complement pathway mediates microglia-dependent
remodeling of spinal motor circuits during development and in spinal muscular
atrophy. *Cell Rep.* **in press**, 3087–3100 (2019).
18. Zusso, M. *et al.* Regulation of postnatal forebrain amoeboid microglial cell proliferation
and development by the transcription factor runx1. *J. Neurosci.* **32**, 11285–11298
(2012).
19. Matcovitch-Natan, O. *et al.* Microglia development follows a stepwise program to
regulate brain homeostasis. *Science (80-.).* **353**, aad8670–aad8670 (2016).
20. Hammond, T. R. *et al.* Single-Cell RNA Sequencing of Microglia throughout the
Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes.
Immunity **50**, 253-271.e6 (2019).
21. Schwarz, J. M., Sholar, P. W. & Bilbo, S. D. Sex differences in microglial colonization
of the developing rat brain. *J. Neurochem.* **120**, 948–963 (2012).
22. De Biase, L. M. *et al.* Local Cues Establish and Maintain Region-Specific Phenotypes
of Basal Ganglia Microglia. *Neuron* **95**, 341-356.e6 (2017).
23. Ayata, P. *et al.* Epigenetic regulation of brain region-specific microglia clearance

activity. *Nat. Neurosci.* **21**, 1049–1060 (2018).

24. Chamessian, A. *et al.* Is Optogenetic Activation of Vglut1-Positive Aβ Low-Threshold
Mechanoreceptors Sufficient to Induce Tactile Allodynia in Mice after Nerve Injury? *J.*
Neurosci. **39**, 6202–6215 (2019).

25. Batti, L. *et al.* TMEM16F Regulates Spinal Microglial Function in Neuropathic Pain
States. *Cell Rep.* **15**, 2608–2615 (2016).

26. Miyazaki, T., Fukaya, M., Shimizu, H. & Watanabe, M. Subtype switching of vesicular
glutamate transporters at parallel fibre-Purkinje cell synapses in developing mouse
cerebellum. *Eur. J. Neurosci.* **17**, 2563–2572 (2003).

27. Minelli, A., Edwards, R. H., Manzoni, T. & Conti, F. Postnatal development of the
glutamate vesicular transporter VGLUT1 in rat cerebral cortex. *Dev. Brain Res.* **140**,
309–314 (2003).

28. Parkhurst, C. N. *et al.* Microglia Promote Learning-Dependent Synapse Formation
through Brain-Derived Neurotrophic Factor. *Cell* **155**, 1596–1609 (2013).

29. Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain. *Nat. Neurosci.* **13**, 133–140 (2010).

30. Taylor-Clark, T. E. *et al.* Thy1.2 YFP-16 Transgenic Mouse Labels a Subset of Large-
Diameter Sensory Neurons that Lack TRPV1 Expression. *PLoS One* **10**, e0119538
(2015).

31. Feng, G. *et al.* Imaging neuronal subsets in transgenic mice expressing multiple
spectral variants of GFP. *Neuron* **28**, 41–51 (2000).

32. Lee, K. Y., Ratté, S. & Prescott, S. A. Excitatory neurons are more disinhibited than
inhibitory neurons by chloride dysregulation in the spinal dorsal horn. *Elife* **8**, (2019).

33. Abraira, V. E. *et al.* The Cellular and Synaptic Architecture of the Mechanosensory
Dorsal Horn. *Cell* **168**, 295-310.e19 (2017).

34. Todd, A. J. Neuronal circuitry for pain processing in the dorsal horn. *Nat. Rev.*
Neurosci. **11**, 823–836 (2010).

35. Todd, A. J. *et al.* The expression of vesicular glutamate transporters VGLUT1 and
VGLUT2 in neurochemically defined axonal populations in the rat spinal cord with
emphasis on the dorsal horn. *Eur. J. Neurosci.* **17**, 13–27 (2003).

36. Fitzgerald, M. & Gibson, S. The postnatal physiological and neurochemical
development of peripheral sensory C fibres. *Neuroscience* **13**, 933–944 (1984).

37. Baccei, M. L., Bardoni, R. & Fitzgerald, M. Development of nociceptive synaptic inputs
to the neonatal rat dorsal horn: Glutamate release by capsaicin and menthol. *J.*
Physiol. **549**, 231–242 (2003).

38. Koch, S. C. & Fitzgerald, M. Activity-dependent development of tactile and nociceptive
spinal cord circuits. *Ann. N. Y. Acad. Sci.* **1279**, 97–102 (2013).

39. Cleveland, L. *et al.* Systematic Review of Skin-to-Skin Care for Full-Term, Healthy
Newborns. *JOGNN - J. Obstet. Gynecol. Neonatal Nurs.* **46**, 857–869 (2017).

40. Perez-Pouchoulen, M., VanRyzin, J. W. & McCarthy, M. M. Morphological and
phagocytic profile of microglia in the developing rat cerebellum. *eNeuro* **2**, (2015).

41. VanRyzin, J. W. *et al.* Microglial Phagocytosis of Newborn Cells Is Induced by
Endocannabinoids and Sculpts Sex Differences in Juvenile Rat Social Play. *Neuron*
102, 435-449.e6 (2019).

42. Salter, E. W. *et al.* Complement C3-dependent glutamatergic synapse elimination in
the developing hippocampus is region- and synapse-specific. *bioRxiv*
2020.05.20.106930 (2020) doi:10.1101/2020.05.20.106930.

43. Zieglgänsberger, W. & Herz, A. Changes of cutaneous receptive fields of spino-
cervical-tract neurones and other dorsal horn neurones by microelectrophoretically
administered amino acids. *Exp. Brain Res.* **13**, 111–126 (1971).

44. Takeda, M., Tanimoto, T. & Matsumoto, S. Change in mechanical receptive field
properties induced by GABA(A) receptor activation in the trigeminal spinal nucleus
caudalis neurons in rats. *Exp. Brain Res.* **134**, 409–416 (2000).

45. Siddoway, B., Hou, H. & Xia, H. Molecular mechanisms of homeostatic synaptic
downscaling. *Neuropharmacology* **78**, 38–44 (2014).

46. Schreyer, D. J. & Jones, E. G. Growth and target finding by axons of the corticospinal
tract in prenatal and postnatal rats. *Neuroscience* **7**, 1837–1853 (1982).
47. Jackman, A. & Fitzgerald, M. Development of peripheral hindlimb and central spinal
cord innervation by subpopulations of dorsal root ganglion cells in the embryonic rat.
J. Comp. Neurol. **418**, 281–298 (2000).
48. Wake, H., Moorhouse, A. J., Jinno, S., Kohsaka, S. & Nabekura, J. Resting microglia
directly monitor the functional state of synapses in vivo and determine the fate of
ischemic terminals. *J. Neurosci.* **29**, 3974–3980 (2009).
49. Liu, Y. U. *et al.* Neuronal network activity controls microglial process surveillance in
awake mice via norepinephrine signaling. *Nat. Neurosci.* **22**, 1771–1781 (2019).
50. Li, Y., Du, X.-F., Liu, C.-S., Wen, Z.-L. & Du, J.-L. Reciprocal regulation between
resting microglial dynamics and neuronal activity in vivo. *Dev. Cell* **23**, 1189–202
(2012).
51. Badimon, A. *et al.* Negative feedback control of neuronal activity by microglia. *Nature*
(2020) doi:10.1038/s41586-020-2777-8.
52. Lehrman, E. K. *et al.* CD47 Protects Synapses from Excess Microglia-Mediated
Pruning during Development. *Neuron* **100**, 120-134.e6 (2018).
53. Scott-Hewitt, N. *et al.* Local externalization of phosphatidylserine mediates
developmental synaptic pruning by microglia. *EMBO J.* **39**, 2020.04.24.059584
(2020).
54. Walker, S. M., Tochiki, K. K. & Fitzgerald, M. Hindpaw incision in early life increases
the hyperalgesic response to repeat surgical injury: Critical period and dependence on
initial afferent activity. *Pain* **147**, 99–106 (2009).
55. Beggs, S., Currie, G., Salter, M. W., Fitzgerald, M. & Walker, S. M. Priming of adult
pain responses by neonatal pain experience: Maintenance by central neuroimmune
activity. *Brain* **135**, 404–417 (2012).
56. Pitulescu, M. E., Schmidt, I., Benedito, R. & Adams, R. H. Inducible gene targeting in
the neonatal vasculature and analysis of retinal angiogenesis in mice. *Nature*
Protocols vol. 5 1518–1534 (2010).
57. Ollion, J., Cochennec, J., Loll, F., Escudé, C. & Boudier, T. TANGO: a generic tool for
high-throughput 3D image analysis for studying nuclear organization. *Bioinformatics*
29, 1840–1841 (2013).
58. Linkert, M. *et al.* Metadata matters: access to image data in the real world. *J. Cell Biol.*
189, 777–782 (2010).
59. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*
Methods **9**, 676–682 (2012).
60. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
image analysis. *Nat. Methods* **9**, 671–675 (2012).
61. Duan, B. *et al.* Identification of spinal circuits transmitting and gating mechanical pain.
Cell **159**, 1417–1432 (2014).
62. Bonin, R. P., Bories, C. & De Koninck, Y. A Simplified Up-Down Method (SUDO) for
Measuring Mechanical Nociception in Rodents Using von Frey Filaments. *Mol. Pain*
10, 1744-8069-10–26 (2014).
63. Ho, J., Tumkaya, T., Aryal, S., Choi, H. & Claridge-Chang, A. Moving beyond P
values: data analysis with estimation graphics. *Nat. Methods* **16**, 565–566 (2019).
64. Chernick, M. R. *Bootstrap Methods*. (John Wiley & Sons, Inc., 2007).
doi:10.1002/9780470192573.
880

Supplementary figures

884   **Figure 1—figure supplement 1. Microglial density changes over postnatal development**

a. Representative images showing an increase in tdT-expression in the dorsal horn with age. Dashed white line indicates border between lamina II and lamina III, white arrow heads indicate tdT-fibres in the superficial laminae I-II. Scale bar = 200µm.

b. Microglial density increases over the postnatal period in both LI-II. P3 vs. P0: mean difference 1.53 [95% CI 0.97, 2.08]; P7 vs. P0: mean difference 3.61 [95% CI 2.58, 4.42]; P10 vs. P0: mean difference 4.46 [95% CI 3.91, 4.98]; P17 vs. P0: mean difference 4.35 [95% CI 3.60, 5.06]; P28 vs. P0: mean difference 3.61 [95% CI 2.79, 4.49]. Field of view (FOV) = 245µm x 65µm.

c. Microglial density increases over the postnatal period in LIII-IV. P3 vs. P0: mean difference 1.83 [95% CI 1.21, 2.42]; P7 vs. P0: mean difference 2.79 [95% CI 1.80, 3.54]; P10 vs. P0: mean difference 3.58 [95% CI 3.09, 4.06]; P17 vs. P0: mean difference 4.24 [95% CI 3.60, 4.81]; P28 vs. P0: mean difference 2.96 [95% CI 2.35, 3.51]. Field of view (FOV) = 245µm x 65µm.

d. Microglia density in rats increases significantly between P3 and P7, with $P < 0:0001$ for all ages compared to P3.

e. Representative images of microglia cells in the dorsal horn (LI-III) stained for Iba1 (green) at postnatal days (P)3, P7, P10, P17. Scale bar = 50 µm.

f. Maximum projected confocal images from the dorsal horn (LI-III) at P3 with representative staining for microglia (Iba1, cyan) and neurons (NeuN, red). Scale bar = 50µm.

g. P7 spinal cord hemisection stained with neuronal marker NeuN (red), white boxes A, B, C indicate the approximate area of laminae I-III, lamina V, and laminae VIII-IX used for analysis.

h. Ratio of neurons per microglia significantly decreases with age in laminae I-II, lamina III, lamina V, and laminae VIII-IX.

i. Representative confocal image of microglial A-fibre engulfment in the spinal dorsal horn of P3 VGLUT1-tdt mice, stained for microglia (Iba1, cyan), microglial lysosomes (CD68, white) and endogenously fluorescent A-fibres (tdTomato, red). Scale bar = 20µm. White inset box show location of higher magnification panels in **i.**

j. High magnification image of microglial A-fibre engulfment in **h** stained for microglia (Iba1, cyan), microglial lysosomes (CD68, white) and endogenously fluorescent A-fibres (tdTomato, red). Cross-hairs show position of the xz and yz side-view panels. Scale bar = 5µm.

k. Further example of microglial A-fibre engulfment, colours and scale as indicated in **i.**

N-numbers as indicated. Black and colourful data points indicate females and males respectively.

917
918
Figure 1—figure supplement 2. Apoptotic cell clearance over the postnatal period
a. Representative confocal image from the P3 dorsal horn of a microglia cell engulfing an apoptotic
cell with a phagocytic cup. Cross-hairs show position of the xz and yz side-view panels. Microglial cell
body is outside of the visible xy-plane. Image shows representative staining for Iba1 (cyan), Cas3
(red), DAPI (blue) at P3. Scale bar = 20µm.
b. Apoptotic cell numbers decrease significantly by P10, with P < 0.001 for P10 and P17 compared to
P3.
c. The percentage of Cas3 positive cells colocalising with microglia significantly increases with age,
with P < 0.05 for all ages compared to P3.
d. Representative image of microglia cell in the dorsal horn with phagocytic cup indicated by white
arrowhead. Image shows representative staining for Iba1 (green), CD68 (red), DAPI (blue) at P3.
Scale bar = 10µm.
e. Phagocytic cup count per microglia decreases significantly between P3 and P7, (P < 0.01 for all
ages compared to P3).
N-numbers as indicated. Black and red data points indicate females and males respectively.

Figure 2—figure supplement 1. Superficial A-fibres form synapses with pre- and post-synaptic densities.

a. TOP: Low magnification image, with white box indicating the location of bottom panel image. Scale bar = 50μm. Bottom: Representative maximum projection of super-resolution image from the superficial dorsal horn laminae showing colocalization of Thy1-GFP labelled A-fibre (yellow) with presynaptic VGLUT1 (magenta) and postsynaptic HOMER (cyan). Scale bar = 5μm.

b, c. Super-resolution images of boxed areas in **a** (bottom panel) with xy, xz, and yz views showing colocalization of Thy1-GFP labelled A-fibre (yellow), presynaptic VGLUT1 (magenta), and postsynaptic HOMER (cyan), as well as surface rendered view on the right. White arrowhead in **b** indicate a Thy1-GFP and VGLUT1 positive presynaptic bouton without postsynaptic HOMER. Scale bars = 1μm.

d. Homer density does not seem affected in *Tmem16f* cKO animals.

e. VGLUT1 density seems higher in *Tmem16f* cKO animals, consistent with results in Fig. 3.

f. Thy1-positive synapse density (as defined by colocalization of Thy1-GFP, VGLUT1, and HOMER) seems to increase in *Tmem16f* cKO animals.

g. Total synapse number (as defined by colocalization of VGLUT1 and HOMER puncta) seems unchanged in *Tmem16f* cKO animals.

FOV= 94μm x 94μm for **d-g**. N-numbers as indicated.

959
960 **Figure 3—figure supplement 1.** *Tmem16f* **cKO does not change vF reflex threshold**
961 Unpaired mean difference -0.29 [95% CI -0.85, 0.39]. N-numbers as indicated. Black and magenta
962 data points indicate females and males respectively.
963
964
965
966

967
Figure 4—figure supplement 1. Recording depth and spontaneous activity in control and
***Tmem16f* cKO animals.**
a Spontaneous activity decreased in adapting WDR neurons from *Tmem16f* cKO animals, unpaired
mean difference -0.072 [95% CI -0.121, -0.039].
b. Spontaneous activity was unchanged in non-adapting WDR neurons from *Tmem16f* cKO animals,
unpaired mean difference 0.042 [95.00% CI -0.092, 0.25].
c. Recording depth of all neurons in control and *Tmem16f* cKO animals, unpaired mean difference -
37.24 [95.00% CI -78.02,   5.73].
N-numbers as indicated. Black and coloured data points indicate females and males.
977
978