# Respiration, heartbeat, and conscious tactile perception

Abbreviated title: Respiration, heartbeat, and tactile perception

Martin Grund[1], Esra Al[1,2], Marc Pabst[1], Alice Dabbagh[1,3], Tilman Stephani[1,4], Till Nierhaus[1,5], and Arno Villringer[1,2]

[1]Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, 04103 Leipzig, Germany

[2]MindBrainBody Institute, Berlin School of Mind and Brain, Charité – Universitätsmedizin Berlin and Humboldt-Universität zu Berlin, 10099 Berlin, Germany

[3]Pain Perception Group, Max Planck Institute for Human Cognitive and Brain Sciences, 04103 Leipzig, Germany

[4]International Max Planck Research School NeuroCom, 04103 Leipzig, Germany

[5]Neurocomputation and Neuroimaging Unit, Department of Education and Psychology, Freie Universität Berlin, 14195 Berlin, Germany

**Corresponding author:** Martin Grund, Max Planck Institute for Human Cognitive and Brain Sciences, Stephanstr. 1A, 04103 Leipzig, Germany (mgrund@cbs.mpg.de)

**Number of pages:** 43

**Number of figures:** 7

**Number of words for** *Abstract*: 243; *Introduction*: 649; *Discussion*: 1375

**Conflict of Interest:** The authors declare no competing financial interest.

**Acknowledgements:** The research was funded by the Max Planck Society. We thank Sylvia Stash for her data acquisition support; Mina Jamshidi Idaji for data analysis advice; and Heike Schmidt-Duderstedt for preparing the figures for publication.

## Abstract

Cardiac activity has been shown to interact with conscious tactile perception: Detecting near-threshold tactile stimuli is more likely during diastole than systole and heart slowing is more pronounced for detected compared to undetected stimuli. Here, we investigated how cardiac cycle effects on conscious tactile perception relate to respiration given the natural coupling of these two dominant body rhythms. Forty-one healthy participants had to report conscious perception of weak electrical pulses applied to the left index finger (yes/no) and confidence about their yes/no-decision (unconfident/confident) while electrocardiography (ECG), respiratory activity (chest circumference), and finger pulse oximetry were recorded. We confirmed the previous findings of higher tactile detection rate during diastole and unimodal distribution of hits in diastole, more specifically, we found this only when participants were confident about their detection decision. Lowest tactile detection rate occurred 250-300 ms after the R-peak corresponding to pulse-wave onsets in the finger. Inspiration was locked to tactile stimulation, and this was more consistent in hits than misses. Respiratory cycles accompanying misses were longer as compared to hits and correct rejections. Cardiac cycle effects on conscious tactile perception interact with decision confidence and coincide with pulse-wave arrival, which suggests the involvement of higher cognitive processing in this phenomenon possibly related to predictive coding. The more consistent phase-locking of inspiration with stimulus onsets for hits than misses is in line with previous reports of phase-locked inspiration

47  to cognitive task onsets which were interpreted as tuning the sensory system for

48  incoming information.

50  **Keywords:** respiration, cardiac cycle, interoception, tactile perception,

51  electrocardiogram, pulse oximetry

## Significance statement

Mechanistic studies on perception and cognition tend to focus on the brain neglecting contributions of the body. Here, we investigated how respiration and heartbeat influence tactile perception: We show that inspiration locked to expected stimulus onsets optimizes detection task performance and that tactile detection varies across the heart cycle with a minimum 250-300 milliseconds after heart contraction, when the pulse reaches the finger. Lower detection was associated with reduced confidence ratings, indicating – together with our previous finding of unchanged early ERPs - that this effect is not a peripheral physiological artifact but a result of higher cognitive processes that model the internal state of our body, make predictions to guide behavior, and might also tune respiration to serve the task.

## Introduction

Our body senses signals from the outer world (exteroception), but also visceral signals from inside the body (interoception) and it has been shown that these two continuous types of perception interact (Critchley and Harrison, 2013; Critchley and Garfinkel, 2015; Babo-Rebelo et al., 2016; Azzalini et al., 2019). For example, we have recently shown that tactile perception interacts with cardiac activity as conscious detection of near-threshold stimuli was more likely towards the end of the cardiac cycle (Motyka et al., 2019; Al et al., 2020) and was followed by a more pronounced deceleration of heart rate as compared to missed stimuli (Motyka et al., 2019). In line with increased detection during later cardiac phases (diastole), late (P300) cortical somatosensory evoked potentials (SEPs) were also higher during diastole as compared to systole (Al et al., 2020). A similar cardiac phase-dependency has also been revealed for visual sampling: microsaccades and saccades were more likely during systole, whereas fixations and blinks during diastole (Ohl et al., 2016; Galvez-Pol et al., 2020). Following an interoceptive predictive coding account, the very same brain model that predicts cardiac-associated bodily changes and suppresses their access to consciousness might suppress perception of external stimuli which coincide with those changes and modulate the generation of actions (Seth and Friston, 2016; Kunzendorf et al., 2019; Allen et al., 2019). Yet, which bodily changes are the main driver for the cardiac-related perceptual suppression effect remain to be determined.

Another dominant body rhythm that can even be regulated intentionally in contrast to cardiac activity is the respiration rhythm (Azzalini et al., 2019). Also for respiration, which naturally drives and is driven by cardiac activity (Kralemann et al.,

2013; Dick et al., 2014), phase-dependency of behavior and perception has been

reported. For instance, self-initiated actions were more likely during expiration,

whereas externally-triggered actions showed no correlation with the respiration

phase (Park et al., 2020). Furthermore, inspiration onsets were reported to lock to

task onsets which resulted in greater task-related brain activity and increased task

performance for visuospatial perception, memory encoding and retrieval, and fearful

face detection (Huijbers et al., 2014; Zelano et al., 2016; Perl et al., 2019). Locked

inspiration was interpreted as tuning the sensory system for upcoming information

(Perl et al., 2019). Thus, inspiration might also be beneficial for conscious perception

and could be timed to paradigms instead of modelled as noise (heartbeat) within an

interoceptive predictive coding framework. While cardiac activity and respiration are

closely interdependent, it remains unclear how they jointly shape perceptual

processes.

Our present study combined the observation of cardiac and respiratory activity

with a paradigm that asked participants to report (a) conscious perception of weak

electrical pulses applied to their left index finger and (b) their decision confidence.

Decision confidence was assessed to identify the potential role of metacognition in

cardiac cycle effects. As we have previously shown that greater tactile detection

during diastole corresponded to increased perceptual sensitivity and not to a more

liberal response criterion (Al et al., 2020), we expected the cardiac cycle effect not to

be a side-effect of unconfident perceptual decisions. Recently, afferent fibers in the

finger have been reported to be modulated by cardiac pressure changes which the

brain has to ignore or filter out (Macefield, 2003). Thus, we measured pulse oximetry

to investigate whether peripheral physiological changes in the finger caused by the

blood pulse wave coincided with lower tactile detection during systole. Furthermore,

we tried to capture early SEPs at the upper arm to rule out differences in (peripheral)

SEP amplitudes as explanation for altered conscious tactile perception across

cardiac or respiratory cycles.

This study setup was intended to address the following research questions:

-   Does the interaction of cardiac activity and conscious tactile perception

depend on decision confidence?

-   How is the relationship between decreased tactile detection and the kinetics

of the pulse wave in the finger?

-   Does conscious tactile perception vary across the respiratory cycle?

## Methods

**Participants**

Forty-one healthy humans (21 women, mean age = 25.5, age range: 19-37) participated in the study. Participants were predominantly right-handed with a mean laterality index of 90, *SD* = 17 (Oldfield, 1971). For four participants, the mean laterality index was not available.

**Ethics statement**

All participants provided an informed consent. The experimental procedure and physiological measurements were approved by the ethics commission at the medical faculty of the University of Leipzig.

**Experimental design and statistical analysis**

The experiment was designed to capture tactile detection of near-threshold stimuli (50% detection) and trials without stimuli (0% detection) across the cardiac and respiratory cycle. This resulted in three main stimulus-response conditions: (a) correct rejections of trials without stimulation, (b) undetected (misses), and (c) detected near-threshold stimuli (hits). False alarms (yes-responses) during trials without stimulation were very rare (mean FAR = 6%, *SD* = 6%) and thus not further analyzed. Additionally, participants reported their decision confidence which allowed us to split trials by confidence.

We applied circular statistics to investigate whether conscious tactile perception was uniformly distributed across the cardiac and respiratory cycle or showed unimodal patterns. For each stimulus onset, the temporal distances to the

preceding R-peak and inspiration onset were calculated and put in relation to its current cardiac and respiratory cycle duration measured in degrees. Following for each participant, these angles were averaged for hits, misses, and correct rejections. For each stimulus-response condition, the resulting distributions of mean angles were tested across participants with the Rayleigh test for uniformity from the R package "circular" (Version 0.4-93). The application of circular statistics had two advantages: First, it accounted for cardiac and respiratory cycle duration variance within and between participants. Second, it allowed us to determine phases when detection differed without having to rely on arbitrary binning. However, it assumed that the different phases of the cardiac and respiratory cycle behave proportionally the same when cycle duration changes. That is why we complemented the circular statistics with a binning analysis that investigated the near-threshold detection rate for fixed time intervals relative to the preceding R-peak and inspiration onset.

In repeated-measures ANOVAs, Greenhouse-Geisser correction was used to adjust for the lack of sphericity. Post-hoc *t*-tests p-values were corrected for multiple comparisons with a false discovery rate (FDR) of 5% (Benjamini and Hochberg, 1995).

**Data and code availability**

The code to run and analyze the experiment is available at http://github.com/grundm/respirationCA. The behavioral and physiological data (electrocardiogram, respiration, and oximetry) can be shared by the corresponding author upon request if data privacy can be guaranteed.

**Stimuli and apparatus**

Somatosensory stimulation was delivered via steel wire ring electrodes to the left index finger with a constant current stimulator (DS5; Digitimer, United Kingdom). The anode was placed on the middle phalanx and the cathode on the proximal phalanx. The stimuli were single square-wave pulses with a duration of 0.2 ms and an individually-assessed near-threshold intensity (mean intensity = 1.95 mA, range: 0.76-3.17 mA). The stimulator was controlled by the waveform generator NI USB-6343 (National Instruments, Austin, Texas) and custom MATLAB scripts using the Data Acquisition Toolbox (The MathWorks Inc., Natick, Massachusetts, USA).

**Behavioral paradigm**

Participants had to report whether they perceived an electrical pulse and whether this yes/no-decision was confident or not. The experiment was separated into four blocks. Each block consisted of 150 trials. Participants received a near-threshold stimulus in 100 trials (mean intensity = 1.96 mA, range: 0.76-3.22 mA). In 50 trials, there was no stimulation (33% catch trials). The order of near-threshold and catch trials was pseudo-randomized for each block and participant. In total, there were 400 near-threshold and 200 catch trials.

Each trial started with a black fixation cross (black "+") for a counterbalanced duration of 1.0-2.0 s (Figure 1). It was followed by a salmon-colored fixation cross (0.62 s) to cue the stimulation at 0.5 s after the cue onset. With the cue offset, the participants had to report the detection of a tactile stimulus (yes/no). After the yes/no-button press, a pause screen was displayed for 0.3 s, before the participants were asked to report their decision confidence (confident/unconfident). With pressing the

button for "confident" or "unconfident", the new trial started. For both reports, the

maximum response time was 1.5 s. Thus, the trial duration was limited to 5.8 s.

Participants indicated their perception and decision confidence with the right

index finger on a two-button box. The buttons were arranged vertically. The four

possible button response mappings were counterbalanced across participants, so

that the top button could be assigned to "yes" or "no", and "confident" or

"unconfident" respectively for one participant.

**Figure 1.** Experimental procedure and physiological parameters visualized for one exemplary trial. The

tiles represent the participant's visual display and the times given below indicate the presentation

duration. The near-threshold electrical finger nerve stimulation was always 0.5 s after the cue onset

(salmon-colored fixation cross). Here only one of four button response mappings is displayed (Y = yes;

N = no; U = unconfident; C = confident). In total, 400 near-threshold trials and 200 trials without

stimulation (33% catch trials) were presented in a randomized order. Exemplary traces of

electrocardiogram (ECG), finger pulse oximetry, and respiration belt below the trial procedure indicate

that stimulus detection was analyzed relative to cardiac and respiratory cycles (0-360°).

Prior to the experiment, participants were familiarized with the electrical finger

nerve stimulation and an automatic threshold assessment was performed in order to

determine the stimulus intensity corresponding to the sensory threshold (50%

detection rate). The threshold assessment entailed an up-and-down procedure

followed by a Bayesian approach (psi method) from the Palamedes Toolbox

(Kingdom and Prins, 2009) and closed with a test block. The visual display of the trials

in the threshold assessment was similar to the trials in the experimental block (Figure

1) but without the confidence rating and a shorter fixed intertrial interval (0.5 s). If

necessary, the threshold assessment was repeated before each block to ensure a

detection rate of 50% throughout the whole experiment. The experimental procedure

was controlled by custom MATLAB scripts using the Psychophysics Toolbox (Kleiner

et al., 2007).

**Electrocardiogram acquisition**

The electrocardiogram (ECG) was recorded with the BrainAmp ExG (Brain Products,

Gilching, Germany) between two adhesive electrodes that were placed on the

sternum and just below the heart on the left side of the thorax. A ground electrode

was placed on the acromion of the right shoulder. The sampling frequency was 5000

Hz for 39 participants. Two participants were recorded with 1000 Hz.

**Respiration acquisition**

Respiration was measured with a respiration belt (BrainAmp ExG; Brain Products, Gilching, Germany). The belt with a pressure-sensitive cushion was placed at the largest expansion of the abdomen during inspiration. The sampling frequency was 5000 Hz for 39 participants. Two participants were recorded with 1000 Hz.

**Peripheral nerve activity acquisition**

To examine the possibility to measure somatosensory evoked potentials (SEP) of peripheral nerve activity in response to near-threshold finger stimulation, two surface electrodes were placed with a distance of 2 cm at the left upper inner arm (below the biceps brachii) above the pathway of the median nerve in a sub-sample of 12 participants. The signal was recorded with a sampling rate of 5000 Hz, low-pass filtered at 1000 Hz, using a bipolar electrode montage (BrainAmp ExG; Brain Products, Gilching, Germany).

**Oximetry acquisition**

The pulse oximetry was recorded with a finger clip $SpO_2$ transducer at the left middle finger at 50 Hz (OXI100C and MP150; BIOPAC Systems Inc., Goleta, California, USA).

**Behavioral data analysis**

The behavioral data was analyzed with R 4.0.3 in RStudio 1.3.10923. First, trials were filtered for detection and confidence responses within the maximum response time of 1.5 s. Second, only blocks were considered with a near-threshold hit rate at least five percentage points above the false alarm rate. These resulted in 37 participants

with 4 valid blocks, 2 participants with 3 valid blocks and 2 participants with 2 valid blocks. The frequencies of the response "confident" for correct rejections, misses, and hits were compared with paired *t*-tests. Furthermore, the detection and confidence response times and resulting trial durations were compared between correct rejections, misses, and hits with paired *t*-tests. The response times for hits and miss were additionally compared between confident and unconfident near-threshold trials.

**Cardiac data analysis**

ECG data was preprocessed with Kubios (Version 2.2) to detect R-peaks. For two participants, the first four and the first twenty-two trials respectively had to be excluded due to a delayed start of the ECG recording. Additionally, one block of one participant and two blocks of another participant were excluded due to low ECG data quality.

First for correct rejections, misses, and hits, the circular distribution within the cardiac cycle was assessed with the Rayleigh test of uniformity. Additionally, this analysis was repeated for confident and unconfident hits and misses.

Second, instead of the relative position within the cardiac cycle, near-threshold trials were assigned to four time intervals based on their temporal distance from the previous R-peak (0-200 ms, 200-400 ms, 400-600 ms, and 600-900 ms). The last interval had a duration of 300 ms to account for interindividual differences of the interbeat interval and aggregate a similar number of trials to the preceding intervals. Then, detection rates for confident and unconfident decisions were calculated in each

time interval and compared with a two-way repeated measures ANOVA and post-hoc

*t*-tests.

Third, we analyzed the interbeat intervals in the course of a trial between the

stimulus-response conditions. For this, two interbeat intervals before, one during, and

two after the stimulus onset were selected and compared with a two-way repeated

measures ANOVA and post-hoc *t*-tests.

**Oximetry data analysis**

Oximetry data was analyzed with custom MATLAB scripts to detect the pulse wave

peaks with a minimum peak distance based on 140 heartbeats per minute and a

minimum peak prominence equal to a tenth of the data range in each block. Pulse

cycles with a duration 1.5 times the median duration of the respective block were

excluded from further processing. In R, the pulse wave cycle data was merged with

the behavioral data to apply the same exclusion criteria and the Rayleigh test of

uniformity. Finally, pulse wave peaks were located in the cardiac cycle to assess the

duration since the previous R-peak (pulse wave transit time, PWTT) and its relative

position in degree within the cardiac cycle.

**Respiration data analysis**

After visual inspection of the respiration traces, respiratory cycle detection was

performed following the procedure by Power et al. (2020). First, outliers were replaced

in a moving 1-s window with linearly interpolated values based on neighboring, non-

outlier values. Local outliers were defined as values of more than three local scaled

median absolute deviations (MAD) away from the local median within a 1-s window

(Power et al., 2020). MAD was chosen for its robustness compared to standard

deviation which is more affected by extreme values. Subsequently, the data was

smoothed with a 1-s window Savitzky-Golay filter (Savitzky and Golay, 1964) to

facilitate peak detection. Traces were then z-scored and inverted to identify local

minima (inspiration onsets) with the MATLAB findpeaks function. Local minima had

to be at least 2 s apart with a minimum prominence of 0.9 times the interquartile range

of the z-scored data. Respiration cycles were defined as the interval from one

inspiration onset to the next inspiration onset. For each participant, respiration cycles

with more than two times the median cycle duration were excluded from further

analysis.

For each stimulus-response condition and participant, the mean angle

direction of stimulus onsets within the respiration cycle and their circular variance

across trials were calculated. The distribution of mean angles of each stimulus-

response condition was tested for uniformity with the Rayleigh test. Circular variance

was defined as $V = 1\text{-}R$, where $R$ is the mean resultant length of each stimulus-

response condition and participant with values between 0 and 1. Differences in

circular variances between stimulus-response conditions were assessed with paired

$t$-tests.

Furthermore, we investigated whether participants gradually aligned their

respiration to the stimulus onset in the beginning of the experiment. For the first 30

trials, the difference between each trial's stimulus onset angle and the mean angle

within the first block was determined ("diff_angle2mean"). The trial angle difference

from the mean was used as a dependent variable in a random-intercept linear

regression based on maximum likelihood estimation with trial number as independent

variable: "diff_angle2mean ~ 1 + trial + (1|participant)". The fit of this model was

compared with a random-intercept only model "diff_angle2mean ~ 1 + (1|participant)"

in a $\chi^2$-test to assess the effect of trial number on the angle difference. This analysis

included only the 37 participants with a valid first block and excluded trials with false

alarms.

Lastly, we compared the respiratory cycle duration between stimulus-

response conditions by performing a one-way repeated-measures ANOVA and post-

hoc $t$-tests.

**Phase-locking analysis between cardiac and respiratory activity**

332 The n:m ($n, m \in N$) synchronization (Lachaux et al., 1999) was calculated in an inter-

333 trial setting for the stimulation onset as the following:

$$PLV_{cross} = \left| \frac{1}{n} \sum_{i=1}^{n} e^{j\phi_i} \right|$$

$$\phi_i = n\, \Phi_{i,resp} - m\, \Phi_{i,ecg}$$

where $\Phi_{i,resp}$ and $\Phi_{i,ecg}$ were the stimulation onset angles for the i-th trial within the

respiratory (*resp*) and cardiac cycle (*ecg*), and *j* was the imaginary number. While $m =$

*1* was chosen for all participants, values for $n$ were selected by calculating the ratio of

the cardiac and respiratory frequency rounded to the nearest integer. The frequencies

were estimated based on the mean cardiac and respiratory cycle durations at stimulus

onset. The inter-trial n:m synchronization at stimulation onset can provide information

about the extent to which the weighted phase difference of the two signals stays

identical over trials. The calculated phase-locking value (PLV) lies between zero and

one, with zero indicating no inter-trial coupling and one showing a constant weighted

phase-difference of the two signals at the stimulation time.

**Somatosensory evoked potential analysis**

For the twelve participants with peripheral nerve recordings, stimulation artefacts were removed with a cubic monotonous Hermite spline interpolation from -2 s until 4 s relative to the trigger. Next, a 70-Hz high-pass filter was applied (4th order Butterworth filter applied forwards and backwards) and the data was epoched -100 ms to 100 ms relative to the trigger with -50 ms to -2 ms as baseline correction. Subsequently, epochs were averaged across for valid trials (yes/no and confidence response within maximum response time) with near-threshold stimuli and without stimulations.

## Results

### Detection and confidence responses

Participants ($N$ = 41) detected on average 51% of near-threshold stimuli ($SD$ = 16%) and correctly rejected 94% of catch trials without stimulation ($SD$ = 6%). On average, 188 catch trials (range: 93-200) and 375 near-threshold trials (range: 191-400) were observed. Participants reported to be "confident" about their yes/no-decision in 88% of the correct rejections ($SD$ = 13%), in 71% of the misses ($SD$ = 21%), and in 62% of the hits ($SD$ = 18%). The confidence rate differed significantly between all conditions in paired $t$-tests (CR vs. miss: $p$ = 3 * $10^{-8}$; CR vs. hit: $p$ = 1 * $10^{-9}$; miss vs. hit: $p$ = 0.019). In total, we observed on average 184 misses (range: 58-303), 192 hits (range: 59-302), 177 correct rejections (range: 72-198) and 11 false alarms (range: 0-36). Two-third of the participants (27) had less than ten false alarms and four participants had zero false alarms. Due to zero or very few observations, false alarms were not further analyzed.

In near-threshold trials, participants reported their yes/no-decision later than for correct rejections (mean ± SD: $RT_{Hit}$ = 641±12 ms, $RT_{Miss}$ = 647±12 ms, $RT_{CR}$ = 594±10 ms; paired $t$-test hit vs. CR: $p$ = 0.02, miss vs. CR: $p$ = 3 * $10^{-8}$). The yes/no-response times for hits and misses did not differ significantly ($p$ = 0.43). Additionally in unconfident compared to confident near-threshold trials, yes/no-responses were on average 221 ms slower (mean ± SD: $RT_{Near\_unconf}$ = 789±11 ms, $RT_{Near\_conf}$ = 569±9 ms; paired $t$-test: $p$ = 2 * $10^{-16}$). Splitting near-threshold trials by confidence resulted in on average 49 unconfident misses (range: 6-143), 135 confident misses (range: 29-289), 70 unconfident hits (range: 9-181), and 122 confident hits (range: 24-277).

## Cardiac cycle

First, we addressed the question whether stimulus detection differed along the cardiac cycle. For hits, mean angles within the cardiac cycle were not uniformly distributed ($R = 0.34$, $p = 0.007$; Figure 2), indicating a relation between cardiac phase and stimulus detection. Sixteen participants had a mean angle for hits in the last quarter of the cardiac cycle (270-360°). The Rayleigh tests were not significant for misses ($R = 0.20$, $p = 0.18$) and correct rejections ($R = 0.05$, $p = 0.91$).

**Figure 2.** Distribution of mean angles (stimulus onset relative to cardiac cycle) for (**A**) correct rejections (green), (**B**) misses (purple), and (**C**) hits (red). Each dot indicates a mean angle by one participant. The line around the inner circle shows the density distribution of these mean angles. The direction of the arrow in the center indicates the mean angle across the participants while the arrow length represents the mean resultant length $R$. The resulting p-value of the Rayleigh test of uniformity is noted below.

Second, we repeated the analysis by splitting near-threshold trials based on the reported decision confidence. The unimodal distribution was also present for confident hits ($R = 0.38$, $p = 0.002$) but not for unconfident hits ($R = 0.10$, $p = 0.69$). Eighteen participants had a mean angle for confident hits in the last quarter of the

cardiac cycle (270-360°). Confident misses also showed a unimodal distribution ($R =$

0.28, $p = 0.04$). Unconfident misses ($R = 0.22$, $p = 0.14$), confident correct rejections

($R = 0.005$ $p = 1.00$) and unconfident correct rejections ($R = 0.01$, $p = 1.00$) did not

allow to reject the uniform distribution. Two participants were excluded from the

analysis of unconfident correct rejections due to zero unconfident correct rejections

(mean $n = 20$, $SD = 22$, range: 0-88).

Third, near-threshold detection was analyzed separately for confident and

unconfident decisions within four time intervals after the R-peak (0-200 ms, 200-400

ms, 400-600 ms, 600-900 ms). First, we calculated a two-way repeated measures

ANOVA with near-threshold detection rate as dependent variable and time interval

and decision confidence as within-participants factors. The ANOVA showed a main

effect between confidence categories ($F(1,40) = 7.84$, $p = 0.008$) and an interaction

effect of time and confidence categories on near-threshold detection ($F(3,120) = 3.09$,

$p = 0.03$). Subsequently, post-hoc $t$-tests were calculated between confidence

categories for each time interval and between time intervals within each confidence

category (Figure 3). The detection rate was significantly lower for confident compared

to unconfident decisions within the first two time intervals (0-200 ms: FDR-corrected

$p = 0.02$; 200-400 ms: FDR-corrected $p = 0.02$). Between time intervals, we only

observed for confident decisions a significant lower detection rate at 200-400 ms

compared to 400-600 ms (FDR-corrected $p = 0.02$) and 600-900 ms (FDR-corrected

$p = 0.02$).

**Figure 3.** Circular distribution within the cardiac cycle of unconfident/confident misses and hits (**A-D**)

and detection rates for unconfident and confident decisions at four time intervals after the R-peak (**E**).

The distributions of mean angles (stimulus onset relative to cardiac cycle) are shown for (**A**) unconfident

misses (red), (**B**) confident misses (blue), (**C**) unconfident hits (red), and (**D**) confident hits (red). In **A-D**,

each dot indicates a mean angle by one participant. The line around the inner circle shows the density

distribution of these mean angles. The direction of the arrow in the center indicates the mean angle

across the participants while the arrow length represents the mean resultant length $R$. The resulting p-

value of the Rayleigh test of uniformity is noted below. $E$, Boxplots show the distribution of detection

rates for unconfident (red) and confident (blue) decisions at four time intervals after the R-peak.

Boxplots indicate the median (centered line), the 25%/75% percentiles (box), 1.5 times the interquartile

range or the maximum value if smaller (whiskers), and outliers (dots beyond the whisker range).

Significant post-hoc $t$-tests are indicated with a black bar and the respective FDR-corrected $p$-value.

**Cardiac interbeat interval**

For each stimulus-response condition (hit, miss, and correct rejection), we extracted

the interbeat interval entailing the stimulus onset, as well as the two preceding and

subsequent interbeat intervals (Figure 4A). We used a two-way repeated measures

ANOVA to test the factors time and stimulus-response condition, and their interaction

on interbeat intervals. The main effect of time ($F(2.73,109.36) = 35.60$, $p = 3 \times 10^{-15}$)

and the interaction of time and stimulus-response condition ($F(4.89,195.55) = 4.92$, $p$

$= 0.0003$) were significant. There was no significant main effect of stimulus-response

condition on interbeat intervals ($F(1.42,56.75) = 2.35$, $p = 0.12$). Following, post-hoc

$t$-tests were calculated at each interbeat interval between the stimulus-response

conditions (5 x 3) and within each stimulus-response condition between subsequent

interbeat intervals (3 x 4), resulting in 27 FDR-corrected $p$-values. At "S+1", the

interbeat intervals (IBI) for hits were significantly longer than for misses ($\Delta IBI = 5.2$

ms, FDR-corrected $p = 0.024$) and correct rejections ($\Delta IBI = 4.4$ ms, FDR-corrected

$p = 0.017$). The interbeat intervals between misses and correct rejections did not differ

significantly (FDR-corrected $p = 0.62$). At "S+2", the interbeat intervals for hits were

still longer compared to correct rejections ($\Delta IBI = 5.3$ ms, FDR-corrected $p = 0.014$)

but not to misses (FDR-corrected $p$ = 0.25). Within each stimulus-response condition

(hit, miss, and correct rejection) subsequent interbeat intervals differed significantly

(FDR-corrected $p < 0.005$).

**Figure 4.** Interbeat intervals before and after the stimulus onset for (**A**) correct rejections (green),

misses (purple), and hits (red), and for (**B**) confident (blue) and unconfident (red) decisions. Confidence

bands reflect within-participant 95% confidence intervals. The label "Stimulus" on the y-axis indicates

the cardiac cycle when the stimulation or cue only were present. The labels "S-1" and "S+1" indicate

the preceding and following intervals respectively. In **A**, the two asterisks "**" at "S+1" indicate

significant FDR-corrected $t$-tests between hits and misses, and between hits and correct rejections.

The one asterisk "*" at "S+2" in **A** indicates a significant FDR-corrected $t$-test between hits and correct

rejections. In **B,** the asterisks at "S+1" and "S+2" indicate significant FDR-corrected $t$-tests between

confident and unconfident decisions. The lines with asterisks on the bottom indicate significant FDR-

corrected $t$-tests for subsequent interbeat intervals within all conditions.

Furthermore, interbeat intervals of trials with confident and unconfident decisions

independent of stimulus presence and yes/no-response (excluding false alarms) were

compared with a two-way repeated measures ANOVA (Figure 4B). The main effects

time ($F(2.71, 108.31) = 42.37$, $p = 2 \times 10^{-16}$) and confidence ($F(1,40) = 5.36$, $p = 0.026$),

as well as the interaction of time and confidence were significant ($F(2.73, 109.05) =$

$30.79$, $p = 2 \times 10^{-16}$). Post-hoc $t$-tests between the two confidence categories for each

interbeat interval (1 x 5) and within each confidence category between subsequent

interbeat intervals (2 x 4) revealed significant longer post-stimulus interbeat intervals

for unconfident compared to confident decisions at "S+1" ($\Delta IBI = 4.6$ ms, FDR-

corrected $p = 0.005$) and "S+2" ($\Delta IBI = 10.6$ ms, FDR-corrected $p = 6 \times 10^{-7}$). All

subsequent interbeat intervals differed significantly within each confidence category

(FDR-corrected $p < 0.05$). When repeated for near-threshold trials only, the difference

between confidence categories was still present within each awareness condition:

unconfident hits and misses showed longer interbeat intervals at "S+2" compared to

confident hits ($\Delta IBI = 5.7$ ms, FDR-corrected $p = 0.047$) and confident misses

respectively ($\Delta IBI = 10.2$ ms, FDR-corrected $p = 0.008$).

**Pulse wave relative to electric cardiac cycle**

Next to the electric cardiac cycle, we assessed whether stimulus detection was

dependent on the pulse wave cycle measured at the left middle finger. Pulse wave

peaks were located in the cardiac cycle by calculating the time to the preceding R-

peak: the pulse wave transit time (PWTT) and the PWTT relative to its current cardiac

cycle in degree. The PWTT was on average 405 ms ($SD = 24$ ms, range: 354-439 ms).

The pulse wave peak occurred on average in the middle of the cardiac cycle (mean

angle $M_{PWTT} = 178°$, $R = 0.91$, $p = 0$) after the mean angle of confident misses ($M_{Confident}$

$_{miss} = 96°$) and before the mean angle of confident hits ($M_{Confident\ hit} = 325°$).

For putting the observed correlations between detection and the cardiac cycle

in relation to the pulse wave peak, the analysis of near-threshold hit rates during

different stimulus onset intervals after the R-peak was repeated limited to 0-400 ms

with shorter intervals (50 ms) and without splitting by confidence. A one-way

repeated-measures ANOVA showed a main effect by time interval on near-threshold

detection rate ($F(5.64, 225.46) = 3.15$, $p = 0.007$). The near-threshold hit rates were

significantly decreased before the pulse wave peak (mean PWTT = 405 ms) during

the interval of 250-300 ms compared to the interval of 0-50 ms (FDR-corrected $p =$

0.038). The interval 250-300 ms was plotted on the average pulse wave locked to the

preceding R-peak and its slope (difference between adjacent samples). It shows that

after 250 ms the pulse wave substantially increases (Figure 5B).

**Figure 5.** Pulse wave and detection relative to cardiac cycle. (**A**) Mean pulse wave measured at the left middle finger for each participant locked to preceding R-peak. (**B**) First derivative of the mean pulse wave. (**C**) Detection rate of near-threshold trials in 50-ms stimulus onset intervals since preceding R-

peak. The black dots indicate the mean across participants. The blue line is the locally smoothed loess

curve with a 95% confidence interval (grey) across these means.

**Respiratory cycle**

First, we investigated whether conscious tactile perception depends on the stimulus

onset relative to the respiratory cycle. Thus, we calculated the mean angles for hits,

misses, and correct rejections for each participant and tested their circular

distribution with the Rayleigh test of uniformity. For all conditions, uniformity was

rejected in favor of an alternative unimodal distribution (correct rejections: $R = 0.58$,

$p = 4 \times 10^{-7}$; misses: $R = 0.54$, $p = 2 \times 10^{-6}$; hits: $R = 0.65$, $p = 1 \times 10^{-8}$; Figure 6). These

unimodal distributions were centered at stimulus onset for the three conditions (mean

angle $M_{correct\ rejection} = 3.2°$, $M_{miss} = 5.0°$, and $M_{hit} = 15.1°$). To assess whether the strength

of the inhale onset locking differed significantly between hits, misses, and correction

rejections, the circular variance of stimulus onset angles across trials was calculated

for each stimulus-response condition and compared with $t$-tests. Hits had a lower

circular variance than misses ($\Delta V = -0.044$, $t(40) = -3.17$, $p = 0.003$) and correct

rejections ($\Delta V = -0.035$, $t(40) = -2.78$, $p = 0.008$), i.e., exhibited a stronger clustering

around the mean direction . There was no significant difference in circular variance

between misses and correct rejections ($p = 0.44$). Additionally, we investigated

whether the detection rate of near-threshold trials varied across the cardiac cycle.

For this purpose, near-threshold trials were assigned to four subsequent 1-s time

intervals relative to the preceding inhale onset (0-4 s). A one-way repeated-measures

ANOVA could not establish an effect by time interval on near-threshold detection rate

($F(1.81, 72.30) = 0.83$, $p = 0.43$).

**Figure 6.** Circular distribution of mean stimulus onsets relative to the respiratory cycle for (**A**) correct rejections (green), (**B**) misses (purple), and (**C**) hits (red). Zero degree indicates the inspiration onset. Each dot indicates a mean angle by one participant. The line around the inner circle shows the density distribution of these mean angles. The direction of the arrow in the center indicates the mean angle across the participants while the arrow length represents the mean resultant length $R$. The resulting p-value of the Rayleigh test of uniformity is noted below.

Second, the distribution of mean angles was assessed for confident and unconfident decisions. Hits, misses, and correct rejections were split by decision confidence and the resulting distributions were evaluated with the Rayleigh test for uniformity. All stimulus-response conditions showed for unconfident and confident decisions a significant unimodal distribution locked around the stimulus onset: unconfident correct rejections (mean angle $M_{unconf\_CR} = 18.3°$; $R = 0.41$, $p = 0.001$), confident correct rejections ($M_{conf\_CR} = 2.2°$; $R = 0.58$, $p = 4 \times 10^{-7}$), unconfident misses ($M_{unconf\_miss} = 13.4°$; $R = 0.45$, $p = 0.0001$), confident misses ($M_{conf\_miss} = 6.7°$, $R = 0.51$; $p = 0.00001$), unconfident hits ($M_{unconf\_hit} = 10.4°$; $R = 0.51$, $p = 0.0001$), and confident

hits ($M_{conf\_hit}$ = 15.2°; $R$ = 0.67, $p$ = 4 x $10^{-9}$). Two participants had zero unconfident

correct rejections and were not considered in the respective Rayleigh test.

Third, in order to examine aforementioned phase effects further, we

investigated whether participants adjusted their respiration rhythm to the paradigm

in the beginning of the experiment. Thus for the first 30 trials of the first block, a

random-intercept linear regression model with maximum likelihood estimation (lmer

function in R) was calculated to evaluate the effect of trial number on trial angle

difference from the mean for each participant. The angle difference was determined

between the stimulus onset angle within the respiratory cycle of each trial and the

mean of all angles in the first block ("diff_angle2mean"). This analysis included 37

participants with a first block and excluded trials with false alarms. Comparing the

model "diff_angle2mean ~ 1 + trial + (1|participant)" with a random intercept-only

model "diff_angle2mean ~ 1 + (1|participant)" revealed an effect of trial on the

difference to the angle mean within the first 30 trials of the first block ($\chi^2$ = 5.84, $p$ =

0.016). The fixed-effect slope was $b_1$ = -0.47 and the mean of the random-intercepts

$b_0$ = 79.7 (*diff_angle2mean* = $b_1$ * *trial* + $b_0$).

**Respiratory cycle duration**

Given the previously reported heart slowing during conscious tactile perception

(Motyka et al., 2019)., we tested whether a similar effect was also present in the

respiratory rhythm. Indeed, the mean duration of respiratory cycles differed between

response categories (Figure 7), as indicated by a one-way repeated-measures

ANOVA ($F$(1.49, 59.61) = 13.11, $p$ = 0.0001). Post-hoc $t$-tests showed that respiratory

cycles accompanying misses (mean $t$ = 3.86 s) were significantly longer than

respiratory cycles with correct rejections (mean $t$ = 3.82 s, $\Delta t$ = 40 ms, FDR-corrected

$p$ = 0.002) and with hits (mean $t$ = 3.77 s, $\Delta t$ = 91 ms, FDR-corrected $p$ = 0.0002).

Respiratory cycles with hits were also significantly shorter than correct rejections ($\Delta t$

= 50 ms, FDR-corrected $p$ = 0.014).

Additionally, we analyzed whether the respiratory cycle duration differed

between confident and unconfident hits and misses. There was a main effect by

detection ($F$(1, 40) = 14.64, $p$ = 0.0004) but not by confidence ($F$(1, 40) = 1.15, $p$ =

0.29) on respiratory cycle duration in a two-way repeated measures ANOVA. The

interaction of detection and confidence was not significant ($F$(1, 40) = 0.83, $p$ = 0.37).

**Figure 7.** Mean respiratory cycle duration in seconds for correct rejections (green), misses (purple),

and hits (red). The boxplots indicate the median (centered line), the 25%/75% percentiles (box), 1.5

times the interquartile range or the maximum value if smaller (whiskers), and outliers (dots beyond the

whisker range). Significant post-hoc $t$-tests are indicated above the boxplot with a black bar and the

respective FDR-corrected $p$-value.

**Phase-locking between cardiac and respiratory activity**

Due to the natural coupling of cardiac and respiratory rhythms (Dick et al., 2014), we investigated whether phase-locking of both rhythms is associated with conscious tactile perception. Phase-locking values (PLVs) were calculated across trials using n:m synchronization (Tass et al., 1998; Lachaux et al., 1999) to account for the different frequency bands of the two signals. PLVs were compared between hits, misses, and correct rejections with a one-way repeated-measures ANOVA. The ANOVA showed no significant main effect of stimulus-response condition on PLVs between cardiac and respiratory activity ($F(1.98,79.15) = 1.72$, $p = 0.19$).

**Peripheral nerve activity**

For the sub-sample of twelve participants with peripheral nerve recordings at the left upper arm, there was no somatosensory evoked potential associated with near-threshold stimuli. Also, the grand mean across participants did not show a difference between trials with and without near-threshold stimulation. We concluded that near-threshold stimulation intensities (in the given sub-sample on average 1.88 mA, range: 0.79-2.50 mA) did not produce sufficiently high peripheral somatosensory evoked potentials to measure them non-invasively from the inner side of the upper arm. Hence, we did not further pursue the analysis of peripheral somatosensory evoked potentials. (Yet note that peripheral somatosensory evoked potentials were observed in a pilot study with the same acquisition setup but applying super-threshold stimulation intensities of 6 mA.)

## Discussion

In a tactile detection paradigm with near-threshold electrical finger nerve stimulation, cardiac and respiratory activity was observed with electrocardiography (ECG), pulse oximetry, and a respiration chest belt. The present study is the third independent study, where we observed a cardiac cycle effect on tactile detection (hits more likely during diastole) and a pronounced heart slowing with conscious tactile perception (Motyka et al., 2019; Al et al., 2020). The previously reported lower tactile detection during systole (Al et al., 2020) and the unimodal distribution of hits in diastole (Motyka et al., 2019; Al et al., 2020) were both only present when decision confidence was high. Using finger pulse oximetry, we identified that the lowest tactile detection rate between 250 and 300 ms after the R-peak corresponded to the pulse wave onset. We furthermore showed that the respiration rhythm adapted to the tactile detection paradigm, such that inhale onsets tended to be locked to expected stimulus onsets. This locking of inhale onsets to stimulus onset was more consistent for detected compared to undetected near-threshold trials. Respiratory cycles accompanying misses were longer in comparison to hits and correct rejections.

### Detection varies across the cardiac cycle only for confident decisions

Relating tactile detection to cardiac activity revealed that hits were not uniformly distributed across the cardiac cycle and more likely to appear in the last quarter, as we have shown previously (Motyka et al., 2019; Al et al., 2020). Splitting hits and misses by decision confidence - not possible in our previous studies - showed that this unimodal distribution was only present for confident but not unconfident hits.

Additionally, also confident but not unconfident misses showed a unimodal

distribution (centered around 90 degrees).

When we analyzed the near-threshold detection rate for four stimulus onset

intervals relative to the preceding R-peak split by decision confidence, we observed

that detection was lower when the decision was confident. Furthermore, only

confident hits were reduced in the second time interval (200-400 ms) compared to

the two later timer intervals (400-900 ms). In our previous study (Al et al., 2020), a

decreased detection rate was observed for the very same time interval.

While replicating the unimodal distribution of hits within the cardiac cycle here

for the third time in an independent study of somatosensory detection (Motyka et al.,

2019; Al et al., 2020), the explanation for this effect still stays speculative. It seems

highly plausible that the decreased detection rate between 200 and 400 ms after the

R-peak is related to the arrival of the pulse wave in the finger at circa 250 ms or

heartbeat sensations (see below discussion on *Pulse wave*). Our finding that the

number of confident hits is reduced (in contrast to unconfident ratings) seems

consistent with a "late" event in the perceptual decision-making process. This is in

line with our previous EEG study which has shown that only late cortical

somatosensory evoked potentials (P300) are increased for stimuli during diastole

compared to systole (Al et al., 2020). Hence, both, the reduction of confident hits and

the modulation of (only) late SEP-components, speak for higher cognitive processes

as meta-cognition or prediction to be involved in the cardiac cycle effect on

conscious tactile perception.

**Pulse wave onset marks decreased detection in cardiac cycle**

Addressing the hypothesis whether decreased near-threshold detection 200-400 ms after the R-peak was caused by the pulse wave peak in the finger (Al et al., 2020), pulse wave peaks were located in the cardiac cycle. As a result, the decreased near-threshold detection rate was located more precisely 250-300 ms after the R-peak, before the pulse wave peak (405 ms) in the middle of the cardiac cycle (178°). Inspecting the slope of the pulse wave showed a take-off around 250 ms after the preceding R-peak, indicating the pulse wave onset.

If a suppressive top-down mechanism for pulse wave related changes in the finger affects near-threshold tactile detection, then it might be already sensitive to initial changes of the pulse wave in the finger than only to its peak (Macefield, 2003). Perception of heartbeats has been reported to occur in the very same time interval of 200-300 ms after the R-peak (Yates et al., 1985; Brener and Kluvitse, 1988; Ring and Brener, 1992). While this temporal judgement is unlikely to be solely based on the pulse wave in the finger - heartbeat sensations were mainly localized on the chest (Khalsa et al., 2009; Hassanpour et al., 2016), it might be possible that the prediction of strongest heartbeat-related changes at 250-300 ms attenuated the detection of weak stimuli presented in the same time window. Temporally locating the lower tactile detection during systole at the pulse wave onset in the finger provides further evidence for the view that the cardiac cycle effect does not correspond to maximal peripheral cardiac changes. It rather suggests that this is a conflict on the perceptual level during an interval with the highest uncertainty whether a weak pulse was generated internally (heartbeat) or applied externally (Allen et al., n.d.).

**Heart slowing with conscious perception and unconfident decisions**

In the present study, previously reported heart slowing with conscious perception (Park et al., 2014; Cobos et al., 2019; Motyka et al., 2019) was replicated and in addition to it observed to be enhanced for unconfident decisions. Greater heart slowing for unconfident decisions suggests the involvement of another mechanism determining heart rate deceleration compared to the unimodal distribution of hits in diastole which was absent for unconfident hits. Increased heart slowing for unconfident decisions might be associated with uncertainty, because heart slowing has been reported for the violation of performance-based expectations in a learning paradigm (Crone et al., 2003), for errors in a visual discrimination task (Łukowska et al., 2018), and for error keystrokes by pianists (Bury et al., 2019).

**Inspiration locking to stimulus onsets**

Localizing stimulus onsets in the respiratory cycle revealed that inspiration onsets locked at the expected stimulus onset independent of stimulus presence, detection, and decision confidence. Circular variance was lower for hits than misses, indicating a more pronounced locking went along with a higher likelihood of hits. For the first thirty trials, the difference to the mean angle of the first block showed a linear decrease, suggesting that participants adapted their respiration rhythm to the paradigm in the beginning. Respiratory cycles accompanying misses had a significant longer duration than hits.

These observations support the notion that inspiration is tuning the sensory system for incoming information (Perl et al., 2019). Recently, it has been shown that inhale onsets were locked to different cognitive task onsets, which increased task-

specific brain activity and performance in a visuospatial perception task (Perl et al., 2019). Neural activity in piriform cortex, amygdala, and hippocampus synchronized with respiration and had its oscillatory power peak during nasal inspiration, when fearful face detection was faster, and memory encoding and retrieval was enhanced compared to expiration (Zelano et al., 2016). While we also observed locking of inhale onsets to our detection task and pronouncedly to stimulus detection, future studies could investigate whether this respiratory pattern also drives the brain activity into an optimal state for conscious tactile perception. For the visuospatial perception task by Perl et al. (2019), decreased alpha power has been observed in postcentral and parahippocampal gyrus during inspiration compared to expiration. Analogously, lower pre-stimulus alpha power in central brain areas (Mu rhythm) has been associated with higher neural excitability reflected in improved conscious tactile perception (Schubert et al., 2009; Nierhaus et al., 2015; Craddock et al., 2017; Forschack et al., 2020; Stephani et al., 2021).

Meanwhile, it is unclear whether longer respiratory cycles for misses in our study are a result of less pronounced inspiration locking or a consequence of missing the stimulus. Future studies with longer interstimulus intervals would be able to dissect whether the respiratory cycle duration is a determinant or consequence of undetected near-threshold trials. For now, we can state that participants adapted their respiratory rhythm to the paradigm and showed this more consistently in detected near-threshold trials.

**Conclusion**

We show that the two predominant body rhythms differentially modulate conscious

tactile perception. Cardiac cycle effects seem to be grounded on higher cognitive

processes compatible with an interoceptive predictive coding account. Phase locking

of respiration seems to facilitate conscious tactile perception by tuning the inspiration

phase to optimize detection task performance.

## References

Al E, Iliopoulos F, Forschack N, Nierhaus T, Grund M, Motyka P, Gaebler M, Nikulin VV, Villringer A (2020) Heart-brain interactions shape somatosensory perception and evoked potentials. Proc Natl Acad Sci USA 7:201915629.

Allen M, Levy A, Parr T, Friston, K (2019) In the body's eye: the computational anatomy of interoceptive inference. bioRxiv 603928. doi: 10.1101/603928

Azzalini D, Rebollo I, Tallon-Baudry C (2019) Visceral Signals Shape Brain Dynamics and Cognition. Trends Cogn Sci 23:488–509.

Babo-Rebelo M, Richter CG, Tallon-Baudry C (2016) Neural Responses to Heartbeats in the Default Network Encode the Self in Spontaneous Thoughts. J Neurosci 36:7829–7840.

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Royal Statistical Society, Series B Methodological 57:289–300

Brener J, Kluvitse C (1988) Heartbeat detection: judgments of the simultaneity of external stimuli and heartbeats. Psychophysiology 25:554–561.

Bury G, García-Huéscar M, Bhattacharya J, Ruiz MH (2019) Cardiac afferent activity modulates early neural signature of error detection during skilled performance. NeuroImage 199:704–717.

Cobos MI, Guerra PM, Vila J, Chica AB (2019) Heart-rate modulations reveal attention and consciousness interactions. Psychophysiology 56:e13295.

Craddock M, Poliakoff E, El-deredy W, Klepousniotou E, Lloyd DM (2017) Pre-stimulus alpha oscillations over somatosensory cortex predict tactile misperceptions. Neuropsychologia 96:9–18.

Critchley HD, Garfinkel SN (2015) Interactions between visceral afferent signaling

and stimulus processing. Front Neurosci 9:305–309.

Critchley HD, Harrison NA (2013) Visceral influences on brain and behavior. Neuron

77:624–638.

Crone EA, van der Veen FM, van der Molen MW, Somsen RJM, van Beek B,

Jennings JR (2003) Cardiac concomitants of feedback processing. Biological

Psychology 64:143–156.

Dick TE, Hsieh Y-H, Dhingra RR, Baekey DM, Galán RF, Wehrwein E, Morris KF

(2014) Cardiorespiratory coupling: common rhythms in cardiac, sympathetic,

and respiratory activities. Prog Brain Res 209:191–205.

Forschack N, Nierhaus T, Müller MM, Villringer A (2020) Dissociable neural

correlates of stimulation intensity and detection in somatosensation.

NeuroImage:116908.

Galvez-Pol A, McConnell R, Kilner JM (2020) Active sampling in visual search is

coupled to the cardiac cycle. Cognition 196:104149.

Hassanpour MS, Yan L, Wang DJJ, Lapidus RC, Arevian AC, Simmons WK,

Feusner JD, Khalsa SS (2016) How the heart speaks to the brain: neural activity

during cardiorespiratory interoceptive stimulation. Phil Trans R Soc Lond B 371.

Huijbers W, Pennartz CMA, Beldzik E, Domagalik A, Vinck M, Hofman WF, Cabeza

R, Daselaar SM (2014) Respiration phase-locks to fast stimulus presentations:

implications for the interpretation of posterior midline "deactivations". Hum

Brain Mapp 35:4932–4943.

Khalsa SS, Rudrauf D, Feinstein JS, Tranel D (2009) The pathways of interoceptive

awareness. Nat Neurosci 12:1494–1496.

Kingdom FAA, Prins N (2009) Psychophysics. London: Academic Press Inc.

Kleiner M, Brainard D, Pelli D, Ingling A, Murray R, Broussard C (2007) What's new

in psychtoolbox-3. Perception 36:1–16.

Kralemann B, Frühwirth M, Pikovsky A, Rosenblum M, Kenner T, Schaefer J, Moser

M (2013) In vivo cardiac phase response curve elucidates human respiratory

heart rate variability. Nat Commun 4:2418–2419.

Kunzendorf S, Klotzsche F, Akbal M, Villringer A, Ohl S, Gaebler M (2019) Active

information sampling varies across the cardiac cycle. Psychophysiology

56:e13322.

Lachaux JP, Rodriguez E, Martinerie J, Varela FJ (1999) Measuring phase

synchrony in brain signals. Hum Brain Mapp 8:194–208.

Macefield VG (2003) Cardiovascular and respiratory modulation of tactile afferents

in the human finger pad. Exp Physiol 88:617–625.

Motyka P, Grund M, Forschack N, Al E, Villringer A, Gaebler M (2019) Interactions

between cardiac activity and conscious somatosensory perception.

Psychophysiology 56:469–13.

Nierhaus T, Forschack N, Piper SK, Holtze S, Krause T, Taskin B, Long X, Stelzer J,

Margulies DS, Steinbrink J, Villringer A (2015) Imperceptible somatosensory

stimulation alters sensorimotor background rhythm and connectivity. J Neurosci

35:5917–5925.

Ohl S, Wohltat C, Kliegl R, Pollatos O, Engbert R (2016) Microsaccades Are

Coupled to Heartbeat. J Neurosci 36:1237–1241.

Oldfield RC (1971) The assessment and analysis of handedness: the Edinburgh

inventory. Neuropsychologia 9:97–113.

Park H-D, Barnoud C, Trang H, Kannape OA, Schaller K, Blanke O (2020) Breathing

is coupled with voluntary action and the cortical readiness potential. Nat

Commun 11:289–8.

Park H-D, Correia S, Ducorps A, Tallon-Baudry C (2014) Spontaneous fluctuations

in neural responses to heartbeats predict visual detection. Nat Neurosci 17:612–

618.

Perl O, Ravia A, Rubinson M, Eisen A, Soroka T, Mor N, Secundo L, Sobel N (2019)

Human non-olfactory cognition phase-locked with inhalation. Nat Hum Behav

3:501–512.

Power JD, Lynch CJ, Dubin MJ, Silver BM, Martin A, Jones RM (2020)

Characteristics of respiratory measures in young adults scanned at rest,

including systematic changes and "missed" deep breaths. NeuroImage

204:116234.

Ring C, Brener J (1992) The temporal locations of heartbeat sensations.

Psychophysiology 29:535–545.

Savitzky A, Golay MJE (1964) Smoothing and differentiation of data by simplified

least squares procedures. Analytical Chemistry 36:1627–1639.

Schubert R, Haufe S, Blankenburg F, Villringer A, Curio G (2009) Now you'll feel it,

now you won't: EEG rhythms predict the effectiveness of perceptual masking. J

Cognitive Neurosci 21:2407–2419.

Seth AK, Friston KJ (2016) Active interoceptive inference and the emotional brain.

Phil Trans R Soc Lond B 371.

Stephani T, Hodapp A, Jamshidi Idaji M, Villringer A, Nikulin VV (2021) Neural

excitability and sensory input determine intensity perception with opposing

directions in initial cortical responses. bioRxiv 2020.11.27.401430.

doi:10.1101/2020.11.27.401430

Tass P, Rosenblum MG, Weule J, Kurths J, Pikovsky A, J V, Schnitzler A, Freund H-

J (1998) Detection of n:m Phase Locking from Noisy Data: Application to

Magnetoencephalography. Phys Rev Lett 81:3291–3294.

Yates AJ, Jones KE, Marie GV, Hogben JH (1985) Detection of the heartbeat and

events in the cardiac cycle. Psychophysiology 22:561–567.

Zelano C, Jiang H, Zhou G, Arora N, Schuele S, Rosenow J, Gottfried JA (2016)

Nasal Respiration Entrains Human Limbic Oscillations and Modulates Cognitive

Function. J Neurosci 36:12448–12467.

Łukowska M, Sznajder M, Wierzchoń M (2018) Error-related cardiac response as

information for visibility judgements. Scientific Reports 8:1131.