Heightened virulence of *Yersinia* is associated with decreased function of the

YopJ protein

Chris A. Mares[1*#], Fernando P. Lugo[1], Mohammad Albataineh[1], Beth A. Goins[2], Irene

G. Newton[3], Ralph R. Isberg[4,5#] , and Molly A. Bergman[1]

[1]Departments of Microbiology & Immunology and [2]Department of Radiology, The

University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Dr, San

Antonio TX 78229, [3]Department of Biology, Indiana University, Jordan Hall, 1001 East

Third Street; Bloomington, IN 47405, [4]Department of Molecular Biology & Microbiology

and [5]Howard Hughes Medical Institute, Tufts University School of Medicine, 150

Harrison Ave, Boston, MA 02111,

*current address: Department of Life Sciences, Texas A&M University-San Antonio, San

Antonio, Texas, United States of America

[#]Corresponding authors

keywords: T3SS, effector, YopJ, YopP, Yersinia, positive selection, Darwinian,

cytotoxicity, macrophage, avirulence

Abstract

Despite the maintenance of YopP/J alleles throughout the human-pathogenic *Yersinia*

lineage, the benefit of YopP/J-induced phagocyte death for *Yersinia* pathogenesis in

animals is not obvious. To determine how sequence divergence of YopP/J has

impacted *Yersinia* virulence, we examined protein polymorphisms in this Type III

secreted effector protein across 17 *Yersinia* species, and tested the consequences of

polymorphism in a murine model of sub-acute systemic yersiniosis. Our evolutionary

analysis revealed that codon 177 has been subjected to positive selection - the *Y.*

enterocolitica residue had been altered from a leucine to a phenylalanine in nearly all *Y.*

pseudotuberculosis and *Y. pestis* strains examined. Despite being a minor change, as

both leucine and phenylalanine have hydrophobic side chains, reversion of YopJ^{F177} to

the ancestral YopJ^{L177} variant yielded a *Y. pseudotuberculosis* strain with enhanced

cytotoxicity towards macrophages, consistent with previous findings. Surprisingly,

expression of YopJ^{F177L} in the mildly attenuated *ksgA*$^-$ background rendered the strain

completely avirulent in mice. Consistent with this hypothesis that YopJ activity indirectly

relates to *Yersinia* pathogenesis *in vivo*, *ksgA*$^-$ strains lacking functional YopJ failed to

kill macrophages but actually regained virulence in animals. Also, treatment with the

anti-apoptosis drug suramin prevented YopJ-mediated macrophage cytotoxicity and

enhanced *Y. pseudotuberculosis* virulence *in vivo*. Our results demonstrate that

Yersinia-induced cell death is detrimental for bacterial pathogenesis in this animal

model of illness, and indicate that positive selection has driven YopJ/P and *Yersinia*

evolution towards diminished cytotoxicity and increased virulence, respectively.

Introduction

Bacterial pathogens induce host cell death by various mechanisms, and with downstream consequences that seem counterintuitive [1]. At face value, death of host cells, particularly cells of the innate immune system, should cripple the host immune response and allow pathogens to escape clearance. However, the opposite often occurs – induction of host cell death can induce a host response that leads to pathogen removal [1]. Mechanisms underlying this outcome are incompletely understood, but likely include removing niches for intracellular replication, or enhancing recruitment of highly activated phagocytes that then ingest and degrade the offending bacteria. Both host and bacterial factors are necessary for pathogen-induced host cell death – caspase-1 being a notable player during macrophage pyroptosis [2] and required for host protection against pyroptosis-inducing pathogens [3, 4]. Thus, pathogen-triggered cytotoxicity is a host defense strategy. But why do pathogens maintain the genes encoding the pro-cytotoxicity factors in their genomes? The existence of such genes throughout the bacterial kingdom suggests that host cell death plays an important role in persistence of individual species.

Three species of mammalian-pathogenic *Yersinia,* the gastroenteritis-causing *Y. enterocolitica* and *Y. pseudotuberculosis* and the plague-causing *Y. pestis*, have maintained such a pro-cytotoxicity factor throughout their evolutionary history, in spite of significant genomic decay [5, 6]. YopJ (so-called in *Y. pestis* and *Y. pseudotuberculosis*, called YopP in *Y. enterocolitica*) is one of the 6-8 Type 3 secretion system (T3SS)

effector proteins (*Yersinia* outer proteins, Yops) injected by the bacteria into host cells

[7-9]. YopJ has been described as having ubiquitin-like protein protease [10],

deubiquitinase [11, 12] and acyl transferase activity [13, 14], with the latter requiring the

host cell co-factor inositol hexaskisphosphate [15]. In macrophages, YopJ activity

inhibits MAPK and NFκB signaling pathways, which in conjunction with TLR4 signaling

results in host cell death (reviewed in [16]), and recent evidence demonstrates that

YopJ-induced phagocyte death occurs by a mechanism distinct from both apoptosis and

pyroptosis but linked to necrosis[17]. Regardless of mechanism, the outcome is clear –

YopJ has profoundly negative effects on macrophage viability.

However, exactly what YopJ does for *Yersinia in vivo* is unclear. YopJ seems minimally

important for virulence, at least compared to most of the other Yops*,* as YopJ$^-$ *Y. pestis,*

YopJ$^-$ *Y. pseudotuberculosis*, and YopP$^-$ *Y. enterocolitica* all display near-wildtype levels

of virulence in various mouse models of acute illness [18-23]. In contrast, there is

evidence indicating that YopJ activity may actually diminish *Yersinia* virulence, as

different isoforms of YopP/J can alter the degree of macrophage death and mouse

illness following exposure to *Yersinia*. *Y. pseudotuberculosis* or *Y. pestis* expressing a

hypersecreted variant of YopP showed enhanced cytotoxicity towards cultured

macrophages but diminished virulence in mice [24, 25]. Moreover, *Y. pestis* strain KIM

also encodes a hypercytotoxic YopJ variant, which when expressed in *Y.*

pseudotuberculosis induced more macrophage cell death due to enhanced IKKβ

binding and thus reduced NFκB signaling; the impact of this variant upon *Y.*

pseudotuberculosis-caused illness in mice was not reported [26].

It is unknown if changes in YopP/J sequences occurred as a consequence of Darwinian

evolution, but given that host-pathogen interactions can exert strong natural selection

pressure upon both organisms [27], it seems likely that the YopJ-target protein

interactions have been guided by evolutionary pressure. Evidence of selection at the

codon level can be detected via phylogenetic analyses across multiple lineages to

detect patterns of protein polymorphisms relative to divergence [28, 29]. The advantage

of this approach is that it can reveal loci affected by positive selection, even if the

selection pressure is obscure. Such analyses of bacterial pathogen effector proteins

have revealed signatures of molecular evolution, with individual codons showing

evidence of directional selection [30, 31].

To address the hypothesis that YopP/J has been subjected to directional selection, we

analyzed 17 sequences from multiple strains of each human-pathogenic *Yersinia*

species for evidence of microevolution, and evaluated how the different isoforms

impacted the outcome of systemic sub-acute yersiniosis in a small animal model. Our

studies reveal that some YopJ residues differing between the species have been

subjected to evolutionary pressure. Reversion of a positively selected residue to the

ancestral one yielded a YopJ isoform with enhanced cytotoxicity, but one that markedly

attenuated *Y. pseudotuberculosis* virulence in animals. Conversely, loss of YopJ due to

catalytic inactivation or deletion rendered *Y. pseudotuberculosis* strains non-cytotoxic

but hypervirulent. Chemical inhibition of macrophage death during *in vivo* infection also

enhanced *Y. pseudotuberculosis* virulence. Our results indicate that that YopJ has

evolved towards diminished function and that YopJ-induced death is detrimental for *Y.*

pseudotuberculosis infection, but also suggest that some level of YopP/J-induced death

is beneficial to the genus, perhaps by allowing persistence in an unknown reservoir.

Results

YopJ has been positively-selected in *Y. pestis* and *Y. pseudotuberculosis* for

reduced cytotoxicity and enhanced virulence

The *yopP/J* gene has been maintained in the genomes of all strains of human

pathogenic *Yersinia* species sequenced to date [5, 32, 33]. This conservation is

suggestive of an important role for the YopJ protein in the natural setting. Although

YopJ homologs are found in the genomes of all 3 human pathogenic *Yersinia* species,

sequence conservation differs across the length of the molecule. Some of the

sequence polymorphisms group *Y. pestis* and *Y. pseudotuberculosis* to the exclusion of

Y. enterocolitica (the expectation), while some other polymorphisms do not cleanly

follow the predicted phylogeny of the organisms (**Figure 1A**). We investigated whether

or not these divergent residues show evidence of positive selection using a likelihood

ratio test comparing support in the data for various models of evolution (either allowing

for selection or imposing a nearly neutral model of evolution) [34, 35]. This analysis

identified sites under positive selection (that is, ancestral sequence reconstruction and

the number and kind of differences at each residue suggested that dN/dS >1,

Supplementary Table 2) and included residue 177, which is a leucine in *Y.*

enterocolitica and *Y. pestis* KIM but a phenylalanine in the other *Y. pestis* homologs

examined and all the *Y. pseudotuberculosis* YopJ homologs (**Figure 1B**). Interestingly,

the F177L isoform was recently described to have enhanced activity relative to the F177

protein, promoting phosphorylation of IκB-α, increased binding to IKKβ, and causing

enhanced macrophage cytotoxicity [26].

The existence of a hypercytotoxic YopJ variant allowed us to query if enhanced YopJ

activity altered *Y. pseudotuberculosis* virulence in a small animal model of systemic

subacute illness. As reported previously, YopJ ^{L177} shows increased cytotoxicity for

macrophages relative to the F177 isoform (Fig. 2A). Using an attenuated strain as our

baseline background strain (*ksgA*$^-$), we observed that KsgA$^-$ bacteria expressing

YopJ^{F177L} were more attenuated for virulence than the parental KsgA$^-$ strain, as

demonstrated by the 100% survival rate of mice exposed to the *ksgA*$^-$ *yopJ*^{F177L} strain

(**Figure 2B**). Increasing the inoculum dose did not reverse the strain's complete

attenuation (data not shown). These results indicate that the L177-containing isoform,

attenuates virulence in the derived *Y. pseudotuberculosis* strains, correlating with the

enhanced cytotoxicity conferred by this variant.

***Y. pseudotuberculosis* strains lacking functional YopJ display increased**

virulence in vivo

Given that a hypercytotoxic YopJ isoform attenuated *Y. pseudotuberculosis* illness in

mice, we predicted that the absence of YopJ would enhance the organism's virulence.

To test this hypothesis, we constructed an isogenic *yopJ* deletion strain in the *ksgA*$^-$

background, and confirmed that YopJ$^-$ KsgA$^-$ bacteria were non-cytotoxic for bone-

marrow-derived macrophages (**Figure 3A**). Consistent with the negative correlation

between cytotoxicity and virulence, mice inoculated with the $\Delta yopJ\ ksgA^-$ strain died

significantly sooner than those challenged with the $ksgA^-$ strain (**Figure 3B**), with

kinetics similar to the parental $ksgA^+$ strain [36]. Confirming that the enhanced virulence

phenotype of the $\Delta yopJ\ ksgA^-$ strain was due to the $yopJ$ deletion, a strain rescued for

the genomic $\Delta yopJ$ lesion by allelic exchange ($yopJ^{repaired}$) caused a mortality rate

identical to the parental $ksgA^-$ strain in animals (**Supplementary Figure 1**).

The enzymatic activity of *Y. pseudotuberculosis* YopJ has been mapped to a triad of

residues – histidine 109, glutamate 128, cysteine 172 [10]. We asked if the virulence

phenotype of a strain carrying the catalytic $yopJ^{C172A}$ mutation mimicked that observed

with the isogenic $\Delta yopJ$ strain. Similar to previous reports, the $yopJ^{C172A}ksgA^-$ strain was

unable to kill cultured macrophages (**Figure 3C**) [37]. Like YopJ$^-$ KsgA$^-$ bacteria,

YopJ^{C172A} KsgA$^-$ *Y. pseudotuberculosis* displayed increased virulence in animals

relative to the parental strain (median time to death day 11 versus 12, respectively).

Interestingly, YopJ^{C172A} KsgA$^-$ bacteria and YopJ$^-$ KsgA$^-$ bacteria consistently displayed

similar virulence and median times to death in multiple mouse survival assays (**Figure**

3D and 3E). Taken together, these results indicate that either the absence of YopJ or

the presence of inactive YopJ renders an attenuated strain of *Y. pseudotuberculosis*

more virulent *in vivo.*

The YopJ^{F177L} isoform attenuates bacterial burden and increases host cell death

in vivo

To further understand the pathogenesis of the different YopJ-expressing strains *in vivo,*

we examined the kinetics of bacterial colonization in the spleen and liver post-challenge,

examining burden at days 1, 3, 9 and 11 post-challenge. This analysis revealed two

trends. Firstly, at later timepoints, mice exposed to the *yopJ^{F177L} ksgA^{-}* strains carried

lower burdens of splenic and hepatic bacteria than mice exposed to the parental *ksgA^{-},*

Δ*yopJ ksgA^{-},* and *yopJ^{C172A} ksgA^{-}* strains (**Figure 4A, B**). Secondly, the number of

YopJ^{F177} and YopJ-deficient bacteria (both YopJ^{-} and YopJ^{C172A} mutants) generally

increased over the 11 day time course in target organs (**Figure 4A, B**), whereas the

YopJ^{F177L} KsgA^{-} bacteria were either slower to accumulate in tissues (liver, **Figure 4A**)

or diminished during this time frame (spleen, **Figure 4B**). These results are consistent

with YopJ triggering host immune responses that inhibit bacterial replication and/or

remove bacteria from the tissue, while the absence of YopJ activity allows the bacteria

to replicate unchecked in tissues.

In addition to causing macrophage cytotoxicity, YopJ can also inhibit macrophage

production of cytokines [38-41], such that the increased mortality of mice exposed to

ksgA^{-} bacteria carrying either a null or inactive *yopJ* allele could result from a massive

overproduction of cytokines, also known as a cytokine storm [42-44]. However, we did

not find any evidence indicating that the absence of functional YopJ triggers a mortality-

inducing cytokine storm *in vivo* (**Supplementary Figure 2**),.

To determine if cell death levels *in vivo* recapitulated what we had observed with

cultured macrophages exposed to the different *Y. pseudotuberculosis* strains, we used

the TUNEL assay to detect dead mammalian cells in tissue sections from colonized

mice. Mice were sacrificed at day 3 post-inoculation for cell death evaluation to ensure

equivalent burden levels between mice inoculated with the different strains (as observed

in **Figure 4**). This analysis revealed that at 3 days post infection, spleens containing

YopJ^{WT} or YopJ^{F177L} KsgA$^-$ bacteria had increased levels of TUNEL$^+$ cells as compared

to spleens containing the YopJ$^-$ KsgA$^-$ bacteria (**Figure 5A-D)**, although the strain

expressing catalytically dead YopJ did not perfectly phenocopy the Δ*yopJ$^-$ ksgA$^-$* strain

(**Figure 5D**). These results indicated that the increased survival of mice infected with

KsgA$^-$ YopJ^{F177L} was significantly correlated with the amount of TUNEL staining present

in their spleens at day 3 post infection. Conversely, a decrease in the amount of TUNEL

staining present in the spleens at this time point was associated with the early death of

mice infected with either *ksgA$^-$ ΔyopJ* or with *ksgA$^-$ yopJ^{C172A}*.

The anti-apoptotic drug suramin inhibits YopJ-induced macrophage death and

enhances bacterial virulence

Our results suggest that YopJ-induced cytotoxicity is detrimental to *Y.*

pseudotuberculosis pathogenesis, such that enhancing or inhibiting of host cell death *in*

vivo should impair or promote *Y. pseudotuberculosis* virulence, respectively. To test this

hypothesis, we first considered if artificially increasing macrophage death in mice

exposed to YopJ-null *Y. pseudotuberculosis* would rescue the mice from illness.

Clodronate-containing liposomes are commonly used to deplete macrophages *in vivo* –

depletion results because the clodronate induces global macrophage death [45]; we

confirmed that mice exposed to clodronate liposomes showed massive levels of

apoptotic cells in tissues (data not shown). We inoculated mice with *ksgA⁻ ΔyopJ Y.*

pseudotuberculosis and 6 hours later, delivered a one-time injection of clodronate

liposomones. The liposome-treated animals showed a slightly increased rate of

mortality as compared to mock-treated animals (**Supplementary Figure 3**), suggesting

that the global depletion of macrophages masked any consequence of increased

macrophage apoptosis.

We next attempted to diminish host cell death levels *in vivo*, to determine if this would

enhance *Y. pseudotuberdulosis* virulence. We turned to the apoptosis inhibitor, suramin,

which is a polysulfonated urea derivative capable of inhibiting death receptor-induced

apoptosis and apoptosis-mediated liver damage [46, 47]. Suramin blocked YopJ-

induced death of cultured macrophages in a dose-dependent manner (**Figure 6A**), at

the same levels previously shown to block hepatic cell cytotoxicity induced by death

receptors CD95, TRAIL-R1 and TRAIL-R2 [46]. To evaluate how suramin treatment

affected *Y. pseudotuberculosis* virulence *in vivo*, mice were inoculated with bacteria,

then injected with suramin 30 minutes to 1 hour post-inoculation. Suramin had no effect

on naïve mice, but dramatically increased mortality of mice exposed to KsgA⁻ bacteria

(**Figure 6B**). This effect was YopJ-independent, however, as mice inoculated with YopJ⁻

KsgA⁻ bacteria and subsequently injected with suramin showed a similar increase in

time-to-death (**Figure 6B**). Strikingly, gross pathological differences were apparent after

examination of H&E stained liver sections taken from KsgA⁻-infected mice treated either

with suramin or PBS. The suramin-treated/infected mice had significantly fewer

microabcesses visible in liver sections when compared to PBS-treated/infected mice

(**Figure 6C-E**). Confirming that suramin successfully impaired host cell death, liver

sections taken from suramin-treated mice at 3 days post-bacterial inoculation showed

significantly fewer TUNEL+ cells within abcesses than untreated mice (**Figure 6F-H**).

These results demonstrate that inhibition of host cell death dramatically worsens the

survival outcome for mice experiencing *Y. pseudotuberculosis*-caused illness, and

strongly supports the premise that host cell death protects the host and checks the

pathogen.

Discussion

Two surprising things emerge from our study. Firstly, positive selection of YopP/J

residues has occurred directionally with *Yersinia* evolution, and has yielded an isoform

with diminished function. Secondly, YopJ function is inversely related to bacterial

pathogenesis – a hypercytotoxic YopJ variant attenuates *Yersinia* virulence in

mammals, while non-functional YopP/J isoforms (or null alleles) confer hypervirulence in

the context of sub-acute infection. The evolutionary forces that selected for bacteria with

reduced YopJ activity or secretion are obscure, but this obscurity does not negate that

positive selection of YopP/J codons has indeed occurred and these polymorphisms

influence the degree of YopP/J-induced macrophage death and/or *Yersinia* virulence

towards animals (**Figure 1**, see also [24, 26, 48, 49]). While we observed the YopJ

phenotype in a model of sub-acute *Y. pseudotuberculosis* illness in inbred mice, we

speculate that the selective niche was/is necessarily a warm-blooded wild animal. This

speculation is based on the knowledge that YopJ and other Yops are not expressed at

the ambient temperatures typical of fleas and soil [50, 51], the other niches known to

harbor *Y. pestis [52]*, such that YopJ likely evolved to target mammalian proteins.

How is YopJ activity diminishing *Y. pseudotuberculosis* virulence? Recent evidence has

shown that YopJ can block innate signaling interfering with TGF-β-activated kinase 1

(TAK1) in addition to interfering with NFκB signaling and MAPK pathways [53-56].

YopJ blockade of the aforementioned signaling events and in the presence of TRIF or

TNF signaling has been proposed to trigger receptor-interacting protein kinase 1

(RIPK1) and caspase-8 dependent apoptosis [55, 57, 58]. Collectively, the above

studies have shown that Yersinia mediated RIPK1/caspase-8 mediated apoptosis can

be protective for the host. Although the precise mechanism of Yersinia induced cell

death was not addressed in this study, our results similarly suggest that tailoring

Yersina's ability to induce host cell cytotoxity may have an impact on the *in vivo*

pathogenesis. We further speculate that the mechanism is similar to the one we

observed previously in our studies of CD8+ T cell-mediated clearance of *Y.*

pseudotuberculosis-associated target cells [59]. In this model, bacteria expressing

wildtype or ancestral/hyperfunctional YopJ attach to the host cell and inject the YopJ

protein, leading to host cell death. As dead host cells are rapidly efferocytosed by

neighboring activated phagocytes, the attached bacteria are also removed in the

process of efferocytosis, leading to pathogen clearance (**Figure 7A**). In contrast,

bacteria expressing no or catalytically dead YopJ would fail to induce host cell death,

meaning the host cell would not be removed and bacteria could persist longer (**Figure**

7B).

Given that the YopJ activity can be detrimental to *Yersinia* virulence, what then is the

purpose of YopJ-induced macrophage death for *Yersinia*, when the absence of YopJ so

clearly can increase *Yersinia* virulence during sub-acute illness? The continued

maintenance of the *yopJ* gene in the *Y. pseudotuberculosis* and *Y. pestis* genomes, the

latter of which has undergone marked decay [5, 6, 32], suggests that the existence of a

niche where YopJ function remains beneficial to *Y. pestis*. One possibility is that YopJ's

beneficial or detrimental effects on bacterial virulence depend on the host tissue

environment. This speculation is supported by prior observations that YopJ-null *Y.*

pseudotuberculosis bacteria have mild defects in virulence via oral inoculation, but no

defects via parental administration [18, 20, 24], suggesting that YopJ promotes

extraintestinal dissemination of *Yersinia* or is irrelevant in systemic tissues during acute

infection, a hypothesis supported by recent observations that YopJ alters intestinal

permeability and promotes *Yersinia* extraintestinal dissemination [60]. Other findings

also point to a role for YopJ in the initial local infection site [25]. Additionally, the role of

YopJ in *Y. pestis* has been shown to be critical for dissemination and in triggering

necroptosis rather than apoptosis. This study also showed that a ΔYopJ strain of *Y.*

pestis led to host survival rather than death as has been observed with ΔYopJ strains of

Y. pseudotuberculosis[57, 61]. Our findings with sub-acute systemic illness, however,

demonstrate that YopJ actually hinders or impairs *Yersinia* virulence in systemic

tissues. We speculate that YopJ activity serves as self-regulating mechanism for

Yersinia pathogenesis by allowing bacterial persistence in a resistant host or tolerant

tissue, possibly in a relationship approaching commensalism or at least mild parasitism.

In other words, an existence comparable to that observed with other chronic illness-

causing pathogens – for example, *Salmonella typhi*, which persists in gall bladders and

can be shed to dissemination to new environments and hosts. Another *Yersinia* Yop can

also subvert host immune recognition to promote bacterial virulence – YopK inhibits

inflammasome recognition of the *Yersinia* type III secretion system and thus prevents

bacterial clearance [7], although it has been proposed that YopK acts by regulating

T3SS translocation, rather than by direct activity towards target proteins within the host

cytoplasm [62, 63]. Additionally, there is one report demonstrating that loss of YopT

renders *Y. enterocolitica* hypervirulent [23]. It may be that some Yops act to indirectly

counter those Yops that promote virulence (i.e. YopE, YopH).

Our observations of mice exposed to non-cytotoxic KsgA$^-$ bacteria, and of mice treated

with the death inhibitory chemical suramin during *Y. pseudutuberculosis* infection,

strongly suggest that host cell death is beneficial for host survival and detrimental to

bacterial replication and/or existence. Our experiments clearly show that suramin-

treated/*ksgA*$^-$-infected animals have a marked reduction in dead host cells (TUNEL$^+$),

suggesting that the diminished apoptosis plays at least a partial role in the increased

susceptibility of suramin-treated animals to infection. Our efforts to reduce host cell

death levels with caspase inhibitors during *in vivo* infection were unsuccessful (data not

shown), perhaps due to functional redundancy in the host cell death program

components or the known consequence of increased necrosis/necroptosis following

caspase inhibition [64].

YopP/J has homologues in other bacterial pathogens of animals[65], including *Vibrio*

parahaemolyticus (VopP/A) [60, 66], *Aeromonas salmonicida* (AopP) [60], and

Bartonella quintana strain Toulouse (YopP) [67]. One interesting YopJ homologue is

AvrA, which is found in numerous species, subspecies and serovars of the genus

Salmonella[33, 68]. Like YopJ, AvrA is an deubiquitinase and acetyltransferase capable

modifying and inhibiting MAPKK and kinases in the JNK and NFκB signaling pathways

[69, 70], leading to modulation of host cell inflammatory responses in the intestine [71].

Interestingly, unlike YopJ, AvrA actually prevents apoptosis in host cells [70],

suggesting that sequence differences between the two proteins may explain the

divergence in function. Although AvrA does possess a leucine at the position that aligns

with YopJ residue 177, suggesting that AvrA should be hyperfunctional relative to

YopJ^{F177}, prior studies have shown that AvrA affects neither cytokine expression nor

plays a role in macrophage killing when expressed by either *Salmonella* or *Yersinia [72]*.

However, it should be noted that the *avrA* alleles used in these studies all possess a

three nucleotide deletion that results in the absence of a leucine residue at position 139

[73], also data not shown), which may explain why AvrA reportedly differs from YopJ in

activity and function. However, regardless of the differences in sequence and tissue

culture cell interactions, AvrA has two major features in common with YopJ – ability to

attenuate virulence [70] and promote chronic infection [69] in mouse models of

salmonellosis.

The ability of YopJ and AvrA to attenuate bacterial virulence is a feature typical of those

YopJ homologues found in plant-associated bacteria. YopJ is a founding member of the

YopJ/HopZ/AvrRxv superfamily – a family of T3SS-translocated proteins, expressed by

both bacterial plant pathogens and symbionts, that are recognized by plant resistance

(R) proteins during a plant defense response called the hypersensitive response (HR)

[65]. Induced host cell death is a critical feature of HR, and serves to limit spread of the

bacterium beyond the initial infection site [74]. YopJ homologues in plant-associated

bacteria induce plant cell death in diverse hosts, including *Arabidopsis*, soybean, rice,

tobacco and pepper plants, among others, by targeting specific plant proteins for

modification [65]. The requirement of a common catalytic triad suggests the plant YopJ

homologues have similar enzymatic activity as YopJ. Supporting this, PopP2 from

Ralstonia solanacearum has been shown to acetylate the *Aradopsis* RRS1-R resistance

protein [71], *Xanthomonas campestris* pathovar *vesicatoria* AvrXv4 is a SUMO protease

[75], and members of the HopZ family from *Pseudomonas syringae* have been shown to

have protease or acetyltransferase activity [31, 76].

Another observation from our studies is reminiscent of plant R protein recognition of

effectors. R proteins sense "altered self" in several ways – 1) direct modification of the R

proteins by injected bacterial effector proteins, 2) indirect recognition of effector protein-

driven modifications of other host proteins or 3) indirect recognition of host cell

molecules released in effector translocation or bacterial infection [77]. We find it

intriguing that bacteria expressing YopJ^{C172A} did not phenocopy the YopJ⁻ bacteria for

mouse mortality, and were <u>slightly</u> attenuated relative to the YopJ⁻ bacteria. This has

been seen before with YopO/YpkA – in a study of YopO/YpkA mutant strains - *Y.*

pseudotuberculosis expressing non-functional YopO was attenuated, whereas a YopO-

null strain was unaffected for virulence [78]. We speculate that a non-functional YopJ

protein may be perceived differently by the host than either a functional YopJ or the

absence of YopJ, perhaps due to host recognition of the YopJ^{C172A} protein itself or

recognition of YopJ^{C172A} binding to target host proteins. This recognition would then

induce a cytosolic innate immune response to the foreign (albeit inactive) YopJ protein.

This speculation remains to be addressed.

Phylogenetic and phylogenomic analyses are powerful approaches to identify

signatures of molecular evolution in the genes and genomes of bacterial pathogens. A

hallmark paper from Ma et. al. determined that the C-terminus of HopZ, the YopJ

homologue found multiple *P. syringae* pathovars, has been under strong positive

selection [31], providing further evidence that bacterial effectors that modulate host cell

death are shaped by strong evolutionary forces. Interestingly, of the 14 pathovar HopZ

sequences examined, all possessed a leucine at the position corresponding to YopJ

177 [31], suggesting that expression of a highly-functional HopZ is evolutionarily

advantageous to multiple *P. syringae* pathovars. A recent phylogenomics study of 60

Chlamydia trachomatis strains detected signatures of positive selection in multiple T3SS

effectors, and also revealed that this selection is driving *C. trachomatis* evolution

towards niche-specific adaptations (colonization of particular cell types or tissues)[30]. It

is almost certain that global or gene-specific analyses of molecular evolution in bacterial

effector proteins will reveal evidence of positive selection, even if the selective forces

and niches remain obscure.

Our work extends the findings of two recent *Yersinia* phylogenomic studies of note. Gu

et. al. examined the evolution and divergence of multigene families from 5 *Yersinia*

strains (4 *Y. pestis*, 1 *Y. pseudotuberculosis*) and found evidence of functional

diversities in genes associated with pathogenicity [79]. While suggestive of positive

selection, the authors did not individually analyze Yops for microevolutionary signatures.

In the second study, Cui et. al. analyzed the genomes of 133 *Y. pestis* strains and

identified 2,326 single-nucleotide polymorphisms (SNPs) that appear to have been fixed

by neutral evolution rather than Darwinian selection [80]. However, the authors only

analyzed SNPs from the core-genome of *Y. pestis*, not the accessory genome that

includes the extrachromosomal plasmid encoding the *yop* genes, meaning that *yopJ*

was not included in their analysis. Others have found at least 1 chromosomally-encoded

Yersinia protein has undergone positive selection at the codon level, as analysis of

OmpF porin sequence from 73 different *Yersinia* strains found positively selected

residues in several surface-exposed OmpF domains [81]. As such, our findings

constitute the first description of diversifying selection in the accessory genome of any

Yersinia species member. Although our analysis was performed on only a small subset

of *Yersinia* YopP/J sequences, the site identified here, codon 177, showed a significant

level of positive selection, indicating the validity of our findings.

Our observations prompt the intriguing speculation that other Yops will also show

evidence of molecular evolution. Given that the number of translocated *Yersinia*

proteins is low (~6-8 proteins), it seems likely that the importance of each individual

effector is therefore increased, such that sequence divergence in response to any

selective pressure would be greater. Microevolution within the sequences of other Yops,

which generally function as pro-virulence factors, could result in Yops with increased

functional abilities that confer increased virulence to individual strains or species. If true,

it would present an interesting tug of war between pro-virulence Yops and anti-virulence

YopP/J, the outcome of which would likely dependent on the niche.

Lastly, the degree of positive selection in host proteins targeted by YopJ or other Yops

is completely unknown. Recent observations of the molecular "arms race" between host

and viral proteins demonstrated that the host anti-viral protein PKR (protein kinase R)

has been subjected to intense episodes of positive selection in primates, in order to

evade viral mimics of the PKR binding partner eIF2α [82]. Regarding the latter, it is

interesting to note that the activities both YopJ and YpkA/YopO converge on a kinase of

eIF2α [83]. Further identification of YopP/J-specific target proteins and evaluation of

sequence divergence through the primate lineage will reveal the degree of YopP/J-

driven evolution of host immune response proteins.

Materials And Methods

Bacterial Strains and Growth Conditions

Strains used in this study are indicated in Table 1, primers used for strain construction

are indicated in Table 2. Lysogeny Broth (LB) and LB agar plates were made using the

Lennox formulation (5g/L NaCl), and when necessary, selective antibiotics included at

the following concentrations: kanamycin 30µg/mL, ampicillin 100µg/mL, irgasan

1μg/mL, chloramphenicol 20μg/mL. The *Y. pseudotuberculosis* strain YPIII pIB1 was

the parent of all strains described here. The *ksgA⁻* strain has been described and used

by us previously [84, 85] and contains a kanamycin-resistance-encoding transposon

inserted into the *ksgA* gene. To generate the *ksgA⁻ ΔyopJ* strain, we used a previously-

described suicide vector construct, pCVD442-Δ*yopJ* (kind gift of Dr. Joan Mecsas, Tufts

University), in which the upstream and downstream DNA flanking *yopJ* open-reading

frame was fused together with only the first and last codon of *yopJ* remaining [86].

Introduction of this unmarked *yopJ* deletion allele into the *ksgA⁻* genome was

accomplished via allelic exchange, using pCVD442-encoded ampicillin and sucrose

resistance as the selectable and counter-selectable markers, respectively. Briefly, the

suicide vector was transferred from *Escherichia coli* strain SM10λpir to *Y.*

pseudotuberculosis strain *ksgA⁻* by conjugation, and ampicillin-resistant/irgasan-

resistant integrants selected. Integrants were grown overnight in the absence of plasmid

selection, then dilutions of the culture placed on counter-selection agar plates

containing 10% sucrose and lacking salt. Sucrose-resistant colonies were confirmed to

be ampicillin-sensitive and the presence or absence of the Δ*yopJ* determined by PCR

screening. To generate *yopJ* alleles carrying mutations in codons 172 and 177, a

fragment of the *yopJ* gene was first cloned by PCR into pACYC184, then site-directed

mutagenesis performed using long-range PCR with primers containing the mutant

sequence and Dpn to remove the vector template. After confirming the sequence

change, the mutated *yopJ* sequence was transferred to pCVD442, allelic exchange

performed as described above, and the presence or absence of the mutant *yopJ*

codons was determined by sequencing amplified genomic DNA. Bacteria were grown

for cultured macrophage experiments as follows. After 2 days growth on LB agar plates

from glycerol stocks, single colonies from the indicated strains were inoculated into 2

mL LB broth and incubated at 26°C for 15-18 hours with rotation. Overnight cultures

were back-diluted to an estimated OD_{600} of 0.2 into low-Ca^{2+} medium (2XYT broth

containing 20 mM of the Ca^{2+} chelator sodium oxalate and 20mM $MgCl_2$, then grown

with constant rotation for 1.5 hours at 26°C and 1.5 hours at 37°C. Cultures were then

adjusted to the desired concentration in macrophage media and used for macrophage

challenges as described below. Bacteria were grown for mouse experiments as follows.

After 2 days growth on LB agar plates from glycerol stocks, single colonies were

inoculated into 2 mL LB broth and incubated at 26°C for 15-18 hours with rotation.

Overnight cultures were then adjusted to the desired concentration in PBS and used for

mouse challenges as described below (after removing a sample for accurate

determination of colony-forming units per mL i.e. CFU/mL).

Codon-level selection analysis of YopJ sequences across multiple

Yersinia lineages.

To investigate the evolutionary history of YopJ, orthologs were found by using reciprocal

blast searches across 15 sequenced genomes, including extrachromosomal elements,

and 2 individual *yopJ* loci (NC_003131.gbk, NC_003132.gbk, NC_003134.gbk,

NC_003143.gbk, NC_004088.gbk, NC_004838.gbk, NC_005810.gbk, NC_005813.gbk,

NC_005814.gbk, NC_005815.gbk, NC_005816.gbk, NC_006153.gbk, NC_006154.gbk,

NC_006155.gbk, NC_008118.gbk, NC_008119.gbk, NC_008120.gbk, NC_008121.gbk,

NC_008122.gbk, NC_008149.gbk, NC_008150.gbk, NC_008791.gbk, NC_008800.gbk,

NC_009377.gbk, NC_009378.gbk, NC_009381.gbk, NC_009704.gbk, NC_009705.gbk,

NC_009708.gbk, NC_010157.gbk, NC_010158.gbk, NC_010159.gbk, NC_010465.gbk,

NC_010634.gbk, NC_010635.gbk, NC_014017.gbk, NC_014022.gbk, NC_014027.gbk,

NC_014029.gbk, NC_015224.gbk, NC_015475.gbk, NC_017153.gbk, NC_017154.gbk,

NC_017155.gbk, NC_017156.gbk, NC_017157.gbk, NC_017158.gbk, NC_017159.gbk,

NC_017160.gbk, NC_017168.gbk, NC_017169.gbk, NC_017170.gbk, NC_017263.gbk,

NC_017264.gbk, NC_017265.gbk, NC_017266.gbk, NC_017564.gbk, NC_017565.gbk).

Protein sequences were aligned using Clustalw and nucleotide sequences for each

YopJ ortholog were aligned based on their amino acid translation (using in house

scripts). These nucleotide alignments were used as input to RAxML[83], where

maximum likelihood phylogenies were generated (GTRGAMMA model of substitution).

In order to identify those YopJ residues under selection and the direction and strength

of selection at the codon level, the frequency of nonsynonymous and synonymous

changes and their ratio (dN/dS or ω) were calculated using Phylogenetic Analysis by

Maximum Likelihood (PAML v4.7) software using the Goldman and Yang amino acid

substitution model (codeML) and these same nucleotide alignments [84], In house Perl

scripts were used to parse the codeML output files to identify residues predicted to have

undergone diversifying selection in YopJ (using maximum likelihood ratio tests before

analyzing the Bayes Empirical Bayes results).

Derivation of Bone Marrow Macrophages

Bone marrow-derived macrophages (BMDM) were generated by culturing bone marrow

cells freshly-harvested from the femurs and tibias of C57BL/6 mice in supplemented

DMEM as previously described [59]. Macrophages were harvested on day 6 of the

derivation protocol and seeded at a density of 5 x 10^4 cells/100µL/well of a 96 well plate

in supplemented DMEM lacking antibiotics and allowed to adhere to the plastic

overnight. Occasionally, at day 6, macrophages were cryofrozen in 5% DMSO/95%

fetal-bovine serum (FBS) at a concentration of 1x10^7 cells/mL and subsequently thawed

for seeding.

Macrophage Infections, Treatments and Cytotoxicity Assay

Y. pseudotuberculosis, grown as described above, was added to adherent

macrophages at a multiplicity of infection (MOI) of 100:1 in DMEM without phenol red +

1% FBS for 45 minutes to 1 hour. Gentamicin was added to a final concentration of 100

µg/mL and the cells were allowed to incubate an additional 2 hours. For suramin

experiments, BMDM were first pretreated with various concentrations of suramin

(Sigma, St. Louis, MO) for 1 hour, then exposed to bacteria at an MOI of 100:1 in the

presence or absence of suramin at the indicated concentrations. Macrophage death

was evaluated by detection of the cytoplasmic enzyme lactate dehyrogenase (LDH) in

the culture supernatants similarly as previously described[59]. Briefly, macrophages

were spun down at 250xg for 2 minutes and 50µL supernatant carefully removed from

each well, transferred to a new 96-well plate, and immediately assayed for the presence

of LDH using the Cytotox 96 Assay kit (Promega, Madison, WI) according to the

manufacturer's recommendations. Note that controls cells were treated with 1% Triton-X

100 to liberate cytoplasmic contents approximately 15 minutes prior to supernatant

harvest. 50µL of reconstituted substrate solution (containing 2-(4-Iodophenyl)-3-(4-

nitrophenyl)-5-phenyl-2H-tetrazolium chloride, or INT) was added to the harvested

supernatants, the reaction incubated at room temperature in the dark for 30 minutes,

then the enzymatic conversion of INT into a red formazan product terminated by

addition of 50μL stop solution (1M acetic acid) and excess bubbles in each well

punctured. Absorbance was measured at 490 nm in a BioTek Synergy H4 plate reader

(BioTek, Winooski, VT). % Cytotoxicity was calculated as follows: 100✕ [(experimental

release – effector T cell spontaneous release – target cell spontaneous release)/(target

cell maximum release – target cell spontaneous release)].

Mouse Infections and Treatments

6-8 week old female C57BL/6 mice were purchased from the National Cancer Institute

colony at Charles River. Mice were allowed to rest for 7 days after arrival, then infected

intravenously via tail vein injection with a 200 μL inoculum containing between 3.5×10^2

and 2×10^3 colony-forming units (CFU) in PBS of the indicated *Y. pseudotuberculosis*

strain. Mice were then visually inspected for signs of illness including lethargy and

piloerection and weighed daily to monitor the disease progression. For suramin

experiments, mice were infected as before, then intraperitoneally injected once with 5

mg suramin/400 μL PBS or 400 μL PBS alone 30 minutes to 1 hour post-infection. For

clodronate experiments, clodronate-containing liposomes were prepared as described

below, then 200 μl of liposomes were injected intravenously 6 hours post infection. All

animal experiments were performed in accordance with the NIH Guide for the Care and

Use of Laboratory Animals and were approved by the UTHSCSA Institutional Animal

Care and Use Committee.

Preparation and Characterization of Clodronate Liposomes

Liposomes containing dichloromethylene bisphosphonate (clodronate) or control empty liposomes containing phosphate buffered saline (PBS) were prepared from reported protocols with slight modifications[45, 87]. In two 500 mL round bottom flasks, 96 mg cholesterol (Calbiochem, San Diego, CA) was added and dissolved with 10 mL chloroform (HPLC grade, Fisher Scientific, Fair Lawn, NJ). Then 4.14 mL of egg phosphatidylcholine in chloroform (25 mg/ mL; Avanti Polar Lipids, Alabaster, AL) was added. Lipid films were formed following removal of chloroform by rotary evaporation. The lipid films were placed under vacuum for 1 h to remove residual chloroform. The lipid films were rehydrated with 12 mL of 0.7 M clodronate (Sigma Aldrich, St Louis, MO) prepared in degassed sterile water for injection (Hospira, Lake Forest, IL) and pH adjusted to 7.0, or degassed Dulbecco's PBS, pH 7.4 (Invitrogen, Carlsbad, CA). The flasks were swirled until the lipids were in suspension. The lipid suspensions were kept at room temperature for 2 h with occasional swirling, sonicated in a water bath for 4 minutes (Branson, Danbury, CT) and then kept at room temperature for 2 h. The liposome samples were then extruded through polycarbonate filters (2 μ, 2 times; 1 μ, 4 times) using an extruder (Lipex, Northern Lipids, Vancouver, Canada) and collected in a new tube. After flushing the liposome samples with argon, the samples were stored overnight at 4° C. The next day the liposomes were transferred to microcentrifuge tubes and centrifuged in an Allegra 21R centrifuge (Beckman Coulter, Brea, CA) at 15,300 rpm for 30 minutes at 10°C. The supernatants were removed and the pellets resuspended with degassed PBS. The samples were washed 4 times. After the last

wash, the liposome pellets were resuspended in degassed PBS to a final volume of 4.8

mL, flushed with argon and stored at 4°C until needed. Liposomes were used for the

study within 2 weeks of preparation. Liposome diameter was measured at 488 nm with

a DynaPro dynamic light scattering system (Wyatt Technology, Santa Barbara, CA).

The diameters for clodronate liposomes and control PBS liposomes were 968 ± 28 nm

and 790 ± 41 nm, respectively. Phospholipid content was measured using

phosphatidylcholine colorimetric assay kit (Cayman Chemical, Ann Arbor, MI) and

determined to be 14.5 mg/mL for the clodronate liposomes and 15.4 mg/mL for the PBS

liposomes. The amount of liposome-encapsulated clodronate was determined to be

12.7 mg/mL [45]. In pilot experiments, clodronate liposomes were confirmed to induce

macrophage death by TUNEL staining in spleens of non-infected mice (data not

shown).

Organ Bacterial Burden Assay

At the noted times post-infection, mice were sacrificed by cervical dislocation, spleens

and livers aseptically removed, transferred to sterile PBS from mice at various time

points after infection and weighed. Tissues were mechanically disrupted using a tissue

homogenizer (Omni, Marietta GA), serially diluted in sterile PBS and 100 μL of the

dilutions plated on nutrient LB agar plates. After 48 hours at 26°C, bacterial colonies

were enumerated and CFU normalized to weight of tissue in grams. Note that as each

animal may have had an organ weight different from the other animals, the CFU values

at or below the limit of detection (10 CFU per mL of organ homogenate) were

occasionally different between individual animals. The limit of detection in the spleen

was (90 CFU/g/spleen or 1.95 Log CFU) and in the liver was (22 CFU/g/Liver or 1.33

Log CFU).

Cytokine Analysis

Cytokine analysis was performed on serum and tissue homogenates from day 3 and

day 9 post-inoculation. Briefly, blood was collected via cardiac puncture and allowed to

clot for 30 minutes. Serum was collected after centrifugation at 7000 rpm for 7 minutes

and stored at -80°C. Cytokines were assayed using the Mouse Cytokine 20-plex panel

strictly adhering the manufacturer's protocol (Life Technologies, Carlsbad CA). Cytokine

levels were detected using a Bio-Plex 200 system and analyzed using Bio-Plex

Manager 6.0 software (Bio-Rad, Hercules CA).

Terminal Deoxynucleotidyl Transferase dUTP Nick End Labeling (TUNEL) Assay

Spleens were harvested at 3 days post-infection and were snap-frozen in the cyrostat

sectioning medium Tissue-Tek® O.C.T.™ (Sakura, Torrance CA). Livers were removed

and placed in 10% buffered formalin for at least three days and then sent to the

UTHSCSA Histopathology Laboratory to be embedded in paraffin and cut into 4 micron

sections. Some slides with liver sections were stained with hemotoxilin and eosin and

others were processed for TUNEL staining with the ApopTag kit as outlined below.

OCT-frozen spleen sections were cut with a cryotome (Thermo Scientific Shandon,

Waltham MA), placed on slides and allowed to air-dry overnight. The slide-affixed tissue

sections were then fixed in acetone for 20 seconds at room temperature, allowed to dry,

then wrapped in foil and stored at -80°C. Subsequently, sections were stained for

nicked DNA, to label TUNEL^{+} cells, using the ApopTag kit with included buffers (EMD

Millipore, Billerica MA). Briefly, slides were removed from -80°C and allowed to warm up

to room temperature. Slides were then fixed in 1% paraformaldehyde for 10 minutes at

room temperature and washed in PBS twice. Tissue sections were not allowed to dry

after this point. Tissue sections were then circled with a hydrophobic PapPen (Kiyota

International, Elk Grove Village, IL) and equilibration buffer was incubated on the

section for at least 10 seconds. TdT enzyme solution (10% enzyme + 20% PBS + 70%

reaction buffer) was applied for 60 minutes in a humidity chamber at 37°C. Stop solution

was then applied for 10 minutes at room temperature followed by three washes with

PBS. A solution containing anti-digoxigenin fluorescein antibody (53% blocking solution

+ 47% anti-digoxigenin fluorescein conjugate) was applied and allowed to incubate for

30 minutes in a dark humidity chamber. Slides were then protected from light as much

as possible from this point forward. Slides were then washed three times in PBS and

mounted with ProLong Gold Antifade reagent containing DAPI (Life Technologies,

Carlsbad CA). Sections from a given experiment were stained on the same day.

Microscopy and Image Capture, Analysis

Stained tissue sections were visualized using either an EC plan neofluar 10x or Plan-

Apochromat 20x/.8 objective on a Zeiss AxioImager Z1 epifluorescent microscope (Carl

Zeiss, Thornwood, NY). Images were captured using a Zeiss AxioXam MRm Rev3

and/or MRc cameras and analyzed using Zeiss AxioVision release 4.5 software.

Sections from a given experiment were imaged on the same day with the same

exposure times for image acquisition to ensure consistency between samples. ImageJ

[88] was used to measure the pixel intensity in each captured image.

Data and Statistical Analysis

Prism 5 (GraphPad Software, La Jolla CA) was used for graphing and statistical

analysis. Survival curves were estimated using the Kaplan Meier method and

significance calculated using the log-rank test. The nonparametric Mann-Whitney U test

and unpaired Student's t test were used to determine statistical differences between

groups of data from animal and tissue culture experiments, respectively. Image

quantification data was evaluated for significance using the Mann-Whitney U test.

Acknowledgements

We thank Ian Morris and Michael Berton for use and instruction of the Zeiss AxioImager

Z1 microscope. We thank Joan Mecsas, Peter Dube and Angel Cantwell for helpful

discussions and critical analysis of the manuscript. This work was supported by National

Institutes of Health, Grant AI085116 from the NIAID to M.A.B., and Grant AI110684

from the National Institute of Allergy and Infectious Diseases and HHMI support of R.R.I.

Figure Legends

Figure 1. YopP/J is divergent between *Yersinia* species and shows evidence of

positive selection

(A) The phylogeny of three *Yersinia* species (*Y. enterocolitica,* 3 strains, *Y. pseudotubercolosis,* 2 strains, and *Y. pestis,* 9 strains) generated using the YopJ

primary sequences. A considerable amount of change in the YopJ homologs has

occurred during the evolution of these pathogens; some of these changes alter protein

encoding (dN, non-synonymous nucleotide substitutions) while some do not (dS,

synonymous substitutions). Evidence of adaptative selection was found across the

aligned ortholog set by calculating the ratio of dN/dS in pairwise comparisons of

orthologs. To increase the sensitivity of our analysis we also calculated variable dN/dS

(ω) over all sites (models M7 vs. M8) and used the likelihood ratio test to identify

statistically significant evidence for diversifying selection ($\chi2 = 14$; df = 2; p < 0.01).

Codon 177 was identified as under positive selection in a Bayes empirical Bayes (BEB)

analysis and amino acid polymorphism at that site is mapped onto the phylogeny of

these *Yersinia* strains ("L" in blue and "F" in red). (B) Alignment of the relevant region of

YopP/J homologs from these 14 *Yersinia* strains from the three human pathogenic

species. YopP/J was identified as having experienced positive selection at residue 177.

Figure 2. Ancestral version of YopP/J confers enhanced macrophage death and

attenuated virulence. (A) Bone marrow derived macrophages were exposed to either

ksgA$^-$, ksgA$^-$ ΔyopJ, or *ksgA$^-$ yopJ^{F177L} Y. pseudotuberculosis* at an MOI of 100:1. LDH

release was measured to determine the cytotoxicity of these strains in macrophages.

(B) C57BL/6 mice were intravenously challenged with *ksgA$^-$* (*n*=12) or *ksgA$^-$ yopJ^{F177L}*

(*n*=11) *Y. pseudotuberculosis*. Morbidity and mortality were followed after challenge and

data shown represents the percent survival of each group. In vivo data is compiled from

two independent experiments.

Figure 3. Loss of YopJ reverses the attenuation phenotype of *ksgA⁻ Y.*

pseudotuberculosis. (A) Bone marrow derived macrophages from C57BL/6 mice were

infected with either *ksgA⁻* or *ksgA⁻ ΔyopJ Y. pseudotuberculosis* at an MOI of 100:1 as

described in the Methods section. LDH release was measured to determine the

cytotoxicity of these strains in macrophages. (B) C57BL/6 mice were intravenously

challenged with 1×10^3 CFU of either *ksgA⁻* (*n*=19) or *ksgA⁻ ΔyopJ* (*n*=22) *Y.*

pseudotuberculosis and morbidity and mortality was followed after infection and data

shown represents the percent survival of each group. (C) Bone marrow derived

macrophages were infected with either *ksgA⁻*, *ksgA⁻ ΔyopJ*, or *ksgA⁻ yopJ^{C172A} Y.*

pseudotuberculosis at an MOI of 100:1. LDH release was measured to determine the

cytotoxicity of these strains in macrophages. (D) C57BL/6 mice were intravenously

challenged with 1×10^3 CFU of either *ksgA⁻* (*n*=18), *ksgA⁻ ΔyopJ* (*n*=22), or *ksgA⁻*

yopJ^{C172A} (*n*=18) *Y. pseudotuberculosis* and scored as in (B) and also scored for (E) the

median time until death for each animal. In vivo data is compiled from 4 independent

experiments.

Figure 4. Enhanced bacterial burden *in vivo* results with *Y. pseudotuberculosis*

lacks functional YopJ. (A) Liver and (B) spleen were isolated at multiple time points

from C57BL/6 mice after post-challenge (1×10^3 CFU) with *ksgA⁻*, *ksgA⁻ ΔyopJ*, *ksgA⁻*

yopJ^{C172A}, or *ksgA⁻ yopJ^{F177L} Y. pseudotuberculosis*. The tissues were homogenized,

serially diluted and plated on LB plates. Burdens recovered in each tissue from each

mouse were above the limit of detection. Data shown is from 5 mice/group and is

representative of two independent experiments.

Figure 5. Ancestral YopP/J variant causes increased levels of cytotoxicity *in vivo*.

6-8 week old female C57BL/6 mice were infected intravenously (1×10^3 CFU) with strains

of *Y. pseudotuberculosis* expressing *ksgA⁻*, *ksgA⁻ ΔyopJ*, *ksgA⁻ yopJ^{C172A}*, or *ksgA⁻*

yopJ^{F177L}. (A and B) Spleens were removed at day 3 post infection snap frozen in OCT

for TUNEL staining. Row A is showing TUNEL only images at 200x and row B is

showing representative merged images with DAPI (blue), TUNEL (FITC) and *Y.*

pseudotuberculosis (Rhodamine Red X). (C and D) Random images were captured (5

from each spleen, 3 mice per group) and TUNEL+ area (total pixels) per field was

quantified using ImageJ software. C and D are results obtained from two independent

experiments.

Figure 6. Suramin inhibits YopJ-induced macrophage cytotoxicity and enhances

virulence *in vivo*. A. Bone marrow derived macrophages from C57BL/6 mice were

exposed to *ksgA⁻ Y. pseudotuberculosis* at an MOI of 100:1 in the presence or absence

of suramin at different concentrations. LDH release was measured to determine

macrophage cytotoxicity. **B.** C57BL/6 mice (n=3-6/group; repeated twice) were infected

intravenously (1×10^3 CFU) with various strains of *Y. pseudotuberculosis* in the presence

or absence of suramin delivered intraperitoneally shortly after infection. Mice were

weighed daily and monitored for signs of morbidity. **C** and **D**. C57BL/6 mice (*n*=3 per

group; repeated twice) were infected intravenously with *Y. pseudotuberculosis ksgA-*

(1x10^3 CFU) and treated intraperitoneally with either 5 mg of suramin or with PBS. After

three days, mice were sacrificed and livers were harvested and stained with H&E, large

white arrows indicate representative lesions found in the sections. **E**. Quantification of

the number of microabscesses from the entirety of the H&E-stained liver sections (*n*=3

per group). **F** and **G**, TUNEL-stained liver sections are shown, with TUNEL+ lesions

highlighted with white solid arrows. **H**. Quantification of the degree of TUNEL positivity

per microabscess (*n*=4-6 microabscesses per liver section; *n*=3 mice per group).

Fig. 7. Proposed model to explain how different YopJ proteins lead to different *in

***vivo* outcomes following *Y. pseudotuberculosis* infection.** (**A**). *Y.*

pseudotuberculosis expressing functional YopJ (harboring either the wild type/positively

selected F177 or the ancestral L177) attach to the host cell and inject YopJ, leading to

host cell death. Efferocytosis of dead host cells by neighboring activated phagocyte also

removes those bacteria attached to dead host cells, leading to pathogen clearance

(**B**). *Y. pseudotuberculosis* expressing no or catalytically dead YopJ fail to induce host

cell death, meaning the host cell would not be removed and bacteria could persist

longer to cause illness. For either scenario, other factors including host innate and

adaptive immune responses (not shown), could also influence the balance between

pathogen clearance and pathogen persistence.

SUPPLEMENTAL FIGURE LEGENDS

S1. Repair of *yopJ in the ksgA⁻ yopJ⁻* strain restores the original phenotype. (A)

RAW macrophages were infected with either *ksgA⁻*, *ksgA⁻ ΔyopJ*, or *ksgA⁻ yopJ^{repaired}* *Y.*

pseudotuberculosis strains at an MOI of 100:1. LDH release was measured to

determine the cytotoxicity of these strains in macrophages. (B and C) C57BL/6 mice

were intravenously challenged with *ksgA⁻*, *ksgA⁻ yopJ⁻* or *ksgA⁻ yopJ^{repaired}* *Y.*

pseudotuberculosis. Morbidity and mortality was followed after infection and data shown

represents the percent survival and weight loss of each group (n=5 mice per group).

S2. A cytokine storm does not characterize illness caused by *ksgA- ΔyopJ*

bacteria. C57BL/6 mice were inoculated intravenously (1×10^3 CFU) with *ksgA⁻*, *ksgA⁻*

ΔyopJ, or *ksgA⁻ yopJ^{C172A}* *ksgA⁻ yopJ^{F177L}* bacteria. Serum was collected from each

mouse at day 9 post inoculation and was assayed for cytokines using a cytokine

multiplex assay.

S3. Systemic clodronate liposome induced macrophage apoptosis in vivo does

not protect mice from *Y. pseudotuberculosis* infection. 6-8 week old female

C57BL/6 mice were infected intravenously (1×10^3 CFU) with ksgA-yopJ- *Y.*

pseudotuberculosis and left untreated or treated with either PBS containing liposomes

or clodronate containing liposomes at 6 hours post infection. Morbidity and mortality

was followed after infection and data shown represents the percent survival of each

group (n=6 mice per group).

Table 1. *Y. pseudotuberculosis* (YPIII pIB1) strains used in this study

Strain	Strain name	Reference
ksgA⁻	500	[80]
ksgA⁻yopJ⁻	MB153	This study
ksgA⁻yopJ^{C172A}	MB219	This study
ksgA⁻yopJ^{F177L}	MB254	This study

Supplementary Table 1. Positively selected sites in *Y. pseudotuberculosis* YopJ

position	Residue	Prob ($\omega > 1$)	Mean ω	Standard deviation
10	I	0.544	5.537	4.754
11	S	0.998**	9.843	1.001
20	S	0.938	9.291	2.394
33	T	0.790	7.904	3.852
40	S	0.581	5.891	4.715
52	M	0.999**	9.851	0.965
54	V	0.791	7.920	3.841
80	L	0.989	9.765	1.311
130	A	0.835	8.324	3.532
139	M	0.740	7.435	4.128
143	R	0.909	9.020	2.795
177	F	0.971	9.592	1.798
205	G	0.838	8.356	3.504
206	E	1.000**	9.861	0.916
212	D	0.806	8.059	3.742

Supplementary Table S2. Primers used in this study

Primer	Sequence	Purpose
F yopJ	5'CAACAAGTTTCTCTACCGGAGAAT3'	screen for ΔyopJ allele
R yopJ	5'CTCATACCACCCGTACTCTAGCA3'	
yopJ F3	5'GATCGATATCCAAGTGCCCCCTAAGCCTTGAGTT3'	PCR clone yopJ orf into pACYC184, screen for yopJ point-mutant alleles
yopJ R2	5'GATCGTCGACCCCATACTGGAGCAAGATTTCC3'	
yopJ* F (sense)	5'GAAATGGATATTCAGCGAAGCTCATCTGAAGCTGGTATTTTTAGTTTTGCAC3'	mutate $yopJ^{WT}$ to $yopJ^{C172A}$
yopJ* R (antisense)	5'GTGCAAAACTTAAAATACCAGCTTCAGATGAGCTTAGCTGAATATCCATTTC3'	
yopJ F4	5'GATCGCATGCCAAGTCCCCCCTAAGCCTTGAGTT3'	PCR clone $yopJ^{C172A}$ into pCVD442
yopJ R4	5'GATCGCATGCCCCATACTGGAGCAAGATTTCC3'	
F177A F (sense)	5'GAAGCTCATCTGAATGTGGTATTTTTAGTTTGGCACTGGCAAAAAAAC3'	mutate $yopJ^{C172A}$ to $yopJ^{F177L}$
F177A R (antisense)	5'GTTTTTTTGCCAGTGCCAAACTAAAAATACCACATTCAGATGAGCTTC3'	

[1]underlined sequence indicates restriction enzyme site used for cloning

References

1.   Fink SL, Cookson BT. Apoptosis, pyroptosis, and necrosis: mechanistic description of dead
and dying eukaryotic cells. Infect Immun. 2005;73(4):1907-16. Epub 2005/03/24. doi:
10.1128/IAI.73.4.1907-1916.2005. PubMed PMID: 15784530; PubMed Central PMCID:
PMCPMC1087413.
2.   Bergsbaken T, Fink SL, Cookson BT. Pyroptosis: host cell death and inflammation. Nat Rev
Microbiol. 2009;7(2):99-109. Epub 2009/01/17. doi: 10.1038/nrmicro2070. PubMed PMID:
19148178; PubMed Central PMCID: PMCPMC2910423.
3.   Lara-Tejero M, Sutterwala FS, Ogura Y, Grant EP, Bertin J, Coyle AJ, et al. Role of the caspase-
1 inflammasome in Salmonella typhimurium pathogenesis. J Exp Med. 2006;203(6):1407-12. Epub
2006/05/24. doi: 10.1084/jem.20060206. PubMed PMID: 16717117; PubMed Central PMCID:
PMCPMC2118315.
4.   Lima-Junior DS, Costa DL, Carregaro V, Cunha LD, Silva AL, Mineo TW, et al. Inflammasome-
derived IL-1beta production induces nitric oxide-mediated resistance to Leishmania. Nat Med.
2013;19(7):909-15. Epub 2013/06/12. doi: 10.1038/nm.3221. PubMed PMID: 23749230.
5.   Chain PS, Hu P, Malfatti SA, Radnedge L, Larimer F, Vergez LM, et al. Complete genome
sequence of Yersinia pestis strains Antiqua and Nepal516: evidence of gene reduction in an
emerging pathogen. J Bacteriol. 2006;188(12):4453-63. Epub 2006/06/03. doi: 10.1128/JB.00124-
06. PubMed PMID: 16740952; PubMed Central PMCID: PMCPMC1482938.
6.   Wren BW. The yersiniae--a model genus to study the rapid evolution of bacterial pathogens.
Nat Rev Microbiol. 2003;1(1):55-64. Epub 2004/03/26. doi: 10.1038/nrmicro730. PubMed PMID:
15040180.
7.   Brodsky IE, Palm NW, Sadanand S, Ryndak MB, Sutterwala FS, Flavell RA, et al. A Yersinia
effector protein promotes virulence by preventing inflammasome recognition of the type III
secretion system. Cell Host Microbe. 2010;7(5):376-87. Epub 2010/05/19. doi:
10.1016/j.chom.2010.04.009. PubMed PMID: 20478539; PubMed Central PMCID:
PMCPMC2883865.
8.   Garcia JT, Ferracci F, Jackson MW, Joseph SS, Pattis I, Plano LR, et al. Measurement of
effector protein injection by type III and type IV secretion systems by using a 13-residue
phosphorylatable glycogen synthase kinase tag. Infect Immun. 2006;74(10):5645-57. Epub
2006/09/22. doi: 10.1128/IAI.00690-06. PubMed PMID: 16988240; PubMed Central PMCID:
PMCPMC1594927.
9.   Trosky JE, Liverman AD, Orth K. Yersinia outer proteins: Yops. Cell Microbiol.
2008;10(3):557-65. Epub 2007/12/18. doi: 10.1111/j.1462-5822.2007.01109.x. PubMed PMID:
18081726.
10.   Orth K, Xu Z, Mudgett MB, Bao ZQ, Palmer LE, Bliska JB, et al. Disruption of signaling by
Yersinia effector YopJ, a ubiquitin-like protein protease. Science. 2000;290(5496):1594-7. Epub
2000/11/25. doi: 10.1126/science.290.5496.1594. PubMed PMID: 11090361.
11.   Sweet CR, Conlon J, Golenbock DT, Goguen J, Silverman N. YopJ targets TRAF proteins to
inhibit TLR-mediated NF-kappaB, MAPK and IRF3 signal transduction. Cell Microbiol.
2007;9(11):2700-15. Epub 2007/07/05. doi: 10.1111/j.1462-5822.2007.00990.x. PubMed PMID:
17608743.
12.   Zhou H, Monack DM, Kayagaki N, Wertz I, Yin J, Wolf B, et al. Yersinia virulence factor YopJ
acts as a deubiquitinase to inhibit NF-kappa B activation. J Exp Med. 2005;202(10):1327-32. Epub
2005/11/23. doi: 10.1084/jem.20051194. PubMed PMID: 16301742; PubMed Central PMCID:
PMCPMC2212976.
13.   Mittal R, Peak-Chew SY, McMahon HT. Acetylation of MEK2 and I kappa B kinase (IKK)
activation loop residues by YopJ inhibits signaling. Proc Natl Acad Sci U S A. 2006;103(49):18574-9.

Epub 2006/11/23. doi: 10.1073/pnas.0608995103. PubMed PMID: 17116858; PubMed Central
PMCID: PMCPMC1654131.
14.   Mukherjee S, Keitany G, Li Y, Wang Y, Ball HL, Goldsmith EJ, et al. Yersinia YopJ acetylates
and inhibits kinase activation by blocking phosphorylation. Science. 2006;312(5777):1211-4. Epub
2006/05/27. doi: 10.1126/science.1126867. PubMed PMID: 16728640.
15.   Mittal R, Peak-Chew SY, Sade RS, Vallis Y, McMahon HT. The acetyltransferase activity of the
bacterial toxin YopJ of Yersinia is activated by eukaryotic host cell inositol hexakisphosphate. J Biol
Chem. 2010;285(26):19927-34. Epub 2010/05/01. doi: 10.1074/jbc.M110.126581. PubMed PMID:
20430892; PubMed Central PMCID: PMCPMC2888404.
16.   Philip NH, Brodsky IE. Cell death programs in Yersinia immunity and pathogenesis. Front
Cell Infect Microbiol. 2012;2:149. Epub 2012/12/12. doi: 10.3389/fcimb.2012.00149. PubMed
PMID: 23226685; PubMed Central PMCID: PMCPMC3510641.
17.   Zheng Y, Lilo S, Mena P, Bliska JB. YopJ-induced caspase-1 activation in Yersinia-infected
macrophages: independent of apoptosis, linked to necrosis, dispensable for innate host defense.
PLoS One. 2012;7(4):e36019. Epub 2012/05/09. doi: 10.1371/journal.pone.0036019. PubMed
PMID: 22563435; PubMed Central PMCID: PMCPMC3338577.
18.   Galyov EE, Hakansson S, Wolf-Watz H. Characterization of the operon encoding the YpkA
Ser/Thr protein kinase and the YopJ protein of Yersinia pseudotuberculosis. J Bacteriol.
1994;176(15):4543-8. Epub 1994/08/01. doi: 10.1128/jb.176.15.4543-4548.1994. PubMed PMID:
8045884; PubMed Central PMCID: PMCPMC196273.
19.   Lemaitre N, Sebbane F, Long D, Hinnebusch BJ. Yersinia pestis YopJ suppresses tumor
necrosis factor alpha induction and contributes to apoptosis of immune cells in the lymph node but
is not required for virulence in a rat model of bubonic plague. Infect Immun. 2006;74(9):5126-31.
Epub 2006/08/24. doi: 10.1128/IAI.00219-06. PubMed PMID: 16926404; PubMed Central PMCID:
PMCPMC1594864.
20.   Monack DM, Mecsas J, Bouley D, Falkow S. Yersinia-induced apoptosis in vivo aids in the
establishment of a systemic infection of mice. J Exp Med. 1998;188(11):2127-37. Epub
1998/12/08. doi: 10.1084/jem.188.11.2127. PubMed PMID: 9841926; PubMed Central PMCID:
PMCPMC2212385.
21.   Straley SC, Bowmer WS. Virulence genes regulated at the transcriptional level by Ca2+ in
Yersinia pestis include structural genes for outer membrane proteins. Infect Immun.
1986;51(2):445-54. Epub 1986/02/01. doi: 10.1128/IAI.51.2.445-454.1986. PubMed PMID:
3002984; PubMed Central PMCID: PMCPMC262351.
22.   Straley SC, Cibull ML. Differential clearance and host-pathogen interactions of YopE- and
YopK- YopL- Yersinia pestis in BALB/c mice. Infect Immun. 1989;57(4):1200-10. Epub
1989/04/01. doi: 10.1128/IAI.57.4.1200-1210.1989. PubMed PMID: 2925246; PubMed Central
PMCID: PMCPMC313251.
23.   Trulzsch K, Sporleder T, Igwe EI, Russmann H, Heesemann J. Contribution of the major
secreted yops of Yersinia enterocolitica O:8 to pathogenicity in the mouse infection model. Infect
Immun. 2004;72(9):5227-34. Epub 2004/08/24. doi: 10.1128/IAI.72.9.5227-5234.2004. PubMed
PMID: 15322017; PubMed Central PMCID: PMCPMC517446.
24.   Brodsky IE, Medzhitov R. Reduced secretion of YopJ by Yersinia limits in vivo cell death but
enhances bacterial virulence. PLoS Pathog. 2008;4(5):e1000067. Epub 2008/05/17. doi:
10.1371/journal.ppat.1000067. PubMed PMID: 18483548; PubMed Central PMCID:
PMCPMC2361194.
25.   Zauberman A, Tidhar A, Levy Y, Bar-Haim E, Halperin G, Flashner Y, et al. Yersinia pestis
endowed with increased cytotoxicity is avirulent in a bubonic plague model and induces rapid
protection against pneumonic plague. PLoS One. 2009;4(6):e5938. Epub 2009/06/17. doi:
10.1371/journal.pone.0005938. PubMed PMID: 19529770; PubMed Central PMCID:
PMCPMC2691952.

26. Zheng Y, Lilo S, Brodsky IE, Zhang Y, Medzhitov R, Marcu KB, et al. A Yersinia effector with
enhanced inhibitory activity on the NF-kappaB pathway activates the NLRP3/ASC/caspase-1
inflammasome in macrophages. PLoS Pathog. 2011;7(4):e1002026. Epub 2011/05/03. doi:
10.1371/journal.ppat.1002026. PubMed PMID: 21533069; PubMed Central PMCID:
PMCPMC3080847.
27. Stavrinides J, McCann HC, Guttman DS. Host-pathogen interplay and the evolution of
bacterial effectors. Cell Microbiol. 2008;10(2):285-92. Epub 2007/11/24. doi: 10.1111/j.1462-
5822.2007.01078.x. PubMed PMID: 18034865.
28. Terauchi R, Yoshida K. Towards population genomics of effector-effector target interactions.
New Phytol. 2010;187(4):929-39. Epub 2010/08/17. doi: 10.1111/j.1469-8137.2010.03408.x.
PubMed PMID: 20707855.
29. Yang Z, Rannala B. Molecular phylogenetics: principles and practice. Nat Rev Genet.
2012;13(5):303-14. Epub 2012/03/30. doi: 10.1038/nrg3186. PubMed PMID: 22456349.
30. Borges V, Nunes A, Ferreira R, Borrego MJ, Gomes JP. Directional evolution of Chlamydia
trachomatis towards niche-specific adaptation. J Bacteriol. 2012;194(22):6143-53. Epub
2012/09/11. doi: 10.1128/JB.01291-12. PubMed PMID: 22961851; PubMed Central PMCID:
PMCPMC3486361.
31. Ma W, Dong FF, Stavrinides J, Guttman DS. Type III effector diversification via both
pathoadaptation and horizontal transfer in response to a coevolutionary arms race. PLoS Genet.
2006;2(12):e209. Epub 2006/12/30. doi: 10.1371/journal.pgen.0020209. PubMed PMID:
17194219; PubMed Central PMCID: PMCPMC1713259.
32. Chain PS, Carniel E, Larimer FW, Lamerdin J, Stoutland PO, Regala WM, et al. Insights into
the evolution of Yersinia pestis through whole-genome comparison with Yersinia
pseudotuberculosis. Proc Natl Acad Sci U S A. 2004;101(38):13826-31. Epub 2004/09/11. doi:
10.1073/pnas.0404012101. PubMed PMID: 15358858; PubMed Central PMCID: PMCPMC518763.
33. Sun J. Pathogenic Bacterial Proteins and their Anti-Inflammatory Effects in the Eukaryotic
Host. Antiinflamm Antiallergy Agents Med Chem. 2009;8(3):214-27. Epub 2010/01/22. doi:
10.2174/187152309789151986. PubMed PMID: 20090866; PubMed Central PMCID:
PMCPMC2808140.
34. Nielsen R, Yang Z. Likelihood models for detecting positively selected amino acid sites and
applications to the HIV-1 envelope gene. Genetics. 1998;148(3):929-36. Epub 1998/04/16. PubMed
PMID: 9539414; PubMed Central PMCID: PMCPMC1460041.
35. Yang Z, Swanson WJ, Vacquier VD. Maximum-likelihood analysis of molecular adaptation in
abalone sperm lysin reveals variable selective pressures among lineages and sites. Mol Biol Evol.
2000;17(10):1446-55. Epub 2000/10/06. doi: 10.1093/oxfordjournals.molbev.a026245. PubMed
PMID: 11018152.
36. Barnes PD, Bergman MA, Mecsas J, Isberg RR. Yersinia pseudotuberculosis disseminates
directly from a replicating bacterial pool in the intestine. J Exp Med. 2006;203(6):1591-601. Epub
2006/06/07. doi: 10.1084/jem.20060905. PubMed PMID: 16754724; PubMed Central PMCID:
PMCPMC2118325.
37. Zhang Y, Ting AT, Marcu KB, Bliska JB. Inhibition of MAPK and NF-kappa B pathways is
necessary for rapid apoptosis in macrophages infected with Yersinia. J Immunol.
2005;174(12):7939-49. Epub 2005/06/10. doi: 10.4049/jimmunol.174.12.7939. PubMed PMID:
15944300.
38. Boland A, Cornelis GR. Role of YopP in suppression of tumor necrosis factor alpha release by
macrophages during Yersinia infection. Infect Immun. 1998;66(5):1878-84. Epub 1998/05/09. doi:
10.1128/IAI.66.5.1878-1884.1998. PubMed PMID: 9573064; PubMed Central PMCID:
PMCPMC108138.
39. Palmer LE, Hobbie S, Galan JE, Bliska JB. YopJ of Yersinia pseudotuberculosis is required for
the inhibition of macrophage TNF-alpha production and downregulation of the MAP kinases p38

and JNK. Mol Microbiol. 1998;27(5):953-65. Epub 1998/04/16. doi: 10.1046/j.1365-
2958.1998.00740.x. PubMed PMID: 9535085.
40.   Schesser K, Spiik AK, Dukuzumuremyi JM, Neurath MF, Pettersson S, Wolf-Watz H. The yopJ
locus is required for Yersinia-mediated inhibition of NF-kappaB activation and cytokine expression:
YopJ contains a eukaryotic SH2-like domain that is essential for its repressive activity. Mol
Microbiol. 1998;28(6):1067-79. Epub 1998/07/29. doi: 10.1046/j.1365-2958.1998.00851.x.
PubMed PMID: 9680199.
41.   Ruckdeschel K, Harb S, Roggenkamp A, Hornef M, Zumbihl R, Kohler S, et al. Yersinia
enterocolitica impairs activation of transcription factor NF-kappaB: involvement in the induction of
programmed cell death and in the suppression of the macrophage tumor necrosis factor alpha
production. J Exp Med. 1998;187(7):1069-79. Epub 1998/05/16. doi: 10.1084/jem.187.7.1069.
PubMed PMID: 9529323; PubMed Central PMCID: PMCPMC2212204.
42.   Inoue S, Sato T, Suzuki-Utsunomiya K, Komori Y, Hozumi K, Chiba T, et al. Sepsis-induced
hypercytokinemia and lymphocyte apoptosis in aging-accelerated Klotho knockout mice. Shock.
2013;39(3):311-6. Epub 2013/02/01. doi: 10.1097/SHK.0b013e3182845445. PubMed PMID:
23364432.
43.   Teijaro JR, Walsh KB, Cahalan S, Fremgen DM, Roberts E, Scott F, et al. Endothelial cells are
central orchestrators of cytokine amplification during influenza virus infection. Cell.
2011;146(6):980-91. Epub 2011/09/20. doi: 10.1016/j.cell.2011.08.015. PubMed PMID:
21925319; PubMed Central PMCID: PMCPMC3176439.
44.   Ulloa L, Tracey KJ. The "cytokine profile": a code for sepsis. Trends Mol Med. 2005;11(2):56-
63. Epub 2005/02/08. doi: 10.1016/j.molmed.2004.12.007. PubMed PMID: 15694867.
45.   van Rooijen N, van Kesteren-Hendrikx E. "In vivo" depletion of macrophages by liposome-
mediated "suicide". Methods Enzymol. 2003;373:3-16. Epub 2004/01/13. doi: 10.1016/s0076-
6879(03)73001-8. PubMed PMID: 14714393.
46.   Eichhorst ST, Krueger A, Muerkoster S, Fas SC, Golks A, Gruetzner U, et al. Suramin inhibits
death receptor-induced apoptosis in vitro and fulminant apoptotic liver damage in mice. Nat Med.
2004;10(6):602-9. Epub 2004/05/18. doi: 10.1038/nm1049. PubMed PMID: 15146177.
47.   Goto T, Takeuchi S, Miura K, Ohshima S, Mikami K, Yoneyama K, et al. Suramin prevents
fulminant hepatic failure resulting in reduction of lethality through the suppression of NF-kappaB
activity. Cytokine. 2006;33(1):28-35. Epub 2006/01/18. doi: 10.1016/j.cyto.2005.11.012. PubMed
PMID: 16413198.
48.   Ruckdeschel K, Richter K, Mannel O, Heesemann J. Arginine-143 of Yersinia enterocolitica
YopP crucially determines isotype-related NF-kappaB suppression and apoptosis induction in
macrophages. Infect Immun. 2001;69(12):7652-62. Epub 2001/11/14. doi:
10.1128/IAI.69.12.7652-7662.2001. PubMed PMID: 11705945; PubMed Central PMCID:
PMCPMC98859.
49.   Zauberman A, Cohen S, Mamroud E, Flashner Y, Tidhar A, Ber R, et al. Interaction of Yersinia
pestis with macrophages: limitations in YopJ-dependent apoptosis. Infect Immun.
2006;74(6):3239-50. Epub 2006/05/23. doi: 10.1128/IAI.00097-06. PubMed PMID: 16714551;
PubMed Central PMCID: PMCPMC1479247.
50.   Bolin I, Norlander L, Wolf-Watz H. Temperature-inducible outer membrane protein of
Yersinia pseudotuberculosis and Yersinia enterocolitica is associated with the virulence plasmid.
Infect Immun. 1982;37(2):506-12. Epub 1982/08/01. doi: 10.1128/IAI.37.2.506-512.1982.
PubMed PMID: 6749681; PubMed Central PMCID: PMCPMC347563.
51.   Portnoy DA, Wolf-Watz H, Bolin I, Beeder AB, Falkow S. Characterization of common
virulence plasmids in Yersinia species and their role in the expression of outer membrane proteins.
Infect Immun. 1984;43(1):108-14. Epub 1984/01/01. doi: 10.1128/IAI.43.1.108-114.1984.
PubMed PMID: 6317562; PubMed Central PMCID: PMCPMC263395.

52.   Easterday WR, Kausrud KL, Star B, Heier L, Haley BJ, Ageyev V, et al. An additional step in
the transmission of Yersinia pestis? ISME J. 2012;6(2):231-6. Epub 2011/08/13. doi:
10.1038/ismej.2011.105. PubMed PMID: 21833036; PubMed Central PMCID: PMCPMC3260498.
53.   Orning P, Weng D, Starheim K, Ratner D, Best Z, Lee B, et al. Pathogen blockade of TAK1
triggers caspase-8-dependent cleavage of gasdermin D and cell death. Science.
2018;362(6418):1064-9. Epub 2018/10/27. doi: 10.1126/science.aau2818. PubMed PMID:
30361383; PubMed Central PMCID: PMCPMC6522129.
54.   Paquette N, Conlon J, Sweet C, Rus F, Wilson L, Pereira A, et al. Serine/threonine acetylation
of TGFbeta-activated kinase (TAK1) by Yersinia pestis YopJ inhibits innate immune signaling. Proc
Natl Acad Sci U S A. 2012;109(31):12710-5. Epub 2012/07/18. doi: 10.1073/pnas.1008203109.
PubMed PMID: 22802624; PubMed Central PMCID: PMCPMC3411953.
55.   Philip NH, Dillon CP, Snyder AG, Fitzgerald P, Wynosky-Dolfi MA, Zwack EE, et al. Caspase-8
mediates caspase-1 processing and innate immune defense in response to bacterial blockade of NF-
kappaB and MAPK signaling. Proc Natl Acad Sci U S A. 2014;111(20):7385-90. Epub 2014/05/07.
doi: 10.1073/pnas.1403252111. PubMed PMID: 24799700; PubMed Central PMCID:
PMCPMC4034241.
56.   Sarhan J, Liu BC, Muendlein HI, Li P, Nilson R, Tang AY, et al. Caspase-8 induces cleavage of
gasdermin D to elicit pyroptosis during Yersinia infection. Proc Natl Acad Sci U S A.
2018;115(46):E10888-E97. Epub 2018/11/02. doi: 10.1073/pnas.1809548115. PubMed PMID:
30381458; PubMed Central PMCID: PMCPMC6243247.
57.   Peterson LW, Philip NH, Dillon CP, Bertin J, Gough PJ, Green DR, et al. Cell-Extrinsic TNF
Collaborates with TRIF Signaling To Promote Yersinia-Induced Apoptosis. J Immunol.
2016;197(10):4110-7. Epub 2016/10/14. doi: 10.4049/jimmunol.1601294. PubMed PMID:
27733552; PubMed Central PMCID: PMCPMC5123756.
58.   Weng D, Marty-Roix R, Ganesan S, Proulx MK, Vladimer GI, Kaiser WJ, et al. Caspase-8 and
RIP kinases regulate bacteria-induced innate immune responses and cell death. Proc Natl Acad Sci U
S A. 2014;111(20):7391-6. Epub 2014/05/07. doi: 10.1073/pnas.1403477111. PubMed PMID:
24799678; PubMed Central PMCID: PMCPMC4034196.
59.   Bergman MA, Loomis WP, Mecsas J, Starnbach MN, Isberg RR. CD8(+) T cells restrict
Yersinia pseudotuberculosis infection: bypass of anti-phagocytosis by targeting antigen-presenting
cells. PLoS Pathog. 2009;5(9):e1000573. Epub 2009/09/05. doi: 10.1371/journal.ppat.1000573.
PubMed PMID: 19730693; PubMed Central PMCID: PMCPMC2731216.
60.   Fehr D, Casanova C, Liverman A, Blazkova H, Orth K, Dobbelaere D, et al. AopP, a type III
effector protein of Aeromonas salmonicida, inhibits the NF-kappaB signalling pathway.
Microbiology. 2006;152(Pt 9):2809-18. Epub 2006/09/02. doi: 10.1099/mic.0.28889-0. PubMed
PMID: 16946275.
61.   Arifuzzaman M, Ang WXG, Choi HW, Nilles ML, St John AL, Abraham SN. Necroptosis of
infiltrated macrophages drives Yersinia pestis dispersal within buboes. JCI Insight. 2018;3(18).
Epub 2018/09/21. doi: 10.1172/jci.insight.122188. PubMed PMID: 30232285; PubMed Central
PMCID: PMCPMC6237236.
62.   Dewoody R, Merritt PM, Houppert AS, Marketon MM. YopK regulates the Yersinia pestis
type III secretion system from within host cells. Mol Microbiol. 2011;79(6):1445-61. Epub
2011/01/06. doi: 10.1111/j.1365-2958.2011.07534.x. PubMed PMID: 21205017; PubMed Central
PMCID: PMCPMC3210821.
63.   Dewoody R, Merritt PM, Marketon MM. YopK controls both rate and fidelity of Yop
translocation. Mol Microbiol. 2013;87(2):301-17. Epub 2012/12/05. doi: 10.1111/mmi.12099.
PubMed PMID: 23205707; PubMed Central PMCID: PMCPMC3545096.
64.   Vandenabeele P, Vanden Berghe T, Festjens N. Caspase inhibitors promote alternative cell
death pathways. Sci STKE. 2006;2006(358):pe44. Epub 2006/10/26. doi:
10.1126/stke.3582006pe44. PubMed PMID: 17062895.

65.   Lewis JD, Lee A, Ma W, Zhou H, Guttman DS, Desveaux D. The YopJ superfamily in plant-
associated bacteria. Mol Plant Pathol. 2011;12(9):928-37. Epub 2011/07/06. doi: 10.1111/j.1364-
3703.2011.00719.x. PubMed PMID: 21726386; PubMed Central PMCID: PMCPMC6640427.
66.   Trosky JE, Mukherjee S, Burdette DL, Roberts M, McCarter L, Siegel RM, et al. Inhibition of
MAPK signaling pathways by VopA from Vibrio parahaemolyticus. J Biol Chem.
2004;279(50):51953-7. Epub 2004/10/02. doi: 10.1074/jbc.M407001200. PubMed PMID:
15459200.
67.   Alsmark CM, Frank AC, Karlberg EO, Legault BA, Ardell DH, Canback B, et al. The louse-
borne human pathogen Bartonella quintana is a genomic derivative of the zoonotic agent Bartonella
henselae. Proc Natl Acad Sci U S A. 2004;101(26):9716-21. Epub 2004/06/24. doi:
10.1073/pnas.0305659101. PubMed PMID: 15210978; PubMed Central PMCID: PMCPMC470741.
68.   Hardt WD, Galan JE. A secreted Salmonella protein with homology to an avirulence
determinant of plant pathogenic bacteria. Proc Natl Acad Sci U S A. 1997;94(18):9887-92. Epub
1997/09/02. doi: 10.1073/pnas.94.18.9887. PubMed PMID: 9275221; PubMed Central PMCID:
PMCPMC23287.
69.   Collier-Hyams LS, Zeng H, Sun J, Tomlinson AD, Bao ZQ, Chen H, et al. Cutting edge:
Salmonella AvrA effector inhibits the key proinflammatory, anti-apoptotic NF-kappa B pathway. J
Immunol. 2002;169(6):2846-50. Epub 2002/09/10. doi: 10.4049/jimmunol.169.6.2846. PubMed
PMID: 12218096.
70.   Jones RM, Wu H, Wentworth C, Luo L, Collier-Hyams L, Neish AS. Salmonella AvrA
Coordinates Suppression of Host Immune and Apoptotic Defenses via JNK Pathway Blockade. Cell
Host Microbe. 2008;3(4):233-44. Epub 2008/04/15. doi: 10.1016/j.chom.2008.02.016. PubMed
PMID: 18407067.
71.   Liu X, Lu R, Xia Y, Wu S, Sun J. Eukaryotic signaling pathways targeted by Salmonella effector
protein AvrA in intestinal infection in vivo. BMC Microbiol. 2010;10:326. Epub 2010/12/25. doi:
10.1186/1471-2180-10-326. PubMed PMID: 21182782; PubMed Central PMCID:
PMCPMC3027599.
72.   Schesser K, Dukuzumuremyi JM, Cilio C, Borg S, Wallis TS, Pettersson S, et al. The Salmonella
YopJ-homologue AvrA does not possess YopJ-like activity. Microb Pathog. 2000;28(2):59-70. Epub
2000/01/25. doi: 10.1006/mpat.1999.0324. PubMed PMID: 10644492.
73.   Du F, Galan JE. Selective inhibition of type III secretion activated signaling by the Salmonella
effector AvrA. PLoS Pathog. 2009;5(9):e1000595. Epub 2009/09/26. doi:
10.1371/journal.ppat.1000595. PubMed PMID: 19779561; PubMed Central PMCID:
PMCPMC2742890.
74.   Coll NS, Epple P, Dangl JL. Programmed cell death in the plant immune system. Cell Death
Differ. 2011;18(8):1247-56. Epub 2011/04/09. doi: 10.1038/cdd.2011.37. PubMed PMID:
21475301; PubMed Central PMCID: PMCPMC3172094.
75.   Roden J, Eardley L, Hotson A, Cao Y, Mudgett MB. Characterization of the Xanthomonas
AvrXv4 effector, a SUMO protease translocated into plant cells. Mol Plant Microbe Interact.
2004;17(6):633-43. Epub 2004/06/16. doi: 10.1094/MPMI.2004.17.6.633. PubMed PMID:
15195946.
76.   Lee AH, Hurley B, Felsensteiner C, Yea C, Ckurshumova W, Bartetzko V, et al. A bacterial
acetyltransferase destroys plant microtubule networks and blocks secretion. PLoS Pathog.
2012;8(2):e1002523. Epub 2012/02/10. doi: 10.1371/journal.ppat.1002523. PubMed PMID:
22319451; PubMed Central PMCID: PMCPMC3271077.
77.   Sanabria NM, Huang JC, Dubery IA. Self/nonself perception in plants in innate immunity and
defense. Self Nonself. 2010;1(1):40-54. Epub 2010/01/01. doi: 10.4161/self.1.1.10442. PubMed
PMID: 21559176; PubMed Central PMCID: PMCPMC3091600.

78.   Prehna G, Ivanov MI, Bliska JB, Stebbins CE. Yersinia virulence depends on mimicry of host
Rho-family nucleotide dissociation inhibitors. Cell. 2006;126(5):869-80. Epub 2006/09/09. doi:
10.1016/j.cell.2006.06.056. PubMed PMID: 16959567.
79.   Gu J, Neary JL, Sanchez M, Yu J, Lilburn TG, Wang Y. Genome evolution and functional
divergence in Yersinia. J Exp Zool B Mol Dev Evol. 2007;308(1):37-49. Epub 2006/07/14. doi:
10.1002/jez.b.21120. PubMed PMID: 16838303.
80.   Cui Y, Yu C, Yan Y, Li D, Li Y, Jombart T, et al. Historical variations in mutation rate in an
epidemic pathogen, Yersinia pestis. Proc Natl Acad Sci U S A. 2013;110(2):577-82. Epub
2012/12/29. doi: 10.1073/pnas.1205750110. PubMed PMID: 23271803; PubMed Central PMCID:
PMCPMC3545753.
81.   Stenkova AM, Isaeva MP, Shubin FN, Rasskazov VA, Rakin AV. Trends of the major porin
gene (ompF) evolution: insight from the genus Yersinia. PLoS One. 2011;6(5):e20546. Epub
2011/06/10. doi: 10.1371/journal.pone.0020546. PubMed PMID: 21655186; PubMed Central
PMCID: PMCPMC3105102.
82.   Elde NC, Child SJ, Geballe AP, Malik HS. Protein kinase R reveals an evolutionary model for
defeating viral mimicry. Nature. 2009;457(7228):485-9. Epub 2008/12/02. doi:
10.1038/nature07529. PubMed PMID: 19043403; PubMed Central PMCID: PMCPMC2629804.
83.   Wiley DJ, Shrestha N, Yang J, Atis N, Dayton K, Schesser K. The activities of the Yersinia
protein kinase A (YpkA) and outer protein J (YopJ) virulence factors converge on an eIF2alpha
kinase. J Biol Chem. 2009;284(37):24744-53. Epub 2009/06/26. doi: 10.1074/jbc.M109.010140.
PubMed PMID: 19553678; PubMed Central PMCID: PMCPMC2757178.
84.   Bergman MA, Cummings LA, Alaniz RC, Mayeda L, Fellnerova I, Cookson BT. CD4+-T-cell
responses generated during murine Salmonella enterica serovar Typhimurium infection are
directed towards multiple epitopes within the natural antigen FliC. Infect Immun.
2005;73(11):7226-35. Epub 2005/10/22. doi: 10.1128/IAI.73.11.7226-7235.2005. PubMed PMID:
16239517; PubMed Central PMCID: PMCPMC1273846.
85.   Mecsas J, Bilis I, Falkow S. Identification of attenuated Yersinia pseudotuberculosis strains
and characterization of an orogastric infection in BALB/c mice on day 5 postinfection by signature-
tagged mutagenesis. Infect Immun. 2001;69(5):2779-87. Epub 2001/04/09. doi:
10.1128/IAI.67.5.2779-2787.2001. PubMed PMID: 11292689; PubMed Central PMCID:
PMCPMC98225.
86.   Fisher ML, Castillo C, Mecsas J. Intranasal inoculation of mice with Yersinia
pseudotuberculosis causes a lethal lung infection that is dependent on Yersinia outer proteins and
PhoP. Infect Immun. 2007;75(1):429-42. Epub 2006/11/01. doi: 10.1128/IAI.01287-06. PubMed
PMID: 17074849; PubMed Central PMCID: PMCPMC1828392.
87.   Van Rooijen N, Sanders A. Liposome mediated depletion of macrophages: mechanism of
action, preparation of liposomes and applications. J Immunol Methods. 1994;174(1-2):83-93. Epub
1994/09/14. doi: 10.1016/0022-1759(94)90012-4. PubMed PMID: 8083541.
88.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
Nat Methods. 2012;9(7):671-5. Epub 2012/08/30. doi: 10.1038/nmeth.2089. PubMed PMID:
22930834; PubMed Central PMCID: PMCPMC5554542.

1148

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

| | PBS | *ksgA⁻* | *ksgA⁻yopJ⁻* | *ksgA⁻yopJ^{C172A}* | *ksgA⁻yopJ^{F177L}* |

A

TUNEL

B

TUNEL
Y. pstb
DAPI

Figure 6.

Figure 7.

A. *Y.pstb* YopJ
Y.pstb YopJ^{F177L}

B. *Y.pstb* YopJ⁻
Y.pstb YopJ^{C172A}

Pathogen clearance

Persistence and death

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3