1

Influence of delayed density and ultraviolet radiation on caterpillar baculovirus infection and

mortality

[1,2,*]Adam Pepi, [2]Vincent Pan, [2]Danielle Rutkowski, [2]Vinay Mase, and [2]Richard Karban

[1]Graduate Group in Ecology, University of California, Davis, CA 95616

[2]Department of Entomology & Nematology, University of California, Davis, CA 95616

* Corresponding author. Email: adampepi@gmail.com

Keywords:

Arctiinae; baculovirus; Bodega Marine Lab; disease dynamics; latitudinal gradient; population cycles

Abstract

1. Infectious disease is an important potential driver of population cycles, but this must occur through delayed density-dependent infection and resulting fitness effects. Delayed density-dependent infection by baculoviruses can be caused by environmental persistence of viral occlusion bodies, which can be influenced by environmental factors. In particular, ultraviolet radiation is potentially important in reducing the environmental persistence of viruses by inactivating viral occlusion bodies.

2. Delayed density-dependent viral infection has rarely been observed empirically at the population level although theory predicts that it is necessary for these pathogens to drive population cycles. Similarly, field studies have not examined the potential effects of ultraviolet radiation on viral infection rates in natural animal populations. We tested if viral infection is delayed density-dependent with the potential to drive cyclic dynamics and if ultraviolet radiation influences viral infection levels.

3. We censused 18 Ranchman's tiger moth (*Arctia virginalis*) populations across nearly 9° of latitude over two years and quantified the effects of direct and delayed density and ultraviolet radiation on baculovirus infection rates, infection severity, and survival to adulthood. Caterpillars were collected from each population in the field and reared in the laboratory. Baculovirus has not previously been described infecting *Arctia virginalis*, and we used genetic methods to confirm the identity of the virus.

4. We found that infection rate, infection severity, and survival to adulthood exhibited delayed density-dependence. Ultraviolet radiation in the previous summer decreased infection severity, and increased survival probability of the virus. Structural equation modelling indicated that the effect of lagged density on moth survival was mediated through infection rate and infection se-

verity, and was 2.5 fold stronger than the effect of ultraviolet radiation on survival through in-

fection severity. We successfully amplified polh, lef-8, and lef-9 viral genes from caterpillar

samples, and BLAST search results confirmed that the virus was a nucleopolyhedrovirus.

5. Our findings provide clear evidence that delayed density dependence can arise through viral in-

fection rate and severity in insects, which supports the role of viral disease as a potential mecha-

nism, among others, that may drive insect population cycles. Furthermore, our findings support

predictions that ultraviolet radiation can modify viral disease dynamics in insect populations,

most likely through attenuating viral persistence in the environment.

Introduction

Infectious disease plays an important role in population dynamics, potentially regulating

populations and driving cycles. Many insect populations undergo dramatic cyclic fluctuations, and

cyclic, delayed density-dependent disease outbreaks have been proposed as an explanation (Anderson

& May, 1980, 1981). However, many other mechanisms have also been proposed to explain insect

population cycles (Myers & Cory, 2013), particularly specialist parasitoids (Berryman, 1996). For

some species of Lepidoptera, empirical evidence has suggested that baculoviruses are most likely the

cause of cyclicity (Myers, 2000; Myers & Cory, 2013, 2016), though this has not been formally tested.

Delayed density-dependent feedbacks are required to drive cyclic population dynamics

(Turchin, 2003). Detection of delayed density-dependence involving many ecological factors has been

common (Turchin, 1990), and there have been several observations that viral epizootics follow high

densities of some insect species (Cory & Myers, 2003; Fuxa, 2004; Myers, 2000; Myers & Cory,

2016). However, direct demonstrations of delayed density-dependent viral infection rate and infection-

induced mortality are less common (Burthe et al., 2006; Fleming et al., 1986; Rothman, 1997).

Observational studies over relatively small areas (>15km) using space-for-time substitutions of local

densities have shown delayed density-dependent incidence of viral infection in voles (Burthe et al.,

2006) and soil-dwelling hepialid caterpillars (Fleming et al., 1986). Experimental manipulation of

western tent caterpillar (*Malacosoma californicum pluviale*) densities at the tree level showed delayed

density-dependent infection rates by a nucleopolyhedrovirus (Rothman, 1997).

Besides host density, other aspects of the local environment may affect viral transmission and

dynamics in the field (Cory & Myers, 2003). In particular, ultraviolet radiation has been shown to

inactivate viruses of all kinds (Sagripanti & Lytle, 2007), including baculovirus occlusion bodies

(Griego et al., 1985; Witt & Stairs, 1975). In field studies, examination of the presence of baculovirus

occlusion bodies on shaded vs. unshaded foliage suggested that sunlight on leaves may inactivate

viruses (Olofsson, 1988). A study of baculovirus transmission in forest tent caterpillars (*Malacosoma*

disstria) found depressed transmission rates on forest edges as opposed to the forest interior, which was

attributed to sunlight inactivating virus on leaves at forest edges (Roland & Kaupp, 1995). A field study

of two strains of gypsy moth NPV showed variable resistance of virus to ultraviolet, in which the

inactivation rate from ambient ultraviolet was greater in a more potent strain than a less potent one

(Akhanaev et al., 2017). In another study of forest tent caterpillars using tree ring analyses, it was

suggested that periods of weaker ultraviolet radiation increased the effect of forest tent caterpillar

outbreaks on tree growth (Haynes et al., 2018). In this example, less ultraviolet radiation may have

allowed better viral persistence and transmission and produced more severe caterpillar outbreaks.

In the present study, we examined the effects of host density, delayed-density dependence, and

ultraviolet radiation on the survival, infection rate, and infection severity of a newly discovered

baculovirus in Ranchman's tiger moth (*Arctia virginalis*; Erebidae:Arctiinae). Ranchman's tiger moth

is a univoltine species with a long larval stage, during which it is a generalist herbivore (Karban et al.,

2010). Caterpillars are parasitized by tachinid parasitoids, though caterpillars sometimes survive

parasitism (English-Loeb et al., 1990). Parasitism has been found to have little role in population

dynamics, leaving the observed oscillatory dynamics largely unexplained (Karban & de Valpine, 2010).

Baculovirus is a dsDNA virus that persists outside of the host in the environment within a protective

protein shell as an occlusion body (OB). Horizontal transmission can occur when caterpillars consume

OBs on food sources, which dissolve in caterpillars' alkaline guts and release infective virus particles

(Vega & Kaya, 2012). Virus particles infect the caterpillar starting from the epithelial tissue and then

move to other parts of the body, including the trachea, fat bodies, and hemolymph (Barrett et al., 1998),

potentially causing death of the caterpillar.

To test the potential role of baculovirus in delayed-density dependent population dynamics in

Ranchman's tiger moth and the influence of ultraviolet radiation on the persistence and transmission of

virus in moth populations, we censused 18 populations over two years along a gradient of latitude and

ultraviolet radiation and reared caterpillars from these populations in the laboratory. To identify the

mechanisms through which baculovirus affects moth population dynamics, we measured baculovirus

infection rate, severity, and survival of caterpillars to adulthood. We analyzed the effect of caterpillar

density and ultraviolet radiation at the study sites on viral infection and survival. We also tested if viral

infection mediated the effects of caterpillar density and ultraviolet radiation on caterpillar survival

using a structural equation model. In addition, we characterized the baculovirus identified in

Ranchman's tiger moths using sequence data and microscopy.

Methods

Description of system

In California, Ranchman's tiger moths eclose as caterpillars in late summer, and feed and

develop until the following spring. Pupation occurs in late spring, and adults emerge in early summer,

mate and lay eggs during the summer. Early instars feed cryptically in litter and undergrowth until they

become larger and climb higher onto vegetation in winter months. Caterpillars are generalists and feed

on a variety of plants, with a preference for alkaloid-containing hosts (Karban et al., 2010), and occur

mostly in open grassland, shrubland, or savannah. At the end of the larval stage, caterpillar movement

increases, and caterpillars disperse away from patches to pupate (Grof-Tisza et al., 2014). Caterpillars

also sometimes show symptoms of baculovirus infection, which include poor growth, inactivity,

watery, opaque frass, regurgitation of milky fluid, and death. Caterpillars do not lyse at death; the

integument of cadavers remains relatively intact. In dissections, viral occlusion bodies have been found

in caterpillars, pupae, and adults. Therefore, it appears that viral infection may not always be lethal – in

some cases, sublethal infections may persist in adults and pass on to the subsequent generation,

although observations suggest that fecundity may be reduced in infected adults.

Identification of baculovirus

To describe the virus, we removed infected fat tissue from four caterpillars from two sites, and

amplified the *polh*, *lef-8*, and *lef-9* genes using oligonucleotides with universal primers attached for the

lef-8 and *lef-9* genes using PCR (following Jehle et al., 2006). PCR products were then sequenced. We

also smeared fat body tissue and stained with buffalo black and examined under a light microscope

with a 400x oil immersion lens. Sequencing results were poor for polh due to short DNA fragments, but

we conducted a BLAST search on NCBI using lef-8 and lef-9 gene sequences.

Censuses

Caterpillars were counted visually along 4 m wide linear transects of varying length (78-300m)

at 18 sites along a ~1000 km length adjacent to the Pacific coast in California, Oregon, and Washington

(Figure 1). Transects were surveyed in 2019 and 2020, starting from 1 March through 30 May. We

visited sites in each year from south to north, in accordance with the phenology of caterpillar

development so that caterpillars were surveyed when most were 4^{th} or 5^{th} instars (sampling dates in

Table S1). Transects were surveyed at a constant walking speed and all by the same observer (A. Pepi).

Density was estimated from caterpillar counts over the transect area (4m x length). During the second

year, sites were revisited within ten days of the calendar date of the first visit; however, this species is a

slow-growing caterpillar (roughly ten month development period), so density estimates were likely not

overly sensitive to small deviations in sampling date. In 2020, up to ca. 30 caterpillars per site (or less

if fewer were found) were collected haphazardly and brought back to the laboratory for rearing.

Rearing

Caterpillars were reared individually in 6 oz plastic souffle cups with fabric lids and kept in an

incubator at 18 ºC and 75% RH. Caterpillars were fed every 3-4 days with washed organic romaine

lettuce and yellow bush lupine (*Lupinus arboreus*) leaves collected from a part of Bodega Marine

Reserve without *Arctia virginalis*. Parasitoids that emerged from caterpillars were counted and

identified to family. Caterpillars that died during rearing were frozen at -20 ºC for subsequent

dissection, except for those that clearly died due to emerged parasitoids. Caterpillars were reared until

pupation (~50 days), and pupae were placed together by site of origin (no more than 10 per site) in 30

cm x 30 cm flight cages and allowed to emerge as moths, at room temperature. Pupae and moths were

reared until one month after the last moth emerged and sprayed once or twice weekly with water to

prevent desiccation. Adults do not feed. Dead pupae and moths were stored dry at room temperature for

subsequent dissection. Individuals that reached adulthood fully formed were counted as having

survived; individuals that died as caterpillars or failed to emerge from pupae or expand wings were

counted as not surviving.

Dissection and viral assays

After death, each individual was dissected to assess infection status. Two to six tissue samples

per insect were taken from the abdomen of adults and pupae or fat bodies of caterpillars, broken up

with forceps, smeared on a glass slide with water, and examined under a light microscope at 200x

magnification with phase contrast. Visible occlusion bodies (OB) ($<0.15\mu m$) were confirmed by

adding a drop of 1 M NaOH to the slide, which dissolves the viral protein coat, turning the OB

transparent (Lacey & Solter, 2012)(Figure S1). Tissue samples were taken until a positive NaOH test

was obtained; if no virus was found after six tissue samples, individuals were classified as uninfected.

Infection severity was rated as uninfected, low, medium, high, or very high, based on the

number of samples required to discover virus, the density of OBs in tissue smear samples, and the

coloration of hemolymph and fat body (Figure S2). Low severity infections took many samples (4-6)

to identify and had low densities of OBs. Medium severity infections took fewer samples (1-2) to

identify and had moderate densities of OBs. High severity samples had infections in all samples with

high densities of OBs, with organ systems in the body still intact and identifiable. Very high severity

infections had infections in all samples with very high densities of OBs, with internal organ systems

unidentifiable.

Ultraviolet irradiation

We calculated the cumulative ultraviolet radiation in 2019 for each site that OBs are likely to

have experienced. Daily values for UV radiation, specifically 'Erythemal Daily Dose' in Watts/m^2 from

OMI/Aura satellite data were accessed from NASA GES DISC (Hovila et al., 2014). Values were

extracted for each sampling location and the average value calculated from 360 to 180 days before the

collection date of caterpillars (~March-September at most sites; see Table S1 for collection dates). This

timing coincided with the season during which viral OBs would have persisted in the environment

outside of hosts.

Statistical analysis

First, we analyzed the density-dependence of infection rate (proportion of infected individuals)

and the influence of ultraviolet radiation on infection rate. The proportion of all individuals (N=208)

that were infected was analyzed in response to log caterpillar density per m^2 from transects in 2019

(density in the previous year) and 2020 (density in the same year) using beta-binomial generalized

linear mixed models with site as a random effect. To test for the effect of ultraviolet radiation, we

implemented a model with both ultraviolet radiation and log density, to control for the effect of density

on infection rate.

Second, we analyzed the density-dependence of infection severity, and the influence of UV

radiation on infection severity.   We analyzed infection severity (N=198 with severity rated) in response

to log density (same year and previous year) using cumulative link mixed models with site as a random

effect. As before, to test for the effect of UV radiation, we implemented a model with both UV

radiation and log density, to control for the effect of density on infection severity. To test for effects of

density and UV on infection severity independent of infection status, we implemented models using

only infected individuals (N=184). This allowed us to test whether different processes influenced

infection rate vs. infection severity. We implemented cumulative link mixed models with only log

density, and both log density and UV radiation as predictors as before.

Third, we tested for density-dependent survival and effects of UV radiation on survival. We

analyzed survival to emergence as fully formed moths (N=208) in response to caterpillar density,

tachinid parasitoid load, and UV radiation in a beta-binomial generalized linear mixed model with site

as a random effect.

Lastly, we tested our structural hypothesis that viral infection and infection severity were the

mechanisms through which lagged density and UV radiation affected survival in moth populations.

There are multiple possible pathways through which density, UV radiation, infection rate, and infection

severity might influence survival and thus population dynamics. To test our specific structural

hypothesis, we constructed a piecewise structural equation model which tested whether infection rate

and infection severity mediated the effects of UV radiation and caterpillar density on survival. To do

this, we transformed infection severity into a continuous variable between 0 and 1, by converting to

numerical categories from 1-5, dividing by 5, and subtracting 0.1. We constructed 3 component sub-

models of our structural equation model: a beta-binomial model of infection rate predicted by log

density and UV radiation, a beta model of infection severity rate predicted by infection rate, log density

and UV radiation, and a beta-binomial model of survival predicted by infection rate and infection

severity (Figure 5). All sub-models included site as a random effect, and only individuals that were not

parasitized and were scored for infection severity were included in the analysis (N=198). Standardized

coefficients were calculated using the latent-theoretic method (Grace et al., 2018).

Beta-binomial and beta generalized linear mixed models were fitted using glmmTMB (Brooks

et al., 2017). Cumulative link mixed models were fitted using the ordinal package (Christensen, 2019).

Piecewise structural equation models were fit using piecewiseSEM v. 1.2.1 (Lefcheck, 2016), and

component models were fit using glmmTMB. UV radiation was logged and scaled to improve model

convergence. We checked for multicollinearity in models using VIFs; multicollinearity was low to

moderate in all models (<6) except for same year density in models of infection severity (VIF from

10.5-11.5). Since we included same year density as a control variable, we kept these models as is,

despite high VIFs. Plots were generated using ggplot2 (Wickham, 2009), ggeffects (Lüdecke, 2018),

and viridis (Garnier, 2018). All analysis and plotting were conducted in R v. 3.6.3 (R Development

Core Team 2020).

Results

Description of virus

We successfully amplified *polh, lef-8* and *lef-9* genes with PCR from four infected individuals.

The BLAST search resulted in top matches of *Perigonia lusca* NPV and *Anticarsia gemmatalis* NPV

using lef-8 and lef-9 genes respectively (Table S2, S3). This result, along with size of occlusion bodies

under the microscope, indicated that the *Arctia virginalis* baculovirus was a nucleopolyhedrovirus since

both BLAST matches were alphabaculoviruses (NPV).

Infection rate

The probability of infection increased with log density in the previous year in models of

infection rate (Table 1, Figure 2a). Infection rate had little relation to log density in the current year or

to UV radiation over the season when environmental OBs could have been exposed (Table 1, Figure

2b).

Infection severity

Log density in the previous year increased infection severity in models of infection severity that

included both infected and uninfected caterpillars but this effect was weakened when UV radiation was

included in the model (Table 1, Figure 3a,b). This effect of UV radiation may have been due to low

power or to a negative correlation between log density in the previous year and UV radiation (Pearson's

r: -0.53). Ultraviolet radiation decreased infection severity when included (Table 1, Figure 3c,d). In

models that included only infected caterpillars, density in the current and previous years had little effect

on infection severity, and UV radiation had a negative effect on infection severity (Table 1).

Survival to adult emergence

Survival to adult emergence decreased with log density in the previous year and with the

number of emerged parasites. Survival increased in response to density in the current year, and in

response to UV radiation (Table 1, Figure 4).

Structural equation model

We hypothesized a causal model linking UV radiation and disease to caterpillar survival (Fig.

5). Shipley's d-separation test (Shipley, 2000) indicated that our structural equation model was

correctly specified (Fig. 5, Fisher's $C = 7.38$, $df = 6$, $P = 0.288$), meaning that there were no missing

paths in our causal model. The model results showed that the negative effect of log density in the

previous year on survival was mediated by infection rate and infection severity and that the positive

effect of UV radiation on caterpillar survival was mediated by infection severity (Fig 5). The relative

indirect negative effect size of log density in the previous year was ~2.5x greater than the positive

effect of UV radiation on survival (-0.085 vs. 0.035, calculated by multiplying path coefficients).

Discussion

Overall, our results show negative delayed density-dependent effects of the virus on caterpillar

population dynamics. Log density during the previous year increased infection rate and infection

severity and decreased survival to adult emergence. The effect of delayed density on both infection

rate and infection severity in univariate models and the result of the structural equation model suggest

that the effect of density on survival acts through infection severity as well as infection rate. This is

likely due to dose-dependent infection severity, in which caterpillars that are exposed to more viral

occlusion bodies develop more severe infections that result in death (Cabodevilla et al., 2011; Eberle et

al., 2008; Matthews et al., 2002).

In contrast, log density in the current year, representing direct density-dependence, had no effect

on infection rate, a marginal negative effect on infection severity, and a positive effect on survival to

adult emergence. Though inconsistent, this positive effect may have been due to ascertainment bias:

sites that had larger viral outbreaks may have already declined in density over the course of the larval

season (summer 2019-spring 2020). Therefore, density at the time of collection had potentially a

reversed relationship to environmental OB density: sites that had declined from earlier high densities

would have higher levels of OB, resulting in more severe infections and a higher probability of death.

Overall, our results indicated negative delayed density dependence and weak or positive direct density-

dependence consistent with the second-order, oscillatory population dynamics observed in the system

over the long term (Pepi et al., 2021).

Ultraviolet radiation by contrast had no detectable effect on infection rate but reduced infection

severity and increased caterpillar survival. The reduction in infection severity was even greater when

only infected individuals were considered. The effect of ultraviolet radiation on infection severity and

survival but not infection rates may represent dose-dependent effects of exposure to OBs. Specifically,

at sites with higher ultraviolet radiation, more viral OBs were likely inactivated before caterpillars

could be exposed to them, so caterpillars received smaller doses of virus and thus became less severely

infected and were more likely to survive.

The observational nature of our study gives it some strengths and weaknesses. Our study is one

of few to show delayed-density dependent viral infection in the field, and the only study to show effects

of radiation on viral infection in a natural population of insects. Our results provide validation in a

field system to laboratory studies of virus dynamics and inactivation effects of UV radiation on viruses

(Bjørnstad et al., 1998; Witt & Stairs, 1975). Despite highly variable conditions between sites, which

were spread across a gradient of over 1000 km, we were able to detect clear effects of delayed density

on infection rate, severity, and survival to adulthood, and of ultraviolet radiation on infection severity

and survival to adulthood. We were also able to use structural equation modelling to show that

infection rate and severity mediate the effect of delayed density and UV radiation on survival and

strengthen the inferences made from our observational study.

Density in the previous year was a strong predictor of infection rates in populations (Fig 2).

This finding makes sense because there must have been a large enough population of hosts in the

previous year for the disease to spread and produce sufficient inoculum to persist into the current year.

Density in the previous year also affected infection severity (Fig 3), but this effect became much

weaker when UV radiation was included in models and even weaker when only infected caterpillars

were considered. In the structural equation model, infection severity was well explained by infection

rate because individuals must be infected to have high infection severity. This result may be due to the

conflation of infection rate and infection severity in the model since the same dose-dependent

mechanism may have caused them. However, in all models of infection severity that included both UV

radiation and delayed density, the effect of delayed density, beyond whether individuals were infected

or not, was weaker than the effects of UV radiation. Thus, when effects on infection severity were

detected, they may have in fact been generated by the spurious negative relationship between delayed

density and UV radiation. In contrast to infection severity, UV radiation had no significant effect on

infection rate. Ultimately, both delayed density and UV radiation had strong and opposing effects on

survival to adulthood, though the effect of density in the previous year was much greater in magnitude.

The effects of delayed density on survival were mediated through infection with limited evidence of

direct effect on infection severity. In contrast the effects of UV radiation on survival were mediated

entirely through effects on infection severity.

A possible explanation for the finding that infection rates were primarily determined by host

density in the previous year but not by ultraviolet radiation is that viral infection may be determined in

part through vertical transmission (Burden et al., 2002; Cabodevilla et al., 2011; Cory, 2015; Cory &

Myers, 2003). In particular, vertical transmission is likely to occur in *A. virginalis* through persistent

sub-lethal infections. We regularly observed OB in egg samples in adult dissections, and thus eggs of

adults with sublethal infections are likely contaminated with virus. Vertical transmission represents a
mechanism through which density in the previous year might influence infection rates in the following
year but is not subject to influence from ultraviolet radiation. In this way, vertical transmission may
maintain higher infection rates after high-density years with epizootics, even if UV radiation attenuates
inoculum in the environment (Cabodevilla et al., 2011). Sublethal infections may represent another
means by which baculovirus affects population dynamics by imposing fitness costs on adults
(Cabodevilla et al., 2011; Matthews et al., 2002; Rothman, 1997), although we did not measure this in
the present study.

In the present study, we also give the first identification of a baculovirus pathogen of *Arctia
virginalis* (AvNPV). Based on lef-8 and lef-9 genes, we conducted a BLAST search which identified
this virus as a nucleopolyhedrovirus, which is consistent with the appearance of stained occlusion
bodies under a light microscope (personal communication, J.S. Cory). However, the likely placement
of AvNPV on the baculovirus phylogeny was not clear, since the two matched species, *Perigonia lusca*
NPV and *Anticarsia gemmatalis* NPV were in the NPV clades II and I, respectively (Thézé et al.,
2018). Further investigation and phylogenetic work will be necessary to identify the phylogeny of
Arctia virginalis NPV. Caterpillars infected with AvNPV exhibited unusual symptoms, such as failure
to lyse, and high survival rates despite heavy infection, which is atypical for NPV infections. In most
well studied species, NPV infection leads to death relatively rapidly, and causes caterpillars to lyse. In
the present case, this could be due to differences between AvNPV and other NPVs, such as in enzymes
that decompose the insect integument, or host differences. *Arctia virginalis* caterpillars are longer lived
and more physically robust than other better-studied lepidopteran baculovirus hosts, such as *Lymantria
dispar* or *Spodoptera* spp.. Further studies should investigate some of the mechanisms of host-pathogen
interactions between AvNPV and *Arctia virginalis* that lead to these unique characteristics, and their
consequences for viral transmission and host-pathogen dynamics.

In summary, we demonstrated population-level delayed density-dependent effects on viral

infection rate, infection severity, and survival to the reproductive stage, showing how viral infection

may drive cyclic dynamics in insects. We also showed for the first time that UV radiation may

influence disease dynamics and ultimately population dynamics of insects, through decreased infection

severity and increased insect survival rates. This provides support for the proposal that viral epizootics

may be an important mechanism driving cyclic dynamics of insects and Lepidoptera in particular

(Myers & Cory, 2013). Our findings also suggest that UV radiation may be an important factor to

consider as a driver of insect viral disease in the context of global change. Long-term changes in

atmospheric transmittance of solar radiation, or "global dimming and brightening," have been observed

and are potentially anthropogenic in origin (Wild, 2016). Such long-term changes may have the

potential to influence insect viral disease and population dynamics through changing levels of UV

attenuation of viral inoculum (Haynes et al., 2018).

Acknowledgements

We would like to thank Marcel Holyoak, Jenny Cory, and Jay Rosenheim for helpful comments

on the manuscript. We would also like to thank California State Parks, Ben Becker at Point Reyes

National Seashore, Kristen Ward at Golden Gate National Recreation Area, Jackie Sones at Bodega

Marine Reserve, Brendan Leigh at Humboldt Bay NWR, Karlee Jewell at North Coast Landtrust,

Diane Steeck at the City of Eugene, Grey Wolf at the City of Salem, Oregon Metro Parks, and Christian

Haaning at the City of Portland for assistance and access to field sites. We would also like to thank

Harry Kaya and Jenny Cory for assistance with identifying the baculovirus, Jay Rosenheim for the use

of incubators, Neal Williams for the use of compound microscopes, and Claire Beck and Jasmine

Daragahi for help with rearing caterpillars. The research was funded by NSF-LTREB-1456225 and an

NSF REU supplement (DEB-2018169).

Author contributions

AP, VP, and RK conceived the study. AP and VP collected the data and conducted the analyses.

DR and VM conducted genetic work. AP wrote the manuscript. All authors contributed critically to the

drafts and approved final publication.

Data Availability

Data will be archived on KNB upon acceptance.

References

Akhanaev, Y. B., Belousova, I. A., Ershov, N. I., Nakai, M., Martemyanov, V. V, & Glupov, V. V.

(2017). Comparison of tolerance to sunlight between spatially distant and genetically different

strains of Lymantria dispar nucleopolyhedrovirus. *Plos One*, *12*(12), e0189992.

Anderson, R. M., & May, R. M. (1980). Infectious Diseases and Population Cycles of Forest Insects.

Science, *210*(4470), 658–661.

Anderson, R. M., & May, R. M. (1981). The population dynamics of microparasites and their

invertebrate hosts. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

291(1054), 451–519.

Barrett, J. W., Brownwright, A. J., Primavera, M. J., & Palli, S. R. (1998). Studies of the

nucleopolyhedrovirus infection process in insects by using the green fluorescence protein as a

reporter. *Journal of Virology*, *72*(4), 3377–3382.

Berryman, A. A. (1996). What causes population cycles of forest Lepidoptera? *Trends in Ecology and*

Evolution, *11*(1), 28–32. https://doi.org/10.1016/0169-5347(96)81066-4

Bjørnstad, O. N., Begon, M., Stenseth, N. C., Falck, W., Sait, S. M., & Thompson, D. J. (1998).

Population dynamics of the indian meal moth: Demographic stochasticity and delayed

regulatory mechanisms. *Journal of Animal Ecology*, *67*(1), 110–126.

https://doi.org/10.1046/j.1365-2656.1998.00168.x

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H.

J., Mächler, M., & Bolker, B. M. (2017). GlmmTMB balances speed and flexibility among

packages for zero-inflated generalized linear mixed modeling. *R Journal*, *9*(2), 378–400.

https://doi.org/10.32614/rj-2017-066

Burden, J. P., Griffiths, C. M., Cory, J. S., Smith, P., & Sait, S. M. (2002). Vertical transmission of
sublethal granulovirus infection in the Indian meal moth, Plodia interpunctella. *Molecular
Ecology*, *11*(3), 547–555.

Burthe, S., Telfer, S., Lambin, X., Bennett, M., Carslake, D., Smith, A., & Begon, M. (2006). Cowpox
virus infection in natural field vole Microtus agrestis populations: Delayed density dependence
and individual risk. *Journal of Animal Ecology*, *75*(6), 1416–1425.

https://doi.org/10.1111/j.1365-2656.2006.01166.x

Cabodevilla, O., Villar, E., Virto, C., Murillo, R., Williams, T., & Caballero, P. (2011). Intra- and
intergenerational persistence of an insect nucleopolyhedrovirus: Adverse effects of sublethal
disease on host development, reproduction, and susceptibility to superinfection. *Applied and
Environmental Microbiology*, *77*(9), 2954–2960. https://doi.org/10.1128/AEM.02762-10

Christensen, R. H. B. (2019). *ordinal—Regression Models for Ordinal Data* (R package version
2019.12-10).

Cory, J. S. (2015). Insect virus transmission: Different routes to persistence. *Current Opinion in Insect
Science*, *8*, 130–135.

Cory, J. S., & Myers, J. H. (2003). The Ecology and Evolution of Insect Baculoviruses. *Annual Review
of Ecology, Evolution, and Systematics*, *34*, 239–272.

https://doi.org/10.1146/annurev.ecolsys.34.011802.132402

Eberle, K. E., Asser-Kaiser, S., Sayed, S. M., Nguyen, H. T., & Jehle, J. A. (2008). Overcoming the
resistance of codling moth against conventional Cydia pomonella granulovirus (CpGV-M) by a
new isolate CpGV-I12. *Journal of Invertebrate Pathology*, *98*(3), 293–298.

https://doi.org/10.1016/j.jip.2008.03.003

English-Loeb, G. M., Karban, R., & Brody, A. K. (1990). Arctiid larvae survive attack by a tachinid

parasitoid and produce viable offspring. *Ecological Entomology*, *15*(3), 361–362.

Fleming, S. B., Kalmakoff, J., Archibald, R. D., & Stewart, K. M. (1986). Density-dependent virus

mortality in populations of Wiseana (Lepidoptera: Hepialidae). *Journal of Invertebrate*

Pathology, *48*(2), 193–198. https://doi.org/10.1016/0022-2011(86)90123-0

Fuxa, J. R. (2004). Ecology of insect nucleopolyhedroviruses. *Agriculture, Ecosystems & Environment*,

103(1), 27–43.

Garnier, S. (2018). *viridis: Default Color Maps from "matplotlib"* (R package version 0.5.1.).

Grace, J. B., Johnson, D. J., Lefcheck, J. S., & Byrnes, J. E. K. (2018). Quantifying relative

importance: Computing standardized effects in models with binary outcomes. *Ecosphere*, *9*(6),

e02283.

Griego, V. M., Martignoni, M. E., & Claycomb, A. E. (1985). Inactivation of nuclear polyhedrosis virus

(Baculovirus subgroup A) by monochromatic UV radiation. *Applied and Environmental*

Microbiology, *49*(3), 709–710. https://doi.org/10.1128/aem.49.3.709-710.1985

Grof-Tisza, P., Holyoak, M., Antell, E. H., & Karban, R. (2014). Predation and associational refuge

drive ontogenetic niche shifts in an arctiid caterpillar. *Ecology*, *96*(1), 140703220532006.

https://doi.org/10.1890/14-1092.1

Haynes, K. J., Tardif, J. C., & Parry, D. (2018). Drought and surface-level solar radiation predict the

severity of outbreaks of a widespread defoliating insect. *Ecosphere*, *9*(8).

https://doi.org/10.1002/ecs2.2387

Hovila, J., Arola, A., & Tamminen, J. (2014). *OMI/Aura Surface UVB Irradiance and Erythemal Dose*

Daily L2 Global Gridded 0.25 degree x 0.25 degree V3,. NASA Goddard Space Flight Center,

Goddard Earth Sciences Data and Information Services Center.

https://doi.org/10.5067/Aura/OMI/DATA2028

Jehle, J. A., Lange, M., Wang, H., Hu, Z., Wang, Y., & Hauschild, R. (2006). Molecular identification

and phylogenetic analysis of baculoviruses from Lepidoptera. *Virology*, *346*(1), 180–193.

https://doi.org/10.1016/j.virol.2005.10.032

Karban, R., & de Valpine, P. (2010). Population dynamics of an Arctiid caterpillar-tachinid parasitoid

system using state-space models. *Journal of Animal Ecology*, *79*(3), 650–661.

https://doi.org/10.1111/j.1365-2656.2010.01664.x

Karban, R., Karban, C., Huntzinger, M., Pearse, I. S., & Crutsinger, G. M. (2010). Diet mixing

enhances the performance of a generalist caterpillar , Platyprepia virginalis. *Ecological*

Entomology, *35*, 92–99. https://doi.org/10.1111/j.1365-2311.2009.01162.x

Lacey, L. A., & Solter, L. F. (2012). Initial handling and diagnosis of diseased invertebrates. In L. A.

Lacey (Ed.), *Manual of techniques in invertebrate pathology* (2nd ed., pp. 1–14). Elsevier Ltd.

Lefcheck, J. S. (2016). PiecewiseSEM: Piecewise structural equation modelling in r for ecology,

evolution, and systematics. *Methods in Ecology and Evolution*, *7*(5), 573–579.

https://doi.org/10.1111/2041-210X.12512

Lüdecke, D. (2018). ggeffects: Tidy data frames of marginal effects from regression models. *Journal of*

Open Source Software, *3*(26), 772.

Matthews, H. J., Smith, I., & Edwards, J. P. (2002). Lethal and sublethal effects of a granulovirus on

the tomato moth Lacanobia oleracea. *Journal of Invertebrate Pathology*, *80*(2), 73–80.

https://doi.org/10.1016/S0022-2011(02)00100-3

Myers, J. H. (2000). Population fluctuations of the western tent caterpillar in southwestern British

Columbia. *Population Ecology*, *42*(3), 231–241. https://doi.org/10.1007/PL00012002

Myers, J. H., & Cory, J. S. (2013). Population Cycles in Forest Lepidoptera Revisited. *Annual Review*

of Ecology, Evolution, and Systematics, *44*(1), 565–592. https://doi.org/10.1146/annurev-

ecolsys-110512-135858

468 Myers, J. H., & Cory, J. S. (2016). Ecology and evolution of pathogens in natural populations of

469  Lepidoptera. *Evolutionary Applications*, *9*(1), 231–247. https://doi.org/10.1111/eva.12328

470 Olofsson, E. (1988). Environmental persistence of the nuclear polyhedrosis virus of the European pine

471  sawfly in relation to epizootics in Swedish Scots pine forests. *Journal of Invertebrate*

472  *Pathology*, *52*(1), 119–129. https://doi.org/10.1016/0022-2011(88)90110-3

473 Pepi, A., Holyoak, M., & Karban, R. (2021). Altered precipitation dynamics lead to a shift in herbivore

474  dynamical regime. *Ecology Letters*, *24*(7), 1400–1407. https://doi.org/10.1111/ele.13756

475 Roland, J., & Kaupp, W. J. (1995). Reduced transmission of forest tent caterpillar (Lepidoptera:

476  Lasiocampidae) nuclear polyhedrosis virus at the forest edge. *Environmental Entomology*,

477  *24*(5), 1175–1178. https://doi.org/10.1093/ee/24.5.1175

478 Rothman, L. D. (1997). Immediate and delayed effects of a viral pathogen and density on tent

479  caterpillar performance. *Ecology*, *78*(5), 1481–1493. https://doi.org/10.1890/0012-

480  9658(1997)078[1481:IADEOA]2.0.CO;2

481 Sagripanti, J. L., & Lytle, C. D. (2007). Inactivation of influenza virus by solar radiation.

482  *Photochemistry and Photobiology*, *83*(5), 1278–1282. https://doi.org/10.1111/j.1751-

483  1097.2007.00177.x

484 Shipley, B. (2000). A New Inferential Test for Path Models Based on Directed Acyclic Graphs A New

485  Inferential Test for Path Models Based on Directed Acyclic Graphs. *Structural Equation*

486  *Modelling*, *7*(2), 206–218. https://doi.org/10.1207/S15328007SEM0702

487 Tatalovich, Z., Wilson, J. P., & Cockburn, M. (2006). A Comparison of Thiessen Polygon, Kriging, and

488  Spline Models of Potential UV Exposure. *Cartography and Geographic Information Science*,

489  *33*(3), 217–231. https://doi.org/10.1559/152304006779077318

490 Thézé, J., Lopez-Vaamonde, C., Cory, J. S., & Herniou, E. A. (2018). Biodiversity, Evolution and

491  Ecological Specialization of Baculoviruses: A Treasure Trove for Future Applied Research.

492  *Viruses*, *10*(7), 366. https://doi.org/10.3390/v10070366

Turchin, P. (1990). Rarity of density dependence or population regulation with lags? *Nature*,

344(6267), 660–663. https://doi.org/10.1038/344660a0

Turchin, P. (2003). *Complex Population Dynamics: A Theoretical/Empirical Synthesis*. Princeton

University Press.

Vega, F. E., & Kaya, H. K. (2012). *Insect pathology*. Academic press.

Wickham, H. (2009). *ggplot2: Elegant graphics for data analysis* (1.0.0). Springer.

Wild, M. (2016). Decadal changes in radiative fluxes at land and ocean surfaces and their relevance for

global warming. *Wiley Interdisciplinary Reviews: Climate Change*, *7*(1), 91–107.

https://doi.org/10.1002/wcc.372

Witt, D. J., & Stairs, G. R. (1975). The effects of ultraviolet irradiation on a Baculovirus infecting

Galleria mellonella. *Journal of Invertebrate Pathology*, *26*(3), 321–327.

https://doi.org/10.1016/0022-2011(75)90229-3

509   **Figures & Tables**

510   **Table 1.** Results of univariate models.

Model structure	Parameter	Logit $\beta \pm 1$ SE	z	P
Infection ~ log density	Log density current year	-0.20±0.38	-0.52	0.603
	Log density previous year	**0.55±0.26**	**2.10**	**0.036**
Infection ~ log density + ultraviolet	Log density current year	-0.10±0.39	-0.25	0.802
	Log density previous year	**0.92±0.46**	**1.998**	**0.046**
	Ultraviolet radiation	0.66±0.58	1.13	0.259

Severity ~ log density	Log density current year	-0.28±0.19	-1.49	0.135
	Log density previous year	**0.26±0.11**	**2.35**	**0.019**
Severity ~ log density + ultraviolet	Log density current year	-0.28±0.18	-1.61	0.101
	Log density previous year	0.16±0.11	1.37	0.170
	Ultraviolet radiation	**-0.35±0.17**	**-2.04**	**0.041**
Severity | infected ~ log density	Log density current year	-0.27±0.19	-1.39	0.163
	Log density previous year	0.018±0.11	1.66	0.096
Severity | infected ~ log density + ultraviolet	Log density current year	-0.28±0.18	-1.52	0.129
	Log density previous year	0.06±0.12	0.48	0.631
	Ultraviolet radiation	**-0.44±0.18**	**-2.40**	**0.017**
Survival ~ log density + ultraviolet	**Log density current year**	**1.15±0.51**	**2.25**	**0.024**
	Log density previous year	**-0.75±0.31**	**-2.39**	**0.017**
	Ultraviolet radiation	**0.64±0.32**	**1.97**	**0.049**
	Emerged parasites	**-5.39±1.58**	**-3.42**	**0.0006**

Figure 1. Map of study site locations in Washington, Oregon, and California (orange points), overlaid

on a map of mean annual UV-B radiation aggregated at the county level, in Wh/m^2 (Tatalovich et al.,

2006).

Figure 2. Infection rate in response to (a) density in the previous year and (b) ultraviolet radiation as

scaled log erythemal daily dose in the summer before collection (2019) from beta-binomial generalized

linear mixed models.

Figure 3. Infection severity in response to (a-b) density in the previous year, and (c-d) ultraviolet radiation as scaled log erythemal daily dose in the summer before collection cumulative link mixed models. The left column (a,c) shows model-predicted rates of infection at each severity class with 95% confidence intervals, and the right column (b,d) shows the proportion of the total population predicted to be infected at each severity class by color.

554   **Figure 4**. Survival to adult emergence in response to (a) density in the previous year, and (b) ultraviolet

555   radiation as scaled log erythemal daily dose in betabinomal generalized linear mixed models. The size

556   of each point is scaled to the number of individuals reared from each site.

566   **Figure 5**. Path diagram of piecewise structural equation model. The size of the arrows is proportional

567   to the standardized path coefficient (shown next to lines), and line type indicates P-value of the path

568   parameter.

579 **Supplement**

580 **Table S1.**

Site	Latitude	Longitude	Number reared	Date Collected
Franklin Point, CA	37.155	-122.35	29	3/1/2020
Pescadero Marsh, CA	37.265	-122.41	16	3/1/2020
Golden Gate NRA, CA	37.844	-122.55	28	3/13/2020
Point Reyes South, CA	38.038	-122.99	15	3/13/2020
Point Reyes North, CA	38.088	-122.97	29	3/13/2020
Point Reyes Kehoe, CA	38.153	-122.94	2	3/13/2020
Bodega Marine Reserve, CA	38.319	-123.06	16	3/27/2020
Manchester State Park, CA	38.959	-123.72	6	3/27/2020
Loyalton, CA	39.700	-120.30	6	5/30/2020
Humboldt NWR, CA	40.670	-124.21	20	4/10/2020
Freshwater Farms, CA	40.786	-124.09	11	4/10/2020
Klamath River, Yreka, CA	41.829	-122.61	11	4/10/2020
Checkermallow Access, OR	44.072	-123.20	12	4/11/2020
Fairview Wetland, OR	44.895	-123.00	20	4/11/2020
Graham Oaks, OR	45.299	-122.80	30	4/11/2020
Powell Butte, OR	45.488	-122.50	33	4/11/2020
Salish Ponds, OR	45.531	-122.45	1	4/11/2020
West Rocky Prairie, WA	46.887	-122.87	2	4/12/2020

596 **Table S2.** Blast results for viral genes amplified from infected caterpillars.

Sample ID	Gene	Matched taxon	Max Score	Total Score	Query cover-age	E value	Percent identity
Loyalton 6	lef-8	Perigonia lusca single NPV	80.5	80.5	9%	2E-10	87.69%
Loyalton 6	lef-9	Anticarsia gemmatalis MNPV	130	130	84%	9E-26	77.16%
Loyalton 24	lef-8	Perigonia lusca single NPV	78.7	78.7	9%	9E-10	87.69%
Loyalton 24	lef-9	Anticarsia gemmatalis MNPV	130	130	84%	9E-26	77.16%
Graham Oaks 10	lef-8	Perigonia lusca single NPV	78.7	78.7	9%	9E-10	87.69%
Graham Oaks 10	lef-9	Anticarsia gemmatalis MNPV	124	124	84%	4E-24	76.86%
Graham Oaks 20	lef-8	Perigonia lusca single NPV	78.7	78.7	9%	9E-10	87.69%
Graham Oaks 20	lef-9	Anticarsia gemmatalis MNPV	117	117	84%	7E-22	76.42%

598 **Table S3.** Primers and sequences for viral genes amplified from infected caterpillars
599

Sample ID	Gene	Primer	Sequences
Loyalton 6	lef-8	**aggaaacagc-tatgacc**caygghgar-atgac	aacgcacgtttacccaccgtcaattacaatatgatggcggacaak-tttaaatttttactgtcacaaaaattaatacagca cgtcgacaatattaacggcgacgacaacagcaacacgttattaatt-gcgttcaacaatcggccgacaaatttaaaatgtt ttaaaaaagatgtgtacacaatcgtgtacacaatgaaac-gtaacatgtcgcccgtcgaattaaaaattaacgataaaatt ttatttgtcaaccatcacgaaggaatggtgtgcattaaaaaagtttt-gatgatcgatgacaacgtgcaagtgaacgcgct gctcaccccctacgagtatcactatcaacaatccatttatcgcaacggccagtg-taaatgtcgcattatcgacgaaaacg acaatgtcacatcgcttatgtccaaattggagcagtattattatagaaattttatacattt-gtttcatacagtgcctgtg cctaaattaattgtgtcgttgaccaaccttaaaaacgccatgcccgtgttcggttacgag-gaaaacaacgccacgttttg gcttcgctgtccactggctattcggtggcggcgcacaagtcaatagtgatgaacaa-taaaatgtttaaactatggacgtt
Loyalton 6	lef-9	**tgtaaaacgac-ggccagt**ttgtcdccrt-crcartc	ggtgcgcgacaacaaactcatgaccgccgargayccgtayrtggtcatagcttgtttcctga aacccacttttacgttccaatctgtaaaactatcgtttgtgcgcacatctgaaactt-gtgtgctcaattggctgatgttg ggatgacgcgtagtccaagcgcgcacattgtccacgtcacgtccataatagcgtt-gcaaactagcgcgcggcggtacaat ttcattaacaccattgaaacattgcacgttggcgtaaaaagaggcatcgttaa-gaaatgtcgaatacgaaaattgtacsg crtacccrttyttggtcatagctgtttccta
Loyalton 24	lef-8	**aggaaacagc-tatgacc**caygghgar-atgac	tcccgagctgacccgacctgcttaagatcatcaccatcccgaaattgccaattatac-ggacatggaaatasctgtttccg

Loyalton 24	lef-9	**tgtaaaacgac-ggccagt**ttgtcdccrt-crcartc	tmaacaagttaaactgacgacgacgccaacacgttcttaattgcgttcggrwtckgscgac-maattaaacatgtttgawc aaagatgtgtacacaatcgtgtacacaatgaaacgtaacatgtcgcccgt-craattaaaaattaacgataaaattttatt tgtcaaccatcacgaaggaatggtgtgcattaaaaaagtttttgatgatcgatgacaacgtg-caagtgaacgcgctgctca ccccctacgagtatcactatcaacaatccatttatcgcaacggccagtgtaaatgtcg-cattatcgacgaaaacgacaat gtcacatcgcttatgtccaaattggagcagtattattatagaaattttatacattt-gtttcatacagtgcctgtgcctaa attaattgtgtcgttgaccaaccttaaaaacgccatgcccgtgttcggttacgag-gaaaacaacgccaacgttttggctt cgctgtccactggctattcggtggcggcgcacaagtcaatagtgatgaacaa-taaaatgtttaaactatggacgttggtg cgcgacaacaaactcatgaccgccgaagayccstayrtggtcatagcttgtttcctga aacccacttttacgttccaatctgtaaaactatcgtttgtgcgcacwtctgaaactt-gtgtgctcaattggctgatgttg ggatgacgcgtagtccaagcgcgcacattgtccacgtcacgtccataatagcgtt-gcaaactagcgcgcggcggtacaat ttcattaacaccattgaaacattgcacgttggcgtaaaaagaggcatcgttaa-gaaatgtcgaatacgaaaattgtacsg crtacccrttyttggtcatagctgtttcctga
Graham Oaks 10	lef-8	**aggaaacagc-tatgacc**caygghgar-atgac	cgaccgyacgtgtkwcmyamcgtgtwttayaatatgatggcg-gacaggtttaaattttactgtcacaaaaattaataca gcacgtcgacaatattaacggcgacgacaacagcaacacgttattaatt-gcgttcaacaatcggccgasaaatttaaaat gttttaaaaaagatgtgtacacaatcgtgtacacaatgaaac-gtaacatgtcgcccgtcgaattaaaaattaacgataaa attttatttgtcaaccatcacgaaggaatggtgtgcattaaaaaagtttt-gatgatcgatgacaacgtgcaagtgaacgc gctgctcacccccctacgagtatcactatcaacaatccatttatcgcaacggccagtg-taaatgtcgcattatcgacgaaa acgacaatgtcacatcgcttatgtccaaattggagcagtattattatagaaattttata-catttgtttcatacagtgcct gtgcctaaattaattgtgtcgttgaccaaccttaaaaacgccatgcccgtgttcggttac-gaggaaaacaacgccacgtt ttggcttcgctgtccactggctattcggtggcggcgcacaagtcaatagtgatgaacaa-taaaatgtttaaactatggac
Graham Oaks 10	lef-9	**tgtaaaacgac-ggccagt**ttgtcdccrt-crcartc	gttggtgcgcgacaacaaactcatgaccgccgaagayccstayrtggtcatasctgtttcctga cactttttacgttccaatctgtaaaactatcgtttgtgcgcacatctgaaactt-gtgtgctcaattggctgatgttgggat gacgcgtagtccaagcgcgcacattgtccacgtcacgtccataatagcgtt-gcaaactagcgcgcggcggtacaatttca ttaacaccattgaaacattgcacgttggcgtaaaaagaggcatcgttaagaaatgtcgaa-tacgaaaattgtacsgcrta cccrttyttggtcatagctgtttccta

Graham Oaks 20	lef-8	**aggaaacagc-tatgacc**caygghgar-atgac	cgatcgtasgtttacacaacggtattaygatatgatggcgkacaggtgtaaatt-gttactgtcacaaaaattaatacwgc acgtcgacmatrtkaacggckacgacaacagcaacacgttattaatt-gcgttcaacaakcggccgacaaatttaaaatgt tttaaaaaagatgtgtacacaatcgtgtacacaatgaaac-gtaacatgtcgcccgtcgaattaaaaattaacgataaart tttatttgtcaaccatcacgaaggaatggtgtgcattaaaaaagtttt-gatgatcgatgacaacgtgcaagtgaacgcgc tgctcacccccctacgagtatcactatcaacaatccatttatcgcaacggccagtg-taaatgtcgcattatcgacgaaaac gacaatgtcacatcgcttatgtccaaattggagcagtattattatagaaattttatacattt-gtttcatacagtgcctgt gcctaaattaattgtgtcgttgaccaaccttaaaaacgccatgcccgtgttcggttacgag-gaaaacaacgccacgtttt ggcttcgctgtccactggctattcggtggcggcgcacaagtcaatagtgatgaacaa-taaaatgtttaaactatggacgt
Graham Oaks 20	lef-9	**tgtaaaacgac-ggccagt**ttgtcdccrt-crcartc	tggtgcgcgacaacaaactcatgaccgccgaagayccstayrtggtcatagctgtttcctga acttttacggttccaatctgtaaaactatcgtttgtgcgcacatctgaaactt-gtgtgctcaattggctgatgttgggat gacgcgtagtccaagcgcgcacattgtccacgtcacgtccataatagcgtt-gcaaactagcgcgcggcggtacaatttca ttaacaccattgaaacattgcacgttggcgtaaaaagaggcatcgttaagaaatgtcgaa-tacgaaaattgtacsgcrta cccrttyttggtcatagctgtttccta

614 **Figure S1**. Fat body tissue smear from a caterpillar infected with GV under a light microscope at 200x

615 magnification with phase contrast. A drop of NaOH was added which moved through the sample. The

616 red arrows point to the boundary between NaOH solution and water. The yellow arrows point to the

617 OBs (Figure A) which turned transparent (Figure B) after they were dissolved by the strong base.

629   **Figure S2.** Ventral dissection of late instar *A. virginalis* larva in the order of infection severity, ranging

630   from (A) uninfected, (B) low severity, (C) medium severity, (D) high severity, to (E) very high severity.

631   The caterpillar's gut in figure A is missing. Caterpillars with higher infection severity have a cloudier

632   hemolymph and a darker and more yellowish fat body. They also often have large clusters of infected

633   tissues next to the midgut (e.g. Figure D).

634